# Title Page

**Manuscript title:** Genetically based variation in fitness and carbon assimilation among bur oak populations

**Author names:** Rea, L.M.S.[1], Ostrowsky, L.[1,2], Mohn, R.[3], Garner, M.[3,4], Worcester, L.[3], Lapadat[1], C., McCarthy, H.R.[5], Hipp, A.L.[3], Cavender Bares, J.[1,2]

**Affiliations:**

[1]Department of Ecology, Evolution, and Behavior, College of Biological Sciences, University of Minnesota, 140 Gortner Laboratory, 1479 Gortner Ave, St Paul, MN 55108

[2]Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford St, Cambridge, MA 02138

[3]Herbarium, The Morton Arboretum, Lisle, IL 60532

[4]Pritzker Lab, The Field Museum, 1400 S Lake Shore Dr, Chicago IL 60605, USA

[5]School of Biological Sciences, University of Oklahoma, Norman, OK 73019

**Corresponding Author:** L. Rea

**Key words/phrases:** local adaptation; reciprocal transplant; tree ecophysiology; Quercus macrocarpa; plasticity

# Abstract

Ongoing climate change will negatively impact tree populations unless they are able to acclimate to the changes in their local environment. Effective planning for climate adaptation management requires an understanding of the current state of local adaptation and physiological performance to assess whether populations are at risk of local extinction, determine if assisted migration is appropriate, and to select appropriate seed sources if intervention is needed. We established a new reciprocal transplant experiment (ACE, Adaptation to Climate and Environment) across a latitudinal gradient in North America to examine the impacts of warming on three bur oak (*Quercus macrocarpa*) populations across much of the species range. We established common gardens in Minnesota, Illinois, and Oklahoma with seedlings grown from seeds collected within 50 km of each of those locations from a total of sixty maternal families . We aimed to 1) assess local adaptation in each of the populations using survival and size as fitness metrics, and 2) evaluate physiological responses to different environments along the latitudinal gradient. We found that northern populations are maladapted to hotter climates as evidenced by their low survival, growth, and photosynthetic rates in the warmest common garden. The southernmost population had the highest survival rate, growth rate, and fitness of the three populations in the southernmost garden, providing evidence for local adaptation to the warmest site. However, conditions in the middle garden resulted in the highest fitness and best physiological performance for all populations. Growth and survival were correlated in the middle garden but were decoupled in the northern and southern gardens. This decoupling is likely due to stress associated with the relatively extreme climates in the gardens at the ends of the gradient that led to greater resource allocation to survival than growth. Our results suggest that southern seed sources may perform well in warmer conditions in the north brought on by climate change,

43   which has important implications for managers assisting broadly ranged tree species in adapting
44   to climate change.

## Introduction

Climate change is rapidly shifting the global climate to historically unprecedented
conditions, as evidenced by the warmest years on record occurring in the past ten years, (Tierney
et al., 2020). With warming comes other related climatic issues, including increasing intensity
and frequency of drought events (Seneviratne et al., 2023). Such shifts in climate are likely to
alter species ranges and ecosystem boundaries (Chen et al., 2011). Ranges of many tree species
have already contracted in response to warming and drought stress (Zhu et al., 2012). Their long
lifespan and sessile nature force tree populations to plastically acclimate or adapt to new climate
conditions, shift to higher latitudes with similar climate conditions that they evolved in, or
become locally extinct in the face of climate change (Aitken et al., 2008). Historical rates of tree
migration based on life history and dispersal data backed by molecular data indicate migration at
a maximum rate of 100 meters per year (McLachlan et al., 2005), though evidence suggests that
modern rates of northern range edge migration for *Quercus* species may average 500 to 1200
meters per year (Sittaro et al., 2017; Tang et al., 2024), which is slower than the current rates of
climate change and may lead to maladaptation at species range edges. Dispersal limitations on
the movement of tree species will necessitate population-level responses to novel climates
through phenotypic plasticity or adaptation (Franks et al., 2014; Hoffmann & Sgrò, 2011;
Huntley, 1989). Knowledge of local adaptation and plasticity in response to changes in climate
are critical to managing tree populations and ecosystems for future climates conditions.

Climate change will likely impact the survival and fitness of individuals, which will have
implications for the long-term persistence of locally adapted populations. Populations that have

higher fitness in their local environment than other populations, or higher fitness in their home

environment than away from their home environment are said to be locally adapted (Kawecki &

Ebert, 2004; Williams, 1966). Local adaptation in large populations of widespread species can

help maintain genetic variation within the species, which is critical in species' persistence

through changing climatic conditions. However, with climate change, populations may perform

better at higher latitudes than in their current environment leading populations to be maladapted

to their home environment (Browne et al., 2019; Etterson et al., 2020; Gorton et al., 2022; Sáenz-

Romero et al., 2017). Such adaptational lags to climate coupled with dispersal limitations may

necessitate anthropogenic intervention to ensure survival of populations that cannot tolerate

novel climate regimes in their current environment.

One conservation management goal to maintain tree species through climate change is to

enhance genetic diversity to improve ecosystem resilience through strategies such as favoring

existing genotypes that are better adapted to future climates (Butler et al., 2012). This

management strategy is commonly referred to as "assisted population migration", where genetic

material (i.e., seeds or pollen) from southern, warm adapted populations is transferred northward

through human intervention to improve persistence of species in a changing climate (Williams &

Dumroese, 2013). For successful assisted migration of populations in the face of climate change,

it is critical to ensure that plasticity and genetic effects that may lead to maladaptation in the

transferred populations are accounted for prior to migration (Nagamitsu & Shuri, 2021). It is

worth noting that what we now call "assisted migration" has been historically practiced by

indigenous people by moving seeds and plant material to create desirable landscapes (Krech et

al., 2004). Assisted migration can be a critical strategy in helping to prevent species extinction

and economic losses and to maintain ecosystem services (Abrams et al., 2021; Krech et al., 2004;

Williams & Dumroese, 2013).

Changes in growing season length along with increased stresses of heat and aridity may

lead to maladaptation in photosynthetic and growth rates and shifts in carbon allocation. Carbon

allocation is a trade-off amongst various carbon pools including growth, reproduction, defense,

symbionts in the rhizosphere, and secondary compounds or increased storage in response to cold

acclimation, and drought response (Herms & Mattson, 1992). At lower latitudes, growing

seasons are longer and cold stress is reduced. Slow life history strategies, in which plants grow

slowly but steadily, are often found at lower latitudes, and have long-lived leaves with slow,

prolonged carbon fixation over a longer time (Chabot & Hicks, 1982; Kikuzawa, 1991; Wright et

al., 2004). Slow life history strategies may be better adapted to drought associated with climate

change, as species with slow strategies are more resistant to drought and can maintain

productivity through intense drought (Oram et al., 2023).

At higher latitudes with shorter growing seasons, growth may compete with investment

in cold tolerance (MacArthur, 1984). Shorter growing season length in higher latitudes is thought

to select for higher carbon assimilation rates, allowing plants to compensate for the shorter

period favorable for growth and reproduction (Bosiö et al., 2014). Cold temperatures in higher

latitudes also increase the allocation to nonstructural carbohydrate reserves, which are critical for

spring bud break after the long, cold winter (Klein et al., 2016; Körner, 2003). The conversion of

photosynthetic carbon to biomass is mediated by other carbon demands that shift with changing

environmental conditions, including drought and freezing, that may affect growth rate (Körner,

2015; Reich & Oleksyn, 2008). Therefore, we expect higher photosynthetic rates and lower

relative growth rates in higher latitudes. These combined expectations lead to contrasting

predictions about how photosynthetic rates and growth rates should vary with growing season

length (see Fig. S1).

Local adaptation can be tested with reciprocal transplant experiments which separate

genetic from environmental (plastic) influences on phenotypes (Kawecki & Ebert, 2004;

Schwinning et al., 2022). We established a reciprocal transplant experiment using the deciduous,

broad-leaved tree species *Quercus macrocarpa L.* across a latitudinal gradient by collecting seeds

from maternal families across the gradient to establish the ACE (Adaptation to Climate and

Environment) experiment. *Quercus macrocarpa* is a useful species for such an experiment

because it has a broad latitudinal and ecological range. Consequently, it is likely to show

adaptive differentiation across its range of contrasting climatic conditions. Furthermore, *Q.*

*macrocarpa* is an ecologically important tree species that is an important food source for birds,

insects, and mammals (Tallamy, 2021). *Quercus macrocarpa* is also an important species in oak

savannas, a rare and threatened ecosystem, due to its fire resistance capabilities (Hoekstra et al.,

2005). We collected acorns from individual trees (half-sib families) within three populations

(Minnesota, Illinois, and Oklahoma) across a climatic gradient which spans a mean annual

temperature gradient of 6-16 °C. ACE was designed to reciprocally plant seedlings from each

family within each population into gardens in those three states using a quantitative genetics

design, such that seedlings from all families were randomized in replicated blocks in each

garden.

Previous predictions of the response to climate change are mixed for *Q. macrocarpa* in

Minnesota: the species is predicted to have high capacity to cope with low levels of climate

change, but high climate change levels are expected to decrease suitable habitat (Iverson et al.,

2019). Climate change may provide some benefit for populations in Minnesota due to the

expected longer growing seasons (Handler et al., 2014) leading to increased growth, as has been

shown in other temperate tree species (Pretzsch et al., 2014). There is a general consensus that

there will be changes in the amount and timing of precipitation in Minnesota, likely leading to

periods of high aridity coupled with increased temperature as well as periods of heavy rainfall

and flooding. Trees in Minnesota may be maladapted to these changes, which could result in

mortality (Galatowitsch et al., 2009; Handler et al., 2014). To develop clear management

strategies for ecologically important tree species such as *Q. macrocarpa*, it is critical to study

seedling survival and function to understand the future of tree populations because seedlings

often have a narrower climatic niche compared to adult trees (Dobrowski et al., 2015). Seedlings

may be more vulnerable to climate change, but their survival and establishment are critical to the

regeneration and survival of tree populations (Williams, 2021).

In this study, we tested for local adaptation and adaptive differentiation by reciprocally

transplanting seedlings from three populations in field conditions. We used components of fitness

and integrated measures of fitness using Aster models (Geyer et al. 2007, Shaw et al. 2008) to

test for adaptative differences among populations across gardens. We acknowledge that common

garden sites vary both climatically and edaphically and in many other respects, and are

experiencing ongoing climate change. We addressed the overarching questions: Do *Q.*

*macrocarpa* progeny from three bur oak tree populations have higher fitness and physiological

performance in the garden closest to their origin, or are they better adapted to a climate that has

shifted northward due to warming with climate change? If populations are locally adapted, we

expect them to have higher fitness and adaptive physiological performance when they are in their

home garden (home vs. away), and to have higher fitness than other populations in their home

garden (local vs. foreign) (Fig. S2). On the other hand, if populations are adapted to a climate

that has shifted northward due to the 1.1 °C warming in the past century (IPCC, 2023), we

expect them to have higher fitness and physiological performance in the more northern (higher

latitude) gardens compared to the hotter and drier southern (lower latitude) gardens, but northern

populations to exhibit lower fitness and higher physiological stress responses in the southern

garden. Because seedling transplants were grown in natural field conditions, there may be

edaphic differences among sites that can offer more favorable soil resource conditions at some

sites relative to others and complicate interpretations of local adaptation in relation to climate.

## Methods

164 Methods

**165 Reciprocal transplant experimental design**

The Adaptation to Climate and Environment (ACE) experiment was designed to test for

local adaptation by reciprocally transplanting bur oak seedlings from known families into three

garden sites across a latitudinal gradient. The design was based on prior work from Etterson and

Shaw (2001), Etterson (2004a, 2004b), Deacon and Cavender-Bares  (2015), and Ramirez-

Valiente et al. (2018). Acorns from Minnesota, Illinois, and Oklahoma populations were

collected in 2018 and 2019 from individual mother trees at least 100 feet apart to minimize the

likelihood of shared pollen among trees. Acorns collected from mothers in the same parkland

area were grouped into sub-populations. Both 2018 and 2019 collections were germinated

outdoors in the ground in a common environment in Vallonia, Indiana at the Vallonia State

Nursery through the Indiana Department of Natural Resources (see appendix S1:Methods for

supplemental details on acorn processing). In March 2021, seedlings were lifted from the

common nursery, sorted, and sent to be planted in the common gardens in tilled former

grasslands in Minnesota (Cedar Creek Ecosystem Science Reserve, East Bethel MN), Illinois

(Morton Arboretum, Lisle IL), and Oklahoma (Kessler Atmospheric and Ecological Field

Station, Purcell OK). In each of the common gardens we planted ten individuals from twenty

families from each of the three populations (Minnesota, Illinois, and Oklahoma) 1 m apart in a

randomized block design, with at least one individual from each family planted in each block

(Fig. 1a). Gardens were weeded or mowed to remove small herbs and grasses that were growing

around the trees, then mulched at the end of the first growing season or beginning of the second

to suppress herbaceous understory growth. The seedlings were irrigated for the first two years

after planting to ensure the gardens were established.

**Climate data**

Monthly temperature data from 1940 to 2023 were acquired from ERA5 data (Hersbach

et al., 2023) using garden coordinates, and extracted the data from the raster file using QGIS

(*QGIS Geographic Information System*, v3.34, 2024). Temperature data for each year were

averaged to calculate the mean annual temperature (MAT) and were averaged over the months of

May to September to calculate the growing season temperature for each year. To find the change

in MAT and growing season temperature since the middle of the 21$^{st}$ century, we calculated the

past mean for the years 1940-1960 and the recent mean 2003-2023 then subtracted the past mean

from the recent mean for each garden. We used a similar approach to extract precipitation data

for the years of 2021-2023, while the garden was planted.

To estimate differences in aridity between the gardens, we calculated an index of

moisture (IM) for each of the gardens following Ramírez-Valiente & Cavender-Bares (2017),

where the index of moisture is equal to the difference between monthly precipitation and

monthly potential evapotranspiration (PET).  The IM reflects the monthly water balance for an

area and indicates water deficit with negative values and water surplus with positive values. We

calculated PET with the Thornthwaite (1948) method, using monthly mean temperature and

average day length per month.

**Soil nutrients and texture**

To document differences in soil properties among sites, we sampled soil from the center

of each garden block to 10 cm depth and conducted tests for soil nutrients (carbon-nitrogen ratios

(C:N), total nitrogen content) and soil texture. To measure soil nutrients, we dried the soil and

sieved it with a 2 mm mesh sieve to remove large rocks and organic matter. To measure total soil

carbon and nitrogen concentration, we ground the soil to a homogenous fine powder and

analyzed with a Costech ECS4010 elemental analyzer (Valencia, California, USA). We

approximated the soil texture using the "texture by feel" method (Thien, 1979).

**Survival and growth surveys**

To determine seedling performance and fitness, the seedlings were surveyed for survival

and size after planting and yearly from 2021 to 2023 at the end of the growing season. The height

of the tallest living bud and basal diameter measured with a caliper 5 cm from the ground were

used as metrics of size. Stem volume (SV) was estimated using a conoid equation as a proxy for

aboveground biomass.

We used conoid volume (CV):

(1)     $CV = H \times \pi \times D_{rc}^2 / 12$

where H is the height of the tree from the root collar to the apical meristem, and $D_{rc}$ is stem

diameter at the root collar.

Relative growth rate was calculated as

(2)     $RGR = \ln(V_{final}) - \ln(V_{initial})/T_{final} - T_{initial}$

where V is stem volume and T is time in total days. We calculated percent survival of maternal

family groups in each garden as the number of trees alive out of the total trees originally planted

in each family group.

**Phenology**

In 2023, to determine growing season length, we tracked spring and fall phenology using

protocols established by the National Phenology Network (Denny & Crimmins, 2023). In the

spring, approximately weekly, we surveyed the trees in each garden and tracked if the trees were

breaking leaf bud. Leaves were considered to be breaking leaf bud if a green leaf tip was visible

at the end of the bud. At similar intervals in the fall, we tracked leaves that were browning and/or

falling after changing color as senescence. We considered senescing leaves to have late-season

color that were actively falling or had recently fallen from the tree.

We used the budburst and the senescence data to estimate the growing season length of

canopy foliage persistence, or how long foliage was present on the trees in a growing season. We

calculated this as the difference between the mean initial budburst date and the mean initial

senescence date to find the total length of the canopy foliage growing season.

**Gas exchange**

To measure photosynthesis rates and stomatal conductance, we conducted measurements

of leaf-level gas exchange on every living tree in each of the three gardens using a LI-6800

Portable Photosynthesis System (LI-COR, Lincoln NE, USA) to determine $CO_2$ assimilation (A)

and intrinsic water use efficiency (iWUE) (see appendix S1:Methods  for supplemental details on

settings and parameters used for measurement). This data was collected in both 2022 and 2023 at

similar points in the growing season and using the same settings and approach. We calculated

iWUE as a ratio of $CO_2$ assimilation over stomatal conductance, which is the rate of

photosynthetic $CO_2$ assimilation against the water loss from open stomata. Because A and iWUE

change over the course of the day with stomatal closure, we found the maximum for A, stomatal

conductance, and iWUE by excluding measurements after stomatal closure occurred (see Fig.

S3 for cutoff times in each garden for each year).

**Spectral indices**

We collected hyperspectral reflectance measurements at the leaf level on every living tree

in each garden to add additional, nondestructive metrics of chlorophyll content and nitrogen

content. We measured leaf reflectance on the uppermost, fully expanded, undamaged leaf on

each tree using a portable field spectrometer (SVC HR-1024i; Spectra Vista) and a leaf clip with

an internal light source (LC-RP PRO; Spectra Vista) (see appendix S1:Methods for supplemental

details on spectral data and processing).  Using the spectra, we calculated the chlorophyll

normalized difference index (ChlNDI)

$\quad$ (3) $\quad (\frac{R750-R705}{R750+R705})$

and the normalized difference nitrogen index (NDNI)

$\quad$ (4) $\quad (\frac{log(1/R1510)-log(1/R1680)}{log(1/R1510)+log(1/R1680)})$.

The ChlNDI is positively correlated with chlorophyll a content with minimal influence of

chlorophyll fluorescence (Gitelson & Merzlyak, 1994). The NDNI is positively correlated with

percent nitrogen of vegetation (Serrano et al., 2002).

**Trait variance partitioning**

We partitioned the variance of relative growth rate, spectral indices, and gas exchange

traits between population, sub-population (the site of acorn collection), and maternal family for

each garden. We used linear mixed models implemented with the lme4 package in R (Bates et

al., 2024). We specified the model with population, sub-population nested within population, and

maternal family nested within sub-population within population. We then extracted the variance

components and the residual variance and calculated the percent of the total variance explained

by each component.

**Statistical analysis**

*Aster modeling of fitness*

We used aster models to estimate fitness for each population in each garden. Aster models

are useful to estimate fitness as they enable integration of fitness components with different

statistical distributions by modeling the dependence of fitness components expressed later in life

on earlier fitness components (Geyer et al., 2007; Shaw et al., 2008). We modeled fitness using

survival to 2023 and height conditional to survival to the last year of measurements, as our

nodes, depicted graphically as:

(5) Planted$\rightarrow$ Survival to 2023$\rightarrow$ Height

We chose height as the fitness estimate because it fit a normally distribution, and is a useful

predictor of eventual fecundity (Cornelius, 1994; Gamache & Payette, 2004). Survival to 2023

was modeled with a binomial distribution and height was modeled with a normal distribution. As

predictors in the aster models, we included population (P), garden (G) and the interaction

between them, as well as the height at the time of planting (S) as a covariate to account for

variation in size that could influence fitness, as:

(6) $F = P \times G + S$

We used the anova() function  in the R software v. 4.3.0 (R Core Team, 2023) to compare models

with and without the interaction effect to test for significance of the interaction effect. Once we

settled on the model (equation 6), we obtained means and standard errors for each population in

each garden using the aster predict function for a dummy dataset with an "average individual"

representing each population-garden combination.

*Statistical testing*

We evaluated differences in $A_{max}$, ChlNDI, survival, and growth among and within

gardens for all populations using ANOVAs followed by Tukey post-hoc tests with a population

by garden interaction effect. Number of individuals in each population in each garden varied due

to survival and the number of measurements we were able to obtain (see supplemental Table S1

for number of individuals included in each measure). We also tested relationships between

photosynthetic physiology ($A_{max}$ and ChlNDI) with fitness components (survival and growth)

using linear models to understand the relationship between physiology and fitness components.

All analyses were conducted in R software v. 4.3.0 (R Core Team, 2023).

# Results

**Site climate differences**

The ERA5 dataset, indicated that the average monthly temperature decreased

latitudinally, with the Minnesota garden having the lowest average temperatures each month and

Oklahoma having the highest (Fig. 1b). Notably, the Oklahoma garden never had a monthly

average below freezing (0 °C), whereas the Minnesota and Illinois gardens did in the winter

months. We found that since 1940, temperatures have increased at each of the garden sites. Mean

annual temperature (MAT) has increased by over 1°C in each garden from averaged 20-year

periods of 1940-1960 to 2003-2023, and the amount of temperature change increased with

latitude, with the greatest change in MAT occurring in the northernmost, Minnesota garden, and

the lowest change in temperature in the southernmost garden, Oklahoma (Table S2). Over the

same period, the average growing season temperature has also increased. Like MAT, the greatest

change in growing season temperature was in Minnesota. However, unlike MAT, the change in

growing season temperature was lowest in Illinois, the middle garden.

The gardens also differed in precipitation and aridity over the year. The Oklahoma garden

peaked in precipitation in the springtime around April and May (Fig. 1c). The Minnesota garden

also had a small peak in precipitation at that time, however the highest monthly precipitation

occurred in late summer for the Minnesota garden around August and September. Precipitation in

the Illinois garden peaked mid-summer in June and July and was the highest precipitation of the

year among the three gardens. The Illinois garden was also the least arid of the three gardens

during the growing season, though the index of moisture was negative for all the gardens,

indicating a water deficit during the growing season (Fig. 1d). Notably, the index of moisture

was negative year-round for the Oklahoma garden, suggesting it was in an atmospheric water

deficit for the entirety of the year while the gardens were planted, though some of the water

stress may have been alleviated by irrigation in the first two years of growth.

**Fitness components**

We assessed two main components of fitness: survival and plant size. Survival rates of

family groups were most disparate in the Oklahoma garden, where survival of the Minnesota

population (MN), was significantly lower than the Illinois (IL) and Oklahoma (OK) populations .

The higher survival of OK compared to MN is in line with local vs. foreign local adaptation. (Fig

2a). Survival for MN and IL was highest overall in the Illinois garden. In the Minnesota garden,

survival was similar across populations and was similar to OK in the Oklahoma garden and to all

populations in the Illinois garden.

Trends in tree size growth varied depending on the aspect of size observed. Stem volume

combines the effects of diameter and height growth. The OK population had the highest stem

volume RGR in the Oklahoma garden but was similar to the IL population in the Illinois garden

(Fig. 2b). In the Minnesota garden, the stem RGR was similar among all populations. Stem

volume RGR was highest in the Illinois garden for all populations.

**Fitness**

We found significant ($p<0.001$) interaction effects of population and garden on fitness

using omnibus tests. Our model showed that aster fitness was highest in the Illinois garden for all

populations, with the OK population having the highest fitness, in line with our hypothesis of

climate change having shifted ranges northward. (Fig. 2c). In the Oklahoma garden, the OK

population had higher fitness than the northern populations, as expected with local vs. foreign

local adaptation. Fitness in the Minnesota garden was about the same for all populations.

**Physiology**

We evaluated photosynthetic responses in the different climates of each garden. The

maximum $CO_2$ assimilation ($A_{max}$) was highest for the MN and IL populations in the Illinois

garden (Fig. 2d). The MN and IL populations had similar $A_{max}$ rates to each other in all gardens

and were lower in the Oklahoma garden than in the northern gardens, as we hypothesized given

the aridity stress in the Oklahoma garden. The OK population had lower $A_{max}$ rates in both the

Minnesota and Illinois gardens compared to the northern populations but had higher $A_{max}$ rates in

the Oklahoma garden compared to the MN population. The OK population also had lower

chlorophyll content (as detected by the chlorophyll normalized difference spectral index

(ChlNDI)) in the Illinois and Minnesota gardens compared to the MN population (Fig. 2e). The

IL population generally was in between the OK and MN populations, with no significant

differences detected between populations. We found that spectral normalized difference nitrogen

index (NDNI) was highest for all populations in the Illinois garden, indicating higher leaf

nitrogen content (Fig. 2f). The OK population had lower leaf nitrogen content than the northern

populations in the Illinois garden. The NDNI was similar for all populations in the Oklahoma

and Minnesota gardens. Data from 2022 was significantly different from 2023 data, but indicated

similar though less strong trends, which may be an effect of irrigation or ontogeny. We include

2023 data here as the effects are stronger and indicate responses in natural conditions without

irrigation (see Fig. S4 for 2022 results).

**Soil quality**

Soils in Oklahoma and the Illinois gardens were very similar in carbon (C) and nitrogen

(N) concentrations, though concentrations in Oklahoma were a bit higher on average (table 2.1).

Soils in the Minnesota garden had low soil percent nitrogen and percent carbon. Minnesota had

the highest soil C:N, followed by the Illinois and then the Oklahoma gardens. The soils also

differed texturally in terms of silt, clay and sand fractions. Oklahoma garden soils were classified

as silty clay loam, Illinois as sandy clay loam, and Minnesota as sandy loam.

**Correlations of survival, growth, and physiology**

Stem volume growth was significantly positively correlated with survival for all

populations in the Illinois garden, but not in the Minnesota or Oklahoma gardens (Fig. 3a). We

found significant positive relationships of $A_{max}$ with stem volume growth for the OK and IL

populations in all the gardens, and for the MN population in the Illinois garden (Fig. 3b). We also

found positive correlations of ChlNDI with stem volume growth in the Illinois garden for all

populations, and for the MN and IL populations in the Minnesota garden (Fig. 3c). We did not

find significant relationships with survival for $A_{max}$ or ChlNDI, except for a negative relationship

between ChlNDI and survival for the MN population in Minnesota.

**Phenology and growing season**

Budburst was recorded earliest in the Oklahoma garden, and latest in the Minnesota

garden, on average. Of the three populations, the OK population tended to break bud earliest on

average, except in the Minnesota garden where all populations broke bud at similar times (Fig.

4a). The Minnesota garden had the earliest date of senescence, whereas the Oklahoma garden

was the latest (Fig. 4b). The OK population also tended to senesce later than the northern

populations, particularly in the Illinois garden. The Oklahoma garden had the longest growing

season length for all three populations, whereas Minnesota was the shortest (Fig. 4c).

**Variance partitioning of traits**

When we partitioned the variance for traits in each garden, we found that variance

attributable to population-level differences was distributed heterogeneously among traits and

differed among gardens. Generally, variance attributable to population-level differences was

greater in the Oklahoma and Illinois gardens, whereas in the Minnesota garden variation was

often more attributed to subpopulation or maternal family-level variance, with the exception of

ChlNDI (Fig. S5). One consistency in the variance partitioning was for $A_{max}$, population-level

variance accounted for the greatest percent of the variance compared to subpopulation or

maternal family-level variance. In the Oklahoma garden the majority of the variance in relative

growth rate (RGR_Vol) was attributed to population-level variation, whereas in the Minnesota

garden most of the variance was attributable to maternal family variation. In the Illinois garden,

stomatal conductance (gswMAX) had the highest population-level variance.

# Discussion

**Evidence for local adaptation**

We established the ACE (Adaptation to Climate and Environment) experiment, a

reciprocal transplant experiment along a latitudinal gradient, to test for local adaptation and

adaptive differentiation among populations of *Q. macrocarpa* from contrasting climates. Results

of this work can inform ongoing efforts to manage forests and assist population migration. For

all populations, we expected that local adaptation would be demonstrated by higher fitness of the

local population relative to other populations (local vs foreign) and higher fitness of each

population in its home location relative to its fitness in other locations (home vs. away). We

found evidence supporting local adaptation based on the local vs. foreign comparison in the

lowest latitude garden with the hottest climate. The OK source population had higher survival,

growth, and fitness in the Oklahoma garden, compared to the MN population. In addition, the

OK population had the highest maximum carbon assimilation rate ($A_{max}$) of the three populations

in the Oklahoma garden. The lower $A_{max}$ rate of the northern populations in the Oklahoma

garden indicate they were stressed in the hotter climate. The relative rates of $A_{max}$ reversed in

both of the northern gardens where the OK population had lower $A_{max}$ than the northern

populations. Drought is known to reduce stomatal conductance, which limits carbon assimilation

(Boyer, 1976). Previous findings from oaks have shown that $A_{max}$ is reduced when exposed to

chronic drought (Gimeno et al., 2008). Higher survival of the OK population in the Oklahoma

garden, which has the hottest climate relative, to the MN population provides evidence that the

OK population is better adapted to warm temperatures. Under the expected 2 °C warming with

climate change, the OK population may be better suited for the emerging warmer climates with

potential to expand northward (IPCC, 2023). Conversely, the decline  in assimilation rates in

seedlings from the IL and MN populations in the hotter and drier Oklahoma climate suggests that

these populations may experience declines in survival and fitness in their current environments.

While we found evidence of a local survival advantage relative to foreign populations in

the Oklahoma garden, all populations expressed highest fitness in the Illinois garden.

Consequently, we did not find evidence for local adaptation based on home vs. away

comparisons. We also did not find clear statistical evidence for local vs. foreign local adaptation

in the two northern populations after three years of growth. However, these patterns of local

adaptation could still emerge through time. The OK population had higher absolute growth rate,

which could be due in part to its larger acorn size (Table S3) with larger seed size supporting

growth and survival in its early years (Bischoff et al., 2006; Deacon & Cavender-Bares, 2015). In

addition, ontological differences in growth may have impacted our ability to observe signal of

local adaptation. We monitored seedlings transplanted to the gardens after germination in a

common nursery. Consequently, we were not able to measure all critical lifetime components of

fitness. For example, we did not account for differences in germination rates of the populations

in different gardens, nor could we observe fecundity as our trees had not reached reproductive

maturity. Observing differences in fitness components relevant at an older, reproductively mature

life stage might still reveal evidence of local adaptation not evident at the seedling life stage

(Wadgymar et al., 2022).

444   Our alternative hypothesis for the observation of high performance outside of

445 populations' local garden was that ongoing climate change has shifted suitable ranges of different

446 populations northward, leading southern populations to be better adapted to northern gardens.

447 While we saw that the OK population did perform better in the Illinois garden, which could be a

448 consequence of both edaphic and climatic factors, we did not see evidence that the IL population

449 performed better in the Minnesota garden compared to in its home garden, nor did it outperform

450 the local MN population in Minnesota. This result indicates that there may be additional

451 conditions or stressors in the Minnesota garden, such as freezing, nutrient deficiencies, and/or

452 insect herbivory, that influence performance of non-local populations.

453 **Adaptive differentiation in phenology and function**

454   We found differences in phenology and plant function in line with adaptive

455 differentiation, though the amount attributable to population-level differences versus

456 subpopulation or maternal family-level differences varied among gardens. Both bud break and

457 senescence differed significantly among populations in the Illinois garden. The OK population

458 broke bud earlier and senesced later in the Oklahoma and in Illinois gardens relative to other

459 populations, consistent with findings in other oak common gardens (Papper & Ackerly, 2021;

460 Wright et al., 2021). In the Minnesota garden, where the growing season was shortest, all

461 populations broke bud later and began senescence earlier compared to southern gardens (Fig. 4).

462   Population explained most of the variance in $A_{max}$, indicating that there are population-

463 level differences in $A_{max}$ that are evident across environments. We found evidence supporting

464 adaptive differentiation in photosynthetic rate in the OK population, which had lower $A_{max}$

465 values than the northern populations in the northern gardens but had highest $A_{max}$ values in its

466 home garden. $A_{max}$ was highest in the least arid Illinois garden, indicating that alleviation of

aridity, cold stress, and soil nitrogen limitation improved photosynthetic rates, as we

hypothesized. On the other hand, low $A_{max}$ in the Oklahoma garden indicated low performance.

The differences in $A_{max}$ likely were due to high temperature stress inhibiting photosynthesis

through increased enzyme inhibition and RuBisCo oxygenation (Moore et al., 2021). The $A_{max}$

was greater for the OK population in its home garden and provides evidence for adaptive

differentiation and heat tolerance in the OK population.

We expected that populations from more arid climates would have greater water use

efficiency and drought tolerance, however $iWUE_{max}$ differed little among populations, and had

more differentiation among gardens, and at the garden level, $iWUE_{max}$ in Oklahoma was

explained less by population-level differences, but rather maternal family-level differences were

most important. Similarly, variation in NDNI was accounted for by subpopulation (mother tree

site) level differences to a higher degree than the population level differences. The variation in

NDNI may be attributed to adaptation to local nutrient availability of the mother tree. Genetic

differentiation in drought related traits in oak common gardens has previously been indicated to

be lower than for phenological traits, consistent with our results (Torres-Ruiz et al., 2019). NDNI

and $iWUE_{max}$ in Oklahoma were explained less by population-level differences, but rather

maternal family-level differences were most important for $iWUE_{max}$, and NDNI was partially

accounted for by subpopulation (mother tree site) level differences. The variation in NDNI may

be attributed to adaption to local nutrient availability of the mother tree.

**Growth and photosynthetic traits were correlated with survival in the middle garden**

Higher survival, growth rates and fitness in the Illinois garden for all populations were

likely a consequence of productive soil conditions and intermediate climate that alleviate stress

found in more extreme ends of the species' range. Previous work in another North American

temperate oak species (*Quercus rubra*) suggests that autumn frost damage at northern range

limits leads to reduced growth in southern seed sources (Lindback et al., 2023), which is likely

the cause of the negative growth observed in *Q. macrocarpa* from Oklahoma planted in the

Minnesota garden, and we did indeed observe top-down dieback of stems in the Minnesota

garden, particularly for the OK population. The lower growth rates of all populations observed in

the Minnesota and Oklahoma gardens compared to Illinois may be also explained by reduced

absolute rates of carbon fixation over the growing season. Photosynthetic rates are lower for the

MN and IL populations in the Oklahoma garden, likely in response to stressful climatic

conditions—such as increased aridity stress (Chaves, 1991).

Stressful conditions can also shift carbon allocation away from growth toward drought

stress responses, and defense (Wang et al., 2024). We found correlations between growth and

survival in Illinois garden, in which all populations expressed highest fitness, likely due to its

relatively mild climate and loamy soils. Previous work in the European pedunculate oak,

*Quercus robur,* indicates that differences in soil conditions and water availability across sites

with similar climate regimes lead to variation in productivity (Buras et al., 2020). Growth and

survival were not correlated in Oklahoma, where drought and high temperature stress were

highest, and the fine textured silty-clay soils in Oklahoma may be less conducive to nutrient

access. Growth was also not correlated with survival in Minnesota, which has the longest,

coldest winters of the three gardens, along with nutrient-poor sandy soils. This decoupling of

growth and survival in the hottest and coldest gardens may be a consequence of resource

allocation trade-offs under increased stress that improves survival at the expense of growth rates

(Bazzaz et al., 1987; Willi & Van Buskirk, 2022).

512 In addition, we found that $A_{max}$ was correlated with growth but not survival in all of the

513 gardens. Leaf chlorophyll content was also correlated with growth in the Illinois garden,

514 indicating that carbon assimilation and light energy acquisition through photosynthesis were

515 important for growth, particularly in Illinois.  In fertile sites, relative growth rates and

516 photosynthetic rates tended to be higher, as was the case in the Illinois garden, where conditions

517 were favorable (Lambers & Poorter, 1992). Survival and growth were higher in the less stressful

518 Illinois garden compared to the Oklahoma or Minnesota garden. Moreover, the two most

519 stressful gardens exhibited little difference from each other in growth or survival between the

520 two, further supporting the Illinois garden as a favorable habitat for bur oaks, where aridity and

521 cold stress are alleviated. In the Illinois garden, the oaks were able to allocate most of their

522 carbon toward aboveground growth, which in turn improved survival. On the other hand, in the

523 less favorable conditions of Minnesota and Oklahoma, survival likely depended on allocating

524 carbon elsewhere, such as root growth or defense against herbivory (Coley et al., 1985; Poorter

525 et al., 2012).

526 **Variation in season length helps explain carbon assimilation and growth patterns**

527 Differences in growing season lengths along the latitudinal gradient also played a role in

528 the populations' photosynthetic and growth responses. The Oklahoma garden has the longest

529 growing season which allowed the trees ample time to acquire carbon and grow. However

530 stresses during the growing season such as drought may reduce carbon acquisition by forcing

531 stomatal closure and decreasing carbon assimilation (Flexas et al., 2004; McDowell et al., 2008).

532 The Oklahoma population had higher fitness in the Oklahoma garden compared to other

533 populations and was likely adapted to the hot conditions during the prolonged growing season

534 there , sustaining carbon assimilation at relatively higher rates compared to the other populations.

On the other hand, the northern populations that originated from shorter growing seasons tended

to have higher $A_{max}$ rates than the Oklahoma population in the northern gardens, which is similar

to findings in a *Betula pendula* providence trial in which northern populations had greater net

photosynthesis (Tenkanen et al., 2021). The northern populations showed reduced growth and

photosynthesis in the high aridity and heat in the Oklahoma garden, which prevented them from

taking advantage of the longer growing season. In this study, we estimated growth in

aboveground biomass, however it is also plausible that the northernmost population from

Minnesota allocated more biomass to root growth than to aboveground biomass, leading to

smaller trees with lower aboveground growth, as has been shown in *B. pendula* (Tenkanen et al.,

2021). The Oklahoma population tended to maintain a relatively low rate of carbon assimilation

across gardens; compared to the other populations in the northern gardens, it showed the lowest

carbon assimilation rate, in accordance with leaf economic expectations where leaves with

longer life spans are theoretically expected and empirically demonstrated to have slower rates of

photosynthesis across species (Chabot & Hicks, 1982; Wright et al., 2004). $A_{max}$ and relative

growth rate peaked at intermediate growing season lengths found in the Illinois garden likely due

to trade-offs in optimizing carbon dynamics for leaf economics and stress tolerance in the other

gardens (Fig. S6).

**Multiple environmental factors vary across gardens**

An outdoor common garden experiment such as ours, with unmanipulated soils, cannot

isolate climate as the sole factor that varies among sites. Edaphic factors, as well as unmeasured

factors such as the microbial environment, also vary substantially. The Illinois garden had high

soil nitrogen  and organic matter content. Leaves from plants across populations in the Illinois

garden had higher spectrally detected nitrogen content (NDNI). Nitrogen is a major component

of photosynthetic chemistry, including RuBisCo, and chlorophyll, and is correlated with higher

photosynthetic rates (Ellis, 1979; Evans, 1989; Luo et al., 2021). Total soil nitrogen

concentration was similar between the Illinois and Oklahoma gardens, but the less arid and

slightly cooler climate in the Illinois garden combined with the high soil nitrogen and soil

organic matter of the loamy soil as an environment that is conducive to bur oak carbon

assimilation and fitness.  We also note that of the three gardens, Illinois had the least change in

growing season temperature since 1940. The fertile soil and relatively stable, mild climate of the

region near the Illinois garden may provide important habitat for bur oak populations to grow

with ongoing climate change. Controlled environment experiments that isolate climate as the

driving variable for performance changes in bur oaks are an important next step in predicting

response of bur oak populations to climate change.

**Conclusion**

Findings from the ACE reciprocal transplant experiment provide context for assisted

migration aimed at improving resilience of northern tree populations in the face of climate

change. Our results indicate that northern bur oak populations may suffer increasing mortality in

the face of climate change. We found evidence that the northern population was poorly adapted

to hotter conditions in the south, but populations originating from warmer southern climates

performed better than the northern population in the hottest garden. These findings indicate that

southern locations may provide valuable seed sources to augment exisiting genetic variation in

northern regions in the face of ongoing climatic warming. Consideration of current climate

conditions alongside future climate conditions are important in decisions to move locally adapted

populations. It may be beneficial to ensure seeds are sourced from locations with varied climates

to accommodate fluctuations in climate. We found that in the coldest, most northern climate, the

southern population performed the worst, suggesting that assisted migration of southern

populations northward must be timed with the pace of climate change. The lack of evidence for

local adaptation in northern populations indicates that the expected change in climate may be

within the tolerance limits of existing populations in the north. Mild climate warming akin to that

experienced by the MN populations in the Illinois garden may even improve performance and

fitness in the case of bur oak. In addition, other environmental factors that can impact growth and

survival must be accounted for in assisted migration planning. For example, limitations in soil

nutrient availability, and variation in competitive ability in different soil types may impact

growth and survival, even if seed sources and climates are well matched. We did not test

performance under competition with other species nor explicitly vary edaphic environments

across climates, however, the performance of oaks across contrasting edaphic environments in

competition with other tree species is an important area of investigation for future studies.

## Acknowledgements

We would like to thank those without whom the work presented here could not have been done.

We thank Kris Bachtell, Cathy Bechtoldt, Brett Fredericksen, Marlene Hahn, Troy Mielke, Maria

Park, Senna Robeson, Allison Scott, Nick Stoynoff, the Cedar Creek Ecosystem Science

Reserve, Morton Arboretum Facilities and Natural Resources staff, The Kessler Atmospheric and

Ecological Field Station staff for help establishing the common garden. We thank the Indiana

Department of Natural Resources Vallonia Nursery for use of their facilities and help with

harvesting seedlings. Field assistants including Iris Cessna, Isaiah Clark, Allison Collins, Nick

Duncan, Rachel Gowett, Jennifer Hamann, Leah Hill, Keion Mackey, and Morton Arboretum

volunteers were assisted with data collection. Ruth Shaw provided instruction on aster modelling

and provided edits on an earlier version of the manuscript. We acknowledge funding from

ASCEND Biological Integration Institute (NSF DBI 2021898), the National Science Foundation

Dimensions of Biodiversity grant (NSF DEB 2129312 (UMN), 2129281 (MOR), and 2129236

(UO)), the Morton Arboretum Center for Tree Science, and Cedar Creek Ecosystem Science

Reserve Long Term Ecological Research Program through NSF DEB 1831944.

**Author Contribution Statement**

All authors contributed intellectually to the study. JCB designed the ACE experiment and

managed its implementation. JCB, AH, HRM managed the collaboration and secured funding for

the study. LMSR lead data collection, analysis and interpretation, and writing of the manuscript.

JCB led nursery planting. CL, LW, and MG led planting efforts in the gardens. LO, RM, MG,

LW, and CL collected data. JCB, AH and HRM edited early versions of the manuscript. All

authors provided critical revision of the article and provided final approval of the version to be

published.

# References

Abrams, M. D., Nowacki, G. J., & Hanberry, B. B. (2021). Oak forests and woodlands as Indigenous

landscapes in the Eastern United States. *The Journal of the Torrey Botanical Society*, *149*(2).

https://doi.org/10.3159/TORREY-D-21-00024.1

Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T., & Curtis-McLane, S. (2008). Adaptation, migration

or extirpation: Climate change outcomes for tree populations. *Evolutionary Applications*, *1*(1),

95–111. https://doi.org/10.1111/j.1752-4571.2007.00013.x

Bates, D., Maechler, M., Bolker [aut, B., cre, Walker, S., Christensen, R. H. B., Singmann, H., Dai, B.,

Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., simulate.formula), P. N. K. (shared

copyright on, Tanaka, E., & Jagan, M. (2024). *lme4: Linear Mixed-Effects Models using "Eigen"*

*and S4* (Version 1.1-35.3) [Computer software]. https://cran.r-

project.org/web/packages/lme4/index.html

Bazzaz, F. A., Chiariello, N. R., Coley, P. D., & Pitelka, L. F. (1987). Allocating Resources to

Reproduction and Defense: New assessments of the costs and benefits of allocation patterns in

plants are relating ecological roles to resource use. *BioScience*, *37*(1), 58–67.

https://doi.org/10.2307/1310178

Bischoff, A., Vonlanthen, B., Steinger, T., & Müller-Schärer, H. (2006). Seed provenance matters—

Effects on germination of four plant species used for ecological restoration. *Basic and Applied*

*Ecology*, *7*(4), 347–359. https://doi.org/10.1016/j.baae.2005.07.009

Bosiö, J., Stiegler, C., Johansson, M., Mbufong, H. N., & Christensen, T. R. (2014). Increased

photosynthesis compensates for shorter growing season in subarctic tundra—8 years of snow

accumulation manipulations. *Climatic Change*, *127*(2), 321–334. https://doi.org/10.1007/s10584-

014-1247-4

Boyer, J. S. (1976). Water deficits and photosynthesis. In T. T. Kozlowski (Ed.), *Soil Water Measurement,*

*Plant Responses, and Breeding for Drought Resistance* (pp. 153–190). Academic Press.

Browne, L., Wright, J. W., Fitz-Gibbon, S., Gugger, P. F., & Sork, V. L. (2019). Adaptational lag to

temperature in valley oak (Quercus lobata) can be mitigated by genome-informed assisted gene

flow. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(50),

25179–25185. https://doi.org/10.1073/pnas.1908771116

Buras, A., Sass-Klaassen, U., Verbeek, I., & Copini, P. (2020). Provenance selection and site conditions

determine growth performance of pedunculate oak. *Dendrochronologia*, *61*, 125705.

https://doi.org/10.1016/j.dendro.2020.125705

Butler, P., Swanston, C., Janowiak, M., Parker, L., St. Pierre, M., & Brandt, L. (2012). Chapter 2:

Adaptation Strategies and Approaches. In *Forest adaptation resources: Climate change tools and*

*approaches for land managers*. en. Tech. Rep. NRS-87. Newtown Square, PA: U.S. Department

of Agriculture, Forest Service, Northern Research Station.

Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P. W., Trisos, C., Romero, J., Aldunce, P.,

Barrett, K., Blanco, G., Cheung, W. W. L., Connors, S., Denton, F., Diongue-Niang, A., Dodman,

D., Garschagen, M., Geden, O., Hayward, B., Jones, C., … Péan, C. (2023). *IPCC, 2023: Climate*

*Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth*

*Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H.*

*Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland.* (First). Intergovernmental Panel on

Climate Change (IPCC). https://doi.org/10.59327/IPCC/AR6-9789291691647

Chabot, B. F., & Hicks, D. J. (1982). The Ecology of Leaf Life Spans. *Annual Review of Ecology and*

*Systematics*, *13*(1), 229–259. https://doi.org/10.1146/annurev.es.13.110182.001305

Chaves, M. M. (1991). Effects of Water Deficits on Carbon Assimilation. *Journal of Experimental*

*Botany*, *42*(1), 1–16. https://doi.org/10.1093/jxb/42.1.1

Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of species

associated with high levels of climate warming. *Science*, *333*(6045), 1024–1026.

https://doi.org/10.1126/science.1206432

Coley, P. D., Bryant, J. P., & Chapin, F. S. (1985). Resource Availability and Plant Antiherbivore Defense.

*Science*, *230*(4728), 895–899. https://doi.org/10.1126/science.230.4728.895

Cornelius, J. (1994). The effectiveness of plus-tree selection for yield. *Forest Ecology and Management*,

*67*(1), 23–34. https://doi.org/10.1016/0378-1127(94)90004-3

Deacon, N. J., & Cavender-Bares, J. (2015). Limited pollen dispersal contributes to population genetic

structure but not local adaptation in Quercus oleoides forests of Costa Rica. *PLoS ONE*, *10*(9), 1–

25. https://doi.org/10.1371/journal.pone.0138783

Denny, E. G., & Crimmins, T. M. (2023). Updates to standardized plant and animal observation protocols

of the USA National Phenology Network. *International Journal of Biometeorology*, *67*(5), 927–

930. https://doi.org/10.1007/s00484-023-02444-0

Dobrowski, S. Z., Swanson, A. K., Abatzoglou, J. T., Holden, Z. A., Safford, H. D., Schwartz, M. K., &

Gavin, D. G. (2015). Forest structure and species traits mediate projected recruitment declines in

western US tree species. *Global Ecology and Biogeography*, *24*(8), 917–927.

https://doi.org/10.1111/geb.12302

Ellis, R. J. (1979). The most abundant protein in the world. *Trends in Biochemical Sciences*, *4*(11), 241–244. https://doi.org/10.1016/0968-0004(79)90212-3

Etterson, J. R. (2004a). Evolutionary potential of Chamaecrista fasciculata in relation to climate change. I. Clinal patterns of selection along an environmental gradient in the Great Plains. *Evolution*, *58*(7), 1446–1456. https://doi.org/10.1111/j.0014-3820.2004.tb01726.x

Etterson, J. R. (2004b). Evolutionary potential of Chamaecrista fasciculata in relation to climate change. II. Genetic architecture of three populations reciprocally planted along an environmental gradient in the Great Plains. *Evolution*, *58*(7), 1459–1471. https://doi.org/10.1111/j.0014-3820.2004.tb01727.x

Etterson, J. R., Cornett, M. W., White, M. A., & Kavajecz, L. C. (2020). Assisted migration across fixed seed zones detects adaptation lags in two major North American tree species. *Ecological Applications*, *30*(5), 1–20. https://doi.org/10.1002/eap.2092

Etterson, J. R., & Shaw, R. G. (2001). Constraint to adaptive evolution in response to global warming. *Science*, *294*(5540), 151–154. https://doi.org/10.1126/science.1063656

Evans, J. R. (1989). Photosynthesis and nitrogen relationships in leaves of Ca plants. *Oecologia*, *78*, 9–19. https://doi.org/10.1109/LSP.2017.2723724

Flexas, J., Bota, J., Loreto, F., Cornic, G., & Sharkey, T. D. (2004). Diffusive and Metabolic Limitations to Photosynthesis under Drought and Salinity in C3 Plants. *Plant Biology*, *6*(3), 269–279. https://doi.org/10.1055/s-2004-820867

Franks, S. J., Weber, J. J., & Aitken, S. N. (2014). Evolutionary and plastic responses to climate change in terrestrial plant populations. *Evolutionary Applications*, *7*(1), 123–139. https://doi.org/10.1111/eva.12112

Galatowitsch, S., Frelich, L., & Phillips-Mao, L. (2009). Regional climate change adaptation strategies for biodiversity conservation in a midcontinental region of North America. *Biological Conservation*, *142*(10), 2012–2022. https://doi.org/10.1016/j.biocon.2009.03.030

Gamache, I., & Payette, S. (2004). Height Growth Response of Tree Line Black Spruce to Recent Climate Warming across the Forest-Tundra of Eastern Canada. *Journal of Ecology*, *92*(5), 835–845.

Geyer, C. J., Wagenius, S., & Shaw, R. G. (2007). Aster models for life history analysis. *Biometrika*, *94*(2), 415–426. https://doi.org/10.1093/biomet/asm030

Gimeno, T. E., Pias, B., Lemos-Filho, J. P., & Valladares, F. (2008). Plasticity and stress tolerance override local adaptation in the responses of Mediterranean holm oak seedlings to drought and cold. *Tree Physiology*, *29*(1), 87–98. https://doi.org/10.1093/treephys/tpn007

Gitelson, A. A., & Merzlyak, M. N. (1994). *Quantitative estimation of chlorophyll-u using reflectance spectra: Experiments with autumn chestnut and maple leaves*.

Gorton, A. J., Benning, J. W., Tiffin, P., & Moeller, D. A. (2022). The spatial scale of adaptation in a native annual plant and its implications for responses to climate change. *Evolution*, evo.14583. https://doi.org/10.1111/evo.14583

Handler, S., Duveneck, M. J., Iverson, L., Peters, E., Scheller, R. M., Wythers, K. R., Brandt, L., Butler, P., Janowiak, M., Shannon, P. D., Swanston, C., Barrett, K., Kolka, R., McQuiston, C., Palik, B., Reich, P. B., Turner, C., White, M., Adams, C., … Ziel, R. (2014). Minnesota forest ecosystem vulnerability assessment and synthesis: A report from the Northwoods Climate Change Response Framework project. *Gen. Tech. Rep. NRS-133. Newtown Square, PA; U.S. Department of Agriculture, Forest Service, Northern Research Station. 228 p.*, *133*, 1–228. https://doi.org/10.2737/NRS-GTR-133

Herms, D. A., & Mattson, W. J. (1992). The Dilemma of Plants: To Grow or Defend. *The Quarterly Review of Biology*, *67*(3), 283–335.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., & Thépaut, J.-N. (2023). *ERA5 monthly averaged data on single levels from 1940 to present.* [Dataset]. Copernicus Climate Change Service (C3S). https://doi.org/DOI: 10.24381/cds.f17050d7

Hoekstra, J. M., Boucher, T. M., Ricketts, T. H., & Roberts, C. (2005). Confronting a biome crisis: Global

      disparities of habitat loss and protection. *Ecology Letters*, *8*(1), 23–29.

      https://doi.org/10.1111/j.1461-0248.2004.00686.x

Hoffmann, A. A., & Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*, *470*(7335),

      479–485. https://doi.org/10.1038/nature09670

Huntley, B. (1989). *European post-glacial vegetation history: A new perspective*. https://durham-

      repository.worktribe.com/output/1693365/european-post-glacial-vegetation-history-a-new-

      perspective

Iverson, L. R., Peters, M. P., Prasad, A. M., & Matthews, S. N. (2019). Analysis of Climate Change

      Impacts on Tree Species of the Eastern US: Results of DISTRIB-II Modeling. *Forests*, *10*(4),

      302. https://doi.org/10.3390/f10040302

Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*(12), 1225–

      1241. https://doi.org/10.1111/j.1461-0248.2004.00684.x

Kikuzawa, K. (1991). A Cost-Benefit Analysis of Leaf Habit and Leaf Longevity of Trees and Their

      Geographical Pattern. *The American Naturalist*, *138*(5), 1250–1263.

Klein, T., Vitasse, Y., & Hoch, G. (2016). Coordination between growth, phenology and carbon storage in

      three coexisting deciduous tree species in a temperate forest. *Tree Physiology*, *36*(7), 847–855.

      https://doi.org/10.1093/treephys/tpw030

Körner, C. (2003). Carbon limitation in trees. *Journal of Ecology*, *91*(1), 4–17.

      https://doi.org/10.1046/j.1365-2745.2003.00742.x

Körner, C. (2015). Paradigm shift in plant growth control. *Current Opinion in Plant Biology*, *25*, 107–

      114. https://doi.org/10.1016/j.pbi.2015.05.003

Krech, S., Kay, C., & Simmons, R. (2004). Wilderness and Political Ecology: Aboriginal Influences and

      the Original State of Nature. *The Western Historical Quarterly*, *35*, 81.

      https://doi.org/10.2307/25442936

Lambers, H., & Poorter, H. (1992). *Inherent Variation in Growth Rate Between Higher Plants: A Search for Physiological Causes and Ecological Consequences. 23*, 187–261. https://doi.org/10.1016/S0065-2504(08)60148-8

Lindback, E. C., Rauschendorfer, J. K., Burton, A. J., Külheim, C., & Cavaleri, M. A. (2023). Common garden study reveals frost-tolerant northern seed sources are best suited to expand range of *Quercus rubra. Forest Ecology and Management*, *539*, 120985. https://doi.org/10.1016/j.foreco.2023.120985

Luo, X., Keenan, T. F., Chen, J. M., Croft, H., Colin Prentice, I., Smith, N. G., Walker, A. P., Wang, H., Wang, R., Xu, C., & Zhang, Y. (2021). Global variation in the fraction of leaf nitrogen allocated to photosynthesis. *Nature Communications*, *12*(1), 1–10. https://doi.org/10.1038/s41467-021-25163-9

MacArthur, R. H. (1984). *Geographical Ecology: Patterns in the Distribution of Species*. Princeton University Press.

McDowell, N., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D. G., & Yepez, E. A. (2008). Mechanisms of plant survival and mortality during drought: Why do some plants survive while others succumb to drought? *New Phytologist*, *178*(4), 719–739. https://doi.org/10.1111/j.1469-8137.2008.02436.x

McLachlan, J. S., Clark, J. S., & Manos, P. S. (2005). Molecular indicators of tree migration capacity under rapid climate change. *Ecology*, *86*(8), 2088–2098. https://doi.org/10.1890/04-1036

Moore, C. E., Meacham-Hensold, K., Lemonnier, P., Slattery, R. A., Benjamin, C., Bernacchi, C. J., Lawson, T., & Cavanagh, A. P. (2021). The effect of increasing temperature on crop photosynthesis: From enzymes to ecosystems. *Journal of Experimental Botany*, *72*(8), 2822–2844. https://doi.org/10.1093/jxb/erab090

Nagamitsu, T., & Shuri, K. (2021). Seed transfer across geographic regions in different climates leads to reduced tree growth and genetic admixture in Quercus mongolica var. Crispula. *Forest Ecology and Management*, *482*, 118787. https://doi.org/10.1016/j.foreco.2020.118787

Oram, N. J., Ingrisch, J., Bardgett, R. D., Brennan, F., Dittmann, G., Gleixner, G., Illmer, P., Praeg, N., &

Bahn, M. (2023). Drought intensity alters productivity, carbon allocation and plant nitrogen

uptake in fast versus slow grassland communities. *Journal of Ecology*, *111*(8), 1681–1699.

https://doi.org/10.1111/1365-2745.14136

Papper, P. D., & Ackerly, D. D. (2021). Partitioning genetic and environmental components of

phenological variation in Quercus douglasii (Fagaceae). *Madroño*, *68*(4).

https://doi.org/10.3120/0024-9637-68.4.425

Poorter, H., Niklas, K. J., Reich, P. B., Oleksyn, J., Poot, P., & Mommer, L. (2012). Biomass allocation to

leaves, stems and roots: Meta-analyses of interspecific variation and environmental control. *New*

*Phytologist*, *193*(1), 30–50. https://doi.org/10.1111/j.1469-8137.2011.03952.x

Pretzsch, H., Biber, P., Schütze, G., Uhl, E., & Rötzer, T. (2014). Forest stand growth dynamics in Central

Europe have accelerated since 1870. *Nature Communications*, *5*(1), 4967.

https://doi.org/10.1038/ncomms5967

QGIS.org. (2024). *QGIS Geographic Information System* [Computer software]. QGIS Association.

R Core Team. (2023). *R: A language and environment for statistical computing* [Computer software]. R

Foundation for Statistical Computing.

Ramírez-Valiente, J. A., & Cavender-Bares, J. (2017). Evolutionary trade-offs between drought resistance

mechanisms across a precipitation gradient in a seasonally dry tropical oak (Quercus oleoides).

*Tree Physiology*, *37*(7), 889–901. https://doi.org/10.1093/treephys/tpx040

Ramírez-Valiente, J. A., Deacon, N. J., Etterson, J., Center, A., Sparks, J. P., Sparks, K. L., Longwell, T.,

Pilz, G., & Cavender-Bares, J. (2018). Natural selection and neutral evolutionary processes

contribute to genetic divergence in leaf traits across a precipitation gradient in the tropical oak

Quercus oleoides. *Molecular Ecology*, *27*(9), 2176–2192. https://doi.org/10.1111/mec.14566

Reich, P. B., & Oleksyn, J. (2008). Climate warming will reduce growth and survival of Scots pine except

in the far north. *Ecology Letters*, *11*(6), 588–597. https://doi.org/10.1111/j.1461-

0248.2008.01172.x

Sáenz-Romero, C., Lamy, J.-B., Ducousso, A., Musch, B., Ehrenmann, F., Delzon, S., Cavers, S.,

Chałupka, W., Dağdaş, S., Hansen, J. K., Lee, S. J., Liesebach, M., Rau, H.-M., Psomas, A.,

Schneck, V., Steiner, W., Zimmermann, N. E., & Kremer, A. (2017). Adaptive and plastic

responses of Quercus petraea populations to climate across Europe. *Global Change Biology*,

*23*(7), 2831–2847. https://doi.org/10.1111/gcb.13576

Schwinning, S., Lortie, C. J., Esque, T. C., & DeFalco, L. A. (2022). What common-garden experiments

tell us about climate responses in plants. *Journal of Ecology*, *110*(5), 986–996.

https://doi.org/10.1111/1365-2745.13887

Seneviratne, S. I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Di Luca, A., Ghosh, S., Iskandar, I.,

Kossin, J., Lewis, S., Otto, F., Pinto, I., Satoh, M., Vicente-Serrano, S. M., Wehner, M., & Zhou,

B. (2023). 2021: Weather and Climate Extreme Events in a Changing Climate. In *Climate Change*

*2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report*

*of the Intergovernmental Panel on Climate Change* (1st ed.). Cambridge University Press.

https://doi.org/10.1017/9781009157896

Serrano, L., Penuelas, J., & Ustin, S. L. (2002). Remote sensing of nitrogen and lignin in Mediterranean

vegetation from AVIRIS data: Decomposing biochemical from structural signals. *Remote Sensing*

*of Environment*.

Shaw, R. G., Geyer, C. J., Wagenius, S., Hangelbroek, H. H., & Etterson, J. R. (2008). Unifying life-

history analyses for inference of fitness and population growth. *American Naturalist*, *172*(1).

https://doi.org/10.1086/588063

Sittaro, F., Paquette, A., Messier, C., & Nock, C. A. (2017). Tree range expansion in eastern North

America fails to keep pace with climate warming at northern range limits. *Global Change*

*Biology*, *23*(8), 3292–3301. https://doi.org/10.1111/gcb.13622

Tallamy, D. W. (2021). *The nature of oaks: The rich ecology of our most essential native trees*. Timber

Press.

Tang, Y., Du, E., Guo, H., Wang, Y., Peñuelas, J., & Reich, P. B. (2024). Rapid migration of Mongolian oak into the southern Asian boreal forest. *Global Change Biology*, *30*(1), e17002. https://doi.org/10.1111/gcb.17002

Tenkanen, A., Suprun, S., Oksanen, E., Keinänen, M., Keski-Saari, S., & Kontunen-Soppela, S. (2021). Strategy by latitude? Higher photosynthetic capacity and root mass fraction in northern than southern silver birch (Betula pendula Roth) in uniform growing conditions. *Tree Physiology*, *41*(6), 974–991. https://doi.org/10.1093/treephys/tpaa148

Thien, S. J. (1979). A flow diagram for teaching texture-by-feel analysis. *Journal of Agronomic Education*, *8*(1), 54–55. https://doi.org/10.2134/jae.1979.0054

Thornthwaite, C. W. (1948). An Approach toward a Rational Classification of Climate. *Geographical Review*, *38*(1), 55–94. https://doi.org/10.2307/210739

Tierney, J. E., Poulsen, C. J., Montañez, I. P., Bhattacharya, T., Feng, R., Ford, H. L., Hönisch, B., Inglis, G. N., Petersen, S. V., Sagoo, N., Tabor, C. R., Thirumalai, K., Zhu, J., Burls, N. J., Foster, G. L., Goddéris, Y., Huber, B. T., Ivany, L. C., Kirtland Turner, S., … Zhang, Y. G. (2020). Past climates inform our future. *Science*, *370*(6517), eaay3701. https://doi.org/10.1126/science.aay3701

Torres-Ruiz, J. M., Kremer, A., Carins Murphy, M. R., Brodribb, T., Lamarque, L. J., Truffaut, L., Bonne, F., Ducousso, A., & Delzon, S. (2019). Genetic differentiation in functional traits among European sessile oak populations. *Tree Physiology*, *39*(10), 1736–1749. https://doi.org/10.1093/treephys/tpz090

Wadgymar, S. M., DeMarche, M. L., Josephs, E. B., Sheth, S. N., & Anderson, J. T. (2022). Local adaptation: Causal agents of selection and adaptive trait divergence. *Annual Review of Ecology, Evolution, and Systematics*, *53*(1), 87–111. https://doi.org/10.1146/annurev-ecolsys-012722-035231

Wang, R., Yun, L., Mao, Y., Yan, T., Wei, W., You, W., & Zhang, H. (2024). Nitrogen deposition alters drought-induced changes in biomass and nonstructural carbohydrates allocation patterns of

*Quercus mongolica* seedlings. *Scientia Horticulturae*, *325*, 112573.

https://doi.org/10.1016/j.scienta.2023.112573

Willi, Y., & Van Buskirk, J. (2022). A review on trade-offs at the warm and cold ends of geographical

distributions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *377*(1848),

20210022. https://doi.org/10.1098/rstb.2021.0022

Williams, G. C. (George C. (with Internet Archive). (1966). *Adaptation and natural selection; a critique*

*of some current evolutionary thought*. Princeton, N.J., Princeton University Press.

http://archive.org/details/adaptationnatura00will

Williams, M. I., & Dumroese, R. K. (2013). Preparing for climate change: Forestry and assisted

migration. *Journal of Forestry*, *111*(4), 287–297. https://doi.org/10.5849/jof.13-016

Williams, N. (2021). *Climate Change Effects on Tree Regeneration*. CLIMATE SMART LAND

NETWORK BULLETIN. https://www.manomet.org/publication/climate-change-effects-on-tree-

regeneration/

Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., Cavender-Bares, J.,

Chapin, T., Cornelissen, J. H. C., Diemer, M., Flexas, J., Garnier, E., Groom, P. K., Gulias, J.,

Hikosaka, K., Lamont, B. B., Lee, T., Lee, W., Lusk, C., … Villar, R. (2004). The worldwide leaf

economics spectrum. *Nature*, *428*(6985), 821–827. https://doi.org/10.1038/nature02403

Wright, J. W., Ivey, C. T., Canning, C., & Sork, V. L. (2021). Timing of bud burst is associated with

climate of maternal origin in quercus lobata progeny in a common garden. *Madroño*, *68*(4).

https://doi.org/10.3120/0024-9637-68.4.443

Zhu, K., Woodall, C. W., & Clark, J. S. (2012). Failure to migrate: Lack of tree range expansion in

response to climate change. *Global Change Biology*, *18*(3), 1042–1052.

https://doi.org/10.1111/j.1365-2486.2011.02571.x

# Tables

883 *Table 2.1.* Soil properties in each of the three gardens.

| Garden | % Nitrogen | % Carbon | C:N | Soil texture |
|--------|-----------|----------|-------|-----------------|
| MN | 0.06 | 1.04 | 21.32 | Sandy loam |
| IL | 0.25 | 3.54 | 14.12 | Sandy clay loam |
| OK | 0.30 | 3.33 | 11.21 | Silty clay loam |

# Figure captions

*Figure 1.* Reciprocal transplant garden design and climate information. In the top panel (a), the
experimental design of the Adaptation to Climate and Environment (ACE) experiment is
depicted. Three common gardens are located across a latitudinal gradient in Minnesota (Cedar
Creek Ecosystem Science Reserve), Illinois (Morton Arboretum), and Oklahoma (Kessler Field
Station). In each garden, 600 seedlings were planted representing the three populations from
Minnesota (blue), Illinois (green) and Oklahoma (orange). Each population was comprised of 20
half-sibling maternal families, with half-sibling acorn progeny collected from one known mother
tree. Ten progeny from each family were planted in each garden. Photos in each site were taken
in summer of 2022 after one year of planting. In the bottom panel, ERA5 monthly climate data
averaged over the years 2021-2023 is shown for temperature (b), total precipitation (c), and the
index of moisture (d). Colored lines indicate gardens: Minnesota (blue), Illinois (green) and
Oklahoma (orange).

*Figure 2.* Fitness, fitness components, and physiology in each garden. Each panel represents the
common garden located in Oklahoma (OK), Illinois (IL), or Minnesota (MN). Progeny from
each source population are represented by orange circles for OK, green triangles for IL, and blue
squares for MN, with larger points indicating population means. Distinct letters denote statistical
significance ($p<0.05$) for each population group. Percent survival is represented by a point for
each maternal family (a), and was highest for the local OK population in OK, but was similar
among all populations in the northern gardens. The relative growth rate of stem volume for
individuals (b) was also highest for the local population in OK, with the lowest growth rates
occurring in MN. Aster expected fitness (c) is represented by a point for each population, and
was highest in the Illinois garden. Aster fitness significance was determined by omnibus tests of
population*garden interactions. Maximum CO2 assimilation ($A_{max}$) (d) of the local populations
was highest for the OK and MN garden, but overall the IL garden had the highest $A_{max}$. Spectral
chlorophyll normalized difference index (ChlNDI) (e) was also higher overall in the IL garden
for all populations, though the local MN population was highest in the MN garden. Normalized
difference of the nitrogen index (NDNI) (f) indicates that leaf nitrogen was highest in the Illinois
garden.

*Figure 3.* Comparisons of growth, survival, and physiology to understand shifts in carbon
allocation. Each panel represents a common garden located in Oklahoma (OK), Illinois (IL), or
Minnesota (MN). The populations are represented by an orange circle for OK, a green triangle
for IL, and a blue square for MN, with significant trendlines (with standard errors) in matching
colors. a) Percent survival by maternal family is significantly positively correlated with relative
stem volume growth for all populations in the Illinois garden, but not in the OK or MN gardens.

$A_{max}$ has significant relationships with growth (b) for OK and IL populations in all gardens, and
for the MN population in the IL garden. Chlorophyll normalized difference index (ChlNDI) was
significantly positively correlated with relative growth rate (c) in IL for all populations, and was
significantly correlated with growth in the MN garden for the IL and MN populations.

*Figure 4.* Phenology and growing season length in each of the gardens. Each panel represents a
common garden located in Oklahoma (OK), Illinois (IL), or Minnesota (MN). The populations
are represented by an orange circle for OK, a green triangle for IL, and a blue square for MN.
Budburst (a) occurs earlier in the OK and IL garden for all populations than in the MN garden,
and senescence (b) occurs latest in the OK garden. The growing season length (c) is shortest in
the MN garden and longest in the OK garden.

*Figure 1.*

*Figure 2.*

*Figure 3.*

*Figure 4.*