1

Full title: Quantifying structural variants in chromosomes using landmark-based

disparity

Short title: Measuring structural variants using genomic landmarks

Authors and author information:

Ashwini V. Mohan[1,2]*, Anjali Goswami[2], and Jeffrey W. Streicher[2]*

1: Institut de Biologie, Université de Neuchâtel, Neuchâtel, Switzerland

2: Natural History Museum, London SW7 5BD, United Kingdom

* Authors for correspondence:

Ashwini V. Mohan (ORCID ID: 0000-0001-9505-0665)

Email: ashwini.venkatanarayana@unine.ch

Jeffrey W. Streicher (ORCID ID: 0000-0002-3738-4162)

Email: j.streicher@nhm.ac.uk

Abstract

Chromosomal architecture has played a key role in the evolution of biodiversity. Detecting structural variants (SVs) on chromosomes has informed the study of speciation, sex determination, adaptation, and some of the earliest divergences in the tree of life. Here we present a computationally non-intensive approach, based on geometric morphometrics, that uses conserved DNA sequences as landmarks to quantify structural disparities of focal chromosomes across multiple species, individuals, or cell types. Based on two approaches, we show that this 'geno-metric' method can be applied at micro- and macroevolutionary scales to discover and diagnose SVs. Using human X-linked genes and ultraconserved elements as landmarks, we provide empirical demonstrations with amniote sex chromosomes, the *Drosophila virilis* group, and placental mammal genomes. Landmark-based structural disparity analysis effectively identifies chromosomal rearrangements and has parallels with traditional morphometrics regarding chromosome size, landmark orientation and landmark availability. Using simulations, we show that structural disparity inferred from ultraconserved elements is correlated with overall levels of chromosome evolution; an attribute which is consistent with observed disparity between and within mammalian orders. We found that the disparity patterns of SVs have significant phylogenetic signal, giving them broad importance for studying evolutionary biology. Structural disparity analyses are a valuable addition to the comparative genomic toolkit in that they offer an intuitive, rapid mechanism for detecting SVs associated with single copy genetic landmarks and the potential to reveal broader patterns of chromosome evolution related to expansions, contractions, rearrangements and phylogeny.

Introduction

We are now firmly in the age of genomics with fully sequenced genomes commonplace for most organismal groups. Because of this, multi-species genomic comparisons are increasingly used in evolutionary and ecological research. This is even the case for organisms with relatively large nuclear genomes [1–3]. Excitingly, the comparison of whole genome sequences has offered valuable evolutionary insights into a range of topics including gene orthology [4], transposable element evolution [5], and adaptation to novel environments [6].

One important area of comparative genomics research is the study of chromosome architecture [7]. Detecting structural variants (SVs) of chromosomes is highly relevant to the evolution of biodiversity and disease [8]. From supergenes [9] to sex chromosomes [10], evolutionary shifts in chromosome structure are known to have a substantial influence on organismal phenotype. Recent research utilizing forward in time simulations has also indicated the important role of SVs in sustaining genomic adaptation [11]. Traditional methods for detecting SVs include chromosome banding and fluorescent in situ hybridization [12]. Several modern methods using genomic sequencing have also been developed including aberrant read alignment patterns, split reads, coverage changes, alignment of long-read sequences, and strand-seq [13–14]. Amongst sequencing-based methods, synteny assessment [15] is increasingly used to compare chromosome structure across multiple species with visual interpretation achieved through complex network diagrams, e.g. [16–19]. Most of these high-throughput DNA sequencing approaches for detecting SVs depend on comparisons to a single reference sequence and/or whole genome alignment, which can be subject to

reference-bias and time consuming, respectively [20]. To complement existing

methods, here we present a method for detecting SVs across chromosomes from

multiple species or individuals simultaneously that is reference-free, computationally

non-intensive, and does not require whole genome alignment. Based on the principles

of morphometrics, our method is applied to chromosomes from high-quality genome

assemblies and facilitates the study of SVs in ordinated micro- and macroevolutionary

contexts.

Within the study of comparative morphology, structural variation in homologous

structures across species is commonly termed 'disparity' [21–22]. Measures of

morphological disparity are routinely used to explore the evolutionary dynamics of

complex phenotypes, e.g. [23–24]. Structural disparity analysis of chromosomes is a

disparity-based method that uses conserved landmark sequences to capture genomic

variation across different species in a single analytical framework. Our 'geno-metric'

method is based on a simple premise: chromosome structure is morphology (i.e., a

shape, form, or structure) and there are well-validated methods for capturing

morphological disparity across multiple species [23–25]. Furthermore, genomes are

filled with homologous sequences (=landmarks) making them amenable for such

analyses (e.g. [26]). To develop and scrutinize landmark-based structural disparity

analyses, we analyzed three empirical datasets. First, we quantified SVs associated

with sex chromosome evolution across amniotes using human X-linked BUSCO

(Benchmarking Universal Single-Copy Orthologs) genes as landmarks which represents

a scenario when the structure of a particular genomic/chromosomal feature is of

interest [sensu 18]. Second, we compared SVs related to a speciation-associated

chromosome in Drosophila [27] using ultraconserved elements (UCEs; [28]) as

landmarks which represents a scenario when chromosome-wide structure may be of

interest. Third, we used genome sequences from placental mammals to develop a

pipeline for discovering chromosomes that share homologous landmarks; a necessary

step when chromosome homology is unclear or partial. We also used the placental

mammal dataset to simulate SVs of chromosomes and determine the effects of indels

and inversions on disparity measures. Finally, we tested all datasets for phylogenetic

signals associated with disparity patterns of SVs.

Results

Pipeline development. We developed two approaches for conducting structural

disparity analysis, [1] using generic feature format (gff3) annotation files as input and [2]

a de novo method that allows users to landmark and analyze whole chromosome

sequences from genome assemblies (Fig. 1). The gff3-formatted pipeline allows users

to extract genes from the chromosome of a single species in the gff3 file and then

identify orthologs in the other species contained within the file. This approach is useful

when chromosome homology is known, and focal landmarks are largely confined to

single chromosomes. The de novo method begins by acquiring chromosome sequences

from NCBI and then reference mapping landmarks to the assemblies using bwa [29].

We tested other approaches for reference mapping including minimap2 using default

settings [30] but found bwa resulted in the most mapped landmarks per chromosome

(Fig. S1). After orthologous landmarks are mapped, landmarked chromosomes are

either prepared for downstream analysis (when chromosomal homology is known), or

clustered based on the presence/absence of landmarks via metric multi-dimensional

scaling (when chromosomal homology is not known). The latter option allows for the

identification of chromosomes that share large numbers of landmarks, and thus likely

large homologous genomic blocks. Following landmarking, both approaches have

several steps in common including; removal of landmarks that are multi-copy or not

shared across homologs, complementarity checks (5'-3' versus 3'-5'), and exportation

of landmark position information into a spreadsheet. The spreadsheet has rows

representing species or individuals (one for each species or individual included in the

input assemblies) and columns are landmark positions on a homologous chromosome

set. The spreadsheet is then used as input for the disparity inference stage. Depending

on user preference, absolute landmark positions can be used, or positions can be

transformed using landmark bounding and/or chromosome size correction. Unless

otherwise indicated, all steps in the pipeline occur using base functions in the R

statistical package [31]. As with other disparity analyses, we use ordination techniques

to reduce the dimensionality of variation in landmark position wherein the spreadsheet

of landmark positions is analyzed using principal component analysis (Fig. 1).

Figure 1. Schematic illustration of the landmarking pipeline for structural disparity analysis of

chromosomes. When using generic feature format (gff3) annotation files, landmark genes are

selected from a reference followed by identification of orthologs from other species in the file and

extraction of chromosome information and landmark positions. In the de novo approach, the first step

is reference mapping landmarks to chromosome-level genome assemblies of different species,

individuals, or cell types. If the homology of chromosomes is known, then the chromosomes are

further prepared for disparity analysis. If homology is unknown, a presence-absence matrix is used to

identify chromosomes with similar landmark compositions using a gap-based threshold for similarity.

Chromosome sets are prepared for disparity analysis by extracting chromosome position information

for each landmark. The resulting data matrix is used for multivariate Principal Component Analysis to

visualize disparity across species.

Structural disparity of human X-linked genes. Lovell et al. [18] examined 300 million

years of vertebrate sex chromosome evolution using synteny-anchoring. We took their

gff3-formatted annotation file and identified 53 human X-linked genes that could be

used as landmarks on a set of single chromosomes for the focal taxa. We found that

principal components (PCs) 1 and 2 explained 69.3% and 14.2% of structural variance,

respectively. Using these first two axes, we found that major phylogenetic groups

occupied distinct areas of geno-metric space (Fig. S2A). PC1 was significantly

correlated with chromosome size (non-parametric Spearman's test, rho = 0.70, p =

0.003), but not PC2 (rho = -0.27, p = 0.3043). Mean structural disparity was highest

amongst those groups where the landmarks were found on sex chromosomes

(placental mammals, marsupial mammals and squamate reptiles) compared to those

where these markers are located on autosomal chromosomes (birds, monotreme

mammals). When the data were analyzed using the reverse complement orientation of

the chromosomes, we found that some chromosomes with clustered landmarks

resulted in exaggerated PC scores for some taxa (e.g., chicken, *Gallus gallus*, Fig. S2B).

We found that landmark bounding corrected for this methodological artifact regardless

of the 5' to 3' orientation of the chromosomes (Fig. 2A, S2C, S2D). A size-corrected

version of the analysis changed the patterns of disparity with birds being the most

disparate group, followed closely by placental mammals (Fig. S2E). The size-corrected

analysis was more robust to chromosome orientation with comparable results (Fig.

S2F). Size-correction of the bounded version of the dataset generated indistinguishable

results in the two directions of analysis (Fig. S2G, S2H). For comparison we generated a

GENESPACE-style synteny link plot (Fig. 2B).

Figure 2. Structural disparity analysis of 53 human X-linked BUSCO gene landmarks in select

amniote vertebrates studied by Lovell et al. (2022). Disparity analysis using bounded, absolute

landmark positions reveals larger disparity in those groups where landmarks are found on the X

chromosomes (placental mammals and marsupial mammals) than in those groups where landmarks

are found on autosomes or Z chromosomes (A). This disparity is largely explained by chromosome

size in that some of these groups maintain high synteny despite also having high structural disparity

(B). Size-corrected disparity analyses capture different aspects of variation in landmark placement

revealing birds as especially disparate after chromosome size is accounted for (Fig. S2E, F, G, H).

Structural disparity among closely related species of *Drosophila*. Poikela et al. [27]

identified multiple chromosomal inversions that are associated with speciation

patterns in the *Drosophila virilis* complex. We selected chromosome 5 for structural

disparity analysis because there are known chromosomal inversions across the focal

species [27]. We identified a set of 191 widely distributed dipteran UCEs [32] on

chromosomes 5 for use as landmarks. Our structural disparity analysis revealed that

three PC axes explained nearly 100% of the variance in landmark placement (PC1, 94%;

PC2 5.2%; PC3, 0.07%). Patterns of disparity were highly polarized with PC1 separating

D. flavomontana from the other species, PC2 separating *D. montana* from the other

species, and PC3 scores separating *D. americana + D. novamexicana* from the two

individuals of *D. virilis* (Fig. 3). Those individuals with highly conserved, syntenic UCE

landmark placements occurred closest in geno-metric space. Notably, this was the

case for the two individuals of *D. virilis* (Fig. 3A) and close relatives *D. americana* and *D.*

novamexicana (Fig. 3B; [33]). In contrast, separation corresponded to distinct SVs. PC1

scores were largely explained by two blocks of UCE landmarks with substantially

different placements in *D. flavomontana* compared to the other taxa (Fig. 3C). PC2

scores largely explained by a pericentric chromosomal inversion in *D. montana* relative

to the other taxa (Fig. 3D). Importantly, when we compare the landmark orientations on

D. montana and *D. flavomontana*, we see evidence (Fig. 3E) of the two blocks of UCE

landmarks (PC1) and the pericentric inversion (PC2). PC3 scores were largely explained

by a paracentric inversion that *D. virilis* has relative to both D. americana (Fig. 3F) and *D.*

novamexicana (Fig. 3G). None of the three most explanatory axes were significantly

correlated with chromosome size (PC1, rho = 0.029, p = 1.000; PC2, rho = -0.371, p =

0.4972; PC3, rho = 0.600, p = 0.2417).

Figure 3. Structural disparity analysis using raw, absolute landmark positions of 191 ultraconserved

elements (UCEs) on chromosome 5 of the *Drosophila virilis* group, different colors correspond to

different species in the complex. Synteny link maps around the main plot demonstrate what

separation in geno-metric space corresponds to, in terms of landmark placement on chromosome 5.

When samples are close to one another in space they are highly syntenic (A, B). Different axes

correspond to discrete differences in landmark placement including two translocations (C), a

pericentric inversion (D, E) and paracentric inversion (E, F). An 'M' was arbitrarily added by plotly (91)

to values on component axes because they are less than one. Synteny link maps were generated

using chromoMap [35].

De novo homology discovery and structural disparity analysis using whole genome

assemblies. Over long periods of evolutionary time, genome structure may change,

substantially eroding the homology of chromosomes. Despite this, large genomic

regions often remain evolutionarily intact on single chromosomes (e.g., [16]) among

which structural disparity can be explored. We used the chromosome clustering

module of our de novo pipeline to identify several chromosome sets containing

homologous tetrapod UCEs [34] from 26 genome assemblies of five placental mammal

orders (Artiodactyla, Carnivora, Perissodactyla, Primates, and Rodentia; Table 1). The

results of the 10-axis chromosome clustering revealed that score thresholding could be

used to identify eight sets of mutually exclusive chromosomes that (a) were highly

similar in their UCE compositions and (b) contained only a single chromosome per

species in the dataset (Fig. 4A, Table 2). Chromosome similarity and threshold gaps

decayed to after the second axis with not all sets containing 26 species and some

instances of overlap with chromosomes belonging to other sets (e.g. *Giraffa*

tippelskirchi chromosomes in sets 7 and 8, Fig 4A, Table 2). Axes 9 and 10 did not reveal

any novel chromosome clusters associated with UCE landmark content (Table 2).

Viewing axes as bivariate plots made some clustering patterns clearer, especially in axis

4 (Fig. 4B, C). Given (a) the clear separation of chromosomes sets 1 and 2, and (b) that

protein content was significantly more similar within these chromosome sets than

between them (W = 6742, p < 0.0001, nonparametric Wilcoxon tests, Fig. S3) we used

these chromosomes as input for a structural disparity analyses based on SVs of their

UCE landmarks.

Figure 4. Metric multi-dimensional scaling (MMDS) of 567 landmarked placental mammal

chromosomes based on patterns of ultraconserved element presence and absence. These

chromosomes represent the genomes of 26 species (Table 1). MMDS axis thresholds are designated

to establish 'chromosome sets' which contain a chromosome for most species included in the dataset.

These chromosome sets likely contain mutually exclusive large homologous genomic blocks that are

evolutionarily conserved and can be compared via structural disparity analysis. An initial assessment

of 10 axes revealed eight sets of chromosomes that may be ideal for structural disparity analysis (A).

Visualizing the clustering patterns in bivariate plots can assist with determining axis thresholds for

defining the sets (B, C).

Table 1. Chromosome sets identified from genome assemblies used to conduct landmark-based

estimates of structural disparity. Those chromosomes appearing in bold text were used in the

simulation/validation section of our study. CN = chromosome number (= reference number used in

genomic assemblies), bp = base pairs, UCE = ultraconserved elements. Following removal of an

outlier landmark, the chromosome sets had 185 and 81 landmarks, respectively.

Species	Order	CN	Chromosome set 1 (185 UCEs)			CN	Chromosome set 2 (81 UCEs)			Reference
			Length	UCE landmarks			Length	UCE landmarks		
Bos indicus	Artiodactyla	2	140,680,885 bp	346		26	51,861,174 bp	235		[67]
Bos taurus	Artiodactyla	2	136,231,102 bp	353		26	51,992,305 bp	239		[68]
Bubalus bubalis	Artiodactyla	2	188,164,321 bp	453		23	52,095,120 bp	237		[69]
Capra aegagrus	Artiodactyla	2	128,023,632 bp	316		26	53,792,517 bp	233		[70]
Capra hircus	Artiodactyla	2	136,510,947 bp	356		26	51,421,553 bp	240		[71]
Giraffa tippelskirchi	Artiodactyla	3	190,444,849 bp	420		2	236,459,614 bp	562		[72]
Ovis aries	Artiodactyla	2	250,202,058 bp	569		22	51,512,491 bp	240		[73]
Felis catus	Carnivora	C1	221,611,373 bp	601		D2	88,083,857 bp	350		[74]
Neomonachus schauinslandi	Carnivora	3	192,837,575 bp	552		6	154,800,178 bp	565		[75]
Panthera tigris + P. leo	Carnivora	C1	219,880,392 bp	601		D2	86,999,983 bp	349		[76]
Ceratotherium simum	Perissodactyla	9	84,132,471 bp	322		6	90,918,853 bp	357		[77]
Equus asinus	Perissodactyla	4	92,916,174 bp	324		2	238,820,862 bp	688		[78]
Equus caballus	Perissodactyla	18	82,641,348 bp	320		1	188,260,577 bp	614		[79]

Gorilla gorilla	Primates	3	127,988,552 bp	368	11	127,481,205 bp	418	[80]
Macaca fascicularis	Primates	12	127,712,166 bp	360	9	129,235,353 bp	426	[81]
Macaca mulatta	Primates	12	130,043,856 bp	361	9	134,124,166 bp	427	[82]
Pan troglodytes	Primates	2B	128,755,405 bp	370	10	129,809,613 bp	421	[83]
Papio anubis	Primates	10	126,462,689 bp	358	11	125,913,696 bp	425	[84]
Cricetulus griseus	Rodentia	6	155,611,870 bp	515	3	282,827,514 bp	720	[85]
Mus caroli	Rodentia	2	171,410,997 bp	536	N/A	N/A	N/A	[86]
Mus musculus	Rodentia	2	181,755,017 bp	543	N/A	N/A	N/A	[87]
Mus pahari	Rodentia	3	164,141,553 bp	519	1	172,151,128 bp	531	[86]
Mus spretus	Rodentia	2	182,815,943 bp	537	N/A	N/A	N/A	[88]
Peromyscus maniculatus	Rodentia	4	154,712,973 bp	504	1	193,310,054 bp	533	[89]
Rattus norvegicus	Rodentia	3	169,034,231 bp	517	1	260,522,016 bp	640	[90]

Table 2. Axis thresholding based on metric multidimensional scaling (MMDS) of 567 placental mammal chromosomes from 26 species. Total number of species are unique chromosomes found to cluster on a given axis that are mutually exclusive when compared to clustering patterns of other axes. Threshold value is the smallest axis value identified in each group that is used to define the different clusters. Threshold gap value is the amount of space between the chromosome set and other chromosomes. An asterisk indicates two chromosomes sets where Macaca mulatta and/or

252　*Giraffa tippelskirchi* chromosomes from other sets were within the threshold range. See Figure 4 for

253　additional information.

MMDS Axis	Number of species	Axis threshold	Threshold gap
1	26	14.52	12.03
2	26	-8.90	2.52
3	25	6.96	0.38
4	24	-8.54	2.78
5	26	9.44	0.03
6	26	3.40	0.72
7	26* (*Giraffa* set 5, *Macaca* other)	4.21	0.01
8	24* (*Giraffa* set 2)	3.60	< 0.01
9	0	0	0
10	0	0	0

Landmarking of UCES on the two chromosome sets resulted in 186 (set 1) and 82 (set 2)

landmarks, respectively. Using absolute landmark positions (Figs. S4A, S5A), we found

that most of the variation in both chromosome sets was explained by PC1. In

chromosome set 1, PC1 described 98.9% of the variation and in chromosome set 2,

PC1 described 99.4% of the variation. We discovered that in both chromosome sets,

PC1 was significantly correlated with chromosome size (set 1, rho = -0.65, p < 0.0001;

set 2, rho = 0.76, p < 0.0001) when using absolute size. However, this was not the case

for PC2 (set 1, rho = 0.19, p = 0.3308; set 2, rho = 0.37, p = 0.066). Bounding the

landmark positions increased the amount of variation explained by PC1 (chromosome

set 1, 99.2%, Fig. S4C; chromosome set 2, 99.9%, Fig. S5C), and produced unexpected

patterns with several notable outliers across the first two PCs (e.g. Cricetulus griseus,

Fig. S5C). Size-corrected analyses of chromosome set 1 resulted in each mammal order

occupying a unique region of geno-metric across PCs 1 and 2 (Fig. S4E), but this was not

the case in chromosome set 2 where orders largely overlapped (Fig. S5E). Size-

correction using bounded landmarks resulted in less variation explained by PC1 for

both chromosome sets (chromosome et 1, 89.2%, Fig. S4G; chromosome set 2, 91.6%,

Fig. S5G), and similar outlier patterns that were observed in the raw bounded position

analysis. Mapping of component loadings onto chromosome plots revealed a single

landmark (UCE 6948) that was heavily loaded on PC1 and PC2 of both chromosome

sets (Fig.S6). Removing this outlier landmark changed the inferred structural disparity

patterns across the PCAs with mammalian orders mostly occupying distinctive regions

of geno-metric space in both chromosome set 1 (Fig. S4B, D, F, H) and, apart from a

single size-corrected analysis (Fig. S5F), chromosome set 2 (Fig. S5B, D, H).

Following the removal of the outlier landmark, we found that PC1 explained most

variation across the analyses. When using raw, absolute landmark positions PC1

explained 99.9% of the variance for both chromosome sets (Fig. S7A, S8A). Unlike the

analyses including the outlier landmark, using bounded landmarks reduced the amount

of variation explained on PC1 (95.3%, chromosome set 1, Fig. S7B; 98.6% chromosome

set 2, Fig. S8B) compared to the raw positions. PC2 scores in the bounded analyses

explained 2.0% (chromosome set 1) and 0.8% (chromosome set 2). Notably, the

bounded landmark PC1 and PC2 separated all mammalian orders into distinct regions

of geno-metric space. Size-correction via chromosome size resulted in different

disparity patterns in both datasets with higher disparity within Artiodactyla in the

chromosome set 1 analysis (Fig. S7E) and substantial overlap across orders in the

chromosome set 2 analysis (Fig. S8E). The size-corrected, bounded analyses featured

tighter clustering patterns within mammalian orders than the raw, size-corrected

analyses in both chromosome sets (Figs. S7G, S8G). As observed in the X-linked BUSCO

analyses, we found that using raw landmark positions on the reverse complemented

version of the chromosome resulted in outlier taxa (*Macaca mulatta*, *Mus musculus*,

chromosome set 1, Fig. S7B; *Equus asininus*, *E. caballus*, *Cricetulus griseus*, *Mus*

pahari, *Peromyscus maniculatus*, *Rattus norvegicus*, chromosome set 2, Fig. S8B).

However, unlike the X-linked BUSCO analyses, these outliers were also observed in the

size-corrected analyses (Fig. S7F, S8F). Reverse complemented versions of the raw and

size-corrected bounded analyses were more consistent with their equivalent original

analyses (Fig. S7D, E, S8D, E). As observed in the X-linked BUSCO analyses, the amount

of variance explained by PCs 1 and 2 in the size-corrected, bounded analyses was

identical for both directions of analysis.

Based on our results, we selected the raw, bounded analysis with outlier landmark

removed for further interpretation and analysis (Fig. 5). Interestingly, although they

explained a small proportion of the overall structural variation, PC3 (0.5%, chromosome

set 1, 0.5% chromosome set 2) and PC4 (0.1% chromosome set 1, 0.09% chromosome

set 2) still clustered mammalian orders in largely distinct regions of geno-metric space

for both chromosome sets (Fig. S9). However, *Capra aegragus* was an outlier on PC3 of

chromosome set 1 (Fig.S9A). Synteny analysis of landmarks using chromoMap [35] to

visualize structural differences captured by UCE landmarks revealed that certain

chromosomal regions are more conserved as indicated by low PC loading values of

landmarks into PC1 (blue links, Fig. 6), compared to other regions where landmarks

show higher differences in their placement (maroon links, Fig. 6). Unlike the *Drosophila*

example where landmark inversions explained most structural disparity (Fig. 3),

insertions and deletion events between the UCE landmarks account for the structural

disparity observed in the two placental mammal chromosomes sets (Fig. 6).

Figure 5. Ordination results from principal component (PC) analysis of UCE landmark positions in two

sets of chromosomes from 26 species of placental mammals using bounded raw positions after the

removal of an outlier landmark (see text). Representatives for five orders (Artiodactyla, Carnivora,

Perissodactyla, Primates and Rodents) were included (Table1). The depicted structural variation is

inferred from 185 (chromosome set 1, A) and 81 (chromosome set 2, B) landmarks, respectively. In

both datasets, the first two PCs explained > 99% of the variation. Comparison of PCs 3 and 4 for

both chromosome sets is provided in Figure S9.

Figure 6. Select synteny link maps between *Bos indicus* and other placental mammals using bounded, absolute landmarking. Principal component loadings for various landmarks are indicated by the color of the links. The chromosome number (Chr) is indicated next to each species' silhouette. The left and right columns are results from chromosome set 1 and set 2, respectively. Comparisons are (from top to bottom) between *Bos indicus* to *Capra aegagrus* (Artiodactyls), *Bos indicus* and *Equus asinus* (Perissodactyla), *Bos indicus* and *Gorilla gorilla* (Primates), *Bos indicus* and *Mus musculus* (Rodentia), and *Bos indicus* and the liger (*Panthera tigris + P. leo*, Carnivora). Synteny link maps were generated using chromoMap [35].

Structural disparity increases through time. In the simulation study, > 99% of the

variation in landmark placement was explained by PC1 with PC2 explaining 0.2% of the

variation. As predicted, pairwise increase in disparity (ID) increased with the amount of

simulated evolution confirming that, at least under neutral (random) conditions,

disparity measures should correspond to relative amounts of genomic evolution (Fig.

7A-D). We compared the variance of log-transformed ID scores for indel versus

inversion simulations using two-way ANOVAs. We found that on both axes, inversions

caused a significantly higher increase in disparity than indels (PC1, $F = 2.88$, $p = 0.0251$;

PC2, $F = 3.34$, $p = 0.0121$; Fig. S10). Using empirical data, we found that inter-order

disparity was significantly higher than intra-order disparity for both PC1 (chromosome

set 1, $W = 63997$, $p < 0.0001$; chromosome set 2, $W = 67457$, $p < 0.0001$, Fig. 7E, F) and

PC2 (chromosome set 1, $W = 67997$, $p < 0.0001$; chromosome set 2, $W = 61201$, $p <$

0.0001, Fig. 7G, H). We found that protein-coding content tended to be more similar

within mammalian orders than between them, however, this was not a significant result

($W = 499.5$, $p = 0.0722$; nonparametric Wilcoxon tests; SI Appendix, Fig. S11).

Figure 7. Log-transformed increase in disparity (ID) scores inferred from principal component (PC)

analysis of ancestral and descendent insertion-deletion simulations for PC1 (A) and PC2 (B). Log-

transformed ID scores inferred from inversion simulations for PC1 (C) and PC2 (D). Insertion-deletion

simulation results are depicted in purple and inversion simulation results are depicted in blue.

Observed empirical disparity scores for within and between mammalian orders for PC1 (E) and PC2

(F) of chromosome set 1 and PC1 (G) and PC2 (H) of chromosome set 2 are depicted in green.

Phylogenetic signal of structural variants. We found that the structural disparity

captured by the most explanatory axes in all datasets had significant phylogenetic

signal in 58% of the test cases (14/24 tests, Table 3). This included some cases where

PC3 (describing < 1% of the overall variation) had significant phylogenetic signal (e.g. X-

linked BUSCO genes, placental mammal UCE chromosome sets). Values of K and λ

were also high (indicative of putative phylogenetic signal) in several non-significant

tests. Ancestral state estimations of disparity revealed an increase in structural

disparity related with the evolution of an X chromosome in the ancestor of placental

and marsupial mammals (Fig. 8A), the evolution of a pericentric inversion on

chromosome 5 in the *Drosophila virilis* group (Fig. 8B), and phylogenetically associated

patterns of structural disparity increase or decrease in placental mammals (Fig. 8C, D).

Figure 8. Ancestral state estimation of structural variants (SV) using select principal component (PC)

scores from disparity analyses conducted herein. Structural variants detected using disparity analysis

often have phylogenetic signal (Table 2) and can be used to quantitatively study the rate of SV

evolution. Examples from this study include human X-linked BUSCO gene disparity of amniote

vertebrates where structural disparity increases following the evolution of the X chromosome (PC1,

A), the evolution of a speciation-associated paracentric inversion in chromosome 5 of the *Drosophila*

virilis group that caused rearrangements in the position of UCE landmarks (PC3, B), and the

identification of placental mammal orders with the least (chromosome set 1, PC1, C) and most

(chromosome set 2, PC1, D) conserved UCE structure, respectively. Phylogenies were obtained from

Kumar et al. [63].

Table 3. Results of phylogenetic signal tests using K [64] and λ [65] statistics for three most

explanatory principal components (PCs) of structural variant disparity from the empirical analyses

conducted herein. P-values are based on 1,000 randomizations (K) and likelihood ratio tests (λ).

Significant tests are indicated with bold text.

Principal component	Variance explained (%)	Phylogenetic signal	
		K	λ
X-linked BUSCO landmarks from amniotes: PC1	71.2	**1.3418 (p = 0.001)**	**1.160 (p < 0.001)**
X-linked BUSCO landmarks from amniotes: PC2	12.1	0.4237 (p = 0.501)	< 0.001 (p = 1.0)
X-linked BUSCO landmarks from amniotes: PC3	7.5	**0.6591 (p = 0.049)**	0.825 (p = 0.454)
UCE landmarks from *Drosophila*: PC1	94.0	0.6935 (p = 0.568)	**1.151 (p = 0.013)**
UCE landmarks from *Drosophila*: PC2	5.2	**0.9949 (p = 0.027)**	**1.151 (p = 0.004)**
UCE landmarks from *Drosophila*: PC3	0.07	0.6831 (p = 0.195)	< 0.001 (p = 1.0)
UCE landmarks placental mammals (set 1): PC1	95.3	**0.2173 (p = 0.022)**	**0.866 (p < 0.001)**
UCE landmarks placental mammals (set 1): PC2	4.1	**0.2600 (p = 0.005)**	**0.818 (p < 0.001)**
UCE landmarks placental mammals (set 1): PC3	0.5	0.0643 (p = 0.717)	**0.476 (p = 0.004)**
UCE landmarks placental mammals (set 2): PC1	98.6	**0.2677 (p = 0.009)**	**0.849 (p < 0.001)**
UCE landmarks placental mammals (set 2): PC2	0.8	0.1104 (p = 0.294)	**0.733 (p < 0.001)**
UCE landmarks placental mammals (set 2): PC3	0.5	0.0573 (p=0.708)	0.659 (p = 0.235)

Discussion

Using a morphometrics-inspired approach we identified SVs associated with

chromosome inversions, insertion-deletion events (=relative size change), and

phylogeny. These qualities mirror the utility of morphological disparity analysis as

applied to complex phenotypic structures. In a phenotypic context, disparity can be

descriptive or serve as a proxy for phylogenetic affinities and ecology [22]. For example,

disparity is widely used to understand adaptive radiation in the paleontological record

[36]. Our simulation and empirical results suggest that, like classic morphological

disparity, structural disparity of chromosomes increases through time (Fig. 7). In a

genomic context, we are optimistic that disparity can play a similar role in

understanding how chromosome architecture varies among species and can be used

for quantitative comparisons to study organismal evolution and adaptation. Below we

discuss the results of our empirical case studies, suggest guidelines for best practice

when conducting structural disparity analysis, and present several future directions for

applications and methods development.

Structural disparity and the evolution of chromosome architecture. Disparity analyses

of SVs enable a broad understanding of how similar (or dissimilar) chromosome

structures are across different species based on the distribution of conserved landmark

sequences like BUSCO genes and UCEs. This analytical quality is particularly

advantageous for studying large-scale shifts in chromosome organization across

organisms relevant to the study of supergenes and homologous synteny blocks (sensu

[16]). In our study we found that following the evolution of a key mammalian sex

chromosome, structural disparity of 53 BUSCO genes increased compared to other

amniotes (Fig. 2A, Fig. 8A). This disparity increase appears to be associated with an

overall increase in chromosome size in those species where the landmark genes are

located on an X chromosome. Due to the cessation of regular recombination, size

increases and decreases in vertebrate sex chromosomes are well-documented [37].

When we scaled the positions by chromosome size, we observed that placental and

marsupial mammals were no longer the most disparate group (Fig. S2E, G), further

supporting chromosome size increase as the key driver of structural disparity among

BUSCO genes in the mammalian X chromosome. Following size-correction, structural

disparity analysis identified high disparity among the bird chromosomes analyzed,

which from viewing the synteny plot (Fig. 2B) appears to be related to several inversions

(within birds) associated with the landmarks. The ability for structural disparity analyses

to capture SVs associated with inversions in closely related species or populations was

also observed in our study of chromosome 5 from the *Drosophila virilis* group based on

191 UCE landmarks (Fig. 3). In this simple example, we observed that the first three

principal components all corresponded to distinctive changes previously linked to

speciation [27]. Although not all axes had statistically significant phylogenetic signal

(Table 3), small sample sizes may have contributed to this outcome (see 38) as even

those axes without significant signal (e.g. PC3) were consistent with phylogenetically

associated changes (Fig. 8B).

The de novo study of structural disparity using tetrapod UCEs with placental mammals

revealed order-specific disparity patterns in PCs 1 and 2 of both chromosome sets (Fig.

5). In addition to revealing clear phylogenetic patterns in UCE-related structural

disparity, these analyses also revealed that disparity patterns are order-specific across

different chromosomes (Fig. 8C, D). Specifically, using the chull function in the R

package 'sp' [39] we found that convex hull values for each mammalian order on PCs 1

and 2 were perfectly correlated across chromosomes set 1 and set 2 (Pearson's

correlation coefficient = 0.996, p = 0.0003). This suggests that order-specific structural

disparity associated with UCEs may be similar across different genomic regions. Given

the importance of UCEs in development and gene regulation [40] these types of

genome-wide patterns are important to explore as additional genomic data become

available.

Best practices for structural disparity analysis. Based on comparative analyses of the X-

linked BUSCO genes from amniote vertebrates and UCEs from placental mammals, we

can recommend a set of best practices for structural disparity that will allow users to

avoid several analytical artifacts that we encountered. First, examining component

loadings for each landmark is important for identifying outlier landmarks that may

obscure more generalized signal of other landmarks. In our study of placental

mammals, we found that a single outlier (UCE 6948) explained most variation in the first

two PCs (Fig. S6). The presence of this UCE in both datasets demonstrates that it is not

single-copy and justifies removing these types of spurious landmarks prior to disparity

analysis. Second, using raw, absolute positions may lead to outlier taxa if the landmarks

are clustered at one end of a large chromosome. This phenomenon was observed in

both the X-linked BUSCO gene dataset and the placental mammals (Fig. S2B, S7B,

S8B). In some cases, scaling the positions based on chromosome size may resolve this

issue (Fig. S2F), whereas in others it does not (Fig. S7F, S8F). We found that the use of

bound landmark positions resolves this analytical artifact associated with the

directionality of the chromosome sequence. If size is an important parameter for

understanding structural disparity, we recommend using bound, absolute positions

(Fig. S2C, S2D, S7C, S7D, S8C, S8D). If patterns like inversions and landmark

translocations are of primary interest, we recommend using size-corrected and bound

landmark positions (Fig. S2G, S2H, S7G, S7H, S8G, S8H). Finally, if using the de novo

approach and clustering chromosomes with MMDS to identify homologous landmark

regions for analysis we recommend viewing various axes as bivariate plots (Fig. 4B, 4C)

where clustering may be more obvious than gap threshold sizes would suggest on the

individual axes (Table 2).

A complementary tool for studying structural changes in chromosomes. Dimensionality

reduction is already in wide use in comparative genomics (e.g., uniform manifold

approximation and projection, [41]). Structural disparity analysis is therefore a natural

companion method for assessing structural differences across the chromosomes of

different species. It complements existing methods like WGA and synteny analysis by

capturing a different dimension of how chromosomes vary across species. Synteny

analysis is a powerful tool for identifying similarities in gene order and arrangement.

While synteny analysis does capture spatial variation (unlike WGA), it is known to be

very sensitive to parameter settings [42]. Furthermore, some methods of synteny

inference are performed on post-WGA datasets meaning that structural information

about the whole chromosomes has been removed a priori [15]. In contrast, landmark-

based estimates of genome structure offer at least two qualities that syntenic analyses

lack (Fig. S12). First, minor chromosomal rearrangements will be interpreted as breaks

in synteny and ignored. In other words, inversions within syntenic regions not involving

all the contiguous markers will break synteny. Second, large gap regions between

contiguous orthologs can also break synteny (depending on parameter settings). The

latter example highlights another important difference; disparity analyses will capture

distance information between landmarks whereas synteny only provides landmark

(ortholog) order information.

Applications beyond diploid genomes and evolutionary biology. Apart from diploid

organisms such as vertebrates and insects, the structural disparity pipeline could also

be implemented on polyploid organisms when chromosome phasing has been

performed. Utilizing the different phased genome assemblies, the pipeline can identify

structural differences between haplotigs of the genome. In addition, the pipeline may

be useful for measuring structural evolution of pangenomes without the need for a

reference input genome. Instead, a class/category of genetic markers could be used as

structural landmarks. As a visualization tool, structural disparity analysis may

complement the routinely used pangenome graph building methods (e.g.[43]). Lastly, a

potential use of structural disparity analysis relates to cancer genomics. The genomes

of somatic cancer cells undergo rapid structural evolution and SVs in cancer cells are

very specific to the type of cancer [44]. SVs in cancer cells have an impact on gene

expression through modifying structural accessibility, DNA methylation (and more) in

the cis-regulatory regions of the genome [44], therefore the ability to identify genomic

rearrangement is valuable (especially when using long-read data).

Avenues to expand the method towards genomic disparity. Structural disparity analyses

could be developed for whole-genome comparisons. At present, the pipeline performs

chromosome to chromosome comparisons, limiting comparisons across genomic

regions that have undergone translocation, fission, or fusion events. With greater

awareness of the frequency of these events, weighting parameters could be defined to

measure global genomic disparity levels. While we limited our validation analyses to

using BUSCOs and UCEs as landmarks and a handful of (largely homologous)

chromosomes, landmarks could be any conserved genetic sequences which can be

accurately mapped. Variation in the placement of conserved sequences can be used to

study colocalization of transcribed genes (e.g. BUSCO [45]), cis-regulatory elements,

enhancers, promoting regions, and more. The ethos of the approach is also scalable to

three-dimensional genome structure once that information is more broadly available for

many species (see [46, 47]). The extensive literature on geometric morphometrics offers

substantial guidance for how 3D comparisons may be performed in the future [25].

Landmark-based estimates of genomic disparity provide a rapid mechanism for

comparisons of genome structure across species and populations, and we anticipate

they will expedite discovery and study of novel structural patterns within comparative

genomics.

Methods

Bioinformatics and identification of landmarks. The bioinformatics and computational

analyses were performed on Crop Diversity HPC, described by Percival-Alwyn et al. [48].

The structural disparity pipelines are modular and can be modified for a variety of other

uses. Three step-by-step tutorials along with custom R and shell scripts are available on

GitHub (https://github.com/nhm-herpetology/genomic-

disparity/blob/main/README.md). Structural disparity analysis can be conducted using

a variety of genomic landmarks assuming they are single copy and orthologous (e.g.

BUSCO genes [45], UCEs [32,34]). In this study, we used human X-linked BUSCO genes

and UCEs as landmarks. We selected human X-linked genes because many of them are

found in homologous blocks on a single chromosome across amniote vertebrate

animals [18] and UCEs because they are highly conserved, lack repeats, and widely

distributed in genomes [34, 49]. UCEs have also been implicated as relevant to the

structure of key developmental and transcriptomic pathways [49, 50], making variation

in their relative chromosomal placements of biological interest.

Sensitivity analyses related to chromosome size, complementarity and landmark

boundaries. Data transformations associated with object size are a well-studied

component of morphometrics [51]. In our datasets where structural disparity was

correlated with chromosome size, we conducted a series of analyses to establish best

practice for structural disparity analysis. We analyzed these datasets in four ways using:

(1) raw positions, (2) bounded, raw positions, (3) chromosome size-corrected positions,

and (4) bounded, size-corrected positions. We repeated each analysis using the reverse

complement version of the four datasets for a total of eight versions of the analysis.

Boundary setting was achieved for each landmarked chromosome using a two-step

procedure: (1) re-positioning all the landmarks relative to the first landmark and (2)

using the position of the last landmark to establish the size of the genomic region being

analyzed. Absolute size correction was performed with R functions that divided the total

chromosome size by each landmark position. Bounded size correction was achieved by

dividing the size of the bounded genomic region by each landmark position with R

functions.

BUSCO genes and amniote vertebrates. Using the BUSCO gene annotation file from

Lovell et al. [18], we extracted all the genes annotated in the human (*Homo sapiens*) X

chromosome. We utilized this list of x-linked genes and identified homologous

chromosomes in other species based on the proportion of the human x-linked genes

mapped on them. The other species included in the study were four placental

mammals (*Mus musculus*, *Rhinolophus ferrumequinum*, *Tursiops truncatus*, and

Choloepus hoffmanni), three marsupial mammals (*Sarcophilus harrisii*, *Monodelphis*

domestica, and *Trichosurus vulpecula*), two monotreme mammals (*Tachyglossus*

aculeatus and *Ornithorhynchus anatinus*), four birds (*Gallus gallus*, *Cygnus olor*,

Calypte anna, and *Taeniopygia guttata*) and two squamate reptiles (*Thamnophis*

elegans and *Lacerta agilis*). When there was more than a single chromosome that

shared human x-linked genes (e.g. *Tachyglossus, Ornithorhynchus*, and *Trichosurus*),

the chromosome with the higher proportion of genes was used to maximize the number

of landmarks available for structural disparity analysis. Lovell et al. (2022) also analyzed

the budgie (*Melopsittacus undulatus*) genome, however, because our pipeline did not

identify focal landmarks on a single chromosome, we did not include this species in the

structural disparity analysis. Using the gff3 file approach (Fig. 1), we identified human x-

linked BUSCO genes that were homologous across all the species' chromosomes.

Detailed instructions for identifying and extracting these genes are available in Step 1 of

the GitHub tutorial. Next, chromosomes are prepared for downstream analysis using a

pipeline which requires the following R statistical software [31] packages: rehshape2

[52], matrixStats [53], tidyr [54], dplyr [55], and ggplot2 [56]). From 838 genes we

identified 53 that were present in all focal species making them suitable for use as

landmarks in a structural disparity analysis. Detailed instructions for identifying and

extracting landmarks are available in Step 2 of the GitHub tutorial. We also conducted

data matrix manipulation and principal component analysis in R, with code available in

Step 3 of the GitHub tutorial. For size-corrections, we obtained chromosome sizes from

NCBI. We tested for correlations between chromosome size and disparity using

Spearman's rho nonparametric tests in R. To create a synteny plot for comparative

purposes, we used the Chromomap package in R [35]. A list of the 53 human X-linked

BUSCO genes used as landmarks can be found in Appendix S1.

Drosophila structural variants, speciation and UCEs. Based on the findings of Poikela et

al. [27], we landmarked chromosome 5 for *D. americana*, *D. flavomontana*, *D. montana*,

D. novamexicana, and two individuals of *D. virilis*. This chromosome has known

inversions and translocations and may have contributed to speciation in the group. We

downloaded chromosome 5 from the *Drosophila virilis* group either via NCBI Entrez

Direct UNIX E-utilities (2) or directly from the literature (1). The following chromosomes

sequence were used: *D. americana* (NCBI accession no: CM061086.1), *D.*

flavomontana (1), *D. montana* (1), *D. novamexicana* (CM061080.1), and two individuals

of *D. virilis* (CM017608.2 and CM061075.1). Detailed instructions for downloading

chromosome sequences are available in Step 1 of the GitHub tutorial. For reference

mapping of ultraconserved elements (UCEs), we utilized a published set of

ultraconserved elements (UCEs) from dipterans [3]. We indexed the chromosome

assemblies and reference mapped UCEs using default settings in BWA-MEM [29] and

SAMtools [57]. Custom shell commands are used to loop the reference mapping and

create a spreadsheet that contains presence and absence information for the UCEs on

each chromosome 5 variant included in the analyses. Detailed instructions on this

reference mapping approach are available in Step 2 of the associated GitHub tutorial.

Next, chromosomes were prepared for downstream analysis using an R pipeline as

described above. The pipeline includes three steps (i) remove mapped UCEs that the

chromosomes do not share, (ii) complementarity checks to ensure all chromosomes

have comparable orientations (5' to 3'), and (iii) merging redundant UCE probes that

target the same UCE locus. The pipeline resulted in the identification of 191

homologous UCE landmarks for chromosome 5 of the *Drosophila virilis* group. Detailed

instructions for the preparation pipeline are available in Step 3 of the GitHub tutorial. In

the final step of the disparity analysis, we conducted a principal component analysis

(PCA) and visualized the results. The input file for the PCA is available on GitHub as

'Drosophila_PCA.csv'. We used R base functions along with dplyr to format the data

matrix and conduct the PCA. Visualization was conducted using ggplot2 [56] and

cowplot [58]. Detailed instructions for this final step are available in Step 4 of the

GitHub tutorial.

De novo landmark discovery and placental mammal genomes. To apply our disparity

pipeline to a group where homology among chromosomes was unknown, we used 26

nuclear genomes from five orders of placental mammals (Table 1). Because placental

mammal orders have diverged over millions of years [59] the homology of their

chromosomes is fragmentary, making them a good candidate for exploring the extent to

which disparity analysis can be applied to single genomic regions in divergent groups.

We downloaded mammal genome assemblies using a custom script

(chromosome_download.sh) that we developed to work with a standard input file

(chromosome_download.config). Setting up a batch download for a subset of the

mammal taxa (26 species) is described in detail in Step 1 of the GitHub tutorial. For

reference mapping, we utilized a published set of tetrapod UCEs [32]. We indexed and

reference-mapped UCEs on the 1,004 mammal chromosomes using default settings in

BWA-MEM [29] and SAMtools [57]. Reference mapping was conducted using a custom

shell script that loops through the chromosome sequences (landmark_mapping.sh).

We tested the utility of minimap2 [30] for mapping conserved regions to the genome

assemblies utilizing default settings on chromosome set 2 of the mammalian genomes.

We then compared these results to those obtained using BWA MEM. We observed that

BWA MEM mapped higher numbers of UCEs than minimap2 (Fig. S1). This difference in

mapping quality could be explained by the short length of the UCE sequences used, at

about 180 bp length [32] because BWA MEM can achieve higher accuracy with short-

read mapping and produces less false negatives than minimap2 [30]. However,

minimap2 may be preferable if larger sequences were used as landmarks, and that

approach could then be easily combined with our post-mapping pipeline steps. In

addition to providing a UCE presence-absence output file for downstream analysis in

the pipeline, landmark_mapping.sh outputs the number of (UCE) landmarks mapped

and length information for each chromosome. Detailed instructions for refence

mapping are available in Step 2 of the GitHub tutorial.

Interestingly, we found that the number of UCEs mapped onto each mammal

chromosome was positively and significantly correlated with chromosome size (r =

0.75, p < 0.05; SI Appendix, Fig. S2). To identify sets of chromosomes that share

homologous landmarks, we applied metric multidimensional scaling (MMDS) on the

presence-absence matrix generated by landmark_mapping.sh. Using MMDS scores

across 10 axes we identified mutually exclusive sets of chromosomes that consisted of

only one chromosome from a species. These chromosome sets were identified by

selecting a threshold value on the relevant axes of the MMDS plot. Detailed instructions

for clustering chromosomes and thresholding are available in Step 3 of the GitHub

tutorial. To determine if protein content was similar on chromosomes sets (implying

that they share functional, protein-coding homologs in addition to UCE landmarks, we

used protein annotations from the Ensembl database (release 110; [60]), to generate

Jaccard similarity coefficients for comparing protein compositions on the two focal

chromosome sets across a subset of placental species where annotations were

available (N = 14). After extracting the protein-coding markers' identities from the

Ensembl annotation files of each species (for the specific chromosomes in set 1 and set

2 separately), we utilized a custom defined Jaccard's similarity index which removed

case sensitivity in the names of the markers (available on GitHub page as

'Jaccard_proteins.R'). This index estimated pairwise similarities between each

chromosome in set 1 and set 2 respectively, as well as for all chromosome pairs in both

the sets. The latter was done to test if chromosomes within a set show higher similarity

in protein-coding markers than chromosomes between the two sets. Following the

identification of chromosome sets, we prepared them for PCA by (i) removing UCEs that

did not occur on all chromosomes, (ii) checking the directional orientation (5' to 3'), and

(iii) merging redundant UCE probes. Unlike the highly conserved chromosomes from the

Drosophila example, determining the directionality of mammal chromosomes was not

always obvious. In other words, sometimes we observed mixed forward and reverse

mappings of UCE landmarks, suggesting that only particular segments of the

landmarked region were sequenced in the opposite direction of most taxa. This is likely

explained by within-chromosome recombination and to account for it we used a

majority rule criteria so that we only changed chromosome orientations if >50% of

landmarks were mapped in the opposite direction of the reference chromosomes.

Detailed instructions for the chromosome preparation pipeline are available in Step 4 of

the GitHub tutorial. Finally, we conducted a PCA on each of the chromosome sets and

visualized the results which are detailed in Step 5 of the GitHub tutorial. The input files

for the PCA analysis are available on GitHub as 'Set_1_mammals_PCA.csv' and

'Set_2_mammals_PCA.csv', respectively. As with the Drosophila example, applying the

disparity pipeline to placental mammal genomes required the following R packages:

rehshape2, matrixStats, dplyr, ggplot2 and cowplot.

Simulation and other tests for increased disparity through time. We simulated two

common evolutionary shifts in chromosome structure that occur as species and

populations diverge: insertion-deletion (indel) events and chromosomal inversions. We

performed these simulations using chromosome set 1 from three species subsampled

from the larger dataset, *Capra aegagrus*, *Mus pahari* and *Neomonachus schauinslandi*

(Table 1). We selected these taxa based on phylogenetic diversity (representing

Artiodactyla, Carnivora and Rodentia) and being outliers in some analyses (*Capra*

aegagrus, Fig. S9). We simulated evolutionary events using SimuG [61]. We designed

the experiment logarithmically, simulating 1, 10, 100, 1,000, and 10,000 random indels

or inversions on the ancestral chromosome of the five focal species. We ran five

replicated simulations for each of the five focal species. To assess the change in

disparity between ancestral chromosomes and their simulated descendants, we used a

simple pairwise metric of increase in disparity (ID):

$$ID = |\, PC^A - PC^B \,|$$

Where PC^A is the principal component score of the ancestral chromosome, PC^B is the

principal component score of the descendent chromosome, and ID is calculated by

taking the absolute difference of these values. As such, ID is an intuitive measurement

of how much landmark placement changed during the simulations, where a value of 0

indicates no change and larger values correspond to increasingly disparate landmark

placements. We predicted that if genomic disparity reflects the degree of evolutionary

change of architecture between two genomes, there should be a positive correlation

between ID and the number of simulated events. While analysis of individual PCs may

be problematic when data have high effective dimensionality [62], here the dominance

of PC1 and the overwhelming correlation with chromosome size on that primary axis,

justifies separate analysis of these components. We compared the original

chromosome (set 1) sequence from three species of mammals, referred to as the

'ancestral chromosome', to simulated 'descendent chromosomes' which we made by

simulating random indel or inversion events (see main text for more detail). Following

the disparity pipeline described above, we mapped UCEs onto the simulated

chromosomes, extracted homologous landmarks and carried out a PCA. PCA also

included ancestral chromosomes from each species. We calculated the disparity

between the ancestral chromosomes and simulated descendent chromosomes using

the equation described in the main manuscript. Disparity was calculated using the

logarithm of the absolute difference between the PC values of the ancestral

chromosome and each simulated chromosome. Furthermore, we tested for differences

between the disparity measures of indel-simulated and inversion-simulated

chromosomes using two-way ANOVAs implemented in R.

To provide empirical context for the simulation results, we tested for the signature of

disparity increase in our placental mammal datasets using nonparametric Wilcoxon

tests in R. If UCE-inferred disparity increases with evolutionary time, we expected inter-

order disparity to be significantly higher than intra-order disparity for the most

explanatory PCs because there has been more evolutionary divergence between

mammalian orders than within them. We also generated Jaccard similarity coefficients

for comparing protein compositions between and within mammalian orders.

Tests for phylogenetic signal and ancestral state estimation. Using the three most explanatory structural disparity PCs, we tested for phylogenetic signal in the (1) amniote vertebrate, (2) *Drosophila virilis* group, and (3) placental mammal (both chromosome sets 1 and 2) datasets. Phylogenies for each dataset were obtained from Kumar et al. [63]. We used two common metrics for phylogenetic signal, K [64] and λ [65]. We implemented test using the phytools package [66] in R. We assessed the significance of the tests using 1,000 randomizations (K) and likelihood ratio tests of the observed data to that estimated under Brownian motion (λ). To place structural disparity of genomic regions in a phylogenetic context, we estimated putative ancestral disparity values we used the 'fastAnc' function in phytools for select PCs in each dataset.

Acknowledgments

We thank members of the Goswami Lab and NHM Herpetology Group for many helpful comments on experimental design and methods development. The authors acknowledge ChatGPT for assistance with some R coding (https://openai.com/index/chatgpt/). We thank Kate Thomas for an introduction to several R packages. Several delegates at the 2023 Society for Integrative and Comparative Biology (Austin, Texas, USA) and 2023 Evolution (Albuquerque, New Mexico, USA) and Young Investigators meeting 2024 (Bhopal, India) conferences provided valuable suggestions on earlier version of this work and putative applications for structural disparity analysis. We thank Jason Head and members of the Department of Zoology at the University of Cambridge for helpful suggestions related to size correction. We thank Christophe Liedtke for discussions on dimensionality reduction in genomics. We are indebted to Brant Faircloth and two anonymous reviewers for

comments that greatly improved the quality of this manuscript. We are grateful to Iain

Milne for bioinformatics support. The authors acknowledge Research Computing at the

James Hutton Institute for providing computational resources and technical support for

the "UK's Crop Diversity Bioinformatics HPC" (BBSRC grants BB/S019669/1 and

BB/X019683/1), use of which has contributed to the results reported within this paper.

This study was part of a project funded by the European Commission through Marie

Skłodowska-Curie Actions to A.V.M. (Grant Agreement ID: 101027832). This work also

received support from the Genomics Research Theme at the Natural History Museum,

London.

References

1.   Genereux DP, Zody MC, Shen W, McGaugh SE, Stemple DO, Roberts AC, et al.

2020. A comparative genomics multitool for scientific discovery and conservation.

Nature 587:240–245. doi:10.1038/s41586-020-2874-7

2.   Feng S, Stiller JW, Deng Y, Armstrong J, Burt DW, Kang H, et al. 2020. Dense

sampling of bird diversity increases power of comparative genomics. Nature 587:252–

257. doi:10.1038/s41586-020-2873-8

3.   Osmanski AB, Maples B, Blahnik J, Griesmann L, Willis WM, Schmitz RJ, et al.

2023. Insights into mammalian TE diversity through the curation of 248 genome

assemblies. Science 380:eabn1430. doi:10.1126/science.abn1430

4.   Rubin GM, Yandell MD, Wortman JR, Clayton DF, Meeker S, Filipski AJ, et al. 2000.

Comparative genomics of the eukaryotes. Science 287:2204–2215.

doi:10.1126/science.287.5461.2204

5.   Zhang X, Wessler SR. 2004. Genome-wide comparative analysis of the

transposable elements in the related species Arabidopsis thaliana and Brassica

oleracea. Proc Natl Acad Sci U S A 101:5589–5594. doi:10.1073/pnas.0401243101

6.   Yuan Y, Liu X, Wang J, Sun Z, Smith AB, Peterson DG, et al. 2021. Comparative

genomics provides insights into the aquatic adaptations of mammals. Proc Natl Acad

Sci U S A 118:e2106080118. doi:10.1073/pnas.2106080118

7.   Lynch M, Walsh B. 2007. The origins of genome architecture. Sunderland (MA):

Sinauer Associates.

8.   Krupina K, Goginashvili A, Cleveland DW. 2024. Scrambling the genome in

cancer: causes and consequences of complex chromosome rearrangements. Nat Rev

Genet 25:196–210. doi:10.1038/s41576-023-00574-6

9.   Hu W, Xie J, Song X, Chen S, Ling H, Li Z, et al. 2020. Nondestructive 3D image

analysis pipeline to extract rice grain traits using X-ray computed tomography. Plant

Phenomics 2020:3414926. doi:10.34133/2020/3414926

10.   Graves JAM. 2016. Evolution of vertebrate sex chromosomes and dosage

compensation. Nat Rev Genet 17:33–46. doi:10.1038/nrg.2015.2

11.   Banse P, et al. Forward-in-time simulation of chromosomal rearrangements: The

invisible backbone that sustains long-term adaptation. Mol Ecol [Internet].

n.d.;(n/a):[page-unspecified].

12.   Chaves R, Adega F, Santos S, Guedes-Pinto H, Heslop Harrison JS. 2002. [Title

missing]. Cytogenet Cell Genet. 96:113–116. doi:10.1159/000063020

13.   Balachandran P, Beck CR. 2020. Structural variant identification and

characterization. Chromosome Res. 28:31–47. doi:10.1007/s10577-019-09623-z

14.   De Coster W, Van Broeckhoven C. 2019. Newest methods for detecting

structural variations. Trends Biotechnol. 37:973–982.

15.   Krasheninnikova K, et al. 2020. halSynteny: a fast, easy-to-use conserved

synteny block construction method for multiple whole-genome alignments.

GigaScience. 9:giaa047. doi:10.1093/gigascience/giaa047

16.   Larkin DM, et al. 2009. Breakpoint regions and homologous synteny blocks in

chromosomes have different evolutionary histories. Genome Res. 19:770–777.

doi:10.1101/gr.087407.108

17.   Chaney L, Sharp AR, Evans CR, Udall JA. 2016. Genome mapping in plant

comparative genomics. Trends Plant Sci. 21:770–780.

doi:10.1016/j.tplants.2016.05.003

18.   Lovell JT, et al. 2022. GENESPACE: syntenic pan-genome annotations for

eukaryotes. bioRxiv. 2022.03.09.483468. doi:10.1101/2022.03.09.483468

19.   Quigley S, Damas J, Larkin DM, Farré M. 2023. syntenyPlotteR: a user-friendly R

package to visualize genome synteny, ideal for both experienced and novice

bioinformaticians. Bioinformatics Adv. 3:vbad161. doi:10.1093/bioadv/vbad161

20.   Armstrong J, Hickey G, Diekhans M, et al. 2020. Progressive Cactus is a multiple-

genome aligner for the thousand-genome era. Nature. 587:246–251.

doi:10.1038/s41586-020-2871-y

21.   Foote M. 1997. The evolution of morphological diversity. Annu Rev Ecol Evol Syst

28:129–152. doi:10.1146/annurev.ecolsys.28.1.129

philpapers.org+1onlinelibrary.wiley.com+1

22.   Guillerme T, Cooper N, Martin RD, Wills MA. 2020. Disparities in the analysis of

morphological disparity. Biol Lett 16:20200199. doi:10.1098/rsbl.2020.0199

23.   Fabre AC, Botero-Castro F, Delsuc F, Douzery EJ, Goswami A. 2020.

Metamorphosis shapes cranial diversity and rate of evolution in salamanders. Nat Ecol

Evol 4:1129–1140. doi:10.1038/s41559-020-1200-4

24.   Briggs DEG, Fortey RA, Wills MA. 1992. Morphological disparity in the Cambrian.

Science 256:1670–1673. doi:10.1126/science.256.5064.1670

25.   Zelditch ML, Lundrigan BL, Garland T Jr. 2004. Developmental regulation of skull

morphology. I. Ontogenetic dynamics of variance. Evol Dev 6:194–206.

doi:10.1111/j.1525-142X.2004.04025.x

26.   Christmas MJ, et al. 2023. Evolutionary constraint and innovation across

hundreds of placental mammals. Science 380:eabn3943.

doi:10.1126/science.abn3943

27.   Poikela N, Laetsch DR, Hoikkala V, Lohse K, Kankare M. 2024. Chromosomal

inversions and the demography of speciation in Drosophila montana and Drosophila

flavomontana. Genome Biol Evol 16:evae024. doi:10.1093/gbe/evae024

28.   Bejerano G, Pheasant M, Makunin I, Stephen S, Kent WJ, Mattick JS, et al. 2004.

Ultraconserved elements in the human genome. Science 304:1321–1325.

doi:10.1126/science.1098119

29.   Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–

Wheeler transform. Bioinformatics 25:1754–1760. doi:10.1093/bioinformatics/btp324

30.   Li H. 2018. Minimap2: pairwise alignment for nucleic acid sequences.

Bioinformatics 34:3094–3100. doi:10.1093/bioinformatics/bty191

31.   R Core Team. 2021. R: A language and environment for statistical computing.

Vienna (Austria): R Foundation for Statistical Computing. Available from:

https://www.R-project.org/ faircloth-

lab.orgbibsonomy.org+10ropensci.org+10scirp.org+10

32.   Faircloth BC. 2017. Identifying conserved genomic elements and designing

universal bait sets to enrich them. Methods Ecol Evol 8:1103–1112. doi:10.1111/2041-

210X.12754 academic.oup.com+9faircloth-lab.org+9pmc.ncbi.nlm.nih.gov+9

33.   Yusuf LH, Tyukmaeva V, Hoikkala A, Ritchie MG. 2022. Divergence and

introgression among the virilis group of Drosophila. Evol Lett 6:537–551.

34.   Faircloth BC, Branstetter MG, White ND, Brady SG, Fisher BL, Joubert MF, et al.

2012. Ultraconserved Elements Anchor Thousands of Genetic Markers Spanning

Multiple Evolutionary Timescales. Syst Biol 61:717–726. doi:10.1093/sysbio/sys004

arxiv.org+6faircloth-lab.org+6pmc.ncbi.nlm.nih.gov+6

35.   Anand L, Rodriguez Lopez CM. 2022. ChromoMap: an R package for interactive

visualization of multi omics data and annotation of chromosomes. BMC Bioinformatics

23:33. doi:10.1186/s12859-021-04516-6

36.   Ruta M, Angielczyk KD, Fröbisch J, Benton MJ. 2013. Decoupling of

morphological disparity and taxic diversity during the adaptive radiation of anomodont

therapsids. Proc R Soc B 280:20131071. doi:10.1098/rspb.2013.1071

37.   Schartl M, Schmid M, Nanda I. 2016. Dynamics of vertebrate sex chromosome

evolution: from equal size to giants and dwarfs. Chromosoma 125:553–571.

doi:10.1007/s00412-015-0569-y

38.   Münkemüller T, et al. 2012. How to measure and test phylogenetic signal.

Methods Ecol Evol 3:743–756. doi:10.1111/j.2041-210X.2012.00196.x

39.   Bivand R, Pebesma E, Gómez Rubio V. 2013. Applied Spatial Data Analysis with

R, 2nd ed. New York: Springer.

40.   Cummins M, Watson C, Edwards RJ, Mattick JS. 2024. The evolution of

ultraconserved elements in vertebrates. Mol Biol Evol 41:1724–1734.

doi:10.1093/molbev/msae146

41.   Dorrity MW, Saunders LM, Queitsch C, Fields S, Trapnell C. 2020. Dimensionality

reduction by UMAP to visualize physical and genetic interactions. Nat Commun

11:1537. doi:10.1038/s41467-020-15351-4

doaj.org+9nature.com+9pubmed.ncbi.nlm.nih.gov+9

42.   Liu D, Hunt M, Tsai IJ. 2018. Inferring synteny between genome assemblies: a

systematic evaluation. BMC Bioinformatics 19:26. doi:10.1186/s12859-018-2008-1

43.   Garrison E, Sirén J, Novak AM, Hickey G, Eizenga JM, Paten B, et al. 2024. Building

pangenome graphs. Nat Methods 21:1–5. doi:10.1038/s41592-024-01450-1

44.   Dubois F, Sidiropoulos N, Weischenfeldt J, Beroukhim R. 2022. Structural

variations in cancer and the 3D genome. Nat Rev Cancer 22:533–546.

doi:10.1038/s41571-022-00698-0

45.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update:

Novel and streamlined workflows along with broader and deeper phylogenetic coverage

for scoring of eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol 38:4647–4654.

doi:10.1093/molbev/msab199

46.   Oluwadare O, Highsmith M, Cheng J. 2019. An overview of methods for

reconstructing 3 D chromosome and genome structures from Hi C data. Biol Proced

Online 21:7. doi:10.1186/s12575-019-0095-0

47.   Lappala A, DeMaere MZ, Dansereau C, Villani TS, Lieberman Aiden E, Dekker J, et

al. 2021. Four dimensional chromosome reconstruction elucidates the spatiotemporal

reorganization of the mammalian X chromosome. Proc Natl Acad Sci U S A

118:e2107092118. doi:10.1073/pnas.2107092118

48.   Percival Alwyn L, Smith J, Patel K, Ahmed A, Zhang Y, Vickers M, et al. 2024.

UKCropDiversity HPC: A collaborative high performance computing resource approach

for sustainable agriculture and biodiversity conservation. Plants People Planet.

doi:10.1002/ppp3.10607

49.   Sandelin A, Klasson L, Medstrand P, Engström U, Fredman D, Lenhard B. 2004.

Arrays of ultraconserved non coding regions span the loci of key developmental genes

in vertebrate genomes. BMC Genomics 5:99. doi:10.1186/1471-2164-5-99

50.   Leclair NK, Anczuków O. 2022. 'Poisoning' of the transcriptome by

ultraconserved elements. Nat Rev Mol Cell Biol 23:777. doi:10.1038/s41580-022-

00527-1 europepmc.org+2pubmed.ncbi.nlm.nih.gov+2researchgate.net+2

51.   Jungers WL, Falsetti AB, Wall CE. 1990. Shape, relative size, and size-

adjustments in morphometrics. J Hum Evol 19:333–348. doi:10.1016/0047-

2484(90)90043-W

52.   Wickham H. 2007. Reshaping data with the reshape package. J Stat Softw 21:1–

20. Available from: http://www.jstatsoft.org/v21/i12/

53.   Müller P, Eddelbuettel D, Bates D. 2025. matrixStats: Functions for Highly

Optimized Computing of Matrix Statistics. Available from: https://cran.r-

project.org/web/packages/matrixStats/matrixStats.pdf

54.   Wickham H, Vaughan D, Girlich M. 2024. tidyr: Tidy messy data. R package

version 1.3.1. Available from: https://github.com/tidyverse/tidyr,

https://tidyr.tidyverse.org

55.   Wickham H, François R, Henry L, Müller K, Vaughan D. 2023. dplyr: A grammar of

data manipulation. R package version 1.1.4. Available from:

https://github.com/tidyverse/dplyr

56.   Wickham H. 2016. ggplot2: Elegant graphics for data analysis. New York:

Springer-Verlag. Available from: https://ggplot2.tidyverse.org

57.   Danecek P, Bonfield JK, Liddle J, Marshall J, et al. 2021. Twelve years of SAMtools

and BCFtools. GigaScience 10:giab008. doi:10.1093/gigascience/giab008

58.   Wilke C. 2024. cowplot: Streamlined plot theme and plot annotations for

'ggplot2'. R package version. Available from: https://cran.r-project.org/package=cowplot

59.   Foley NM, et al. 2023. A genomic timescale for placental mammal evolution.

Science 380:1239–1245. doi:10.1126/science.abl8189

60.   Martin FJ, et al. 2023. Ensembl 2023. Nucleic Acids Res 51:D933–D941.

doi:10.1093/nar/gkac1040

61.   Yue JX, Liti G. 2019. simuG: a general-purpose genome simulator. Bioinformatics

35:4442–4444. doi:10.1093/bioinformatics/btz366

62.   Uyeda JC, Caetano DS, Pennell MW. 2015. Comparative analysis of principal

components can be misleading. Syst Biol 64:677–689. doi:10.1093/sysbio/syv019

63.   Kumar S, Stecher G, Suleski M, Hedges SB. 2022. TimeTree 5: an expanded

resource for species divergence times. Mol Biol Evol 39:msac174.

doi:10.1093/molbev/msac174

64.   Blomberg SP, Garland T Jr, Ives AR. 2003. Testing for phylogenetic signal in

comparative data: behavioral traits are more labile. Evolution 57:717–745.

doi:10.1111/j.0014-3820.2003.tb00285.x

65.   Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature

401:877–884. doi:10.1038/44766

66.   Revell LJ. 2024. phytools 2.0: an updated R ecosystem for phylogenetic

comparative methods (and other things). PeerJ 12:e16505. doi:10.7717/peerj.16505

67. Canavez FC, Luche DD, Stothard P, Leite KR, Sousa-Canavez JM, Plastow G, et al.

2012. Genome sequence and assembly of *Bos indicus*. J Hered 103:342–348.

doi:10.1093/jhered/esr150

68. Rosen BD, Bickhart DM, Koren S, Schnabel RD, Hall R, Zimin A, et al. 2020. De

novo assembly of the cattle reference genome with single-molecule sequencing.

Gigascience 9:giaa021. doi:10.1093/gigascience/giaa021

69. Ananthasayanam S, Kothandaraman H, Nayee N, Saha S, Baghel DS,

Gopalakrishnan K, et al. 2019. First near complete haplotype phased genome assembly

of River buffalo (*Bubalus bubalis*). bioRxiv 618785. doi:10.1101/618785

70. Dong Y, Zhang X, Xie M, Arefnezhad B, Wang Z, Wang W, et al. 2015. Reference

genome of wild goat (*Capra aegagrus*) and sequencing of goat breeds provide insight

into genic basis of goat domestication. BMC Genomics 16:1–11. doi:10.1186/s12864-

015-1925-6

71. Bickhart DM, Rosen BD, Koren S, Sayre BL, Hastie AR, Chan S, et al. 2017. Single-

molecule sequencing and chromatin conformation capture enable de novo reference

assembly of the domestic goat genome. Nat Genet 49:643–650. doi:10.1038/ng.3802

72. Farré M, Li Q, Darolti I, Zhou Y, Damas J, Proskuryakova AA, et al. 2019. An

integrated chromosome-scale genome assembly of the Masai giraffe (*Giraffa*

camelopardalis tippelskirchi). GigaScience 8:giz090. doi:10.1093/gigascience/giz090

73. Davenport KM, Bickhart DM, Worley K, Murali SC, Salavati M, Clark EL, et al.

2022. An improved ovine reference genome assembly to facilitate in-depth functional

annotation of the sheep genome. Gigascience 11:giab096.

doi:10.1093/gigascience/giab096

74.   Bredemeyer KR, Harris AJ, Li G, Zhao L, Foley NM, Roelke-Parker M, et al. 2021.

Ultracontinuous single haplotype genome assemblies for the domestic cat (*Felis catus*)

and Asian leopard cat (*Prionailurus bengalensis*). J Hered 112:165–173.

doi:10.1093/jhered/esaa047

75.   Mohr DW, Naguib A, Weisenfeld NI, Kumar V, Shah P, Church DM, et al. 2017.

Improved de novo genome assembly: linked-read sequencing combined with optical

mapping produce a high-quality mammalian genome at relatively low cost. bioRxiv

128348. doi:10.1101/128348

76.   Bredemeyer KR, Murphy WJ. 2021. Unpublished Liger assemblies. Veterinary

Integrative Biosciences, Texas A&M University, College Station, TX. Submitted 11 Jan

2021.

77.   Wang G, Brändl B, Rohrandt C, Hong K, Pang A, Lee J, et al. 2021. Chromosome-

level genome assembly of the functionally extinct northern white rhinoceros

(*Ceratotherium simum cottoni*). bioRxiv. doi:10.1101/2021.12.01.470802

78.   Wang C, Li H, Guo Y, Huang J, Sun Y, Min J, et al. 2020. Donkey genomes provide

new insights into domestication and selection for coat color. Nat Commun 11:6014.

doi:10.1038/s41467-020-19702-6

79.   Kalbfleisch TS, Rice ES, DePriest MS Jr, Walenz BP, Hestand MS, Vermeesch JR,

et al. 2018. Improved reference genome for the domestic horse increases assembly

contiguity and composition. Commun Biol 1:197. doi:10.1038/s42003-018-0209-z

80.   Gordon D, Murali SC, Ventura M, Mercuri L, Eichler EE. De novo PacBio assembly

of a female gorilla (Kamilah). Genome Sciences, University of Washington, Seattle, WA,

USA. Unpublished.

81.   Jayakumar V, Nishimura O, Kadota M, Hirose N, Sano H, Murakawa Y, et al. 2021.

Chromosomal-scale de novo genome assemblies of cynomolgus macaque and

common marmoset. Sci Data 8:159. doi:10.1038/s41597-021-00911-0

82.   Dutcher S, Fulton R, Lindsay T. The rhesus macaque (*Macaca mulatta*) genome

assembly. The McDonnell Genome Institute, Washington University School of Medicine;

University of Washington, Seattle, WA. Unpublished.

83.   Kronenberg ZN, Fiddes IT, Gordon D, Murali S, Cantsilieris S, Meyerson OS, et al.

2018. High-resolution comparative analysis of great ape genomes. Science

360:eaar6343. doi:10.1126/science.aar6343

84.   Batra SS, Levy-Sakin M, Robinson J, Guillory J, Durinck S, Vilgalys TP, et al. 2020.

Accurate assembly of the olive baboon (*Papio anubis*) genome using long-read and Hi-C

data. Gigascience 9:giaa134. doi:10.1093/gigascience/giaa134

85.   Rupp O, MacDonald ML, Li S, Dhiman H, Polson S, Griep S, et al. 2018. A

reference genome of the Chinese hamster based on a hybrid assembly strategy.

Biotechnol Bioeng 115:2087–2100. doi:10.1002/bit.26809

86.   Roller M, Navarro FCP, Fiddes I, Streeter I, Feig C, Martin-Galvez D, et al. Repeat

associated mechanisms of genome evolution and function revealed by the *Mus caroli*

and *Mus pahari* genomes. Available from: http://bura.brunel.ac.uk/handle/2438/16068

87.   Church DM, Goodstadt L, Hillier LW, Zody MC, Goldstein S, She X, et al. 2009.

Lineage-specific biology revealed by a finished genome assembly of the mouse. PLoS

Biol 7:e1000112. doi:10.1371/journal.pbio.1000112

88.   Lilue J, Doran AG, Fiddes IT, Abrudan M, Armstrong J, Bennett R, et al. 2018.

Sixteen diverse laboratory mouse reference genomes define strain-specific haplotypes

and novel functional loci. Nat Genet 50:1574–1583. doi:10.1038/s41588-018-0223-8

89.   Lassance JM, Hoekstra HE. Improved assembly of the deer mouse (Peromyscus

maniculatus) genome. Organismic and Evolutionary Biology / Molecular and Cellular

Biology, Harvard University / Howard Hughes Medical Institute, Cambridge, MA, USA.

Unpublished.

90.   Howe K, Dwinell M, Shimoyama M, Corton C, Betteridge E, Dove A, et al. 2021.

The genome sequence of the Norway rat, *Rattus norvegicus* Berkenhout 1769.

Wellcome Open Res 6:60. doi:10.12688/wellcomeopenres.17242.1

91.   Sievert C. 2020. Interactive web-based data visualization with R, plotly, and

shiny. Chapman and Hall/CRC. ISBN 9781138331457. Available from: https://plotly-

r.com

1007

1008

1009

1010

1011

1012 **Supplementary Information**

1013

1014 **Figure S1.** Number of ultraconserved elements (UCEs) mapped utilizing two different mapping

1015 software: minimap2 (blue) and bwa-mem (grey). Demonstrated using a subset of chromosome

1016 assemblies picked from the 'chromosome set 2' of the placental mammals UCE dataset discussed in

1017 the main text.

1018

1019

Figure S2. Principal component (PC) sensitivity analyses of 53 X-linked BUSCO landmarks in amniote vertebrates taken from Lovell et al. (2022). The first row are PCAs conducted using raw, absolute positions for the original direction (A) and reverse complement (B), respectively. The second row are PCAs conducted using bounded, absolute positions for the original direction (C) and the reverse complement (D). The third row are PCAs conducted using size-correction with chromosome size for the original direction (E) and the reverse complement (F). The fourth row of plots are PCAs conducted using size-correction using bounded chromosome sizes and positions for the original direction (G) and reverse complement (H). The chicken (*Gallus gallus*) is highlighted in plot B to illustrate an analytical artifact associated with clustered landmarks on large chromosomes. Organism silhouettes downloaded from https://www.phylopic.org/.

1030
1031   Figure S3. Jaccard similarity coefficients for within and between protein-coding gene content in the

1032   two focal chromosome sets used in the study.

1033

Figure S4. Principal component analyses (PCAs) of ultraconserved element (UCE) landmarks in
placental mammals for chromosome set 1 before and after the removal of an outlier landmark (UCE
6948). The first row of plots are PCAs conducted using raw, absolute positions for the original
direction (A) and reverse complement (B), respectively. The second row of plots are PCAs conducted
using bounded, absolute positions for the original direction (C) and the reverse complement (D). The
third row of plots are PCAs conducted using size-correction with chromosome size for the original
direction (E) and the reverse complement (F). The fourth row of plots are PCAs conducted using size-
correction using bounded chromosome sizes and positions for the original direction (G) and reverse
complement (H). The first column of plots are PCAs conducted with all 186 UCE landmarks and the
second column of plots are PCAs based on 185 UCE landmarks (outlier removed).

1045

1046

1047

1048

Figure S5. Principal component analyses (PCAs) of ultraconserved element (UCE) landmarks in

placental mammals for chromosome set 2 before and after the removal of an outlier landmark (UCE

6948). The first row of plots are PCAs conducted using raw, absolute positions for the original

direction (A) and reverse complement (B), respectively. The second row of plots are PCAs conducted

using bounded, absolute positions for the original direction (C) and the reverse complement (D). The

third row of plots are PCAs conducted using size-correction with chromosome size for the original

direction (E) and the reverse complement (F). The fourth row of plots are PCAs conducted using size-

correction using bounded chromosome sizes and positions for the original direction (G) and reverse

complement (H). The first column of plots are PCAs conducted with all 82 UCE landmarks and the

second column of plots are PCAs based on 81 UCE landmarks (outlier removed).

1059

1060

1061

1062

1063

Figure S6. ChromoMap plots visualizing principal component analysis (PCA) loadings of landmarks

on PC1, PC2, and PC3 in the raw landmark PCA, bounded landmark PCA, and size corrected

landmark PCA of chromosome set 1 (A) and set 2 (B) from the placental mammal visualized using

links between *Bos indicus* and *Equus asinus*. Respective variances explained by each principal

component are listed in the plot. A single landmark (UCE 6948) was found to be uniquely weighed on

the various PCs identifying it as an outlier landmark.

1070

1071

1072

Figure S7. Principal component (PC) sensitivity analyses of outlier removed UCE landmarks in

placental mammals chromosome set 1. The first row are PCAs conducted using raw, absolute

positions for the original direction (A) and reverse complement (B), respectively. The second row are

PCAs conducted using bounded, absolute positions for the original direction (C) and the reverse

complement (D). The third row are PCAs conducted using size-correction with chromosome size for

the original direction (E) and the reverse complement (F). The fourth row are PCAs conducted using

size-correction using bounded chromosome sizes and positions for the original direction (G) and

reverse complement (H).

1081

1082

1083

Figure S8. Principal component (PC) sensitivity analyses of outlier removed UCE landmarks in

placental mammals chromosome set 2. The first row are PCAs conducted using raw, absolute

positions for the original direction (A) and reverse complement (B), respectively. The second row are

PCAs conducted using bounded, absolute positions for the original direction (C) and the reverse

complement (D). The third row are PCAs conducted using size-correction with chromosome size for

the original direction (E) and the reverse complement (F). The fourth row are PCAs conducted using

size-correction using bounded chromosome sizes and positions for the original direction (G) and

reverse complement (H).

1092

1093

Figure S9. Ordination results from principal component (PC) analysis of UCE landmark positions in

two sets of chromosomes from 26 species of placental mammals using bounded raw positions after

the removal of an outlier landmark (see text). Representatives for five orders (Artiodactyla, Carnivora,

Perissodactyla, Primates and Rodents) were included (Table1). The depicted structural variation is

inferred from PCs 3 and 4 of 185 (chromosome set 1, A) and 81 (chromosome set 2, B) landmarks,

respectively. In both datasets, the first two PCs explained > 99% of the variation (see **Figure 5**, main

text). While explaining < 1% of the overall variation, PCs 3 and 4 mostly separated mammalian

orders in geno-metric space. *Carpa aegragus* was a notable outlier in chromosome set 1.

1102

1103
1104
1105 **Figure S10.** Boxplots comparing increase in disparity (ID) between indel and inversion simulations for

1106 principal component (PC)1 (A) and PC2 (B), respectively. Note that inversions cause significantly

1107 greater ID than indels.

1108

1109
1110
1111   **Figure S11.** Jaccard similarity coefficients for within and between protein-coding gene content in

1112   mammalian orders used in the study.

1113

1114
1115
Figure S12. Schematic diagram illustrating sensitivities of disparity analysis and synteny analysis to

different landmark scenarios. Scenarios are described from top to bottom of the figure. First, both

analytical approaches will report highly conserved landmark/ortholog order, indicating when two

species have highly conserved genomic architecture. Second, both approaches will report large

inversions among landmark positions. Third, depending on parameter settings syntenic analysis may

ignore small landmark rearrangements whereas disparity analysis captures this variation. Finally,

synteny analysis ignores spatial variation in landmark placement and 'synteny' can be broken if

chromosomal gaps are large enough whereas disparity analysis captures this variation in landmark

placement.

Appendix S1. List of 53 BUSCO genes identified as landmarks for the human X-linked study of

amniote vertebrates.

1127

1128   eda, dlg3, snx12, gjb1, zmym3, nono, shroom4, amot, lrch2, pls3, plp1, gla, btk, drp2, sytl4, srpx2,

1129   tspan6, dach2, apool, rps6ka6, lpar4, magt1, fgf16, zdhhc15, uprt, nsdhl, vma21, cd99l2, mtmr1,

1130   mtm1, aff2, fmr1, slitrk4, atp11c, cd40lg, vgll1, htatsf1, gpc3, rap2c, frmd7, bcorl1, utp14a, smarca1,

1131   stag2, cul4b, lamp2, tmem255a,