Chronic pancreatitis patient-derived organoids reveal new paths to precision therapeutics

Victoria Osorio-Vasquez[1], Jonathan Zhu[1], Jan C. Lumibao[1], Kathryn Lande[1,2], Kristina L. Peck[1], McKenna K. Stamp[1], Shira R. Okhovat[1], Hyemin Song[1,3], Satoshi Ogawa[1], Casie Kubota[1], Vasiliki Pantazopoulou[1], Yang Dai[1], Angelica Rock[1], Chelsea Bottomley[1], Ethan Thomas[1], Jasper Hsu[1,3], Araceli Herrera Morales[1], Alexandra Fowler[1], T'Onj McGriff[1], K. Garrett Evensen[1], Siri Larsen[4], Muhamad Abdulla[4], Phil Greer[5], Jessica Gibson[5], Michael Downes[1], Ronald Evans[1], Andrew M. Lowy[6], David C. Whitcomb[5,7], Jingjing Zou[8], Alfredo Molinolo[9], Tae Gyu Oh[10], Rebekah White[6], Melena Bellin[4,11], Herve Tiriac[6], Dannielle D. Engle[1,3,*]

[1]Salk Institute for Biological Studies, La Jolla, CA, USA.
[2]Sanford Burnham Prebys Medical Discovery Institute, La Jolla, CA, USA
[3]School of Biological Sciences Department of Molecular Biology, University of California San Diego, School of Medicine, La Jolla, CA, USA
[4]Schulze Diabetes Institute, Department of Surgery, University of Minnesota Medical School, Minneapolis, MN, USA.
[5]Ariel Precision Medicine, Pittsburgh, PA, USA.
[6]Department of Surgery, Division of Surgical Oncology, Moores Cancer Center, University of California San Diego, San Diego, CA, USA.
[7]University of Pittsburgh, Pittsburgh, PA, USA.
[8]Herbert Wertheim School of Public Health and Human Longevity Science, University of California San Diego, CA, USA.
[9]Department of Pathology, Biorepository and Tissue Technology Shared Resource, Moores Cancer Center, UC San Diego, San Diego, CA, USA.
[10]University of Oklahoma, College of Medicine, Oklahoma City, OK, USA.
[11]Departments of Pediatrics, Division of Pediatric Endocrinology, University of Minnesota Medical School Minneapolis, MN, USA.

*Correspondence: engle@salk.edu

Summary

Chronic pancreatitis (CP) affects ~3 million people worldwide, yet altering the course of disease is challenging. We developed a patient-derived organoid (PDO) platform to investigate the molecular pathogenesis of this disease and identify therapeutic strategies. We generated 36 PDOs from patients with idiopathic, hereditary, and alcohol-related CP with high genetic concordance. PDOs retained inflammation-associated transcriptional and proteomic features. Transcriptomic profiling revealed three molecular subtypes of CP independent of etiology. We discovered widespread dysfunction of the cystic fibrosis transmembrane conductance regulator (CFTR) in half of the CP PDOs, including those with wildtype CFTR. Clinically available CFTR modulators stabilized mutant or wildtype CFTR, restored CFTR function, and decreased mitogenic and inflammatory signaling. This work provides the first comprehensive PDO platform for modeling CP. We demonstrate the utility of this platform for precision therapeutic investigations. Our findings reveal CFTR modulators as a broadly applicable and effective therapeutic strategy.

Keywords: chronic pancreatitis, patient derived organoids, CFTR, molecular subtypes, CFTR correctors, CFTR potentiators

Introduction

Pancreatitis, or pancreatic inflammation, is a continuum of diseases that may progress from acute to chronic stages. More than 300,000 Americans and 3 million individuals globally have chronic pancreatitis (CP)[1-3]. CP is characterized by atrophy, fibrosis, pancreatic exocrine insufficiency, diabetes mellitus, and intractable pain. Current treatment is largely supportive and often fails to modify disease progression, sometimes requiring surgical interventions such as partial or total pancreatectomy for pain management. CP may arise from numerous environmental and genetic factors. It is classified using the Toxic-metabolic, Idiopathic, Genetic, Autoimmune, Recurrent and severe acute pancreatitis and Obstructive (TIGAR-O v2) framework[4]. The most common identifiable causes of CP include tobacco use (46-60%)[5,6], gallstones (14-37%)[7,8], alcohol (7-34%)[5,9], and genetics (10%)[5], though a striking 28-80% of cases remain idiopathic[10]. Hereditary CP is associated with pathogenic germline mutations in protease serine 1 (*PRSS1*) (68-80%)[11-14], serine protease inhibitor kazal type 1 (*SPINK1*) (1-8.8%)[10-13], cystic fibrosis transmembrane conductance regulator (*CFTR*) (16.5%)[13], and other genes.

Pancreatitis initiation can be traced to dysfunction in both acinar and ductal cells. *PRSS1* and *SPINK1* mutations lead to hyperactivation of digestive enzymes, leading to acinar cell injury and inflammation[15-20]. Mutations in *CFTR* disrupt bicarbonate secretion in the pancreas. Ductal cells normally secrete 1-2L of 140mM sodium bicarbonate per day[21], which is essential for mucin production, flushing digestive enzymes into the duodenum, preventing premature digestive enzyme activation, and neutralizing acidic chyme from the stomach[21-23]. Unlike cystic fibrosis, CFTR variants associated with hereditary CP are often heterozygous[24]. Up to 37% of hereditary CP patients harbor *CFTR* mutations[25-32]. Importantly, 50% of idiopathic CP patients harbor *CFTR* alterations when extensive genotyping is performed[21,25-34]. Many *CFTR* variants associated with hereditary CP selectively impair bicarbonate, but not chloride transductance. Altogether, CFTR dysfunction in the pancreas, distinct from classical cystic fibrosis, may underlie a significant proportion of CP and represents an underexplored therapeutic target.

With more than 2,000 mutations annotated for *CFTR*, the response of most *CFTR* variants to modulators are unknown. Of those with response information, this testing has been primarily performed for cystic fibrosis-related variants in non-pancreas relevant contexts. Further, cystic fibrosis patient responses to CFTR therapies are highly variable due to undefined mechanisms[35-40]. Limited information is available regarding CFTR modulator response in the context of hereditary pancreatitis associated *CFTR* mutations. Delineating the genotypic and phenotypic relationships of pancreatitis-associated CFTR alterations with modulator response is essential. But these investigations have been hampered by the limited availability of renewable human models. Unfortunately, despite sharing 78% homology, mouse CFTR does not respond to human CFTR modulators, complicating mechanistic investigations in mouse models[41]. Further, mouse bicarbonate secretion is notably different than humans, reaching only 50mM in pancreatic secretions[37], limiting our ability to make conclusions relevant to human physiology.

Three-dimensional patient-derived organoids (PDOs) have emerged as transformative models for studying pancreas biology. PDOs can be generated from normal human pancreas and pancreatic ductal adenocarcinoma with high efficiency [42,43], far surpassing earlier monolayer or iPSC-based platforms[44]. PDOs are expandable, cryopreservable, and genetically stable, enabling in-depth molecular and functional analyses[43,45-50]. In pancreatic ductal adenocarcinoma (PDA), PDOs faithfully recapitulate transcriptomic and genetic subtypes, predict treatment response, and inform mechanisms of resistance[43,45-50]. Despite lacking a microenvironment, PDOs retain remarkable predictive power due to the central role of epithelial-intrinsic signaling in disease behavior[51,52]. PDO models of CP remain rare. A pediatric cohort (n=9) has been described[53], but generation efficiency, molecular fidelity, and translational potential remain unknown. To address this gap, we created a comprehensive biobank of CP PDOs encompassing all major etiologies and benchmarked their ability to model disease features and guide treatment strategies.

We report a high PDO generation success rate (70.5%, *n*=36) and strong concordance with

matched primary tissues at the DNA, RNA, and protein levels. Genetic analyses of 24 PDOs
identified germline pathogenic variants in 2 out of 8 idiopathic cases. In addition, rare somatic
variant analyses identified 2 CP PDOs with alterations associated with pancreatic ductal
adenocarcinoma, including *KRAS* and *TP53*. These oncogenic alterations were not detectable in
primary samples in which low cellularity limits the ability to detect somatic mutations in the
epithelial compartment. CP PDOs exhibit elevated fibroinflammatory gene and protein expression
profiles despite being removed from the pancreatic microenvironment, suggesting that ductal
epithelial cell intrinsic features contribute to inflammation. Strikingly, half of the PDOs tested
displayed CFTR functional impairment despite 2 out of the 4 dysfunctional lines harboring wildtype
CFTR. Treatment with clinically available CFTR modulators restored ductal function and
suppressed pro-inflammatory and mitogenic signaling, suggesting that CFTR modulation may
benefit a broad segment of CP patients. Together, this study presents a robust and scalable
platform for dissecting CP pathophysiology and accelerating precision medicine efforts.
116
117

Methods

Human Specimens

Normal pancreata were obtained from the Lifesharing organ donation program. The average age of the hNP PDOs is 56 years of age. Pancreatitis tissues were obtained from the University of Minnesota (UMN) and University of California San Diego (UCSD) Biorepository and Tissue Technology Shared Resource (BTTSR). De-identified pancreatitis tissues were obtained from the University of Minnesota (UMN) or the UCSD BTTSR with approval from their respective Institutional Review Boards. Written and informed consent or parental assent, as age appropriate, was obtained prior to acquisition of tissue from all patients. Age and/or age ranges will not be provided for pediatric patients as the rarity of the procedure in this patient population could itself be identifying. Specimens were obtained from patients with chronic pancreatitis at UCSD undergoing either partial pancreatectomy (distal pancreatectomy or pancreaticoduodenectomy) or pancreatic head debridement during pancreatic drainage procedures for symptomatic relief. Excess tissue was collected fresh into cold media for organoid generation. When available, pancreatic juice was collected in parallel by needle aspiration of pancreatic duct during drainage procedures. Specimens were obtained from patients undergoing total pancreatectomy and islet auto-transplantation at UMN. During pancreatectomy, the blood supply was preserved to the pancreas as long as possible to avoid prolonged ischemia. The entire pancreas, spleen, and duodenum were removed, and the gastrointestinal and biliary continuity is restored by a Roux-en-Y approach. Islet isolation was performed by infusing the pancreas with collagenase and neutral protease for enzymatic digestion, followed by mechanical disruption using the semi-automated method of Ricordi. The islets were then transported back to the operating room where they were infused via the portal vein on the same day as the pancreatectomy procedure. Residual exocrine pancreas tissue following islet isolation was collected for tissue research. Tissue specimens were catalogued, imaged, and processed (i.e. histology, organoid generation, flash frozen, and viably cryopreserved).

Histology and immunohistochemistry

Tissues were fixed in 10% neutral buffered formalin for 48 hours. Tissues were embedded in paraffin and stained with hematoxylin and eosin or Masson's trichrome by the Tissue Technology Shared Resource at UCSD supported by the National Cancer Institute Cancer Center Support Grant (CCSG Grant P30CA23100). The following primary antibodies were used for immunohistochemical staining Ki-67 (8D5, Cell Signaling Technologies# 9449), CK19 (Troma-III, Developmental studies hybridoma bank), CFTR (M3A7, Thermo Fischer #MA5-11768), Cancer Antigen 19-9 (MyBioSource #MBS850562). Secondary antibodies used were ImmPRESS® HRP Horse Anti-Mouse IgG Polymer Detection Kit (MP-7402) or ImmPRESS® HRP Goat Anti-Rat IgG Polymer Detection Kit (MP-7404) with ImmPACT® DAB Substrate Kit (HRP) (SK-4105). Staining intensity was semi quantitatively scored as: low, medium, or high using QuPath 0.5.0[54] and ranking percent positive staining. Slides were imaged using the Olympus VS-120 virtual slide scanning microscope, Olympus BX43 Microscope, or Zeiss Axioscan 7. Zen lite or CellSens was used to visualize and annotate images.

Pathologist Review of Patient Primary Tissue Specimen
Primary tissue specimen stained with hematoxylin and eosin or Masson's trichrome were scanned (Aperio-Leica AT2 Scanner, 40x) analyzed and imaged. Severity of fibrosis, acinar atrophy, inflammation and ductal changes were categorized as mild, moderate, or severe as denoted by dilation of ducts, expansion of fibroblast compartment, collagen deposition, and infiltration of

168 immune cells[55]. Alterations in histology due to autolytic changes or signs of cautery from an
169 electro-scalpel.
170
171 *Organoid generation and culture*
172
Prior to processing, visible fat, mesentery, and calcifications were removed. Patient specimens
were minced and aliquoted for viable tissue cryopreservation for potential regeneration of PDOs,
flash frozen for primary tissue DNA extraction, and further dissociation for PDO generation. For
organoid generation, tissue was minced and incubated in digestion media (3000 U/mL
Collagenase XI + 1000 U/mL hyaluronidase- Stem Cell #07919 in DMEM/F12 diluted 1:4 in
Advanced DMEM (Thermo Fischer #12634010) with 1% (v/v) glutamax (Thermo Fischer
#35050061), 1% (v/v) penicillin/streptomycin (Life Technologies #15140122), and 1%(v/v)
HEPES (1M, pH 7.5)) at 35°C while rotating. If cell lysis became apparent, DNAse I (10 μg/mL,
Roche #1128493200) was added. Fractions were collected every 20 minutes for up to 1 hour by
allowing large tissue pieces to sediment before supernatant was collected. Dissociation of the
remaining large tissue fragments continued by replacing the digestion media. Supernatant was
centrifuged at 300x RCF at 4°C for 5 minutes. The first fraction cell pellet collected was
resuspended in 1 mL of HBSS (Life Technologies #14175095) with 2% FBS and placed on ice
while collecting additional fractions from the digested tissue. After 3 fractions were collected, the
supernatant fractions were washed with Advanced DMEM with 1% (v/v) glutamax, 1% (v/v)
penicillin/ streptomycin, and 1% (v/v) HEPES prior to resuspension in Matrigel for plating. If larger
tissue pieces remained at the end of dissociation, they were either viably cryopreserved
(CrySOfree- Sigma-Aldrich #C9249) or plated in Matrigel (Corning #356231) in separate wells.
Cells were plated in Corning Matrigel from lots 28323001, 2011001, 0287001, and 1062004
during growth, expansion and experiments. Following Matrigel solidification, complete feeding
media was applied to each well.
194
Murine normal pancreas organoids were cultured in medium previously described[43,50]. Human
normal pancreas and human pancreatitis were cultured in human complete feeding medium.
Human complete feeding medium was prepared using Advanced DMEM/F12 containing HEPES
(1 % (v/v), 1M, pH 7.5), glutamax 1 % (v/v), B27 supplement (1x ,Life Technologies #A3582801),
and 1% (v/v) penicillin/streptomycin; N-acetylcysteine (1.25 mM, Sigma-Aldrich #A9165),
nicotinamide (10 mM, Thermo Fischer #125271000); hEGF (50 ng/mL, R&D Systems #236-EG),
FGF-10 (100 ng/mL, R&D Systems #345-FG), gastrin (10 nM, R&D #3006), and A83-01 (500 nM,
Tocris Bioscience #2939); rNoggin (100 ug/mL, Qkine LTD #Qk034) and rRSPO1 (10nM Qkine
LTD # Qk006); Wnt surrogate (300 pM, ImmunoPrecise Antibodies #N001); PGE2 (1 μM, Tocris
Bioscience #2296) and Y-27632 dihydrocholoride (10.5 μM, Tocris Bioscience #1254). Mouse
organoid media lacked Wnt3a, PGE2, and Y-27632 and utilized 10% RSPO1 conditioned media
and 10% Noggin conditioned media. Organoid nomenclature is defined as: hNP, human normal
pancreas; hCP, human chronic pancreatitis, mNP, mouse normal pancreas. All organoid models
were routinely tested for *mycoplasma* at Salk Institute.
209
210 *Whole mount immunofluorescence and EdU incorporation*
211
Organoids were cultured as described above in μ-slide (Ibidi #80826). Reagents for Click-iT Plus
EdU Cell Proliferation Kit for Imaging (Thermo Fischer #C10639) were used in accordance with
manufacturer's protocol. Organoids were treated with EdU for 25 hours by removing half of the
media and adding EdU containing media (20 μM). Matrigel domes containing organoids were
washed using PBS prior to fixing with 4% PFA (v/v) at 37°C for 10 minutes. After fixation rinsed
with excess 3% BSA and removed wash solution. Permeabilized with 0.5% Triton X-100 in PBS
for 10 minutes at room temperature. Steps subsequent to Click-it reaction cocktail addition were

protected from light. Remaining organoids were washed with 3% BSA in PBS and blocked with
IF buffer (0.1% Bovine serum albumin, 0.2% goat serum, 0.2% triton x-100, 0.05% Tween
20:sterile filtered + 10% Goat serum in PBS) for 1 hour. Organoids were incubated with a dilution
of 1:200 for CK-19 (Invitrogen #MA5-12663) overnight at room temperature. Organoids were
washed for 1 hour prior to incubation with secondary antibody (Jackson ImmunoResearch
Laboratories Inc. #115-545-003) and Hoechst for 2 hours at room temperature. Washed
organoids for at least 1 hour with IF buffer prior to mounting using Prolong Glass Antifade
Mountant (Thermo Fisher P36980) and dried overnight. Imaged using the Yokogawa CQ3000 at
20x. Nuclei and EdU positive nuclei were quantified using Cell Profiler (3.8.1)[56].
228
229 *Immunoblot*
230
Standard techniques were employed for immunoblotting (IB) of human organoids. Organoids
were isolated from Matrigel using cell recovery solution (10 mM HEPES 7.5 pH, 1x glutamax (v/v),
12.5 mM EDTA, 12.5 mM EGTA, 4.5 g/L dextrose in PBS w/o Ca and Mg) on ice with 0.5x
protease and phosphatase inhibitors (Thermo Fischer # PIA32961). After incubation on ice for 20
minutes, organoids were centrifuged at 0.8 x 1000 RCF at 4°C to pellet organoids and remove
cell recovery solution. Cell pellet was resuspended in TNET (1% (v/v) Triton X100, 150 mM NaCl,
5 mM EDTA, 50 mM Tris pH 7.5) with 1.0X protease and phosphatase inhibitors (Life
Technologies #A32961) and incubated on ice for 10 minutes before clarifying lysates for whole
cell lysate isolation. Protein quantification was carried out using BCA assay (Thermo Scientific
#23227). Protein lysates used to probe for CFTR were warmed to 37°C for 1 hour to prevent
insolubility, all other blots were performed using lysates heated to 95°C for 5 minutes. Protein
lysates were separated on a 4%-12% Bis-Tris NuPage gel (Invitrogen). Western blots were
probed with at a dilution of 1:1000 for CFTR (D6W6L, Cell Signaling Technologies #78335),
CA19-9 (MyBioSource #MBS850562), pEGFR (Y1068, Cell Signaling Technologies #3777),
EGFR (D38B1, Cell Signaling Technologies #4267), pSTAT3 (Y705, Cell Signaling Technologies
#D4XC3), STAT3 (D3Z2G, Cell Signaling Technologies #63585), pAKT (S473, Cell Signaling
Technologies #4060), AKT (11 E 7,Cell Signaling Technologies #4685), pERK (T202/Y204,Cell
Signaling Technologies #4370S), ERK2 (137F5, Cell Signaling Technologies #4695), CA2
(EPR5195,Abcam #ab124687), SPINK1 (CGQG032302A R&D #AF7496), and Cofilin (D3F9, Cell
Signaling Technologies #5175). Secondary antibodies used were Peroxidase AffiniPure™ Goat
Anti-Mouse IgG (H+L) (Jackson ImmunoResearch #115-035-003), Peroxidase AffiniPure™
Donkey Anti-Rabbit IgG (H+L) (Jackson ImmunoResearch #711-035-152), Peroxidase
AffiniPure™ Goat Anti-Rabbit IgG (H+L) (Jackson ImmunoResearch #111-035-003), Sheep IgG
Horseradish Peroxidase-conjugated Antibody (R&D Systems #HAF016), IRDye 800 CW Goat
Anti-Rabbit IgG (Licor Bio #926-32211) and IRDye 680 CW Goat Anti-Mouse IgG (Licor Bio #926-
68070).
257
258 hCP35 and hCP44 organoids at 50-70% confluency were treated with Ivacaftor (3 µM-
259 MedChemExpress #HY-13017), combination of Ivacaftor, Elexacaftor (3 µM -MedChemExpress
260 #HY 111772), and Tezacaftor (3 µM- Selleck Chemicals #S7059), or vehicle control for 48 hours
261 in human complete feeding media prior to whole cell lysate collection, as described above.
262
263 Western blots were developed using either a film processor (Mini-Med 90- AFP Manufacturing
264 Corporation) or Odyssey M (LiCor). Quantification of immunoblots was conducted using either
265 ImageJ or Empiria Studio.
266
267 *WGS library preparation, sequencing, variant calling, and primary tissue- organoid SNP*
268 *concordance*
269

DNA was isolated from PDOs after at least 3 passages. Isolation of primary tissue DNA was conducted from flash frozen or viably cryopreserved samples. Genomic DNA was isolated from organoids or primary tissue using DNeasy Blood and Tissue Kit (Qiagen #69504). WGS libraries were prepared using Nextera XT (Illumina DNA library) in accordance with the manufacturer's instructions. Final libraries were evaluated using TapeStation (Agilent). Libraries were sequenced on an Illumina NovaSeq X sequencer for 30 x coverage for primary tissues and PDO.

Raw data was aligned to the hg38 reference genome with bwa mem v0.7.12-r1039. PCR duplicates were marked with Picard v1.130, and germline variants were called with GATK's haplotypecaller, v4.0.8.1. SNPs and INDELs were filtered with vcftools v0.1.13 the parameters: max-missing=0.5, mac=2, min-meanDP=15, minQ=32. For the purposes of concordance testing, multi-allelic sites and sites with missing genotype calls were discarded.

Concordance was determined using R 'irr' package (version 0.84.1) using intraclass correlation coefficient (ICC) using a two-way model with a 0.95 confidence level. As a control, comparison of PDOs from two different patients and found a match rate of 72% with an ICC of 0.4. When comparing a PDO from one patient to the primary tissue of another, the match rate is 74% with an ICC of 0.43.

Germline pathogenic variants were investigated in: ABCG8, APOA5, APOB, CASR, CFTR, CLDN2, CTRC, GGT1, HLA-DQA1, HLA-DQB1, LPL, PRSS1, PRSS1-PRSS2, SLC10A2, SLC26A9 and SPINK1

RNA isolation and RNA-seq library construction

For RNA-seq experiment, organoids in Matrigel were lysed directly with 1 mL of TRIzol reagent (Thermo Fischer #15596018), and total RNA was extracted using RNeasy Mini Kit (Qiagen #74104). RNA quality control was performed on all samples using Qubit Fluorometer (Invitrogen) and TapeStation (Agilent) before RNA-seq analyses. A total of 200 ng of high-quality RNA (RIN > 8, assessed by Agilent TapeStation) was used for library construction. mRNA was enriched using poly(A) selection, followed by fragmentation. First- and second-strand cDNA synthesis was performed, and 3′ adenylation was carried out prior to ligation of Illumina-compatible adapters containing unique dual barcodes. Adapter-ligated fragments were enriched by limited-cycle PCR amplification. Final libraries were assessed for quality and fragment size distribution using the Agilent Bioanalyzer DNA 1000 kit, and equal amounts of libraries were pooled for sequencing on an Illumina platform. Novaseq 6000 sp reagent kit v 1.5 was used to sequence samples for 300 cycles for 150 bp paired end reads using an Illumina Novoseq 6000.

snRNA-seq analysis, RNA-seq analysis, PCA visualization, and differential gene expression analysis

Raw reads were quality checked with FASTQC (v0.12.1)[57]. Poor quality 3' ends and sequencing adapters were trimmed with Cutadapt (v4.9)[58] and reads shorter than 20nt were filtered. Trimmed reads were again checked with FASTQC, then aligned to human reference genome (hg38 from UCSC with gene annotation gencode.v45.basic.annotation.gtf)[59,60] and genes quantified with STAR[61] (2.7.11b). Subsequent analysis was performed in R (version shown below, R Core Team 2024).

The column with 2nd read strand aligned with RNA from the STAR generated expression tables were used since the library prep was strand specific. Expressions were normalized with DESeq2[62] and the top 500 most variable genes after variance stabilization were used for the PCA plot and

heatmap. DESeq2 was also used for differential gene analysis, with cutoffs padj <0.05 and |log$_2$FoldChange| >1. These values plotted in the volcano plot for every gene.

Nonnegative matrix factorization exclusively for hCP was performed with the NMF library[63,64]. Ranks 2 through 7 were tested, each with 500 runs to determine the optimal rank. Based on consensus throughout 500 runs, as well as the rank survey scores from the best run each rank, rank 3 was picked as the optimal rank. The consensus shows the best run from the 500 and which samples clustered together but basis was the parameter used to determine the most stable subtypes. Differential gene analysis was performed again, this time separated by basis found with NMF.

Normalized counts from DESeq2 were analyzed with GSEA comparing either hCP with hNAT and hNP, or between hCP bases to determine enriched gene sets[65,66]. GSEA was also used to find gene markers specific to ductal or acinar samples from BioProject PRJNA924346[67]. Each individual PDO transcriptome profile was projected onto the set of ductal or acinar gene markers by single sample GSEA to assess their similarity[68] .

Tumor PDO sample transcriptomes were obtained from NCBI dbGAP accession phs001611.v1.p1[43]. Differentially expressed genes with padj < 0.05 from tumor (hT) vs. normal, and hCP vs hNP and hNAT were integrated to find concordant or discordant expression patterns.

Heatmaps are generated with pheatmap or Graphpad Prism, all other figures are generated with ggplot2. R version and library versions are below.

EGAC00001001732 snRNA-seq data was analyzed[69], and the ductal cell cluster was interrogated. CP ductal cells contained 903 nuclei from 1 male and 1 female CP patient with unknown etiology, who are 71 and 52 years of age, respectively. NP ductal cells contained 4,432 nuclei from 3 NAT patients (2 female, 1 male, 46, 59, and 77 years of age respectively).

Fluorescent in situ hybridization

Sections were formalin fix and paraffin embedded. Reagents for the RNAscope Multiplex Fluorescent Reagent kit V2 (#323280) along with RNAscope 4-plex Ancillary Kit for Multiplex Fluorescent Reagent kit V2 (#323120) from Advanced Cell Diagnostics (Newark, CA, USA) were used in accordance with the manufacturer's instructions using the HybEZ Hybridization system (#241000). Probes were all designed by Advanced Cell Diagnostics: *Gpx3* (#470591-C3), *Aqp1* (#421831-C2), *Bpifb1* (#576511-C4). We used the manual target retrieval method described in appendix B of the protocol for 15 minutes with protease plus treatment for 15 minutes. Fluorescent dyes used TSA Vivid fluorophore 520 (323271- 1:3000), 570 (323272- 1:3000), 650 (323273-1:3000), and Opal 780 (# FP1501001KT- Akoya Biosciences; Opal TSA-DIG 1:3000; Opal Polaris 780 1:750). Sections were counterstained with DAPI, sealed with a coverslip using ProLong Gold Antifade reagent (Invitrogen #P36930), and then dried overnight covered from light. Imaged using either Fluoview FV 3000 confocal microscope at 30x or whole slide Zeiss Axioscan 7 at 20x.

Forskolin induced swelling assay

Organoids were seeded into 96-well clear bottom plates in 8 µL domes at a confluency between 50-70%. Organoids were fed human complete medium after seeding. 18-12 hours prior to treatments, media was changed and PGE$_2$ was removed from media. For dependency studies, organoids were treated with CFTR and PKA inhibitors for 2 hours prior to forskolin administration

(10 μM CFTRinh-172- Selleck Chemicals #S7139; 5 μM PPQ-102- Selleck Chemicals #S1565;
20 μM H-89- Fisher #501873404; 20 μM forskolin- Selleck Chemicals #S2449). Restoration of
CFTR function was investigated using CFTR potentiators (10 μM Icenticaftor (QBW251)- Selleck
Chemicals #S9973; 10 μM Ivacaftor- MedChemExpress #HY-13017; 10 μM Ivacaftor- Selleck
Chemicals #S1144), CFTR corrector (10 μM Lumacaftor- Selleck Chemicals #S1565; 10 μM
Elexacaftor- MedChemExpress #HY 111772, Selleck Chemicals #S8851; 10 μM Tezacaftor-
Selleck Chemicals #S7059). Mouse normal organoids treated with 20 μM forskolin were used as
a positive control for assay. All compounds were dispensed using a Tecan HP D300e in DMSO.
Organoids were imaged using brightfield imaging on a Tecan Spark Cyto. Images were taken at
time of treatment and 6 hours post treatment with CFTR correctors, potentiators, or forskolin. The
area of 10 organoids was measured at these two time points using either QuPath 0.5.0 or ImageJ.
The relative area at 6 hours was used for comparison across treatment groups. T-tests and
ANOVA statistical tests were conducted.
385
386 *Data visualization*
387
388 Data was visualized using R studio (2024.04.2+764) and GraphPad Prism (9 & 10). Figures were
389 arranged in Adobe Illustrator.
390
391 *Data Availability*
392
393 Raw data for whole genome sequencing and RNA-seq will be publicly available on NCBI the
394 database of genotype and phenotypes (dbGAP) upon final publication of manuscript.
395
396

Results

Generation of organoids from patients with human chronic pancreatitis (CP) with high efficiency

We received 51 CP surgical specimens (**Table 1, S1, Figures 1A-B, S1A-C, S2A-B**). The patient cohort was predominantly female (65%), with a mean age of 38 years. The mean age was 45 years for patients with idiopathic CP, 25 years for genetic CP, and 47 years for alcohol-related CP. Most patients self-identified as white and non-Hispanic. Idiopathic CP was the most common diagnosis, accounting for 41% of the cohort (**Figure 1B**). Among the 51 patient specimens received, 24 were from patients who underwent genetic testing. Pathogenic variants were reported in CFTR (20%, n=10), PRSS1 (14%, n=7), and SPINK1 (12%, n=6) (**Figure 1C**).

Most CP specimens were obtained following total pancreatectomy islet auto transplantation (TPIAT, 75%), and the remainder were from partial pancreatectomies (25%) (**Figures 1B, S1A-C, Table S1**). Pancreatectomy specimens were obtained within 2 hours post-resection with intact parenchyma but were often heavily cauterized (**Figure S1B-hCP1, hCP3, S1C- hCP10**). TPIAT specimens were partially dissociated and then received typically 24-48 hours post-surgery, often with disrupted parenchyma. When sufficient intact tissue was available, a portion of the specimens were fixed for histology (**Figures 1A, S1A**). Histologic assessment of the primary specimens by a pathologist confirmed classic CP features, including parenchymal atrophy, fibrosis, and ductal abnormalities (**Table 1, S1, Figures 1A, S2A-B**). Fibrosis scoring and quantification of collagen deposition revealed substantial desmoplasia, with some specimens containing little to no remaining acinar epithelial tissues (**Table 1, S1, Figures 1B, S2A-B**).

Organoids were observed after 1-7 days in culture and could be passaged every 4-7 days, 12-25 times, similar to the passaging potential of normal pancreas (NP) organoids (**Figure S1D-E**). PDO generation efficiency was determined after 5 passages to exclude lines that did not expand. Generation of 3 PDO lines are still in process and have not yet reached 5 passages. Out of the 48 specimens that have reached evaluation criteria, 36 yielded PDO lines. We successfully generated organoids from 70% of CP specimens (**Table 1, Figure 1D**). Failures occurred due to a lack of viable epithelial cells when tissue was frozen upon arrival, extensively cauterized, or composed primarily of fibrosis (**Figure S2B, Table S1**). PDO generation efficiency was higher for TPIAT specimens (77%) relative to partial pancreatectomy samples (56%) (**Figure 1D**).

CP organoids displayed a consistent, homogenous cystic morphology with a single cell layer (**Figure 1E**). Organoid morphology, passage time, and passaging potential did not differ across CP etiologies (**Figure S1D-E**). CP PDOs are comprised of rapidly dividing ductal epithelial cells as demonstrated by incorporation of EdU and immunofluorescence (IF) for CK19 (**Figure 1F-G, S1F**).

PDOs have high genetic concordance and reveal important and previously unannotated alterations

We assessed the genetic fidelity of PDOs by performing genomic analyses on matched primary and PDO specimens. Genomic alterations were compared between PDOs and primary tissue by whole genome sequencing (WGS), whole exome sequencing (WES), or targeted sequencing (**Figures 2, S3A, Table S1**). More than 9 million single nucleotide polymorphisms (SNPs) and almost 2 million insertions and deletions (indels) were compared by WGS, revealing an average 98% match rate with an intra correlation coefficient (ICC) of 0.96 between 14 pairs of PDOs and primary tissue (**Figures 2, S3A**). More than 80,000 SNPs and 7,000 indels were compared by WES, revealing an average 91% match rate with ICC of 0.83 between 6 pairs of PDOs and primary tissue (**Figures 2**, denoted by [#] in **S3A**). These data demonstrate the high genetic concordance between patient-matched primary tissue and PDOs. NP PDOs did not

contain any pancreatitis-associated pathogenic mutations nor PDA-associated oncogenic alterations.

We then queried germline variants in 14 genes associated with hereditary pancreatitis. All clinically annotated PRSS1, SPINK1, and CFTR germline variants are retained in our PDOs (**Figure 2**). PRSS1 has five paralogs with 90-91% nucleotide similarity, resulting in reads aligning to multiple locations with low mapping quality scores (MapQ)[70,71]. The PRSS1 N91I mutation was called with high confidence (hCP49), while the R122H (hCP24, hCP35) and R116C mutations (hCP49) were identified with low MapQ scores (**Figure 2**). Mutations in SPINK1 were confirmed in both PDO and pancreatic tissue from hCP08, hCP11, hCP30, hCP36, and hCP49. Interestingly, in 2 PDOs from patients with idiopathic CP with no previous genetic testing, we found SPINK1 N34S mutation (hCP07, hCP48) (**Figure 2**). In addition, we identified an unannotated SPINK1 P55S mutation (hCP35) and LPL N318S mutation (hCP32), which is associated with hyperlipidemia[72] and hereditary CP[73] (**Figure 2**).

Intronic variants of CFTR are common[30] and were confirmed in agreement with patient genetic testing results (hCP25, hCP28, and hCP32) (**Figure 2**). 9T variants are included in our assessment because their impact on exon 9 skipping in the pancreas in unknown despite minimal effect in nasal epithelium[74]. We confirmed the PDOs recapitulate coding sequence variants of CFTR, including R75Q, R117H, E217G, R347P, W1282*, and Q1352H mutations (**Figure 2**)[75-81]. Additionally, 75% of the CP PDO cohort harbors the CFTR V470M variant, which has conflicting reports of pathogenicity in association with CP and is likely a passenger mutation[75,80-84]. Additional G576A and R668C CFTR mutations were detected in hCP07, which have been found to be co-mutated in hereditary pancreatitis[85-87]. Also, a mild V201M CFTR variant was found in hCP32[88]. CFTR mutations associated with hereditary CP are distinct from cystic fibrosis and often do not overlap[25-32]. None of the cystic fibrosis associated CFTR mutations (CFTR2 database, n=1092) are found in the CP PDO cohort (**Figure S3B**).

Importantly, CP is a risk factor for developing pancreatic cancer and some hereditary CP patients have up to a 55% lifetime risk. The PDO platform enables identification of potential somatic oncogenic variants that are present in a small percentage of epithelial cells that would be obscured by the low epithelial cellularity of primary tissues. Here, we identified oncogenic KRAS G12D mutation in hCP30 with an allele frequency of 0.21 that was not detected in the primary specimen. We also found oncogenic TP53 R175H mutation in hCP28 at an allele frequency of 0.05 that was not detected in the primary specimen. These data suggest that the CP PDO models may be useful in identifying patients at higher risk of progressing to PDA.

Gene expression analyses demonstrate distinct cell signaling programs in CP organoids

To understand the specific programs differentially regulated in CP PDOs, we performed RNA-seq on 24 CP, 9 NP, and 4 normal adjacent to tumor (NAT) PDOs. Ductal lineage identity is maintained by all PDOs with no evidence of non-epithelial cell contamination (**Figure S3C**), concordant with positivity for the ductal epithelial marker CK19 by IF (**Fig. 1F**). CP PDOs have a distinct transcriptomic profile as compared to NP and NAT PDOs (**Figures 3A-C**). Gene set enrichment analyses (GSEA) reveal that CP PDOs are enriched in gene sets associated with inflammatory response, TNFA signaling via NFKB, KRAS signaling, Epithelial to Mesenchymal Transition (EMT), and TGF beta signaling while having depressed Oxidative Phosphorylation relative to NP and NAT PDOs (**Figure 3D**). Compared to NP and NAT PDOs, CP expressed higher levels of fibrosis and inflammatory cytokine associated genes (e.g., *COL1A1*, *TGFB2*, *IL1A*, *CSF1*) (**Figure 3E**), suggesting that CP PDOs maintain the fibrosis and inflammatory programs present in CP patients[89].

Principal component analysis of CP PDOs suggested the presence of 3 distinct clusters (**Figure S4A**). The PDO transcriptomes were independently clustered using nonnegative matrix factorization (NMF)[63,64], revealing 3 stable subtypes in CP PDO cultures (basis 1-3) (**Figures 3F**,

S4B). The molecular subtypes identified using this approach do not cluster according to etiology
(**Figure 3F**), suggesting distinct subtypes independent of TIGAR-O classification. We performed
GSEA on each CP subtype relative to NP and NAT PDOs (**Figure S4C**). The basis 1 subtype is
depleted in oxidative phosphorylation and enriched in EMT and apoptosis gene sets. The
apoptosis gene set includes proteins that involve cellular stress response pathways, such as
glutathione peroxidase 3 (GPX3). The basis 2 subtype is enriched in E2F targets, MYC targets,
and myogenesis. The basis 3 subtype is enriched in E2F targets. All subtypes trend towards
enrichment of the inflammatory response gene set.
To validate these subtypes in matched primary tissue specimens, we identified the genes
that were differentially expressed between the 3 subtypes of CP and have the highest Kim-score,
and therefore drive these distinct clusters (**Figure S4D**). Subtype defining genes were evaluated
by RNA *in situ* hybridization (**Figure S5A**). We found that a patient specimen from the basis 1
subtype PDO hCP1 expressed the highest levels of GPX3 as expected. The patient matched
specimen for the basis 2 subtype PDO hCP3 exhibited the highest level of aquaporin 1 (AQP1*)*.
The basis 3 subtype was defined by BPI-family member B 1 (BPIFB1*)*, which exhibited the highest
levels in patient specimen from the basis 3 subtype PDO hCP48. This indicates that the genes
driving CP PDO transcriptional subtypes can also be detected in the corresponding primary
tissue.
Transcriptomic analyses on primary CP tissues are challenging due to the high degree of
fibrosis and the high levels of RNases. We further explored the expression of subtype-defining
genes using single-nucleus RNA sequencing (snRNA-seq) data from human NAT and CP
samples[69]. In the ductal cell cluster, we observed that subtype defining genes were expressed in
primary pancreatic ductal epithelium (**Figure S5B-C**), but insufficient sample and cell numbers
made identification of subtypes in these data challenging. These findings confirm that the genes
driving transcriptional subtypes in the CP PDOs are present in an independent patient cohort.
Additional analysis comparing the transcriptomes of our CP PDO cohort to human
pancreatic cancer organoids revealed genes that are uniquely expressed in either disease
(**Figure S5D**). This along with the discovery of somatic oncogenic variants in 2 of our CP PDOs
demonstrate that this model can be used to study patients with heightened risk of cancer and to
identify early detection biomarkers.
527
CP organoids recapitulate protein expression from primary tissue specimens and exhibit elevated
inflammatory and mitogenic signaling

Immunohistochemistry (IHC) and immunoblotting (IB) were used to evaluate proteins and
glycan levels associated with pancreatitis, as well as mitogenic and inflammatory signaling. CP
and NP PDOs exhibited protein alterations consistent with matched patient specimens (**Figures
4A-B**). We previously found that up to ~90% of adult CP has local elevation of CA19-9[90]. Patient
specimens had a range of CA19-9 that matched CP PDOs (**Figures 4A-B**). Activation of EGFR
has been reported in human CP[91]. A subset of CP PDOs have elevated levels of phospho-EGFR
and downstream effectors AKT and ERK relative to NP PDOs (**Figure 4A**), suggesting there are
"mitogenic" and "non-mitogenic" CP subtypes. NP PDOs primarily express mature CFTR by IB
(upper band, C-band, 150-160KD[92]) with very little immature CFTR (lower band, B-band, 135-
140KD[92]) (**Figures 4A, S6A**). Of those with CFTR mutations, hCP11 and hCP44 had low levels
of mature CFTR by IB, which was replicated in the primary patient specimen IHC (**Figures 4A-
B**). This corresponds with what is known about the pre-mature stop codon mutation present in
hCP11 and compound CFTR R347P; R117H mutation in hCP44. These mutations have been
reported to lead to major folding defects, and thus, as expected, do not produce mature CFTR[79].
E217G and R75Q CFTR mutations found in hCP08 and hCP17, respectively, impact anion
channel function[75,92], but not protein maturation. These two lines have mature CFTR, as predicted.
hCP15, a non-hereditary case attributed to alcohol and hypothyroidism, also demonstrated loss

of mature CFTR protein despite having a benign V470M mutation (data not shown). Decreased
CFTR expression and function has been shown to occur secondary to alcohol[93,94], tobacco[95], and
hypothyroidism[96]. These findings suggest that CFTR modulators that increase protein activity may
help even CP patients without pathogenic CFTR mutations. Carbonic anhydrase 2 (CA2), which
catalyzes the conversion of CO_2 into HCO_3^-, was decreased in most CP PDOs when compared
to NP, pointing to another possible mechanism through which HCO_3^- secretion may be impaired
in CP (**Figure 4A**).
To identify fibroinflammatory mediators in CP PDOs at the protein level, we evaluated
conditioned medium from 14 CP and 9 NP PDOs. IL1A, TGFB2, CX3CL1, and other
fibroinflammatory secreted factors were elevated in CP PDOs and reflected the changes
observed in gene expression and published changes in CP patients[97,98] (**Figure 4C, S6B**),
demonstrating the ability of CP PDOs to model inflammatory changes seen in patients. Together,
these data support that PDOs retain key molecular features of CP, including alterations in cell
signaling proteins.
561
CFTR function in CP organoids is restored upon pharmacologic modulation

The forskolin-induced swelling (FIS) assay was developed to measure CFTR dysfunction
and previously deployed in intestinal and rectal organoids from cystic fibrosis patients to predict
response to CFTR modulators[39,40]. Intestinal current measurements and FIS activity were tightly
correlated, demonstrating that FIS assays can serve as a surrogate for more complex
electrophysiological measurements[40]. We optimized the FIS assay in murine and human NP
PDOs (**Figure 5A**). Vehicle treated organoids had no response, but forskolin or CFTR potentiator
treatment caused organoid swelling (**Figures 5B-F, S7A-C**). Swelling was inhibited when mouse
and human NP organoids were treated with CFTR inhibitors (CFTRi-172, PPQ-102) or a PKA
inhibitor (H-89), demonstrating that swelling is dependent on forskolin induced activation of CFTR
via PKA (**Figures 5D-F, S7B**). Mouse normal organoids served as a positive control while testing
the CFTR function of CP PDOs (**Figure S7A-B)**. Half of the CP PDOs evaluated (n=4 out of 8)
were FIS impaired across multiple experimental replicates (**Figure 5G**, **S7D**).
Most CFTR wild type PDOs (hCP1, hCP3, hCP48, hCP49) swelled in response to forskolin
treatment as expected (**Figure 5G**). Unexpectedly, hCP16, which is CFTR wildtype and exhibits
normal CFTR protein levels (**Figure 4A**), is FIS deficient (**Figure 5G, S7D**). We hypothesized that
organoids deficient in FIS could be rescued by treatment with CFTR potentiators, which increase
the basal activity of CFTR (**Figure 6A**). Treatment of hCP16 with CFTR potentiators did not
restore swelling (**Figure S7D**), suggesting that the dysfunction in FIS in this PDO is independent
of CFTR.
hCP11 did not swell, which can be attributed to the lack of mature CFTR due to a
premature stop codon mutation (**Figures 4A**)[99]. Given the lack of mature CFTR protein, treatment
with CFTR potentiators did not restore swelling (**Figure 6B**). However, treatment with the read
through agent, Ataluren, was able to restore FIS response (**Figure 6C**) and increase levels of
mature CFTR (**Figure 6D**). These data demonstrate that we can successfully achieve genotype-
driven precision therapeutics in CP PDOs.
hCP35 is CFTR wildtype and exhibits normal CFTR protein levels. Unexpectedly, hCP35
had a limited FIS response (**Figures 5G, 6E**). Swelling was restored by treatment with CFTR
potentiators, Ivacaftor and Icenticaftor (**Figure 6E**). This demonstrates ability of potentiators to
increase CFTR function in PDOs even with wild type CFTR.
hCP44 was also deficient in FIS (**Figure 6F, 6I**), which was expected given the CFTR
R347P, R117H mutations that impair protein folding and anion transport[100] and because the low
levels of mature CFTR protein (**Figure 4A**). The protein folding defect can be ameliorated with
CFTR correctors. This class of drugs increases the correct folding and plasma membrane
localization of CFTR and the combination of correctors and potentiators is FDA approved for the

treatment of cystic fibrosis (e.g., Orkambi: Ivacaftor, Lumacaftor)[101]. hCP44 FIS was rescued by
treatment with a combination of correctors and potentiators (Orakambi, and Ivacaftor +
Elexacaftor) (**Figure 6G**), but not monotherapies (**Figure 6F**). This demonstrates that CP PDO
models can used to identify the most effective combinatorial treatment strategies in the
background of complex allelic combinations.
We tested if treatment with CFTR correctors and potentiators increases mature CFTR
protein levels in FIS deficient hCP35 and hCP44. Single agent treatment of hCP35 with the
potentiator Ivacaftor marginally increased mature CFTR, which was further increased by ETI
treatment (**Figure 6H**). As expected from our FIS results, ETI treatment increases mature CFTR
protein levels in hCP44 (**Figure 6I**). Importantly, treatment of hCP44 with single agent CFTR
potentiator caused a decrease in mature CFTR protein levels (**Figure 6I**). This is in line with
previous findings that Ivacaftor treatment can interfere with CFTR stability in the background of
certain mutations, such as the R347P, R117H alterations found in hCP44[102]. These findings stress
the importance of investigating CFTR modulator response in patient-specific contexts, especially
in the setting of complex allelic combinations. Altogether these data indicate that patients with
and without pathogenic *CFTR* alterations could benefit from ETI treatment to increase mature
CFTR expression. Treatment with ETI in both hCP35 and hCP44 also led to a decrease in pERK
and pSTAT3 levels (**Figures 6H-I**), suggesting that restoration of CFTR function can decrease
mitogenic and inflammatory signaling. Our findings suggest that restoring normal ductal function
can contribute to the resolution of inflammation.
617
Discussion

Surgical samples from chronic pancreatitis patients are rare, making this CP PDO biobank
a unique resource. Importantly, while our PDO biobank encompasses the major etiologies of CP,
the patients are a highly selected cohort that required surgical management. In the future, this
limitation may be addressed through generation of CP PDOs from pancreatic juice, which we
accomplished in 2 cases and is a more accessible sample source.
The identification of pathogenic germline alterations in 3 patients diagnosed with idiopathic
CP suggests that genetic testing may have broader benefit in CP clinical care. Two of the CP
PDOs also contain oncogenic mutations in TP53 or KRAS, known drivers of pancreatic ductal
adenocarcinoma. This is expected given the high rate of sporadic KRAS mutations that have been
found in autopsy studies of the normal human pancreas[103]. Work has also shown that even in the
context of wild type KRAS, TP53 mutation promotes injury in a non-cancer context[104]. CP PDO
models will facilitate our understanding of which patients are at risk of severe disease and
progression to PDA. Identification of the KRAS and TP53 mutations were uniquely possible from
analysis of the PDOs and not the tissue. Further work is needed to explore how CP PDOs can be
leveraged to model the transition from a chronic inflammatory state to invasive carcinoma.
CP is primarily classified by etiology, but our transcriptomic analyses reveal molecular
subtypes that do not correlate with etiology, suggesting that distinct cell signaling changes drive
CP subtypes. The validation of these subtypes remains challenging due to the limited availability
of these surgical tissue specimens and the high abundance of RNases and proteases in the
pancreas. Future work will focus on creating primary CP specimen data resources for
corroboration of these findings and for investigation of clinical correlates. These subtypes offer an
initial framework to explore potential signaling pathways driving CP pathogenesis.
Surprisingly, the CP PDOs maintain a fibroinflammatory gene and protein expression
signature despite being removed from the microenvironment. These data suggest that ductal cells
can contribute to maintenance of the fibroinflammatory program and opens several lines of
questioning. When are these ductal gene expression programs established and how are they
maintained? Is the intrinsic epithelial fibroinflammatory program sufficient to cause CP and
promote PDA initiation? Would reversing this program restore pancreatic function? CP PDOs

enable exploration of these important questions at unprecedented depth as well as the appropriate physiologic context.

The use of PDOs for functional precision medicine has been previously established in pancreas cancer[43,46,50,52]. Here, we show that CP PDOs can be similarly used as a platform for studying CFTR function in different complex genetic contexts and delineating treatment response. Our study shows that CP PDOs are commonly deficient in FIS. This deficiency can be effectively reversed by treatment with FDA-approved CFTR modulators, suggesting potential targeted therapeutic strategies for the treatment of CFTR-mutant CP. An important consideration when testing CFTR modulators to treat CP includes organ context and species. CFTR mutations in cystic fibrosis patients primarily impacts chloride secretion whereas in the pancreas, it contributes substantially to bicarbonate transport. This highlights the importance of investigating CP CFTR mutations in pancreatic cell types. In addition, mutant mouse CFTR does not respond to human CFTR modulators[41]. This is further complicated by reduced levels of bicarbonate in mouse pancreatic juice (50 mM) relative to human (140 mM)[37], suggesting a lower level of reliance on bicarbonate in the mouse pancreas. Finally, as evidenced in our cohort, the genetics of CP is complex. There are more than 2,000 different CFTR mutations and limited information regarding drug response to most of these alterations, especially those associated with hereditary CP. Even in the setting of the same pathogenic alterations, patient response to CFTR modulators is very heterogeneous, highlighting the need for precision medicine approaches[35-40]. The physiologically relevant context of human CP PDOs enables direct investigation of genotypic relationships to therapeutic responses. We further demonstrate CFTR dysfunction in CP PDOs with wildtype CFTR that can be restored with potentiator treatment, raising the possibility that this treatment approach could be widely applicable to a large patient population. Our work also highlights the utility of restoring normal homeostatic function to reduce inflammatory signaling as evidenced by decreased STAT3 activation following CFTR modulator treatment. Altogether, we demonstrate the power of our PDO platform to study the underlying molecular pathogenesis of human CP and pave the way for precision therapeutics.

Acknowledgements
The authors would like to extend our deepest gratitude to all the patients who generously consented to participate in this study. The authors would like to thank all the members of the Engle laboratory, the UCSD Biorepository and Tissue Technology Shared Resource (BTTSR), the UCSD Department of Surgery, the University of Minnesota Departments of Surgery and Pediatrics, April Williams, Ling Ouyang, and Tzu-when Wang at the NGS Core Facility of the Salk Institute, Elsa Molina at the Single-Cell and Spatial Omics core of the Salk Institute, Daniela Boassa, Sammy Weiser Novak and Elsie Quansah at the Biophotonics Core Facility of the Salk Institute, and the Lifesharing Donate Life organization.

Funding Statement
V.O.V is supported through the Salk Institute Cancer Training Grant (T32CA009370). J.C.L. was supported through the Salk Institute Cancer Training Grant (T32CA0093790). This work was supported by the Razavi Newman Integrative Genomics and Bioinformatics Core Facility of the Salk Institute with funding from NIH-NCI CCSG: P30 CA01495, NIH-NIA San Diego Nathan Shock Center P30 AG068635, and the Helmsley Trust. This work was supported by the Waitt Advanced Biophotonics Core Facility of the Salk Institute with funding from NIH-NCI CCSG: P30 CA01495, NIH-NIA San Diego Nathan Shock Center P30 AG068635, and the Waitt Foundation. This work was supported by the NGS Core Facility of the Salk Institute with funding from NIH-NCI CCSG: P30 CA01495, NIH-NIA San Diego Nathan Shock Center P30 AG068635, the Chapman Foundation and the Helmsley Charitable Trust. The Biorepository and Tissue Technology Shared Resource is supported by a National Cancer Institute Cancer Center Support Grant (CCSG P30CA23100) and by recharges from users. This research program was supported by Mission

Land Acknowledgement
The Salk community holds great respect for the land and the original people of the area where
our campus is located. The Institute is built on the territory of the Kumeyaay Nation. Today, the
Kumeyaay people continue to maintain their political sovereignty and cultural traditions as vital
members of the San Diego community. We acknowledge their tremendous contributions to our
region and thank them for their stewardship.
706

707 **Declaration of Interests**
708 J. Gibson, and D. Whitcomb are founders of Ariel Precision Medicine. D. Whitcomb is a member
709 of its scientific advisory board at Ariel Precision Medicine. P. Greer is an employee of Ariel
710 Precision Medicine.
711
712

References

713
714
1.   Kleeff, J., Whitcomb, D.C., Shimosegawa, T., Esposito, I., Lerch, M.M., Gress, T.,
Mayerle, J., Drewes, A.M., Rebours, V., Akisik, F., et al. (2017). Chronic
pancreatitis. Nature Reviews Disease Primers *3*, 17060. 10.1038/nrdp.2017.60.
2.   ZM, S., D, M., H, Y., M, D., KY, Z., R, B., JJ, W., C, K., and KT, P. (2018 Aug).
Nationwide Trends in Acute and Chronic Pancreatitis Among Privately Insured
Children and Non-Elderly Adults in the United States, 2007-2014 - PubMed.
Gastroenterology *155*. 10.1053/j.gastro.2018.04.013.
3.   Jorge D Machicado (1), V.R., Dhiraj Yadav (1) (28/07/2016). Epidemiology of
Chronic Pancreatitis. Pancreapedia: The Exocrine Pancreas Knowledge Base.
10.3998/panc.2016.13.
4.   Whitcomb, D.C., and Group, N.A.P.S. (2019). Pancreatitis: TIGAR-O Version 2
Risk/Etiology Checklist With Topic Reviews, Updates, and Use Primers. Clinical
and   Translational   Gastroenterology   *10*,   e00027.
10.14309/ctg.0000000000000027.
5.   Vege, S.S., and Chari, S.T. (2022). Chronic Pancreatitis. New England Journal of
Medicine *386*, 869-878. 10.1056/nejmcp1809396.
6.   JS, T., L, K., U, B., and M, G. (03/23/2009). Smoking and risk of acute and chronic
pancreatitis among women and men: a population-based cohort study - PubMed.
Archives of internal medicine *169*. 10.1001/archinternmed.2008.601.
7.   Yan, M.-X., and Li, Y.-Q. (2006/04/01). Gall stones and chronic pancreatitis: the
black   box   in   between.   Postgraduate   Medical   Journal   *82*.
10.1136/pgmj.2005.037192.
8.   Otsuki, M., and Otsuki, M. (2003/04). Chronic pancreatitis in Japan: epidemiology,
prognosis, diagnostic criteria, and future problems. Journal of Gastroenterology
2003 38:4 *38*. 10.1007/s005350300058.
9.   Yadav, D., Whitcomb, D.C., Yadav, D., and Whitcomb, D.C. (2010-02-02). The role
of alcohol and smoking in pancreatitis. Nature Reviews Gastroenterology &
Hepatology 2010 7:3 *7*. 10.1038/nrgastro.2010.6.
10.   Singh, V.K., Yadav, D., and Garg, P.K. (2019). Diagnosis and Management of
Chronic Pancreatitis. JAMA *322*, 2422-2434. 10.1001/jama.2019.19411.
11.   AE, P., J, L., P, G., J, A., S, A., ST, A., MA, A., J, B., PA, B., RE, B., et al. (2018
Jul). Known genetic susceptibility factors for chronic pancreatitis in patients of
European ancestry are rare in patients of African ancestry - PubMed.
Pancreatology : official journal of the International Association of Pancreatology
(IAP) ... [et al.] *18*. 10.1016/j.pan.2018.05.482.
12.   J, C., R, L., DI, W., A, C., and R, C. (07/03/1999). Mutations of the cationic
trypsinogen gene in patients with chronic pancreatitis - PubMed. Lancet (London,
England) *354*. 10.1016/S0140-6736(99)01814-0.
13.   Schwarzenberg, S.J., Uc, A., Zimmerman, B., Wilschanski, M., Wilcox, C.M.,
Whitcomb, D.C., Werlin, S.L., Troendle, D., Tang, G., Slivka, A., et al.
(2019/04/01). Chronic Pancreatitis. Journal of Pediatric Gastroenterology and
Nutrition *68*. 10.1097/MPG.0000000000002279.
14.   LaRusch, J., and Whitcomb, D.C. (2011 Sep). Genetics of pancreatitis. Current
opinion in gastroenterology *27*. 10.1097/MOG.0b013e328349e2f8.

15.   Whitcomb, D.C., Gorry, M.C., Preston, R.A., Furey, W., Sossenheimer, M.J.,
Ulrich, C.D., Martin, S.P., Gates, L.K., Amann, S.T., Toskes, P.P., et al. (1996).
Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene.
Nature Genetics *14*, 141-145. 10.1038/ng1096-141.
16.   Howes, N., Lerch, M.M., Greenhalf, W., Stocken, D.D., Ellis, I., Simon, P.,
Truninger, K., Ammann, R., Cavallini, G., Charnley, R.M., et al. (2004). Clinical and
genetic characteristics of hereditary pancreatitis in Europe. Clinical
Gastroenterology and Hepatology *2*, 252-261. 10.1016/s1542-3565(04)00013-8.
17.   Schnúr, A., Beer, S., Witt, H., Hegyi, P., and Sahin-Tóth, M. (2014). Functional
effects of 13 rare PRSS1 variants presumed to cause chronic pancreatitis. Gut *63*,
337. 10.1136/gutjnl-2012-304331.
18.   Whitcomb, D.C. (2010). Genetic Aspects of Pancreatitis. Annual Review of
Medicine *61*, 413-424. 10.1146/annurev.med.041608.121416.
19.   Threadgold, J., Greenhalf, W., Ellis, I., Howes, N., Lerch, M.M., Simon, P., Jansen,
J., Charnley, R., Laugier, R., Frulloni, L., et al. (2002). The N34S mutation of
SPINK1 (PSTI) is associated with a familial pattern of idiopathic chronic
pancreatitis but does not cause the disease. Gut *50*, 675. 10.1136/gut.50.5.675.
20.   Grocock, C.J., Rebours, V., Delhaye, M.N., Andrén-Sandberg, A., Weiss, F.U.,
Mountford, R., Harcus, M.J., Niemczyck, E., Vitone, L.J., Dodd, S., et al. (2010).
The variable phenotype of the p.A16V mutation of cationic trypsinogen (PRSS1)
in pancreatitis families. Gut *59*, 357. 10.1136/gut.2009.186817.
21.   Ooi, C.Y., Dorfman, R., Cipolli, M., Gonska, T., Castellani, C., Keenan, K.,
Freedman, S.D., Zielenski, J., Berthiaume, Y., Corey, M., et al. (2011). Type of
CFTR Mutation Determines Risk of Pancreatitis in Patients With Cystic Fibrosis.
Gastroenterology *140*, 153-161. 10.1053/j.gastro.2010.09.046.
22.   Moniaux, N., Andrianifahanana, M., Brand, R.E., and Batra, S.K. (2004). Multiple
roles of mucins in pancreatic cancer, a lethal and challenging malignancy. British
Journal of Cancer *91*, 1633-1638. 10.1038/sj.bjc.6602163.
23.   Viktoria Venglovecz, P.H., 2 Michael A. Gray3 (07/04/2021). Ion Channels in
Pancreatic Duct Epithelial Cells in Health and Disease. Pancreapedia: The
Exocrine Pancreas Knowledge Base. 10.3998/panc.2021.08.
24.   Richter, L.D., Ruderfer, D.M., Peterson, J.F., and Bastarache, L.A. (2025). Beyond
Carrier Status: CFTR Heterozygosity as an Overlooked Clinical Risk Factor for
Pancreatitis. Clinical Genetics *n/a*. https://doi.org/10.1111/cge.14780.
25.   Zoller, H., Egg, M., Graziadei, I., Creus, M., Janecke, A.R., Löffler-Ragg, J., and
Vogel, W. (2007). CFTR gene mutations in pancreatitis: Frequency and clinical
manifestations in an Austrian patient cohort. Wiener klinische Wochenschrift *119*,
527-533. 10.1007/s00508-007-0849-5.
26.   Witt, H. (2003). Chronic pancreatitis and cystic fibrosis. Gut *52*, ii31.
10.1136/gut.52.suppl_2.ii31.
27.   Truninger, K., Malik, N., Ammann, R.W., Muellhaupt, B., Seifert, B., Müller, H.J.,
and Blum, H.E. (2001). Mutations of the cystic fibrosis gene in patients with chronic
pancreatitis. The American journal of gastroenterology *96*, 2657-2661.
10.1111/j.1572-0241.2001.04047.x.
28.   Kiesewetter, S., Macek, M., Davis, C., Curristin, S.M., Chu, C.S., Graham, C.,
Shrimpton, A.E., Cashman, S.M., Tsui, L.C., Mickle, J., et al. (1993). A mutation in

805   CFTR produces different phenotypes depending on chromosomal background.
806   Nature Genetics *5*, 274-278. 10.1038/ng1193-274.

807 29.   Noone, P.G., Zhou, Z., Silverman, L.M., Jowell§, P.S., Knowles, M.R., and Cohn§,
808   J.A. (2001). Cystic fibrosis gene mutations and pancreatitis risk: Relation to
809   epithelial ion transport and trypsin inhibitor gene mutations. Gastroenterology *121*,
810   1310-1319. 10.1053/gast.2001.29673.

811 30.   Audrézet, M.-P., Chen, J.-M., Maréchal, C.L., Ruszniewski, P., Robaszkiewicz, M.,
812   Raguénès, O., Quéré, I., Scotet, V., and Férec, C. (2002). Determination of the
813   relative contribution of three genes–the cystic fibrosis transmembrane
814   conductance regulator gene, the cationic trypsinogen gene, and the pancreatic
815   secretory trypsin inhibitor gene–to the etiology of idiopathic chronic pancreatitis.
816   European Journal of Human Genetics *10*, 100-106. 10.1038/sj.ejhg.5200786.

817 31.   Lukacs, G.L., and Verkman, A.S. (2012). CFTR: folding, misfolding and correcting
818   the ΔF508 conformational defect. Trends in Molecular Medicine *18*, 81-91.
819   10.1016/j.molmed.2011.10.003.

820 32.   Cohn, J.A., Friedman, K.J., Noone, P.G., Knowles, M.R., Silverman, L.M., and
821   Jowell, P.S. (1998-09-03). Relation between Mutations of the Cystic Fibrosis Gene
822   and Idiopathic Pancreatitis. New England Journal of Medicine *339*.
823   10.1056/NEJM199809033391002.

824 33.   Wiseman, A.J., and Ooi, C.Y. (2021). Clinical Pancreatology for Practising
825   Gastroenterologists and Surgeons 10.1002/9781119570097.ch47.

826 34.   Ooi, C.Y., Dupuis, A., Ellis, L., Jarvi, K., Martin, S., Ray, P.N., Steele, L., Kortan,
827   P., Gonska, T., Dorfman, R., et al. (2014). Does extensive genotyping and nasal
828   potential difference testing clarify the diagnosis of cystic fibrosis among patients
829   with single-organ manifestations of cystic fibrosis? Thorax *69*, 254.
830   10.1136/thoraxjnl-2013-203832.

831 35.   Clancy, J.P., Rowe, S.M., Accurso, F.J., Aitken, M.L., Amin, R.S., Ashlock, M.A.,
832   Ballmann, M., Boyle, M.P., Bronsveld, I., Campbell, P.W., et al. (2012). Results of
833   a phase IIa study of VX-809, an investigational CFTR corrector compound, in
834   subjects with cystic fibrosis homozygous for the F508del-CFTR mutation. Thorax
835   *67*, 12-18. 10.1136/thoraxjnl-2011-200393.

836 36.   Flume, P.A., Liou, T.G., Borowitz, D.S., Li, H., Yen, K., Ordoñez, C.L., and Geller,
837   D.E. (2012). Ivacaftor in subjects with cystic fibrosis who are homozygous for the
838   F508del-CFTR mutation. Chest *142*, 718-724. 10.1378/chest.11-2672.

839 37.   Ishiguro, H., Yamamoto, A., Nakakuki, M., Yi, L., Ishiguro, M., Yamaguchi, M.,
840   Kondo, S., and Mochimaru, Y. (2012). Physiology and pathophysiology of
841   bicarbonate secretion by pancreatic duct epithelium. Nagoya J Med Sci *74*, 1-18.

842 38.   Ramsey, B.W., Davies, J., McElvaney, N.G., Tullis, E., Bell, S.C., Dřevínek, P.,
843   Griese, M., McKone, E.F., Wainwright, C.E., Konstan, M.W., et al. (2011). A CFTR
844   potentiator in patients with cystic fibrosis and the G551D mutation. N Engl J Med
845   *365*, 1663-1672. 10.1056/NEJMoa1105185.

846 39.   van Mourik, P., Michel, S., Vonk, A.M., Beekman, J.M., van der Ent, C.K., van
847   Mourik, P., Michel, S., Vonk, A.M., Beekman, J.M., and van der Ent, C.K. (2020-
848   06-03). Rationale and design of the HIT-CF organoid study: stratifying cystic
849   fibrosis patients based on intestinal organoid response to different CFTR-

850   modulators. Translational Medicine Communications 2020 5:1 *5*. 10.1186/s41231-
851   020-00060-3.

40.   Dekkers, J.F., Wiegerinck, C.L., de Jonge, H.R., Bronsveld, I., Janssens, H.M., de
      Winter-de Groot, K.M., Brandsma, A.M., de Jong, N.W.M., Bijvelds, M.J.C.,
      Scholte, B.J., et al. (2013-06-02). A functional CFTR assay using primary cystic
      fibrosis intestinal organoids. Nature Medicine 2013 19:7 *19*. 10.1038/nm.3201.

41.   McCarron, A., Parsons, D., and Donnelley, M. (2021). Animal and Cell Culture
      Models for Cystic Fibrosis: Which Model Is Right for Your Application? The
      American Journal of Pathology *191*, 228-242. 10.1016/j.ajpath.2020.10.017.

42.   Boj, Sylvia F., Hwang, C.-I., Baker, Lindsey A., Chio, Iok In C., Engle,
      Dannielle D., Corbo, V., Jager, M., Ponz-Sarvise, M., Tiriac, H., Spector, Mona S.,
      et al. (2015). Organoid Models of Human and Mouse Ductal Pancreatic Cancer.
      Cell *160*, 324-338. 10.1016/j.cell.2014.12.021.

43.   Tiriac, H., Belleau, P., Engle, D.D., Plenker, D., Deschênes, A., Somerville, T.,
      Froeling, F.E.M., Burkhart, R.A., Denroche, R.E., Jang, G.-H., et al. (2018).
      Organoid profiling identifies common responders to chemotherapy in pancreatic
      cancer. Cancer Discovery *8*, CD-18-0349. 10.1158/2159-8290.cd-18-0349.

44.   Kim, J., Hoffman, J.P., Alpaugh, R.K., Rhim, A.D., Rhimm, A.D., Reichert, M.,
      Stanger, B.Z., Furth, E.E., Sepulveda, A.R., Yuan, C.-X., et al. (2013). An iPSC
      Line from Human Pancreatic Ductal Adenocarcinoma Undergoes Early to Invasive
      Stages of Pancreatic Cancer Progression. Cell Reports *3*, 2088-2099.
      10.1016/j.celrep.2013.05.036.

45.   Grossman, J.E., Muthuswamy, L., Huang, L., Akshinthala, D., Perea, S., Gonzalez,
      R.S., Tsai, L.L., Cohen, J., Bockorny, B., Bullock, A.J., et al. (2021). Organoid
      Sensitivity Correlates with Therapeutic Response in Patients with Pancreatic
      Cancer. Clinical Cancer Research *28*, clincanres.4116.2020. 10.1158/1078-
      0432.ccr-20-4116.

46.   Gulay, K.C.M., Zhang, X., Pantazopoulou, V., Patel, J., Esparza, E., Babu, D.S.P.,
      Ogawa, S., Weitz, J., Ng, I., Mose, E.S., et al. (2023). Dual inhibition of KRASG12D
      and pan-ERBB is synergistic in pancreatic ductal adenocarcinoma. Cancer
      Research. 10.1158/0008-5472.can-23-1313.

47.   Ponz-Sarvise, M., Corbo, V., Tiriac, H., Engle, D.D., Frese, K.K., Oni, T.E., Hwang,
      C.-I., Öhlund, D., Chio, I.I.C., Baker, L.A., et al. (2019). Identification of Resistance
      Pathways Specific to Malignancy Using Organoid Models of Pancreatic Cancer.
      Clinical Cancer Research *25*, 6742-6755. 10.1158/1078-0432.ccr-19-1398.

48.   Hou, S., Tiriac, H., Sridharan, B.P., Scampavia, L., Madoux, F., Seldin, J., Souza,
      G.R., Watson, D., Tuveson, D., and Spicer, T.P. (2018). Advanced Development
      of Primary Pancreatic Organoid Tumor Models for High-Throughput Phenotypic
      Drug Screening. SLAS Discovery *23*, 574-584. 10.1177/2472555218766842.

49.   Wolff, R.A., Wang-Gillam, A., Alvarez, H., Tiriac, H., Engle, D., Hou, S., Groff, A.F.,
      Lucas, A.S., Bernard, V., Allenson, K., et al. (2018). Dynamic changes during the
      treatment of pancreatic cancer. Oncotarget *9*, 14764-14790.
      10.18632/oncotarget.24483.

50.   Lumibao, J.C., Okhovat, S.R., Peck, K.L., Lin, X., Lande, K., Yomtoubian, S., Ng,
      I., Tiriac, H., Lowy, A.M., Zou, J., and Engle, D.D. (2023). The impact of

extracellular matrix on the precision medicine utility of pancreatic cancer patient-
derived organoids. JCI Insight *9*, e172419. 10.1172/jci.insight.172419.

51. Letai, A., Bhola, P., and Welm, A.L. (2022). Functional precision oncology: Testing
tumors with drugs to identify vulnerabilities and novel combinations. Cancer Cell
40, 26-35. 10.1016/j.ccell.2021.12.004.

52. Boilève, A., Cartry, J., Goudarzi, N., Bedja, S., Mathieu, J.R.R., Bani, M.-A.,
Nicolle, R., Mouawia, A., Bouyakoub, R., Nicotra, C., et al. (2024). Organoids for
functional precision medicine in advanced pancreatic cancer. Gastroenterology.
10.1053/j.gastro.2024.05.032.

53. Shik Mun, K., Arora, K., Huang, Y., Yang, F., Yarlagadda, S., Ramananda, Y.,
Abu-El-Haija, M., Palermo, J.J., Appakalai, B.N., Nathan, J.D., et al. (2019-07-16).
Patient-derived pancreas-on-a-chip to model cystic fibrosis-related disorders.
Nature Communications 2019 10:1 *10*. 10.1038/s41467-019-11178-w.

54. Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G.,
Dunne, P.D., McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al. (2017).
QuPath: Open source software for digital pathology image analysis. Scientific
Reports *7*, 16878. 10.1038/s41598-017-17204-5.

55. Esposito, I., Hruban, R.H., Verbeke, C., Terris, B., Zamboni, G., Scarpa, A.,
Morohoshi, T., Suda, K., Luchini, C., Klimstra, D.S., et al. (2020/06/01). Guidelines
on the histopathology of chronic pancreatitis. Recommendations from the working
group for the international consensus guidelines for chronic pancreatitis in
collaboration with the International Association of Pancreatology, the American
Pancreatic Association, the Japan Pancreas Society, and the European
Pancreatic Club. Pancreatology *20*. 10.1016/j.pan.2020.04.009.

56. Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A.,
Goodman, A., Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E.,
et al. (2021-09-10). CellProfiler 4: improvements in speed, utility and usability.
BMC Bioinformatics 2021 22:1 *22*. 10.1186/s12859-021-04344-9.

57. SW, W., and S, A. (08/24/2018). FastQ Screen: A tool for multi-genome mapping
and quality control - PubMed. F1000Research *7*.
10.12688/f1000research.15931.2.

58. Martin, M. (2011/05/02). Cutadapt removes adapter sequences from high-
throughput sequencing reads. EMBnet.journal *17*. 10.14806/ej.17.1.200.

59. JM, M., S, C.-S., M, D., JG, M., T, H., I, J., JE, L., C, A., I, B., R, B., et al.
(01/06/2025). GENCODE 2025: reference gene annotation for human and mouse
- PubMed. Nucleic acids research *53*. 10.1093/nar/gkae1078.

60. VA, S., T, G.-L., K, H., N, B., HC, C., PA, K., TD, M., KD, P., F, T.-N., D, A., et al.
(2017 May). Evaluation of GRCh38 and de novo haploid genome assemblies
demonstrates the enduring quality of the reference assembly - PubMed. Genome
research *27*. 10.1101/gr.213611.116.

61. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq
aligner. Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.

62. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. Genome Biology *15*, 550.
10.1186/s13059-014-0550-8.

63.   Gaujoux, R., Seoighe, C., Gaujoux, R., and Seoighe, C. (2010-07-02). A flexible R
package for nonnegative matrix factorization. BMC Bioinformatics 2010 11:1 *11*.
10.1186/1471-2105-11-367.
64.   Kim, H., and Park, H. (2007/06/15). Sparse non-negative matrix factorizations via
alternating non-negativity-constrained least squares for microarray data analysis.
Bioinformatics *23*. 10.1093/bioinformatics/btm134.
65.   Mootha, V.K., Lindgren, C.M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar,
J., Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-
1α-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. Nature Genetics *34*, 267-273. 10.1038/ng1180.
66.   A, S., P, T., VK, M., S, M., BL, E., MA, G., A, P., SL, P., TR, G., ES, L., and JP, M.
(10/25/2005). Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles - PubMed. Proceedings of the
National Academy of Sciences of the United States of America *102*.
10.1073/pnas.0506580102.
67.   Xu, Y., Nipper, M.H., Dominguez, A.A., Ye, Z., Akanuma, N., Lopez, K., Deng, J.J.,
Arenas, D., Sanchez, A., Sharkey, F.E., et al. (2024-01-27). Reconstitution of
human PDAC using primary cells reveals oncogenic transcriptomic features at
tumor onset. Nature Communications 2024 15:1 *15*. 10.1038/s41467-024-45097-
2.
68.   DA, B., P, T., JS, B., SY, K., SE, M., IF, D., AC, S., P, S., E, M., C, S., et al.
(11/05/2009). Systematic RNA interference reveals that oncogenic KRAS-driven
cancers require TBK1 - PubMed. Nature *462*. 10.1038/nature08460.
69.   Tosti, L., Hang, Y., Debnath, O., Tiesmeyer, S., Trefzer, T., Steiger, K., Ten, F.W.,
Lukassen, S., Ballke, S., Kühl, A.A., et al. (2021). Single-Nucleus and In Situ RNA–
Sequencing Reveal Cell Topographies in the Human Pancreas. Gastroenterology
160, 1330-1344.e1311. 10.1053/j.gastro.2020.11.010.
70.   Mastromatteo, S., Chen, A., Gong, J., Lin, F., Thiruvahindrapuram, B., Sung,
W.W., Whitney, J., Wang, Z., Patel, R.V., Keenan, K., et al. (2022 Oct 20). High-
quality read-based phasing of cystic fibrosis cohort informs genetic understanding
of disease modification. Human Genetics and Genomics Advances *4*.
10.1016/j.xhgg.2022.100156.
71.   Lou, H., Xie, B., Wang, Y., Gao, Y., and Xu, S. (2023-01-01). Improved NGS
variant calling tool for the PRSS1–PRSS2 locus. Gut *72*. 10.1136/gutjnl-2022-
327203.
72.   Surendran, R.P., Visser, M.E., Heemelaar, S., Wang, J., Peter, J., Defesche, J.C.,
Kuivenhoven, J.A., Hosseini, M., Péterfy, M., Kastelein, J.J.P., et al. (2012/08/01).
Journal of Internal Medicine | Wiley Online Library. Journal of Internal
Medicine *272*. 10.1111/j.1365-2796.2012.02516.x.
73.   Hang, S.M., Hegele, R.A., and Berberich, A.J. (2023 Feb 1). Genetic determinants
of pancreatitis: relevance in severe hypertriglyceridemia. Current Opinion in
Lipidology *34*. 10.1097/MOL.0000000000000866.
74.   Cuppens, H., Lin, W., Jaspers, M., Costes, B., Teng, H., Vankeerberghen, A.,
Jorissen, M., Droogmans, G., Reynaert, I., Goossens, M., et al. (1998/01/15).
Polyvariant mutant cystic fibrosis transmembrane conductance regulator genes.
The polymorphic (Tg)m locus explains the partial penetrance of the T5

polymorphism as a disease mutation. The Journal of Clinical Investigation *101*.
10.1172/JCI639.

75. JH, L., JH, C., W, N., JW, H., J, C., SY, S., SW, P., DS, K., JH, Y., Y, S., et al.
(09/15/2003). A haplotype-based molecular analysis of CFTR mutations
associated with respiratory and pancreatic diseases - PubMed. Human molecular
genetics *12*. 10.1093/hmg/ddg243.

76. A, S., J, L., X, S., A, A., J, L., R, H., P, C., RE, B., MA, A., ME, M., et al. (2011
Jan). Combined bicarbonate conductance-impairing variants in CFTR and SPINK1
variants are associated with chronic pancreatitis in patients without cystic fibrosis
- PubMed. Gastroenterology *140*. 10.1053/j.gastro.2010.10.045.

77. J, L., J, J., IJ, G., MD, L., HW, P., RE, B., A, G., MA, A., PA, B., D, C., et al.
(07/17/2014). Mechanisms of CFTR functional variants that impair regulated
bicarbonate permeation and increase risk for pancreatitis but not for cystic fibrosis
- PubMed. PLoS genetics *10*. 10.1371/journal.pgen.1004376.

78. Haggie, P.M., Phuan, P.-W., Tan, J.-A., Xu, H., Avramescu, R.G., Perdomo, D.,
Zlock, L., Nielson, D.W., Finkbeiner, W.E., Lukacs, G.L., and Verkman, A.S. (2016
Nov 28). Correctors and Potentiators Rescue Function of the Truncated W1282X-
Cystic Fibrosis Transmembrane Regulator (CFTR) Translation Product. The
Journal of Biological Chemistry *292*. 10.1074/jbc.M116.764720.

79. van Willigen, M., Vonk, A.M., Yeoh, H.Y., Kruisselbrink, E., Kleizen, B., van der
Ent, C.K., Egmond, M.R., de Jonge, H.R., Braakman, I.A.-O., Beekman, J.M., and
van der Sluijs, P.A.-O. (2019). Folding-function relationship of the most common
cystic fibrosis-causing CFTR conductance mutants. LID - 10.26508/lsa.201800172
[doi] LID - e201800172.

80. Modiano, G., Bombieri, C., Ciminelli, B.M., Belpinati, F., Giorgi, S., Georges, M.d.,
Scotet, V., Pompei, F., Ciccacci, C., Guittard, C., et al. (2004-11-10). A large-scale
study of the random variability of a coding sequence: a study on the CFTR gene.
European Journal of Human Genetics 2005 13:2 *13*. 10.1038/sj.ejhg.5201306.

81. BM, C., A, B., C, B., F, P., M, G., C, C., A, B., A, H., P, Z., T, P., et al. (2007 Jan).
Highly preferential association of NonF508del CF mutations with the M470 allele -
PubMed. Journal of cystic fibrosis : official journal of the European Cystic Fibrosis
Society *6*. 10.1016/j.jcf.2006.04.003.

82. Zielenski, J., Patrizio, P., Corey, M., Handelin, B., Markiewicz, D., Asch, R., and
Tsui, L.C. (1995). CFTR gene variant for patients with congenital absence of vas
deferens. Am J Hum Genet *57*, 958-960.

83. D, Z., R, B., and L, W. (2020 Jan). The Cystic Fibrosis Transmembrane
Conductance Regulator 470 Met Allele Is Associated with an Increased Risk of
Chronic Pancreatitis in Both Asian and Caucasian Populations: A Meta-Analysis -
PubMed. Genetic testing and molecular biomarkers *24*. 10.1089/gtmb.2019.0199.

84. Iso, M., Suzuki, M., Yanagi, K., Minowa, K., Sakurai, Y., Nakano, S., Satou, K.,
Shimizu, T., Kaname, T., Iso, M., et al. (2019-04-11). The CFTR gene variants in
Japanese children with idiopathic pancreatitis. Human Genome Variation 2019 6:1
6. 10.1038/s41439-019-0049-7.

85. Angyal, D., Kleinfelder, K., Ciciriello, F., Groeneweg, T.A., Marchi, G.D., Pretis,
N.d., Bernardoni, L., Rodella, L., Tomba, F., Angelis, P.D., et al. (2024). CFTR

function is impaired in a subset of patients with pancreatitis carrying rare CFTR
variants. Pancreatology *24*, 394-403. 10.1016/j.pan.2024.03.005.

86. El-Seedy, A., Girodon, E., Norez, C., Pajaud, J., Pasquet, M.-C., Becdelièvre, A.d., Bienvenu, T., Georges, M.d., Cabet, F., Lalau, G., et al. (2012/11/01). CFTR mutation combinations producing frequent complex alleles with different clinical and functional outcomes. Human Mutation *33*. 10.1002/humu.22129.

87. Doss, C.G.P., Rajasekaran, R., Sudandiradoss, C., Ramanathan, K., Purohit, R., and Sethumadhavan, R. (2008 May 14). A novel computational and structural analysis of nsSNPs in CFTR gene. Genomic Medicine *2*. 10.1007/s11568-008-9019-8.

88. Meléndez-Montañez, J.M., Jesús-Rojas, W.D., Meléndez-Montañez, J.M., and De Jesús-Rojas, W. (2023-01-31). CFTR-Related Metabolic Syndrome: Genetic Variants Increasing Pancreatitis Risk in the Pediatric Puerto Rican Population. Children 2023, Vol. 10, Page 280 *10*. 10.3390/children10020280.

89. S, R., I, V., A, M., S, B., J, R., RM, C., and SM, R. (2016 Sep-Oct). Chronic pancreatitis: Do serum biomarkers provide an association with an inflammageing phenotype? - PubMed. Pancreatology : official journal of the International Association of Pancreatology (IAP) ... [et al.] *16*. 10.1016/j.pan.2016.08.004.

90. Engle, D.D., Tiriac, H., Rivera, K.D., Pommier, A., Whalen, S., Oni, T.E., Alagesan, B., Lee, E.J., Yao, M.A., Lucito, M.S., et al. (2019). The glycan CA19-9 promotes pancreatitis and pancreatic cancer in mice. Science *364*, 1156-1162. 10.1126/science.aaw3145.

91. CM, A., BM, G., KK, T., C, L.-M., N, T., PK, M., KE, D., ES, C., CJ, H., JC, H., et al. (09/11/2012). EGF receptor is required for KRAS-induced pancreatic tumorigenesis - PubMed. Cancer cell *22*. 10.1016/j.ccr.2012.07.024.

92. Cheng, S.H., Gregory, R.J., Marshall, J., Paul, S., Souza, D.W., White, G.A., O'Riordan, C.R., and Smith, A.E. (1990/11/16). Defective intracellular transport and processing of CFTR is the molecular basis of most cystic fibrosis. Cell *63*. 10.1016/0092-8674(90)90148-8.

93. Wilson, J.S., Pirola, R.C., and Apte, M.V. (2018). Epidemiology and Etiology of Alcohol-Induced Pancreatitis. The Pancreas. 10.1002/9781119188421.ch13.

94. Maléth, J., Balázs, A., Pallagi, P., Balla, Z., Kui, B., Katona, M., Judák, L., Németh, I., Kemény, L.V., Rakonczay, Z., et al. (2015). Alcohol Disrupts Levels and Function of the Cystic Fibrosis Transmembrane Conductance Regulator to Promote Development of Pancreatitis. Gastroenterology *148*, 427-439.e416. 10.1053/j.gastro.2014.11.002.

95. MY, P., and ZM, S. (2022 Jun). Current clinical opinion on CFTR dysfunction and patient risk of pancreatitis: diagnostic and therapeutic considerations - PubMed. Expert review of gastroenterology & hepatology *16*. 10.1080/17474124.2022.2084072.

96. Gullo, L., Pezzilli, R., Bellanova, B., D'Ambrosi, A., Alvisi, V., and Barbara, L. (1991). Influence of the thyroid on exocrine pancreatic function. Gastroenterology *100*, 1392-1396.

97. Komar, H.M., Hart, P.A., Cruz-Monserrate, Z., Conwell, D.L., and Lesinski, G.B. (2017). Local and Systemic Expression of Immunomodulatory Factors in Chronic Pancreatitis. Pancreas *46*, 986-993. 10.1097/mpa.0000000000000896.

98. Yasuda, M., Ito, T., Oono, T., Kawabe, K., Kaku, T., Igarashi, H., Nakamura, T.,
and Takayanagi, R. (2008 Nov 14). Fractalkine and TGF-β1 levels reflect the
severity of chronic pancreatitis in humans. World Journal of Gastroenterology :
WJG *14*. 10.3748/wjg.14.6488.
99. Bedwell, D.M., Kaenjak, A., Benos, D.J., Bebok, Z., Bubien, J.K., Hong, J.,
Tousson, A., Clancy, J.P., Sorscher, E.J., Bedwell, D.M., et al. (1997/11).
Suppression of a CFTR premature stop mutation in a bronchial epithelial cell line.
Nature Medicine 1997 3:11 *3*. 10.1038/nm1197-1280.
100. Mourik, P.V., Haaren, P.v., Kruisselbrink, E., Korkmaz, C., Janssens, H.M., Groot,
K.M.d.W.d., Ent, C.K.v.d., Hagemeijer, M.C., and Beekman, J.M. (2020). R117H-
CFTR function and response to VX-770 correlate with mRNA and protein
expression in intestinal organoids. Journal of Cystic Fibrosis *19*, 728-732.
10.1016/j.jcf.2020.02.001.
101. J, R., C, P.-S., IAI, S., C, M., S, A., R, F., and G, O. (05/21/2022). New drugs in
cystic fibrosis: what has changed in the last decade? - PubMed. Therapeutic
advances in chronic disease *13*. 10.1177/20406223221098136.
102. DM, C., NL, Q., ML, F., CR, E., J, D., NV, D., SH, R., RC, B., and M, G.
(07/23/2014). Potentiator ivacaftor abrogates pharmacological correction of ΔF508
CFTR in cystic fibrosis - PubMed. Science translational medicine *6*.
10.1126/scitranslmed.3008680.
103. Carpenter, E.S., Elhossiny, A.M., Kadiyala, P., Li, J., McGue, J., Griffith, B.D.,
Zhang, Y., Edwards, J., Nelson, S., Lima, F., et al. (2023/06/01). Analysis of Donor
Pancreata Defines the Transcriptomic Signature and Microenvironment of Early
Neoplastic Lesions. Cancer Discovery *13*. 10.1158/2159-8290.CD-23-0013.
104. Mello, S.S., Flowers, B.M., Mazur, P.K., Lee, J.J., Müller, F., Denny, S.K., Ferreira,
S., Hanson, K., Kim, S.K., Greenleaf, W.J., et al. (2023-2-27). Multifaceted role for
p53 in pancreatic cancer suppression. Proceedings of the National Academy of
Sciences *120*. 10.1073/pnas.2211937120.
1106
1107

Tables

1109

1110   **Table 1.** Characteristics of patients (n=51).

Demographics	PDO established (n=36)	PDO not established (n=12)/ in progress (n=3)
Age, y, average (SD)	37(18%)	41 (17%)/ 37(21%)
Female sex, n (%)	24 (66%)	7 (58%)/ 2 (66%)
Male sex, n (%)	12 (33%)	5 (42%)/ 1 (33%)
Race		
White, n (%)	28 (77%)	7 (58%) / 1 (33%)
Hispanic, n (%)	4 (11%)	1 (8.3%) / 2 (66%)
Black, n (%)	1 (2.7%)	1 (8.3%)
Multi-racial, n (%)	1 (2.7%)	2 (16.6%)
Not reported	2 (5.5%)	0 (0%)
Procedure		
Partial Pancreatectomy (total =16), n (%)	9 (25%)	7 (58%)/ 0 (0%)
TPIAT (total = 35), n (%)	27 (75%)	5 (42%)/ 3 (100%)
Etiology		
Genetic	16 (44%)	2 (16.6%)/ 0(0%)
Idiopathic	12 (33%)	8 (66%)/ 1(33%)
Alcohol	3 (8.3%)	0(0%)/ 0(0%)
Obstruction	1 (2.7%)	0(0%)/ 0(0%)
Solid-pseudopapillary neoplasm	1 (2.7%)	0(0%)/ 0(0%)
Divisium + Genetic	2 (5.5%)	2 (16.6%) / 1(33%)
Groove	1 (2.7%)	0(0%)/ 0(0%)
Sphincter of Oddi dysfunction	0 (0%)	0(0%)/1(33%)
Mutations		
PRSS1	7 (19%)	2 (16%)/ 0(0%)
SPINK1	6 (17%)	0 (0%)/ 0(0%)
CFTR	10 (28%)	2 (17%)/ 0(0%)

1111

1112   **Table S1. Patient sample information and downstream sequencing technologies applied to patient derived organoids (PDOs).**

Sample	Patient	Sex	Age	Race	Type	Etiology	Procedure	PDO	Targeted sequencing	WES-PDO	WGS-PDO	WGS-PT	RNA seq
hCP01	1	F	ADT	W,NH	CP	I	PP	Y	Y	N	N	Y	Y
hCP02	2	F	ADT	W,NH	SPN	SPN	PP	Y	N	N	N	N	N
hCP03	3	F	ADT	W,NH	CP	I	PP	Y	Y	N	N	Y	Y
hCP04	3	F	ADT	W,NH	CP	I	PP	N	NA	NA	NA	N	NA
hCP05	4	F	ADT	W,NH	CP	I	PP	N	NA	NA	NA	N	NA
hCP06	5	F	ADT	W	RAP	G, D	TPIAT	N	NA	NA	NA	N	NA
hCP07	6	F	ADT	W,NH	CP	I	PP*	Y	N	N	Y	Y	Y
hCP08	7	F	PED	W,NH	RAP to CP	G	TPIAT	Y	Y	Y	N	Y	Y
hCP09	8	F	ADT	W,NH	CP	I	PP	Y	N	N	N	N	N
hCP10	9	M	ADT	W,NH	CP	I	PP	N	NA	NA	NA	N	NA
hCP11	10	F	ADT	W,NH	RAP to CP	G	TPIAT	Y	Y	Y	N	Y	Y
hCP12	11	F	PED	W,NH	RAP to CP	G	TPIAT	Y	N	N	N	N	N
hCP13	12	F	ADT	W,NH	RAP to CP	G	TPIAT	Y	N	N	N	N	N
hCP14	13	M	PED	W,NH	RAP to CP	G	TPIAT	N	NA	NA	NA	N	NA
hCP15	14	M	ADT	W,NH	RAP to CP	A	TPIAT	Y	Y	Y	N	Y	Y
hCP16	15	F	ADT	W,NH	RAP to CP	O	TPIAT	Y	Y	N	Y	Y	Y
hCP17	16	F	PED	W,NH	RAP to CP	G	TPIAT	Y	Y	Y	N	Y	Y
hCP18	17	F	ADT	W,NH	CP	I	TPIAT	Y	N	N	Y	Y	Y
hCP19	18	F	ADT	Multi	CP	I	PP	N	NA	NA	NA	N	NA
hCP20	19	F	PED	W,NH	RAP to CP	I	TPIAT	Y	Y	Y	N	Y	Y

hCP21	20	M	ADT	W,NH	RAP to early CP	D	TPIAT	Y	N	N	N	N	N
hCP22	21	F	ADT	W,NH	CP	I	PP	Y	N	N	N	N	N
hCP23	22	F	PED	W,NH	RAP to CP	G	TPIAT	Y	N	N	N	N	N
hCP24	23	F	PED	W,NH	RAP	G	TPIAT	Y	N	N	Y	Y	Y
hCP25	24	F	PED	H	RAP to CP	I	TPIAT	Y	N	N	Y	Y	Y
hCP26	25	M	ADT	W,NH	CP	I	PP	Y	Y	N	N	Y	Y
hCP27	25	M	ADT	W,NH	CP	I	PP*	Y	N	N	Y	Y	Y
hCP28	26	F	ADT	H	RAP to CP	G	TPIAT	Y	N	N	Y	Y	Y
hCP29	27	M	ADT	W,NH	RAP to CP	A	TPIAT	Y	N	N	Y	Y	Y
hCP30	28	M	ADT	W,NH	RAP to early CP	G	TPIAT	Y	N	N	Y	Y	Y
hCP31	29	M	ADT	Multi	CP	I	PP	N	NA	NA	NA	N	NA
hCP32	30	M	ADT	W,NH	RAP to CP	G	TPIAT	Y	N	N	Y	Y	Y
hCP33	31	F	ADT	W	RAP to CP	I	PP	N	NA	NA	NA	N	NA
hCP34	32	M	ADT	W,NH	RAP to CP	I	TPIAT	N	NA	NA	NA	N	NA
hCP35	33	M	ADT	H	RAP to CP	G	TPIAT	Y	Y	N	N	Y	Y
hCP36	34	F	ADT	W,NH	RAP to CP	G	TPIAT	Y	N	N	Y	Y	Y
hCP37	35	F	PED	H	RAP to CP	G	TPIAT	N	NA	NA	NA	N	NA
hCP38	36	F	ADT	B	RAP to CP	Groove	TPIAT	Y	N	N	N	N	N
hCP39	37	M	ADT	B	CP	I	PP	N	NA	NA	NA	N	NA
hCP40	38	F	ADT	W,NH	RAP to CP	G	TPIAT	Y	N	N	N	N	N

hCP41	39	F	ADT	W,NH	RAP to CP	I	TPIAT	Y	N	N	N	N	N
hCP42	40	M	ADT	W,NH	RAP to CP	G	TPIAT	Y	N	N	N	N	N
hCP43	41	F	ADT	W,NH	RAP to CP	A	TPIAT	Y	N	N	N	N	N
hCP44	42	M	ADT	W,NH	RAP to early CP	G	TPIAT	Y	Y	N	N	Y	Y
hCP45	43	F	ADT	W,NH	RAP to CP	D	TPIAT	N	NA	NA	NA	N	NA
hCP46	44	M	ADT	H	RAP to early CP	I	TPIAT	IP	NA	NA	NA	N	NA
hCP47	45	F	ADT	W,NH	CP	SOD	TPIAT	IP	NA	NA	NA	N	NA
hCP48	46	F	ADT	Multi	CP	I	PP	Y	N	N	Y	Y	Y
hCP49	47	M	ADT	W,NH	RAP to CP	D, VATER	TPIAT	Y	N	N	Y	Y	Y
hCP50	48	F	ADT	H	RAP to early CP	D	TPIAT	IP	NA	NA	NA	N	NA
hCP51	49	F	ADT	H	RAP to early CP	D	TPIAT	Y	N	N	Y	Y	Y
hNAT02	50	M	ADT	W,NH	normal	Distal to NET		Y	N	N	N	N	Y
hNAT04	51	M	ADT	W,NH	normal	Adj to PDAC		Y	N	N	N	N	Y
hNAT07	53	M	ADT	W,NH	normal	Adj to ampullary tumor		Y	N	N	N	N	Y
hNAT09	54	M	ADT	W,NH	normal	Adj to PDAC		Y	N	N	N	N	Y
hNP03 Body	55	F	ADT	W,NH	normal	COD: anoxia	Organ donation	Y	N	N	N	N	Y
hNP03 Head	55	F	ADT	W,NH	normal	COD: anoxia	Organ donation	Y	N	N	Y	N	Y
hNP03 Tail	55	F	ADT	W,NH	normal	COD: anoxia	Organ donation	Y	N	N	Y	N	Y

hNP05 Body	56	M	ADT	H	normal	COD: cerebrovascular stroke	Organ donation	Y	N	N	N	N	Y
hNP05 Head	56	M	ADT	H	normal	COD: cerebrovascular stroke	Organ donation	Y	N	N	Y	N	Y
hNP05 Tail	56	M	ADT	H	normal	COD: cerebrovascular stroke	Organ donation	Y	N	N	Y	N	Y
hNP16 Body	57	M	ADT	W,NH	normal	COD: cerebrovascular stroke	Organ donation	Y	Y	N	N	N	Y
hNP16 Head	57	M	ADT	W,NH	normal	COD: cerebrovascular stroke	Organ donation	N	N	N	Y	N	Y
hNP16 Tail	57	M	ADT	W,NH	normal	COD: cerebrovascular stroke	Organ donation	Y	N	N	Y	N	Y

Abbreviations: Race: W-White, NH- non-Hispanic, H- Hispanic, B- Black, Multi- multi-racial; Sex: M-Male, F-Female; Age: ADT- Adult,
PED- Pediatric Type: CP- chronic pancreatitis, SPN- Solid-pseudopapillary neoplasm, RAP- recurrent acute pancreatitis; Etiology: I-
idiopathic, SPN- Solid-pseudopapillary neoplasm, G- genetic, D- divisum, A- alcohol, O- obstruction, SOD- Sphincter of Oddi
Dysfunction, VATER- ampulla of Vater, NET- neuroendocrine tumor, PDAC- pancreatic ductal adenocarcinoma, Adj- adjacent, COD-
cause of death; Procedure: PP- partial pancreatectomy, TPIAT- total pancreatectomy with islet auto transplantation; IP- in progress,
Y- Yes, N- No, NA- not applicable, *- from pancreatic juice.

Figure 1. Generation of organoids from patients with human chronic pancreatitis (CP) with high effciency.
(A) Representative human normal pancreas (hNP) section displays characteristic ductal, acinar, and islet cells. Masson's trichrome (MT) staining highlights low degree of fibrosis in NP sample and varying severity in human CP (hCP) samples in blue. NP sample has high levels of cytokeratin 19 staining (CK19) in ductal epithelial cells, while CP samples have an altered ductal cell compartment. Ki-67 expression is low in NP sample and is altered or completely lost in hCP samples. Scale bar 100 µm. For diagram outlining generation procedure and histological features of samples that did not produce organoids see Figure S1.
(B) Patient sample cohort demographic, etiology, procedure for acquisition (total pancreatectomy with auto islet transplantation (TPIAT) and partial pancreatectomy (PP), organoid generation, and pathologist fibrosis scoring (Fibrosis) information. For additional patient sample information see also Table S1, S2. Pancreas juice samples do not have histological sections for review and samples deemed histological failures could not be reviewed (N/A).
(C) Mutational profile of patient samples as annotated through patient medical history for which PDOs were generated. Mutations in cystic fibrosis transmembrane conductance regulator (CFTR), serine protease inhibitor kazal type 1 (SPINK1), and protease serine 1 (PRSS1) are present in our patient samples representing some of the most common alterations in CP. Mutation - a mutation was annotated in patient medical history but the specific alteration was not reported.
(D) Generation efficiency of patient derived organoids (PDOs); cumulative generation efficiency of 70.5%.
(E) Representative images of human organoid cultures established from normal (hNP) or hCP after at least 4 passages. Scale bar 500 µm.
(F) Whole mount immunofluorescence of PDOs stained with Hoechst (nuclei), EdU (pulsed for 25 hours), and CK19 (ductal marker). Scale bar 100 µm.
(G) Quantification of EdU positive nuclei in a panel of NP and CP PDOs across two or three independent experiments and multiple passages. For representative images of immunofluorescence see Figure S1D.

Figure 2. PDOs have high genetic concordance and reveal important and previously unannotated alterations.
CP PDO and primary tissue coding sequence variants in PRSS1, SPINK1, LPL and CFTR determined from either WES (whole exome sequencing), targeted genetic sequencing, or WGS (whole genome sequencing). Genotype matches reference or is annotated as a heterozygous (Het) or homozygous (Homo) alteration as compared to reference. Clinical annotations from medical notes annotated if not found in PDO or primary tissue. Concordance is annotated for each pair of primary tissue and PDO as match rate (%). For concordance between unmatched pairs of primary tissue and PDO or PDOs see Figure S4A. For cystic fibrosis associated CFTR mutations annotated in our cohort see Figure S3B.

Figure 3. Gene expression analyses demonstrate distinct cell signaling programs in CP organoids.

(A) Principal component analysis (PCA) of top 500 most variable genes from organoids isolated from NP, normal adjacent to tumor (NAT), and CP samples.

(B) Top 500 variable genes from variance stabilized gene expression in hCP , hNAT, and hNP samples. Hierarchical clustering separates NP and NAT from most CP organoids.

(C) Volcano plot of CP versus NP and NAT samples. (p value <0.05, |log2fold change| > 1 colored) . Top 10 enriched or depleted genes annotated.

(D) Gene set enrichment analysis of CP PDOs versus NP and NAT PDOs.

(E) Relative normalized transcript expression of select inflammatory, collagens, and TGFb genes from RNAseq. Sample disease origin and cluster are denoted. For more information on cell lineage genes see Figure S4.

(F) Clustering using non-matrix factorization (NMF) defines three distinct subtypes of hCP PDO cultures; basis 1, 2, and 3. Patient etiology (A- alcoholic, D- divisum + groove, G- genetic, I- idiopathic, O- obstruction), silhouette, and consensus is indicated. For NMF rank survey see Figure S4.

Figure 4. CP organoids recapitulate protein expression from primary tissue specimens and exhibit elevated inflammatory and mitogenic signaling.

(A) Immunoblot of PDOs generated from NP and CP samples. CFTR mutations annotated for CP samples. For normalized mature CFTR expression across cohort see Figure S7A.

(B) Immunohistochemistry (IHC) staining for cancer antigen 19-9 (CA19-9), signal transducer and activator of transcription 3 phosphorylation (pSTAT3), mitogen-activated protein kinase 1 phosphorylation (pERK2), and cystic fibrosis transmembrane conductance regulator (CFTR) of NP and CP samples. Scale bar 100 μm.

(C) Normalized secreted protein expression of inflammatory cytokines in NP and CP PDOs. Average concentration of two technical replicates per sample. P-value determined by Mann-Whitney test. Sample type and cluster is shown. For additional cytokines assayed see Figure S7B.

Figure 5. Forskolin induced swelling of CP PDOs is CFTR dependent.
(A) Schematic of forskolin induced swelling assay. Created in BioRender. Osorio Vasquez, V. (2025) https://BioRender.com/oahkr5p
(B) Representative brightfield images of organoids used for the quantification of forskolin induced swelling assay. Upto 10 organoids are selected per condition.
(C) Forskolin induced swelling (FIS) assay of hNP organoids with forskolin, CFTR potentiators (icenticaftor or ivacaftor), or a combination of both for 6 hours.
(D) Forskolin induced swelling (FIS) assay of hNP organoids treated with forskolin for 6 hours and pre-treated with CFTR inhibitors (CFTRi-172, PPQ-102), or protein kinase A inhibitor (H-89) for 2 hours. Swelling is inhibited with the addition of CFTR and PKA inhibitors.
(E) Forskolin induced swelling (FIS) assay of hNP organoids treated with ivacaftor for 6 hours and pre-treated with CFTR inhibitors (CFTRi-172, PPQ-102), or protein kinase A inhibitor (H-89) for 2 hours. Swelling is inhibited with the addition of CFTR and PKA inhibitors.
(F) Forskolin induced swelling (FIS) assay of hNP organoids treated with icenticaftor for 6 hours and pre-treated with CFTR inhibitors (CFTRi-172, PPQ-102), or protein kinase A inhibitor (H-89) for 2 hours. Swelling is inhibited with the addition of CFTR and PKA inhibitors.
(G) FIS assay of hCP organoids treated with forskolin for 6 hours. Data represented as swelling deficient or proficient compared to DMSO control at 6 hours. CFTR mutations are shown for each sample. Data represented as mean ± SD for C, D, E. Concentrations for forskolin treatments 20 μM, CFTR potentiators (ivacaftor, icenticaftor) or correctors (elexacaftor, lumacaftor) 10 μM for FIS assays. Data represented as swelling deficient or proficient compared to DMSO control of at least 2 independent experiments. ETI treatment concentrations are 3 μM for each compound.

Figure 6. CFTR function in CP organoids is restored upon pharmacologic modulation.

(A) Schematic of rescuing impaired forskolin induced swelling in PDOs. Created in BioRender. Osorio Vasquez, V. (2025) https://BioRender.com/a9tuea3

(B) hCP11 (CFTR W1282*) is FIS deficient upon treatment with forskolin or CFTR potentiators (icenticaftor or ivacaftor) for 6 hours.

(C) Treatment with CFTR readthrough compound (Ataluren- 10 μM) and 6 hour treatment with forskolin in hCP11 (CFTR W1282*) induces swelling.

(D) IB of mature CFTR expression upon 48 hour Ataluren (10 μM) treatment in hCP11 (CFTR W1282*). Quantification of IB directly adjacent.

(E) FIS is rescued upon 6 hour CFTR potentiator (Ivacaftor and Incenticaftor) and forskolin treatment in hCP35 PDOs. Data representative of at least 2 independent experiments.

(F) hCP44 (CFTR R347,R117H) organoids did not swell after 6 hours of treatment with DMSO, forskolin, CFTR potentiator (Ivacaftor), or CFTR corrector (lumacaftor).

(G) hCP44 (CFTR R347,R117H) organoids swelled in response to combination CFTR potentiator (Ivacaftor) and CFTR corrector (lumacaftor or elexacaftor) treatment after 6 hours.

(H) Mature CFTR expression is increased in hCP35 organoids treated with CFTR potentiator (Ivacaftor) or further increased with ETI (CFTR correctors-Elexacaftor and Tezacaftor, CFTR potentiator-Ivacaftor) for 48 hours. pERK2 and pSTAT3 expression decreases upon ETI treatment. Cofilin is used as loading control. Representative blot of 3. Quantification of IB relative to DMSO adjacent.

(I) Mature CFTR expression increases upon ETI treatment for 48 hours (CFTR correctors-Elexacaftor and Tezacaftor, CFTR potentiator-Ivacaftor) in hCP44 (CFTR R347,R117H) organoids. pERK1/2 and pSTAT3 is signaling is decreased upon ETI treatment. Cofilin is used as loading control. Quantification of IB relative to DMSO adjacent. Data represented as mean ± SD for C, D, E. Concentrations for forskolin treatments 20 μM, CFTR potentiators (ivacaftor, icenticaftor) or correctors (elexacaftor, lumacaftor) 10 μM for FIS assays. Data represented as swelling deficient or proficient compared to DMSO control of at least 2 independent experiments. ETI treatment concentrations are 3 μM for each compound.