Virulence factor candidalysin impairs epithelial barrier integrity and initiates inflammation in a gut-on-chip model

Moran Morelli[1*], Karla Queiroz[1]

[1]Mimetas B.V., Oegstgeest, The Netherlands

* Correspondence:
m.morelli@mimetas.com

Keywords: organ-on-a-chip; microfluidics; host–microbial interactions; organoids; drug
screening; microbiome; Candida albicans; candidiasis

Graphical Abstract

Abstract

Candida albicans is an opportunistic pathogenic yeast commonly found in the gastrointestinal tract,
vagina, and oral cavity of healthy humans. Under certain conditions, it can become invasive, causing
mucosal or life-threatening systemic infections. One mechanism used by *C.albicans* to breach the
epithelial barrier is the secretion of candidalysin, a cytolytic peptide toxin. Candidalysin damages
epithelial membranes but also activates the innate epithelial immune response, making it a key player
in the pathogenicity of *C.albicans* and a promising therapeutic target. Although candidalysin is
known to mediate transcellular translocation of *C. albicans* through intestinal layers, the epithelial
response to candidalysin remains an area of active research. This study aims to characterize this
response using gut-on-chip models with Caco-2 and colon organoid tubules. We used the
OrganoPlate, a microfluidic platform supporting the culture of up to 64 perfused, membrane-free
intestinal epithelial tubes. We exposed Caco-2 tubes to candidalysin and evaluated their response
with trans-epithelial electrical resistance (TEER), protein detection, and immunostaining. We then
validated our findings in a proof-of-concept experiment using human intestinal organoid tubules.
Candidalysin impaired barrier integrity, as indicated by decreased TEER and increased permeability
in a fluorescent dye assay. It also induced actin remodeling and DRAQ7 uptake, a marker of cell
permeability. This disruption was associated with the release of LDH, cytokines, and the

antimicrobial peptide LL37, suggesting cellular damage, inflammation, and antimicrobial activity.
This study strengthens our understanding of candidalysin's role in *C. albicans* pathogenesis and
suggests new therapeutic strategies targeting this toxin. Moreover, the use of patient-derived
organoids in toxin response studies shows promise for capturing patient heterogeneity and
developing personalized treatments.

1   Introduction

The intestinal mucosa is the first line of defense against microbial invasion. Intestinal epithelial cells
(IECs) form a barrier, which separates the sterile host environment from the gut microbiota.
Perturbations of the host-microbial interplay such as an imbalance in the microbial community, a
breach in the barrier, or an impairment of the host's immune system can lead to disease[1].

Candida species are among the leading fungal pathogens, significantly contributing to morbidity and
mortality[2]. Among these, *C. albicans* remains the most prevalent cause of life-threatening systemic
candidiasis and is a major contributor to nosocomial infections, particularly in Intensive Care Units
(ICUs)[2,3]. *C. albicans* is an opportunistic pathogen that frequently inhabits the mucosal surfaces of
humans, with its main reservoir being the gastrointestinal tract[4–7]. A compromised immune system
and the use of broad-spectrum antibiotics can induce the transition of *C. albicans* from
commensalism to pathogenicity, where it can breach the intestinal barrier and reach the bloodstream
to cause systemic candidiasis[8–10]. One of the key mechanisms mediating the translocation of *C.*
albicans through the intestinal layers is the release of the virulence factor candidalysin[11,12].

Candidalysin is a cytolytic peptide playing a dual role in *C. albicans* infection. On one hand, it acts
as a virulence factor by directly damaging the host's epithelium through the formation of membrane
pores[11,13,14]. On the other hand, candidalysin acts as an immunomodulatory molecule recognized by
the host to initiate an immune response through p38 and EGFR-ERK signaling[11,15,16]. Characterizing
the epithelial response to candidalysin is therefore of critical importance to better manage
candidiasis.

So far, most of the *in vitro* research exploring the mechanisms of candidalysin has been done in oral
epithelial cell lines, where candidalysin has been shown to trigger the release of cytokines,
chemokines, alarmins, and antimicrobial peptides[16–19]; to induce epithelial stress such as
mitochondrial dysfunction, ATP depletion, and epithelial necrosis[20]; and to trigger Ca^{2+} influx and
breakdown of F-actin[21]. Nevertheless, although it is known that the main reservoir of *C. albicans* is
the gut[4,22], not many studies explored the effect of candidalysin on intestinal epithelial cells (IECs).

Allert and colleagues investigated the translocation process of *C. albicans* in IECs *in vitro* by
screening for >2000 mutants with deleted genes coding for translocation and cellular damage. They
showed that the mutant without candidalysin displayed moderate loss of barrier integrity, caused
almost no damage, and exhibited a reduced translocation in the Caco-2 brush border subclone
C2BBe1, indicating that the toxin is essential for translocation of the fungus through the intestinal
wall[10].

A follow up study investigated the mechanisms of *C. albicans* translocation across the intestinal
epithelial barrier focusing on fungal and host activities involved in the process. They found that
intestinal translocation was promoted by acquisition of host-cell zinc by *C. albicans* while at the
same time host NFκB signaling was protecting epithelial integrity, mitigating candidalysin-induced
damage and limiting fungal translocation[23].

Here, we used a membrane-free gut-on-chip model to evaluate the intestinal response to candidalysin in the cell line Caco-2. We characterized the response to the toxin by studying the barrier integrity, morphology, permeability, and release of epithelial mediators at the luminal side of the model. To validate our findings in a more relevant gut-on-chip model, we included a proof-of-concept experiment using patient-derived colon organoids to study barrier integrity, morphology, and permeability after candidalysin exposure.

In summary, our study presents a novel platform and readouts for evaluating the epithelial response of intestinal epithelial cells to candidalysin with the potential to run phenotypic screenings to identify therapeutic molecules. Our system can further evolve to incorporate other relevant cell types and create a more comprehensive and physiologically relevant model for personalized drug screening and the identification of drugs targeting microbial virulence factors.

2   Materials and Methods

2.1   Cell Culture

2.1.1 OrganoReady Colon Caco-2

OrganoReady Colon Caco-2 3-lane 40 (MI-OR-CC-01, MIMETAS B.V.) were cultured according to the manufacturer's instructions (Figure 1). Medium was replaced with Caco-2 medium on the day of receiving. OrganoReady Colon Caco-2 3-lane 40 are ready to use Caco-2 tubes in OrganoPlates that follow a similar process than what was described by Triestsch and colleagues[24]. Perfusion flow was maintained by placing the plate on OrganoFlow rocker (Mimetas B.V., MI-OFPR-L) set at 7 degrees with 8-minute intervals optimized for the 3-lane 40. On the second day after receiving (day 6 after seeding), Caco-2 medium was refreshed. Exposures were performed on day 4 after receiving (day 8 after seeding).

Figure 1

Figure 1. OrganoReady Caco-2 model. (A) Picture of an OrganoPlate 3-Lane 40 with one microfluidic chip enlarged. (**B**) Illustration of a microfluidic chip. Apical channel can be accessed through A1 and A3. Middle channel can be accessed through B1 and B3. Basal channel can be accessed through C1 and C3. (**C**) Schematic of a Caco-2 tubule (apical channel) against a collagen ECM (middle channel). Basal channel contains culture medium. Arrows indicate flow direction. (**D**) Phase-contrast image of a Caco-2 tubule in the OrganoPLate 3-lane 40, scale bar is 100 µM. (**E**) OrganoPlate on an OrganoFlow rocker. The rocker angle is set at 7 degrees and oscillates every 8 minutes. (**F**) Illustration of bi-directional flow in the OrganoPlate.

2.1.2 OrganoReady Colon Organoid

OrganoReady colon organoid plates (OrganoReady Colon Organoid 3-lane 64, MI-OR-CORG-02, MIMETAS B.V.) were used to study the effect of candidalysin in colon organoids. The plates consisted of 64 chips containing colon organoid tubules and were cultured with OrganoMedium Colon Recovery and Culture A and B according to manufacturer's instructions. Perfusion flow was maintained by placing the plate on OrganoFlow rocker (Mimetas B.V., MI-OFPR-L) set at 14 degrees with 8-minute intervals optimized for the 3-lane 64. The biopsy used to generate the tubules comes from the healthy portion of a sigmoid biopsy of a 58-years-old caucasian female with colorectal cancer.

2.2 Candidalysin Exposure

Candidalysin (SIIGIIMGILGNIPQVIQIIMSIVKAFKGNK) was synthesized by Peptide Protein Research Ltd (UK) and reconstituted in sterile purified water to 5 mg/mL (1.5 mM) for storage at -20°C, prior to further dilution for individual experiments.

For exposures in Caco-2 tubules, medium was replaced with serum-free medium in all channels 24h
prior to exposure. Then, the medium of the apical channel was replaced with serum-free medium
containing candidalysin at the specified concentration.

Due to their increased sensitivity and the limitation of modifying their culture medium, colon
organoids tubules were not serum starved. Instead, all channels were washed with HBSS (Hank's
Balanced Salt Solution) prior to adding HBSS containing the specified concentration of candidalysin
in the apical channel.

2.3   Protein Detection

Supernatant were collected from the apical channel and centrifuged for 10 min at 1500 rpm prior to
aliquoting and storage at -80 °C.

LL37 ELISA kits were purchased from Elabscience (UK) and performed according to the
manufacturer's instructions. Absorbance was measured using a Spark Cyto plate reader (TECAN,
Switzerland).

G-CSF, GM-CSF, IL-1 alpha, IL-1 beta, IL-6, IL-8, IP-10, MCP-1 (CCL2), MIP-3 alpha (CCL20),
S100A8/A9 were measured using a Luminex MAGPIX instrument (Thermo, USA) and a custom
Procartaplex assay kit (Thermo, USA), following the manufacturer's instructions. Plates were
measured using a Luminex MAGPIX® instrument and the Luminex xPONENT® software (version
4.3). ProcartaPlex Analysis App was used to determine analyte concentrations.

2.4   Trans-Epithelial-Electrical-Resistance (TEER)

Trans-epithelial electrical resistance (TEER) was measured using an automated multichannel
impedance spectrometer (OrganoTEER, Figure 2A-C) to evaluate the integrity of the gut barrier[25].
An electrode board, designed to fit the 3-lane OrganoPlate was sterilized with 70% ethanol at least 1
hour before measurement. The OrganoPlate was taken out of the incubator and equilibrated at room
temperature for 30 min prior to the measurement to eliminate any temperature or flow effect.
Baseline measurement was performed right after equilibration, then TEER was further measured at
the indicated timepoints after candidalysin exposure.

For Caco-2 the OrganoTEER was set on "high TEER" and we used a TEER threshold of 350 Ω/cm^2
at baseline, meaning that all tubules below 350 Ω/cm^2 before exposure were excluded from the
analysis. For colon organoids the OrganoTEER was set on "medium TEER" and we used a TEER
threshold of 100 Ω/cm^2 at baseline, meaning that all tubules below 100 Ω/cm^2 before exposure were
excluded from the analysis.

2.5   Barrier Integrity Assay

To further investigate the barrier integrity, leakage of fluorescent molecules from the lumen to the
ECM compartment was evaluated after exposure (Figure 3A-C). First, medium from all inlet and
outlets was removed. Then 25 µL serum-free medium was added to the inlet and outlets of the middle
and basal channel. Next, a dye solution consisting of Sodium fluorescein (0.4kDa, 10µg/ml, Sigma-
Aldrich, USA) and TRITC-dextran (4.4kDa, 500µg/ml, Sigma-Aldrich, USA) in serum-free medium
was added to the inlet and outlet of the apical channel (40 and 30 µL, respectively). Then, the
OrganoPlate was placed in an ImageXpress XLS Micro (Molecular Devices, USA) and imaged every

minutes for 7 timepoints. Leakage from the lumen into the ECM compartment was quantified
according to the protocol described by Soragni et al.[26]

2.6   Cell Damage (LDH) Assay

Lactate dehydrogenase (LDH) activity was measured using the LDH-Glo™ Cytotoxicity Assay kit
(Promega, USA), following the manufacturer's instructions. Supernatant were collected from the
lumen and centrifuged for 10 min at 1500 rpm, prior to dilution in LDH assay buffer (1:100) and
storage at -80 °C. Luminescence was measured in 96W half area white microplates (Greiner bio-one,
Austria) using a Spark Cyto plate reader (TECAN, Switzerland).

2.7   Immunohistochemistry

For visual characterization of the effect of candidalysin on actin cytoskeleton and cell permeability,
tubules were directly fixed and stained on the OrganoPlate based on the protocol described by
Trietsch et al.[24]

2.7.1 DRAQ7 staining

Before fixation, medium in the tubule inlets and outlets was replaced by 20μL of DRAQ7 dye
(Biostatus, DR71000) diluted 1:100 in serum-free EMEM (Caco-2) or HBSS (colon organoids), and
cells were incubated during 30 min under continuous perfusion inside the incubator. OrganoPlates
were then fixed and stained following the previously explained protocol.

2.7.2 Fixation

Intestinal tubules were fixed with 3.7% formaldehyde (Sigma, 252,549) in HBSS with Calcium and
Magnesium (Thermo Scientific, 14,025,092) for 15 min, washed twice with phosphate-buffered
saline (PBS; Gibco, 70,013,065) for 5 min and then stored with 50μL PBS per well at 4 °C until
further staining.

2.7.3 Actin and Nuclear Staining

Intestinal tubules cells were permeabilized with 0.03% Triton X-100 (Sigma, T8787) in PBS for 10
min and washed twice with 4% FBS in PBS solution. Actin and nuclear stainings were performed
using the direct stains ActinGreen 488 ReadyProbes Reagent (Invitrogen, R37110) and NucBlue
Fixed Cell ReadyProbes Reagent (Invitrogen, R37606). Direct staining was performed following the
manufacturer's instructions and under constant flow.

For ActinGreen™ and NucBlue™ staining, two drops/ml were added in PBS and 20μl of the staining
solution was added to the tube inlet and outlet (20μL each) and incubated for 30min at room
temperature under continuous perfusion. Plates were then washed twice for 5 minutes with PBS and
the bottom of the tubules was imaged as max projection using an ImageXpress Micro Confocal
microscope (Molecular Devices, USA).

2.8   Intestinal tubules visualization and imaging

Tubules were imaged using the ImageXpress Micro XLS (Molecular Devices) and Micro XLS-C
High Content Imaging Systems (Molecular Devices) and processed using Fiji34 to enhance contrast
and improve visualization. To monitor the integrity of the tubules, phase-contrast images were
recorded before and after exposure to candidalysin. This was routinely performed immediately after
measurement of baseline TEER and after the required exposure time. Fixed and stained OrganoPlates

189 were stored at 4 °C until imaging and equilibrated at room temperature at least 30 min before
190 imaging. Maximum intensity projection images were saved as TIFF files after confocal imaging of
191 stained cells.

2.9 Staining quantification

Open-source cell image analysis software CellProfiler35 (version 4.2.5) was used to process visual
immunofluorescence images. An image analysis pipeline was designed to quantify the number and
area of actin clumps and the number of cells, by segmentation of the actin clumps and nuclei and
measurement of the total actin clump area. This pipeline was used to process the TIFF files that
captured actin staining or nuclei staining, correspondingly. Parallelly, another pipeline was designed
to process the TIFF files capturing DRAQ7 and nuclei staining, that allowed segmentation of
DRAQ7-positive cells and nuclei, to quantify the percentage of DRAQ7-positive (DRAQ7 +) cells.

2.10 Statistics

Data analysis and visualization were conducted using R version 4.2.2 and RStudio version
2023.03.0-386 (details on the software packages used are available in the supplementary material).

All experiments were performed three times in an independent manner (N = 3), and the number of
technical replicates per concentration (n) is indicated in the caption of each figure. Normality of data
distribution and homogeneity of variances were assessed using the Shapiro-Wilk test and Levene's
test, respectively. Differences among two groups were compared using independent T-test or
Wilcoxon Rank-Sum Test (Mann-Whitney U Test). Differences among three groups or more were
compared using one-way ANOVA followed by post-hoc Tukey's HSD, or Kruskal–Wallis tests
followed by Dunn's test, to obtain pairwise comparisons between the effect of exposure to the
various candidalysin concentrations and exposure. P values were adjusted for multiple comparisons
using Bonferroni method. Adjusted P values of 5% or lower were considered statistically significant
and were reported with asterisks: $*p \leq 0.05$; $**p \leq 0.01$; $***p \leq 0.001$, $****p \leq 0.0001$. P values
above 5% were considered not statistically significant (ns) and were not reported.

3 Results

3.1 Candidalysin impairs barrier integrity of Caco-2 tubules and increases their permeability

To determine candidalysin-mediated effects on the intestinal barrier integrity, we used the
OrganoTEER (Figure 2A-C) to measure TEER of the Caco-2 tubules after 30min, 1h, and 2h
candidalysin exposure (37 and 75µM). Both concentrations significantly decreased TEER after all
timepoints compared to vehicle (Figure 2C, D). When comparing the effect over time, although 37
µM candidalysin exposure showed a trend of recovering TEER over time, and 75 µM candidalysin
showed a trend of decreasing TEER over time, these changes were not statistically significant (p >
0.05) (Figure 2C, D).

Figure 2

Figure 2. TEER measurements of Caco-2 tubules exposed to candidalysin. (**A**) Photograph of the OrganoTEER setup: the plate holder (**1**) holds the OrganoPlate (**2**) in which the electrode board (**3**) is positioned and connected to the measuring device (**4**). An expanded view of the electrodes is shown. (**B**) Illustration of a cross-section of a microfluidic chip and positioning of the electrodes. PhG: phaseguide technology, allowing membrane-free separation of the channels. ECM: extracellular matrix. (**C**) OrganoTEER software used to measure TEER and analyze the results. (**D**) TEER measurements after 30min, 60min, and 120min exposure to 37 µM candidalysin. (**E**) TEER measurements after 30min, 60min, and 120min exposure to 75 µM candidalysin. Results are expressed as boxplot with individual measurements for each chip. Each chip is normalized to its baseline. Data was analyzed using Kruskal Wallis test with Dunn´s post-hoc test (*$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$, ****$p \leq 0.0001$, N=3, n=3-8).

We next investigated the permeability of the Caco-2 tubules after 37 and 75 µM candidalysin exposure, using fluorescent molecules of two different sizes. Permeability to sodium fluorescein (0.4kDa) was significantly increased after 30, 60, and 120min incubation with 37 µM candidalysin (Figure 3**Error! Reference source not found.**D), while permeability to dextran 4.4 kDa was significantly increased only after 30min 37 µM candidalysin exposure (Figure 3E). 75 µM candidalysin increased permeability to both dyes after all incubation times (Figure 3F, G). Following evaluation of permeability across various candidalysin exposure conditions, we selected the 2-hour incubation with 75 µM candidalysin for further analysis.

Figure 3

Figure 3. Caco-2 tubule permeability to fluorescent dyes after candidalysin exposure. (**A**) Illustration of a chip with the dye added to the lumen of the Caco-2 tubule. (**B**) Enlarged view of a chip with dye perfused in the lumen of the tubule. Top illustration shows a leak-tight tubule; bottom illustration shows a leaky tubule. (**C**) Cross-section of a chip to show a leak-tight tubule at the top and a leaky tubule at the bottom. (**D**) Permeability of Caco-2 tubule to 0.4kDa dye after 30, 60, and 120min exposure to 37 µM candidalysin. (**E**) Permeability of Caco-2 tubule to 0.4kDa dye after 30, 60, and 120min exposure to 75 µM candidalysin. (**F**) Permeability of Caco-2 tubule to 4.4kDa dye after 30, 60, and 120min exposure to 37 µM candidalysin. (**G**) Permeability of Caco-2 tubule to 4.4kDa dye after 30, 60, and 120min exposure to 75 µM candidalysin. Results are expressed as boxplot with individual measurements for each chip. Data was analyzed using Kruskal Wallis test with Dunn´s post-hoc test (*$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$, ****$p \leq 0.0001$, N=3, n=3-7).

We then assessed whether the expression of the permeability marker DRAQ7 was increased after 75 µM candidalysin exposure for 2h when compared to vehicle. Figure 4A shows images of representative chips exposed to vehicle or candidalysin and stained for DRAQ7 and Hoechst.

254 Permeable cells' nuclei and cell count (total nuclei) quantification are shown in middle and bottom
255 rows, respectively. 75 µM candidalysin significantly increased the number of DRAQ7 positive cells
256 compared to vehicle (Figure 4B).

Figure 4

Figure 4. Candidalysin increases permeability of IECs. Caco-2 tubules exposed to candidalysin for 2h were stained
with DRAQ7 nuclear dye, which only penetrates permeabilized or dead cells. Nuclei were also stained with Hoechst. (**A**)
A CellProfiler pipeline was designed to identify and quantify DRAQ7-positive and Hoechst-positive nuclei from confocal
images. DRAQ7 + objects were filtered to only consider those overlapping with Hoechst positive objects. Pictures are
representative of 3 independent experiments (N=3). Scalebar is 100 µM. (**B**) The number of DRAQ7 + nuclei was
normalized against the total number of nuclei (Hoechst positive) to give a percentage of DRAQ7 positive cells.) Results
are expressed as boxplot with individual measurements for each chip. Data was analyzed using Wilcoxon Rank-Sum Test
(****p ≤ 0.0001, N=3, n=4-5).

3.2   Candidalysin induces actin remodeling in Caco-2 tubules

Next, we investigated the morphology of the tubules after candidalysin exposure. Cells were exposed
to vehicle or 75 µM candidalysin for 2h. Figure5A shows images of representative chips exposed to
vehicle or candidalysin and stained for actin and nuclei. Actin positive objects and cell count (nuclei)
quantification are shown in middle and bottom rows, respectively. 75 µM candidalysin significantly
increased the number of actin positive objects (Figure 5B), their area (Figure 5C), and their perimeter
(Figure 5D) compared to vehicle. All parameters were normalized to the total cell number.

Figure 5

Figure 5. Candidalysin induces actin remodeling. Caco-2 tubules exposed to candidalysin for 2h were stained with
actin and Hoechst nuclear dye. (**A**) A CellProfiler pipeline was designed to identify and quantify actin-positive and
Hoechst-positive objects from confocal images. Pictures are representative of 3 independent experiments (N=3). Scalebar
is 100 µM. The number (**B**), area (**C**), and perimeter (**D**) of Actin-positive objects were normalized against the number of
nuclei (Hoechst positive). Results are expressed as boxplot with individual measurements for each chip. Data was
analyzed using independent T-test or Wilcoxon Rank-Sum Test (****p ≤ 0.0001, N=3, n=4-5).

3.3 Exposure to candidalysin induces cytotoxicity and secretion of inflammatory markers in Caco-2 tubules

To evaluate the effect of candidalysin on the cellular activation of Caco-2 tubules, the production of epithelial inflammatory mediators LL37, S100A8/A9, MCP-1, IP-10, IL-8, IL-6, IL-1beta, IL-1alpha, GM-CSF, MIP-3 alpha, and G-CSF was quantified using ELISA and Luminex technology. Caco-2 cells produced no or low amounts of these analytes in non-triggered conditions. After 2h 75 µM candidalysin, secretion of all mediators increased significantly (Figure 6), except S100A8/A9, which was not detected. We used the lactate dehydrogenase (LDH) assay to measure cytotoxicity after candidalysin exposure. LDH was significantly higher in chips treated with 2h 75µM candidalysin compared to vehicle (Figure 6).

Figure 6

Figure 6. Candidalysin induces cytotoxicity and secretion of inflammatory markers in Caco-2 tubules.
Candidalysin induced the secretion of MIP-3 alpha, G-CSF, GM-CSF, Il1-alpha, Il-1 beta, IL-6, IL-8, IP-10, MCP-1, LL37, and LDH in the apical side (tubule lumen). Results are expressed as boxplot with individual measurements for each chip. Data was analyzed using independent T-test or Wilcoxon Rank-Sum Test (*$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$, ****$p \leq 0.0001$, N=3, n=3-5).

3.4 Assessment of response to candidalysin in human colon organoids

To validate our results obtained from Caco-2 tubules and to explore future applications in a patient-derived model, we conducted a proof-of-concept experiment using human colon organoid tubules. We used the OrganoTEER to measure the barrier integrity of colon organoids following a 2-hour exposure to 0, 18, 37, and 75 µM candidalysin. Subsequently, we fixed the plates and performed DRAQ7 and actin staining (Figure 7A) and quantification. Compared to the vehicle, only 75 µM

candidalysin significantly decreased TEER after the 2-hour exposure (Figure 7B). Both 37 and 75 µM candidalysin increased permeability to DRAQ7 compared to the vehicle (Figure 7C). Furthermore, 75 µM candidalysin led to significant actin remodeling, as evidenced by increases in the number (Figure 7D), area (Figure 7E), and perimeter of actin objects (Figure 7F).

Figure 7

Figure 7. Candidalysin exposures in human colon organoid tubules. (**A**) colon organoid tubules exposed to 0, 18, 37, and 75 µM candidalysin for 2h were stained with actin (green), DRAQ7 (red), and Hoechst nuclear dye (blue). Representative images are max projections taken at the bottom of the tubules. Scale bar is 100 µM. (**B**) TEER measurement after candidalysin exposure. Each chip is normalized to its value at baseline (before exposure). (**C**) A CellProfiler pipeline was designed to identify and quantify DRAQ7-positive and Hoechst-positive nuclei from confocal images. DRAQ7 + objects were filtered to only consider those overlapping with Hoechst positive objects. The number of DRAQ7 + nuclei was normalized against the total number of nuclei (Hoechst positive) to give a percentage of DRAQ7 positive cells. Another CellProfiler pipeline was designed to identify and quantify actin-positive and Hoechst-positive objects from confocal images. The area (**D**), perimeter (**E**), and number (**F**) of Actin-positive objects were normalized

316 against the number of nuclei (Hoechst positive). Results are expressed as boxplot with individual measurements for each
317 chip. Data was analyzed using Kruskal Wallis test with Dunn´s post-hoc test or one-way ANOVA with Tukey's HSD
318 post-hoc test (*p ≤ 0.05; **p ≤ 0.01; ***p ≤ 0.001, ****p ≤ 0.0001, N=3, n=3-6).

4 Discussion

This study aimed to evaluate the phenotypic response of IECs to candidalysin, focusing on barrier
integrity and inflammation. Given that candidalysin forms pores and damages the epithelium[13,14], we
hypothesized it would impair barrier integrity and increase permeability in our model. We also
expected to detect toxicity and inflammation mediators, as described in studies using oral epithelial
cells[17,19].

To test these hypotheses, we used the OrganoPlate technology, allowing the culture of up to 64
perfused intestinal Caco-2 tubules in a membrane-free system[24]. This platform allowed the rapid
assessment of the barrier integrity of the gut tubules[25], along with easy imaging and sampling of
luminal medium to detect inflammatory markers. We found that candidalysin impaired barrier
integrity of Caco-2 tubules (measured by TEER and barrier integrity assay), increased permeability
(measured by DRAQ7 staining), and induced actin remodeling and secretion of inflammatory
mediators. We then validated our results in a more complex model using patient-derived organoid
tubules exposed to candidalysin, measuring TEER, DRAQ7 permeability, and actin remodeling.

Disruption of barrier integrity is a hallmark of mucosal inflammation. Using a combination of
readouts, we observed that candidalysin impaired barrier integrity and increased permeability. In
Caco-2 tubules, only TEER detected significant differences at all timepoints after 37 µM
candidalysin exposure, suggesting that TEER is more sensitive than barrier integrity assay[24,25]. In
colon organoid tubules, TEER was less sensitive than in Caco-2, with only 75 µM of candidalysin
showing a significant decrease after 2-hour exposure. This reduced sensitivity may be due to the
increased complexity of colon organoids compared to Caco-2 and could be influenced by the
washing step before exposure, as indicated by decreased TEER in the vehicle control.

While several studies reported loss of barrier integrity after infection with *C. albicans* in IECs[7,10,27]
and other epithelial cell type[28,29], to date, only one study used exogenous candidalysin in IECs.
Contrary to our observations, they did not observe a decrease in TEER or an increase in 4-kDa
dextran permeability after 24 hours of 70 µM candidalysin exposure using the Caco-2 subclone
C2BBe1 in Transwells[10]. This discrepancy likely arises from model differences (dynamic gut-on-chip
vs. static Transwells), exposure duration (acute 2-hour vs. prolonged 24-hour), cell line sensitivity,
and measurement sensitivity (more responsive TEER in the gut-on-chip[25]). These differences
highlight how model choices impact findings, suggesting the gut-on-chip system could better capture
acute barrier disruption, influencing interpretations of candidalysin's role in epithelial damage.

The actin cytoskeleton mediates disruption of mucosal barriers in inflamed tissues[30]. We therefore
looked at the actin network after candidalysin exposure and observed remodeling in Caco-2 and
organoids, confirming results observed in oral epithelial cells[21]. Notably, in colon organoids, actin
clumps were found to co-localize with Draq7-positive nuclei, suggesting a localized effect of
candidalysin in these areas.

Similarly to observations in other cell types[12,17–19], we observed the induction of several
candidalysin-induced proteins in Caco-2. Along with the release of LL-37, G-CSF, GM-CSF, IL-6,
IL-1a, IL-1b, MCP-1, and IL-8, we also observed candidalysin-induced release of IP-10 and MIP-3

in our cultures. Notably, S100A8/A9 release was not induced by candidalysin in our system, in line
with findings in TR146 cells[17]. The secretion of LL-37, an antimicrobial peptide, is induced by
candidalysin, highlighting its role in the host defense mechanism. However, further experiments with
additional stimuli are needed to determine the specificity of this response. Future studies could
address this by testing other toxins or stimuli to compare their effects on LL-37 release and clarify
whether this response is specific to candidalysin. These observations support the notion that the
epithelial barrier is not just a physical barrier but actively participates in the host's early response to
pathogens[31,32].

A limitation of our study is the use of exogenous candidalysin. While it allows for the evaluation of
its direct effect on IECs and provides insights into the early stages of *C. albicans* infection, it is
important to note that the delivery of candidalysin to host cell membranes is critical[33] and is not
recapitulated in our system. Additionally, our model does not address direct fungal-host interactions,
such as the use of host zinc by the yeast to promote translocation and virulence[23]. Another limitation
is the lack of immune components. Gut-on-chip models of *C. albicans* infection including living
yeast and immune cells have been reported[34] but still present technical challenges to be scaled up for
testing potential therapies[35]. Moreover, advanced gut-on-chip models that enable the study of
immune migration have been described[36,37], and their adaptation to investigate candidiasis
pathogenicity could be a promising direction. Another point to consider is the interactions between *C.*
albicans and the rest of the microbiota, as it has been shown that candidalysin can inhibit bacterial
species[38].

Due to the pilot nature of this study, we did not measure protein detection in organoid tubules. Future
experiments should incorporate this approach to leverage the more physiologically relevant
environment of organoids, potentially providing more accurate insights into protein expression
responses. Furthermore, the response to candidalysin might be donor-specific, highlighting the need
to test in more donors to capture the variability in responses. This research represents a starting point
for understanding the physiological responses to candidalysin and underscores the necessity of
screening more donors while examining various response angles.

In summary, our gut-on-chip model effectively evaluated the phenotypic response to candidalysin in
intestinal epithelial cells, providing insights into the toxin's mechanisms. This system's ability to
multiplex readouts offers a valuable tool for screening potential therapeutic inhibitors and modulators
of candidalysin-induced responses. It could be used to screen for candidalysin inhibitors, such as
nanobodies neutralizing candidalysin[39], or modulators of candidalysin-induced responses.
Furthermore, this model can be adapted to study the effects of candidalysins produced by other non-
albicans Candida species, which are also prevalent fungal pathogens[19]. It could also be adapted to
study the phenotypic response of other molecules like other toxins[40], micro-organisms, and microbial
metabolites. Lastly, the use of patient-derived organoids allows for personalized modeling and the
use of cells coming from patients with candidiasis or other diseases. Overall, our findings highlight
the potential of our platform to study the effects of various virulence factors produced by different
microorganisms and microbial communities. This paves the way for developing targeted therapeutic
strategies that modulate these factors or the host responses they trigger.

5   Conflict of Interest

M.M., and K.Q. are employees of Mimetas B.V., which markets OrganoPlate, OrganoTEER,
OrganoReady and OrganoFlow, and holds the registered trademarks OrganoPlate, OrganoTEER,
OrganoReady, and OrganoFlow.

6   Author Contributions

M.M. designed and performed the experiments, analyzed the data, and wrote the manuscript. KQ provided guidance and revision. All authors have read and agreed to the published version of the manuscript.

7   Funding

M.M. is supported by European Union's Horizon 2020 research and innovation program under the Marie Sklodowska-Curie action, Innovative Training Network: FunHoMic; Grant No: 812969.

K.Q. is supported by an innovation credit (IK17088) from the Ministry of Economic Affairs and Climate of the Netherlands.

Research reported in this publication was supported by Oncode Accelerator, a Dutch National Growth Fund project under grant number NGFOP2201.

8   Acknowledgments

The authors would like to thank Bernhard Hube, Mark Gresnigt, and Selene Mogavero of the Hans-Knöll institute in Jena, Germany for their insights on candidalysin usage and for helpful discussions on this work. We thank Frederik Schavemaker at Mimetas for designing some of the illustrations. We also thank our colleagues at Mimetas and in the FunHoMic consortium for the many fruitful discussions.

12   Data Availability Statement

The datasets generated during this study are available from the corresponding authors on reasonable request.

13   References

1. Zheng, D., Liwinski, T. & Elinav, E. Interaction between microbiota and immunity in health and disease. *Cell Res* **30**, 492–506 (2020).

2. Brown, G. D. *et al.* Hidden killers: human fungal infections. *Sci Transl Med* **4**, 165rv13 (2012).

3. d'Enfert, C. *et al. The Impact of the Fungus-Host-Microbiota Interplay upon Candida Albicans Infections: Current Knowledge and New Perspectives. FEMS Microbiology Reviews* (2020). doi:10.1093/femsre/fuaa060.

4. Alonso-Monge, R., Gresnigt, M. S., Román, E., Hube, B. & Pla, J. Candida albicans colonization of the gastrointestinal tract: A double-edged sword. *PLoS Pathog* **17**, e1009710 (2021).

5. Gouba, N. & Drancourt, M. Digestive tract mycobiota: a source of infection. *Med Mal Infect* **45**,

9–16 (2015).

6. Miranda, L. N. *et al.* Candida colonisation as a source for candidaemia. *J Hosp Infect* **72**, 9–16

(2009).

7. Graf, K. *et al.* Keeping Candida commensal: how lactobacilli antagonize pathogenicity of

Candida albicans in an in vitro gut model. *Disease Models & Mechanisms* **12**, dmm039719

(2019).

8. Naglik, J. R., Richardson, J. P. & Moyes, D. L. Candida albicans Pathogenicity and Epithelial

Immunity. *PLoS Pathogens* **10**, 8–11 (2014).

9. Mayer, F. L., Wilson, D. & Hube, B. Candida albicans pathogenicity mechanisms. *Virulence* **4**,

119–128 (2013).

10. Allert, S. *et al.* Candida albicans-induced epithelial damage mediates translocation through

intestinal barriers. *mBio* **9**, 1–20 (2018).

11. Naglik, J. R., Gaffen, S. L. & Hube, B. Candidalysin: discovery and function in Candida albicans

infections. *Current Opinion in Microbiology* **52**, 100–109 (2019).

12. Moyes, D. L. *et al.* Candidalysin is a fungal peptide toxin critical for mucosal infection. *Nature*

532, 64–68 (2016).

13. Russell, C. M. *et al.* The Candida albicans virulence factor candidalysin polymerizes in solution

to form membrane pores and damage epithelial cells. *eLife* **11**, 1–21 (2022).

14. Russell, C. M., Rybak, J. A., Miao, J., Peters, B. M. & Barrera, F. N. Candidalysin: Connecting

the pore forming mechanism of this virulence factor to its immunostimulatory properties. *J Biol*

Chem **299**, 102829 (2023).

15. Ho, J. *et al.* Candidalysin activates innate epithelial immune responses via epidermal growth

factor receptor. *Nature Communications* **10**, (2019).

16. Nikou, S. A. *et al.* The Candida albicans toxin candidalysin mediates distinct epithelial

inflammatory responses through p38 and EGFR-ERK pathways. *Science Signaling* **15**, (2022).

17. Ho, J. *et al.* Candidalysin Is a Potent Trigger of Alarmin and Antimicrobial Peptide Release in

Epithelial Cells. *Cells* **9**, 699 (2020).

18. Hanaoka, M. & Domae, E. IL-1α released from oral epithelial cells upon candidalysin exposure

initiates an early innate epithelial response. *International Immunology* (2020)

doi:10.1093/intimm/dxaa070.

19. Richardson, J. P. *et al.* Candidalysins Are a New Family of Cytolytic Fungal Peptide Toxins.

mBio **13**, 1–13 (2022).

20. Blagojevic, M. *et al.* Candidalysin triggers epithelial cellular stresses that induce necrotic death.

Cellular Microbiology **23**, 1–25 (2021).

21. Westman, J. *et al.* Calcium-dependent ESCRT recruitment and lysosome exocytosis maintain

epithelial integrity during Candida albicans invasion. *Cell Reports* **38**, 110187 (2022).

22. Nucci, M. & Anaissie, E. Revisiting the source of candidemia: Skin or gut? *Clinical Infectious*

Diseases **33**, 1959–1967 (2001).

23. Sprague, J. L. *et al.* Candida albicans translocation through the intestinal epithelial barrier is

promoted by fungal zinc acquisition and limited by NFκB-mediated barrier protection. *PLoS*

Pathog **20**, e1012031 (2024).

24. Trietsch, S. J. *et al.* Membrane-free culture and real-time barrier integrity assessment of perfused

intestinal epithelium tubes. *Nature Communications* **8**, 1–7 (2017).

475 25. Nicolas, A. *et al.* High throughput transepithelial electrical resistance (TEER) measurements on

476  perfused membrane-free epithelia. *Lab on a Chip* **21**, 1676–1685 (2021).

477 26. Soragni, C. *et al.* Quantify permeability using on-a-chip models in high-throughput applications.

478  *STAR Protocols* **4**, 102051 (2023).

479 27. Fusco, A., Savio, V., Donniacuo, M., Perfetto, B. & Donnarumma, G. Antimicrobial Peptides

480  Human Beta-Defensin-2 and -3 Protect the Gut During Candida albicans Infections Enhancing

481  the Intestinal Barrier Integrity: In Vitro Study. *Frontiers in Cellular and Infection Microbiology*

482  **11**, 1–11 (2021).

483 28. Kumar, R., Rojas, I. G. & Edgerton, M. Candida albicans Sap6 Initiates Oral Mucosal

484  Inflammation via the Protease Activated Receptor PAR2. *Front Immunol* **13**, 912748 (2022).

485 29. Eyerich, S. *et al.* IL-22 and TNF-α represent a key cytokine combination for epidermal integrity

486  during infection with Candida albicans. *Eur J Immunol* **41**, 1894–1901 (2011).

487 30. Lechuga, S. & Ivanov, A. I. Actin cytoskeleton dynamics during mucosal inflammation: a view

488  from broken epithelial barriers. *Curr Opin Physiol* **19**, 10–16 (2021).

489 31. Naglik, J. R. & Moyes, D. Epithelial cell innate response to Candida albicans. *Advances in dental*

490  *research* **23**, 50–55 (2011).

491 32. Bals, R. Epithelial antimicrobial peptides in host defense against infection. *Respir Res* **1**, 141–

492  150 (2000).

493 33. Mogavero, S. *et al.* Candidalysin delivery to the invasion pocket is critical for host epithelial

494  damage induced by Candida albicans. *Cell Microbiol* **23**, e13378 (2021).

495 34. Maurer, M. *et al.* A three-dimensional immunocompetent intestine-on-chip model as in vitro

496  platform for functional and microbial interaction studies. *Biomaterials* **220**, 119396 (2019).

35. Morelli, M., Kurek, D., Ng, C. P. & Queiroz, K. Gut-on-a-Chip Models: Current and Future

Perspectives for Host-Microbial Interactions Research. *Biomedicines* **11**, (2023).

36. de Haan, L. *et al.* A Microfluidic 3D Endothelium-on-a-Chip Model to Study Transendothelial

Migration of T Cells in Health and Disease. *International Journal of Molecular Sciences* **22**,

8234 (2021).

37. Gjorevski, N. *et al.* Neutrophilic infiltration in organ-on-a-chip model of tissue inflammation.

Lab on a Chip **20**, 3365–3374 (2020).

38. Liang, S.-H. *et al.* The hyphal-specific toxin candidalysin promotes fungal gut commensalism.

Nature 1–8 (2024) doi:10.1038/s41586-024-07142-4.

39. Valentine, M. *et al.* Nanobody-mediated neutralization of candidalysin prevents epithelial

damage and inflammatory responses that drive vulvovaginal candidiasis pathogenesis. *mBio* **15**,

e03409-23 (2024).

40. Morelli, M., Cabezuelo Rodríguez, M. & Queiroz, K. A high-throughput gut-on-chip platform to

study the epithelial responses to enterotoxins. *Sci Rep* **14**, 5797 (2024).