Reaction norm tracking of fitness peaks

may explain time series in the fossil record

Rolf Ergon

University of South-Eastern Norway

29.10.2024

Abstract

Species that have exixted over millions of years have done so because of high mean fitness through the ages. This means that such species continuously have been able to evolve mean phenotypic traits that have tracked the optimal phenotypic values that maximize fitness. These optimal values are functions of the environment, for example the sea temperature. The mean phenotypic traits over time will thus be determined by the environment over time, and the trait history may be found in the fossil record. Here, I simulate such a tracking system by use of a univariate intercept-slope reaction norm model and show how both linear and nonlinear mean phenotype vs. environment functions can be found also from quite short time series with irregular sampling, and I also discuss how this methodology can be extended to multivariate systems. The methodology is finally applied on two plastic traits in a record of *Microporella agonistes* fossils spanning 2.3 million years, using the $\partial^{18}O$ measure as environmental variable. From few irregular samples of phenotypic mean values found in the fossil record, it was possible to identify linear mean phenotype vs. environment functions, and thus to predict the continuous mean phenotypic values as function of time.

Keywords: bryozoans, evolution, fossil record, optimal phenotypes, parameter estimation, phenotypic plasticity, reaction norm tracking.

Introduction

Evolutionary time series in the fossil record provide information on phenotypic changes over millions of years. Such time series are often short, and they have traditionally been modeled by various forms of random walk processes (Hunt, 2006; Voje, 2023), or by Ornstein-Uhlenbeck (OU) processes describing evolution of phenotypic traits toward fixed or moving peaks in the adaptive landscape (Voje, 2023). In an example, Voje (2023) shows how an OU model describes the evolution in *Eucyrtidium calvertense* from one optimal size to another (Kellogg,1975). Neither random walk processes nor OU processes have explicit means for handling of phenotypic plasticity, but one cannot exclude the possibility that plasticity plays an implicit part in such models. Such possibilities are discussed in Sætre et al. (2017), where an OU model was used in the analysis of changed body size measurements of individuals of *Acrocephalus scirpaceus* over a 19-year period.

Reaction norm analysis of fossil data under the explicit assumption of phenotypic plasticity would in many cases appear to be more appropriate, especially for species that have adapted to environmental changes over millions of years with no apparent loss of fitness. In such cases the mean phenotypic traits have been tracking the optimal values that maximize fitness as determined by moving fitness peaks in the adaptive landscape (Bell, 2013).

Phenotypic plasticity is ubiquitous in nature and generally regarded as a key mechanism for enabling organisms to survive in the face of environmental change (Williams et al., 2008). Phenotypic changes in natural populations caused by climate change have been widely studied, and a fundamental difficulty is then to disentangle the effects of microevolution and phenotypic plasticity. Merilä and Hendry (2013) discussed this problem and summarized the results of 11 review articles. Among the review papers were Charmantier and Gienapp (2014)

(birds), Boutin and Lane (2014) (mammals), and Crozier and Hutchings (2014) (fish), which

summarize the findings in altogether 66 studies of responses on climate change. Merilä and

Hendry (2013) wrote that "Summarizing and relying on the results of these reviews, we arrive at

the conclusion that evidence for genetic adaptation to climate change has been found in some

systems, but is still relatively scarce." They further wrote that "Indeed, a number of phenotypic

differences originally thought to be genetically based were subsequently attributed to

phenotypic plasticity (e.g., James 1983; Charmantier et al. 2008; Teplitsky et al. 2008). In the

wake of such demonstrations phenotypic plasticity is increasingly adopted as a parsimonious

model that is to be rejected only if direct evidence is obtained for genetic change: That is,

plasticity is treated as a null model. The reality, however, is that inferences regarding

phenotypic plasticity also should be supported by a specific set of evidentiary criteria". The

disentanglement problem was also discussed in Lister (2021), who concluded that "The role of

phenotypic plasticity is particularly challenging to identify in the fossil record as the

environmental and genetic influences on morphological variation and change are difficult to

separate".

In a more recent review regarding temporal variation in selection on breeding date across 39

populations of 21 wild animals, de Villemereuil et al. (2020) found that "plastic changes track

movements of the optimum phenotype across years, especially in bird species, reducing

temporal variation in directional selection. This suggests that phenological plasticity has

evolved to cope with fluctuations in the optimum, despite their currently modest contribution to

variation in selection".

In general, a reaction norm model should be multivariate, allowing for correlated traits (Ergon,

2023). For simplicity I will here make use of a univariate individual intercept-slope reaction

norm model with parameters $a(t)$ and $b(t)$ as given by

$$y(t) \, = \, a(t) + b(t)\big(u(t) - u_{ref}\big) + \varepsilon(t), \tag{1}$$

where $u(t) - u_{ref}$ is the environmental cue, and where $\varepsilon(t)$ is assumed to be normally

distributed with constant mean 0 and variance σ_ε^2 among individuals in every generation

(Lande, 2009, although Lande assumed $u_{ref} = 0$). See comments on the definition of an

environmental cue in Discussion. I will use simulations based on the model in Eq. (1) to see

how the mean phenotypic trait $\bar{y}(t)$ in a population evolves in a changing environment under

the fundamental assumption that the mean fitness is kept close to the top of a moving peak in

an adaptive landscape. In the simulations I will make use of the $\partial^{18}O(t)$ measure, which is a

widely used proxy for seawater temperature in paleontological studies (Lisiecki and Raymo,

2005; Liow et al., 2024). Since both $\partial^{18}O(t)$ and $\bar{y}(t)$ have large variances, and since data from

the fossil record may give mean values over both populations and time (Liow et al., 2024), I will

in the simulations use filtered versions of these variables.

Basic theory is given in the Theory section, while simulations are presented in the Simulation

section. In the Real data section, I will apply the theoretical results on two phenotypic traits of

Microporella agonistes with a 2.3 million long fossil record (Liow et al., 2024). Finally follows a

Discussion section.

Theory

A univariate case

88   The individual intercept-slope reaction norm model with parameters $a(t)$ and $b(t)$ in Eq. (1) will

89   result in the mean reaction norm

$$\bar{y}(t) = \bar{a}(t) + \bar{b}(t)\big(u(t) - u_{ref}\big). \tag{2}$$

Provided that $\bar{a}(t)$ and $\bar{b}(t)$ evolves such that the mean fitness is at a near maximum level, $\bar{y}(t)$

will be kept in the vicinity of moving fitness peaks in the adaptive landscape. For simplicity and

for use in simulations I will initially assume that mean fitness is determined by the single

phenotypic mean trait $\bar{y}(t)$ in Eq. (2), and that the adaptive landscape thus is two-dimensional.

I will later extend some essential results to multivariate cases.

The phenotypic value that maximizes fitness is in general a function of the environment,

$$\theta(t) = \alpha + f\big(u(t) - u_{ref}\big), \tag{3}$$

96   in the linear case $\theta(t) = \alpha + \beta\big(u(t) - u_{ref}\big)$, where α and β are constant parameters. In the

97   simulations I will assume an individual fitness function

$$W(t) = W_{max} \exp\left(-\frac{\big(y(t) - \theta(t)\big)}{2\omega^2}\right), \tag{4a}$$

98   which gives the mean fitness in a population (Lande, 2009)

$$\overline{W}(t) = W_{max}\sqrt{\frac{\omega^2}{\omega^2 + \sigma_y^2}} \exp\left(-\frac{\big(\bar{y}(t) - \theta(t)\big)}{2\omega^2}\right), \tag{4b}$$

99   where σ_y^2 is the variance of $y(t)$ according to Eq. (1).

100   Under the assumptions of nonoverlapping generations and random mating in a large

101   population the evolution of the mean reaction norm parameters is governed by (Lande, 1979)

$$\begin{bmatrix} \bar{a}(t+1) - \bar{a}(t) \\ \bar{b}(t+1) - \bar{b}(t) \end{bmatrix} = \begin{bmatrix} G_{aa} & G_{ab} \\ G_{ab} & G_{bb} \end{bmatrix} \begin{bmatrix} G_{aa} + \sigma_\varepsilon^2 & G_{ab} \\ G_{ab} & G_{bb} \end{bmatrix}^{-1} \begin{bmatrix} cov(w(t), a(t) + \varepsilon(t)) \\ cov(w(t), b(t)) \end{bmatrix}, \tag{5}$$

where G_{aa}, G_{bb} and G_{ab} are variances and the covariance of the reaction norm parameters, and

where $w(t) = W(t)/\overline{W}(t)$ is the relative fitness. In a stationary stochastic environment, the

optimal reaction norm slope is $\bar{b}_{opt} = cov(\theta(t), u(t))/var(u(t))$ (Lande, 2009; Ergon, 2022).

When the mean phenotypic trait in a population is tracking a moving value $\theta(t)$ according to

Eq. (3) it will become $\bar{y}(t) = \theta(t) + e(t)$, where $e(t)$ is a residual term. This means that the

parameter values $\bar{a}(t)$ and $\bar{b}(t)$ are forced to evolve such that $\bar{y}(t) \approx \theta(t) = \alpha + f\big(u(t) - u_{ref}\big)$,

i.e., such that the mean fitness according to Eq. (4b) is kept at a near maximum level and thus

the residual $e(t)$ at a low level. It is thus the mean trait following error $\theta(t) - \bar{y}(t)$ that is the

driving force in the adaptive process. As shown in simulations as well as in the real data case the

parameters in such a $\theta(t)$ model can be estimated also from rather few fossil data points at

irregular sampling times. Note that even though $\bar{y}(t) \approx \alpha + f\big(u(t) - u_{ref}\big)$, such a model is not

a reaction norm. As verified by simulations, the reaction to a sudden change Δu in $u(t)$ will still

be a sudden change $\Delta \bar{y} = \bar{b}(t)\Delta u$.

Multivariate cases

When the mean phenotypic value in a population is tracking a moving fitness peak, the mean

phenotypic value will in the univariate case be determined by

$\bar{y}(t) \approx \theta(t) = \alpha + f\big(u(t) - u_{ref}\big)$, i.e., by the fitness peak vs. environment function. Here, it is

important to note that we do not need to know the details of the underlying reaction norm

model and fitness function.

The univariate theory can be extended to multivariate cases, and although we in practice do

not need to know the underlying fitness function, we may for a theoretical discussion use an

example with two phenotypic traits and the bivariate fitness function (Ch. 4, Johnson and

Wichern, 2008)

$$W(t) = W_{max} \exp\left(-[y_1(t) - \theta_1(t) \quad y_2(t) - \theta_2(t)]\Sigma^{-1}\begin{bmatrix} y_1(t) - \theta_1(t) \\ y_2(t) - \theta_2(t) \end{bmatrix}/2c\right), \qquad (6)$$

125   where Σ is the variance-covariance matrix and where c is a scaling factor controlling the 'width'

126   of the fitness function. In order to keep the mean fitness at a near maximum level, the mean

127   traits must in this case continuously evolve such that $\bar{y}_1(t) \approx \theta_1(t)$ and $\bar{y}_2(t) \approx \theta_2(t)$, and we

128   will thus find

$$\bar{y}_1(t) \approx \alpha_1 + f_1\big(u(t) - u_{ref}\big) \qquad (7a)$$

$$\bar{y}_2(t) \approx \alpha_2 + f_2\big(u(t) - u_{ref}\big). \qquad (7b)$$

where the fitness peak vs. environment functions may be linear or nonlinear, and as in the

univariate case the parameters can be estimated from rather few fossil data points at irregular

sampling times. Here, the essential point is that $\bar{y}_1(t)$ and $\bar{y}_2(t)$ are tracking a peak in a three-

dimensional adaptive landscape, and that this tracking process can be decomposed into two

univariate tracking processes. We may thus consider the projections of the bivariate fitness

function onto orthogonal axes for the two phenotypic traits, and such a projection approach

may be extended to systems with higher dimensions, as discussed in Goodnight (2013).

Simulations

A system with a linear mean reaction norm $\bar{y}(t) = \bar{a}(t) + \bar{b}(t)\big(u(t) - u_{ref}\big)$ was simulated with

use of the 2,115 available samples of $u(t) = \partial^{18}O(t)$, and with population size 1,000. These

samples go 5.3 million years back in time with various sampling intervals, but here I just

assume 2,115 nonoverlapping generations (however, see Discussion on relevance for fossil

data). The reaction norm parameter variances and covariance were chosen as $G_{aa} = \sigma_\varepsilon^2 = 0.5$,

$G_{bb} = 0.05$ and $G_{ab} = 0$, and at each new generation random individual samples around $\bar{a}(t)$,

$\bar{\varepsilon}(t) = 0$ and $\bar{b}(t)$ were drawn from normal populations with the given variances. The fitness

function in Eq. (4a) had parameter values $W_{max} = 1$ and $\omega^2 = 50$. The phenotypic value that

maximizes fitness was in the first linear case chosen as $\theta(t) = 0.5 + u(t) - 2.9$, where $\theta(t)$

thus and for clarity of presentation is assumed to be a deterministic function, and where 2.9 is

the mean value of $\partial^{18}O(t)$ around $t = 0$. From this follows that the optimal plasticity slope is

$\bar{b}_{opt} = cov(\theta(t), u(t))/var(u(t)) = 1$. The initial intercept value was $\bar{a}(0) = 0$, and simulations

were performed with three different initial values $\bar{b}(0)$. Since $\bar{y}(t)$, $\theta(t)$ and $u(t) = \partial^{18}O(t)$

have large variances, I also used filtered signals $\bar{y}_{filt}(t)$, $\theta_{filt}(t)$ and $u_{filt}(t)$ in the plots. These

were obtained by moving window filters with window size 100, in MATLAB notation

$\bar{y}_{filt}(t) = movmean(\bar{y}(t), 100)$ etc.

Simulation results are shown in Fig. 1. Note that $\bar{b}(0) = \bar{b}_{opt} = 1$ results in a constant

plasticity slope, and that also $\bar{a}(t)$ settles on a constant value (panel E). With an initial mean

plasticity slope value $\bar{b}(0) \neq \bar{b}_{opt}$, the mean intercept evolves all the time, while the mean

plasticity slope very slowly evolves towards $\bar{b}_{opt} = 1$. Note that $\bar{y}_{filt}(t)$ is tracking $\theta_{filt}(t)$ quite

well (panels C, G and K). Also note that $\bar{y}_{filt}(t)$ as functions of $u_{filt}(t)$ are very similar straight

lines as predicted by theory, with seven discrete values marked by stars (panels D, H and L).

These discrete values could be used to determine the slope of this line by an ordinary least

squares method. With $\bar{b}(0) = 1$ it appears that $\bar{a}(t)$ and $\bar{b}(t)$ are constant over time, once $\bar{a}(t)$

has settled (panel E), but a closer look reveals slight fluctuations around fixed values. This is

because the individual values around $\bar{a}(t)$, $\bar{\varepsilon}(t) = 0$ and $\bar{b}(t)$ at each time step are drawn from

a random Gaussian distribution.

Fig. 2 shows result for a system with a nonlinear function

$\theta(t) = 0.5 + (u(t) - 2.9) + 2(u(t) - 2.9)^2$, combined with the linear reaction norm

$\bar{y}(t) = \bar{a}(t) + \bar{b}(t)(u(t) - 2.9)$. Note that $\bar{y}_{filt}(t)$ also now is tracking $\theta_{filt}(t)$ pretty well

(panels C, G and K) and that $\bar{y}_{filt}(t)$ as function of $u_{filt}(t)$ are the same second order functions

in the different cases (panels D, H and L). See Discussion for results with a nonlinear reaction

norm.

Figure 1. Simulation results with a linear $\theta(t) = \alpha + f(u(t))$ function and a linear

reaction norm for three different initial values $\bar{b}(0)$, while $\bar{a}(0) = 0$. Panels A, E and I

show $\bar{a}(t)$ (blue lines) and $\bar{b}(t)$ (red lines). Panels B, F and J show mean fitness $\overline{W}(t)$.

Panels C, G and K show $\theta_{filt}(t)$ (blue lines) and $\bar{y}_{filt}(t)$ (red lines). Panels D, H and L

show that $\bar{y}_{filt}(t)$ as function of $u_{filt}(t)$ in all cases is the same straight line, with values

for samples 200, 400, 600, 800, 1000, 1200 and 1400 shown by stars.

Figure 2. Simulation results as in Fig. 1, but with a nonlinear $\theta(t) = \alpha + f(u(t))$

function. The discrete values in panels D, H or L could here be used to determine the

nonlinear function $\hat{\bar{y}}_{filt}(t) = f\left(u_{filt}(t)\right)$ by means of any suitable fitting method.

Real data case

Liow et al. (2024) recently presented results from a field study of bryozoans

(genus *Microporella*), based on fossil records over 2.3 million years. The fossils were collected

from colonies in New Zealand cliffs, and it is especially interesting that the same species still

live in the waters near those cliffs. In a project running over several years, Liow and her

coworkers collected 985 fossil colonies with a large total number of individuals and developed

an artificial intelligence tool called DeepBryo, and they then analyzed the data using scanning

electron microscope pictures. This image analysis system can rapidly assess three well defined

phenotypic traits; autozooid size, ovicell size and autozooid shape, and as seen from time

series in Fig. S2 in Liow et al. (2024) the two size traits are plastic. Liow et al. (2024) used

average fecundity and competitive outcome as two different fitness components, where

competitive outcome is a proxy for partial survival.

I will here use the log AZ and log OV area mean data in Fig. S2 in Liew et al. (2024) as

examples. The last data points in these time series are based on an extreme $\partial^{18}O(t)$ value and

are therefore omitted, see Fig. 6 in Liow et al. (2024). Fig. 3 shows $\partial^{18}O$ (panel A),

$log\ AZ\ area\ mean$ (panel B) and $log\ OV\ area\ mean$ (panel C) as functions of time. Panel A also

shows a moving window filtered version $u_{filt}(t)$ of $\partial^{18}O$, with window size 300. Note that the

values of $log\ AZ\ area\ mean$ and $log\ OV\ area\ mean$ are mean values of mean values in several

colonies in time windows of various sizes, i.e., highly filtered information (Liow et al., 2024).

Figure 3. Raw $\partial^{18}O(t)$ data (panel A, blue line) and the filtered version $u_{filt}(t)$ (panel A,

red line), as well as *log AZ area mean* (panel B, circles) and *log OV area mean* (panel C,

circles) as in Fig. S2 in Liow at al. (2024). Present time is used as zero point.

From Fig. 3 in Liow et al. (2024) follows that fitness measured as ovicell density (average

fecundity) is fairly constant for all log OV area mean values, and Fig. S8 shows that also

competitive outcome is fairly constant for all of the three phenotypic traits. Since the bryozoan

species under study have existed for 2.3 million years and still is living it can be inferred that the

mean fitness through the ages has been high enough for tracking of the optimal phenotypic

values that maximize fitness (compare with panels B, F and J in Figs. 1 and 2).

Fig. 4, panel A, shows *log AZ area mean* as function of $u_{filt}(i)$ for the 10 data points in Fig. 3,

panel B. This plot should be compared with the corresponding plots in Figs. 1 and 2 (panels D, H

and L), and it is rather obvious that it is not possible to fit more than a straight line

$log\ AZ\ area\ mean\ = \alpha + \beta\big(u_{filt}(i) - u_{ref}\big)$. With $u_{ref} = u_{filt}(-2.3) = 3.6$ as found from Fig.

3, panel A, the ordinary least squares method results in

$log\ AZ\ area\ mean = 11.39 + 0.48\big(u_{filt}(t) - 3.6\big)$, as shown in Fig. 4, panel B. With unfiltered

environmental data we thus find $log\ AZ\ area$ mean $= 11.39 + 0.48(\partial^{18}O(t) - 3.6)$.

Figure 4. Panel A shows $log\ AZ\ area\ mean$ as function of $u_{filt}(i)$ for 10 data points as in

Fig. S2 in Liow at al. (2024) (circles), as well as a straight line fitted by the ordinary least

squares method (red line). Panel B shows $log\ AZ\ area\ mean$ as function of time (blue

circles), and the fitted reaction norm response (red line).

Finally, Fig. 5 shows results as in Fig. 4, but with the fitted predictions

$log\ OV\ area\ mean = 10.20 + 0.43(u_{filt}(t) - 3.6)$. Note that compared with Fig. 4, the number

of data points is reduced from 10 to 8 owing to a limited number of raw data points (Liow et al.,

2024). A closer look reveals that the predictions of $log\ OV\ area\ mean$ for all practical purposes

are identical to $0.896 \times log\ AZ\ area\ mean$, which cannot be understood without a detailed look

at the data collection procedures.

Figure 5. Panel A shows $log\ OV\ area\ mean$ as function of $u_{filt}(i)$ for 8 data points as in

Fig. S2 in Liow at al. (2024) (circles), as well as a straight line fitted by the ordinary least

squares method (red line). Panel B shows $log\ OV\ area\ mean$ as function of time (blue

circles), and the fitted reaction norm response (red line).

Discussion

The basic insight behind the results in this article is that the mean phenotypic traits will stay in the vicinity of moving peaks in the adaptive landscape defined by the phenotypic value $\theta(t)$ that maximizes fitness, provided that the mean fitness is upheld at a high level. This can for example be seen in the results in Lande (2009), where $\bar{y}(t)$ long after a sudden change from one stationary stochastic environment to another, settles on $\theta(t)$. From this follows the quite simple result that the function $\bar{y}(t) = f(u(t) - u_{ref})$ for a given phenotypic trait approximately is $\theta(t) = f(u(t) - u_{ref})$ for the same trait, and that $\bar{y}(t)$ thus can be predicted by fitting also quite short fossil data.

I have here used $u(t) - u_{ref}$ as environmental cue, where $u(t)$ for example is the temperature on any scale (°C, °F or K). It could be argued that the cue is the environment $u(t)$ itself, but in a mathematical model that must necessarily be on a scale with a defined zero-point. As Lande (2009) I thus use the environment the population is fully adapted to as zero-point, such that the cue is $u_{cue} = u(t) - u_{ref}$ (where Lande (2009) assumed $u_{ref} = 0$). For the real data case in Liow et al. (2024) I assume that u_{ref} is the mean $\partial^{18}O(t)$ value 2.3 million years back in time. Since the data spans 2.3 million years there is a possibility that u_{ref} evolves, as discussed in Ergon and Ergon (2017), but since $\bar{y}_{filt}(i)$ as function of $u_{filt}(i)$ is a straight line such a change in u_{ref} has no effect on the results.

In the simulations with a linear fitness peak vs. environment function $\theta(t) = 0.5 + u(t) - 2.9$ and use of the linear reaction norm in Eq. (2) we find linear $\bar{y}_{filt}(t)$ vs. $f(u_{filt}(t))$ plots. This is so with constant mean intercept and plasticity parameters (Fig. 1, panel H), which is in line with the findings in de Villemereuil et al. (2020) regarding breeding date in populations of wild animals. But the same linear result is also found in cases with time varying parameters (Fig. 1, panels D and L), and there is thus no need for disentanglement of the effects of microevolution and plasticity, as discussed in Merilä and Hendry (2013) and Lister (2021). This is so because

the function $\bar{y}_{filt}(t) = f(u_{filt}(t))$ only depends on how $\theta(t)$ varies with the environment, and

that $\bar{y}_{filt}(t) \approx \theta_{filt}(t)$ (Figs. 1, panels C, G and K). From the $\bar{y}_{filt}(t)$ vs. $f(u_{filt}(t))$ plots it is

possible to fit a straight line also from quite few known values $u_{filt}(i)$ and $\bar{y}_{filt}(i)$, which is a

valuable property considering the effort needed to obtain fossil data. Fig. 1 also shows that the

mean fitness in all three cases is slowly reduced over time. This is because u_{ref} is chosen as

$u_{filt}(0)$, such that the individual phenotypic values owing to plasticity are increasingly spread

around the fitness peak as time increases.

In the nonlinear case simulations with $\theta(t) = 0.5 + u(t) - 2.9 + 2(u(t) - 2.9)^2$ and use of

the linear reaction norm in Eq. (2) we find that the mean fitness is somewhat reduced (Fig. 2),

and that $\bar{y}_{filt}(t)$ vs. $f(u_{filt}(t))$ plots indicate a nonlinear function. As in the linear case we find

very similar plots for the three different initial mean slope values (Fig. 2, panels D, H and L). In

these plots it is here possible to fit a second order function from quite few known values $u_{filt}(i)$

and $\bar{y}_{filt}(i)$. As the combination of a linear reaction norm and a nonlinear $\theta(t)$ function leads to

reduced mean fitness, it is natural to consider a nonlinear reaction norm, and the mean fitness

is in fact considerably improved with a reaction norm

$\bar{y}(t) = \bar{a}(t) + \bar{b}(t)(u(t) - 2.9 + 2(u(t) - 2.9)^2)$, where $\bar{a}(t)$ and $\bar{b}(t)$ as in the linear case are

evolvable parameters. When the nonlinear $\theta(t)$ function was changed into

$\theta(t) = 0.5 + u(t) - u_{ref} + 8(u(t) - u_{ref})^2$ the mean fitness in simulations with a linear

reaction norm turned out to be zero in several periods around $t = 1,500$, which illustrates that

there are limits to how much nonlinearities in the $\theta(t)$ function an evolutionary process is able

to track.

In the simulations I assumed 2,115 nonoverlapping generations and found $\bar{a}(t)$ and $\bar{b}(t)$ for

different initial values $\bar{b}(0)$. With a different time scale, similar $\bar{a}(t)$ and $\bar{b}(t)$ values could

plausibly be underlying samples of true and continuously varying parameter values, just as the

$\partial^{18}O(t)$ data used as $u(t)$ are samples of a continuous variable.

The time series in the real data case have samples from around 2 million years ago and

samples for the last 0.9 million years, and although it would have been nice with additional data

from around 1.4 million years ago the available data indicate a linear fitness peak vs.

environment function. It is in any case no indications of a very nonlinear function, and it seems

reasonable to assume that long existing species in the quite variable $\partial^{18}O$ environment in

general have fitness peak vs. environment functions that are not too difficult to track. This

would again be in accordance with the findings in de Villemereuil et al. (2020) regarding

breeding dates of 21 wild animals.

If the unfiltered $\partial^{18}O(t)$ data are used as inputs to the model for *log AZ area mean* found by

use of filtered $\partial^{18}O(t)$ data we would in the real data case obtain the results in Fig. 6, which

illustrates the sampling diffficulties involved in the Liow et al. project, and thus underlines the

need for filtering. For the $\partial^{18}O(t)$ filtering, I used a moving window with a fixed width of 300

samples, but since the data is collected with various window sizes a variable filtering window

size could possibly improve the predictions.

Figure 6. Predictions as in Fig. 4, panel B (red line), and with unfiltered $\partial^{18}O(t)$ data

used as inputs to the same prediction model (green line).

Finally, as pointed out in Lister (2021) adaptation to a changing environment can be

accomplished without plasticity. This has been verified by setting $G_{bb} = 0$ and $\bar{b}(0) = 0$ in the

simulations, which only resulted in some losses of mean fitness and a very minor change in

$\bar{y}_{filt}(t)$ as function of $u_{filt}(i)$. However, Liow et al. (2024) reports that plasticity plays a role in

the real data case, although it is unclear to which extent.

Acknowledgement

306   I thank University of South-Eastern Norway for support and funding.

References

Bell, M.A. 2013. Adaptive Landscapes, Evolution, and the Fossil Record. In The Adaptive

Landscape in Evolutionary Biology, Eds. Svensson, E. and R. Calsbeek. Oxford University

Press. https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

Boutin, S., and J. E. Lane. 2014. Climate change and mammals: evolutionary versus plastic

responses. Evolutionary Applications 7: 29–41

Charmantier, A., R. H. McCleery, L. R. Cole, C. Perrins, L. E. B. Kruuk, and B. Sheldon. 2008.

Adaptive phenotypic plasticity in response to climate change in a wild bird population.

Science 320:800–803

Charmantier, A., and P. Gienapp. 2014. Climate change and timing of avian breeding

and migration: evolutionary versus plastic changes. Evol. Appl. 7, 15–28.

https://doi.org/10.1111/eva.12126

Crozier, L. G., and J. A. Hutchings. 2014. Plastic and evolutionary responses to climate change

in fish. Evolutionary Applications 7:68– 87

de Villemereuil, P. et al. 2020. Fluctuating optimum and temporally variable

selection on breeding date in birds and mammals. Proc. Natl Acad. Sci. USA 117,

31 969–31 978. doi:10.1073/pnas.2009003117)

Ergon T. and R. Ergon. 2017. When three traits make a line: evolution of phenotypic plasticity

and genetic assimilation through linear reaction norms in stochastic environments.

J. Evol. Biol. 30:486-500. doi: 10.1111/jeb.13003

Ergon, R. 2022. The important choice of reference environment in microevolutionary

climate response predictions. Ecology and Evolution 12(4):e8836.

https://doi.org/10.1002/ece3.8836

Ergon, R. 2023. Dynamical BLUP modeling of reaction norm evolution, accommodating

changing environments, overlapping generations, and multivariate data. Ecology and

Evolution. 2023;13:e10194. https://doi.org/10.1002/ece3.10194

Goodnight, C. J. 2013. Wright's Shifting Balance Theory and Factors Affecting the Probability of

Peak Shifts. In The Adaptive Landscape in Evolutionary Biology, Eds. Svensson, E. and

R. Calsbeek. Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

Hunt, G. 2006. Fitting and comparing models of phyletic evolution: random walks and

beyond. Paleobiology 32:578–601.

James, F. C. 1983. Environmental component of morphological differentiation in birds.

Science 221:184–186

Johnson, R.A. and D.W. Wichern. 2008. Applied Multivariate Statistical Analysis.

Prentice-Hall, London

Kellogg, D. E. 1975. Character displacement in the radiolarian genus, Eucyrtidium.

Evolution 4:736–749.

Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation. J. Evol. Biol. 22, 1435-1446.

doi: 10.1111/j.1420-9101.2009.01754.x

Lisiecki, L. E. and M. E. Raymo. 2005. A Pliocene–Pleistocene stack of 57 globally distributed

benthic δ18O records. Paleoceanography 20, A1003. doi:10.1029/ 2004PA001071

Liow, L. H., A. Porto, and E. Di Martino. 2024. Trait-Fitness Associations via Fecundity and

Competition in a Two-Million-Year-Long Fossil Record. Am. Nat. 204/3.

https://doi.org/10.1086/73133

Lister, A. M. 2021. Phenotypic Plasticity in the Fossil Record. In Phenotypic Plasticity &

Evolution: Causes, Consequences, Controversies. Ed. *David W. Pfennig*.Imprint CRC Press

https://doi.org/10.1201/9780429343001

Merilä, J. and A. P. Hendry. 2013. Climate change, adaptation, and phenotypic

plasticity: the problem and the evidence. Evol. Appl. 7, 1-14.

doi: 10.1111/eva. 12137.

Teplitsky, C., J. A. Mills, J. S. Alho, J. W. Yarall, and J. Merilä. 2008. Bergmann's Rule and climate

change: disentangling environmental and genetic responses in a wild bird population.

Proceedings of the National Academy of Sciences U. S. A 105:13492–13496

Voje, K. L. 2023. Fitting and evaluating univariate and multivariate models

of within-lineage evolution. Paleobiology, 49(4), 2023, pp. 747–764 DOI: 10.1017/pab.2023.1

Williams, S. E., L. P. Shoo, J. L. Isaac, A. A. Hoffmann, and G. Langham. 2008. Towards an

Integrated Framework for Assessing the Vulnerability of Species to Climate Change. PLoS

Biol 6(12): e325. doi:10.1371/journal. pbio.0060325