Adaptive peak tracking as explanation of sparse fossil data across fluctuating ancient environments

Rolf Ergon

rolf.ergon@usn.no

University of South-Eastern Norway

May 8. 2025

Abstract

Species that have existed over millions of years have done so because they have been able to track peaks in an adaptive landscape well enough to survive and reproduce. Such optima are defined by the mean phenotypic values that maximize mean fitness, and they are predominantly functions of the environment, for example the sea temperature. The mean phenotypic values over time will thus predominantly be determined by the environment over time, and the trait history may be found in the fossil record. Here, I simulate such a tracking system, using both a basic non-plastic selection model and a univariate intercept-slope reaction norm model. I show how both linear and nonlinear mean phenotype vs. environment functions can be found also from quite sparse and short time series from the fossil record, and I discuss how this methodology can be extended to multivariate systems. The simulations include cases with a constraint on the individual trait values and with other factors than environment influencing the position of the adaptive peak. The methodology is finally applied on a time series of mean phenotypic values in a record of bryozoan *Microporella agonistes* fossils spanning 2.3 million years, using the $\partial^{18}O$ measure as proxy for sea water temperature.

From as few as eleven samples of mean phenotypic values found in the fossil record it was

possible to identify a linear mean phenotype vs. environment function, and to predict the

continuous mean phenotypic values as functions of time with prediction errors within or just

outside the standard errors of the observations. Leave-one-out cross validation gave

satisfactory results. It remains to verify predictions for longer time periods without known or

investigated fossil data.

Keywords: adaptive landscape, adaptive peak tracking, bryozoans, fitness landscape, fossil

record, optimal phenotypic values

1 Introduction

Evolutionary time series in the fossil record provide information on phenotypic changes over

millions of years. Such time series may consist of quite sparse samples, and they are

traditionally modeled by various forms of random walk processes (Hunt, 2006; Voje, 2023), or

by Ornstein-Uhlenbeck (OU) processes describing evolution of mean phenotypic values toward

fixed or moving peaks in the adaptive landscape (Voje, 2023). Here, the adaptive landscape and

adaptive peak metaphors assume that the mean population fitness is a function of possibly

multivariate mean phenotypic values, see Pigliucci (2013) for the historical background and a

clarifying discussion.

As an alternative to the traditional fossil time series modeling, I propose that adaptive peak

tracking may be a useful modeling approach, and that this requires two types of landscape

metaphors. First, I assume a fitness landscape where individual fitness is a function of a

possibly multivariate set of individual phenotypic values. Second, I assume an adaptive

landscape where the mean population fitness is a function of a possibly multivariate set of

mean phenotypic values. For simplicity I assume that the position of the individual fitness peak

is a function of a dominating environmental factor, and that the adaptive peak may have a

somewhat different position owing to non-symmetrical individual fitness or trait distribution

functions. I am thus not attempting to estimate the effect of various selective or other factors

on the position of adaptive optima (Hansen, 1997), but I just assume that various selective

factors over time keep the mean phenotypic values in the vicinity of the adaptive peak, and as

an example I will use the relative size of feeding zooids (autozooids) in encrusting cheilostome

bryozoans. Liow et al. (2017) studied communities of such organisms spanning more than 2

million years, and an interesting result was that six different species showed temporally

coordinated changes in average zooid sizes, suggesting that the different species responded to

a common external driver. Liow et al. pointed to sea temperature as a potential common driver,

but they found that the available time series of changes in mean zooid size lacked sufficient

length to rigorously test such a hypothesis. In my view, a study by means of an adaptive peak

tracking model as introduced above is clarifying.

For an introduction, Fig. 1 shows data from Liow et al. (2024) (see a more detailed

presentation in Section 4), where the $\partial^{18}O(t)$ measure is a widely used proxy for seawater

temperature (Lisiecki and Raymo, 2005). A natural question is now which type of model we

need for explanation of the *log AZ area mean* data in panel B given the $\partial^{18}O(t)$ data in panel A,

and as already mentioned an adaptive peak tracking model is in my view a viable alternative.

Figure 1. Raw $u(t) = \partial^{18}O(t)$ data (panel A, green line) and a centered moving average signal $u_{filt}(t)$ computed from a sliding window of length 100 samples along the time axis (panel A, red line, see Section 4 for details), as well as *log AZ area mean* (panel B, circles and dots) with standard errors as in Fig. S2 in Liow at al. (2024). Present time is used as zero point. Note that the moving average signal cannot capture the sharp decline of the $\partial^{18}O(t)$ value towards time zero.

A block diagram for the proposed adaptive peak tracking model is shown in Fig. 2, and as described above it is based on two types of landscape metaphors. The optimum phenotype that maximizes individual fitness is assumed to be a function $\theta_i(u, t) = f(u(t))$, where $u(t)$ is the dominating environmental variable (in the bryozoan example the $\partial^{18}O(t)$ measure). The optimum mean phenotypic value that maximizes mean fitness is assumed to be a function $\theta_m(u, t) = f(u(t))$, deviating from $\theta_i(u, t)$ when either the individual fitness function or the trait distribution is non-symmetrical. When both the fitness function and the trait distribution are symmetrical, we will thus have $\theta_m(u, t) = \theta_i(u, t)$ (see Section 2 for an example with Gaussian functions). The tracking process in Fig. 2 is attempting to keep the mean phenotypic value $\bar{y}(t)$ (in the bryozoan example the *log AZ area mean* value) in the vicinity of the optimal value

$\theta_m(u,t)$, such that the tracking error $e(t)$ is kept small. The basic assumption is thus that the

tracking process keeps the tracking error small, regardless of the details of this process.

Figure 2. Block diagram for fitness peak tracking system, with the environmental

variable $u(t)$ as input and the mean phenotypic value $\bar{y}(t)$ as output. The phenotypic

values $\theta_i(u,t)$ and $\theta_m(u,t)$ that maximize individual and mean fitness, respectively, are

together with the tracking error $e(t)$ internal signals in the system. See text for

explanations of blocks marked by $f(\cdot)$ symbols.

A fluctuating environment as shown in Fig. 1 corresponds to an adaptive landscape where the

adaptive peak position fluctuates, and how well the peak is tracked depends on the properties

of the evolutionary system. For a population with phenotypic plasticity the tracking error may be

kept small at all times, while a population without plasticity may only be able to track a moving

average as indicated in Fig. 1. However, a population without plasticity may also keep the

tracking error small, provided that additive genetic variances are large. Sparse fossil data will in

any case not be informative enough for predictions of strongly fluctuating mean phenotypic

values, and a remaining possibility is therefore to predict moving averages over a window size

that may be used as prediction parameter. In cases where the fossil phenotypic data are mean

values from given time periods, a natural choice is also to use mean environmental data from

these time periods as input to the tracking model (see Section 4 for example).

The assumption that the mean phenotypic value stays in the vicinity of the adaptive peak is

supported by findings regarding multiple species and traits indicating that "fossil lineages have

ascended adaptive peaks and remained at their summits as they shift position through time"

(Bell, 2013). An important reference is here Estes and Arnold (2007), who based on a large data

set found that "the underlying process causing phenotypic stasis is adaptation to an optimum

that moves within an adaptive zone with stable boundaries", which applied to the data in Fig. 1,

panel B, and considering the standard errors given, means that *log AZ area mean* moves

between approximately 11 and 12. Using a database over many contemporary populations,

Estes and Arnold also found that phenotypic means are typically close to the adaptive peak

(46% within 1 phenotypic standard deviation of the optimum, and 65% within 2 standard

deviations). Estes and Arnold (2007) based their results on tests with a single displacement of

the optimum, using an extensive set of data (Gingerich, 2001), but they acknowledged that the

actual optimum might move many times with a small total net displacement. That is a good

description of the optimal mean phenotype corresponding to the environmental data in Fig. 1,

panel A, with large short-term variations and a rather limited long-time net variation. I thus

assume that short-term environmental fluctuations as in, e.g., Fig. 1, panel A, result in

variations in optima within a stable adaptive zone (Uyeda et al., 2011). However, contrary to

Uyeda et al. I assume a causal tracking system and not a purely stochastic model. In real cases

there will of course be stochastic variations in the environmental variable and in the position of

the individual fitness peak, and additional stochastic factors that forces the adaptive peak to

deviate from the individual fitness peak. Such stochastic variations are included in simulations,

but further discussion of this topic is beyond the scope of this paper.

    A persistent system excitation in the form of short-term environmental fluctuations as

exemplified in Fig. 1, panel A, makes it unlikely that a tracking system as shown in Fig. 2 gets

stuck in a local optimum or in a valley in the adaptive landscape. As illustrated in simulations it

is more likely that the mean phenotypic values fluctuate around global optima, see however an

example of slow adaptation in subsection 3.5 and a discussion in Section 5.

In simulations I assume that the tracking process block in Fig. 2 can be described by use of

Lande's selection theory for populations without plasticity (Lande, 1979), which was also used

by Estes and Arnold (2007), but I also include examples with phenotypic plasticity (Lande,

2009). In real cases, however, other mechanisms such as, e.g., adaptive walks in response to

mutations may also be involved (Ch. 27, Walsh and Lynch, 2018). It is as already mentioned

essential to realize that although knowledge of the details of the tracking process may be

interesting, such knowledge is not necessarily needed for finding continuous predictions $\hat{\bar{y}}(t)$

based on sparse measurements of mean phenotypic values $\bar{y}(t)$.

Under the assumption that the mean trait values have remained centered around the mean

fitness peaks as they have shifted position through time, such that the following error $e(t)$ in

Fig. 2 has remained small relative to the width of the underlying mean fitness function, the

position of the mean fitness peak as function of the environment, $\theta_m(u, t) = f(u(t))$, can be

reconstructed from fossil samples of $\bar{y}(t)$, and provided that this function is sufficiently smooth

this can be done also when the fossil samples are sparse and at irregular time intervals. For the

bryozoan data in Fig. 1, it turns for example out that the mean trait value

$\bar{y} = log\ AZ\ area\ mean$ as function of the environment $u = \partial^{18}O$ can be modeled by a straight

line, from which follows continuous predictions of $\hat{\bar{y}}$ as function of time (see Section 4 for

details).

The prediction method outlined above is supported by simulations in Section 3, with the

theoretical background given in Section 2. The simulations are based on a reaction norm model

where I use the $\partial^{18}O(t)$ measure as environmental cue, and where the phenotypic plasticity

may be set to zero. This gives a connection between the simulations and the real bryozoan data

case in Section 4, in the sense that the same type of large but short-time fluctuations in the

input signal is used. Since both $\partial^{18}O(t)$ and $\bar{y}(t)$ in the simulations have large variances, and

since data from the fossil record often are mean values over both populations and time (Liow et

al., 2024), I will in the simulations use moving average filtered versions of these variables. The

simulations are intended to show that mean phenotype vs. environment functions can be

predicted from sparse data also in cases where the population in shorter time periods have

quite low mean fitness. The simulations also show that $\bar{y}(t)$ fluctuates around the optimal

value $\theta_m(u, t)$, as discussed above, although in some cases with a minor time lag. In order to

show that $\theta_m(u, t)$ may be different from $\theta_i(u, t)$, a case with a constraint on individual trait

values is included. In addition to theory necessary for the simulations, as given in Section 2, the

theory is extended to cover multivariate cases.

In Section 4, the prediction method is applied on recently presented data from an already

introduced field study of the bryozoan species *Microporella agonistes*, based on a fossil record

over 2.3 million years (Liow et al., 2024). Parameters in a linear mean phenotype vs.

environment function are here estimated from as few as 11 data points at irregular sampling

times. This is done using mean values of the $\partial^{18}O(t)$ measure in the time periods of collected

fossil data, but also, and with better results, using moving average mean environmental data.

Since the moving average mean as shown in Fig. 1, panel A, cannot capture the sharp decline of

the $\partial^{18}O(t)$ measure towards time zero, the best results are found by using a hybrid model with

the moving average mean for the last 0.02 million years replaced by the mean value of $\partial^{18}O(t)$

in that time period. Finally follows a summary and discussion in Section 5.

2 Theory

2.1 A basic univariate reaction norm case

For simplicity and for use in simulations I will initially assume that mean fitness is determined

by a univariate phenotypic mean value $\bar{y}(t)$, and that the adaptive landscape thus is two-

dimensional. I will later extend some essential results to multivariate cases, and the

simulations will also include examples where other factors than the physical environment are

involved.

In the simulations I will make use of an individual intercept-slope reaction norm model with

parameters $a(t)$ and $b(t)$ as given by

$$y(t) = a(t) + b(t)\big(u(t) - u_{ref}\big) + \varepsilon(t), \tag{1}$$

181   where $b(t)$ and the variance $G_{bb} = var(b(t))$ in some cases are set to zero. Here, $u(t)$ is the

182   environmental cue, while $\varepsilon(t)$ is assumed to be normally distributed with mean 0 and variance

183   σ_ε^2 among individuals in every generation. In Eq. (1), u_{ref} is a reference environment defined as

184   the environment at which the phenotypic variance is at a minimum (Lande, 2009). For a given

185   population this will result in the mean reaction norm

$$\bar{y}(t) = \bar{a}(t) + \bar{b}(t)\big(u(t) - u_{ref}\big). \tag{2}$$

186   The optimal individual phenotypic value is in general a function of the environment,

$$\theta_i(u, t) = f\big(u(t) - u_{ref}\big), \tag{3}$$

187   in the linear case $\theta_i(u, t) = \alpha + \beta\big(u(t) - u_{ref}\big)$, where α and β are constant parameters.

188   In the simulations I will assume an individual fitness function

$$W(t) = W_{max} \exp\left(-\frac{\big(y(t) - \theta_i(u, t)\big)^2}{2\omega^2}\right), \tag{4a}$$

189   which assuming that $y(t)$ is normally distributed gives the mean fitness in a population (Lande,

190   2009; Estes and Arnold, 2007)

$$\bar{W}(t) = W_{max} \sqrt{\frac{\omega^2}{\omega^2 + \sigma_y^2(t)}} \exp\left(-\frac{\big(\bar{y}(t) - \theta_i(u, t)\big)^2}{2\big(\omega^2 + \sigma_y^2(t)\big)}\right), \tag{4b}$$

where $\sigma_y^2(t)$ is the variance of $y(t)$ according to Eq. (1). The mean fitness will thus be

maximized when $\bar{y}(t) = \theta_i(u, t)$. Note that this is an example with symmetrical individual

fitness and trait distribution functions, such that optimal individual and mean phenotypic

values are equal, i.e., $\theta_m(u, t) = \theta_i(u, t)$. In Subsection 3.3 I will, however, also include a case

where $\theta_m(u, t) \neq \theta_i(u, t)$.

Under the assumptions of non-overlapping generations and random mating in a large

population the evolution of the mean reaction norm parameters is governed by (Lande, 1979)

$$\begin{bmatrix} \bar{a}(t+1) \\ \bar{b}(t+1) \end{bmatrix} = \begin{bmatrix} \bar{a}(t) \\ \bar{b}(t) \end{bmatrix} + \begin{bmatrix} G_{aa} & G_{ab} \\ G_{ab} & G_{bb} \end{bmatrix} \begin{bmatrix} G_{aa} + \sigma_\varepsilon^2 & G_{ab} \\ G_{ab} & G_{bb} \end{bmatrix}^{-1} \begin{bmatrix} cov(w(t), a(t) + \varepsilon(t)) \\ cov(w(t), b(t)) \end{bmatrix}, \quad (5)$$

where G_{aa}, G_{bb} and G_{ab} are variances and the covariance of the reaction norm parameters, and

where $w(t) = W(t)/\overline{W}(t)$ is the relative fitness. In a stationary stochastic environment, the

optimal reaction norm slope is $\bar{b}_{opt} = cov(\theta_i(u,t), u(t))/var(u(t))$ (Lande, 2009).

When the mean phenotypic trait in a population is tracking a moving optimum $\theta_m(u,t)$ it will

be given by $\bar{y}(t) = \theta_m(u,t) - e(t)$, where $e(t)$ is the tracking error (see block diagram in Fig. 2).

This means that the parameter values $\bar{a}(t)$ and $\bar{b}(t)$ are forced to evolve such that

$\bar{y}(t) \approx \theta_m(u,t) = f\big(u(t) - u_{ref}\big)$, i.e., such that the mean fitness according to Eq. (4b) is kept

close to the optimum and that the tracking error is small. In feedback control terminology it is

thus the mean trait tracking error $e(t) = \theta_m(u,t) - \bar{y}(t)$ that via mean fitness is the driving

force in the adaptive process. It follows from Eq. (5) that the tracking system in Fig. 2 has what

in feedback control terminology is called integral action (Åström and Murray, 2008), which

means that the tracking error in a stationary case with a fixed adaptive peak will asymptotically

go to zero. As shown in simulations in Section 3 the parameters in the

$\bar{y}(t) \approx \theta_m(u,t) = f\big(u(t) - u_{ref}\big)$ function can be estimated also from rather few fossil data

points $\big[u_{filt}(t), \bar{y}_{filt}(t)\big]$ at irregular sampling times, where $\bar{y}_{filt}(t)$ and $u_{filt}(t)$ are moving

average filtered versions of often quite noisy signals. In the real data case in Section 4 the

parameters in $\theta_m(u,t)$ are also found from data points $[u_{mean}(t), \bar{y}_{mean}(t)]$ for the time periods

of collected fossil data. Note that even though $\bar{y}(t) \approx f\big(u(t) - u_{ref}\big)$, such a model is not a

reaction norm, i.e., the reaction to a sudden change Δu in $u(t)$ will still be a sudden change

$\Delta \bar{y} = \bar{b}(t)\Delta u$.

The details in the tracking process in Fig. 2 as used in the simulations are described by Eqs.

(2), (4) and (5), but also if the parameters in those equations were known they could not be used

for computation of predictions $\hat{\bar{y}}(t)$ based on only a few fossil samples at irregular time points.

The remaining possibility for prediction of $\bar{y}(t)$ is to assume good enough tracking such that

$\bar{y}(t) \approx \theta_m(u,t) = f\big(u(t) - u_{ref}\big)$, which as shown in the simulations as well as in the real data

case makes it possible to estimate the parameters in the $\theta_m(u,t)$ function also from few

$\big[u_{filt}(t), \bar{y}_{filt}(t)\big]$ or $[u_{mean}(t), \bar{y}_{mean}(t)]$ data points. Based on those estimates and known

values of $u(t)$ it is then possible to predict $\bar{y}(t)$.

2.2 Influence of other factors than the environment

Eqs. (2) and (3) assume that $u(t)$ is the only factor that affects the mean phenotypic value $\bar{y}(t)$

and the optimal mean phenotypic value $\theta_m(u,t) = \theta_i(u,t)$, respectively. When other factors

affect $\bar{y}(t)$, the tracking system in Fig. 2 will force the parameters $\bar{a}(t)$ and $\bar{b}(t)$ in Eq. (2) to

evolve such that these factors are compensated for, and the result will be seen in the mean

phenotype vs. environment function that can be found from data, and then most likely in the

form of nonlinearities as shown in a simulation example in Subsection 3.3. When other factors

affect $\theta_m(u,t)$ it is again the $\bar{a}(t)$ and $\bar{b}(t)$ parameter values that must compensate, also then

most likely in the form of nonlinearities as shown in Subsection 3.4. A linear mean phenotype

vs. environment function thus indicates that no other factors than the environment affect the

fitness vs. environment function, although there may of course be several other factors that

compensates the effects of each other. And in cases where $\theta_i(u,t)$ is a nonlinear function there

may be factors that make $\theta_m(u,t)$ linear.

2.3 Multivariate cases

When the mean phenotypic trait in a univariate population is tracking the optimal phenotypic

value, the mean phenotypic trait can as shown above be described by

$\bar{y}(t) \approx \theta_m(u,t) = f\big(u(t) - u_{ref}\big)$, i.e., by the adaptive peak vs. environment function. Here, it is

again important to note that we do not need to know the details of the tracking process in Fig. 2.

The univariate theory can be extended to multivariate cases, and although we in practice do

not need to know the underlying fitness function, we may for a theoretical discussion use an

example with two phenotypic traits and the bivariate fitness function (Ch. 4, Johnson &

Wichern, 2008)

$$W(t) = W_{max} \exp\left(-[y_1(t) - \theta_{i,1}(u,t) \quad y_2(t) - \theta_{i,2}(u,t)]\Sigma^{-1}(t)\begin{bmatrix} y_1(t) - \theta_{i,1}(u,t) \\ y_2(t) - \theta_{i,2}(u,t) \end{bmatrix}/2c\right), \quad (6)$$

248   where $\Sigma(t) = \text{cov}(y_1(t), y_2(t))$ is the variance-covariance matrix and where c is a scaling factor

249   controlling the 'width' of the fitness function. In order to keep the mean traits in the vicinity of

250   the adaptive peak, they must in this case continuously evolve such that $\bar{y}_1(t) \approx \theta_{m,1}(u,t)$ and

251    $\bar{y}_2(t) \approx \theta_{m,2}(u,t)$, and we will thus find

$$\bar{y}_1(t) \approx f_1\big(u(t) - u_{ref}\big) \qquad\qquad (7a)$$

$$\bar{y}_2(t) \approx f_2\big(u(t) - u_{ref}\big), \qquad\qquad (7b)$$

where the moving adaptive peak vs. environment functions may be linear or nonlinear, and

where the parameters can be estimated from rather few fossil data points at irregular sampling

times. Here, the essential point is that $\bar{y}_1(t)$ and $\bar{y}_2(t)$ in this example are tracking a peak in a

three-dimensional adaptive landscape, and that this tracking process can be decomposed into

two univariate tracking processes. We may thus consider the projections of the bivariate fitness

function onto orthogonal axes for the two phenotypic traits, and such a projection approach

may be extended to systems with higher dimensions, as discussed in Goodnight (2013). Note

that we here are only interested in the $\theta_{m,1}(u,t)$ and $\theta_{m,2}(u,t)$ functions that define the position

of the adaptive peak, and that we do not need to know the shape of the adaptive landscape

around the peak.

3 Simulations

3.1 Linear reaction norm example

A system with a linear mean reaction norm $\bar{y}(t) = \bar{a}(t) + \bar{b}(t)\big(u(t) - u_{ref}\big)$ was simulated with use of the 2,115 available samples of $u(t) = \partial^{18}O(t)$, and with population size 400. These samples go 5.3 million years back in time with various sampling intervals (Lisiecki and Raymo, 2005), but here I just assume 2,115 nonoverlapping generations (however, see Section 5 for a discussion on relevance for fossil data). At each new generation random individual samples around $\bar{a}(t)$, $\bar{\varepsilon}(t) = 0$ and $\bar{b}(t)$ were drawn from normal populations with the given variances. The fitness function in Eq. (4a) had parameter values $W_{max} = 1$ and $\omega^2 = 50$. The optimal mean phenotypic value was in a first linear case chosen as $\theta_m(u, t) = \theta_i(u, t) = u(t) - 2.9$, where $\theta_m(u, t)$ thus and for clarity of presentation is assumed to be a deterministic function, and where 2.9 is the mean value of $\partial^{18}O(t)$ around $t = 0$ (however, see results with added white noise below).

As it is not realistic that $u(t)$ and $\theta_m(u, t)$ are purely deterministic signals, correlated white noise with variances $\sigma_u^2 = 0.05$ and $\sigma_\theta^2 = 0.2$, respectively, were added to $\theta_m(u, t)$ and $u(t)$. The covariance was $\sigma_{\theta,u} = 0.025$, see Lande (2009) for a possible explanation.

In a first case without phenotypic plasticity, the reaction norm parameter variances and covariance were chosen as $G_{aa} = 2$, $\sigma_\varepsilon^2 = 0.5G_{aa} = 1$, $G_{bb} = 0$ and $G_{ab} = 0$, while the initial parameter values were $\bar{a}(0) = \bar{b}(0) = 0$. Since $\bar{y}(t)$, $\theta(u, t)$ and $u(t) = \partial^{18}O(t)$ have large variances, I used moving average filtered signals $\bar{y}_{filt}(t)$, $\theta_{filt}(u, t)$ and $u_{i,filt}(t)$ in some of the plots. These were obtained by moving window filters with window size 100, in MATLAB notation $\bar{y}_{filt}(t) = movmean(\bar{y}(t), 100)$, $\theta_{i,filt}(u, t) = movmean(\theta_i(u, t), 100)$ and

$u_{filt}(t) = movmean(u(t), 100)$. Simulation results in the linear case without plasticity are

shown in Fig. 3, panels A to E. Note that $\bar{y}(0) = \bar{a}(t)$ follows the main trend in $u(t)$, and that the

fluctuations are somewhat delayed compared to the fluctuations in $u(t)$ (panel A).

In a second case with phenotypic plasticity, the reaction norm parameter variances and

covariance were chosen as $G_{aa} = 2$, $\sigma_\varepsilon^2 = 1$, $G_{bb} = 2$ and $G_{ab} = 0$, while the initial parameter

values were $\bar{a}(0) = 0$ and $\bar{b}(0) = 0.8$. Simulation results in the linear case with plasticity are

shown in Fig. 3, panels F to J. As follows from the reaction norm equations (1) and (2), the

variance $\sigma_y^2(t)$ will increase with $u(t) - u_{ref}$, and as a result the mean fitness will be reduced

(panel H).

In both cases (without and with plasticity) $\bar{y}_{filt}(t)$ tracks $\theta_{m,filt}(u,t) = \theta_{i,filt}(u,t)$ closely

(panels D and I), such that the mean fitness is kept at a high level (panels C and H). This results

in linear functions $\bar{y}_{filt}(t) = f\left(u_{filt}(t)\right)$ (panels E and J), with seven discrete values marked by

circles. Using $u_{ref} = u_{filt}(0) = 2.9$, these discrete values could be used for determination of

the parameters $\hat{\alpha}$ and $\hat{\beta}$ of this line by an ordinary least squares method, and thus to find

predictions $\hat{\bar{y}}_{filt}(t) = \hat{\alpha} + \hat{\beta}\left(u_{filt}(t) - u_{ref}\right)$, see examples in Section 4.

Figure 3. Simulation results with the linear $\theta_m(u,t) = \theta_i(u,t) = u(t) - 2.9$ function,

without plasticity (panels A to E) and with plasticity (panels F to J). Panels A and F show

$u(t)$ (green lines) and $\bar{y}(t)$ (blue lines), while the red line is $u_{filt}(t)$. Panels B and G show

$\bar{a}(t)$ (solid lines) and $\bar{b}(t)$ (dashed lines), while panels C and H show mean fitness $\bar{W}(t)$.

Panels D and I show $\theta_{i,filt}(u,t)$ (blue lines) and $\bar{y}_{filt}(t)$ (red lines). Panels E and J show

that $\bar{y}_{filt}(t)$ as function of $u_{filt}(t)$ in both cases is the same straight line, with values for

samples 200, 400, 600, 800, 1000, 1200 and 1400 shown by circles.

3.2 Nonlinear reaction norm example

Fig. 4 shows results as in Fig. 3, but with the nonlinear adaptive peak vs. environment function

$\theta_m(u,t) = \theta_i(u,t) = (u(t) - 2.9) + 4(u(t) - 2.9)^2$. Note that the mean fitness now is reduced

(panels B and F), especially in the case without plasticity. The signal $\bar{y}_{filt}(t)$ is also in this case

tracking $\theta_{filt}(u,t)$ well, although with a noticeable time lag (panels C and G). Here, $\bar{y}_{filt}(t)$ as

function of $u_{filt}(t)$ is the same second order function in the different cases (panels D and H).

314   **Figure 4**. Simulation results as in Fig. 3, but with

315     $\theta_m(u,t) = \theta_i(u,t) = (u(t) - 2.9) + 4(u(t) - 2.9)^2$. The discrete values in panels D and

316   H could here be used to determine the nonlinear function $\hat{\bar{y}}_{filt}(t) = f\left(u_{filt}(t)\right)$ by

317   means of any suitable fitting method.

3.3 Example with phenotypic constraint

With a constraint on the individual phenotypic values the distribution will no longer be normal,

and the adaptive peak $\theta_m(u,t)$ will therefore deviate from the individual fitness peak such that

Eq. (4b) no longer is correct. The mean trait $\bar{y}(t)$ will, however, still track the adaptive peak that

maximizes mean fitness. When there for example is an upper limit for the individual phenotypic

trait values as shown in Fig. 5, the $\bar{a}(t)$ and $\bar{b}(t)$ parameters will evolve such that

$\bar{y}(t)$ deviates from $\theta_m(u,t)$ with results as shown in Fig. 6, panels A to D. The effect of the

constraint is clearly seen in panel C in Fig. 6, in that $\bar{y}_{filt}(t)$ does not follow $\theta_{i,filt}(u,t)$, but

instead follows $\theta_{m,filt}(u,t)$ (not shown) such that the mean fitness as shown in panel B is

maximized. The effect is also seen by a comparison of panels A in Fig. 6 and panel G in Fig. 3, in

that $\bar{a}(t)$ evolves into a large value, such that $\bar{y}(t)$ without constraints on $y(t)$ would become

larger than $\theta_m(u, t)$. Note that the mean fitness is increased compared to the results in Fig. 3,

panel H, because the variance $\sigma_y^2(t)$ is reduced. Also note that the mean phenotype vs.

environment functions in panel D now is nonlinear in a way that with knowledge of the

biological system possibly could be interpreted in a meaningful way. Note that individual trait

values will have a larger variance than the mean trait value, such that the constraint in Fig. 5 has

an effect before $\bar{y}(t) \approx \theta(u, t)$ reaches the upper limit.

Figure 5. The linear function $\theta_m(u, t) = u(t) - 2.9$ as used for the results in Fig. 3 (blue

line) and an upper constraint for the individual trait values (red line).

3.4 Example with additional term in $\theta(u, t)$

The optimal value $\theta_m(u, t)$ that maximizes mean fitness may not be a function of only the

environment, and as an example Fig. 6, panels E to H, shows results as in Fig. 3, panels F to J,

but with $\theta_m(u, t) = \theta_i(u, t) = u(t) - 2.9 + 2t/2115$. Note that panel H now shows a nonlinear

function.

Figure 6. Panels A to D show simulation results as with plasticity in Fig. 3, but with a constraint on the individual phenotypic trait values as shown in Fig. 5. Panels E to H show simulation results as with plasticity in Fig. 3, but with

$$\theta_m(u,t) = \theta_i(u,t) = u(t) - 2.9 + 2t/2115.$$

3.5 Example without plasticity, but with changed parameter values

Simulations as in Fig. 3, panels A to E, were repeated but with changed parameter values. First, G_{aa} and σ_ε^2 were reduced from 2 and 1 to 0.2 and 0.1, with results as in Fig. 7, panels A to E. As seen in panels A and F the mean phenotypic value $\bar{y}(t)$ cannot now follow the adaptive peak $\theta_m(u,t)$), and as a result the mean phenotype vs. environment function in panel E becomes non-linear. However, owing to the broad fitness function with $\omega^2 = 50$, the mean fitness is still close to one.

  Second, with $G_{aa} = 2$ and $\sigma_\varepsilon^2 = 1$ as in Fig. 3, the width of the fitness function was reduced by reducing ω^2 from 50 to 5, with results as in Fig. 7, panels F to J. The mean phenotypic value still

358   follows the adaptive peak but owing to the narrow fitness function the mean fitness is clearly

359   reduced (panel H).

361   **Figure 7**. Panels A to E show simulation results as in Fig. 3 with plasticity, but with G_{aa}

362   and σ_{ε}^2 reduced by a factor 10. Panels F to J show simulation results as in Fig. 3 with

363   plasticity, but with the width of the fitness function reduced from $\omega^2 = 50$ to $\omega^2 = 5$.

4 Real data case

Liow et al. (2024) recently presented results from a field study of the bryozoan species

Microporella agonistes, based on a fossil record over 2.3 million years (see Fig. 1 for some

results). The fossils were collected from colonies in New Zealand cliffs, and it is especially

interesting that the same species still live in the waters near those cliffs. In a project running

over several years, Liow and her coworkers collected 985 fossil colonies with a large total

number of individuals and developed an artificial intelligence tool for data retrieval by use of

pictures from a scanning electron microscope. This image analysis system can rapidly assess

three well defined phenotypic traits; autozooid (AZ) area, ovicell (OV) area and autozooid

shape, and as seen from time series in Fig. S2 in Liow et al. (2024) the two AZ and OV area mean

trait values increase with time except for a drop in recent time owing to an extreme $\partial^{18}O(t)$

value.

Since the *log AZ area mean* and *log OV area mean* data in Fig. S2 are linearly dependent, I will

here focus on the former. The last two data points in this time series are according to Table S1

found from fossils collected at time zero, with time span (duration) zero, but that is not

consistent with Fig. 2, panel A, in Liow et al. (2024). After comparison with data from other time

periods I therefore assume that the samples behind these two mean values are within time

spans of approximately 0.05 and 0.02 million years. The fossil data are thus collected from

eleven periods back in time according to Table 1, which for the purpose of comparison with

moving average mean data also includes the time window size for 100 samples. The reason for

use of the natural logarithm of the AZ area is that it gives normally distributed data (Di Martino

and Liow, 2021).

Table 1. Fossil collection time data

sample number	mean sample time point	window size	window size for 100 samples
1	−2.1895	0.203	0.25
2	−1.9660	0.102	0.25
3	−1.8505	0.049	0.25
4	−0.9265	0.019	0.2
5	−0.6485	0.055	0.2
6	−0.5480	0.030	0.2
7	−0.5055	0.055	0.1
8	−0.4510	0.054	0.1
9	−0.3990	0.050	0.1
10	−0.025	0.050	0.1
11	−0.010	0.020	0.1

Since the *log AZ area mean* values are given as mean values from samples within time

windows as given in Table 1, it is natural to use mean values $u_{mean}(t)$ of $\partial^{18}O$ in the same

windows. Fig. 8 shows $u(t) = \partial^{18}O(t)$ and $y(t) = log\ AZ\ area\ mean\ (t)$, as well as predictions

$\hat{y}(t)$ based on least squares fitting of a straight line in a plot over $y(t)$ as function of $u_{mean}(t)$.

The slope of the prediction line is 0.20, and the mean squares prediction error

$MSE = \left(\bar{y}(t) - \hat{y}(t)\right)^2/(number\ of\ samples)$ is $MSE = 0.0075$ (Table 2).

Fig. 9 shows the corresponding results based on centered moving average filtering of $\partial^{18}O(t)$.

The optimal moving window size was 90, resulting in prediction line slope 0.39 and prediction

error $MSE_{min} = 0.0051$. However, note that the way the $\partial^{18}O(t)$ data are organized in Lisiecki

and Raymo (2005) means that a fixed window size given by number of samples corresponds to

different time durations as shown in Table 1.

As seen in especially Fig. 9, it is the very last value $\bar{y}(t)$ that is difficult to predict (sample 11 in Table 1, population 19.2 in Liow et al., 2024). However, when this sample is removed the results with use of mean $u(t)$ for the various time periods (as in Fig. 8) changed very little because the last sample has little statistical leverage. The MSE value with use of moving average filtering of $u(t)$ (as in Fig. 9), on the other hand, changed a lot (Table 2) because the last sample has large statistical leverage.

A special problem with the last sample in Table 1 is that moving average filtering of $u(t)$ cannot capture the sharp decline in $u(t)$ towards time zero. If $u_{filt}(t)$ for the last 0.02 million years is replaced by the mean value of $u(t)$ in that period, i.e., a hybrid model, the last sample is predicted with the very small error 0.01, while MSE_{min} is improved from 0.0049 to 0.0022 (Table 2). Fig, 10 shows results for the hybrid model.

Figure 8. Panel A shows raw $\partial^{18}O(t)$ data (green line) with values in fossil collection time windows marked by red and blue, and with mean values marked by circles with dots. Panel B shows *log AZ area mean* with error bars as found from Fig. S2 in Liow at al. (2024) with corrections for the two last points as given in Table 1 (blue circles with dots and blue line), as well as predictions (red circles with dots and red line). Panel C shows

prediction line (red) computed by the ordinary least squares method, with the two last

samples marked by solid circles. See prediction results in Table 2.

Figure 9. Results as in Fig. 8, but based on centered moving average filtering of $\partial^{18}O(t)$

and with continuous predictions of *log AZ mean area* values (red line in panel B). See

prediction results in Table 2.

The prediction results for the hybrid model described above and shown in Fig. 10, i.e., with

$u_{filt}(t)$ for the last 0.02 million years replaced by the mean value of $u(t)$ in that period, were

validated by leave-one-out cross validation, with results as in Fig. 10 and Table 2.

Figure 10. Results for hybrid model with $u_{filt}(t)$ for the last 0.02 million years replaced

by the mean value of $u(t)$ in that period, and for leave-one-out cross validation. Solid

red line shows prediction results, while dashed red line and red circled with dots show

validation results for the left-out samples. See prediction and validation results in Table

2.

Table 2. Results with different prediction models.

	Model with use of u_{mean} Fig. 8	Model with use of u_{mean}, without sample 11	Model with use of u_{filt} Fig. 9	Model with use of u_{filt}, without sample 11	Hybrid Model, Fig. 10
b	0.20	0.20	0.39	0.51	0.48
MSE	0.0075	0.0081	0.0049	0.0015	0.0015
b_{CV}	-	-	-	-	0.48
MSE_{CV}	-	-	-	-	0.0022

As shown in Figs. 1, 8, 9 and 10 the measurements of *log AZ area mean* have various standard

errors. Repeated simulations of the final hybrid model as used for Fig. 10, with discrete values

drawn from normal distributions with standard deviations equal to the standard errors, gave

results as in Fig. 11. As can be seen in the upper panel there is ample space for nonlinear

prediction functions, although they would be less likely than the fitted linear function marked by

blue line. From the histogram in the lower panel follows that the prediction line slope value

$b = 0.48$ has a standard error of approximately $\mathrm{SE} = 0.1$.

Figure 11. Upper panel shows linear prediction lines for *log AZ area mean* from 100

simulations with measurement values drawn from normal distributions of $\bar{y}(t)$ with the

given standard errors (cyan lines), and for the mean prediction line corresponding to Fig.

10 (blue line with error bars). Lower panel shows histogram over slopes of fitted

prediction lines as in upper panel, but for 4,000 simulations, indicating that the

prediction line slope is $b = 0.48 \pm 0.1$.

5 Summary and Discussion

5.1 Main points

In this paper I make use of two landscape metaphors. First, an individual fitness landscape

where the peak of the individual fitness as function of individual phenotypic values moves with

changes in the environment. Second, an adaptive landscape where the peak of mean fitness as

function of mean phenotypic values moves with changes in the environment. The adaptive peak

will essentially follow the individual fitness peak, but with deviations caused by non-

symmetrical fitness and trait distribution functions and possibly also owing to other causes.

I also make some other fundamental assumptions. First, species that have existed for

millions of years in a fluctuating environment have by means of various selection mechanisms

been tracking a limited movement of an optimum in the adaptive landscape, and they have

done so well enough to survive and reproduce. This assumption is supported by results in Estes

and Arnold (2007) and other references in the introduction. Second, the location of the

individual fitness peak is a simple continuous function of a dominating environmental variable,

e.g., the temperature, such that the movement of the peak in the adaptive landscape is

continuous without sudden jumps. Third, the dominating environmental variable fluctuates in a

way that makes it unlikely that the mean phenotypic value gets stuck in local minima or valleys

in the adaptive landscape.

Under the assumptions given, two main conclusions can be drawn from the results in this

article: First, adaptive vs. environment functions, and thus approximate mean phenotype vs.

environment functions can be found from sparse and short fossil data, provided that they are

sufficiently spread over the time period of interest. As demonstrated in the real data example in

Section 4, the environmental data used for prediction of mean phenotypic values may be

means from time periods of fossil collection or a moving average mean. Second, prediction of

mean phenotypic values does not require detailed knowledge of the tracking process in Fig. 2,

i.e., neither parameter values in an underlying reaction norm or adaptive walk model nor the

fitness functions need to be known. Also note that it is not necessary to find explicit

mathematical adaptive peak vs. environment functions, predictions may instead be found by

use of spline interpolation and similar methods.

Even if the position of the individual fitness peak is a linear or other simple function of the

environment, the adaptive peak vs. environment function may be less simple. This is because

non-symmetrical fitness functions and/or trait distributions may drive the adaptive peak away

from the individual fitness peak. A closer look at the tracking process might thus be necessary

for interpretations of nonlinear mean phenotype vs. environment results. A deeper

understanding of the adaptive landscape may also be needed in cases with slow adaptation,

where the mean phenotypic value can track the adaptive peak only with a large time lag.

5.2 Discussion of simulation results

The main purpose of the simulations is to show that the mean phenotypic values over time as

functions of mean environment over time (panels E and J in Fig. 3 and panels D and H in Fig. 4)

are determined by the adaptive peak vs. environment function, and thus independent of the

details of the tracking process in Fig. 2. The variances of the reaction norm parameters are

chosen such that there is a clear difference in the evolutionary responses between cases

without and with plasticity. If these variances are increased by a factor ten (or a factor one

hundred) this difference very much disappears in that $\bar{y}(t)$ in both cases appear to follow the

adaptive peak $\theta_m(t) = f(u(t))$ instantaneously, but the mean phenotypic values over time as

functions of mean environment over time are essentially unchanged.

Figs. 3 and 4 show that although a population without plasticity may not be able to track rapid

changes in environment it may track a moving average quite well, and that the resulting mean

phenotype prediction functions, linear or non-linear, will be the same as for a population with

plasticity. Note that the mean plasticity slope with a linear fitness peak vs. environment

function approaches an optimal value (Fig. 3, panel F), such that the linear mean phenotype vs.

environment function seen in panel J can be interpreted as the result of adaptive phenotypic

plasticity. However, a comparison with results without plasticity (panel E) shows that this

interpretation may be wrong.

Fig. 6, left panels, illustrates that non-symmetrical trait distributions result in non-linear

mean phenotype vs. environment functions also in cases where the individual fitness peak

position is a linear function of a dominating environmental factor. Fig. 6, right panels, illustrates

that other factors than the environment may contribute to non-linearities in the mean

phenotype vs. environment function.

Panel 7, left panels, illustrates a case with slow adaptation as in, e.g., Toljagic et al. (2018),

where a mean trait value lags behind changes in the adaptive peak. The result is a non-linear

mean phenotype vs. environment function, but also in this case mean phenotypic values over

time can be predicted from a limited number of samples. Panel 7, right panels, shows that a

narrow fitness function, as must be expected results in reduced mean fitness.

5.3 Discussion of the real data case

The time series in the real data case in Section 4 has samples from around 2 million years ago

and samples for the last 0.9 million years, and although it would have been nice with additional

data from around 1.4 million years ago the available data indicate a linear adaptive peak vs.

environment function. Considering the large standard errors in the *log AZ area mean* data it is in

any case no clear indications of a very nonlinear function. It also seems reasonable to assume

that long existing species in the quite variable $\partial^{18}O$ environment in general have adaptive peak

vs. environment functions that are not too difficult to track, at least in the sense that the mean

phenotypic value is fluctuating around a moving average as in Fig. 3, panel A.

The results in Fig. 9 show that the mean phenotypic values can be predicted quite well with

use of a linear adaptive peak vs. environment function and moving average means of the $\partial^{18}O$

data. The exception is the very last point in the time series (population 19.2 in Liow et al., 2024),

found after a minor correction of a data inconsistency in Liow et al. (2024). Note that Fig. 9 gives

continuous predictions. Use of mean $\partial^{18}O$ values from the time periods of fossil collection

worked less well (Fig. 8), indicating that the raw data points in each sample period (as shown in

Fig. 2 in Liow et al., 2024) should not all be given the same weight.

There is a special problem with the very last sample in Fig. 9 in that moving average filtering of

$u(t)$ cannot capture the sharp decline in $u(t)$ towards time zero. If $u_{filt}(t)$ for the last 0.02

million years is replaced by the mean value of $u(t)$ in that period (a hybrid solution), the last

sample is predicted with a very small error while MSE is very much improved. As shown in Fig.

10, this solution gives predictions that fall within or for one sample just outside of the error

bars. The prediction results for the hybrid model were validated by leave-one-out cross

validation, with results included in Fig. 10. Validation gave $MSE_{CV} = 0.0022$, which should be

compared with $MSE_{min} = 0.0015$ for predictions. As shown in Fig. 10, the validation results are

thus quite satisfactory in that all predictions of the left-out samples except two fall within the

error bars.

Note that although the predictions in Fig. 10 are quite good as compared with the error bars

there are other possible errors that are not covered by these bars, such as fluctuations around

the moving adaptive peak due to imperfect tracking, and random fluctuations of the adaptive

peak itself.

It is of course an underlying assumption that the continuous mean trait predictions in Fig. 10

not only give values that can be compared to mean values from collected data, but that also

mean values in time periods without a known or investigated fossil record are predicted. That

remains to be verified, and it would be especially interesting if the pronounced variations

around 1.5 million years ago could be reproduced.

Liow et al. (2024) report that plasticity plays a role in the real data case, although in a complex

way owing to the clonal and colonial nature of the data. There is in fact a theoretical possibility

that the apparently linear mean phenotype vs. environment function is caused by adaptive

plasticity such that the mean phenotypic value tracks the fluctuations in sea temperature with

very small errors. However, as pointed out above, a linear mean phenotype vs. environment

function does not necessarily imply plasticity.

It is finally worth noting that just by looking at Fig. 1 it might be tempting to approximate the

mean phenotypic values as a linear function of the moving average of the environment. The

results in Fig. 10 could thus be found by an *ad hoc* least squares approach, without any

consideration of the theoretical background. However, the discussion of the tracking model in

Fig. 2 shows that the prediction results have a more general theoretical foundation.

Acknowledgement

559 I thank two anonymous reviewers for constructive criticism, and I thank University of South-

560 Eastern Norway for support and funding.

Author contribution

562 Rolf Ergon is the sole author of this article.

Data availability statement

564 MATLAB code and data are archived on bioRxiv, doi.org/10.1101/2024.10.30.621046.

Conflict of interest statement

566 The first author of Liow et al. (2024) which is a main reference, Lee Hsiang Liow, is my daughter-

567 in-law. Otherwise, there are no conflicts of interest.

ORCID

569 *Rolf Ergon* https://orcid.org/0000-0003-2393-2997

References

Åström, K. J., & Murray, R. M. (2008). Feedback systems: an introduction for scientists and

engineers. Princeton, NJ: Princeton University Press.

Bell, M. A. (2013). Adaptive Landscapes, Evolution, and the Fossil Record. In The Adaptive

Landscape in Evolutionary Biology, Eds. Svensson, E. and R. Calsbeek. Oxford University

Press. https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

Di Martino, E., & Liow, L. H. (2021). Larger offspring associated with lower temperatures across

species of Microporella, a widespread colonial invertebrate. Marine Ecology Progress Series

662:1–13

Estes, S. & Arnold, S. J. (2007). Resolving the paradox of stasis: models with stabilizing selection

explain evolutionary divergence at all timescales. American Naturalist, 169, 227–244.

Gingerich, P. D. (2001). Rates of evolution on the time scale of the evolutionary process.

Genetica 112/113:127–144.

Goodnight, C. J. (2013). Wright's Shifting Balance Theory and Factors Affecting the Probability of

Peak Shifts. In The Adaptive Landscape in Evolutionary Biology, Eds. Svensson, E. and

R. Calsbeek. Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

Hansen, T. F. (1997). Stabilizing selection and the comparative analysis of adaptation.

Evolution 51:1341–1351.

Hunt, G. (2006). Fitting and comparing models of phyletic evolution: random walks and

beyond. Paleobiology 32:578–601.

Johnson, R. A., & Wichern, D. W. (2008). Applied Multivariate Statistical Analysis.

Prentice-Hall, London

Lande, R. (1979). Quantitative genetic analysis of multivariate evolution, applied to

brain:body size allometry. Evolution 33:402-416

Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic

plasticity and genetic assimilation. J. Evol. Biol. 22, 1435-1446.

doi: 10.1111/j.1420-9101.2009.01754.x

Lisiecki, L. E., & Raymo, M. E. (2005). A Pliocene–Pleistocene stack of 57 globally distributed

benthic δ18O records. Paleoceanography 20, A1003. doi:10.1029/ 2004PA001071

Liow, L. H., Porto, A. & Di Martino, E. (2024). Trait-Fitness Associations via Fecundity and

Competition in a Two-Million-Year-Long Fossil Record. Am. Nat. 204/3.

https://doi.org/10.1086/73133

Liow, L. H., Di Martino, E., Krzeminska, M., Ramsfjell, M. H., Rust, S., & Taylor, P. D.

(2017). Relative size predicts competitive outcome through 2 million years.

Ecology Letters. ISSN 1461-023X. 20(8), p. 981–988. doi: 10.1111/ele.12795.

Lister, A. M. (2021). Phenotypic Plasticity in the Fossil Record. In Phenotypic Plasticity &

Evolution: Causes, Consequences, Controversies. Ed. *David W. Pfennig*.Imprint CRC Press

https://doi.org/10.1201/9780429343001

Pigliucci, M. (2013). Landscapes, Surfaces, and Morphospaces: What Are They Good For? In

The Adaptive Landscape in Evolutionary Biology, Eds. Svensson, E. and R. Calsbeek. Oxford

University Press. https://doi.org/10.1093/acprof:oso/9780199595372.001.0001

Uyeda, J. C., Hansen, T. F., Arnold, S. J., & Pienaar, J. (2011). The million-year wait for

macroevolutionary bursts. Proceedings of the National Academy of Sciences.

doi:10.5061/dryad.7d580

Voje, K. L. (2023). Fitting and evaluating univariate and multivariate models

of within-lineage evolution. Paleobiology, 49(4), (2023), pp. 747–764

DOI: 10.1017/pab.2023.1

Walsh, B., & Lynch, M. (2018). Evolution and selection of quantitative traits.

Oxford University Press