Functionally Important Residues from Graph Analysis of Coevolved Dynamic couplings

Manming Xu,[1*] Sarath Chandra Dantu,[2*] James A Garnett,[3] Robert A Bonomo,[4,5,6,7,8,9] Alessandro Pandini ,[2,10^] and Shozeb Haider[1,10,11^]

[1] UCL School of Pharmacy, London WC1N 1AX UK
[2] Department of Computer Science, Brunel University London, Uxbridge UB8 3PH, UK
[3] Centre for Host-Microbiome Interactions, Faculty of Dentistry, Oral & Craniofacial Sciences, King's College London, London, UK
[4] Research Service, Louis Stokes Cleveland Department of Veterans Affairs Medical Center, Cleveland, OH, USA;
[5] Department of Molecular Biology and Microbiology, Case Western Reserve University School of Medicine, Cleveland, OH, USA;
[6] Department of Medicine, Case Western Reserve University School of Medicine, Cleveland, OH, USA;
[7] Clinician Scientist Investigator, Louis Stokes Cleveland Department of Veterans Affairs Medical Center, Cleveland, OH, USA;
[8] Departments of Pharmacology, Biochemistry, and Proteomics and Bioinformatics, Case Western Reserve University School of Medicine, Cleveland, OH, USA;
[9] CWRU-Cleveland VAMC Center for Antimicrobial Resistance and Epidemiology (Case VA CARES) Cleveland, OH, USA;
[10] University of Tabuk (PFSCBR), Tabuk, Saudi Arabia
[11] UCL Center for Advanced Research Computing, University College London, WC1H 9RL, U.K.

* Joint first authors

^ To whom correspondence should be addressed:
Prof Shozeb Haider (Shozeb.haider@ucl.ac.uk)
Dr Alessandro Pandini (alessandro.pandini@brunel.ac.uk)

ORCID:
Shozeb Haider: 0000-0003-2650-2925
Alessandro Pandini: 0000-0002-4158-233X

Abstract

The relationship between protein dynamics and function is essential for understanding biological processes and developing effective therapeutics. Functional sites within proteins are critical for activities such as substrate binding, catalysis, and structural changes. Existing computational methods for the predictions of functional residues are trained on sequence, structural and experimental data, but they do not explicitly model the influence of evolution on protein dynamics. This overlooked contribution is essential as it is known that evolution can fine tune protein dynamics through compensatory mutations, either to improve the proteins' performance or diversify its function while maintaining the same structural scaffold. To model this critical contribution, we introduce DyNoPy, a computational method that combines residue coevolution analysis with molecular dynamics (MD) simulations, revealing hidden correlations between functional sites. DyNoPy constructs a graph model of residue-residue interactions, identifies communities of key residue groups and annotates critical sites based on their roles. By leveraging the concept of coevolved dynamical couplings—residue pairs with critical dynamical interactions that have been preserved during evolution—DyNoPy offers a powerful method for predicting and analysing protein evolution and dynamics. We demonstrate the effectiveness of DyNoPy on SHV-1 and PDC-3, chromosomally encoded β-lactamases linked to antibiotic resistance, highlighting its potential to inform drug design and address pressing healthcare challenges.

Introduction

Quantifying the contribution of individual residues or residue groups to protein function is important to estimate the pathogenic effect of mutations (1). Identifying the functional roles of individual residues has primarily been done through mutagenesis experiments (2). Bioinformatics methods have complemented these approaches through analysis of multiple sequence alignments (MSA) of homologous proteins and structural data (3-8). Among these methods, computational techniques that can decode inter-residue evolutionary relationships from MSAs have paved the way for machine learning (ML) based strategies that can predict protein structure (9-12), stability (13), and function (7) and extend the scope of computational protein design (14-16). A most recent approach has combined experimental data from three proteins, NUDT15, PTEN and CYP2C9, on stability and function with sequence and structural features to train a ML model to predict functional sites (17).

Functional sites are often regulated by both, local and global interactions. Changes in these interactions are instrumental for functional events like substrate binding, catalysis, and conformational changes (18). The development of physical models of protein dynamics and the increase in available computational power has stimulated the adoption of computational techniques (19, 20) to investigate the conformational dynamics of proteins, an essential component of the many biological functions (21, 22). Different models have been proposed to describe the interactions between residues during simulations and network models have been particularly popular, including methods on single structures and MD simulations data built by analysing the response to external forces on residue networks (23), by estimating the prevalence of non-covalent energy interaction networks in homologous proteins (24), or by analysing linear or non-linear correlation in atomic fluctuations (25, 26). These techniques have demonstrated their usefulness in extracting allosteric networks from structural data with applications in enzyme design (26).

However, none of these techniques incorporate information on residue evolution into the computational approach, While it has been established that evolution through compensatory mutations in dynamic regions, like hinges and loops, can fine tune protein structural dynamics and introduce promiscuity, thereby diversifying biological function. Assuming that protein functional

dynamics is conserved during evolution, significant information on dynamic regions and substrate recognition sites should be recoverable using inter residue coevolution scores extracted from MSAs (27, 28). Coevolution analysis and Molecular Dynamics (MD) simulations have independently (29) and synergistically been combined in the past to identify important residues for function (30-34). Yet a method that combines hidden information on dynamics from evolution with direct information on local and global dynamics from conformational ensembles from MD is not yet available.

Here, we present DyNoPy, a computational method that can extract hidden information on functional sites from the combination of pairwise residue coevolution data and powerful descriptors of dynamics extracted from the analysis of MD ensembles. The method can detect coevolved dynamic couplings, i.e. residue pairs with critical dynamical interactions that have been preserved during evolution. These pairs are extracted from a graph model of residue-residue interactions. Communities of important residue groups are detected, and critical sites are identified by their eigenvector centrality in the graph (Fig. 1). We demonstrate the power of this approach on SHV-1 and PDC-3 β-lactamases of major clinical importance (35, 36). DyNoPy successfully detects residue couplings that align with previous studies, guide in the explanations of mutation sites with previously unexplained mechanisms and provide predictions on plausible important sites for the emergence of clinically relevant variants.

2　　**Fig. 1** – Overview of the DyNoPy workflow

Results and Discussion

β-lactamases are a group of enzymes capable of hydrolysing β-lactams, conferring resistance to β-lactam antibiotics (37). These enzymes are evolving rapidly, as single amino acid substitutions are sufficient to drive their evolution and increase their catalytic spectrum and inhibitor resistance profile (38). The widespread dissemination of β-lactamases across different bacterial species and their extensive emergence highlight their global impact on antibiotic resistance (39). The rapid evolution of β-lactamases and their clinical significance (38) makes them an ideal target for evaluating the robustness of DyNoPy.

In this study, we applied DyNoPy to two model enzymes from different β-lactamase families: class A β-lactamase SHV-1 (a chromosomally encoded enzyme in *Klebsiella pneumoniae*) and class C β-lactamase PDC-3 (a chromosomally encoded enzyme in *Pseudomonas aeruginosa*) (35, 36) (Supplementary Fig. S1 and S2). Both class A and class C β-lactamases comprise an α/β domain and an α helical domain, with the active site situated in between (40, 41). Moreover, both enzymes target the carbonyl carbon of the β-lactams using a highly conserved serine residue (42, 43). Despite these similarities, the structures of class A and class C β-lactamases are remarkably different (Fig. 2).

SHV-1 is a very well characterised enzyme with wealth of information on mutations and their corresponding effects on protein function. In contrast, the information available on PDC-3 remains limited. Detailed structural information on these enzymes can be found in the supplementary materials. Essential catalytic residues in SHV-1 are: S_{70}, K_{73}, S_{130}, E_{166}, N_{170}, K_{234}, G_{236}, and A_{237} (44) and conserved catalytic residues in PDC-3 include S_{64}, K_{67}, Y_{150}, N_{152}, K_{315}, T_{316}, and G_{317}. Highly conserved stretches of 3-9 hydrophobic residues, annotated as hydrophobic nodes, exists in class A β-lactamases and have been proven to be essential for protein stability (45). Residues defined as belonging to hydrophobic nodes within SHV-1 are listed in Supplementary Table S1.

In SHV-1, the predominant extended spectrum β-lactamase (ESBL) substitutions occur at L_{35}, G_{238}, and E_{240}, while R_{43}, E_{64}, D_{104}, A_{146}, G_{156}, D_{179}, R_{202}, and R_{205} appear in ESBLs with lower frequency (46). Mutations at M_{69}, S_{130}, A_{187}, T_{235}, and R_{244} are known to induce inhibitor resistance in the enzyme (47). In PDC-3, substitutions primarily occur on the Ω-loop, enhancing its flexibility to accommodate the bulky side chains of antibiotics, while deletions are more

common in the R2-loop (43). The predominant Ω-loop mutations isolated from clinics are found at

positions V_{211}, G_{214}, E_{219}, and Y_{221} (48).

Fig. 2 – Structural Comparison of SHV-1 (PDB ID: 3N4I) and PDC-3 (PDB ID: 4HEF) β-Lactamases. Catalytic serine S_{70} (SHV-1) and S_{64} (PDC3) are highlighted using stick representation. Important loops surrounding the active site are highlighted in red. In SHV-1, highlighted loops are the α3-α4 loop (residues 101-111), the Ω-loop (residues 164-179), and the hinge region (residues 213-218). In PDC-3, highlighted loops are the Ω-loop (residues 183-226) and the R2-loop (residues 280-310).

Emergence of highly conserved dynamic couplings

DyNoPy builds a pairwise model of conserved dynamic couplings detected by combining

coevolution scores and information on functional motions into a score J_{ij} (see Methods and Fig. 1).

To this end a dynamics descriptor should be selected. When the descriptor is associated with

functional conformational changes, it is expected that functionally relevant couplings will report

higher scores. Dynamics descriptors can be selected from commonly used geometrical collective

variables (CVs) for the analysis of MD trajectories (see Methods). As expected, the average ***J***

matrix score varies across the different CVs, with some of them showing no signal of dynamic

coupling (Supplementary Fig. S4C).

SHV-1 and PDC-3 exhibit distinct dynamics, requiring a different choice of the CV that best

captures the functional dynamics. For SHV-1, the global first principal component (PC1) proved to be the most effective feature, identifying 571 residue pairs with a J_{ij} value greater than 0. Conversely, PDC-3 requires selection of more localized features that can extract the Ω-loop dynamics from the overall protein motion. Among the dynamic descriptors, the partial first time-lagged component (TC1_partial) performed best for PDC-3, detecting 216 residue pairs with a J_{ij} value greater than 0. Consequently, PC1 and TC1_partial were selected to build the J matrix for SHV-1 and PDC-3, respectively. The performance of all 12 CVs for each protein was assessed and listed in the Supplementary Table S2.

The importance of dynamical information is evident when coevolution couplings (γ_{ij}) and conserved dynamic couplings (J_{ij}) are compared: the number of non-zero couplings decrease from 40% to <2% of total residue pairs in the protein (Supplementary Fig. S4D) when information from the dynamics descriptor is added. Thus, the inclusion of protein dynamics in coevolution studies acts as an effective filter that rules out residue pairs that do not have significant correlations with functional motions. Moreover, when relying only on γ_{ij}, all the residues in SHV-1 and PDC-3 are included within four identified communities (Supplementary Table S3), suggesting that coevolution scores (γ_{ij}) alone do not effectively discriminate residues relevant for protein functions. Furthermore, it would be hard to distinguish critical core residues for each community using only γ_{ij}, as the eigenvector centrality (EVC) values for the residues do not show remarkable differences (Supplementary Fig. S5A and S5B). This means that detailed dynamic investigation of the top residues is needed to determine which pairs should be picked up and further analysed. On the other hand, it is much easier to identify essential residues based on J scores calculated, as clear outliers with significantly higher EVC values could be seen for almost all communities (Supplementary Fig. S5C and S5D) (29, 49). In conclusion, the lack of specificity in the statistically based coevolution analysis supports the choice of incorporating a score for the correlation between residue interactions and dynamic behaviours that enables deconvolution of community information.

DyNoPy reveals critical residues and predicts evolutionary pathways in SHV-1

DyNoPy identified eight meaningful communities, each consisting of at least three strongly coupled residues within SHV-1 (Supplementary Fig. S4A). All crucial catalytic residues and

critical substitution sites previously mentioned participating in one of these communities with the exceptions of R_{43}, R_{202}, and S_{130}. Residues previously known to have critical role in function or conferring ESBLs/IRBLs phenotype are either directly coupled to protein dynamics or act as a central hub. The hubs interact with residues with either a role in catalysis or structural stability through their membership of hydrophobic nodes (35). Furthermore, DyNoPy identified key positions (L_{162} and N_{136}) within some communities that are known to undergo substitutions, conferring an ESBL phenotype in other class A β-lactamases. These substitutions have not yet emerged in the SHV family, providing insightful predictions about the potential future evolution of the enzyme. Detailed description of communities with secondary importance for protein function (community 3, 8, and 9) is provided in the supplementary information (Supplementary Fig. S6).

DyNoPy predicts mutation hotspots in SHV-1

DyNoPy detects critical mutation sites (L_{162} and N_{136}) that are known to extend the range of substrates in other class A β-lactamases but have not yet emerged as variants in the SHV family. These sites have not been modified in SHV family because of their plausible central role within the communities as they are mediating couplings with key functional residues essential for catalytic activity and structural stability, indicating their critical role in protein function and the potential lower mutation rate. These findings provide insightful predictions about the potential future evolution of the enzyme, as well as plausible explanations for why these mutations have not yet appeared.

$L_{162,}$ positioned at the start of the Ω-loop and adjacent to the crucial catalytic residue $E_{166,}$ is assigned as the core residue for community 1 (Fig. 3A). While it remains conserved in SHV family, variants of L_{162} have been isolated in other class A β-lactamase and are known to expand the enzyme catalytic spectrum. Single amino acid substitution at L_{162} can intensify antibiotic resistance in BEL-1 (50), a class A ESBL clinical variant, exhibiting robust resistance to ticarcillin and ceftazidime (51). BEL-2 diverges from BEL-1 by single amino acid substitution ($L_{162}F$) which alters the kinetic properties of the enzyme significantly and increases its affinity towards expanded-spectrum cephalosporins (52). The relationship between L_{162} and protein catalytic

functions can be explained using DyNoPy model, as there are couplings with catalytic important residues M_{69}, K_{73}, E_{166}, and K_{234}. Moreover, the BEL case has confirmed that $L_{162}F$ mutation significantly destabilizes the overall protein structure, highlighting the crucial role of L_{162} in maintaining protein stability (50). DyNoPy accurately identifies the centrality of L_{162} by reporting its connections with 28 backbone residues, including nine hydrophobic node residues critical for protein stability. Among these, five hydrophobic residues are part of the α2 node: V_{75}, L_{76}, G_{78}, V_{80}, and L_{81}, highlighting the contribution of L_{162} to the stability of the α2 helix (35).

Just like L_{162}, N_{136} undergoes advantageous mutations in other class A β-lactamases while remains highly conserved within the SHV family. It is the core residue for community 7 (Fig. 4B). This residue forms a hydrogen bond with E_{166}, stabilizing the Ω-loop (53). Although DyNoPy did not detect this direct interaction between N_{136} and E_{166}, the established relationship between N_{136} and N_{170} highlights the role of N_{136} in influencing E_{166}. N_{170}, an essential catalytic residue located on the Ω-loop, contributes to priming the water molecule for the deacylation step with E_{166} (54) and is directly coupled with N_{136}. Due to the essential contribution of N_{136} in facilitating E_{166} to maintain its proper orientation, it was previously thought to be intolerant to mutations as substitution of Asparagine to Alanine at this position would make the enzyme lose its function completely (55). However, $N_{136}D$ substitution has emerged as a new clinical variant very recently in PenL, a class A β-lactamase, by increasing its ability in hydrolysing ceftazidime (55), suggesting that this site has potential to mutate. This gain of function is mainly triggered by the increased flexibility of the Ω-loop (55). DyNoPy correctly detect a dynamical relationship between N_{136} and the Ω-loop (residues 164-179). Six residues present in the Ω-loop participate within this community, including R_{164} and D_{179}. These two residues are critical as they are forming the 'bottleneck' of the Ω-loop which is essential for the correct position of E_{166} (56). D_{179} is also a critical mutation site for SHV-1. Single amino acid substitutions like $D_{179}A$, $D_{179}N$, and $D_{179}G$ are enough for the extended spectrum phenotype (46).

DyNoPy detects residue couplings essential for protein stability

DyNoPy identifies residue couplings critical for protein functional motions, particularly associated with protein stability. These residue pairs exhibit strong relationships as they are not

only directly coupled with each other, but also forms various indirect couplings via other residues.

As a result, both residues are considered as core residues inside these communities. It is expected

that disruption of these couplings through mutation could compromise collective motions essential

for enzyme activity.

Fig. 3 – Community 1, 4, and 5 of SHV-1 β-Lactamase. All the residues are depicted as spheres on

the protein structure. The core residue for each community is highlighted in red, while purple is

used to emphasize the secondary core residue. Residues that interact with both cores are coloured in light yellow. Functional important residues are marked in cyan. Hydrophobic nodes are enclosed with cyan boxes. A. Community 1 of SHV-1, comprising 33 residues with L_{162} being the primary core residue. B. Community 4 of SHV-1, containing 12 residues and is centred by G_{156}. G_{156} and A_{146} are two functional important residues distant from the active site. G_{156} is 21.3□ away from the catalytic S_{70}. A_{146} is 16.8□ away from S_{70}. C. Community 5 of SHV-1, embracing 48 residues and showing a strong correlation between V_{103} and S_{106}.

As the secondary core residues in community 1 (Fig. 3A) F_{72} is showing a strong coupling with the primary core residue L_{162} and also forms nine indirect couplings with L_{162}, including via the catalytic K_{234}. This network of direct and indirect relationships reveals the importance of F_{72} and L_{162} coupling in maintaining protein functional motions. Interestingly, previous studies identified a small hydrophobic cavity formed by L_{162} and F_{72}, together with L_{139}, and L_{148}, which is essential for the stability of the active site (50). Notably, DyNoPy successfully recovers the key residues of this local hydrophobic cavity (L_{162}, F_{72}, and L_{148}).

The strong interplay between V_{103} and S_{106}, which are both residues on the α3-α4 loop, is seen in community 5 (Fig. 3C). These residues not only interact with each other directly but are also indirectly coupled via 22 other residues. This community emphasizes the significance of hydrophobic nodes in SHV stability and dynamics. Within the analysed 48 residues, 27 are hydrophobic, out of which 15 residues act as nodes critical for enzyme stabilization. Hydrophobic nodes stabilize their own secondary structures and interconnect to stabilize the overall protein (57). V_{103} and S_{106} themselves are hydrophobic nodes, stabilizing α3 helix and α4 helix respectively, and are strongly coupled with each other. In CTX-M, another class A enzyme, $N_{106}S$ is a common substitution that results in improved thermodynamic stability and compensate for the loss in stability of the variants (58). Interestingly, this residue is already a Serine in SHV, but still implies its pivotal role in protein stability.

DyNoPy provides valid explanations for mutation sites

During the evolution of β-lactamases, single mutations on specific sites that are distant from the functional sites have been observed to significantly alter protein catalytic functions. Additionally, single mutations on some surface exposed residues can dramatically increase protein stability. Understanding how these distant mutations impact function and stability becomes a

major challenge in understanding protein evolutionary pathways. Communities extracted by DyNoPy show these residues linked with functional important residues, providing a rational for these mutation sites with unknown functions.

Fig. 4 – Community 6 and 7 of SHV-1 β-Lactamase. All the residues are depicted as spheres on the protein structure. The core residue for each community is highlighted in red, while purple is used to emphasize the secondary core residue. Residues that interact with both cores are coloured in light yellow. Functional important residues are marked in cyan. A. Community 6 of SHV-1, comprising 30 residues with Y_{105} being the primary core residue. R_{205} is a functional important residue that is 20.6 away from the active site S_{70}. B. Community 7 of SHV-1, containing 34 residues and is centred by N_{136}.

Mutations of G_{156} are limited but they lead to ESBL phenotype in the SHV family (46). G_{156} is the central residue for community 4 (Fig. 3B), but it is distant from the active site, over 20Å away from the catalytic serine S_{70}. Clinical variant SHV-27, has extended resistance ability towards cefotaxime, ceftazidime, and aztreonam (59). It differs from SHV-1 by single amino acid substitution $G_{156}D$, suggesting that it has directly evolved from SHV-1 (59). Limited research has been done on position G_{156}, and the understanding of how it affects the enzyme catalytic properties given that it is far away from the active site is still unclear. Based on our results, we suggest that this residue is essential for the overall protein function because of its 11 coevolved dynamic couplings with protein dynamics, including A_{146}, another ESBL substitution site.

SHV-38, another ESBL that is capable of hydrolysing carbapenems, harbours a single $A_{146}V$ substitution compared to SHV-1 (60). Like G_{156}, A_{146} is 16.8 Å away from S_{70} but shows an ability in altering protein catalytic function. The A_{146}-G_{156} residue pair shows a strong coevolutionary signal and strong correlation with protein overall dynamics, implying that there may compensatory mutations at these sites with potential to emerge in the SHV family in the future. These two residues are not connected to any catalytic residues but their coupling to functional dynamics can offer plausible explanation to ESBL activity of these two mutations.

Unlike other substitution sites that are adjacent to the active site, R_{205} is situated more than 20 Å away from catalytic serine S_{70}. Its side chain points outward from the protein, exposing to the solvent. The $R_{205}L$ substitution often co-occurs with other ESBL mutations and is thought to indirectly contribute to the ESBL phenotype by compensating for stability loss induced by other mutations (61). SHV-3 is an ESBL that exhibits significant resistance to cefotaxime and ceftriaxone (62). Two substitutions in this enzyme, $R_{205}L$ and $G_{238}S$, extend its resistance profile (62). Thus, it is promising to see that DyNoPy detected these two mutation sites together within community 6 (Fig. 4A).

Y_{105} and R_{266} are the core residues for community 6. Y_{105} is situated on the α3-α4 loop positioned at the left side of the binding pocket. It is an important catalytic residue that recognizes and binds to the thiazolidine ring of penicillins or β-lactamase inhibitors (63). There is very limited information on the role of R_{266}, except that it may stabilize the Ω-loop in the SHV family similar to the analogous T_{266} in TEM (64). G_{238} is coupled with an essential catalytic residue Y_{105},

which further links with other catalytic functional residues: S_{70} and A_{237}, and R_{266}, a residue that

known to stabilize the Ω-loop. This indicates that mutations on G_{238} would result in an alteration

on protein catalytic function, as well as an increased flexibility of the protein, which strongly

aligns with previous finding (62). Its linked mutation site R_{205} does not showing direct coupling

with any catalytic residues. Instead, it is directly coupled with $R_{266,}$ which we mentioned as an

Ω-loop stabilizer. Thus, it is not surprising that R_{205} substitution alone is never observed in nature

(65), as it would not give significant evolutionary advantage to the protein.

Insights into unexplained functional sites of PDC-3

Unlike the extensively studied SHV-1, the functional roles of individual amino acids in

PDC-3 remains largely unexplored. This gap in understanding serves as welcome challenge for

interpreting the effects of mutations and the dynamic behaviour of PDC-3 from our results.

Although several mutation hotspots, such as those on the Ω-loop (48), have been identified, very

little is known about the specific contributions of individual amino acids on the functionality of

PDC-3.

In PDC-3, mutations have primarily been reported in the Ω-loop. They enhance its flexibility

to accommodate the bulky side chains of antibiotics, while deletions are more common in the

R2-loop (43). DyNoPy detected five communities in total (Supplementary Fig. S4B) with all the

four predominant Ω-loop mutations appeared in these communities. Community 3, 4 and 5 are

explained in the Supplementary information (Fig. S7). Furthermore, DyNoPy also detected several

previously unexplored Ω-loop residues.

G_{214}, a known mutation site in PDC-3, is the core residue in community 1. Another two

essential mutation sites: E_{219} and Y_{221} also participate in this community, directly coupled with

G_{214} (Fig. 5A). G_{214} also has direct couplings with four other Ω-loop residues: A_{195}, A_{197}, G_{212}, and

L_{216}. Previous results have demonstrated that substitutions of Glycine to Alanine or Arginine at

214 significantly destabilizes the Ω-loop (36). The strong correlation between G_{214} and these

Ω-loop residues emphasizes the significant contribution of G_{214} towards the stability of the Ω-loop,

which corroborates with previous results (36). Moreover, substitutions such as $G_{214}A$ and $G_{214}R$

and mutations on E_{219} and Y_{221} do not affect R2 loop flexibility, resulting in the smaller active site

volume among variants (36) because none of the residues from the R2 loop are detected in this

community offering plausible explanation to previously unexplained phenomenon.

Fig. 5 - Community 1 and 2 of PDC-3 β-Lactamase. All the residues are depicted as spheres on
the protein structure. The core residue for each community is highlighted in red. Functional
important residues are marked in cyan. A. Community 1 of PDC-3, comprising 36 residues with
G_{214} being the primary core residue. B. Community 2 of PDC-3, containing 74 residues and is
centred by G_{204}.

G_{204} is the core residue of community 2, coupled with 73 other residues, most of which are

distant from the catalytic site, suggesting plausible crucial role in overall protein stability like L_{162}

in SHV-1 (Fig. 5B). G_{204}, a newly emerged mutation site in the PDC family (66), is located on the

short β-sheet β5a within the Ω-loop, near the hinge region between β8 and β9 just above the active

site. The only known variant of G_{204} is PDC-466, which was derived from PDC-462 ($A_{89}V$, $Q_{120}K$,

$V_{211}A$, $N_{320}S$), with an addition of $G_{204}D$ (66). Coupling of G_{204} to several catalytically important

residues, including K_{67}, K_{315}, and T_{316} can suggest that mutations at this site can negatively impact catalytic power. This offers a plausible explanation of seeing fewer variants at this site and mutations at this site could have impact on hydrolysing capabilities of PDC variants. This should be confirmed by further experimental studies of variants of G_{204}. Unlike G_{214}, E_{219} and Y_{221} mutations which do not influence the dynamics of the R2 loop, substitutions on V_{211}, a member of Ω-loop, has impact on dynamics of R2 loop because of its indirect couplings, through G_{204} to R2-loop residues (36). Two less critical substitution sites, H_{188} and V_{329}, were also observed in community 2.

Conclusions

DyNoPy offers two distinct advantages over existing computational tools (24, 26): a) information on residue-residue coevolution can be directly used to detect the components of protein dynamics that have been preserved during evolution b) dynamic descriptors extracted from the MD ensembles can be used to identify the function-specific conserved dynamic couplings. These couplings are then easily modelled as a graph and network analysis is used to extract epistatic communities and assign roles to residues based on their importance in the graph model. The choice of a relevant descriptor of functional dynamics has an impact on the ability to detect couplings that are involved in functional dynamics. In systems where there is limited role of dynamics in the function, the analysis done with DyNoPy is equivalent to conventional coevolution analysis, which can be consider one limitation of our method.

Here we demonstrated how the choice of relevant global and local descriptors returns a higher number of effective couplings (greater than 0), and in turn leads to interpretable graph models and communities. In other systems, when multiple descriptors can be used to quantify functional conformational change, it is expected that they will differently modulate the effect of coevolution coupling, which will be reflected in a different structure of the associated graph models. This suggests the use of DyNoPy to generate comparative models in proteins with multiple functions associated to distinct dynamical changes.

Mutations of L_{162} and N_{136} have not yet emerged in SHV-1, but they are detected by DyNoPy as core residues for communities. These residues are strongly coupled with other functional

important residues, which play critical roles in protein stability and catalytic activity. The identification of these couplings shows high consistency with previous studies and highlights the importance of L_{162} and N_{136} in SHV-1 functional dynamics. Given their central role in these communities, mutations in L_{162} and N_{136} can significantly alter protein function, suggesting their potential for future evolutionary changes. However, their strong relationships with these critical functional residues also suggest that mutation at these sites would need to be balanced to maintain protein function, providing an explanation for why such mutations have not yet emerged in SHV-1 (67). The ability of DyNoPy in detecting functionally important mutation sites was demonstrated via well-characterized mutation sites including R_{205} and G_{238} from SHV-1. Moreover, DyNoPy shows predictive ability on less-studied mutation sites such as G_{156} and $A_{146,}$ by detecting critical residue couplings that coevolved with functional motions.

Based on the knowledge we have gained from analysis of SHV-1 functional protein dynamics we suggest that in PDC-3, mutations at G_{204} because of its significant conserved dynamic couplings can lead to new ESBL/IRBL clinical variants. We suggest that DyNoPy can be used as a predictive tool to identify potential functional residues within this enzyme and guide future mutagenesis studies.

In summary, by integrating hidden evolutionary information with direct dynamic interactions, DyNoPy provides a powerful framework for identifying and analysing functional sites in proteins. The tool not only identifies key residues involved in local and global interactions, but also improves our ability to predict silent residues with previously unknown roles for future experimental testing. Our application of DyNoPy to broad-spectrum β-lactamases ESBLs and IRBLs demonstrates its potential to address key medical challenges such as antibiotic resistance by providing valid predictions on protein evolution.

1 Methodology

DyNoPy generates a graph representation of the protein structure that captures the couplings

between amino acid residues contributing to the functional dynamics of the protein. Residues are

represented as graph nodes, and conserved dynamic couplings are recorded as edges. Edge

weights quantify the strength of these couplings. The model is built on two assumptions: residue

pairs should have i) coevolved and their ii) time-dependent interactions correlate with a functional

conformational change.

Therefore, edge weights (J_{ij}) for residue i and j are calculated as:

$$J_{ij} = \alpha\gamma_{ij} + \beta\rho_{ij} \quad where \ \alpha + \beta = 1 \tag{1}$$

11 where γ_{ij} is the scaled coevolution score and ρ_{ij} is the degree of correlation with the selected

12 functional conformational change. α and β are weights assigned to γ_{ij} and ρ_{ij} that have a sum

13 of one. The relative weight of the scaled coevolution score (α) is set to 0.5 in this study. When

14 either of the assumptions listed above is not met, J_{ij} is set to zero.

Scaled coevolution scores

The occurrence of residue-residue coevolution can be estimated and quantified using probabilistic

models of correlated mutations from deep multiple sequence alignments (MSA). DyNoPy

supports generation of the MSA using the HH-Suite package (68) and calculation of scaled

coevolution score (γ_{ij}) using CCMpred (69) as per the protocol described in Bibik et al. (70). For

SHV-1 and PDC-3 hhblits returned 18,174 sequences (N$_{eff}$: 11.082) and 27,892 sequences (N$_{eff}$:

9.951). Sequences were detected from the UniRef30 (v2022_02) database (71). First a pairwise

residue coevolution matrix (C) is calculated, then these raw scores (c_{ij}) are divided by the matrix

mean (Equation 2). All scores (s_{ij}) smaller than 1 are set to zero, and the remaining values are

normalised by the maximum value (Equation 3):

$$s_{ij} = \frac{c_{ij}}{\langle C \rangle} \tag{2}$$

$$\gamma_{ij} = \begin{cases} 0, & s_{ij} < 1 \\ \dfrac{s_{ij}}{s_{max}}, & s_{ij} \geq 1 \end{cases} \tag{3}$$

Correlation with functional motions

The contribution of a residue pair to a selected functional motion is estimated by how much the change in interaction energy between the two residues over time is correlated with a collective variable (CV) describing the functional motion:

$$r_{ij} = cor(\varepsilon_{ij}(t), d(t)) \tag{4}$$

$$\rho_{ij} = \begin{cases} 0, & r_{ij} \leq 0.5 \\ r_{ij}, & r_{ij} > 0.5 \end{cases} \tag{5}$$

where $\varepsilon_{ij}(t)$ is the pairwise non-bonded interaction energy (see details in Supplementary Information) and $d(t)$ is the time-dependent value of the CV. Examples of CV and a discussion on the choice of the most relevant CV is presented in the results section. Correlation values smaller than 0.5 are set to 0. In absence of detectable contributions to the functional dynamics of the system, the couplings extracted by DyNoPy will describe a pure evolutionary model, and the community detection method presented below will be equivalent to a direct decomposition of the residue coevolution network into units.

Graph representation and analysis of conserved dynamic couplings

All pairwise conserved dynamic couplings (Equation 1) are collected into a square matrix J. A graph is built from J, using python-igraph v0.11 library (72). Nodes represent residues, and edges are drawn between nodes with positive J_{ij}. Edge weights are set to J_{ij}. The relative importance of the residues in this model of protein dynamics is calculated as eigenvector centrality of the nodes (73). Residues involved in extensive correlated dynamics with other highly connected residues have higher eigenvector centrality (EVC) scores. Groups of residues contributing to important collective motions are detected by community analysis of the graph structure. The

Girvan-Newman algorithm is used to extract the community structure (74). A meaningful community should contain at least three residues. Applying network analysis on the combined dynamics-coevolution matrix helps us extracting higher-order interactions beyond pairwise coupling and detecting critical residues, which show multiple interactions with each other. Moreover, indirect long-range relationships, which would be hard to identify from numerical data, could be detected through community clustering. Community-based analysis offers a more comprehensive understanding of residue relationships and enables the visualization of residue couplings on the protein structure.

Adaptive Sampling Molecular Dynamics Simulations

MD simulation data was sourced from our previous studies (35, 36). To summarise, SHV-1 structural coordinates (PDB ID: 3N4I) were obtained from the Protein Data Bank and modified to the wild type by introducing the E104D mutation. Similarly, the PDC-3 structure was derived from PDC-1 (PDB ID: 4HEF) by a T105A substitution. Both enzymes were protonated at pH 7.0 using PropKa from the PlayMolecule platform (75). One disulfide bond between C_{77} and C_{123} was specified in SHV-1. Both structures were solvated with TIP3P water molecules in a periodic box with a box size of 10 Å. Ions were added to neutralize the overall charge of each system at 150mM KCl. Amber force field ff14SB was used for all MD simulations (76). After an initial minimisation of 1000 steps, both the enzymes were equilibrated for 5 ns in the NPT ensemble at 1 atmospheric pressure using the Berendsen barostat (77). The initial velocities for each simulation were sampled from the Boltzmann distribution at 300 K. Multiple Markov State Model (MSM)-based adaptively sampled simulations were performed for both proteins based on the ACEMD engine (78, 79). A canonical (NVT) ensemble with a Langevin thermostat (80) (damping coefficient of 0.1 ps−1) and a hydrogen mass repartitioning scheme were employed to achieve time steps of 4 fs. For SHV-1, each trajectory spanned 60 ns with a time step of 0.1 ns, with a total of 593 trajectories. In the case of PDC-3, 100 trajectories were collected, each containing 3000 frames, lasting 300 ns. To manage the extensive datasets efficiently, trajectories were strategically stridden to ensure that a minimum of 30,000 frames were preserved for each system. The resulting trajectories are summarized in Supplementary Table S4.

Calculation and Selection of Collective Variables

DyNoPy works on the assumption that time-dependent interactions between critical residues,

either having significant structural change or not will correlate with functional conformational

motions. Since MD simulation data is high-dimensional, a time-dependent collected variable (CV)

is required to extract the most relevant information for the process under study. The usefulness of

DyNoPy is dependent on the choice of the CVs. To guide the selection of CVs, we selected 12

distinct features: radius of gyration (R_g), the first principal component (PC1), partial PC1

(PC1_partial), the first time-lagged independent component (TC1), partial TC1 (TC1_partial),

global root mean square deviation (gRMSD), partial RMSD (pRMSD), dynamical RMSD

(dRMSD), global solvent accessible surface area (gSASA), partial SASA (pSASA), active site

pocket volume, and the number of hydrogen bonds (hbond). A description of the CVs, including

the calculation methods and the residues used to calculate the partial variables, is detailed in the

Supplementary information. CVs were subsequently used as input features for DyNoPy. A good

collective variable (CV) should appropriately describe protein functional motions. Thus, a CV that

detects the highest number of residue couplings is expected to be the most suitable descriptor. The

length of the MD simulations should be appropriate to effectively sample the desired functional

process as described by the selected CV.

Data Availability

All files required to run the simulations (topology, coordinates, input), processed trajectories (xtc), corresponding coordinates (pdb), can be downloaded from the DOI https://doi.org/10.57760/sciencedb.15876 (PDC-3) and 10.5281/zenodo.13693144 (SHV-1). DyNoPy is available at https://github.com/alepandini/DyNoPy.

Acknowledgements

SCD was supported by Leverhulme Trust grant RPG-2017-222 awarded to AP and JAG. The authors would like to thank Arianna Fornili for insightful suggestions on the design of DyNoPy methodology.

Conflict of Interests

All authors declare no conflict of interests.

References

1. P. D. Stenson *et al.*, The Human Gene Mutation Database: towards a comprehensive repository of inherited mutation data for medical research, genetic diagnosis and next-generation sequencing studies. *Hum Genet* **136**, 665-677 (2017).

2. K. A. Matreyek *et al.*, Multiplex assessment of protein variant abundance by massively parallel sequencing. *Nat Genet* **50**, 874-882 (2018).

3. F. J. Poelwijk, V. Krishna, R. Ranganathan, The Context-Dependence of Mutations: A Linkage of Formalisms. *PLoS Comput Biol* **12**, e1004771 (2016).

4. M. H. Høie, M. Cagiada, A. H. Beck Frederiksen, A. Stein, K. Lindorff-Larsen, Predicting and interpreting large-scale mutagenesis data using analyses of protein stability and conservation. *Cell Rep* **38**, 110207 (2022).

5. L. M. Blaabjerg *et al.*, Rapid protein stability prediction using deep learning representations. *Elife* **12** (2023).

6. A. S. Dunham, P. Beltrao, Exploring amino acid functions in a deep mutational landscape. *Mol Syst Biol* **17**, e10305 (2021).

7. T. A. Hopf *et al.*, Mutation effects predicted from sequence co-variation. *Nat Biotechnol* **35**, 128-135 (2017).

8. P. Radivojac *et al.*, A large-scale evaluation of computational protein function prediction. *Nat Methods* **10**, 221-227 (2013).

9. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021).

10. Z. Lin *et al.*, Evolutionary-scale prediction of atomic-level protein structure with a

language model. *Science* **379**, 1123-1130 (2023).

11. M. Baek *et al.*, Accurate prediction of protein structures and interactions using a three-track neural network. *Science* **373**, 871-876 (2021).

12. D. S. Marks, T. A. Hopf, C. Sander, Protein structure prediction from sequence variation. *Nat Biotechnol* **30**, 1072-1080 (2012).

13. A. Broom, K. Trainor, Z. Jacobi, E. M. Meiering, Computational Modeling of Protein Stability: Quantitative Analysis Reveals Solutions to Pervasive Problems. *Structure* **28**, 717-726 e713 (2020).

14. D. Ding *et al.*, Protein design using structure-based residue preferences. *Nat Commun* **15**, 1639 (2024).

15. Z. Wu, S. B. J. Kan, R. D. Lewis, B. J. Wittmann, F. H. Arnold, Machine learning-assisted directed protein evolution with combinatorial libraries. *Proc Natl Acad Sci U S A* **116**, 8852-8858 (2019).

16. W. P. Russ *et al.*, An evolution-based model for designing chorismate mutase enzymes. *Science* **369**, 440-445 (2020).

17. M. Cagiada *et al.*, Discovering functionally important sites in proteins. *Nat Commun* **14**, 4175 (2023).

18. S. J. Wodak *et al.*, Allostery in Its Many Disguises: From Theory to Applications. *Structure* **27**, 566-578 (2019).

19. P. Campitelli, T. Modi, S. Kumar, S. B. Ozkan, The Role of Conformational Dynamics and Allostery in Modulating Protein Evolution. *Annu Rev Biophys* **49**, 267-288 (2020).

20. C. H. M. Rodrigues, D. E. V. Pires, D. B. Ascher, DynaMut2: Assessing changes in

stability and flexibility upon single and multiple point missense mutations. *Protein Sci* **30**, 60-69 (2021).

21. K. Henzler-Wildman, D. Kern, Dynamic personalities of proteins. *Nature* **450**, 964-972 (2007).

22. L. C. James, D. S. Tawfik, Conformational diversity and protein evolution--a 60-year-old hypothesis revisited. *Trends Biochem Sci* **28**, 361-368 (2003).

23. Z. Nevin Gerek, S. Kumar, S. Banu Ozkan, Structural dynamics flexibility informs function and evolution at a proteome scale. *Evol Appl* **6**, 423-433 (2013).

24. D. Yehorova, R. M. Crean, P. M. Kasson, S. C. L. Kamerlin, Key interaction networks: Identifying evolutionarily conserved non-covalent interaction networks across protein families. *Protein Sci* **33**, e4911 (2024).

25. O. F. Lange, H. Grubmuller, Generalized correlation for biomolecular dynamics. *Proteins* **62**, 1053-1061 (2006).

26. S. Osuna, The challenge of predicting distal active site mutations in computational enzyme design. *WIREs Computational Molecular Science* **11** (2020).

27. D. Granata, L. Ponzoni, C. Micheletti, V. Carnevale, Patterns of coevolving amino acids unveil structural and dynamical domains. *Proc Natl Acad Sci U S A* **114**, E10612-E10621 (2017).

28. Y. Liu, I. Bahar, Sequence evolution correlates with structural dynamics. *Mol Biol Evol* **29**, 2253-2263 (2012).

29. D. J. Parente, J. C. Ray, L. Swint-Kruse, Amino acid positions subject to multiple coevolutionary constraints can be robustly identified by their eigenvector network

centrality scores. *Proteins* **83**, 2293-2306 (2015).

30. L. Ponzoni, G. Polles, V. Carnevale, C. Micheletti, SPECTRUS: A Dimensionality Reduction Approach for Identifying Dynamical Domains in Protein Complexes from Limited Structural Datasets. *Structure* **23**, 1516-1525 (2015).

31. L. Sutto, S. Marsili, A. Valencia, F. L. Gervasio, From residue coevolution to protein conformational ensembles and functional dynamics. *Proc Natl Acad Sci U S A* **112**, 13567-13572 (2015).

32. R. A. Estabrook *et al.*, Statistical coevolution analysis and molecular dynamics: identification of amino acid pairs essential for catalysis. *Proc Natl Acad Sci U S A* **102**, 994-999 (2005).

33. Z. Chen, S. Rappert, J. Sun, A. P. Zeng, Integrating molecular dynamics and co-evolutionary analysis for reliable target prediction and deregulation of the allosteric inhibition of aspartokinase for amino acid production. *J Biotechnol* **154**, 248-254 (2011).

34. J. Wang, Y. Zhao, Y. Wang, J. Huang, Molecular dynamics simulations and statistical coupling analysis reveal functional coevolution network of oncogenic mutations in the CDKN2A-CDK6 complex. *FEBS Lett* **587**, 136-141 (2013).

35. E. Olehnovics *et al.*, The Role of Hydrophobic Nodes in the Dynamics of Class A beta-Lactamases. *Front Microbiol* **12**, 720991 (2021).

36. S. Chen *et al.*, Omega-Loop mutations control the dynamics of the active site by modulating a network of hydrogen bonds in PDC-3 beta-lactamase. *bioRxiv* 10.1101/2024.02.04.578824 (2024).

1   37.   K. Poole, Resistance to beta-lactam antibiotics. *Cell Mol Life Sci* **61**, 2200-2223

2   (2004).

3   38.   K. Bush, Past and Present Perspectives on beta-Lactamases. *Antimicrob Agents*

4   *Chemother* **62** (2018).

5   39.   K. Bush, Proliferation and significance of clinically relevant beta-lactamases. *Ann N Y*

6   *Acad Sci* **1277**, 84-90 (2013).

7   40.   A. Matagne, J. Lamotte-Brasseur, J. M. Frere, Catalytic properties of class A

8   beta-lactamases: efficiency and diversity. *Biochem J* **330 (Pt 2)**, 581-598 (1998).

41.   A. Philippon, G. Arlet, R. Labia, B. I. Iorga, Class C beta-Lactamases: Molecular

Characteristics. *Clin Microbiol Rev* **35**, e0015021 (2022).

42.   T. Palzkill, Structural and Mechanistic Basis for Extended-Spectrum Drug-Resistance

Mutations in Altering the Specificity of TEM, CTX-M, and KPC beta-lactamases. *Front*

Mol Biosci **5**, 16 (2018).

43.   G. A. Jacoby, AmpC beta-lactamases. *Clin Microbiol Rev* **22**, 161-182, Table of

Contents (2009).

44.   R. P. Ambler *et al.*, A standard numbering scheme for the class A beta-lactamases.

Biochem J **276 (Pt 1)**, 269-270 (1991).

45.   I. Galdadas *et al.*, Defining the architecture of KPC-2 Carbapenemase: identifying

allosteric networks to fight antibiotics resistance. *Sci Rep* **8**, 12916 (2018).

46.   A. Liakopoulos, D. Mevius, D. Ceccarelli, A Review of SHV Extended-Spectrum

beta-Lactamases: Neglected Yet Ubiquitous. *Front Microbiol* **7**, 1374 (2016).

47.   D. Pagan-Rodriguez *et al.*, Tazobactam inactivation of SHV-1 and the

inhibitor-resistant Ser130 -->Gly SHV-1 beta-lactamase: insights into the mechanism of inhibition. *J Biol Chem* **279**, 19494-19501 (2004).

48. M. D. Barnes *et al.*, Deciphering the Evolution of Cephalosporin Resistance to Ceftolozane-Tazobactam in Pseudomonas aeruginosa. *mBio* **9** (2018).

49. C. F. A. Negre *et al.*, Eigenvector centrality for characterization of protein allosteric pathways. *Proc Natl Acad Sci U S A* **115**, E12201-E12208 (2018).

50. C. Pozzi *et al.*, Crystal Structure of the Pseudomonas aeruginosa BEL-1 Extended-Spectrum beta-Lactamase and Its Complexes with Moxalactam and Imipenem. *Antimicrob Agents Chemother* **60**, 7189-7199 (2016).

51. P. Bogaerts, C. Bauraing, A. Deplano, Y. Glupczynski, Emergence and dissemination of BEL-1-producing Pseudomonas aeruginosa isolates in Belgium. *Antimicrob Agents Chemother* **51**, 1584-1585 (2007).

52. L. Poirel *et al.*, BEL-2, an extended-spectrum beta-lactamase with increased activity toward expanded-spectrum cephalosporins in Pseudomonas aeruginosa. *Antimicrob Agents Chemother* **54**, 533-535 (2010).

53. F. Bo☐s, J. Pleiss, Conserved water molecules stabilize the Omega-loop in class A beta-lactamases. *Antimicrob Agents Chemother* **52**, 1072-1079 (2008).

54. V. Agarwal, T. C. Yadav, A. Tiwari, P. Varadwaj, Detailed investigation of catalytically important residues of class A beta-lactamase. *J Biomol Struct Dyn* **41**, 2046-2073 (2023).

55. T. P. Cao *et al.*, Non-catalytic-Region Mutations Conferring Transition of Class A beta-Lactamases Into ESBLs. *Front Mol Biosci* **7**, 598998 (2020).

56. D. Parwana *et al.*, The Structural Role of N170 in Substrate-Assisted Deacylation in

KPC-2 beta-Lactamase. *Angew Chem Int Ed Engl* **63**, e202317315 (2024).

57. I. Galdadas *et al.*, Allosteric communication in class A beta-lactamases occurs via

cooperative coupling of loop dynamics. *Elife* **10** (2021).

58. S. Lu *et al.*, An active site loop toggles between conformations to control antibiotic

hydrolysis and inhibition potency for CTX-M beta-lactamase drug-resistance enzymes.

Nat Commun **13**, 6726 (2022).

59. J. E. Corkill, L. E. Cuevas, R. Q. Gurgel, J. Greensill, C. A. Hart, SHV-27, a novel

cefotaxime-hydrolysing beta-lactamase, identified in Klebsiella pneumoniae isolates

from a Brazilian hospital. *J Antimicrob Chemother* **47**, 463-465 (2001).

60. L. Poirel *et al.*, Emergence in Klebsiella pneumoniae of a chromosome-encoded SHV

beta-lactamase that compromises the efficacy of imipenem. *Antimicrob Agents*

Chemother **47**, 755-758 (2003).

61. N. Ben Achour, P. S. Mercuri, M. Ben Moussa, M. Galleni, O. Belhadj,

Characterization of a novel extended-spectrum TEM-type beta-lactamase, TEM-164,

in a clinical strain of Klebsiella pneumoniae in Tunisia. *Microb Drug Resist* **15**,

195-199 (2009).

62. M. H. Nicolas, V. Jarlier, N. Honore, A. Philippon, S. T. Cole, Molecular

characterization of the gene encoding SHV-3 beta-lactamase responsible for

transferable cefotaxime resistance in clinical isolates of Klebsiella pneumoniae.

Antimicrob Agents Chemother **33**, 2096-2100 (1989).

63. C. R. Bethel *et al.*, Role of Asp104 in the SHV beta-lactamase. *Antimicrob Agents*

Chemother **50**, 4124-4131 (2006).

64. A. P. Kuzin *et al.*, Structure of the SHV-1 beta-lactamase. *Biochemistry* **38**, 5720-5727 (1999).

65. S. Neubauer *et al.*, A Genotype-Phenotype Correlation Study of SHV beta-Lactamases Offers New Insight into SHV Resistance Profiles. *Antimicrob Agents Chemother* **64** (2020).

66. C. A. Colque *et al.*, Development of antibiotic resistance reveals diverse evolutionary pathways to face the complex and dynamic environment of a long-term treated patient. *bioRxiv* 10.1101/2021.05.14.444257 (2021).

67. M. Soskine, D. S. Tawfik, Mutational effects and the evolution of new protein functions. *Nat Rev Genet* **11**, 572-582 (2010).

68. M. Remmert, A. Biegert, A. Hauser, J. Soding, HHblits: lightning-fast iterative protein sequence searching by HMM-HMM alignment. *Nat Methods* **9**, 173-175 (2011).

69. S. Seemayer, M. Gruber, J. Soding, CCMpred--fast and precise prediction of protein residue-residue contacts from correlated mutations. *Bioinformatics* **30**, 3128-3130 (2014).

70. P. Bibik, S. Alibai, A. Pandini, S. C. Dantu, PyCoM: a python library for large-scale analysis of residue-residue coevolution data. *Bioinformatics* **40** (2024).

71. M. Mirdita *et al.*, Uniclust databases of clustered and deeply annotated protein sequences and alignments. *Nucleic Acids Res* **45**, D170-D176 (2017).

72. G. Csárdi, T. Nepusz (2006) The igraph software package for complex network research.

73. M. E. J. Newman, Detecting community structure in networks. *The European Physical*

Journal B - Condensed Matter **38**, 321-330 (2004).

74. M. E. Newman, Finding community structure in networks using the eigenvectors of

matrices. *Phys Rev E Stat Nonlin Soft Matter Phys* **74**, 036104 (2006).

75. G. Martinez-Rosell, T. Giorgino, G. De Fabritiis, PlayMolecule ProteinPrepare: A Web

Application for Protein Preparation for Molecular Dynamics Simulations. *J Chem Inf*

Model **57**, 1511-1516 (2017).

76. J. A. Maier *et al.*, ff14SB: Improving the Accuracy of Protein Side Chain and Backbone

Parameters from ff99SB. *J Chem Theory Comput* **11**, 3696-3713 (2015).

77. H. J. C. Berendsen, J. P. M. Postma, W. F. van Gunsteren, A. DiNola, J. R. Haak,

Molecular dynamics with coupling to an external bath. *The Journal of Chemical*

Physics **81**, 3684-3690 (1984).

78. S. Doerr, M. J. Harvey, F. Noe, G. De Fabritiis, HTMD: High-Throughput Molecular

Dynamics for Molecular Discovery. *J Chem Theory Comput* **12**, 1845-1852 (2016).

79. M. J. Harvey, G. Giupponi, G. D. Fabritiis, ACEMD: Accelerating Biomolecular

Dynamics in the Microsecond Time Scale. *J Chem Theory Comput* **5**, 1632-1639

(2009).

80. R. L. Davidchack, R. Handel, M. V. Tretyakov, Langevin thermostat for rigid body

dynamics. *J Chem Phys* **130**, 234101 (2009).