Characterisation of the historic demographic decline of the British European
polecat population

R. Shaw [1,2], J. MacPherson [3], A, C. Kitchener[4], G, J. Etherington[1], W. Haerty [1,2]

[1] Earlham Institute, Norwich Research Park, Colney Lane, Norwich, NR4 7UZ
rebecca.shaw@earlham.ac.uk

[2] University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ

[3] Vincent Wildlife Trust, Ledbury, Herefordshire, HR8 1EP

4 Department of Natural Sciences, National Museums Scotland, Edinburgh EH1 1JF,
UKUK and School of Geosciences, University of Edinburgh, Drummond Street,
Edinburgh EH8 9XP, UK

Keywords: population genomics, demographic history, range expansion

Abstract

The European polecat (*Mustela putorius*) has a widespread distribution across many
countries of mainland Europe but is documented to be declining within these ranges. In
Britain, direct persecution led to a severe decline of the polecat population during the 19th
century. Unlike European mainland populations, it is now recovering across much of its
former British range. The genomic and conservation implications of such a severe
demographic decline, followed by the current recovery, have still to be characterised in the
European polecat in Britain. Here we carry out population-level whole-genome analyses of
65 polecats from Britain (Wales and England) and the European mainland. Our analyses
reveal that Welsh polecats show genetic variability from both English and European
polecats, while British polecats as a whole exhibit signs of genetic isolation from mainland
European populations. We also reconstructed the demographic history of the Welsh polecat
to quantify the magnitude of the bottleneck. Our analyses confirmed the drastic decline of
the Welsh polecat's effective population size, with a severe genetic bottleneck around 30-40
generations ago (1854-1894). We investigated whether whole genome diversity reflected
this demographic event and found that Welsh polecats had significantly less genetic diversity
than English polecats, but not European polecats. Runs of homozygosity and genetic load
present in Welsh and English polecat genomes also indicated recent historic inbreeding. Our
findings suggest that the increase in the British polecat population size may be attributed to
admixture events. Additionally, we demonstrate that the Welsh polecat constitutes a
genetically distinct population, which could be crucial for the overall conservation of
European polecats by preserving unique genetic diversity.

1.0 Introduction

Human-mediated actions are considered some of the most serious threats to the survival of
small carnivore populations (Willcox 2020; C. Marneweck et al. 2021; C. J. Marneweck et al.

2022; Prugh et al. 2023). This includes active persecution and habitat fragmentation, which
reduces connectivity between individuals, thereby limiting gene flow and further contributing
to reducing effective population size (N_e) (Croose et al. 2018; C. Marneweck et al. 2021).
The ensuing decline in genetic diversity further decreases future resilience to arising
challenges including environmental changes (Frankham et al. 2017). Smaller populations
are therefore highly susceptible to inbreeding and genetic drift, which can limit the species'
ability to purge deleterious mutations due to reduced efficacy of purifying selection at low N_e,
coupled with increasing efficacy of drift which increases their probability of fixation. This
phenomenon allows for weakly deleterious mutations to segregate at higher frequencies
which can negatively affect the population and lead to an increased risk of extinction (Lande
1994; Spielman, Brook, and Frankham 2004; Khan et al. 2021).

Similarly, populations that have recently undergone rapid expansion often display genetic
signatures of low effective population size, partly due to allele surfing, where certain gene
variants increase in frequency among individuals at the forefront of the expansion (Welles
and Dlugosch 2018; Cisternas-Fuentes and Koski 2023; Excoffier and Ray 2008; Graciá et
al. 2013). Likewise, the 'founder effect' can leave a distinct genetic imprint by reducing
genetic diversity, often resulting in greater genetic differentiation between populations
(Szewczyk et al. 2019; Jarausch et al. 2021). These signals are often associated with
historical (post-glacial), contemporary (often human-mediated) introductions of species
(Sendell-Price, Ruegg, and Clegg 2020) or by range expansions. The latter has been well
documented in the recent, rapid range expansion of the grey wolf (*Canis lupus*) across
Europe, where recolonisation of these areas by expanding populations can lead to altering
genetic structure across the landscape (Gustafson et al. 2018; Szewczyk et al. 2019;
Jarausch et al. 2021).

Habitat fragmentation or population displacement can also threaten the genetic integrity of
species through introgression and hybridisation. As the boundaries between species
populations overlap, or through the introduction of invasive species, populations may
become more susceptible to interbreeding with closely related sister species. Hybridisation
with closely related taxa has been shown to have negative consequences, particularly in
cases for endangered species (Allendorf et al. 2001). These arise through outbreeding
depression which causes a reduction in the fitness of offspring (P. W. Hedrick et al. 2019) or
genomic swamping whereby parental genetics are overwhelmed by the hybrid, potentially
leading to their extinction (Howard-McCombe et al. 2023). Hybridisation can also impact how
a species is managed, with there often being uncertainty over the classification and
protection of hybrids in the wild (Senn et al. 2019; Moroni et al. 2022).

Adaptive introgression, which occurs through successful hybridisation between two species,
can be particularly beneficial for populations recovering from a bottleneck (Philip W. Hedrick
2013). In such cases, genetic diversity is low, and introgression can introduce new variants
into the offspring's genome. This phenomenon has been observed in the Alpine ibex (*Capra
ibex*) (Brambilla et al. 2024) and Scottish wildcat (*Felis silvestris*) (Howard-McCombe et al.
2023), where adaptive introgression has replenished genetic diversity in previously depleted
genomic regions, such as immune-related genes, thereby increasing the populations'
adaptive potential. This can be beneficial for recovering populations, yet adds to the
complexity of managing hybrid individuals, as distinguishing signals of introgression are
complex and require well-defined local ancestry estimates (Howard-McCombe et al. 2023;
Brambilla et al. 2024).

Therefore, it is crucial to monitor and investigate genomic signals to understand how
populations adapt to and recover from environmental change. This knowledge can inform
conservation strategies, such as prioritising populations for protection (von Takach et al.

2024), monitoring reintroductions (Gajdárová et al. 2023) or translocations (Farquharson et
al. 2022), identifying hybrid individuals to inform action plans (Senn et al. 2019) and
contributing to ex-situ breeding programmes (Speak et al. 2024). Additionally, capturing the
temporal fluctuations of genomic changes in a recovering population undergoing range
expansion is essential to inform conservation management strategies to prevent long-term
genetic fragmentation and monitor loss (or gain) of adaptive potential (Thomas et al. 2022).

Several of the aforementioned environmental threats are known to severely impact
populations of the European polecat (*Mustela putorius*), a medium-sized carnivore of the
Mustelidae family. This family is a great example of adaptive radiation, as species belonging
to this family occupy a wide range of niches within different climates and habitats (Schluter
2000; Koepfli et al. 2008). Despite this, and despite the polecat inhabiting a wide
geographical range across mainland Europe (Virgós 2003; Mestre, Ferreira, and Mira 2007;
Brzeziński, Marzec, and Żmihorski 2010), in several of these countries their populations are
listed as or suspected to be declining (Maran et al. 2016; Croose et al. 2018) due to habitat
fragmentation, prey availability, poisoning, and persecution in continental Europe (Eeraerts
et al. 2022; Barrientos et al. 2024; Szapu et al. 2024). Hybridisation has also been known to
occur between the European polecat and other closely related mustelid species, such as the
Steppe polecat (*Mustela eversmanii*) and the severely endangered European mink (*Mustela
lutreola*) (Cabria et al. 2011; Cserkész et al. 2021).

Polecat population in Britain

In Britain, the European polecat has undergone an extensive population reduction and range
contraction, followed by a subsequent rapid range and population expansion over the last 70
years (Langley and Yalden 1977; Birks and Kitchener 1999; S. 2008; Costa et al. 2013;
Croose 2016). The population experienced a severe decline in numbers during the late 19th
century as the species was persecuted heavily by gamekeepers and regarded as pests
which impacted gamebird populations (Birks and Kitchener 1999; Costa et al. 2013). This led
to a contraction in the species range, with small pockets of populations in Cumbria and
Scotland persisting until the early 20th century and a stronghold in central Wales and the
neighbouring English border counties of Herefordshire and Shropshire (Langley and Yalden
1977; Harris et al. 1995).

The existing stronghold in Wales started to recover in population size after the end of the
First World War This resulted in a period of reduced persecution and coincided with the
recovery of rabbit populations in the post myxomatosis period, a key prey species in the
polecats diet (Langley and Yalden 1977). To aid its protection, the polecat was listed on
schedule 6 of the UK Wildlife and Countryside Act in 1981, and since 2007 has been listed
as a priority species on the UK Biodiversity Action Plan (BRIG 2007). As the population
began to expand eastwards across its former range, it came into contact with populations of
feral domestic ferrets (*Mustela putorius furo*) (Davison et al. 1999; Costa et al. 2013;
Etherington et al. 2022). Ferrets are thought to be derived from the European polecat
through domestication that occurred over 2000 years ago (Blandford 1987; Davison et al.
1999) and have long been used for hunting purposes. In Britain, they are known to have
been used for predominantly hunting rabbits, and are thought to have been released into the
wild and accidentally introduced. Owing to the short evolutionary distance between the two
species, the European polecat and the domestic ferret readily hybridise and this has been
well-documented to occur in Britain (Costa et al. 2013; Etherington et al. 2022).

Previous work on the British population of polecats has focussed on assessing genome
introgression between European polecats and feral domestic ferrets (Davison et al. 1999;

Costa et al. 2013; Etherington et al. 2022) or has aimed to model their demographic history
(Costa et al. 2013). These studies identified high degrees of genome introgression in British
polecats outside of their stronghold, emphasising the importance of the Welsh population for
conservation of the British European polecat. However, genetic assessments for the Welsh
population have either been conducted using microsatellites and mitochondrial genes, which
did not provide enough power to confidently assess the size of the genetic bottleneck (Costa
et al. 2013), or were limited in the number of Welsh individuals used in their assessment
(Etherington et al. 2022).

Here, we are able to characterise the demographic history of the British European polecat
population through using whole genome resequencing of English, Welsh and mainland
European polecats. We also further mine the genome for signatures of the population
bottleneck, documenting what impact the decline and rapid expansion had on the population
and assessing the conservation implications for British polecats in comparison to mainland
European polecat populations.

152
153 **2.0 Methods**
154
2.1 Sample Collection
For this study, 31 roadkill British polecat samples were resourced from the Vincent Wildlife
Trust 2014-2015 survey (Croose 2016) (Figure 1b). Carcasses were collected by members
of the public who were requested to freeze the carcass where possible. These were then
sent to the Centre for Ecology and Hydrology (CEH) via a pre-paid postage box and the
carcasses were frozen upon receipt. Samples were selected based on geographical location
from areas around Wales and from a trajectory across England from west to east to capture
any diversity across the predicted range expansion of the polecat in Britain (Figure 1c),
(Supplementary table sheet 1 (S1)). The home ranges of polecats in Britain are known to
vary seasonally, and male polecats tend to disperse further than females (Birks and
Kitchener 1999), with males travelling as far as 8 km between dens (Blandford 1987). To
avoid incorporating familial bias into our selection through potentially selecting related
individuals, we selected carcasses that were located greater than 10 km apart. We further
increased our sample size through the inclusion of publicly available data consisting of eight
domestic ferrets genomes, 19 British polecats, and 15 European mainland polecat samples
(Etherington et al. 2022).

171
2.2 DNA Extraction
DNA extraction from tissue samples was carried out using a Qiagen^{TM} DNEasy Blood &
Tissue kit according to the manufacturer's protocol. Samples were quality checked using a
Nanodrop Spectrophore and DNA quantity was checked using a Qubit Fluorometer.
Samples with a low yield, and with 260/280 and 260/230, absorbance values outside the
desired range (260/280 <1.80, 260/230 <2.0) were further cleaned up using Beckman
Coulter^{TM} AMPureXP Beads. A total of 31 samples were selected for library preparation and
sent for sequencing. Figure 1c shows the distribution of the samples sent for sequencing
and the location of British polecats already sequenced.

181
182 **2.3 Library preparation & sequencing**
183 Samples were sequenced by The Earlham Institute Technical Genomics with LITE library
184 preparation (Perez-Sepulveda et al. 2021) and sequenced using an Illumina NovaSeq 6000

185 generating paired-end (2 x 150bp) reads. Samples were run on one lane with a predicted
186 coverage of 10x.
187
2.4 Sequencing QC, Read mapping & SNP calling
Raw read quality for all British and European samples were first assessed using fastQC
version 0.11.9 (Andrews 2010), and any reads below the threshold (Q20) were trimmed
using TRIMMOMATIC version 0.39 (Bolger, Lohse, and Usadel 2014). Raw reads were
mapped to a new version of the domestic ferret genome MusPutFur1.0 scaffolded with
Bionano data ([GCA_920103865.1](#)) (Peng et al. 2014; Etherington et al. 2022) using BWA
MEM version 0.7.17 (Li and Durbin 2009). Duplicate reads were then marked using
PicardTools version 2.1.1 ("Picard Toolkit" 2019). Single nucleotide polymorphisms (SNPs)
were then called using GATK version 4.5.0 (McKenna et al. 2010) following the best practice
workflow for SNP discovery (van der Auwera and O'Connor 2020). Briefly, BaseRecalibrator
was run on each sample bam file to detect and correct patterns of systematic errors in the
base quality scores. The recalibration metric was then applied to each sample bam file, and
HaplotypeCaller was run for each sample before consolidating all of them together using
GATK GenomicsDBImport. For downstream analysis, BCFtools version 1.15.1 (Danecek et
al. 2021) was used to select only biallelic SNPs with a quality score (QUAL) of at least 20
and a minor allele count of at least three (AC3). A further step of pruning variants over a
linkage disequilibrium (LD) threshold of r2 ≥0.6 within 50kb windows was carried out using
the bcftools prune command. The resulting SNP dataset was used for all downstream
analyses unless stated. For GONE analyses, all variants called before pruning were used.
207
2.5 Relatedness
We assessed relatedness between samples using the KING algorithm in PLINK version 1.9
(Purcell et al. 2007; Chang et al. 2015). KING kinship coefficients determine first-degree
relations (parent-child, full siblings) as values between 0.17-0.35, second-degree relations
correspond to 0.08-0.17 and third-degree relations as between 0.04-0.08 (Manichaikul et al.
2010). One pairing between samples S17 and S0012_VWT665 had a kinship coefficient of
0.07, suggesting a third-degree level of relatedness. Both of these individuals were English
polecats but were found in very different geographical locations (~291km apart). We
removed S17 from all of the downstream analysis as this sample was found in Waterrow,
England, and was therefore outside of the west-to-east trajectory that we were hypothesising
was the route of the Welsh polecat expansion.
219
2.6 Population Structure
We first generated a principal component analysis (PCA) using the whole-genome SNP
dataset of 65 polecats and 8 domestic ferrets in PLINK version 1.9 (Purcell et al. 2007;
Chang et al. 2015) to determine the British population structure. A Spearman correlation test
was implemented in R version 4.1.1 (R Core Team 2024), using values from PC1 and
sample longitude coordinates, to ascertain whether population structure in Britain could be
correlated with geographical location. We used ADMIXTURE version 1.3 (Alexander,
Novembre, and Lange 2009) to perform ancestry runs on the SNP dataset for K values 1-12,
and a CV error was used to compare all K values. Replicates for each ancestry run were
carried out using the custom python wrapper (admixturewrapper.py). Pairwise F_{st} values
between populations were computed using VCFtools version 0.1.16 (Danecek et al. 2011).
231

To further detect structure within the British population, we tested for isolation by distance
between individuals. We did this by first calculating the genetic distance (Hamming distance)
between British samples using PLINK version 1.9 (Purcell et al. 2007; Chang et al. 2015).
We then used a geographic distance matrix between British samples using the geosphere
package in R version 4.1.1 (R Core Team 2024; Hijmans 2024), inputting longitudinal and
latitudinal coordinates to output distance in km. A mantel test was used to test for isolation
by distance between the two matrices.
239
240   We used RAXML-ng version 0.9.0 (Stamatakis 2014) to infer a phylogenetic tree from our
241   whole-genome SNP dataset. The phylogeny was generated using the GTRGAMMA model of
242   evolution with the Lewis bias ascertainment correction and 100 rapid bootstraps.
243

2.7 Genomic Diversity

To assess the genetic diversity between populations, we used VCFtools version 0.1.16
(Danecek et al. 2011) to detect nucleotide diversity in 20 Mb windows across all samples.
We tested for significant differences in the nucleotide diversity estimates between
populations by using an analysis of variance test (ANOVA) implemented in R version 4.1.1(R
Core Team 2024). Genome-wide SNPs that were private to one population, or that were
shared across populations, were identified using VCFtools. The observed proportion of
heterozygous sites and the inbreeding coefficient (f) for each individual per population were
also calculated using VCFtools.
253

2.8 Detection of historic bottleneck: Runs of homozygosity and identification of deleterious mutations

To detect signatures associated with a genetic bottleneck, we calculated the distribution of
runs of homozygosity (ROH) using PLINK. Since calling ROH can be sensitive to low
sequencing depth in areas of the genome, which can introduce genotype errors (Ceballos et
al. 2018), we only calculated ROH for individuals per population that had an average read
depth of 6.0 or higher. We used the following recommended parameters to call ROH for low
coverage samples - plink --geno 0.5 --maf 0.1 --homozyg --homozyg-density 50 --homozyg-
gap 1000 --homozyg-kb 300 --homozyg-snp 50 --homozyg-window-het 4 --homozyg-
window-missing 5 --homozyg-window-snp 50 --homozyg-window-threshold 0.05 --aec --
double-id (Ceballos et al. 2018). We then categorised ROH into three length classes; 300-
500kb, 500 kb-1Mb, and >1Mb.
266

SnpEff version 5.1 (Cingolani et al. 2012) was used to detect the functional impacts of the
biallelic SNPs on the genome of the European polecat. SnpEff compares a set of variants to
a known annotation database and compares the derived alleles within genic regions to
predict the impact of that variant. In this case, we used the domestic ferret annotation as the
input database. We assessed whether mutational load differed between populations and this
was calculated based on homozygous and heterozygous mutational load within missense
and loss-of-function (LoF) categories.
274

Due to the potential bias that could arise by aligning reads to the domestic ferret reference
genome to infer deleterious mutations within the polecat genome landscape, we polarised
our variant calls using the Steppe polecat (*Mustela eversmanii*) and weasel (*Mustela nivalis*)
as outgroups to infer the ancestral allele at sites called by SnpEff version 5.1 (Cingolani et
al. 2012) as deleterious.

To infer the ancestral allele within outgroups, we selected species from the Mustelidae

family with available genomes: European polecat (*Mustela putorius*) (GCA_902207235.1)

(Etherington et al. 2020), domestic ferret (*Mustela putorius furo*) (GCA_920103865.1) (Peng

et al. 2014; Etherington et al. 2022), Steppe polecat (*Mustela eversmanii*)

(GCA_963422785.1), Black-footed ferret (*Mustela nigripes*) (GCF_022355385.1) (Kliver et

al. 2023), stoat (*Mustela erminea*) (GCA_009829155.1), stone marten (*Martes foina*)

(GCA_040938555.1) (Tomarovsky et al. 2025), and Eurasian otter (*Lutra lutra*)

(GCA_902655055) (Mead et al. 2020). Additionally, we included three mammalian

outgroups: human (*Homo sapiens*) (GCA_000001405.15) (Schneider et al. 2017), domestic

cat (*Felis catus*) (GCF_000181335.3) (Buckley et al. 2020), and domestic dog (*Canis lupus*

familiaris) (GCF_011100685.1) (Wang et al. 2021) to construct a whole-genome alignment.

We generated a whole-genome alignment using Progressive Cactus v2.2.0 (Armstrong et al.

2020) without specifying branch lengths. Next, we extracted ancestral repeats from the

alignment and inferred a phylogenetic tree under a neutral model using phyloFit (part of the

PHAST toolkit) (Siepel and Haussler 2004). This tree was then used as a guide for rerunning

Progressive Cactus to refine the alignment.

We subsequently extracted variants from this alignment for four sister species (domestic

ferret, European polecat, Steppe polecat, and weasel), using the domestic ferret as the

reference, via halSNPs v2.2 (Hickey et al. 2013). Finally, we intersected the extracted

variants with our called SnpEff variants to identify alleles derived in the European polecat.

2.9 Historic effective population size

Recent historic effective population size (N_e) was estimated using GONE (Santiago et al.

2020). GONE estimates recent effective population size by calculating LD between pairs of

SNPs and finds the N_e that best fits the observed LD spectrum. GONE assumes that the

population being tested is closed, with no recent admixture to allow for unbiased estimates in

the calculation of effective population size. Therefore, we ran GONE for Welsh, English, and

European populations separately. We restricted GONE runs for the English population to

those individuals in the south, central and eastern regions of England. This was to account

for any potential gene flow between individuals located within the border counties of Wales

and Welsh individuals, as well as lack of gene flow due to the geographic isolation of

individuals within the north of England. We further removed an individual (S0024_VWT1661)

from the Welsh population run due to apparent recent hybridisation (F1 hybrid) identified by

PCA and ADMIXTURE analyses. We accounted for low sequencing depth by further

removing individuals from the analyses that had a mean sequencing depth at sites of less

than 5x coverage. For each population, we simulated 2000 generations and calculated 400

bins. GONE takes up to 200 chromosomes as input data, so we removed scaffolds that

overlapped with the X chromosome, using the annotated X chromosome of the black-footed

ferret (*Mustela nigripes*) for reference (Kliver et al. 2023) and used the largest 100

autosomal scaffolds as input, constituting 78% of the whole genome assembly. GONE

calculated LD for up to 10,000 SNPs per scaffold. Each simulation was run 100 times, with a

new subsample of 10,000 SNPs for each run. For each population, we plotted the mean Ne

for the 100 runs.

3.0 Results
327

3.1 Whole Genome Resequencing
We generated whole-genome sequence data for 31 European polecats from Wales and
England. Samples had been chosen from the location of the known refugium (Figure 1a),
across a trajectory of the historic range expansion following the population's recovery from a
genetic bottleneck. These were combined with an already published dataset of 34 European
polecats from Italy (5), Germany (2), Austria (3), France (3), Spain (2), Wales (3) and
England (16), along with 8 domestic ferrets. Sequence reads were aligned to the domestic
Ferret genome. Initial exploratory analyses of called variants determined that there was a
presence of batch effect (Lou and Therkildsen 2022) across the different sequencing runs,
and steps taken to alleviate this effect are described in the Supplemetary Methods. Higher-
coverage samples were downsampled to account for coverage biases, and SNPs were then
called after downsampling (Supplementary table S2). We identified a total of 2,941,545
SNPs.
341

3.2 Population Structure Using Genomic Data
Principal component analysis (PCA) of whole-genome SNPs revealed 3 distinct clusters of
samples of polecats from (1) mainland Europe, (2) Britain, and (3) domestic ferrets (Figure
2a). Within this, PCs 1 (26%) & 2 (15.8%) revealed the English and Welsh samples
clustered together and away from mainland European samples. To assess structure within
the British population, we repeated the PCA only using the 49 British samples, which
revealed a geographical separation of samples from West (Welsh) and East (English)
(Figure 2b). Longitudinal coordinates strongly correlated with PC1 (r_s = -0.754, p <0.01).
Outliers to this relationship that clustered with the Eastern individuals are from northwestern
and southern England, suggesting that Welsh samples were distinct from English samples.
353

ADMIXTURE analyses were best supported by the cross-validation error of *K=3*
(Supplementary Figure S3 & S5), indicating the existence of three genetically distinct
populations across all of the sampled populations; domestic ferret, British (encompassing
English and Welsh populations) and European polecats. These results appear to resolve the
British population as a homogenous group distinct from the European mainland population,
however this could also be due to the small sample size representing different European
populations (Figure 2c). Under these parameters, the Welsh population demonstrated a
lower degree of admixture with the domestic ferret than English individuals (Figure 2c)
(Supplementary table S3), with the exception of one individual, which is likely a polecat-ferret
hybrid (S0024_VWT1661). These results suggest that there is evidence of hybridisation and
introgression occurring between English polecats and feral domestic ferrets at varying
degrees across the polecats distribution.
366

367 We performed a mantel test to examine the correlation between genetic distance and
368 geographic distance across individuals from Wales and England. The analysis revealed a
369 positive statistically significant correlation (r = 0.333, p = 0.001), suggesting that genetic
370 differentiation increases with geographic distance, demonstrating that individuals in Britain
371 are isolated by distance.
372

373 Pairwise F_{st} values were computed to detect the genetic divergence between populations.
374 Our results showed a comparison of English and Welsh populations having an F_{st} value of
375 0.036, compared to 0.106 between English and European and 0.145 between Welsh and
376 European (Supplementary Figure S4).
377
3.3 Genomic Diversity in Britain in Comparison to Mainland Europe
As the British populations have undergone a severe population contraction prior to their
current recovery, we investigated the impact of these demographic events on the genomic
diversity of the different populations. European populations demonstrated a higher number
of private SNPs (2,290,307) in comparison to English (838,471) and Welsh (578,869)
(Figure 3a). This could be attributed to population structure and also demonstrates the
genetic divergence between British and European mainland polecats. Welsh individuals
showed a lower number of private SNPs in comparison to English individuals, although still
demonstrating unique genomic diversity that still resides within the stronghold population.
387
388 Nucleotide diversity estimates (Tajima's π) (Supplementary table S4) revealed a significant
389 difference amongst populations (H = 7.14, df = 2, p = 0.02), with the English population
390 having significantly higher diversity when compared to the Welsh population (p <0.001) and
391 European population (p = 0.01) . However, there was no significant difference between the
392 Welsh and European populations (Figure 3c).
393
Inbreeding coefficients (F2,60 = 17.6, p <0.001) and proportion of heterozygous sites (F2,61
= 18.16, p <0.001) significantly differed across the three populations (Figure 3b & 3d). Post-
hoc comparisons found that the European population had a significantly higher inbreeding
coefficient (p <0.001) and significantly lower proportion of heterozygous sites (p <0.001) in
comparison to individuals in the English population. However, we did not account for
population structure in the grouping of individuals due to the total sample size. French and
Austrian populations had higher proportions of heterozygous sites and lower inbreeding
coefficients than German, Italian, and Spanish populations (Supplementary Figure S7a &
S7b). There was no significant difference between the Welsh and European populations for
either inbreeding coefficient or the proportion of heterozygous site estimates. English
individuals did show a significantly higher proportion of heterozygous sites (p <0.001) and
significantly lower inbreeding coefficient (p <0.001) than those in the Welsh population.
406
3.4 Runs of Homozygosity (ROH) & Deleterious Mutations
To assess whether the Welsh population had undergone prolonged periods of inbreeding
due to the population bottleneck, we investigated the presence of ROH within polecat
genomes. We calculated the realised inbreeding coefficient (F_{ROH}), the proportion of the
genome that fell within ROH, which ranged from 0.33 - 0.76. Post-hoc comparisons showed
that F_{ROH} values were significantly different between European and English individuals (W =
2.19, p = 0.0143) and English and Welsh (W = 2.42, p = 0.0077) individuals, but not between
European and Welsh (W = 0.23, p = 0.4074) (Figure 4a & Supplementary table S5).
415
We separated ROH lengths into different classes to visualise short (300-500kb), medium
(500-1Mb), and long (>1Mb) ROH across the genomes of individuals belonging to the
Welsh, European, and English populations. Longer ROH are indicative of recent inbreeding
events, with shorter ROH being indicative of historic events. Individuals belonging to the
Welsh population were seen to have the lowest number of ROH segments in all length

classes. Individuals belonging to the English population were seen to have a greater
proportion of their genome within longer ROH segments (>1mb) (58.3% ± 1.50 SE) in
comparison to the European (53.8% ± 2.14 SE) and Welsh (55.0% ± 1.90 SE) individuals
(Figure 4b).
425
We investigated whether the mutation landscape differed among individuals belonging to
European, Welsh, and English populations. We assessed derived mutational load across
two functional categories as reported by SnpEff: missense and loss of function (LoF). We
calculated both homozygous and heterozygous mutations per individual per category.
Homozygous mutations in both categories were expected to be the most deleterious, and we
also considered the load of heterozygous mutations.
432
Homozygous LoF mutation load differed significantly between populations ($F_{2,60} = 16.81$, p
<0.01), with both Welsh and English populations showing a significantly higher homozygous
load within LoF mutations in comparison to the European populations (p <0.01) (Figure 5a).
There was also a significant difference amongst populations in LoF masked load ($F_{2,60} =$
5.584, p <0.01), with individuals in the English population having a significantly higher
masked load than Welsh individuals (p <0.01).
439
Missense homozygous mutations significantly differed between populations ($F_{2,60} = 31.39$,
p < 0.01). Post-hoc comparisons revealed that Welsh individuals had significantly more
missense homozygous mutations than English and European individuals (p <0.01) (Figure
5c).
Masked, heterozygous missense load also significantly differed among populations, ($F_{2,60} =$
14.77, p < 0.01), with English individuals showing a higher masked load than both Welsh (p
<0.01) and European individuals (p <0.01) (Figure 5d).
447
3.4 Historic effective population size (N$_e$)
We used GONE to estimate the recent historic effective population size for the British
population of European polecats. Our estimates using GONE correlate with a demographic
decline in population size occurring around 30-40 generations ago - which using an estimate
for generation time of the polecat of 4 years (Di Marco et al. 2013), corresponds to the mid-
late 19th century (1854-1894) (Figure 6). Recovery of the polecat population in Wales based
on GONE estimates fits with previous estimates of 10-12 generations ago (1966-1974)
(Supplementary table S7).
456
4.0 Discussion
Here, we provide the first evidence of a genetic bottleneck having occurred within the British
population of polecats. Our analyses indicate that the population drastically decreased in
effective population size from the late 1800s, with this bottleneck persisting for ca. 100
years. Our analyses also indicate that the population began recovering from 1960-70
onwards. This coincides with a prolonged period of relaxed persecution, and the introduction
of protective legislation. Our results demonstrate that the British polecat population is distinct
from European mainland populations, and is recovering from the genetic bottleneck.
Conservation efforts should focus on increased protection measures and monitoring of the
Welsh population, as it represents unique genetic integrity in comparison to admixed,
English populations. Our study also highlights the importance of genomic data in being able

to detect signatures within populations to fully detect and monitor the success of their
recovery.
470
4.1 Comparison of British to Mainland European populations
Our analyses indicate that the British polecat population is genetically diverged from
European mainland populations. This is highlighted in pairwise F_{st} estimates per population
where a higher F_{st} value (closer to 1.0) suggests a population has low levels of breeding and
therefore is more genetically distinct from populations compared to lower F_{st} values. Our
results indicate that English and Welsh populations have a low fixation index (0.03) and are
therefore likely to share genetic variation. F_{st} values between English and European (0.106)
and European and Welsh populations (0.145) are slightly higher, suggesting a greater
degree of genetic distinction between mainland European and British polecats, which is also
demonstrated by the degree of private and shared polymorphisms across the populations
(Figure 3a).
482
This is also evident from our ADMIXTURE analyses which further corroborates the genetic
divergence seen between Welsh and European polecats (Etherington et al. 2022). British
and European polecats are thought to have been separated for ca. 8ky (Montgomery et al.
2014), since the flooding of the Doggerland bridge after the Last Glacial Maximum. Several
widespread mammal populations that are native to Britain demonstrate genetic distinction
due to isolation over time from mainland European populations. For example, Irish and
British populations of red fox (*Vulpes vulpes*) demonstrate unique genetic diversity that
distinguishes them from mainland European populations (McDevitt et al. 2022). Likewise,
native British populations of roe deer (*Capreolous capreolus*) demonstrate genetic
differentiation from mainland European populations, most likely due to colonisation and
isolation events (de Jong et al. 2020). Populations of British red deer (*Cervus elaphus*) also
demonstrate genetic distinction from mainland European populations, but show a slightly
lower genetic distance to Iberian populations, potentially suggesting retention of genetic
diversity within populations from similar migration waves during postglacial colonisation
(Carranza et al. 2024). Therefore, the genetic distinction we observe between Welsh and
European mainland polecats likely results from their recent isolation, mirroring the pattern of
genetic divergence seen across other British populations of wide-ranging mammals
(Montgomery et al. 2014; McDevitt et al. 2022; Carranza et al. 2024).
501
4.2 Population Structure
Our analyses indicate population structuring based on geographic location within polecats in
Britain (Figure 2b). Ancestry estimates also demonstrate a higher incidence of introgression
in polecats that are found at the furthest edges of the polecat range from the Welsh refugium
population (Figure 2c). We also observe a similar relationship in our maximum likelihood
phylogeny, whereby English individuals that are found at the more eastern and southern
edge of the polecat's range are clustered more closely to domestic ferrets than to the Welsh
individuals (Supplementary Figure S6). In agreement with (Costa et al. 2013), we find the
least introgression of domestic ferret genes within individuals found in Wales and English
border counties. This observed pattern fits the narrative that as the polecat population
expanded eastwards, individuals hybridised with feral domestic ferrets within those regions.
However, the presence of introgressed polecats within these regions may also have been
exacerbated by covert reintroductions of individuals from hybrid stocks (Birks and Kitchener
1999).

4.3 Demographic Analyses

Using whole genome sequence data, it was possible to detect and analyse historic trends in effective population size in the British population of the European polecat. The British polecat bottleneck was thought to have occurred between the late 1800s to the early 1900s, with populations showing recovery since the 1970s (Langley and Yalden 1977). The most recent historic genetic bottleneck was evident in our results using GONE, with a sharp decrease in N_e occurring in the Welsh population around 30-40 generations (120-160 years ago), from the sampling time of 2014-2015. This coincides with the known history of persecution of polecats from gamekeepers, as with other populations of terrestrial carnivorans in Britain, which subsided for a period during and following World War I, and again after legal protections came into place (Langley and Yalden 1977; Sainsbury et al. 2019). Alongside legal protections, it is thought that increasing rabbit (*Oryctolagus cuniculus*) populations, an important prey for polecats (Sainsbury et al. 2020), which had declined due to myxomatosis and began recovering from the mid-20th century, are also closely linked with the recovery of the polecat (Langley and Yalden 1977; Sumption and Flowerdew 1985; Sainsbury et al. 2019).

We observed that historic N_e estimates were lower for the English population than the Welsh (Figure 6, Supplementary table S8). This could be explained both by the rapid expansion of individuals from Wales to England and by the hybridisation and subsequent introgression of those individuals with populations of feral domestic ferrets. Our results also highlight that modelling demographic histories can be impacted by hybridisation and population structure (Novo et al. 2023). Previous work (Costa et al. 2013), using microsatellites in British polecats could not detect any evidence of a genetic bottleneck. This could be due to the lower resolution from the smaller representation of genetic markers used but could also be due to the admixture from domestic ferret populations as suggested by the authors. We observe this pattern within the English population having greater levels of introgression than the Welsh population and therefore observe not only a lower historic effective population size but also a lower present day effective population size when just modelled using the English population. We see that population structure can obscure the recent historic effective population size when trying to model the European population's demographic histories (Supplementary Figure S8). Modelling these as a whole population is difficult due to the different environmental pressures that each population faces. GONE authors advise not to model demographic histories with less than 5 individuals (Novo et al. 2023), of which only the Italian population was viable from our European samples. However, our model reconstruction demonstrated the unreliability of this with low power and possible population structure (Supplementary Figure S8).

Census data in 2018 estimated that the polecat population in Britain was currently 88,300 (Mathews et al. 2018). Our estimates corroborate survey data from the Vincent Wildlife Trust that the Welsh population is expanding, with more 'true polecat' sightings in Welsh counties and border counties such as Herefordshire and Shropshire (Croose 2016), which are areas that were thought to contribute to the historic stronghold of the polecat (Langley and Yalden 1977), and these areas also having the individuals with the least admixture (Figure 2c).

4.4 Genomic signatures associated with the population bottleneck

We hypothesised that we would see genomic signatures within the population associated with the prolonged genetic bottleneck period. Our results demonstrate the presence of potentially harmful deleterious mutations within both Welsh and English individuals. It also shows that heterozygous, potentially masked deleterious variants are more prevalent in admixed English individuals, and that the Welsh population has a higher unmasked load of missense variants. This is as expected, as periods of inbreeding in Welsh populations would lead to a higher presence of homozygous deleterious variants within the population. English individuals appear to have higher heterozygosity, due to the introgression with domestic ferret. This is also confirmed in the higher nucleotide diversity across the genomes of English individuals. However, no significant difference amongst populations was found in mutations that were categorised as homozygous LoF, suggesting these mutations may have been purged from the Welsh population due to their expansion and purifying selection acting upon them. Further investigation into historic genomic signatures through the addition of historical samples would provide an insight into temporal changes of mutational load over time, which would assist in resolving potential signatures of purifying selection within the Welsh population, and would be consistent with other studies that have observed this (van der Valk et al. 2019; Grossen et al. 2020; Dussex et al. 2021)

Individuals belonging to the Welsh population had higher overall ROH in their genomes in comparison to the English population (Figure 4a). This could be due to the long-term, low effective population size within the Welsh refugium. Welsh individuals also demonstrated a higher mean of shorter ROH within the genome in comparison to the English populations. This is indicative of more ancestral inbreeding events, and is a strong indicator that the population underwent a bottleneck further back in time (Saremi et al. 2019). Longer ROH lengths can indicate recent inbreeding events, as recombination has not yet acted on breaking apart these segments (Ceballos et al. 2018). Individuals belonging to the English population were shown to have a higher mean number of long ROH in their genome than Welsh. This could be associated to the introgression with feral domestic ferrets that have been subjected to years of selective, human-mediated breeding. However, it may also be indicative of the historic bottleneck, followed by the expansion of some individuals to low areas of genetic connectivity and therefore increased occurrence of inbreeding events ((Manunza et al. 2016; Colpitts, McLoughlin, and Poissant 2022).

European individuals overall showed a higher mean F_{ROH} than both English and Welsh individuals, as well as a higher mean proportion of medium length ROH. However, across the length classes and F_{ROH} values there was a varied ROH landscape per population (Supplementary Figure 7c). This could be due to population-level pressures persisting in their environment at the time (Croose et al. 2018) and may indicate the necessity for similar genomic studies in polecat populations in regions of Europe where the population is declining.

4.5 Conservation Implications

Here, we provide the first evidence of the severity of the population bottleneck experienced by the Welsh polecat population from the late 1800s to the mid-1950s. We demonstrate that the population started to expand when persecution pressures were lessened and after protective legislation came into action, thus supporting legal protection as a conservation management tool.

612
We present a unique scenario of population recovery in a once heavily persecuted mammal
in Britain. The recovery can largely be attributed to protective legislation and the resurgence
of rabbit populations following the myxomatosis outbreak, as well as to the introgression with
feral domestic ferrets, which has facilitated the eastern range expansion of the population.
Genomic introgression with a domesticated species is not a typical conservation
management tool but has been witnessed to assist in the recovery of other species, such as
the American bison (*Bison bison*) (Stroupe et al. 2022), and Iberian ibex (*Capra pyrenaica*)
(Münger et al. 2024). However, the decline of the wildcat (*Felis silvestris*) population in
Scotland has been severely hampered by hybridisation and subsequent introgression with
the domestic cat (Howard-McCombe et al. 2023). It could be that due to the relatively recent
nature of the hybridisation of the European polecat and domestic ferret in Britain, the harmful
consequences are yet to surface. However, (Costa et al. 2013) demonstrated that the lack of
F1 hybrids observed within the British population suggests that these hybridisation events
are rare. Feral ferret populations are hard to distinguish in mainland Britain but are known to
thrive on surrounding islands where there is a lack of or reduction in competition from
polecats and other predatory mammals (Kitchener and Birks 2008). This indicates that F1
hybridisation events between feral ferrets and polecats in wild populations are likely to be
rare, yet they still represent a potential threat to the genetic integrity of the recovering
polecat population.
632
Our analyses also demonstrate that the Welsh population holds unique genomic diversity
that could be important for the resilience of the population. This could have important
conservation implications for reintroducing polecats from Wales to other parts of the country.
Additionally, conservation concerns are underscored by the unmasked missense mutational
load persisting in the genome as a result of the prolonged low effective population size
(Robinson et al. 2023). The monitoring of this elusive species is hard, due to its low
population density and nocturnal habits (Croose et al. 2018; Barrientos et al. 2024).
Monitoring through non-invasive methods, such as the Vincent Wildlife Trust survey, where
sightings and suspected roadkill are reported, has proven successful and has helped to
document the polecat's recovery within Wales and England (S. 2008; Croose 2016).
However, these efforts are not matched in some parts of mainland Europe and therefore the
polecat's status in some countries needs to be clarified (Kominos and Galanaki 2018;
Hofmeester et al. 2019).
646
In a study aiming to document the status of the polecat across its known range, of the 34
countries where data were found, 20 countries reported that the polecat population was
declining (Croose et al. 2018). Only two countries reported the polecat population as
increasing, i.e., Estonia and Britain, although there is some evidence that populations in
Poland could be genetically stable (Martínez-Cruz, Zalewska, and Zalewski 2022).
Therefore, conservation priorities should be to continue monitoring the Welsh population to
protect its genetic legacy and ensure the population's continued expansion. Given the
reported declines in other known polecat ranges (Baghli and Verhagen 2003; Costa,
Fernandes, and Santos-Reis 2014; Barrientos 2015; Szatmári et al. 2021; Frantz et al.
2022), conserving the Welsh population is crucial for supporting the global population. British
polecats face threats not only from hybridisation with feral ferrets, but also from road traffic
collisions, which hinder movement during peak dispersal (Barg, MacPherson, and Caravaggi

2022). Conservation efforts should therefore prioritise reducing habitat fragmentation and
minimising the need for polecats to cross major roads.
661
It should also be noted that the genetic recovery of a population can often lag behind its
demographic recovery (Adams and Edmands 2023; Thomas et al. 2022; Dussex et al.
2023). In the case of the British polecat population, we do see genetic structuring due to
admixture across parts of England. Another British mustelid, the Eurasian Otter (*Lutra lutra*),
has subsequently undergone a recent historic genetic bottleneck and is recovering across its
previous ranges (Thomas et al. 2022; du Plessis et al. 2023). However, barriers, such as
landscape changes, have potentially contributed to the fragmentation and low genetic
connectivity between populations. We recommend that continued monitoring be
complemented with continued analysis of genetic data, to ensure that polecat populations in
Wales do not become genetically isolated due to landscape change and continue to recover
both demographically and genetically (Thomas et al. 2022).
673
This study examined the recovery of polecats across much of Great Britain, however it did
not include data on the population in Scotland. According to survey records, it was thought
that by 1915, polecat populations were eradicated throughout Scotland. Covert re-
introductions of polecat were thought to have occurred across some areas in southern
England, Cumbria and Argyll (Davison et al. 1999; Birks and Kitchener 1999). Previous work
has highlighted the more 'ferret-like' ancestry of polecats found within regions of Scotland
(Costa et al. 2013), with feral ferret populations being found on offshore Scottish islands.
The 2014-15 survey conducted by the Vincent Wildlife Trust recorded only 29 sightings of
polecats, representing just 2% of the total records from Britain (Croose 2016). However,
some of these sightings occurred in a newly identified distribution area in Dumfries, which
may indicate a range expansion of the Cumbrian populations. Further sampling and genomic
analyses, in addition to the continued monitoring efforts, would provide greater insights into
the population structure and conservation status of the remaining polecat population within
Scotland.
688
689

Acknowledgments
RS was supported by a NERC PhD studentship through the ARIES DTP (NE/S007334/1,
2465312). WH and GE acknowledge the support of the Biotechnology and Biological
Sciences Research Council (BBSRC), part of UK Research and Innovation; Earlham
Institute Strategic Programme Grants Cellular Genomics BBX011070/1, Decoding
Biodiversity BBX011089/1, Delivering Sustainable Wheat BBX011003/1 and their constituent
work packages (BBS/E/ER/230001A, BBS/E/ER/230001B, BBS/E/ER/230001C,
BBS/E/ER/230002A, BBS/E/ER/230002B, BBS/E/ER/230003A, BBS/E/ER/230003C). The
author(s) acknowledge support from the Biotechnology and Biological Sciences Research
Council (BBSRC), part of UK Research and Innovation, Core Capability Grant
BB/CCG1720/1 supporting the work delivered via the Scientific Computing group, as well as
support for the physical HPC infrastructure and data centre delivered via the NBI Computing
infrastructure for Science (CiS) group.and the National Capability BBS/E/T/000PR9816. ACK
thanks the Negaunee Foundation for its continued support of a curatorial preparator who
sampled many of the polecats used in this study. We are also grateful to Federica Di Palma
for her early input to the project.

References

Adams, Nicole E., and Suzanne Edmands. 2023. "Genomic Recovery Lags behind Demographic Recovery in Bottlenecked Populations of the Channel Island Fox, Urocyon Littoralis." *Molecular Ecology* 32 (15): 4151–64.

Alexander, David H., John Novembre, and Kenneth Lange. 2009. "Fast Model-Based Estimation of Ancestry in Unrelated Individuals." *Genome Research* 19 (9): 1655–64.

Allendorf, Fred W., Robb F. Leary, Paul Spruell, and John K. Wenburg. 2001. "The Problems with Hybrids: Setting Conservation Guidelines." *Trends in Ecology & Evolution* 16 (11): 613–22.

Andrews, S. 2010. *FastQC: Quality Control Tool High Throughput Sequence Data*.

Armstrong, Joel, Glenn Hickey, Mark Diekhans, Ian T. Fiddes, Adam M. Novak, Alden Deran, Qi Fang, et al. 2020. "Progressive Cactus Is a Multiple-Genome Aligner for the Thousand-Genome Era." *Nature* 587 (7833): 246–51.

Auwera, Geraldine van der, and Brian D. O'Connor. 2020. *Genomics in the Cloud: Using Docker, GATK, and WDL in Terra*. Sebastopol, CA: O'Reilly Media.

Baghli, Adil, and Ronald Verhagen. 2003. "The Distribution and Status of the polecat*Mustela Putorius*in Luxembourg: Polecats in Luxembourg." *Mammal Review* 33 (1): 57–68.

Barg, Allison, Jenny MacPherson, and Anthony Caravaggi. 2022. "Spatial and Temporal Trends in Western Polecat Road Mortality in Wales." *PeerJ* 10 (e14291): e14291.

Barrientos, Rafael. 2015. "Adult Sex-Ratio Distortion in the Native European Polecat Is Related to the Expansion of the Invasive American Mink." *Biological Conservation* 186 (June):28–34.

Barrientos, Rafael, Guillermo Carmona, Tamara Burgos, Sara Martin-Garcia, Marta Vargas-Ramírez, Javier Hernández-Hernández, Victor J. Bandeira, et al. 2024. "The Genetics of the European Polecat in the Iberian Peninsula." *The Journal of Wildlife Management* 88 (7): e22628.

Birks, J. D. S., and A. C. Kitchener. 1999. *The Distribution and Status of the Polecat Mustela Putorius in Britain in the 1990s*. Vincent Wildlife Trust.

Blandford, P. R. S. 1987. "Biology of the Polecat Mustela Putorius: A Literature Review." *Mammal Review* 17 (4): 155–98.

Bolger, Anthony M., Marc Lohse, and Bjoern Usadel. 2014. "Trimmomatic: A Flexible Trimmer for Illumina Sequence Data." *Bioinformatics (Oxford, England)* 30 (15): 2114–20.

Brambilla, Alice, Noel Zehnder, Bruno Bassano, Luca Rossi, and Christine Grossen. 2024. "Genetic Evidence of a Hybrid Swarm between Alpine Ibex (Capra Ibex) and Domestic Goat (C. Hircus)." *Evolutionary Applications* 17 (8): e13761.

BRIG. 2007. "Report on the Species and Habitat Review." https://data.jncc.gov.uk/data/bdd8ad64-c247-4b69-ab33-19c2e0d63736/UKBAP-Species-HabitatsReview-2007.pdf.

Brzeziński, Marcin, Magdalena Marzec, and Michał Żmihorski. 2010. "Spatial Distribution, Activity, Habitat Selection of American Mink (Neovison Vison) and Polecats (Mustela Putorius) Inhabiting the Vicinity of Eutrophic Lakes in NE Poland." *Folia Zoologica* 59 (3): 183–91.

Cabria, M. T., J. R. Michaux, B. J. Gómez-Moliner, D. Skumatov, T. Maran, P. Fournier, J. López de Luzuriaga, and R. Zardoya. 2011. "Bayesian Analysis of Hybridization and Introgression between the Endangered European Mink (Mustela Lutreola) and the Polecat (Mustela Putorius): HYBRID IDENTIFICATION BETWEEN TWO MUSTELID SPECIES." *Molecular Ecology* 20 (6): 1176–90.

Carranza, Juan, Javier Pérez-González, Gabriel Anaya, Menno de Jong, Camilla Broggini, Frank E. Zachos, Allan D. McDevitt, et al. 2024. "Genome-Wide SNP Assessment of Contemporary European Red Deer Genetic Structure Highlights the Distinction of Peripheral Populations and the Main Admixture Zones in Europe." *Molecular Ecology*, August, e17508.

Ceballos, Francisco C., Peter K. Joshi, David W. Clark, Michèle Ramsay, and James F. Wilson. 2018. "Runs of Homozygosity: Windows into Population History and Trait Architecture." *Nature Reviews. Genetics* 19 (4): 220–34.

Chang, Christopher C., Carson C. Chow, Laurent Cam Tellier, Shashaank Vattikuti, Shaun M. Purcell, and James J. Lee. 2015. "Second-Generation PLINK: Rising to the Challenge of Larger and Richer Datasets." *GigaScience* 4 (1): 7.

Cingolani, Pablo, Adrian Platts, Le Lily Wang, Melissa Coon, Tung Nguyen, Luan Wang, Susan J. Land, Xiangyi Lu, and Douglas M. Ruden. 2012. "A Program for Annotating and Predicting the Effects of Single Nucleotide Polymorphisms, SnpEff: SNPs in the Genome of Drosophila Melanogaster Strain w1118; Iso-2; Iso-3: SNPs in the Genome of Drosophila Melanogaster Strain w1118; Iso-2; Iso-3." *Fly* 6 (2): 80–92.

Cisternas-Fuentes, Anita, and Matthew H. Koski. 2023. "Drivers of Strong Isolation and Small Effective Population Size at a Leading Range Edge of a Widespread Plant." *Heredity* 130 (6): 347–57.

Colpitts, Julie, Philip Dunstan McLoughlin, and Jocelyn Poissant. 2022. "Runs of Homozygosity in Sable Island Feral Horses Reveal the Genomic Consequences of Inbreeding and Divergence from Domestic Breeds." *BMC Genomics* 23 (1): 501.

Costa, M., C. Fernandes, J. D. S. Birks, A. C. Kitchener, M. Santos-Reis, and M. W. Bruford. 2013. "The Genetic Legacy of the 19th-Century Decline of the British Polecat: Evidence for Extensive Introgression from Feral Ferrets." *Molecular Ecology* 22 (20): 5130–47.

Costa, M., C. Fernandes, and M. Santos-Reis. 2014. "Ecology and Conservation of the Polecat Mustela Putorius (Linnaeus, 1758) in Portugal: A Review." *Munibe Monographs Nature Series* 3:79–87.

Croose, Elizabeth. 2016. "The Distribution and Status of the Polecat (Mustela Putorius) in Britain 2014-2015." https://www.vwt.org.uk/wp-content/uploads/2016/04/Polecat-Report-2016.pdf.

Croose, Elizabeth, J. W. Duckworth, Sandrine Ruette, Dmitry V. Skumatov, Vyacheslav V. Kolesnikov, and Alexander P. Saveljev. 2018. "A Review of the Status of the Western polecat*Mustela Putorius*: A Neglected and Declining Species?" *Mammalia* 82 (6): 550–64.

Cserkész, Tamás, Csaba Kiss, Zoltán Barkaszi, Tamás Görföl, Igor Zagorodniuk, Gábor Sramkó, and Gábor Csorba. 2021. "Intra- and Interspecific Morphological Variation in Sympatric and Allopatric Populations of Mustela Putorius and M. Eversmanii (Carnivora: Mustelidae) and Detection of Potential Hybrids." *Mammal Research* 66 (1): 103–14.

Danecek, Petr, Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A. DePristo, Robert E. Handsaker, et al. 2011. "The Variant Call Format and VCFtools." *Bioinformatics (Oxford, England)* 27 (15): 2156–58.

Danecek, Petr, James K. Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O. Pollard, Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools." *GigaScience* 10 (2): giab008.

Davison, A., J. D. S. Birks, H. I. Griffiths, A. C. Kitchener, D. Biggins, and R. K. Butlin. 1999. "Hybridization and the Phylogenetic Relationship between Polecats and Domestic Ferrets in Britain." *Biological Conservation* 87 (2): 155–61.

Di Marco, Moreno, Michela Pacifici, Luca Santini, Daniele Baisero, Lucilla Francucci, Gabriele Grottolo Marasini, Piero Visconti, and Carlo Rondinini. 2013. "Generation Length for Mammals." *Nature Conservation* 5 (November):89–94.

Dussex, Nicolas, Hernán E. Morales, Christine Grossen, Love Dalén, and Cock van Oosterhout. 2023. "Purging and Accumulation of Genetic Load in Conservation." *Trends in Ecology & Evolution* 38 (10): 961–69.

Dussex, Nicolas, Tom van der Valk, Hernán E. Morales, Christopher W. Wheat, David Díez-Del-Molino, Johanna von Seth, Yasmin Foster, et al. 2021. "Population Genomics of the Critically Endangered Kākāpō." *Cell Genomics* 1 (1): 100002.

Eeraerts, Maxime, Pieter Verschelde, Jan Gouwy, Sanne Van Den Berge, and Koen Van Den Berge. 2022. "Shifting Dietary Habits Linked with Contrasting Population Trends of Two Sympatric Mustelids (Martes Foina and Mustela Putorius) over 16 Years." *Global*

Ecology and Conservation 35 (e02108): e02108.
Etherington, Graham J., Adam Ciezarek, Rebecca Shaw, Johan Michaux, Elizabeth Croose,
Wilfried Haerty, and Federica Di Palma. 2022. "Extensive Genome Introgression
between Domestic Ferret and European Polecat during Population Recovery in Great
Britain." *The Journal of Heredity* 113 (5): 500–515.
Excoffier, Laurent, and Nicolas Ray. 2008. "Surfing during Population Expansions Promotes
Genetic Revolutions and Structuration." *Trends in Ecology & Evolution* 23 (7): 347–51.
Farquharson, Katherine A., Elspeth A. McLennan, Yuanyuan Cheng, Lauren Alexander,
Samantha Fox, Andrew V. Lee, Katherine Belov, and Carolyn J. Hogg. 2022. "Restoring
Faith in Conservation Action: Maintaining Wild Genetic Diversity through the Tasmanian
Devil Insurance Program." *iScience* 25 (7): 104474.
Frankham, Richard, Jonathan D. Ballou, Katherine Ralls, Mark D. B. Eldridge, Michele R.
Dudash, Charles B. Fenster, Robert C. Lacy, and Paul Sunnucks. 2017. *Loss of Genetic*
Diversity Reduces Ability to Adapt. Oxford University Press.
Frantz, Alain C., Lisette Cantú Salazar, Franz Müller, Peter Steinbach, Julian Wittische, and
Mike Heddergott. 2022. "Interactions of Cranial Helminths in the European Polecat
(Mustela Putorius): Implications for Host Body Condition." *International Journal for*
Parasitology. Parasites and Wildlife 18 (August):273–82.
Gajdárová, Barbora, Elisa Belotti, Luděk Bufka, Josefa Volfová, Sybille Wölfl, Tereza
Mináriková, Laura Hollerbach, et al. 2023. "Long-Term Genetic Monitoring of a
Reintroduced Eurasian Lynx Population Does Not Indicate an Ongoing Loss of Genetic
Diversity." *Global Ecology and Conservation* 42 (e02399): e02399.
Graciá, Eva, Francisco Botella, José Daniel Anadón, Pim Edelaar, D. James Harris, and
Andrés Giménez. 2013. "Surfing in Tortoises? Empirical Signs of Genetic Structuring
Owing to Range Expansion." *Biology Letters* 9 (3): 20121091.
Grossen, Christine, Frédéric Guillaume, Lukas F. Keller, and Daniel Croll. 2020. "Purging of
Highly Deleterious Mutations through Severe Bottlenecks in Alpine Ibex." *Nature*
Communications 11 (1): 1001.
Gustafson, Kyle D., Michelle G. Hawkins, Tracy L. Drazenovich, Robert Church, Susan A.
Brown, and Holly B. Ernest. 2018. "Founder Events, Isolation, and Inbreeding:
Intercontinental Genetic Structure of the Domestic Ferret." *Evolutionary Applications* 11
(5): 694–704.
Harris, Stephen, Pat Morris, Stephanie Wray, and Derek Yalden. 1995. "A Review of British
Mammals: Population Estimates and Conservation Status of British Mammals Other
Than Cetaceans," January.
https://www.researchgate.net/publication/242583185_A_Review_of_British_Mammals_
Population_Estimates_and_Conservation_Status_of_British_Mammals_Other_Than_C
etaceans.
Hedrick, Philip W. 2013. "Adaptive Introgression in Animals: Examples and Comparison to
New Mutation and Standing Variation as Sources of Adaptive Variation." *Molecular*
Ecology 22 (18): 4606–18.
Hedrick, P. W., J. A. Robinson, R. O. Peterson, and J. A. Vucetich. 2019. "Genetics and
Extinction and the Example of Isle Royale Wolves." *Animal Conservation* 22 (3): 302–9.
Hickey, Glenn, Benedict Paten, Dent Earl, Daniel Zerbino, and David Haussler. 2013. "HAL:
A Hierarchical Format for Storing and Analyzing Multiple Genome Alignments."
Bioinformatics (Oxford, England) 29 (10): 1341–42.
Hijmans, R. 2024. "Geosphere: Spherical Trigonometry. R Package Version 1.5-20,
Https://github.com/rspatial/geosphere."
Hofmeester, Tim R., Vilmar Dijkstra, Jasja J. A. Dekker, Tom van der Meij, and Sim
Broekhuizen. 2019. "The Status of the Dutch Polecat Population: Correction of a
Recently Published Error." *Mammalia* 83 (5): 453–54.
Howard-McCombe, Jo, Alexandra Jamieson, Alberto Carmagnini, Isa-Rita M. Russo,
Muhammad Ghazali, Ruairidh Campbell, Carlos Driscoll, et al. 2023. "Genetic
Swamping of the Critically Endangered Scottish Wildcat Was Recent and Accelerated
by Disease." *Current Biology: CB* 33 (21): 4761–69.e5.

Jarausch, Anne, Verena Harms, Gesa Kluth, Ilka Reinhardt, and Carsten Nowak. 2021.
"How the West Was Won: Genetic Reconstruction of Rapid Wolf Recolonization into
Germany's Anthropogenic Landscapes." *Heredity* 127 (1): 92–106.
Jong, Menno J. de, Zhipeng Li, Yanli Qin, Erwan Quéméré, Karis Baker, Wen Wang, and A.
Rus Hoelzel. 2020. "Demography and Adaptation Promoting Evolutionary Transitions in
a Mammalian Genus That Diversified during the Pleistocene." *Molecular Ecology* 29
(15): 2777–92.
Khan, Anubhab, Kaushalkumar Patel, Harsh Shukla, Ashwin Viswanathan, Tom van der
Valk, Udayan Borthakur, Parag Nigam, et al. 2021. "Genomic Evidence for Inbreeding
Depression and Purging of Deleterious Genetic Variation in Indian Tigers." *Proceedings*
of the National Academy of Sciences of the United States of America 118 (49).
https://doi.org/10.1073/pnas.2023018118.
Kitchener, Andrew C., and J. D. S. Birks. 2008. "Feral Ferret Mustela Furo."
https://scholar.google.com/citations?user=b6MOxGsAAAAJ&hl=en&oi=sra.
Kliver, Sergei, Marlys L. Houck, Polina L. Perelman, Azamat Totikov, Andrey Tomarovsky,
Olga Dudchenko, Arina D. Omer, et al. 2023. "Chromosome-Length Genome Assembly
and Karyotype of the Endangered Black-Footed Ferret (Mustela Nigripes)." *The Journal*
of Heredity 114 (5): 539–48.
Koepfli, Klaus-Peter, Kerry A. Deere, Graham J. Slater, Colleen Begg, Keith Begg, Lon
Grassman, Mauro Lucherini, Geraldine Veron, and Robert K. Wayne. 2008. "Multigene
Phylogeny of the Mustelidae: Resolving Relationships, Tempo and Biogeographic
History of a Mammalian Adaptive Radiation." *BMC Biology* 6 (1): 10.
Kominos, T., and A. Galanaki. 2018. "Western Polecat Mustela Putorius Distribution in
Greece." *IUCN Small Carniv. Conserv* 57:5–7.
Lande, Russell. 1994. "Risk of Population Extinction from Fixation of New Deleterious
Mutations." *Evolution; International Journal of Organic Evolution* 48 (5): 1460–69.
Langley, P. J. W., and D. W. Yalden. 1977. "The Decline of the Rarer Carnivores in Great
Britain during the Nineteenth Century." *Mammal Review* 7 (3-4): 95–116.
Li, Heng, and Richard Durbin. 2009. "Fast and Accurate Short Read Alignment with
Burrows-Wheeler Transform." *Bioinformatics (Oxford, England)* 25 (14): 1754–60.
Lou, Runyang Nicolas, and Nina Overgaard Therkildsen. 2022. "Batch Effects in Population
Genomic Studies with Low-Coverage Whole Genome Sequencing Data: Causes,
Detection and Mitigation." *Molecular Ecology Resources* 22 (5): 1678–92.
Manichaikul, Ani, Josyf C. Mychaleckyj, Stephen S. Rich, Kathy Daly, Michèle Sale, and
Wei-Min Chen. 2010. "Robust Relationship Inference in Genome-Wide Association
Studies." *Bioinformatics (Oxford, England)* 26 (22): 2867–73.
Manunza, A., M. Amills, A. Noce, B. Cabrera, A. Zidi, S. Eghbalsaied, E. Carrillo de
Albornoz, et al. 2016. "Romanian Wild Boars and Mangalitza Pigs Have a European
Ancestry and Harbour Genetic Signatures Compatible with Past Population
Bottlenecks." *Scientific Reports* 6 (1): 29913.
Maran, T., D. Skumatov, A. V. Abramov, and A. Kranz. 2016.
https://www.researchgate.net/profile/Alexei-
Abramov/publication/308778184_Mustela_eversmanii/links/57ef8f5208ae886b897417c
4/Mustela-eversmanii.pdf.
Marneweck, C., A. R. Butler, L. C. Gigliotti, S. N. Harris, A. J. Jensen, M. Muthersbaugh, B.
A. Newman, et al. 2021. "Shining the Spotlight on Small Mammalian Carnivores: Global
Status and Threats." *Biological Conservation* 255 (109005): 109005.
Marneweck, Courtney J., Benjamin L. Allen, Andrew R. Butler, Emmanuel Do Linh San,
Stephen N. Harris, Alex J. Jensen, Elizabeth A. Saldo, et al. 2022. "Middle☐out
Ecology: Small Carnivores as Sentinels of Global Change." *Mammal Review* 52 (4):
471–79.
Martínez-Cruz, Begoña, Hanna Zalewska, and Andrzej Zalewski. 2022. "The Genetic
Diversity and Structure in the European Polecat Were Not Affected by the Introduction
of the American Mink in Poland." *PloS One* 17 (9): e0266161.
Mathews, Fiona, Laura Kubasiewicz, John Gurnell, Colin Harrower, Robbie Mcdonald, and

Richard Shore. 2018. "A Review of the Population and Conservation Status of British
Mammals." Unpublished. https://doi.org/10.13140/RG.2.2.28059.31521.
McDevitt, Allan D., Ilaria Coscia, Samuel S. Browett, Aritz Ruiz-González, Mark J. Statham,
Iwona Ruczyńska, Liam Roberts, et al. 2022. "Next-Generation Phylogeography
Resolves Post-Glacial Colonization Patterns in a Widespread Carnivore, the Red Fox
(Vulpes Vulpes), in Europe." *Molecular Ecology* 31 (3): 993–1006.
McKenna, Aaron, Matthew Hanna, Eric Banks, Andrey Sivachenko, Kristian Cibulskis,
Andrew Kernytsky, Kiran Garimella, et al. 2010. "The Genome Analysis Toolkit: A
MapReduce Framework for Analyzing next-Generation DNA Sequencing Data."
Genome Research 20 (9): 1297–1303.
Mestre, F. M., J. P. Ferreira, and A. Mira. 2007. "Modelling the Distribution of the European
Polecat Mustela Putorius in a Mediterranean Agricultural Landscape." *Revue D'écologie*
62 (1): 35–47.
Montgomery, W. Ian, Jim Provan, A. Marshal McCabe, and Derek W. Yalden. 2014. "Origin
of British and Irish Mammals: Disparate Post-Glacial Colonisation and Species
Introductions." *Quaternary Science Reviews* 98 (August):144–65.
Moroni, Barbara, Alice Brambilla, Luca Rossi, Pier Giuseppe Meneguz, Bruno Bassano, and
Paolo Tizzani. 2022. "Hybridization between Alpine Ibex and Domestic Goat in the Alps:
A Sporadic and Localized Phenomenon?" *Animals: An Open Access Journal from MDPI*
12 (6): 751.
Münger, Xenia, Mathieu Robin, Love Dalén, and Christine Grossen. 2024. "Facilitated
Introgression from Domestic Goat into Alpine Ibex at Immune Loci." *Molecular Ecology*
33 (14): e17429.
Novo, Irene, Pilar Ordás, Natalia Moraga, Enrique Santiago, Humberto Quesada, and
Armando Caballero. 2023. "Impact of Population Structure in the Estimation of Recent
Historical Effective Population Size by the Software GONE." *Genetics, Selection,*
Evolution: GSE 55 (1): 86.
Peng, Xinxia, Jessica Alföldi, Kevin Gori, Amie J. Eisfeld, Scott R. Tyler, Jennifer Tisoncik-
Go, David Brawand, et al. 2014. "The Draft Genome Sequence of the Ferret (Mustela
Putorius Furo) Facilitates Study of Human Respiratory Disease." *Nature Biotechnology*
32 (12): 1250–55.
Perez-Sepulveda, Blanca M., Darren Heavens, Caisey V. Pulford, Alexander V. Predeus,
Ross Low, Hermione Webster, Gregory F. Dykes, et al. 2021. "An Accessible, Efficient
and Global Approach for the Large-Scale Sequencing of Bacterial Genomes." *Genome*
Biology 22 (1): 349.
"Picard Toolkit." 2019. *Broad Institute, GitHub Repository*. Broad Institute.
https://broadinstitute.github.io/picard/.
Plessis, Sarah J. du, Mark Blaxter, Klaus-Peter Koepfli, Elizabeth A. Chadwick, and Frank
Hailer. 2023. "Genomics Reveals Complex Population History and Unexpected Diversity
of Eurasian Otters (Lutra Lutra) in Britain Relative to Genetic Methods." *Molecular*
Biology and Evolution 40 (11). https://doi.org/10.1093/molbev/msad207.
Prugh, Laura R., Calum X. Cunningham, Rebecca M. Windell, Brian N. Kertson, Taylor R.
Ganz, Savanah L. Walker, and Aaron J. Wirsing. 2023. "Fear of Large Carnivores
Amplifies Human-Caused Mortality for Mesopredators." *Science (New York, N.Y.)* 380
(6646): 754–58.
Purcell, Shaun, Benjamin Neale, Kathe Todd-Brown, Lori Thomas, Manuel A. R. Ferreira,
David Bender, Julian Maller, et al. 2007. "PLINK: A Tool Set for Whole-Genome
Association and Population-Based Linkage Analyses." *The American Journal of Human*
Genetics 81 (3): 559–75.
R Core Team. 2024. "A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing.https://www.R-Project.org/."
Robinson, Jacqueline, Christopher C. Kyriazis, Stella C. Yuan, and Kirk E. Lohmueller. 2023.
"Deleterious Variation in Natural Populations and Implications for Conservation
Genetics." *Annual Review of Animal Biosciences* 11 (February):93–114.
Sainsbury, Katherine A., Richard F. Shore, Henry Schofield, Elizabeth Croose, Ruairidh D.

Campbell, and Robbie A. Mcdonald. 2019. "Recent History, Current Status,
Conservation and Management of Native Mammalian Carnivore Species in Great
Britain." *Mammal Review* 49 (2): 171–88.
Sainsbury, Katherine A., Richard F. Shore, Henry Schofield, Elizabeth Croose, Georg
Hantke, Andrew C. Kitchener, and Robbie A. McDonald. 2020. "Diets of European
Polecat Mustela Putorius in Great Britain during Fifty Years of Population Recovery."
Mammal Research 65 (2): 181–90.
Santiago, Enrique, Irene Novo, Antonio F. Pardiñas, María Saura, Jinliang Wang, and
Armando Caballero. 2020. "Recent Demographic History Inferred by High-Resolution
Analysis of Linkage Disequilibrium." *Molecular Biology and Evolution* 37 (12): 3642–53.
Saremi, Nedda F., Megan A. Supple, Ashley Byrne, James A. Cahill, Luiz Lehmann
Coutinho, Love Dalén, Henrique V. Figueiró, et al. 2019. "Puma Genomes from North
and South America Provide Insights into the Genomic Consequences of Inbreeding."
Nature Communications 10 (1): 4769.
S., Birks J. D. 2008. "The Polecat Survey of Britain 2004-2006: A Report on the Polecats
Distribution, Status and Conservation."
Schluter, Dolph. 2000. *The Ecology of Adaptive Radiation*. Oxford Series in Ecology and
Evolution. London, England: Oxford University Press.
Sendell-Price, Ashley T., Kristen C. Ruegg, and Sonya M. Clegg. 2020. "Rapid
Morphological Divergence Following a Human-Mediated Introduction: The Role of Drift
and Directional Selection." *Heredity* 124 (4): 535–49.
Senn, Helen V., Muhammad Ghazali, Jennifer Kaden, David Barclay, Ben Harrower,
Ruairidh D. Campbell, David W. Macdonald, and Andrew C. Kitchener. 2019.
"Distinguishing the Victim from the Threat: SNP-Based Methods Reveal the Extent of
Introgressive Hybridization between Wildcats and Domestic Cats in Scotland and Inform
Future in Situ and Ex Situ Management Options for Species Restoration." *Evolutionary
Applications* 12 (3): 399–414.
Siepel, Adam, and David Haussler. 2004. "Phylogenetic Estimation of Context-Dependent
Substitution Rates by Maximum Likelihood." *Molecular Biology and Evolution* 21 (3):
468–88.
Speak, Samuel A., Thomas Birley, Chiara Bortoluzzi, Matthew D. Clark, Lawrence Percival-
Alwyn, Hernán E. Morales, and Cock van Oosterhout. 2024. "Genomics-Informed
Captive Breeding Can Reduce Inbreeding Depression and the Genetic Load in Zoo
Populations." *Molecular Ecology Resources* 24 (7): e13967.
Spielman, Derek, Barry W. Brook, and Richard Frankham. 2004. "Most Species Are Not
Driven to Extinction before Genetic Factors Impact Them." *Proceedings of the National
Academy of Sciences of the United States of America* 101 (42): 15261–64.
Stamatakis, Alexandros. 2014. "RAxML Version 8: A Tool for Phylogenetic Analysis and
Post-Analysis of Large Phylogenies." *Bioinformatics (Oxford, England)* 30 (9): 1312–13.
Stroupe, Sam, David Forgacs, Andrew Harris, James N. Derr, and Brian W. Davis. 2022.
"Genomic Evaluation of Hybridization in Historic and Modern North American Bison
(Bison Bison)." *Scientific Reports* 12 (1): 6397.
Sumption, K. J., and J. R. Flowerdew. 1985. "The Ecological Effects of the Decline in
Rabbits *(Oryctolagus Cuniculus L.)* due to Myxomatosis." *Mammal Review* 15 (4): 151–
86.
Szapu, Julianna Szulamit, Tamás Cserkész, Zsolt Pirger, Csaba Kiss, and József Lanszki.
2024. "Exposure to Anticoagulant Rodenticides in Steppe Polecat (Mustela Eversmanii)
and European Polecat (Mustela Putorius) in Central Europe." *The Science of the Total
Environment* 948 (174282): 174282.
Szatmári, Lajos, Tamás Cserkész, Levente Laczkó, József Lanszki, Cino Pertoldi, Alexei V.
Abramov, Morten Elmeros, Barnabás Ottlecz, Zsolt Hegyeli, and Gábor Sramkó. 2021.
"A Comparison of Microsatellites and Genome-Wide SNPs for the Detection of
Admixture Brings the First Molecular Evidence for Hybridization between Mustela
Eversmanii and M. Putorius (Mustelidae, Carnivora)." *Evolutionary Applications* 14 (9):
2286–2304.

Szewczyk, Maciej, Sabina Nowak, Natalia Niedźwiecka, Pavel Hulva, Renata Špinkytė-Bačkaitienė, Klára Demjanovičová, Barbora Černá Bolfíková, et al. 2019. "Dynamic Range Expansion Leads to Establishment of a New, Genetically Distinct Wolf Population in Central Europe." *Scientific Reports* 9 (1): 19003.

Takach, Brenton von, Skye F. Cameron, Teigan Cremona, Mark D. B. Eldridge, Diana O. Fisher, Rosemary Hohnen, Chris J. Jolly, et al. 2024. "Conservation Prioritisation of Genomic Diversity to Inform Management of a Declining Mammal Species." *Biological Conservation* 291 (110467): 110467.

Thomas, Nia E., Frank Hailer, Michael W. Bruford, and Elizabeth A. Chadwick. 2022. "Country-Wide Genetic Monitoring over 21 Years Reveals Lag in Genetic Recovery despite Spatial Connectivity in an Expanding Carnivore (Eurasian Otter, Lutra Lutra) Population." *Evolutionary Applications* 15 (12): 2125–41.

Valk, Tom van der, David Díez-Del-Molino, Tomas Marques-Bonet, Katerina Guschanski, and Love Dalén. 2019. "Historical Genomes Reveal the Genomic Consequences of Recent Population Decline in Eastern Gorillas." *Current Biology: CB* 29 (1): 165–70.e6.

Virgós, Emilio. 2003. "Association of the Polecat Mustela Putorius in Eastern Spain with Montane Pine Forests." *Oryx* 37 (4): 484–87.

Welles, Shana R., and Katrina M. Dlugosch. 2018. "Population Genomics of Colonization and Invasion." In *Population Genomics*, 655–83. Cham: Springer International Publishing.

Willcox, D. 2020. "Conservation Status,ex Situpriorities and Emerging Threats to Small Carnivores." *The International Zoo Yearbook* 54 (1): 19–34.

Author Contribution

The language used to recognise author contributions is described by CRediT Taxonomy (https://credit.niso.org/).

Conceptualization: GE, WH; formal analysis: RS; funding acquisition: GE, WH, FDP; investigation: RS; methodology: RS, GE, WH; project administration: RS, GE, WH; resources: JM, ACK, GE, WH; software: RS; supervision: JM, GE, WH; visualisation: RS; writing - original draft: RS; writing review & editing: RS, JM, ACK, GE, WH

Conflict of Interest

The authors declare no conflict of interest.

Data Availability

All raw reads for the sequences generated during this study have been uploaded to ENA with the project number PRJEB76468. Bioinformatic scripts used to carry out analyses can be found at this repository (https://github.com/TGAC/EuropeanPolecatPopGen/).

1076

Supplementary material

1077

1078   Supplementary methods, figures and tables are available in the links provided.

1079

1080   **Figures**

1081
1082

Figure 1. (a) The range of the European polecat in Britain in 1915. Dark green areas
indicate the population stronghold, light green indicates known occurrences, and "?" refers to
uncertain occurrences of polecats in those areas (Langley and Yalden 1977). (b) Records
received during the Vincent Wildlife Trust 2014-15 survey of verifiable sightings, based on
their phenotype, of true polecats (dark-green), polecat-ferret hybrids (yellow), both true
polecats and hybrids (lime green), and unverifiable (grey) (Croose 2016). (c) Samples
selected to be sequenced in this study (dark yellow) and (red) samples that were previously
sequenced by (Etherington et al. 2022). Figures 1(a) and 1(b) were adapted from Croose,
2016.
1092

1093

1094

1095
1096
Figure 2. Population structure of the European polecat in Great Britain and across mainland
Europe. (a) PCA based on whole genome SNPs for all European polecats and domestic
ferrets. (b) Admixture results for whole-genome SNPs between the four groups (Domestic
ferret, English, Welsh and mainland European polecats). (c) Isolation by distance of samples
plotted using a pairwise Hamming genetic distance and geographic distance matrix (d) Map
of Great Britain demonstrating longitude coordinates of samples and PCA based on whole
genome SNPs for all British polecats, which is supported by a significant correlation between
PC1 and longitude coordinates.

1105

1106   **Figure 3.** Genomic diversity estimates and comparisons across British and European
1107   populations (a) Shared and private SNPs between populations. (b) Proportion of
1108   heterozygous sites calculated using VCFtools (c) Population nucleotide diversity values (π),
1109   calculated in 20Mb windows across the genome and (d) Population-level inbreeding
1110   coefficient values
1111
1112
1113

1114
1115   **Figure 4.** Population estimates of higher coverage individuals (mean depth >6x) and (a) the
1116   calculated proportion of the genome in ROH (F_{ROH}) and (b) the distribution of ROH across
1117   different length categories. Data points represent the percentage of an individual's total ROH
1118   that falls within each length category.
1119

1120
1121

1122
1123
1124 **Figure 5.** Mutation load landscape across populations within different functional categories.
1125 Loss of function (LoF) (a) homozygous mutational load per population and (b) heterozygous
1126 mutational load and missense (c) homozygous mutational load and (d) heterozygous
1127 mutational load.
1128
1129

1130
1131
1132 **Figure 6.** Estimated mean effective population size (solid line) with 95% confidence intervals
1133 (shaded area) for English (green) and Welsh (orange) populations across 200 generations
1134 (generation time = 4 years) based on whole genome SNP data computed in GONE.
1135