Consistent hierarchies of single-neuron timescales in mice, macaques and humans

Authors: Zachary R. Zeisler[1], Marques Love[1], Ueli Rutishauser[2,3,4,5], Frederic M. Stoll[1†], and Peter H. Rudebeck[1†*]

Author affiliations:

1. Nash Family Department of Neuroscience, Lipschultz Center for Cognitive Neuroscience, and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York, NY 10029
2. Department of Neurosurgery, Cedars-Sinai Medical Center, Los Angeles, CA 90048
3. Department of Neurology, Cedars-Sinai Medical Center, Los Angeles, CA 90048
4. Center for Neural Science and Medicine, Department of Biological Sciences, Cedars-Sinai Medical Center, Los Angeles, CA 90048
5. Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125

[†] Denotes equal contribution

* Corresponding author

Correspondence should be addressed to:

Dr. Peter Rudebeck – peter.rudebeck@mssm.edu

Lipschultz Center for Cognitive Neuroscience, Nash Family Department of Neuroscience and Friedman Brain Institute

Icahn School of Medicine at Mount Sinai

One Gustave L. Levy Place

New York, NY 10029

Author contributions: ZRZ, FMS, and PHR conceived the study. ZRZ and FMS performed the analyses. FMS collected the macaque ventral frontal cortex dataset, and FMS and ML localized the macaque ventral frontal cortex recordings to cytoarchitectonic regions. UR provided access to the human datasets and key feedback on analysis approaches throughout. ZRZ, FMS, and PHR wrote the paper. All authors edited and approved the final version of the paper.

Acknowledgements: The authors thank the individuals who publicly provided their data for our use in these analyses: Tomas Aquino, Nand Chandravadia, and Juri Minxha; Vincent Fontanier and Emmanuel Procyk; Sylvia Wirth and Jean-Rene Duhamel; Megan Young and Clayton Mosher; and Nick Steinmetz. ZRZ, ML, FMS and PHR are supported by grants from NIMH and the BRAIN Initiative (R01MH110822, R01MH132064). UR is supported by the BRAIN Initiative (U01NS117839).

Figures: 3 (plus 1 supplemental)

Tables: 2

Abstract

The intrinsic timescales of single neurons are thought to be hierarchically organized across the cortex. This conclusion, however, is primarily based on analyses of neural responses from macaques. Whether hierarchical variation in timescales is a general brain organizing principle across mammals remains unclear. Here we took a cross-species approach and estimated neuronal timescales of thousands of single neurons recorded across multiple areas in mice, monkeys, and humans using a task-agnostic method. We identify largely consistent hierarchies of timescales in frontal and limbic regions across species: hippocampus had the shortest timescale whereas anterior cingulate cortex had the longest. Within this scheme, variability across species was found, most notably in amygdala and orbitofrontal cortex. We show that variation in timescales is not simply related to differences in spiking statistics nor the result of cytoarchitectonic features such as cortical granularity. Thus, hierarchically organized timescales are a consistent organizing principle across species and appear to be related to a combination of intrinsic and extrinsic factors.

Highlights

- Similar hierarchies of single neuron timescales exist across humans, macaques, and mice
- Humans diverge from an otherwise conserved hierarchy in amygdala and OFC
- Timescale variability in ventral frontal cortex is not entirely related to cytoarchitecture

Introduction:

Receptive fields are a foundational concept in psychophysics and neuroscience, a classic example being the spatial receptive fields of single neurons in primary visual cortex (Hubel and Wiesel, 1962; Spillmann, 2014) and higher-order parietal and frontal areas (Viswanathan and Nieder, 2017). An influential line of work has extended this framework to the temporal domain (Murray et al., 2014). Using the rate of exponential decay of the spike-count autocorrelation, Murray and colleagues extracted the intrinsic temporal receptive field, or timescale, of a single neuron or population of neurons. These neural timescales, which are hypothesized to reflect the amount of time over which a neuron integrates information, appear to be hierarchically organized across the brain (Murray et al., 2014; Nougaret et al., 2021). Neurons within sensory cortex have shorter timescales, while neurons within higher-order associative areas, particularly in frontal cortex, have longer timescales. Convergent evidence from fMRI in humans (Hasson et al., 2008; Honey et al., 2012; Raut et al., 2020), fMRI and neurophysiology recordings in non-human primates (Murray et al., 2014; Manea et al., 2022, 2024), and modelling work (Gao et al., 2020; Perez-Nieves et al., 2021; Fontanier et al., 2022; Li and Wang, 2022) suggest that this hierarchical organization of timescales may be a fundamental principle of brain organization. Despite this, analyses of single neuron timescales across humans and animal models utilizing the same data modality and fitting procedure have been lacking. This is a key consideration given the differences in temporal resolution between fMRI and single neuron activity.

Neural timescales are hypothesized to be driven by the physical properties as well as connectivity of single neurons (Soltani et al., 2021). The divergence in brain anatomy across humans, non-human primates, and rodents suggests that there could be corresponding changes in timescales across brain areas that mirror anatomical differences (Cavanagh et al., 2020). For instance, the dorsal portion of anterior cingulate cortex (ACC) is largely conserved across humans, macaques and mice as it is agranular cortex (Wise, 2008; van Heukelum et al., 2020). While the timescales of ACC neurons across primate and rodent species have not been directly compared, their shared anatomy suggests that they may be similar. By contrast,

amygdala, an area which sends strong projections to ACC and other frontal cortex areas

(Ghashghaei et al., 2007; Hintiryan et al., 2021; Zeisler et al., 2023), is markedly different across

humans, macaques and mice – particularly in the basolateral subdivision (Chareyron et al.,

2011; Zeisler et al., 2024). Whether these differences in amygdala anatomy are mirrored by

differences in timescale is unknown.

To address these issues, we compared the timescale of tens of thousands of neurons

within frontal cortex, amygdala, and hippocampus of mice, macaque monkeys, and humans.

We identified largely consistent hierarchies across species within this amygdala-hippocampus-

frontal cortex system, with key exceptions particularly in amygdala. Importantly, divergences

from the common hierarchy were not related to known differences in frontal cortex

cytoarchitecture.

Results

We analyzed the firing activity of 17,458 single neurons recorded in orbitofrontal cortex

(OFC), medial prefrontal cortex (mPFC), anterior cingulate cortex (ACC), amygdala and

hippocampus in three species: humans, macaque monkeys, and mice (**Figure 1A**, **Table 1,** see

Methods for detailed explanation on anatomical mapping). Multiple methods have been put

forth to calculate the temporal timescales of single neurons. Timescale estimation originally

relied only on the spiking activity during periods in which animals were not actively performing

a task but rather sitting in restful attentiveness (Murray et al., 2014). While well-suited for

analysis of data recorded from non-human primates, who often perform tasks with rigid

temporal structure, such an approach is not optimal for recordings from rodents or humans,

where tasks are often performed with more continuous and variable time courses. One method

based on the distribution of inter-spike intervals (ISI) allows for the estimation of timescales

irrespective of task events and could therefore be readily applied to the activity of neurons

recorded in different species (Fontanier et al., 2022). Importantly, the estimated timescales

from the approach that uses the ISI and the more standard approach (Murray et al., 2014) have

been shown to be highly correlated with one another, suggesting that they are modeling the

same underlying physiological properties.

To estimate the intrinsic timescale of each of the neurons we analyzed, we employed

this latter ISI-based method from Fontanier, Sarazin and colleagues (Fontanier et al., 2022)**.** In

brief, we fit an exponential decay function to the descending portion of the distribution of ISIs

according to the following equation:

82
$$R(k\Delta) = A\left[e^{-\left(\frac{k\Delta}{\tau}\right)} + B\right]$$

where $k\Delta$ is the time lag between bins, τ represents the neuron's *timescale*, describing the rate

of decay of the autocorrelation distribution, and A and B are free parameters which together

describe the peak and asymptotic limit of the autocorrelation. We also extracted the time bin at

which the autocorrelation begins decreasing, which we refer to as *latency*. Only neurons for

which the model fit well ($R^2 > 0.5$) and exhibiting timescales within the range $10 < \tau < 1000$ were

included for further analysis (n=13,755 single neurons, 78.7% success; **Figure 1B**).

Mice, macaques, and humans exhibited largely similar hierarchies of single neuron

timescales and latencies across areas (**Figure 1C, D**). Across all three species, ACC neurons

showed the longest timescales and latencies while hippocampus neurons showed the shortest

timescales. Other areas were characterized by greater across-species variability. Notably,

human amygdala neurons had a significantly shorter timescale compared to macaques and

mice (**Figure 1E**; middle). All three species also had different amygdala latencies. Furthermore,

the timescales of single neurons in OFC were statistically distinguishable in all three species,

with humans exhibiting the longest timescale. Taken together, we found numerous similarities

between the timescale and latency hierarchies across these three species, with a notable

exception for OFC and amygdala neurons in humans.

Figure 1. Single neuron timescale and latency hierarchies are largely conserved across species.
A) Schematic representing the approximate position of brain areas from which recordings were
collected in mice (left), macaques (middle) and humans (right); ACC in pink, amygdala in purple,
hippocampus in cyan, mPFC in light blue, OFC in orange (see **Methods** for detailed explanation of
area assignments). B) Number of neurons considered (light colors) and successfully fit (dark
colors) across all brain areas. C) Average [±95% CI] timescales and D) latencies for each area in
humans (red), macaques (green) and mice (blue). There was a main effect of both species ($F_{(2)}$ =
48.37, p = 1.2*10^{-21}) and area ($F_{(4)}$ = 422.15, p = 0) on timescale as well as a significant
interaction ($F_{(8)}$ = 31.53, p = 1.7*10^{-49}). All three species had different OFC timescales (OLS
regression, mouse vs monkey: $t_{(2)}$ = 1.961, p = 0.05; monkey vs human $t_{(2)}$ = 7.332, p = 2.6*10^{-13}), OFC latency (mouse vs monkey: $t_{(2)}$ = 5.672, p = 1.5*10^{-8}; monkey vs human $t_{(2)}$ = 4.788, p =
1.7*10^{-6}), and amygdala latency (mouse vs monkey: $t_{(2)}$ = 2.096, p = 0.036; monkey vs human
$t_{(2)}$ = 6.379, p = 2.1 *10^{-10}). Human amygdala timescale was different from the other species
(mouse vs monkey $t_{(2)}$ = 0.338, p = 0.735; monkey vs human $t_{(2)}$ = 13.331, p = 2.6*10^{-39}). E)
Average fits for neurons from each area and species. Lines are plotted by the mean fit
parameters from successfully fit neurons, with shading indicating the standard error.

While the above analyses focused on the average timescales for each area and species,

timescales were not necessarily uniform within a given area. Indeed, we found large variability

in the distributions of timescale and latency within each area and across all three species

(**Figure 2 A, C-D)**. In fact, both timescale and latency distributions across most considered areas

and species had unequal variances. This within-area variation is particularly prominent in ACC,

where mice, macaques, and humans all had a bimodal distribution of timescale: a small

population of neurons with short timescale and a larger population of neurons with longer

timescales. Such consistent heterogeneity across species within an area potentially provides

insight into the organizing role that timescales might play in the temporal processing of

information. Indeed, it could be the case that the distribution of timescales within an area is

more critical for supporting that area's function than the average timescale.

Across the different areas and species, there was also significant variability in the

latency measure that was fitted to each neuron's spiking activity (**Figure 1D**). Here ACC and

mPFC had the longest latency whereas hippocampus generally had the shortest. Further, within

each area there was also a high degree of variability in the latency (**Figure 2D**). Importantly,

while timescale and latency exhibit some degree of correlation (**Figure 2B**), each explains only

half of the variance in the other. This point highlights that latency is a factor that is partially

separable from the timescale and likely contributes to how information is temporally processed

within a brain area.

One potential issue with fitting exponential decay functions to the activity of single

neurons is that they might be influenced solely by the firing rates of the neurons. Thus, we next

checked that our timescale and latency estimates reflected distinct neuronal functional

properties, independent from the influence of firing rate. To that end, we built regression

models to predict timescales based on spiking statistics, areas and species (**Supplemental**

Figure 1). We found that the number of spikes entered into the analysis only had a significant

influence on a neurons' timescale while the number of spikes, mean firing rate, and Fano factor

all influenced the neural latencies. These parameters, however, offered little additional

explanatory power compared to species and area, explaining significantly less variance than

area and species alone. Thus, the neural timescale and latency that we reported above (**Figures**

1, 2) represent features of neuronal dynamics that are distinct from classical spiking statistics.

Figure 2: Distributions of timescale and latency for individual neurons. A) Two example single-neuron fits from macaque OFC with relatively short (left; light green) and long (right; dark green) timescales. Grey traces indicate normalized ISI distribution, the black line is the smoothed autocorrelation, and the green lines are the fit autocorrelation function, specified by timescale (τ) and latency (*lat*) shown in the top right of each panel. B) Density plot of timescale and latency, with regression line overlaid. Timescale and latency are correlated with each other (OLS regression: R^2 = 0.501, p < 0.001). C) Density histograms showing distribution of timescales across neurons recorded within each brain area. The timescales of neurons from OFC, amygdala, and mPFC all have unequal variance across species (Brown-Forsythe test: p < 0.004). Green arrows indicate the position of the example cells shown in panel A. D) Same as C for latencies (bottom). All areas with the exception of mPFC had unequal variance regarding latencies (p < 10^{-6}).

The relationship between cytoarchitecture and timescales within ventral frontal cortex

As previously noted, ACC in humans, macaques, and mice is agranular cortex (van

Heukelum et al., 2020). By contrast, OFC in macaques (**Figure 3A**) and humans is composed of

granular, dysgranular, and agranular cortex (Carmichael and Price, 1994; Wise, 2008). OFC in

mice, by contrast, is entirely agranular. One possibility is that these differences in granularity

might underlie the timescale and latency differences we observed in OFC across species (**Figure**

1). To assess this, we concentrated our analysis on recordings from 12,043 single neurons in

macaque ventral frontal cortex, of which 5,230 were included in the previous analyses of

macaque OFC in **Figures 1** and **2**. This dataset included the activity of single neurons recorded

using 157-channel semi-chronic microdrives in two monkeys, where each electrode trajectory

was precisely reconstructed (Stoll and Rudebeck, 2024b, 2024c). Using standard histological

techniques, the ventral frontal cortex was subdivided into 8 cytoarchitectonic areas based on

analysis of Nissl-, SMI-32- and calbindin-stained tissue series (**Figure 3A**). This allowed us to

pinpoint the precise location of recorded neurons according to previous parcellations of the

ventral frontal cortex of macaques (Carmichael and Price, 1994; Rapan et al., 2023). Thus, we

could compare the timescales of neurons from neighboring subdivisions with distinct cortical

granularity; fully granular areas 11m/l, 12l, 12m, and 12o, dysgranular areas 12r, 13m, and 13l

areas, and agranular insula (AI; the number of neurons in each subdivision is shown in **Figure 3B**

and **Table 2**).

We found that timescale variability across macaque ventral frontal cortex could not be

fully explained by differences in granularity (**Figure 3C,** left). While there was a main effect of

granularity, such that neurons in granular areas tended to exhibit longer timescales than those

in dysgranular areas, hierarchical model comparison indicated that granularity did not

significantly improve a model predicting neural timescales based on area alone. Notably,

adjacent areas 12m and 12r were most different from other granular and dysgranular areas,

respectively as they had relatively longer timescales. Neuronal latencies were found to have a

different relationship to granularity (**Figure 3D**, right). Specifically, granular and agranular areas

demonstrated longer latencies than dysgranular – and a model predicting latency with

granularity performed better than a model containing area alone. Thus, the presence or

absence of an internal granule cell layer does not appear to be a strong predictor of the

timescale of single neurons in an area. Granularity, however, does significantly affect neuronal

latency, which may be related to variation in the cortical microstructure (Gao et al., 2020).

Finally, we investigated whether single neuron timescales and latencies could be related

to the spatial location of neurons on the ventral surface of the frontal lobe. Across frontal

cortex there are both antero-posterior (AP) and medio-lateral (ML) gradients in the connections

from other parts of the brain, especially the amygdala (Ghashghaei et al., 2007) and

hippocampus (Aggleton et al., 2015). We hypothesized that such differences in long-range

inputs could drive differences in neural timescales. To this end, we assessed whether the AP

and ML coordinates of each recorded neuron influenced neuronal timescales and latencies. In

both animals, there were main effects of the AP position ($F(1) = 20.16$, $p = 7.3*10^{-6}$), ML

position ($F(1) = 26.97$, $p = 2.1*10^{-7}$), and a significant interaction ($F(1) = 47.12$, $p = 7.5*10^{-12}$),

such that timescales were longest in neurons recorded in posterior/lateral and anterior/medial

regions. Thus, the variation in timescales in ventral frontal cortex could be influenced by the

patterns of projections into the frontal cortex from limbic (Cavada et al., 2000; Ghashghaei et

al., 2007) and/or sensory systems (Carmichael and Price, 1996).

Figure 3. Timescales and latencies within macaque ventral frontal cortex are not purely driven by cortical granularity. A) Schematic representations of the orbital surface of macaque's brain showing the relative location and cytoarchitectonic subdivisions of the ventral frontal cortex included in this analysis. B) Number of recorded neurons (lighter bars) and number of neurons successfully fit (darker bars) across areas. C) Average (+/-95% confidence intervals) timescale of single neurons in granular (green), dysgranular (orange) and agranular (blue) subdivisions of ventral frontal cortex. There is a significant main effect of granularity (OLS regression, dysgranular < granular: $t(2) = 10.58$, $p = 4.7*10^{-26}$; dysgranular < agranular: $t(2) = 7.34$, $p = 2.36*10^{-13}$; agranular ≈ granular: $t(2) = .19$, $p = 0.85$); although a model which includes area and granularity is no better at predicting timescale than one which includes area alone ($F(2) = 1.36$,

Discussion:

Here, we computed the timescales of single neurons recorded in parts of the fronto-limbic system in humans, macaques, and mice. We found that across the different species,

single-neuron timescales followed a similar hierarchical organization; hippocampus had the

shortest timescale whereas ACC had the longest. Further, while timescales in hippocampus,

ACC, and medial frontal cortex were highly consistent across species, the timescales of

amygdala and OFC neurons were more variable. We further found that differences in

timescales across macaque ventral frontal cortex followed specific spatial patterns but were

independent of the underlying variation in cytoarchitecture. Taken together, we report that

hierarchical organization of brain systems is common between three mammalian species

frequently studied in neuroscience, despite differences in some areas.

Timescales in subcortical areas

One of the most striking results from these data was the large divergence in amygdala

timescale across the species that we studied (**Figure 1**). We observed that mice had the longest

amygdala timescales while human amygdala timescales were the shortest. It is important to

note that within these mammals, large anatomical differences between the amygdala of

different species exist (Chareyron et al., 2011). Such differences in amygdala structure are most

dramatic within the basolateral subdivision, the target of the recordings considered in our

study. Other parts of amygdala like the central nucleus, by contrast, are more similar across

primates and rodents and it is possible that timescales could be more conserved in these nuclei.

Why this difference in basolateral timescales exists between the species is unknown, but work

by Nougaret et al. reported gradients of timescales within the basal ganglia, a circuit with well-

defined input and output structure (Nougaret et al., 2021). This then suggests that an area's

outputs might contribute to determining its timescale. Thus it follows that the basolateral

amygdala, which primarily projects to frontal cortex, might have longer timescales than central

nucleus, which primarily projects to brainstem and hypothalamus (Ghashghaei et al., 2007).

Future experimental work and analyses could intentionally characterize neurons from separate

amygdala subdivisions to determine whether anatomical and functional organization might also

extend to differences in timescales.

The macaque amygdala has a higher glia to neuron ratio, more sparse distribution of

neurons, and greater density of neuropil compared to rats (Chareyron et al., 2011). This

difference is even more stark when comparing to humans, where neurons in amygdala are even

more sparsely distributed than in macaques. Chareyron and colleagues speculate that these

specifications predispose the primate amygdala to a preferential role in integrating information

coming from a dramatically expanded cortex compared to rats. The basal and lateral nuclei

receive much of the input from the cortex and send intra-amygdala projections to other nuclei

(Freese and Amaral, 2009), placing them in a key position to perform such an integrative role. In

sensory areas, short timescales are hypothesized to be critical to quickly process and update

incoming information. Such shorter timescales in amygdala in humans and macaques compared

to mice could therefore predispose neurons in these species to be able to quickly integrate

information from the cortex which would in return enable the amygdala to drive learning in

areas with longer timescales, such as ACC (Stolyarova et al., 2019). That the primate amygdala

serves a more integrative role is further supported by recent insights about the organization of

projections from this area. Specifically, basolateral amygdala neurons in macaques exhibit

highly-organized patterns of collateral projections to the frontal lobe (Zeisler et al., 2023), while

those in mouse amygdala project more broadly in a less specific fashion (Zeisler et al., 2024).

While technically challenging, comparing the timescale of individual neurons whose anatomical

connections are known could bring important insight in our understanding of how neuronal

branching is related to temporal dynamics.

Finally, the cross-species variability in amygdala timescales stands in stark contrast to

hippocampus, where timescales were shorter than other brain areas measured here and highly

similar across species. The hippocampus exhibits much less differentiation across mammalian

species (Manns and Eichenbaum, 2006), with little about its anatomy and possibly the

computation it subserves changing throughout the course of evolution (Clark and Squire, 2013).

Thus, our finding that hippocampal timescales are similar across species appears to correlate

with such conservation. It's also striking that hippocampus had such short timescales given that

previous reports refer to primary sensory cortices as the prototype of an area with a short

timescale (Murray et al., 2014; Manea et al., 2022). Perhaps such similarity is driven by the fact

that most hippocampal inputs arise from the entorhinal cortex (Witter et al., 2000), in the same

way that primary sensory cortex inputs are predominantly from their respective sensory

organs.

Anatomy, granularity, commonality

It remains unclear what exactly about a brain area or a neuron sets its intrinsic

timescale. Possible contributing factors include region-wide variation in receptor expression,

genetic identity, intrinsic recurrence, or inputs from distant areas (Cavanagh et al., 2020;

Froudist-Walsh et al., 2021; Rapan et al., 2023). Indeed, we identified differences across

monkey cytoarchitectonic ventral frontal subdivisions (**Figure 3**). Such variation could also be

more significantly driven by network connectivity. Carmichael and Price have described two

largely non-overlapping networks within these regions (Carmichael and Price, 1996): the orbital

or so-called sensory network, and the medial or viscero-motor network. Recent functional

connectivity work (Stoll and Rudebeck, 2024b) has also explored information flow between

these regions and revealed that both areas 13m and 13l lack strong functional connectivity with

other OFC subregions during this particular decision-making paradigm. This low intra-frontal

connectivity might be influencing the relatively low timescale in these two subdivisions of area

13. By contrast, Stoll and Rudebeck show that the nearby area 12o is very highly interconnected

with other prefrontal regions; 12o is also more likely to receive information from many other

areas than to send it (Stoll and Rudebeck, 2024b). These results suggest that the highly

integrative role of area 12o might contribute to its relatively long timescale. Further evidence

for this relationship between connectivity and timescale could come from future experiments

targeting the visual system, with its well-documented organization of feed-forward and

feedback connections (Vanni et al., 2020), as one would expect shorter timescales in earlier

visual areas like V1 and longer timescales further up the visual hierarchy in V4 or parietal

cortex.

Within OFC, we identified spatial gradients of timescale that could not be accounted for

by cortical granularity alone (**Figure 3**). Rather, we found that timescales varied in a patchy

organization, with no clear pattern along a single anatomical axis. What drives this variation is

unclear, although it could be related to the projections coming into the ventral frontal cortex.

Inputs from the amygdala and hippocampus exhibit a sparse distribution in ventral frontal

cortex (Aggleton et al., 2015). Hippocampal inputs to OFC preferentially target more medial and

posterior areas, while amygdala inputs project more strongly to lateral areas (Ghashghaei et al.,

2007). Perhaps this difference in long-range innervation could drive differences in timescale

across the ventral surface of macaques. Within this view, the short timescale of hippocampal

neurons might entrain the neurons to which they project, although we note that the precise

effect of the presynaptic neurons' timescale on its downstream targets has yet to be assessed.

While often overlooked in analyses of timescales, we found that the presence or

absence of granule cells in layer IV modulated the latency of the exponential decay (**Figure 3B**).

Prior work by Fontanier, Sarazin and colleagues found that timescale and latency tend to vary

together and biophysical modeling revealed that the latency of a particular neuron was closely

related to potassium and other cationic currents (Fontanier et al., 2022). By contrast, we found

that variation in latency was related to granularity across macaque ventral frontal cortex,

whereas such a relationship was not present for timescale. Such a difference supports the idea

that the latency measure is distinct from timescale. Indeed, while there is consensus that a

longer timescale indicates a longer time interval over which a neuron might integrate

information (Soltani et al., 2021), a functional role for latency remains to be described. Further

work both *in vivo* and *in silico* is needed to fully understand how variations in latency might

alter computation.

Our analyses revealed a diversity of within-area timescales that were in some cases

consistent across species (**Figure 2**). The extent of these within-area differences has not been

previously reported. Such heterogeneity is hypothesized to be important for optimal

functioning both in brains and in artificial neural networks (Cavanagh et al., 2020; Stern et al.,

2023). Artificial neural network (ANN) modelling of prefrontal activity during a working memory

task suggests that timescale diversity is an adaptation to facilitate task performance; the

network had a largely homogenous timescale distribution before training which widened

significantly after repeated training iterations. Using brain-inspired spiking ANNs, Perez-Nieves

and colleagues successfully determined that heterogeneity is a feature of brain that is critical to

enable problem-solving (Perez-Nieves et al., 2021). ANN modeling therefore highlights the

computational significance of diverse timescales as well as the need for such heterogeneity to

both accurately model the brain during task performance.

While the timescales of neurons varied within an area we found that they were not

uniform but often exhibited a bimodal distribution (**Figure 2**). Importantly, a recent analysis of

intrinsic timescales in the macaque frontal eye field (FEF; Soyuhos et al., 2024), an area critical

for directing attention and coordinating eye movements, also identified bimodal distributions

of timescales. By characterizing these two populations using their functional properties during a

passive viewing task, they reported that short timescale FEF neurons primarily processed rapid

visual features of the task, while long timescale FEF neurons were related to sustaining

attention. Although a bimodal distribution was not directly reported, other studies have also

identified a relationship between variation in timescale and functional properties (Cavanagh et

al., 2020; Spitmaan et al., 2020; Fontanier et al., 2022). Since we did not characterize the

functional properties of the neurons included in our analysis, it remains an open question

whether bimodality in other brain areas is driven by related features. Further, while nearly all

timescale distributions considered in this analysis were bimodal, some species had uniquely

unimodal distributions: for example, macaque OFC or human amygdala. Whether this effect is a

result of the precise areas sampled or the task performed by subjects, understanding to what

extent bimodal distributions of timescales are ubiquitous across brain areas and behaviors is an

exciting next direction both for neuroscience and for computational modeling. Future research

should strive to further elucidate how the structure of variation of within-area timescales

across the cortex and subcortical structures contributes to the computational roles of those

areas in behavior.

Taken together, our data show that there are consistent timescale and latency

hierarchies across three commonly utilized model species: humans, macaque monkeys, and

mice. While we were able to reveal clear differences, particularly in OFC and amygdala, the high

degree of consistency across species suggests that a hierarchy of timescales might be a broad

organizing principle of mammalian brains. The implementation of such principles in neural

networks could provide improved sensitivity to both short- and long- term variation in

environmental inputs.

Methods:

Analyses were performed using MATLAB R2021a (MathWorks) and Python 3.9. All

figures made in Python used Seaborn (Waskom, 2021) and Matplotlib (Hunter, 2007) packages.

Other specific packages used will be cited in the methods below. Analysis code is available on

GitHub: https://github.com/RudebeckLab/timescales.

Neurophysiological datasets

Here, we analyzed the activity of single neurons recorded in humans, macaques and

mice across both open-source and unpublished datasets. **Table 1** illustrates the sources,

primary authors, and web links to these datasets as well as the papers in which they were

originally presented, if applicable. Unpublished datasets are indicated as such.

Table 1. Datasets used for cross-species analysis.

First Author	Species	Brain areas	# single neurons	Number of subjects	Analysis Paper Link	Dataset link
Chandra-vadia (Chandravadia et al., 2020)	Human	Amygdala Hippocampus	1086 759	59	https://doi.org/10.1038/s41597-020-0415-9	https://doi.org/10.17605/OSF.IO/HV7JA
Minxha (Minxha et al., 2020)	Human	Amygdala ACC Hippocampus	460 329 203	13	https://doi.org/10.1126/science.aba3313	https://osf.io/u3kcp/

Aquino (Aquino et al., 2024)	Human	OFC	320	9	https://doi.org/10.1523/JNEUROSCI.1628-23.2024	https://osf.io/5rv8f/
Fontanier (Fontanier et al., 2022)	Macaque	ACC	298	2	https://doi.org/10.7554/eLife.63795	https://doi.org/10.5281/zenodo.3947745
Stoll (Stoll and Rudebeck, 2024a)	Macaque	Amygdala OFC ACC	1691 5230 343	2	https://doi.org/10.1016/j.neuron.2024.03.020	Unpublished
Wirth (Wirth et al., 2017)	Macaque	Hippocampus	290	2	https://doi.org/10.1371/journal.pbio.2001045	http://crcns.org/datasets/hc/hc-12/about-hc-11
Young/ Mosher (Young et al., 2023)	Macaque	Amygdala mPFC	229 230	2	https://doi.org/10.1016/j.neuron.2023.07.012	https://zenodo.org/records/8164440
Steinmetz (Steinmetz et al., 2019)	Mouse	ACC Amygdala Hippocampus	841 317 2745	9	https://doi.org/10.1038/s4	https://figshare.com/articles/dataset/Dataset_fro

		mPFC	1274		1586-019-	m_Steinmet
		OFC	813		1787-x	z_et_al_201
						9/9598406

Operational Definitions of Brain Regions

For the sake of clearer comparisons, both within and across species, we combined
recordings into larger brain regions.

From mice (Steinmetz et al., 2019), hippocampal subfields CA1, CA2, and Dentate Gyrus
were all combined into the hippocampus category; Anterior cingulate area (ACA) was assigned
to ACC; Basolateral amygdala (BLA) to amygdala; Infralimbic area (ILA) and prelimbic cortex (PL)
to the mPFC category (Wise, 2008; Wallis et al., 2017); Orbital cortex (ORB) to OFC.

From macaque monkeys, amygdala recordings primarily targeted BLA, though precise
anatomical placement within amygdala was not formally assessed; OFC recordings were taken
from Brodmann's areas 11 and 13; subcallosal ACC (Walker's area 25) was assigned to the
mPFC category (Wise, 2008; Wallis et al., 2017); hippocampal recordings (Wirth et al., 2017)
were taken from across the extent of the hippocampus both in the antero-posterior and medio-
lateral planes; mid-cingulate cortex (MCC) recordings (Fontanier et al., 2022; Stoll and
Rudebeck, 2024a) were assigned to the ACC category, as this area has been referred both as
anterior MCC or dorsal ACC (Procyk et al., 2016).

From humans (Chandravadia et al., 2020; Minxha et al., 2020; Aquino et al., 2024), all
recordings were performed during seizure monitoring in cases of drug-resistant epilepsy;
recording targets were exclusively chosen according to clinical criteria. Amygdala recordings
preferentially targeted the BLA; ACC recordings were from Brodmann areas 24 and 32;
hippocampal recordings covered the extent of the hippocampus both in the A/P and M/L
planes; OFC recordings were taken primarily from Brodmann's areas 14, 13, and 10.

Macaque ventral frontal cortex dataset

Neural recordings were collected from two male rhesus macaques performing one of two behavioral tasks: either an instrumental preference task described in Stoll and Rudebeck (Stoll and Rudebeck, 2024a), or a learning task in which animals associated novel visual stimuli with different reward probabilities which changed over the course of the session (unpublished).

Recording procedures are described in more detail in Stoll and Rudebeck (Stoll and Rudebeck, 2024a). Briefly, animals were implanted with 157-channel semi-chronic recording microdrives (Gray Matter Research). During recordings, animals were seated in primate chairs while electrophysiological signals were recorded using the Plexon Omniplex system. Putative single neurons were determined using MountainSort (Chung et al., 2017) and manually curated based on cluster isolation metrics.

Electrodes were moved each day, and their cumulative adjustments tallied to determine their precise location in the brain. CT images collected throughout the recordings and histological assessment after perfusion were also used to verify these locations. Specifically, following standard perfusion with 4% formaldehyde (Electron Microscopy Sciences) in phosphate-buffered saline (PBS, Fisher), brains were extracted and post-fixed for 24-48 hours. Brains were then cryoprotected in increasing sucrose solutions (10%, 20%, 30% sucrose in PBS) before being shipped to FD NeuroTechnologies, Inc for processing, the details of which are also described in detail(Stoll and Rudebeck, 2024b). Brains were further cryoprotected, frozen in isopentane, and sectioned at 50um on a sliding microtome. Then equally spaced series of sections were prepared for either Nissl staining (one series) or immunohistochemistry (three series). Immunohistochemistry was performed for calbindin and SMI-32 in order to delineate boundaries between cortical areas.

Boundaries between cortical areas were determined according to features described previously (Carmichael and Price, 1994; Rapan et al., 2023; Stoll and Rudebeck, 2024b). In the most anterior sections, areas 11m/l, 12m, and 12r were separated primarily based on SMI-32 staining (11m and 11l were not consistently distinguishable in these animals and so have been

left grouped together). 11m/l is denser with cells than the more medial 13b, and it is denser

with superficial neurites in layer II/III than the more lateral 12m. Then, in Nissl-stained sections,

the 12r and 12m can be separated based on the dysgranular layer IV in the former compared to

the fully granular later IV in the latter. More posteriorly, area 13l was distinguished from 13m

by the presence of vertical processes in calbindin-stained sections, particularly in layers III and

V. Area 12o was defined by its stark lack of neurite staining in SMI-32-labeled sections

compared to the more medial 13m. 12l was distinguished from 12o by significantly increased

density of SMI-32-labeled cells. Finally, agranular insular areas were distinguished from area 13l

on the medial side and 12o on the lateral by the distinct lack of a granular layer IV; instead,

significant vertical processes were observed alongside a dense and thick layer IV in SMI-32-

stained sections.

Timescale estimation using inter-spike interval-based approach

Single-neuron timescales were estimated using an inter-spike interval-based (ISI)

method (Fontanier et al., 2022). Fitting was performed using MATLAB 2021a. Briefly, the time

difference between every spike in a neuron's time series and its following 100 spikes was

calculated. The lagged differences were then binned at 3.33ms from 0 to 1000ms. This

distribution was smoothed, after which we fitted the following equation:

399
$$R(k\Delta) = A\left[e^{-\left(\frac{k\Delta}{\tau}\right)} + B\right]$$

where $k\Delta$ is the time lag between bins, τ represents the neuron's *timescale*, describing

the rate of decay of the autocorrelation distribution, and A and B are free parameters which

together describe the peak and asymptotic limit of the autocorrelation. We also extracted the

time bin at which the autocorrelation begins decreasing, which we refer to as *latency*. As

before, fitting was performed using only the decreasing portion of the autocorrelation

distribution (after the latency).

Post-processing and plotting was performed in Python 3.9 using the following packages:

NumPy, Pandas, Matplotlib, Seaborn, SciPy, and statsmodels. Only neurons for which the model

fit well ($R^2 > 0.5$) and exhibiting timescales within the range $10 < \tau < 1000$ were included for

further analysis. To assess the effects of species and brain region on neuronal timescales and

latencies, we first used an ordinary least squares regression (OLS; statsmodels *ols* function). We

fitted the following linear model to describe a neuron's timescale and latency in terms of the

species and brain region from which it was recorded:

$$\tau = a + b_1 * species + b_2 * brain\ region$$

Post-hoc tests were performed within brain regions to compare species directly with FDR

corrections for multiple comparisons. Timescale and latency distributions within each brain

area were compared using a Brown-Forsythe test for equality of variances.

Firing rate analyses

To ensure our timescale and latency estimates were distinct from differences in other

firing rate statistics, we calculated the following measures for each neuron:

- Number of spikes (*n spikes*): number of spikes in timeseries
- Mean firing rate (*mean FR*): number of spikes divided by total length of timeseries
- Normalized FR (*norm FR*): spikes were binned at 500ms, and averaged with a sliding window of 1 second (2 bins); this firing rate was then z-scored using the mean firing rate above
- Fano: *norm FR* divided by the variance of *norm FR*
- Proportion of bursts (*prop burst*): proportion of spikes included in a burst, defined as the top 0.5 percentile of the ISI distribution(Ko et al., 2012)
- Proportion of pause (*prop pause*): proportion of spikes included in a pause, defined as the bottom 0.5 percentile of the ISI distribution(Ko et al., 2012)

All parameters were z-scored within each species and brain region. Then, we performed

OLS regressions predicting both timescale and latency using each z-scored parameter

individually alongside species and brain region. We extracted the percent explained variance for

each parameter and computed p-values for each. Models were directly compared using ANOVA

(statsmodels *anova_lm*).

Analysis of single neuron timescales in macaque ventral frontal cortex

To understand whether timescales were influenced by cortical granularity, we analyzed

an extended dataset containing single unit activity recorded from multiple subdivisions of the

ventral frontal cortex in monkeys, including the previously defined OFC neurons (**Table 2**) (Stoll

and Rudebeck, 2024b). We then used hierarchical model comparison to determine the relative

contributions of both cortical granularity and area. For both timescale and latency, two OLS

model were built, using either the specific area or the granularity as predictors. We then

compared the fit of those models using *anova_lm* to determine whether the differences in

degrees of freedom resulted in a better fit. Neurons' position in 3D space were also computed

using the recording depth, MRI and CT scans as well as using histological confirmation. This

allowed us to assess the relationship between spatial position and timescale/latency. 3D

coordinates were projected into 2D space; because our AP and DV coordinates were highly

correlated with one another (more posterior neurons are more ventral), we focused our

analyses only on the A/P and M/L planes. OLS regression models were built to explain the

extracted timescales and latencies using the AP and ML coordinates (including the interaction

term).

Table 2. Data used for within-frontal cortex analysis

Cytoarchitectonic Area	Larger Brain Region	Granularity	# single neurons
11m/l	OFC (orbitofrontal cortex)	Fully granular	2199
12m	vlPFC (ventrolateral prefrontal cortex)	Fully granular	770
12l	vlPFC	Fully granular	1115

12o	vlPFC	Fully granular	2268
12r	vlPFC	Dysgranular	774
13m	OFC	Dysgranular	1425
13l	OFC	Dysgranular	1606
AI (agranular insula)	n/a	Agranular	1886

453 Neurons located in areas 11m/l, 13m, and 13l comprise the OFC category used in the cross-

454 species analysis.

Supplemental Figure 1. Latency and timescale are weakly related to spiking statistics.

A) Percent explained variance (log scale) of timescale using a linear model containing area and species (black), and additional variance explained when including each firing rate statistic (grey). While the number of spikes was a significant predictor of timescale, the variance explained by this parameter was nearly two orders of magnitude less than the variance explained by area and species. P-values are also shown for each predictor.

B) Same as A for latency. Again, while the number of spikes, mean firing rate, and Fano factor were significant predictors of latency, together, they explain less than 1% of the total variance in latency.

References

References

Aggleton JP, Wright NF, Rosene DL, Saunders RC (2015) Complementary Patterns of Direct
Amygdala and Hippocampal Projections to the Macaque Prefrontal Cortex. Cereb Cortex
25:4351–4373.

Aquino TG, Courellis H, Mamelak AN, Rutishauser U, O'Doherty JP (2024) Encoding of Predictive
Associations in Human Prefrontal and Medial Temporal Neurons During Pavlovian
Appetitive Conditioning. J Neurosci 44 Available at: https://www-jneurosci-
org.us1.proxy.openathens.net/content/44/17/e1628232024 [Accessed July 5, 2024].

Carmichael ST, Price JL (1994) Architectonic subdivision of the orbital and medial prefrontal
cortex in the macaque monkey. Journal of Comparative Neurology 346:366–402.

Carmichael ST, Price JL (1996) Connectional networks within the orbital and medial prefrontal
cortex of macaque monkeys. Journal of Comparative Neurology 371:179–207.

Cavada C, Compañy T, Tejedor J, Cruz-Rizzolo RJ, Reinoso-Suárez F (2000) The Anatomical
Connections of the Macaque Monkey Orbitofrontal Cortex. A Review. Cerebral Cortex
10:220–242.

Cavanagh SE, Hunt LT, Kennerley SW (2020) A Diversity of Intrinsic Timescales Underlie Neural
Computations. Front Neural Circuits 14 Available at:
https://www.frontiersin.org/articles/10.3389/fncir.2020.615626/full?utm_source=S-
TWT&utm_medium=SNET&utm_campaign=ECO_FNINS_XXXXXXXX_auto-dlvrit
[Accessed January 4, 2021].

Chandravadia N, Liang D, Schjetnan AGP, Carlson A, Faraut M, Chung JM, Reed CM, Dichter B,
Maoz U, Kalia SK, Valiante TA, Mamelak AN, Rutishauser U (2020) A NWB-based dataset
and processing pipeline of human single-neuron activity during a declarative memory
task. Scientific Data 7:78.

Chareyron LJ, Lavenex PB, Amaral DG, Lavenex P (2011) Stereological analysis of the rat and
monkey amygdala. Journal of Comparative Neurology 519:3218–3239.

Chung JE, Magland JF, Barnett AH, Tolosa VM, Tooker AC, Lee KY, Shah KG, Felix SH, Frank LM,
Greengard LF (2017) A Fully Automated Approach to Spike Sorting. Neuron 95:1381-
1394.e6.

Clark RE, Squire LR (2013) Similarity in form and function of the hippocampus in rodents,
monkeys, and humans. Proc Natl Acad Sci U S A 110:10365–10370.

Fontanier V, Sarazin M, Stoll FM, Delord B, Procyk E (2022) Inhibitory Control of Frontal
Metastability Sets the Temporal Signature of Cognition Behrens TE, ed. eLife 11:e63795.

Freese JL, Amaral DG (2009) Neuroanatomy of the primate amygdala. In: The human amygdala,
pp 3–42. New York, NY, US: The Guilford Press.

Froudist-Walsh S, Xu T, Niu M, Rapan L, Margulies DS, Zilles K, Wang X-J, Palomero-Gallagher N
(2021) Gradients of receptor expression in the macaque cortex. Neuroscience. Available
at: http://biorxiv.org/lookup/doi/10.1101/2021.02.22.432173 [Accessed April 15, 2021].

Gao R, van den Brink RL, Pfeffer T, Voytek B (2020) Neuronal timescales are functionally
dynamic and shaped by cortical microarchitecture Vinck M, Colgin LL, Womelsdorf T,
eds. eLife 9:e61277.

Ghashghaei HT, Hilgetag CC, Barbas H (2007) Sequence of information processing for emotions
based on the anatomic dialogue between prefrontal cortex and amygdala. NeuroImage
34:905–923.

Hasson U, Yang E, Vallines I, Heeger DJ, Rubin N (2008) A Hierarchy of Temporal Receptive
Windows in Human Cortex. J Neurosci 28:2539–2550.

Hintiryan H et al. (2021) Connectivity characterization of the mouse basolateral amygdalar
complex. Nat Commun 12:2859.

Honey CJ, Thesen T, Donner TH, Silbert LJ, Carlson CE, Devinsky O, Doyle WK, Rubin N, Heeger
DJ, Hasson U (2012) Slow Cortical Dynamics and the Accumulation of Information over
Long Timescales. Neuron 76:423–434.

Hubel DH, Wiesel TN (1962) Receptive fields, binocular interaction and functional architecture
in the cat's visual cortex. The Journal of Physiology 160:106–154.

Hunter JD (2007) Matplotlib: A 2D Graphics Environment. Computing in Science & Engineering
9:90–95.

Ko D, Wilson CJ, Lobb CJ, Paladini CA (2012) Detection of bursts and pauses in spike trains.
Journal of Neuroscience Methods 211:145–158.

Li S, Wang X-J (2022) Hierarchical timescales in the neocortex: Mathematical mechanism and
biological insights. Proceedings of the National Academy of Sciences 119:e2110274119.

Manea AM, Zilverstand A, Ugurbil K, Heilbronner S, Zimmermann J (2022) Intrinsic timescales as
an organizational principle of neural processing across the whole rhesus macaque brain
Lee D, ed. eLife 11:e75540.

Manea AMG, Maisson DJ-N, Voloh B, Zilverstand A, Hayden B, Zimmermann J (2024) Neural
timescales reflect behavioral demands in freely moving rhesus macaques. Nat Commun
15:2151.

Manns JR, Eichenbaum H (2006) Evolution of declarative memory. Hippocampus 16:795–808.

Minxha J, Adolphs R, Fusi S, Mamelak AN, Rutishauser U (2020) Flexible recruitment of
533   memory-based choice representations by the human medial frontal cortex. Science 368
534   Available at: https://science.sciencemag.org/content/368/6498/eaba3313 [Accessed
535   September 9, 2020].

536   Murray JD, Bernacchia A, Freedman DJ, Romo R, Wallis JD, Cai X, Padoa-Schioppa C, Pasternak
537   T, Seo H, Lee D, Wang X-J (2014) A hierarchy of intrinsic timescales across primate
538   cortex. Nature Neuroscience 17:1661–1663.

539   Nougaret S, Fascianelli V, Ravel S, Genovesio A (2021) Intrinsic timescales across the basal
540   ganglia. Sci Rep 11:21395.

541   Perez-Nieves N, Leung VCH, Dragotti PL, Goodman DFM (2021) Neural heterogeneity promotes
542   robust learning. Nat Commun 12:5791.

543   Procyk E, Wilson CRE, Stoll FM, Faraut MCM, Petrides M, Amiez C (2016) Midcingulate Motor
544   Map and Feedback Detection: Converging Data from Humans and Monkeys. Cerebral
545   Cortex 26:467–476.

546   Rapan L, Froudist-Walsh S, Niu M, Xu T, Zhao L, Funck T, Wang X-J, Amunts K, Palomero-
547   Gallagher N (2023) Cytoarchitectonic, receptor distribution and functional connectivity
548   analyses of the macaque frontal lobe Badre D, Baker CI, Thiebaut de Schotten M, eds.
549   eLife 12:e82850.

550   Raut RV, Snyder AZ, Raichle ME (2020) Hierarchical dynamics as a macroscopic organizing
551   principle of the human brain. PNAS 117:20890–20897.

552   Soltani A, Murray JD, Seo H, Lee D (2021) Timescales of cognition in the brain. Current Opinion
553   in Behavioral Sciences 41:30–37.

554   Soyuhos O, Moore T, Chaudhuri R, Chen X (2024) Posterior Parietal Cortex Regulates Intrinsic
555   Neural Timescales and Attentional Processing in Frontal Eye Field. :2024.09.30.615928
556   Available at: https://www.biorxiv.org/content/10.1101/2024.09.30.615928v1 [Accessed
557   October 9, 2024].

558   Spillmann L (2014) Receptive Fields of Visual Neurons: The Early Years. Perception 43:1145–
559   1176.

560   Spitmaan M, Seo H, Lee D, Soltani A (2020) Multiple timescales of neural dynamics and
561   integration of task-relevant signals across cortex. PNAS 117:22522–22531.

562   Steinmetz NA, Zatka-Haas P, Carandini M, Harris KD (2019) Distributed coding of choice, action
563   and engagement across the mouse brain. Nature 576:266–273.

Stern M, Istrate N, Mazzucato L (2023) A reservoir of timescales emerges in recurrent circuits
with heterogeneous neural assemblies Ostojic S, Frank MJ, Schwalger T, eds. eLife
12:e86552.

Stoll FM, Rudebeck PH (2024a) Preferences reveal dissociable encoding across prefrontal-limbic
circuits. Neuron 112:2241-2256.e8.

Stoll FM, Rudebeck PH (2024b) Decision-making shapes dynamic inter-areal communication
within macaque ventral frontal cortex. Current Biology 34:4526-4538.e5.

Stoll FM, Rudebeck PH (2024c) Dissociable Representations of Decision Variables within
Subdivisions of the Macaque Orbital and Ventrolateral Frontal Cortex. J Neurosci 44
Available at: https://www.jneurosci.org/content/44/35/e0464242024 [Accessed
October 28, 2024].

Stolyarova A, Rakhshan M, Hart EE, O'Dell TJ, Peters M a. K, Lau H, Soltani A, Izquierdo A (2019)
Contributions of anterior cingulate cortex and basolateral amygdala to decision
confidence and learning under uncertainty. Nat Commun 10:4704.

van Heukelum S, Mars RB, Guthrie M, Buitelaar JK, Beckmann CF, Tiesinga PHE, Vogt BA,
Glennon JC, Havenith MN (2020) Where is Cingulate Cortex? A Cross-Species View.
Trends in Neurosciences 43:285–299.

Vanni S, Hokkanen H, Werner F, Angelucci A (2020) Anatomy and Physiology of Macaque Visual
Cortical Areas V1, V2, and V5/MT: Bases for Biologically Realistic Models. Cereb Cortex
30:3483–3517.

Viswanathan P, Nieder A (2017) Comparison of visual receptive fields in the dorsolateral
prefrontal cortex and ventral intraparietal area in macaques. European Journal of
Neuroscience 46:2702–2712.

Wallis CU, Cardinal RN, Alexander L, Roberts AC, Clarke HF (2017) Opposing roles of primate
areas 25 and 32 and their putative rodent homologs in the regulation of negative
emotion. PNAS 114:E4075–E4084.

Waskom ML (2021) seaborn: statistical data visualization. Journal of Open Source Software
6:3021.

Wirth S, Baraduc P, Planté A, Pinède S, Duhamel J-R (2017) Gaze-informed, task-situated
representation of space in primate hippocampus during virtual navigation. PLOS Biology
15:e2001045.

Wise SP (2008) Forward frontal fields: phylogeny and fundamental function. Trends in
Neurosciences 31:599–608.

Witter MP, Wouterlood FG, Naber PA, Van Haeften T (2000) Anatomical Organization of the
Parahippocampal-Hippocampal Network. Annals of the New York Academy of Sciences
911:1–24.

Young ME, Spencer-Salmon C, Mosher C, Tamang S, Rajan K, Rudebeck PH (2023) Temporally
specific patterns of neural activity in interconnected corticolimbic structures during
reward anticipation. Neuron 111:3668-3682.e5.

Zeisler ZR, Heslin KA, Stoll FM, Hof PR, Clem RL, Rudebeck PH (2024) Comparative basolateral
amygdala connectomics reveals dissociable single-neuron projection patterns to frontal
cortex in macaques and mice. Current Biology 34:3249-3257.e3.

Zeisler ZR, London L, Janssen WG, Fredericks JM, Elorette C, Fujimoto A, Zhan H, Russ BE, Clem
RL, Hof PR, Stoll FM, Rudebeck PH (2023) Single basolateral amygdala neurons in
macaques exhibit distinct connectional motifs with frontal cortex. Neuron 111:3307-
3320.e5.

610