Title

Constrained Open Chromatin Regions Reveal Functional Elements Shaping Human Traits

Kohei Tomizuka[1]†, Masaru Koido[1,2]†,*, Akari Suzuki[3]†, Soichiro Yoshino[1], Nao Tanaka[1],

Yuki Ishikawa[1], Xiaoxi Liu[1], Satoshi Koyama[4], Kazuyoshi Ishigaki[5], Yasuhiro Murakawa[6],

Immune transcript/enhancer consortium (ITEC)[#], Kazuhiko Yamamoto[3,7], Chikashi

Terao[1,8,9,*]

Affiliations

[1]Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical

Sciences, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa, 230-0045, Japan.

[2]Laboratory of Complex Trait Genomics, Department of Computational Biology and Medical

Sciences, Graduate School of Frontier Sciences, The University of Tokyo, 4-6-1,

Shirokane-dai, Minato-ku, Tokyo, 108-8639, Japan.

[3]Laboratory for Autoimmune Diseases, RIKEN Center for Integrative Medical Sciences, 1-

7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa, 230-0045, Japan.

[4]Laboratory for Cardiovascular Genomics and Informatics, RIKEN Center for Integrative

Medical Sciences, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa, 230-0045,

Japan.

[5]Laboratory for Human Immunogenetics, RIKEN Center for Integrative Medical Sciences,

1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa, 230-0045, Japan.

[6]RIKEN-IFOM Joint Laboratory for Cancer Genomics, RIKEN Center for Integrative Medical

Sciences, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa, 230-0045, Japan.

[7]Drug Discovery Antibody Platform Unit, RIKEN Center for Integrative Medical Sciences, 1-

7-22 Suehiro-cho, Tsurumi-ku, Yokohama City, Kanagawa, 230-0045, Japan.

[8]Clinical Research Center, Shizuoka General Hospital, 4 Chome-27-1 Kitaando, Aoi Ward,

Shizuoka, 420-8527, Japan.

[9]Department of Applied Genetics, The School of Pharmaceutical Sciences, University of

Shizuoka, 52-1 Yada, Suruga Ward, Shizuoka, 422-8526, Japan.

[#]Immune transcript/enhancer consortium (ITEC) members

Matteo Guerrini, Hiroaki Hatano, Keiko Hikino, Kazuyoshi Ishigaki, Yuki Ishikawa, Michihiro

Kono, Xiaoxi Liu, Yasuhiro Murakawa, Masahiro Nakano, Akiko Oguchi, Raku Son, Akari

Suzuki, Kazuhiro Takeuchi, Nao Tanaka, Chikashi Terao, Kohei Tomizuka, Kazuhiko

Yamamoto, Soichiro Yoshino

†These authors contributed equally: Kohei Tomizuka, Masaru Koido, Akari Suzuki

*Corresponding authors:

Chikashi Terao (chikashi.terao@riken.jp)

Masaru Koido (masaru.koido@a.riken.jp)

Abstract

Open chromatin regions (OCRs) define cell-type-specific regulatory elements across the

genome, yet their functional significance varies, making it challenging to pinpoint

biologically essential regions. Here, we introduce CAMBUS (Chromatin Accessibility

Mutation Burden Score), a machine-learning framework that identifies active and

evolutionarily constrained OCRs by leveraging surrounding DNA sequences. Applying

CAMBUS to 29 immune cell types, we identified 66,043 constrained OCRs, which were

substantially enriched in the known constraint genome (odds ratio=11.45 (95% confidence

interval 9.33–14.05), P=4.7 × 10^{-68}), while 90% of these OCRs were not prioritized by existing constraint metrics. These OCRs were highly enriched for known enhancers and super-enhancers, independent of known epigenetic markers and annotated regions, and overlapped with regulatory elements implicated in immune-mediated diseases and experimentally validated functional variants, including rare variants, particularly in leukocyte-related traits. Furthermore, CAMBUS revealed cell-type-specific transcriptional regulatory landscapes, linking genetic constraint with gene regulation in immune cells and identifying plausible connections between 1,533 causal variants and 70 complex traits. By defining biologically constrained regulatory elements at high resolution, CAMBUS provides a framework for understanding the selective pressures shaping the non-coding genome and its role in human health and disease.

Introduction

Genome sequences define species identities and evolutionary trajectories by shaping cellular, organ, and functional complexities (O'Bleness, Searles et al. 2012, Pollen, Kilik et al. 2023). Given that protein-coding regions account for only a small fraction of vertebrate genomes (Leypold and Speicher 2021), understanding the role of non-coding regions—particularly active regulatory elements—is critical for deciphering the molecular basis of species identity and function (Ward and Kellis 2012, Franchini and Pollard 2017). Active regulatory regions, such as enhancers, regulate gene expression in a cell-type- and state-specific manner (Andersson, Gebhard et al. 2014, Soskic, Cano-Gamez et al. 2019, Gupta, Weinand et al. 2023, Oguchi, Suzuki et al. 2024).

Several genome-wide experimental methods, including chromatin immunoprecipitation followed by sequencing (ChIP-seq) for histone modifications, DNase I

hypersensitive site sequencing (DNase-seq), and assay for transposase-accessible

chromatin using sequencing (ATAC-seq), have been developed to identify candidate

regulatory regions. Among them, ATAC-seq has become widely adopted due to its

technical simplicity and robust performance (Buenrostro, Giresi et al. 2013, Yan, Powell et

al. 2020). Compared with ChIP-seq peaks, ATAC-seq-defined open chromatin regions

(OCRs) provide high positional specificity for identifying regulatory elements (Consortium,

Moore et al. 2020). Because chromatin accessibility is a prerequisite for gene transcription,

mapping and characterizing OCRs—particularly those that are cell-type-specific or

conserved across multiple lineages—offers crucial insights into general and context-

specific regulatory mechanisms, evolutionary constraint, and the basics of complex traits

(Boyle, Davis et al. 2008, Zhang, Hocker et al. 2021).

Although genome-wide association studies (GWAS) in human have highlighted

the enrichment of disease-associated variants within non-coding regions (Maurano,

Humbert et al. 2012, Claussnitzer, Cho et al. 2020), and particularly within regulatory

elements (Trynka, Sandor et al. 2013, Soskic, Cano-Gamez et al. 2019, Zeng, Bendl et al.

2024), distinguishing active regulatory elements from non-active OCRs remains challenging

(Maurano, Humbert et al. 2012, Visscher, Wray et al. 2017, Schaid, Chen et al. 2018), as

ATAC-seq measures chromatin openness but does not directly assess regulatory activity

(Thibodeau, Uyar et al. 2018, Klemm, Shipony et al. 2019). Therefore, integrating additional

layers of information is necessary to systemically predict active and functional regulatory

regions.

Constrained genomic regions, which show evidence of purifying selection, serve

as important markers of functionally active elements (Ward and Kellis 2012, Radke, Sul et

al. 2021). The systematic identification of constrained, cell-type-specific active regulatory

regions across diverse tissues remains an open challenge. Recent large-scale human

whole-genome sequencing efforts have enabled the development of constraint metrics

such as Gnocchi (Chen, Francioli et al. 2024), which leverage local mutability to infer constrained sites. Consortia efforts such as ENCODE Phase 3 have combined chromatin accessibility with histone modification ChIP-seq signals to annotate candidate cis-regulatory elements (cCREs), but cCREs cannot fully capture functional regulatory regions.

In addition to evolutionary constraint and epigenomic signals, chromatin accessibility quantitative trait loci (caQTLs), which identify genetic variants that alter chromatin openness, provide further evidence that genetic variation at regulatory regions influences complex traits (Kumasaka, Knights et al. 2016, Gate, Cheng et al. 2018). However, caQTL mapping requires large cohorts and is limited in its ability to assess rare variants. Recently, machine learning models trained on DNA sequences have emerged as a powerful alternative, capable of predicting regulatory activity and inferring variant effects without requiring population-level data (Zhou, Theesfeld et al. 2018, Eraslan, Avsec et al. 2019, Koido, Hon et al. 2023). These models have enabled the tissue- and cell-type-specific prediction of gene expression and chromatin features from sequence alone (Zhou and Troyanskaya 2015, Koido, Hon et al. 2023).

Together, these observations highlight the need for methods that integrate evolutionary constraint, chromatin accessibility, and DNA sequence features to prioritize functional regulatory elements. Motivated by this gap, we developed CAMBUS (Chromatin Accessibility Mutation Burden Score), a new framework for prioritizing active and evolutionarily constrained OCRs. By leveraging sequence-embedded constraint information (Fig. 1), CAMBUS provides a principled approach to identify regulatory elements with likely functional importance across diverse cell types. We assessed the characteristics of CAMBUS and their cell-type specificity using OCRs in various cell types in peripheral blood mononuclear cells (PBMCs). By leveraging CAMBUS, we identified 1,533 cell-type-specific regulatory variants, including rare variants, with potential functional significance in complex traits across European (EUR) and East Asian (EAS) ancestries (Fig. 1D and 1E). We

demonstrated how integrating genetic constraint with chromatin accessibility can refine the

biological interpretation of non-coding regulatory variants and their roles in transcriptional

regulation in a cell-type-specific manner.

Results

Comprehensive ATAC-seq profiling of 29 immune cell types

We sorted PBMCs (peripheral blood mononuclear cells) and Neutrophils from 55

Japanese individuals (Methods). PBMCs were sorted into 27 immune cell types by cell

sorter. We conducted ATAC-seq for PBMC, Neutrophil, and the 27 cell types (hereafter, 29

cell types) in eight subgroups (CD4; CD4$^+$ T-cells, CD8; CD8$^+$ T-cells, B; B cells, Mono;

Monocytes, DC; dendritic cells, NK; Natural Killer cells, Neutrophils, and PBMC)

(Supplementary Table 1). The number of cell types in this study is comparable to that in

recent scATAC-seq studies using PBMC (Zhang, Wang et al. 2020, Benaglio, Newsome et

al. 2023). Notably, the 29 cell types include minor cell types, such as double-negative B-

cells (~5% in B-cells) (Li, Li et al. 2021). After stringent quality control, we obtained

7,939,714 OCRs that covered 498 Mb (16.1 %) of the human genome (Methods). The

median number and size of OCRs were 252,668 OCRs and 276 bp, respectively. Among

the 7,939,714 OCRs, 330,367 (4.2%) were observed in only one cell type, and 437,368

(5.5%) were observed in only one subgroup. Both hierarchical clustering and dimension

reduction using principal component analysis (PCA) and uniform manifold approximation

and projection (UMAP) showed that these OCRs could clearly distinguish the eight

subgroups but not all 29 immune cell types (Extended Data Fig. 1).

Detecting constrained OCRs by CAMBUS

To infer constraint levels in the OCRs, we first developed machine learning models, CAMBUS ML (Chromatin Accessibility Mutation Burden Score machine learning), to predict the OCRs of each of 29 cell types from DNA sequence patterns, and calculated *in silico* saturated mutation effects- quantifying the predicted changes in accessibility probability for all possible single nucleotide mutations within 2kb of each OCR (whether or not the mutations were reported) (Methods). We aggregated and took the mean of mutation effects within 2kb of each OCR and called this value CAMBUS. Hereafter, we defined OCRs with negative CAMBUS as CAMBUS+ (where the '+' denotes positive constraint) based on the local false discovery rate (FDR) method (Fig. 2A, 2B; Method). Besides, we defined relaxed CAMBUS+ OCRs, a more relaxed CAMBUS+ set by extracting the bottom 5% of the overall CAMBUS distribution. The median number of CAMBUS+ OCRs was 1,584, covering 0.056% of the human genome, while that of relaxed CAMBUS+ OCRs was 12,632, covering 0.33%. Notably, read counts from ATAC-seq were not a determinant factor for CAMBUS+ OCRs (Extended Data Fig. 2A), reinforcing that our machine-learning approach incorporated information from DNA sequence patterns into OCRs, not from technical fluctuations or noises.

To confirm our key assumption that CAMBUS+ OCRs (or relaxed CAMBUS+ OCRs) are under positive constraint, we assessed whether CAMBUS+ OCRs overlapped with orthogonal seven constrained metrics (Methods). As expected, five out of the seven constraint metrics supported the constraint hypothesis for CAMBUS+ OCRs across 29 cell types (Method; Supplementary Table 3). Notably, we found that CAMBUS+ OCRs significantly cover the strongly constrained regions (one-sided Wilcoxon-test P = 2.5 × 10^{-69} for CAMBUS+ OCRs in NaiveCD4). Although CAMBUS+ OCRs contained a greater proportion of coding regions than control OCRs (Extended Data Fig. 2B), the same trend of enrichment was observed even when the analysis was restricted to non-coding regions

(one-sided Wilcoxon-test P = 9.1 × 10^{-46} for CAMBUS+ OCRs in NaiveCD4; Fig. 2C;

Extended Data Fig. 2C). Of note, when we analyzed each of CAMBUS+ OCRs, 59,605

(90%) of CAMBUS+ OCRs were not prioritized by existing constraint metrics (Methods).

Nevertheless, the CAMBUS+ OCRs showed the strongest constraint defined by Gnocchi

(odds ratio (OR)=11.45 (95% confidence interval (CI) 9.33–14.05), P=4.7 × 10^{-68} by

Fisher's Exact test) among the compared regulatory elements (Fig. 2D; Supplementary

Table 4). Meta-analysis demonstrated the consistent tendency across the 29 cell types

(OR=8.71 (95% CI 7.85–9.66), P=3.1 × 10^{-365} from random-effect meta-analysis; Fig. 2E),

outperforming the ENCODE promoter-like regulatory elements (OR=7.3), the strongest

constraint candidate regulatory elements in the previous study (Chen, Francioli et al. 2024).

Collectively, these indicate that CAMBUS+ OCRs could uniquely find regions under very

strong positive constraints. The relaxed CAMBUS+ OCRs also showed the same patterns

with slightly reduced magnitude (Extended Data Fig. 2D).

Previous studies (Silvert, Quintana-Murci et al. 2019) demonstrated that enhancers,

especially activated in T-cells, are enriched in archaic genome sequences due to positive

selection. Consistently, we observed that CAMBUS+ OCRs showed significant enrichment

in the selection signals (Nait Saada, Kalantzis et al. 2020) (OR=1.4 (95% CI 1.3–1.5),

P=1.1 × 10^{-26} from random-effect meta-analysis; Extended Data Fig. 2E), the Neanderthal

genome sequences observed in the Japanese population (Liu, Koyama et al. 2024)

(OR=1.4 (95% CI 1.4–1.5), P=4.8 x 10^{-32} from random-effect meta-analysis; Extended Data

Fig. 2F), and the Neanderthal and Denisovans' sequences observed in 1000 Genomes

populations (Supplementary Table 5) (Browning, Browning et al. 2018). Besides, we

observed the enrichment of CAMBUS+ in human ancestor quickly evolved regions

(HAQER) (Mangan, Alsina et al. 2022) (OR=2.35 (95% CI 1.91–2.89), P=8.3 × 10^{-16}).

Taken together, CAMBUS+ OCRs can define novel constrained regions, well supported by

other constraint-relevant metrics.

Enrichment of active regulatory elements by selecting CAMBUS+ OCRs

As expected, the OCRs obtained in this study were well overlapped with the known

candidate regulatory elements; for example, 73.66% of all OCRs in NaiveCD4 were

overlapped with super-enhancers (Jiang, Qian et al. 2019) (the median of 29 cell type is

74.70%) and 4.49% with FANTOM5 enhancers (Andersson, Gebhard et al. 2014) (the

median of 29 cell type is 5.10%; Supplementary Table 7). Then we explored whether

CAMBUS+ OCRs could aid in prioritizing functional regulatory elements in comparison with

control OCRs (Methods). As a result, CAMBUS+ OCRs significantly enriched for the known

regulatory elements, especially active enhancers, like super-enhancers (NaiveCD4:

OR=9.07 (95% CI 6.80–12.11), Fisher's exact test $P = 2.0 \times 10^{-102}$) and FANTOM5

enhancers (NaiveCD4: OR=6.16 (95% CI 5.39–7.03), Fisher's exact test $P = 6.3 \times 10^{-111}$)

(Supplementary Table 8). Meta-analysis showed a consistent trend across 29 cell types

(OR=6.63 (95% CI 5.50–7.99), $P = 1.1 \times 10^{-87}$ from random-effect meta-analysis for super-

enhancers; Extended Data Fig. 2H, and OR=5.19 (95% CI 4.87–5.54), $P = 4.9 \times 10^{-547}$ from

random-effect meta-analysis for FANTOM5 enhancers; Extended Data Fig. 2I).

Remarkably, after adjusting for known constraint metrics, we still observed statistically

significant enrichment for regulatory elements in CAMBUS+ OCRs across the 29 cell types

(Fig. 2F; Supplementary Table 9). Reflecting the characteristics of the analyzed cell types

(blood and immune cell types), a comparison of enrichment across four tissues

(blood/hematopoietic, brain, kidney, and lung) revealed the strongest prioritization of

blood's regulatory elements (CD4 as a representative: super-enhancers; OR=6.83 (95% CI

6.07–7.71), $P=1.1 \times 10^{-217}$; FANTOM5 enhancers; OR=7.64 (95% CI 6.64–8.78), $P=8.7 \times$

10^{-179} from logistic regression; Fig. 2G; Supplementary Table 10). This pattern was

consistently observed across 29 cell types (Supplementary Table 11). We also confirmed

that the relaxed CAMBUS+ OCRs exhibited the same tendency (Extended Data Fig. 2J;

Supplementary Table 12). Taken together, CAMBUS+ OCRs (and relaxed CAMBUS+

OCRs) effectively capture active regulatory elements under positive constraints in a cell-

type-specific manner, particularly enhancers, that were not well captured by the previous

methods.

Heritability enrichment of CAMBUS+ in blood-relevant diseases and traits

Since the majority of trait heritability is captured by regulatory regions, we computed

heritability enrichments in CAMBUS+ OCRs using stratified LD score regression (S-LDSC)

(Finucane, Bulik-Sullivan et al. 2015) conditioning on known functional annotations and all

OCRs from this study (Methods). We used GWAS summary statistics for 33 complex traits,

including 14 immune-mediated traits (e.g., multiple sclerosis (MS), rheumatoid arthritis (RA),

and Atopic dermatitis (AD)), from European (EUR) and East Asian (EAS) populations

(Supplementary Table 13).

As a result, CAMBUS+ OCRs demonstrated the highly significant heritability enrichment

in a wide variety of immune-mediated diseases (FDR < 5%; Fig. 3A; Supplementary Table

14). Particularly, we found plausible pairs of immune traits and cell types supported by the

strong enrichment of heritability: MS and T_h17 (Moser, Akgun et al. 2020), and systemic

lupus erythematosus (SLE) and B cells (Jarvinen, Hellquist et al. 2012, Fillatreau, Manfroi

et al. 2021) (Extended Data Fig. 3A). Other plausible pairs included pairs of RA with CD4 or

CD8[+] T cells, and AD with CD4 (particularly Th2 (Honda and Kabashima 2020)), or CD8[+] T

cells. Remarkably, these enrichments were not observed in other constraint metrics (Fig.

3B).

Based on these results, we hypothesized that polygenic risk scores (PRS) prioritized by

CAMBUS+ OCRs could enhance their predictive ability. To evaluate this hypothesis, we

focused on CAMBUS+ OCRs in double-negative B cells (DN B), which showed significant

enrichment for the heritability of SLE in EAS ancestry (Extended Data Fig. 3A) and was

also known for its functional associations with SLE (Chung, Gong et al. 2023). We

constructed the PRS based on variants overlapping CAMBUS+ OCRs and evaluated the

additional performance of the conventional PRS by five-fold cross-validation using 137,620

individuals (Methods). A fixed effect meta-analysis demonstrated that the additional

variants significantly improved the predictive performance of PRS ($P = 3.12 \times 10^{-6}$;

Supplementary Table 15; Methods). These suggest that CAMBUS+ OCRs help us to

understand the basics of the complex traits in a polygenic manner and in a cell-type

specific manner and help us to construct better PRS by prioritizing target cell types.

Putative causal variants were enriched in CAMBUS+ OCRs

To further search the potential usefulness of CAMBUS+ OCRs in a single variant

resolution, we assessed whether CAMBUS+ OCRs could prioritize or pinpoint the putative

causal variants for complex traits. As expected, the median regional length of CAMBUS+

(845 bp) was much shorter than that of 95% credible sets (3.0M bp for EUR ancestry and

1.1M bp for EAS ancestry), which may potentially pinpoint a variant. We observed

CAMBUS+ OCRs showed significant enrichment in the pathogenic regulatory variants

cataloged in the Human Gene Mutation Database (HGMD) (Stenson, Ball et al. 2003)

(OR=2.70 (95% CI 2.43–2.99), $P = 1.7 \times 10^{-77}$; from random-effect meta-analysis; Extended

Data Fig. 2G; Supplementary Table 6). This is in line with overlap between CAMBUS+

OCRs and super-enhancers or FANTOM5 enhancers, both of which are known to be

enriched for causal variants (Hnisz, Abraham et al. 2013). We further analyzed the large-

scale statistical fine-mapping studies in 94 traits in EUR ancestry (Kanai, Ulirsch et al.

2021) and 62 traits in EAS ancestry (Koyama, Liu et al. 2024). The CAMBUS+ OCRs were

significantly enriched for the putative causal variants in 95% credible sets (median OR:3.85

and 3.94 in EUR and EAS, respectively, Extended Data Fig. 4A). In line with a data source

of OCRs, CAMBUS+ OCRs exhibited significantly higher enrichment for the putative causal

variants of hematopoietic traits, especially leukocyte-related traits (Methods,

Supplementary Table 21) than non-hematopoietic traits (two-sided t-test P = 1.0×10^{-11} in

EUR ancestry and 1.4×10^{-2} in EAS ancestry for leukocyte-related traits; Extended Data

Fig. 4B). As an example, CAMBUS+ OCRs in type 1 helper T-cells demonstrated stronger

enrichment for variants with high posterior inclusion probability (PIP) (>0.9) for

hematopoietic traits in EAS ancestry (OR=7.62 (95% CI 3.60–16.18), P = 1.2×10^{-7} from

logistic regression; Fig. 4A), and this enrichment was further increased when restricted to

the leukocyte-related traits (OR=11.0 (95% CI 4.04–29.96), P = 2.7×10^{-6}; Fig. 4A). Meta-

analysis showed that this trend of leukocyte-related traits was consistent across 29 cell

types (OR=8.99 (95% CI 7.85–10.29), P = 8.8×10^{-221} for EUR ancestry; OR=19.24 (95%

CI 16.03 – 23.09), P = 3.9×10^{-221} for EAS ancestry; Fig. 4B). Taken together, CAMBUS+

OCRs can pinpoint or at least narrow down causal variants.

Notably, even for rare variants (minor allele frequency (MAF) < 0.01), the putative causal

variants of hematopoietic traits were significantly enriched in CAMBUS+ OCRs (Methods;

OR=4.68 (95% CI 2.66–7.99), two-sided Fisher's exact test P = 2.23×10^{-6}; Fig. 4C). This

suggested that CAMBUS+ OCRs can potentially prioritize even rare causal variants which

are difficult to prioritize in the conventional statistical fine-mapping (Schaid, Chen et al.

2018).

We found plausible examples of putative causal variants that were not covered by known

constraint regions (Gnocchi) but were identified by CAMBUS+ OCRs, leading to the

biological interpretation of GWAS findings. First, rs13122305 was the only one overlapping

with CAMBUS+ OCRs in monocytes among seven candidate causal variants in a credible

set for monocyte count (Fig. 4D). The nearest gene of this variant is *COQ2*, whose anti-

inflammatory functions via regulating gene expression were reported (Armanfar, Jafari et al.

2015, Mantle, Heaton et al. 2021). Additionally, we found that the Activity-by-Contact (ABC)

model's prediction (ABC enhancer) in monocytes only overlapped with rs13122305 and

that *COQ2* was also supported by the ABC model with the second-highest ABC score

(0.025) among the eight target genes. Second, among the three candidate causal variants

in a credible set in the FCGR region for Albumin/Globulin ratio (AG), only rs2487451

overlapped with CAMBUS+ OCR. Non-classical monocytes, known to display antigen

presenting properties (Mukherjee, Kanti Barman et al. 2015), were the only cell type whose

OCR overlapped with the candidate causal variant. The nearest gene was *FCGR3B*,

encoding a receptor (FcγRIIIB) that binds to the Fc region of IgG (Immunoglobulin)

antibodies, which is known to be expressed in non-classical monocytes (Nassir, Tambi et al.

2021). Third, rs55646091, a putative causal variant for asthma (PIP = 0.99), overlapped

with CAMBUS+ OCR of regulatory T-cells (T-reg) (Fig. 4E). The nearest gene of this

variant was *LRRC32* (leucine-rich repeat-containing 32), which regulates TGF-beta.

LRRC32 was potentially involved in asthma (Li, Wang et al. 2023) and whose encoding

protein was known as a surface marker for T-reg. Furthermore, this variant-gene pair was

supported by the ABC model and *LRRC32* displayed the highest ABC score (ABC score =

0.041 that is much higher than the threshold corresponding to 70% recall and 59%

precision) (Fulco, Nasser et al. 2019) among the nine genes connected with the regulatory

element overlapping with the OCR and the variant. Fourth, we found that rs8009224 was

the only one that overlapped with CAMBUS+ OCRs of plasmablast (Extended Data Fig.

5A) among the seven putative causal variants for hemoglobin A1c levels. This is an intronic

variant of *MIDEAS* (Mitotic Deacetylase Associated SANT Domain Protein), which is low

frequency in both 1KG EUR (MAF = 3.4%) and EAS (MAF = 1.2%) and also associated

with mean corpuscular hemoglobin concentration (Barton, Sherman et al. 2021). The

overlapping regulatory element was connected to the 16 genes by the ABC enhancer in B-

cell, and the putative target gene with the second highest ABC score is *MIDEAS* (ABC

score = 0.097, much higher than the proposed threshold corresponding to 70% recall and

59% precision). Lastly, we demonstrated that CAMBUS+ in T-reg could prioritize only one

putative causal variant (rs9814991) from numerous variants (fifty-seven) in the credible set

for eosinophil count (Extended Data Fig. 5B). These indicate that CAMBUS+ OCRs,

together with statistical fine-mapping, could pinpoint putative causal variants of complex

traits in a single base resolution and in a cell type-specific manner.

Discussion

In this study, we developed CAMBUS based on the hypothesis that DNA

sequencing patterns surrounding OCRs contain hidden information defining OCR

constraints. Although CAMBUS+ OCRs were derived from a fundamentally different

approach, they were well supported by conventional constraint metrics as expected, while

also capturing high-resolution, cell-type-specific regions that are both active and

constrained, beyond existing annotations. An alternative potential approach using ATAC-

seq (or DNase-seq) could prioritize regions with high accessibility (i.e., regions with many

read counts). However, previous studies have suggested no relationship between

accessibility and functionality or constraints (Roadmap Epigenomics, Kundaje et al. 2015,

Kuderna, Ulirsch et al. 2024, Martens, Fischer et al. 2024). Consistently, accessibility was

not a determinant factor in defining CAMBUS+ OCRs. Thus, leveraging surrounding DNA

sequence patterns through machine learning offers a unique and effective strategy for

identifying biologically important OCRs.

CAMBUS+ OCRs can mark active and functional regulatory elements, particularly

tissue-specific active enhancers, even after being adjusted for known constraint metrics

(Gnocchi, phyloP46way, phyloP100way, cactus241way, phyloPPrimates, GERP and DR).

Furthermore, CAMBUS+ OCRs captured a significant proportion of the heritability for many

complex traits, highlighting their biological and genetic relevance. CAMBUS+ OCRs

effectively prioritized regulatory causal variants for immune-related traits, even rare causal

variants. By defining biologically constrained regulatory elements, CAMBUS+ OCRs will

provide a foundation for understanding the regulatory architecture of complex traits and

their biological insights into genetic variations especially in non-coding regions.

We demonstrate that CAMBUS+ OCRs can effectively prioritize causal rare

variants for complex traits. A previous study showed that rare variants tend to be enriched

in the constraint genome (Karczewski, Francioli et al. 2020), consistent with our study as

CAMBUS+ OCRs can mark such constrained regions. In GWAS using SNP arrays, low

statistical power and low imputation accuracy for rare variants make it difficult to detect rare

variant associations and identify causal rare variants through statistical fine-mapping.

Despite these challenges, CAMBUS+ OCRs can facilitate the rare variant prioritization that

might otherwise be missed.

Eosinophilic asthma is a subtype characterized by elevated levels of eosinophils

(Nelson, Bush et al. 2020). In this analysis, we identified a variant (rs55646091) associated

with asthma linked to increased eosinophil counts (PIP = 0.03). This variant exhibited

positive effect sizes in both traits (0.20 for asthma, 0.072 for eosinophil counts), suggesting

that carriers of this variant demonstrate elevated eosinophil levels, which contribute to

inflammation and the onset of asthma (Harb and Chatila 2023). Notably, this variant

overlaps specifically with the OCR in T-reg cells among 29 blood cell types analyzed. The

nearest gene of this variant is *LRRC32*, which regulates TGF-beta, a known surface marker

of T-reg cells, which has been previously implicated in immune regulation (Tran, Andersson

et al. 2009). The findings from this analysis provide additional evidence supporting a

biological mechanism in which the variant, through its effects on T-reg cells, leads to

increased eosinophil counts, thereby contributing to the development of asthma. This

insight deepens our understanding of the role of T-reg cells and *LRRC32* in asthma

pathogenesis, particularly in eosinophilic inflammation.

In this study, we focused on various immune cell types; however, our approach is

broadly applicable to other tissues and cell types. Because calculating CAMBUS relies on

in silico saturated mutagenesis across millions of OCRs, our CNN-based framework is

practical for large-scale regulatory annotation across diverse cell types, achieving

substantially greater computational efficiency than recent large-scale generative AI models

(e.g., Evo 2 (Brixi, Durrant et al. 2025) will require more than 200 years for the task on a

single H100 GPU (Supplementary Note)). As GWAS results continue to accumulate,

integrating CAMBUS+ OCRs with these expanding resources has the potential to uncover

further insights into the regulatory mechanisms underlying complex traits. Moreover, this

method can be applicable to OCRs identified at the single-cell level, further extending its

applicability. Taken together, CAMBUS+ OCRs not only capture constrained and functional

regulatory elements but also facilitate the interpretation of non-coding genetic variants by

prioritizing causal variants in a cell-type-specific manner without relying on LD information

and allele frequencies.

Method

Sample and library preparation

We collected peripheral blood (PB) cells from 55 healthy Japanese volunteers and sorted

them into 29 cell types. PBMCs were isolated by density gradient separation with Ficoll-

Paque (GE healthcare) immediately after blood drawing. Erythrocytes were lysed with a

potassium ammonium chloride buffer, and non-specific binding was blocked with Fc-

gamma receptor antibodies. PBMC is stained with cell surface specific antibodies

(Supplementary Table 22). 27 immune cell populations were sorted by FACS (BDAria III)

using the gating strategy (Supplementary Table 1; Extended Data Fig. 6). In addition,

neutrophils were isolated by MACSxpress Neutrophil Isolation Kits human (Miltenyi Biotec)

from 4 × 106 PBMC cells. Sorted cells, isolated PBMC, and Neutrophils were stored at

−80°C in CytoStor-10 (Charles River Laboratories Cell Solutions). Libraries for ATAC-seq

were prepared using a modified protocol of Omni-ATAC (Corces, Trevino et al. 2017).

Genomic DNA from whole blood was isolated using QIAamp DNA Blood Mini Kits,

respectively. ATAC-seq was performed using 150bp paired-end on Illumina NovaSeq6000.

In total, we obtained a median of 13.34M (+/- 7.56M) reads per sample. This study was

approved by the Ethics Committees of RIKEN. Written informed consent was obtained from

each volunteer.

Mapping ATAC-seq data

ATAC-seq data were processed following the ENCODE ATAC-seq pipeline (version 1.8.0)

as follows: Raw reads were trimmed using cutadapt (version 2.5) and aligned to human

genome assembly GRCh38 analysis set using bowtie2 (version 2.3.4.3) for each sample.

After alignment, duplicated reads were removed using MarkDuplicates (Picard version

2.20.7-SNAPSHOT), retaining uniquely mapped reads and improperly paired reads using

samtools (version 1.9), and removed reads that mapped to the mitochondrial genome.

In total, we obtained a median of 5.39M (s.d. 3.28M) QCed reads per sample.

ATAC-seq peak calling

For each 29 immune cell type, we used the samtools merge command to merge the QCed

reads for each cell type and used MACS2 (version 2.1.1) to identify the ATAC-seq peaks

(OCRs) following a recent study (Gate, Cheng et al. 2018) with parameters (-p 0.01 --shift -

75 –extsize 150 --nomodel). We obtained a median of 253,553 (±66,738) OCRs with a

median peak width of 271bp ($\pm$18bp) (FDR<0.05). Using the alignment file for each cell

type, we calculated the read counts from the first mates for each OCR with bedtools

(version 2.30.0)'s coverage function.

Quality control for ATAC-seq dataset

We first assessed the quality of ATAC-seq data using Pearson correlations of chromatin

accessibility among the samples for each cell type, using 614,516 peaks that were

identified from across all samples using the same parameters described above (these

peaks were only used for quality control purposes). We filtered OCRs out when the

Pearson correlations were not greater than 0.65 (determined by manual inspection of the

distribution of these correlations). Additionally, we excluded outlier samples with low

Pearson correlations by visual inspection (Supplementary Table 16). Samples with a

mapped rate of less than 0.5 were further excluded. Finally, we obtained 1,386 samples

(the median number of donors per cell type is 48; Supplementary Table 1).

443   **Hierarchical clustering and dimension reduction**

444   We first chose the top 5,000 accessible OCRs (defined by TPM) from each cell type and

445   merged them across 29 cell types, resulting in a total of 145,000 unique regions. We then

446   created a binary matrix (0, 1) to indicate whether each 300bp bin across the genome was

447   overlapped by the corresponding OCRs. Hierarchical clustering was performed using

448   Ward's method of Euclidean distance using the binary matrix. PCA for the binary matrix

449   was performed using the binaryPCA function from the scBFA package (version 1.16.0) (Li

450   and Quon 2019) in R with default parameters. UMAP was performed with the top five PCs

451   using the umap function from the scikit-learn package (version 0.18.1) in Python with

452   default parameters.

453   **Strategy for the CAMBUS ML model**

454   The CAMBUS ML model was built upon the DeepSEA architecture, a deep CNN model

455   (Zhou and Troyanskaya 2015). We used the Selene library (v0.4.7) to train the CAMBUS

456   ML model. A 5kb reference sequence, randomly chosen with Selene, is used as input

457   features, five times longer than the sequence used in the original DeepSEA model to

458   capture broader genomic contexts. The model outputs a sigmoid function-derived score,

459   which can be interpreted as a proxy for the likelihood that the OCRs overlap with the

460   central 200bp of the input sequence (see Supplementary Table 17 for the configuration

461   details). We referred to this predicted probability for chromatin accessibility as the

462   accessibility probability.

463   We combined 7,939,714 OCRs across 29 cell types using the `bedtools merge` function

464   and obtained 1,349,934 non-overlapping unique input sequences. According to Selene's

465   default settings, we used chromosomes 8 and 9 as testing, chromosomes 6 and 7 as

validation, and the other autosomal chromosomes as training data. The details of Selene's
parameters are described in Supplementary Table 17.

***In silico* saturated mutagenesis using the trained CAMBUS ML model**

For all bases within 1kb from the center of each OCR, we used the trained CAMBUS ML
model to calculate the predictions for all possible single-based mutations. This
comprehensive approach, inspired by the previous study working on transcript expression
levels (Zhou, Theesfeld et al. 2018), provides an unbiased analysis of mutational effects
across the entire sequence, rather than focusing on only specific known polymorphisms. To
quantify the mutation effect, we calculated the difference in log fold changes of predictions
between the reference and alternative alleles for each site and averaged them across all
bases. In total, we obtained 505 million mutation effects (median across the 29 cell types).

Identification of putative constraint open chromatin region

We modeled the distribution of the mean mutation effect (*in silico* saturated mutagenesis)
within 2kb of each OCR (CAMBUS) as a mixture of a Gaussian null distribution. We
hypothesized that the OCRs with negative CAMBUS are likely positive constraint regions
and OCRs with neither negative nor positive CAMBUS as control OCRs. We calculated
local FDR with the locfdr function in the locfdr R package (version 1.1-8) for each cell type.
This analysis identified a median of 1,584 putative positive constraint OCRs across 29 cell
types with FDR < 0.5, referred to as CAMBUS+ OCRs, and a median of 248,749 control
OCRs across 29 cell types (Zhou, Theesfeld et al. 2018). We also defined relaxed
CAMBUS+ OCRs by extracting the bottom 5% of the CAMBUS distribution for each cell
type (Supplementary Table 19).

Enrichment of CAMBUS+ OCRs in constraint regions

We evaluated whether CAMBUS+ OCRs are enriched in constraint regions of the genome

using Gnocchi. We used the 1,984,900 QCed Gnocchi (see URLs) and calculated the OR

following the methodology described in the previous paper (Chen, Francioli et al. 2024) for

1,843,559 non-coding 1-kb windows. In brief, we counted the number of overlapping 1kb

windows within the following bins: (∞, -4), [−4,−3), [−3,−2), [−2,−1), [−1,−0), [0,1), [1,2), [2,3),

[3,4), [4,∞) with each CAMBUS+ OCR. ORs were calculated by comparing these overlaps

with all 1kb windows. Fisher's exact test was conducted to estimate the statistical

significance (p-value). The OR of the known regulatory elements for each bin was obtained

from the gnomAD download site (see URLs) and used for the comparison.

To calculate the overlapping of the constraint metrics in CAMBUS+ OCRs, we used the

seven constraint metrics (see URLs): (1) The Gnocchi, estimated by comparing observed

allele counts in diverse human populations with the estimated counts in each 1kb region of

the human genome (Chen, Francioli et al. 2024), (2) phyloP, defined by phylogenetic P-

values (phyloP) scores derived from conservation across various species: phyloP46way

(Pollard, Hubisz et al. 2010), phyloP100way, phyloP241way (Zoonomia 2020), based on

alignments of 46, 100, and 241 species, respectively, (3) Primate-specific PhastCons

(phyloPPrimates), defined by phylogenetic hidden Markov model (phylo-HMM) across 100

primate species (Zoonomia 2020), (4) Depletion Rank (DR), estimated by comparing

observed allele counts in EUR populations with the estimated counts in each 500bp region

of the human genome (Halldorsson, Eggertsson et al. 2022), (5) Genomic Evolutionary

Rate Profiling (GERP), defined by rejected substitution scores (RS) that quantify

evolutionary constraint using alignments of diverse mammalian genomes (Cooper, Stone et

al. 2005). The phyloP46way mapped the positions of those datasets to the hg38/GRCh38

coordinate using liftOver as described previously. For the Gnocchi and DR, the overlapping

regions were considered to be overlapped if they had more than 50% overlap with each

other.

To determine whether CAMBUS+ OCRs were prioritized by existing constraint metrics, we

employed the following approach: First, we calculated the mean values for each of the

seven constraint metrics overlapping for each CAMBUS+ OCR. Second, a CAMBUS+ OCR

was considered not prioritized by existing constraint metrics if all seven mean values failed

to meet their recommended thresholds; Gnocchi > 4 (Chen, Francioli et al. 2024),

phyloP100way with score > 3.0075 (FDR5%, following the previous approach (Sullivan,

Meadows et al. 2023)), phyloPPrimates with score > 0.961 (Sullivan, Meadows et al. 2023),

cactus241way > 2.27 (Sullivan, Meadows et al. 2023), DR < 0.01 (Halldorsson, Eggertsson

et al. 2022), GERP > 2 (a RS score threshold of 2 provides high sensitivity while still

strongly enriching for truly constrained sites, as described in the UCSC Genome Browser

documentation at 2025/7/22) (Fu, Liu et al. 2014)). Regarding the phyloP46way, we used

top10% score (0.533) as a threshold since even the maximum score (0.655)is not

nominally significant (corresponding P value is 0.221) due to lack of power since the

number of species is small (Pollard, Hubisz et al. 2010).

Enrichment of CAMBUS+ OCRs in the positive selection signals

We evaluated whether CAMBUS+ OCRs are enriched in the positive selection signals

previously reported (Nait Saada, Kalantzis et al. 2020). For each CAMBUS+ OCR, we

counted overlapping annotations for the positive selection signals observed in the EUR

population and estimated the OR compared with all OCRs using a logistic regression model.

To perform enrichment analysis of HAQERs (Mangan, Alsina et al. 2022), we first created

merged OCRs by combining CAMBUS+ OCRs from 29 different cell types (merged

CAMBUS+ OCRs). The number of overlapped merged CAMBUS+ OCRs in the HAQERs

was counted, and OR was estimated compared with all OCRs using a logistic regression

model.

Also, we evaluated whether CAMBUS+ OCRs are enriched in the archaic genome. For

each CAMBUS+ OCR, we counted overlapping annotations for the Neanderthal genome

sequences observed in the Japanese population (Liu, Koyama et al. 2024) and the

Neanderthal and Denisovans' sequences observed in 1000 Genomes populations

(Browning, Browning et al. 2018), and estimated the OR compared with all OCRs using a

logistic regression model.

Correlation between CAMBUS and regulatory elements

We evaluated the functional enrichment of CAMBUS+ OCRs in known regulatory elements,

including ENCODE cCREs, FANTOM5 enhancers, and super-enhancers (see URLs). For

each CAMBUS+ OCR, we counted overlapping annotations for each known regulatory

element and calculated the OR compared with all OCRs, and tested it using Fisher's exact

test, as described in the previous section. For tissue-expressed FANTOM5 enhancers, we

obtained count per million (CPM) data for 249,599 enhancers across 347 tissue/cell types

with hg19 coordinates (see URLs) and selected enhancers expressed in hematopoietic_cell

(N = 163), brain (N = 85), kidney (N = 16), and lung (N = 8) based on a threshold of CPM

>= 0.1 in at least 10% of individuals in each enhancer. As described previously, the

positions were mapped to the hg38/GRCh38 coordinate using liftOver. For tissue-

expressed super-enhancers, we obtained Blood (N = 115), brain (N = 203), kidney (N = 34),

and lung (N = 119) enhancers with hg38 coordinates (see URLs) and merged these

enhancers by tissue with `bedtools merge` function. The adjusted OR with Gnocchi,

phyloP46way, phyloP100way, cactus241way, phyloPPrimates, GERP, and DR was

estimated using a logistic regression model, adjusting for the average of these seven

constraint metrics overlapping CAMBUS+ OCRs and control CAMBUS OCRs.

For all analyses, regions were considered to be overlapped if they had more than 50%

overlap with each other.

Heritability enrichment analysis

To estimate heritability enrichment in each OCR, we conducted stratified-linkage

disequilibrium score regression (S-LDSC) with LDSC (ver.1.0.1) using the baseline model

(version 1.2; 53 categories). The p-value of the coefficient for CAMBUS+ OCR (or relaxed

CAMBUS+ OCR) was obtained by adjusting for the 53 categories in the baseline model

and all OCRs in each cell type. To avoid false positives (Tashman, Cui et al. 2021), we

used the relaxed CAMBUS+ OCRs (the median proportion of single nucleotide

polymorphism (SNPs) across the 29 cell types was 0.5%) for primary analyses instead of

the CAMBUS+ OCRs (the median was 0.07%). Multiple testing correction across all trait-

cell type pairs was performed to estimate FDR using the Storey method (Storey 2002). The

summary statistics were obtained from Alkes Price Lab (See URLs and our previous study

(Oguchi, Suzuki et al. 2024)), and five GWAS results in EAS ancestry: systemic lupus

erythematosus (Yin, Kim et al. 2021), rheumatoid arthritis (Ishigaki, Sakaue et al. 2022),

Systemic sclerosis (Ishikawa, Tanaka et al. 2024), Atopic dermatitis (Tanaka, Koido et al.

2021), and Takayasu arteritis (Terao, Yoshifuji et al. 2018). For this analysis, CAMBUS

OCRs' positions were mapped to the hg19 coordinate using the UCSC liftOver tool with

hg38ToHg19.over.chain file to be consistent with the summary statistics. After mapping to

hg19, regions mapped to different chromosomes were excluded from the analysis. For the

other constraint metrics, we extracted the strong constraints region: Gnocchi Z > 4,

phyloP46way score > 0.655, phyloP100way score > 9, phyloPPrimates score > 0.999,

cactus241way score > 8.8, GERP RS > 5, DR rank <= 0.005. These thresholds were

adjusted so that the proportion of SNPs matched that of Gnocchi (1.18%) because using

the recommended thresholds (FDR 5% for phyloP (Sullivan, Meadows et al. 2023), 0.961

for phyloPPrimates (Sullivan, Meadows et al. 2023), 2 for GERP (a RS score threshold of 2

provides high sensitivity while still strongly enriching for truly constrained sites, as

described in the UCSC Genome Browser documentation at 2025/7/22) (Fu, Liu et al. 2014),

0.01 for DR (Halldorsson, Eggertsson et al. 2022), cactus241way with score > 2.27 (FDR

5%)) resulted in a proportion of SNPs that was too large, leading to lower resolution to

assess heritability enrichment. For example, the recommended parameter for

cactus241way captured 66% of the proportion of SNPs and yielded only modest SNP

heritability enrichment (e.g., 1.37 for SLE), while the threshold that we used (8.8)

substantially improved the enrichment (7.32 for SLE).

PRS analysis

The PRS was constructed using PLINK v1.9, following a systematic approach with p-value

thresholds using SLE GWAS in EAS ancestry (1,178 cases and 136,442 controls, in total,

137,620 individuals) (Yin, Kim et al. 2021). To select relevant variants for the PRS, we

selected SNPs that met 20 types of p-value thresholds: 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3,

0.2, 0.1, 0.05, 0.01, 0.005, 5.0×10^{-4}, 5.0×10^{-5}, 1.0×10^{-5}, 5.0×10^{-6}, 1.0×10^{-7},

5.0×10^{-7} and 5.0×10^{-8}. After selecting SNPs based on these thresholds, we applied

LD clumping to remove SNPs in high linkage disequilibrium (LD) using nine types of r²

thresholds (0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9) and a 250kb physical distance. This

process yielded 180 PRS models. To determine the optimal PRS model, we applied a 5-

fold cross-validation strategy. The dataset was divided into five folds, with four folds used

for training and one for testing in each iteration. We computed PRSs for all 180 models

using variants from SLE GWAS in EAS ancestry for each training set. We calculated the

AUC with pROC (v1.18.4) to select the combination of p-value and r² parameters that

yielded the best AUC for each training set. We then took the two variant sets; (a) variants

filtered with the best parameters, and (b) all variants overlapping with DN B CAMBUS+

OCRs as prioritized variants. To mitigate overfitting due to LD structures, we removed the

variants with LD>0.7 with variants in set (b). We then calculated the PRS for both sets (a)

and (b) for each of the five test sets, and estimated the effect size of PRS from set (b),

controlling for the PRS from set (a) using a logistic regression model. Finally, we performed

a fixed-effects meta-analysis to evaluate the improvement in predictive power with the

prioritized variants.

Fine-mapping enrichment analysis

We downloaded the UKB fine-mapped variants of 94 complex diseases and traits from the

Finucane lab (see URLs) and obtained BBJ fine-mapped variants of 62 quantitative traits

from a recent study (Koyama, Liu et al. 2024). Regarding the UKB fine-mapped variants,

we used PIP from the SuSiE. The positions of those datasets were mapped to the

hg38/GRCh38 coordinate using liftOver as described previously. In total, we obtained

830,654 trait-variant pairs for UKB and 544,745 trait-variant pairs for BBJ. The number of

overlapped variants for each trait in 29 cell types of CAMBUS+ OCRs was counted, and the

OR was estimated using all OCRs in each 29-cell type as the background, and the P-value

was calculated from the logistic regression model. After Bonferroni correction (P=0.05

divided by the number of tested phenotypes), the median of OR is calculated from the

significant enrichment in UKB and BBJ (Extended Data Fig. 4A, B; Supplementary Table

20).

We categorized each trait into hematopoietic and non-hematopoietic. Furthermore,

hematopoietic traits are categorized into leukocyte and non-leukocyte traits. The

classification of traits and the number of putative causal variants in each category are

described in Supplementary Table 21.

Then, we counted the overlap of variants for each leukocyte and non-leukocyte and non-

hematopoietic for each 29 cell-type CAMBUS+ OCRs. We calculated the OR using the

method described above.

To perform enrichment analysis of rare variants, we first created merged OCRs by

combining CAMBUS+ OCRs from 29 different cell types (merged CAMBUS+ OCRs). The

number of overlapped variants with MAF < 0.01 and PIP > 0.1 in merged CAMBUS+ OCRs

was counted for all traits and only hematopoietic traits, and the OR was calculated using

the variants with MAF < 0.01 and PIP <= 0.1 as the background and calculated one-sided

Fisher's exact P-value.

Catalog cell-type-specific putative causal variants

We chose variants as putative causal variants from each of the credible sets overlapping

only one variant with CAMBUS+ OCRs. Then, we filtered the variants that overlapped with

only one cell type of CAMBUS+ OCRs. In total, we obtained 186 and 62 cell-type-specific

putative causal variants. Furthermore, we chose the variants that overlapped only one

subgroup (one of the CD4, CD8, B, Mono, DC, NK, Neutrophil; Supplementary Table 1).

Finally, we obtained 1,210 and 323 for trait-related cell-type-specific putative causal

variants from both EUR and EAS ancestries, respectively.

Statistical analysis

For p-value calculation with R, in case of insufficient precision with standard floating-point

arithmetic, the Rmpfr package (ver 0.9.2) was used to handle high-precision floating-point

arithmetic to calculate p-value with z-score. For the meta-analysis, the pooled OR was

calculated using a random effect model with metafor R package (version 2.0-0).

Data availability

The CAMBUS will be publicly available online soon after acceptance.

Code availability

The CAMBUS and its calculation scripts will be publicly available online soon after

acceptance.

URLs

Gnocchi is at https://gnomad.broadinstitute.org/downloads#v3-genomic-constraint.

Enrichment for Gnocchi with the known cCREs is at gs://gnomad-nc-constraint-v31-

paper/fig_tables/constraint_z_genome_1kb.annot.txt. ENCODE cCREs V3 is at

https://screen.encodeproject.org/. FANTOM5 enhancer is at

https://zenodo.org/record/556775/. Super enhancer is at

https://bio.liclab.net/sedb/download.php. phylop46way is at

https://hgdownload.cse.ucsc.edu/goldenPath/hg19/phyloP46way/primates. phyloP100way

is at

https://hgdownload.cse.ucsc.edu/goldenpath/hg38/phyloP100way/hg38.phyloP100way.bw

phyloPPrimates is at https://cgl.gi.ucsc.edu/data/cactus/zoonomia-2021-track-

hub/hg38/phyloPPrimates.bigWig. cactus241way is at

https://hgdownload.soe.ucsc.edu/goldenPath/hg38/cactus241way/cactus241way.phyloP.b

w. GERP is at

ftp.ensembl.org/pub/current_compara/conservation_scores/91_mammals.gerp_conservatio

n_score/gerp_conservation_scores.homo_sapiens.GRCh38.bw. DR is at https://static-

content.springer.com/esm/art%3A10.1038%2Fs41586-022-04965-

x/MediaObjects/41586_2022_4965_MOESM3_ESM.gz.GWAS summary statistics are at

https://alkesgroup.broadinstitute.org/sumstats_formatted, and

https://alkesgroup.broadinstitute.org/UKBB/. ABC predictions are at

https://mitra.stanford.edu/engreitz/oak/public/Nasser2021/AllPredictions.AvgHiC.ABC0.015.

minus150.ForABCPaperV3.txt.gz. UKB fine-mapped variants are at

https://www.finucanelab.org/data. FANTOM5 tissue-expressed data is at

https://zenodo.org/records/5348471.

Acknowledgment

We would like to thank all the participants in the study. We deeply thank all the members of
the Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative
Medical Sciences for the technical support. We acknowledge the use of Biorender for
aspects of the figures. This work was supported by Japan Agency for Medical Research
and Development (AMED) grants 21ek0109555, 21tm0424220, 21ck0106642,
23ek0410114, and 23tm0424225. Japan Society for the Promotion of Science (JSPS)
KAKENHI grant JP20H00462 and JP23K14164, Takeda Hosho Grants for Research in
Medicine.

Author contributions

K.T., M.K., and C.T. conceived the study design. A.S. performed ATAC-sequencing and

DNA genotyping. K.T. and N.T. performed whole genome imputation. K.T. analyzed this

study with the help of S.Y., N.T., Y.I., X.L., S.K., and M.K. M.K. and C.T. supervised this

study. S.Y., N.T., S.K., K.I., Y.M., Immune transcript/enhancer consortium, and K.Y.

provided helpful comments and discussion. K.T., M.K., A.S., and C.T. wrote the manuscript.

Ethics declarations

Competing interests

The authors declare no competing financial interests.

Figure legends

Fig. 1 | Study design. Overview of the study design. **A,** Dataset comprising 29 cell types

was used to validate our approach, and a machine learning model was developed to

predict OCRs. **B,C,** Identification of putative constraint OCRs (CAMBUS+). **D,**

Characteristics of CAMBUS+ OCRs based on CAMBUS. **E,** Characterization of CAMBUS+

OCRs and prioritization for putative causal variants.

Fig. 2 | CAMBUS+ OCRs can mark constraint regions and active enhancers. A,

Distribution of CAMBUS. The red part represents putative constraint OCRs. **B,** Genome

browser view at representative constraint region around CAMBUS+ in monocytes

(chr3:141,217,899-141,218,844). The black box represents Gnocchi with Z > 4, and the

grey box represents Gnocchi with Z < 4. **C,** Distribution of non-coding Gnocchi in

CAMBUS+ OCRs and CAMBUS control in NaiveCD4. The one-sided Wilcoxon-tests were

performed between CAMBUS+ OCRs and CAMBUS+ control. **D,E,** Enrichment (odds ratio;

OR) of CAMBUS+ in non-coding Gnocchi (n = 1,843,599), comparing with the known

regulatory elements (using the previous study's results, see URLs) in non-coding Gnocchi

(n = 1,843,599). We calculated the OR of CAMBUS+ (n = 1,266) and CAMBUS control (n =

251,807) for NaiveCD4 using various Gnocchi's thresholds in (**D**) and OR for each

CAMBUS+ OCR in 29 cell types and their pooled OR through random effects meta-analysis

using Gnocchi > 4 in (**E**). The blue dashed line represents the OR of ENCODE cCRE-PLS

(**E**). **F,** Enrichment of CAMBUS+ OCRs for NaiveCD4 (n = 1,266) in the known cCREs

(ENCODE cCRE-PLS; n = 40,891, ENCODE cCRE-pELS; n = 172,027, ENCODE cCRE-

dELS; n = 789,200, ENCODE CTCF-only; n = 35,839, FANTOM5 enhancers; n = 63,280,

Super enhancers; n = 16,472), adjusted for Gnocchi, phyloP46way, phyloP100way,

cactus241way, phyloPPrimates, GERP and DR. **G,** Enrichment of CAMBUS+ for NaiveCD4

(n = 1,266) in the tissue expressed FANTOM5 enhancers (hematopoietic cells; n = 64,225,

Brain; n = 62,978, Kidney; n = 35,177, Lung; n = 26,630), and Super enhancers (Blood; n =

8,055, Brain; n = 9,372, Kidney; n = 4,712, Lung; n = 9,878). The dashed line represents no

enrichment (OR=1.0; **D, E, F, G**). CAMBUS control in NaiveCD4 (n = 251,807) is used as

background in (**F, G**). The error bars represent 95% confidence intervals, with only lower

bounds shown in **D** (See Supplementary Table 4 for higher bounds).

Fig. 3 | Heritability enrichment for immune-mediated diseases. A, The heritability enrichment of various GWAS traits (in total, 33 GWAS for EUR and EAS ancestries; Supplementary Table 13) in CAMBUS+ OCRs for each of 29 cell types. The heatmap colors indicate -log10-scale p-value of the coefficient for CAMBUS+ annotation in S-LDSC (coefficient P). **B,** The -log10-scale coefficient P of CAMBUS+ OCRs in double-negative B-cells (DN B) and seven known constraint metrics for SLE GWAS in EAS ancestry. '*' indicates significance (qvalue < 0.05) after FDR correction (**A, B**).

Fig. 4 | Prioritization of putative causal variants using CAMBUS+ OCRs. A,
Enrichment of putative causal variants in leukocyte (leuko), non-leukocytes (non_leuko),
hemato (leuko + non_leuko), and non-hematopoietic (non_hemato) traits in CAMBUS+
OCRs for Th1. Putative causal variants were obtained from statistical fine-mapping analysis
of many complex traits' GWAS results for EAS and EUR populations (Kanai, Ulirsch et al.
2021, Koyama, Liu et al. 2024). **B,** Forest plot showing the enrichment of CAMBUS+ OCRs
across 29 cell types and the pooled OR in highly confident putative causal variants (PIP >
0.9). Variants with PIP <= 0.1 were used as the background for OR calculation. **C,**
Enrichment of CAMBUS+ OCRs for any of 29 cell types in putative causal rare variants
(maf < 0.01 in EAS), evaluated by comparing the proportion of CAMBUS+ OCRs
overlapping with putative causal rare variants with PIP > 0.1. The left dot represents the OR
calculated using all the putative causal variants, while the right dot represents the OR
based only on the putative causal variants for hematopoietic phenotypes. The dashed line
represents no enrichment (OR=1.0). **D,E,** Representative examples of variants associated
with monocyte counts prioritized with CAMBUS+ in Mono (**D**) and asthma prioritized with
CAMBUS+ in regulatory T-cells (**E**) in putative causal variants in EUR ancestry and
surrounding genomic annotations. The navy boxes represent putative causal variants in
this credible set. The blue dashed line represents an enhancer-gene connection to the
nearest protein-coding gene predicted by the ABC model for $CD14^+$ monocytes (**D**) and
HUVEC (**E**). The gene with the highest ABC score is also shown in (**D**). All the other
CAMBUS+ OCRs around putative causal variants (rs13122305 (**D**), rs55646091 (**E**)) are
also shown in the figures (The purple box represents PBMC).

Extended Data Figure

Extended Data Fig. 1 | Clustering analysis and dimension reduction using PCA and

UMAP. A-C, (**A**) Hierarchical clustering, (**B**) PCA, and (**C**) UMAP of 1,386 samples. The

top 5,000 most accessible OCRs were used for these analyses. The colors indicate the

classification of samples into eight subgroups across 29 cell types. **D,** UMAP of 1,386

samples (**C**) colored by eleven types of CD4; other subgroups are colored with grey.

Extended Data Fig. 2 | CAMBUS+ can mark constraint regions and active enhancers.

A, Scatter plot of median TPM and CAMBUS in Naïve CD4. The vertical dashed line

represents the top 5% of the median TPM. **B,** The number of OCRs in coding and non-

coding regions for each CAMBUS+ OCRs and CAMBUS control. The regions of coding and

non-coding are referred to previous study (Chen, Francioli et al. 2024). **C,** The distribution

of non-coding Gnocchi (median) for each CAMBUS+ OCRs and CAMBUS control in each

29 cell type. **D,** OR of relaxed CAMBUS+ OCRs across 29 cell types in Gnocchi> 4 and

pooled OR estimated through random effect meta-analysis. The blue dashed line

represents the OR of ENCODE promoter-like elements. **E-I,** OR and the pooled OR of the

CAMBUS+ OCRs in UKB selection signals (**E**), Neanderthal genome sequences derived

from samples of EAS ancestry (**F**), pathogenic regulatory variants (**G**), super-enhancers (**H**)

and FANTOM5 enhancers (**I**). **J,** ORs of relaxed CAMBUS+ (NaiveCD4; n = 12,736) in cell-

type-specific regulatory elements: the tissue expressed FANTOM5 enhancers

(hematopoietic cells; n = 64,225, Brain; n = 62,978, Kidney; n = 35,177, Lung; n = 26,630),

and super-enhancers (Blood; n = 8,055, Brain; n = 9,372, Kidney; n = 4,712, Lung; n =

9,878). The dashed line represents no enrichment (OR=1.0; **D-J**)

Extended Data Fig. 3 | Heritability enrichment for immune-mediated diseases. A, The
-log10-scale coefficient P of CAMBUS+ OCRs in 29 cell types for SLE GWAS in EAS
ancestry. The dashed line indicates a coefficient P = 0.05. '*' indicates significance (qvalue
< 0.05) after FDR correction.

Extended Data Fig. 4 | Prioritization of putative causal variants using CAMBUS+
OCRs. A, Median OR of all variants in 95% credible sets in CAMBUS+ OCRs across 29
cell types. We used putative causal variants from statistical fine-mapping studies in EUR
and EAS ancestries (Kanai, Ulirsch et al. 2021, Koyama, Liu et al. 2024). The solid line
represents the median OR across 29 cell types. **B,** Enrichment of the putative causal
variants in leukocyte (leuko), non-leukocytes (non_leuko), and non-hematopoietic
(non_hemato) traits in CAMBUS+ OCRs across 29 cell types. Only cell types with P < 0.05
are shown (27 cell types in BBJ leuko; 28 in UKB non_hemato; 29 in BBJ non_leuko, BBJ
non_hemato, UKB leuko, and UKB non_leuko), and dots indicate -log10-scale p-value. t-

tests were performed between leuko and non_hemato traits, with significance levels

denoted as P <0.05 ('*') and P < 0.001 ('***').

Extended Data Fig. 5 | Prioritization of putative causal variants using CAMBUS+

OCRs. A,B, representative example of variants associated with Hemoglobin A1c level

prioritized with CAMBUS+ in plasma blast in putative causal variants in EUR ancestry (**A**)

and eosinophils prioritized with CAMBUS+ in regulatory T-cell (**B**) and their surrounding

genomic annotations. The navy boxes represent putative causal variants in the credible set.

The blue dashed line represents an enhancer-gene connection to the nearest protein-

coding gene predicted by the ABC model in B-cells (**A**). All the other CAMBUS+ OCRs

around putative causal variants (rs8009224 (**A**)) are also shown in the figures (The purple

box represents PBMC).

Extended Data Fig. 6 | Sorting strategy for immune cell subsets. A, Representative

FACS plots describing gating strategies for 27 cell types. See Supplementary Table 22 for

flow cytometry antibodies used in this study.

References

Andersson, R., C. Gebhard, I. Miguel-Escalada, I. Hoof, J. Bornholdt, M. Boyd, Y. Chen, X.
Zhao, C. Schmidl, T. Suzuki, E. Ntini, E. Arner, E. Valen, K. Li, L. Schwarzfischer, D. Glatz, J.
Raithel, B. Lilje, N. Rapin, F. O. Bagger, M. Jorgensen, P. R. Andersen, N. Bertin, O. Rackham,
A. M. Burroughs, J. K. Baillie, Y. Ishizu, Y. Shimizu, E. Furuhata, S. Maeda, Y. Negishi, C. J.
Mungall, T. F. Meehan, T. Lassmann, M. Itoh, H. Kawaji, N. Kondo, J. Kawai, A. Lennartsson, C.
O. Daub, P. Heutink, D. A. Hume, T. H. Jensen, H. Suzuki, Y. Hayashizaki, F. Muller, A. R. R.
Forrest, P. Carninci, M. Rehli and A. Sandelin (2014). "An atlas of active enhancers across
human cell types and tissues." <u>Nature</u> **507**(7493): 455-461.
Armanfar, M., A. Jafari, G. R. Dehghan and L. Abdizadeh (2015). "Effect of coenzyme Q10
supplementation on exercise-induced response of inflammatory indicators and blood lactate in
male runners." <u>Med J Islam Repub Iran</u> **29**: 202.

Barton, A. R., M. A. Sherman, R. E. Mukamel and P. R. Loh (2021). "Whole-exome imputation
within UK Biobank powers rare coding variant association and fine-mapping analyses." Nat
Genet **53**(8): 1260-1269.
Benaglio, P., J. Newsome, J. Y. Han, J. Chiou, A. Aylward, S. Corban, M. Miller, M. L. Okino, J.
Kaur, S. Preissl, D. U. Gorkin and K. J. Gaulton (2023). "Mapping genetic effects on cell type-
specific chromatin accessibility and annotating complex immune trait variants using single
nucleus ATAC-seq in peripheral blood." PLoS Genet **19**(6): e1010759.
Boyle, A. P., S. Davis, H. P. Shulha, P. Meltzer, E. H. Margulies, Z. Weng, T. S. Furey and G. E.
Crawford (2008). "High-resolution mapping and characterization of open chromatin across the
genome." Cell **132**(2): 311-322.
Brixi, G., M. G. Durrant, J. Ku, M. Poli, G. Brockman, D. Chang, G. A. Gonzalez, S. H. King, D.
B. Li, A. T. Merchant, M. Naghipourfar, E. Nguyen, C. Ricci-Tam, D. W. Romero, G. Sun, A.
Taghibakshi, A. Vorontsov, B. Yang, M. Deng, L. Gorton, N. Nguyen, N. K. Wang, E. Adams, S.
A. Baccus, S. Dillmann, S. Ermon, D. Guo, R. Ilango, K. Janik, A. X. Lu, R. Mehta, M. R. K.
Mofrad, M. Y. Ng, J. Pannu, C. Ré, J. C. Schmok, J. S. John, J. Sullivan, K. Zhu, G. Zynda, D.
Balsam, P. Collison, A. B. Costa, T. Hernandez-Boussard, E. Ho, M.-Y. Liu, T. McGrath, K.
Powell, D. P. Burke, H. Goodarzi, P. D. Hsu and B. L. Hie (2025). "Genome modeling and
design across all domains of life with Evo 2." bioRxiv: 2025.2002.2018.638918.
Browning, S. R., B. L. Browning, Y. Zhou, S. Tucci and J. M. Akey (2018). "Analysis of Human
Sequence Data Reveals Two Pulses of Archaic Denisovan Admixture." Cell **173**(1): 53-61 e59.
Buenrostro, J. D., P. G. Giresi, L. C. Zaba, H. Y. Chang and W. J. Greenleaf (2013).
"Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin,
DNA-binding proteins and nucleosome position." Nat Methods **10**(12): 1213-1218.
Chen, S., L. C. Francioli, J. K. Goodrich, R. L. Collins, M. Kanai, Q. Wang, J. Alfoldi, N. A. Watts,
C. Vittal, L. D. Gauthier, T. Poterba, M. W. Wilson, Y. Tarasova, W. Phu, R. Grant, M. T.
Yohannes, Z. Koenig, Y. Farjoun, E. Banks, S. Donnelly, S. Gabriel, N. Gupta, S. Ferriera, C.
Tolonen, S. Novod, L. Bergelson, D. Roazen, V. Ruano-Rubio, M. Covarrubias, C. Llanwarne, N.
Petrillo, G. Wade, T. Jeandet, R. Munshi, K. Tibbetts, C. Genome Aggregation Database, A.
O'Donnell-Luria, M. Solomonson, C. Seed, A. R. Martin, M. E. Talkowski, H. L. Rehm, M. J.
Daly, G. Tiao, B. M. Neale, D. G. MacArthur and K. J. Karczewski (2024). "A genomic
mutational constraint map using variation in 76,156 human genomes." Nature **625**(7993): 92-
100.
Chung, M. K. Y., L. Gong, D. L. Kwong, V. H. Lee, A. W. Lee, X. Y. Guan, N. W. Kam and W.
Dai (2023). "Functions of double-negative B cells in autoimmune diseases, infections, and
cancers." EMBO Mol Med **15**(9): e17341.
Claussnitzer, M., J. H. Cho, R. Collins, N. J. Cox, E. T. Dermitzakis, M. E. Hurles, S. Kathiresan,
E. E. Kenny, C. M. Lindgren, D. G. MacArthur, K. N. North, S. E. Plon, H. L. Rehm, N. Risch, C.
N. Rotimi, J. Shendure, N. Soranzo and M. I. McCarthy (2020). "A brief history of human
disease genetics." Nature **577**(7789): 179-189.
Consortium, E. P., J. E. Moore, M. J. Purcaro, H. E. Pratt, C. B. Epstein, N. Shoresh, J. Adrian,
T. Kawli, C. A. Davis, A. Dobin, R. Kaul, J. Halow, E. L. Van Nostrand, P. Freese, D. U. Gorkin,
Y. Shen, Y. He, M. Mackiewicz, F. Pauli-Behn, B. A. Williams, A. Mortazavi, C. A. Keller, X. O.
Zhang, S. I. Elhajjajy, J. Huey, D. E. Dickel, V. Snetkova, X. Wei, X. Wang, J. C. Rivera-Mulia, J.
Rozowsky, J. Zhang, S. B. Chhetri, J. Zhang, A. Victorsen, K. P. White, A. Visel, G. W. Yeo, C.
B. Burge, E. Lecuyer, D. M. Gilbert, J. Dekker, J. Rinn, E. M. Mendenhall, J. R. Ecker, M. Kellis,
R. J. Klein, W. S. Noble, A. Kundaje, R. Guigo, P. J. Farnham, J. M. Cherry, R. M. Myers, B.
Ren, B. R. Graveley, M. B. Gerstein, L. A. Pennacchio, M. P. Snyder, B. E. Bernstein, B. Wold,
R. C. Hardison, T. R. Gingeras, J. A. Stamatoyannopoulos and Z. Weng (2020). "Expanded
encyclopaedias of DNA elements in the human and mouse genomes." Nature **583**(7818): 699-
710.
Cooper, G. M., E. A. Stone, G. Asimenos, N. C. S. Program, E. D. Green, S. Batzoglou and A.
Sidow (2005). "Distribution and intensity of constraint in mammalian genomic sequence."
Genome Res **15**(7): 901-913.

Corces, M. R., A. E. Trevino, E. G. Hamilton, P. G. Greenside, N. A. Sinnott-Armstrong, S.
Vesuna, A. T. Satpathy, A. J. Rubin, K. S. Montine, B. Wu, A. Kathiria, S. W. Cho, M. R.
Mumbach, A. C. Carter, M. Kasowski, L. A. Orloff, V. I. Risca, A. Kundaje, P. A. Khavari, T. J.
Montine, W. J. Greenleaf and H. Y. Chang (2017). "An improved ATAC-seq protocol reduces
background and enables interrogation of frozen tissues." Nat Methods **14**(10): 959-962.
Eraslan, G., Z. Avsec, J. Gagneur and F. J. Theis (2019). "Deep learning: new computational
modelling techniques for genomics." Nat Rev Genet **20**(7): 389-403.
Fillatreau, S., B. Manfroi and T. Dorner (2021). "Toll-like receptor signalling in B cells during
systemic lupus erythematosus." Nat Rev Rheumatol **17**(2): 98-108.
Finucane, H. K., B. Bulik-Sullivan, A. Gusev, G. Trynka, Y. Reshef, P. R. Loh, V. Anttila, H. Xu,
C. Zang, K. Farh, S. Ripke, F. R. Day, C. ReproGen, C. Schizophrenia Working Group of the
Psychiatric Genomics, R. Consortium, S. Purcell, E. Stahl, S. Lindstrom, J. R. Perry, Y. Okada,
S. Raychaudhuri, M. J. Daly, N. Patterson, B. M. Neale and A. L. Price (2015). "Partitioning
heritability by functional annotation using genome-wide association summary statistics." Nat
Genet **47**(11): 1228-1235.
Franchini, L. F. and K. S. Pollard (2017). "Human evolution: the non-coding revolution." BMC
Biol **15**(1): 89.
Fu, Y., Z. Liu, S. Lou, J. Bedford, X. J. Mu, K. Y. Yip, E. Khurana and M. Gerstein (2014).
"FunSeq2: a framework for prioritizing noncoding regulatory variants in cancer." Genome Biol
15(10): 480.
Fulco, C. P., J. Nasser, T. R. Jones, G. Munson, D. T. Bergman, V. Subramanian, S. R.
Grossman, R. Anyoha, B. R. Doughty, T. A. Patwardhan, T. H. Nguyen, M. Kane, E. M. Perez,
N. C. Durand, C. A. Lareau, E. K. Stamenova, E. L. Aiden, E. S. Lander and J. M. Engreitz
(2019). "Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR
perturbations." Nat Genet **51**(12): 1664-1669.
Gate, R. E., C. S. Cheng, A. P. Aiden, A. Siba, M. Tabaka, D. Lituiev, I. Machol, M. G. Gordon,
M. Subramaniam, M. Shamim, K. L. Hougen, I. Wortman, S. C. Huang, N. C. Durand, T. Feng,
P. L. De Jager, H. Y. Chang, E. L. Aiden, C. Benoist, M. A. Beer, C. J. Ye and A. Regev (2018).
"Genetic determinants of co-accessible chromatin regions in activated T cells across humans."
Nat Genet **50**(8): 1140-1150.
Gupta, A., K. Weinand, A. Nathan, S. Sakaue, M. J. Zhang, R. A. S. L. E. P. Accelerating
Medicines Partnership, Network, L. Donlin, K. Wei, A. L. Price, T. Amariuta and S.
Raychaudhuri (2023). "Dynamic regulatory elements in single-cell multimodal data implicate key
immune cell states enriched for autoimmune disease heritability." Nat Genet **55**(12): 2200-2210.
Halldorsson, B. V., H. P. Eggertsson, K. H. S. Moore, H. Hauswedell, O. Eiriksson, M. O.
Ulfarsson, G. Palsson, M. T. Hardarson, A. Oddsson, B. O. Jensson, S. Kristmundsdottir, B. D.
Sigurpalsdottir, O. A. Stefansson, D. Beyter, G. Holley, V. Tragante, A. Gylfason, P. I. Olason, F.
Zink, M. Asgeirsdottir, S. T. Sverrisson, B. Sigurdsson, S. A. Gudjonsson, G. T. Sigurdsson, G.
H. Halldorsson, G. Sveinbjornsson, K. Norland, U. Styrkarsdottir, D. N. Magnusdottir, S.
Snorradottir, K. Kristinsson, E. Sobech, H. Jonsson, A. J. Geirsson, I. Olafsson, P. Jonsson, O.
B. Pedersen, C. Erikstrup, S. Brunak, S. R. Ostrowski, D. G. Consortium, G. Thorleifsson, F.
Jonsson, P. Melsted, I. Jonsdottir, T. Rafnar, H. Holm, H. Stefansson, J. Saemundsdottir, D. F.
Gudbjartsson, O. T. Magnusson, G. Masson, U. Thorsteinsdottir, A. Helgason, H. Jonsson, P.
Sulem and K. Stefansson (2022). "The sequences of 150,119 genomes in the UK Biobank."
Nature **607**(7920): 732-740.
Harb, H. and T. A. Chatila (2023). "Regulatory T-cells in asthma." Curr Opin Allergy Clin
Immunol **23**(2): 151-157.
Hnisz, D., B. J. Abraham, T. I. Lee, A. Lau, V. Saint-André, A. A. Sigova, H. A. Hoke and R. A.
Young (2013). "Super-enhancers in the control of cell identity and disease." Cell **155**(4): 934-
947.
Honda, T. and K. Kabashima (2020). "Reconciling innate and acquired immunity in atopic
dermatitis." J Allergy Clin Immunol **145**(4): 1136-1137.

Ishigaki, K., S. Sakaue, C. Terao, Y. Luo, K. Sonehara, K. Yamaguchi, T. Amariuta, C. L. Too, V.
A. Laufer, I. C. Scott, S. Viatte, M. Takahashi, K. Ohmura, A. Murasawa, M. Hashimoto, H. Ito,
M. Hammoudeh, S. A. Emadi, B. K. Masri, H. Halabi, H. Badsha, I. W. Uthman, X. Wu, L. Lin, T.
Li, D. Plant, A. Barton, G. Orozco, S. M. M. Verstappen, J. Bowes, A. J. MacGregor, S. Honda,
M. Koido, K. Tomizuka, Y. Kamatani, H. Tanaka, E. Tanaka, A. Suzuki, Y. Maeda, K.
Yamamoto, S. Miyawaki, G. Xie, J. Zhang, C. I. Amos, E. Keystone, G. Wolbink, I. van der
Horst-Bruinsma, J. Cui, K. P. Liao, R. J. Carroll, H. S. Lee, S. Y. Bang, K. A. Siminovitch, N. de
Vries, L. Alfredsson, S. Rantapaa-Dahlqvist, E. W. Karlson, S. C. Bae, R. P. Kimberly, J. C.
Edberg, X. Mariette, T. Huizinga, P. Dieude, M. Schneider, M. Kerick, J. C. Denny, P. BioBank
Japan, K. Matsuda, K. Matsuo, T. Mimori, F. Matsuda, K. Fujio, Y. Tanaka, A. Kumanogoh, M.
Traylor, C. M. Lewis, S. Eyre, H. Xu, R. Saxena, T. Arayssi, Y. Kochi, K. Ikari, M. Harigai, P. K.
Gregersen, K. Yamamoto, S. Louis Bridges, Jr., L. Padyukov, J. Martin, L. Klareskog, Y. Okada
and S. Raychaudhuri (2022). "Multi-ancestry genome-wide association analyses identify novel
genetic mechanisms in rheumatoid arthritis." Nat Genet **54**(11): 1640-1651.
Ishikawa, Y., N. Tanaka, Y. Asano, M. Kodera, Y. Shirai, M. Akahoshi, M. Hasegawa, T.
Matsushita, K. Saito, S. I. Motegi, H. Yoshifuji, A. Yoshizaki, T. Kohmoto, K. Takagi, A. Oka, M.
Kanda, Y. Tanaka, Y. Ito, K. Nakano, H. Kasamatsu, A. Utsunomiya, A. Sekiguchi, H. Niiro, M.
Jinnin, K. Makino, T. Makino, H. Ihn, M. Yamamoto, C. Suzuki, H. Takahashi, E. Nishida, A.
Morita, T. Yamamoto, M. Fujimoto, Y. Kondo, D. Goto, T. Sumida, N. Ayuzawa, H. Yanagida, T.
Horita, T. Atsumi, H. Endo, Y. Shima, A. Kumanogoh, J. Hirata, N. Otomo, H. Suetsugu, Y.
Koike, K. Tomizuka, S. Yoshino, X. Liu, S. Ito, K. Hikino, A. Suzuki, Y. Momozawa, S. Ikegawa,
Y. Tanaka, O. Ishikawa, K. Takehara, T. Torii, S. Sato, Y. Okada, T. Mimori, F. Matsuda, K.
Matsuda, T. Amariuta, I. Imoto, K. Matsuo, M. Kuwana, Y. Kawaguchi, K. Ohmura and C. Terao
(2024). "GWAS for systemic sclerosis identifies six novel susceptibility loci including one in the
Fcgamma receptor region." Nat Commun **15**(1): 319.
Jarvinen, T. M., A. Hellquist, M. Zucchelli, S. Koskenmies, J. Panelius, T. Hasan, H. Julkunen,
M. D'Amato and J. Kere (2012). "Replication of GWAS-identified systemic lupus erythematosus
susceptibility genes affirms B-cell receptor pathway signalling and strengthens the role of IRF5
in disease susceptibility in a Northern European population." Rheumatology (Oxford) **51**(1): 87-
92.
Jiang, Y., F. Qian, X. Bai, Y. Liu, Q. Wang, B. Ai, X. Han, S. Shi, J. Zhang, X. Li, Z. Tang, Q.
Pan, Y. Wang, F. Wang and C. Li (2019). "SEdb: a comprehensive human super-enhancer
database." Nucleic Acids Res **47**(D1): D235-D243.
Kanai, M., J. C. Ulirsch, J. Karjalainen, M. Kurki, K. J. Karczewski, E. Fauman, Q. S. Wang, H.
Jacobs, F. Aguet, K. G. Ardlie, N. Kerimov, K. Alasoo, C. Benner, K. Ishigaki, S. Sakaue, S.
Reilly, T. B. J. Project, FinnGen, Y. Kamatani, K. Matsuda, A. Palotie, B. M. Neale, R. Tewhey,
P. C. Sabeti, Y. Okada, M. J. Daly and H. K. Finucane (2021). "Insights from complex trait fine-
mapping across diverse populations." medRxiv: 2021.2009.2003.21262975.
Karczewski, K. J., L. C. Francioli, G. Tiao, B. B. Cummings, J. Alfoldi, Q. Wang, R. L. Collins, K.
M. Laricchia, A. Ganna, D. P. Birnbaum, L. D. Gauthier, H. Brand, M. Solomonson, N. A. Watts,
D. Rhodes, M. Singer-Berk, E. M. England, E. G. Seaby, J. A. Kosmicki, R. K. Walters, K.
Tashman, Y. Farjoun, E. Banks, T. Poterba, A. Wang, C. Seed, N. Whiffin, J. X. Chong, K. E.
Samocha, E. Pierce-Hoffman, Z. Zappala, A. H. O'Donnell-Luria, E. V. Minikel, B. Weisburd, M.
Lek, J. S. Ware, C. Vittal, I. M. Armean, L. Bergelson, K. Cibulskis, K. M. Connolly, M.
Covarrubias, S. Donnelly, S. Ferriera, S. Gabriel, J. Gentry, N. Gupta, T. Jeandet, D. Kaplan, C.
Llanwarne, R. Munshi, S. Novod, N. Petrillo, D. Roazen, V. Ruano-Rubio, A. Saltzman, M.
Schleicher, J. Soto, K. Tibbetts, C. Tolonen, G. Wade, M. E. Talkowski, C. Genome Aggregation
Database, B. M. Neale, M. J. Daly and D. G. MacArthur (2020). "The mutational constraint
spectrum quantified from variation in 141,456 humans." Nature **581**(7809): 434-443.
Klemm, S. L., Z. Shipony and W. J. Greenleaf (2019). "Chromatin accessibility and the
regulatory epigenome." Nat Rev Genet **20**(4): 207-220.
Koido, M., C. C. Hon, S. Koyama, H. Kawaji, Y. Murakawa, K. Ishigaki, K. Ito, J. Sese, N. F.
Parrish, Y. Kamatani, P. Carninci and C. Terao (2023). "Prediction of the cell-type-specific

transcription of non-coding RNAs from genome sequences via machine learning." <u>Nat Biomed Eng</u> **7**(6): 830-844.

Koyama, S., X. Liu, Y. Koike, K. Hikino, M. Koido, W. Li, K. Akaki, K. Tomizuka, S. Ito, N. Otomo, H. Suetsugu, S. Yoshino, M. Akiyama, K. Saito, Y. Ishikawa, C. Benner, P. Natarajan, P. T. Ellinor, T. Mushiroda, M. Horikoshi, M. Ikeda, N. Iwata, K. Matsuda, K. Matsuda, S. Niida, K. Ozaki, Y. Momozawa, S. Ikegawa, O. Takeuchi, K. Ito, C. Terao and P. Biobank Japan (2024). "Population-specific putative causal variants shape quantitative traits." <u>Nature Genetics</u>.

Kuderna, L. F. K., J. C. Ulirsch, S. Rashid, M. Ameen, L. Sundaram, G. Hickey, A. J. Cox, H. Gao, A. Kumar, F. Aguet, M. J. Christmas, H. Clawson, M. Haeussler, M. C. Janiak, M. Kuhlwilm, J. D. Orkin, T. Bataillon, S. Manu, A. Valenzuela, J. Bergman, M. Rouselle, F. E. Silva, L. Agueda, J. Blanc, M. Gut, D. de Vries, I. Goodhead, R. A. Harris, M. Raveendran, A. Jensen, I. S. Chuma, J. E. Horvath, C. Hvilsom, D. Juan, P. Frandsen, J. G. Schraiber, F. R. de Melo, F. Bertuol, H. Byrne, I. Sampaio, I. Farias, J. Valsecchi, M. Messias, M. N. F. da Silva, M. Trivedi, R. Rossi, T. Hrbek, N. Andriaholinirina, C. J. Rabarivola, A. Zaramody, C. J. Jolly, J. Phillips-Conroy, G. Wilkerson, C. Abee, J. H. Simmons, E. Fernandez-Duque, S. Kanthaswamy, F. Shiferaw, D. Wu, L. Zhou, Y. Shao, G. Zhang, J. D. Keyyu, S. Knauf, M. D. Le, E. Lizano, S. Merker, A. Navarro, T. Nadler, C. C. Khor, J. Lee, P. Tan, W. K. Lim, A. C. Kitchener, D. Zinner, I. Gut, A. D. Melin, K. Guschanski, M. H. Schierup, R. M. D. Beck, I. Karakikes, K. C. Wang, G. Umapathy, C. Roos, J. P. Boubli, A. Siepel, A. Kundaje, B. Paten, K. Lindblad-Toh, J. Rogers, T. Marques Bonet and K. K. Farh (2024). "Identification of constrained sequence elements across 239 primate genomes." <u>Nature</u> **625**(7996): 735-742.

Kumasaka, N., A. J. Knights and D. J. Gaffney (2016). "Fine-mapping cellular QTLs with RASQUAL and ATAC-seq." <u>Nat Genet</u> **48**(2): 206-213.

Leypold, N. A. and M. R. Speicher (2021). "Evolutionary conservation in noncoding genomic regions." <u>Trends Genet</u> **37**(10): 903-918.

Li, R. and G. Quon (2019). "scBFA: modeling detection patterns to mitigate technical noise in large-scale single-cell genomics data." <u>Genome Biol</u> **20**(1): 193.

Li, Y., Z. Li and F. Hu (2021). "Double-negative (DN) B cells: an under-recognized effector memory B cell subset in autoimmunity." <u>Clin Exp Immunol</u> **205**(2): 119-127.

Li, Y. K., H. Y. Wang, Y. Chen, X. Q. Shi, X. X. Zhang, K. Li, W. P. Fu and C. Sun (2023). "Single‐nucleotide polymorphism rs6592645 confers asthma risk through regulating LRRC32 expression." <u>Exp Ther Med</u> **26**(3): 451.

Liu, X., S. Koyama, K. Tomizuka, S. Takata, Y. Ishikawa, S. Ito, S. Kosugi, K. Suzuki, K. Hikino, M. Koido, Y. Koike, M. Horikoshi, T. Gakuhari, S. Ikegawa, K. Matsuda, Y. Momozawa, K. Ito, Y. Kamatani and C. Terao (2024). "Decoding triancestral origins, archaic introgression, and natural selection in the Japanese population by whole-genome sequencing." <u>Sci Adv</u> **10**(16): eadi8419.

Mangan, R. J., F. C. Alsina, F. Mosti, J. E. Sotelo-Fonseca, D. A. Snellings, E. H. Au, J. Carvalho, L. Sathyan, G. D. Johnson, T. E. Reddy, D. L. Silver and C. B. Lowe (2022). "Adaptive sequence divergence forged new neurodevelopmental enhancers in humans." <u>Cell</u> **185**(24): 4587-4603 e4523.

Mantle, D., R. A. Heaton and I. P. Hargreaves (2021). "Coenzyme Q10 and Immune Function: An Overview." <u>Antioxidants (Basel)</u> **10**(5).

Martens, L. D., D. S. Fischer, V. A. Yepez, F. J. Theis and J. Gagneur (2024). "Modeling fragment counts improves single-cell ATAC-seq analysis." <u>Nat Methods</u> **21**(1): 28-31.

Maurano, M. T., R. Humbert, E. Rynes, R. E. Thurman, E. Haugen, H. Wang, A. P. Reynolds, R. Sandstrom, H. Qu, J. Brody, A. Shafer, F. Neri, K. Lee, T. Kutyavin, S. Stehling-Sun, A. K. Johnson, T. K. Canfield, E. Giste, M. Diegel, D. Bates, R. S. Hansen, S. Neph, P. J. Sabo, S. Heimfeld, A. Raubitschek, S. Ziegler, C. Cotsapas, N. Sotoodehnia, I. Glass, S. R. Sunyaev, R. Kaul and J. A. Stamatoyannopoulos (2012). "Systematic localization of common disease-associated variation in regulatory DNA." <u>Science</u> **337**(6099): 1190-1195.

Moser, T., K. Akgun, U. Proschmann, J. Sellner and T. Ziemssen (2020). "The role of TH17 cells in multiple sclerosis: Therapeutic implications." <u>Autoimmun Rev</u> **19**(10): 102647.

Mukherjee, R., P. Kanti Barman, P. Kumar Thatoi, R. Tripathy, B. Kumar Das and B. Ravindran
(2015). "Non-Classical monocytes display inflammatory features: Validation in Sepsis and
Systemic Lupus Erythematous." Sci Rep **5**: 13886.
Nait Saada, J., G. Kalantzis, D. Shyr, F. Cooper, M. Robinson, A. Gusev and P. F. Palamara
(2020). "Identity-by-descent detection across 487,409 British samples reveals fine scale
population structure and ultra-rare variant associations." Nat Commun **11**(1): 6130.
Nassir, N., R. Tambi, A. Bankapur, S. Al Heialy, N. Karuvantevida, H. H. Khansaheb, B. Zehra,
G. Begum, R. A. Hameid, A. Ahmed, Z. Deesi, A. Alkhajeh, K. M. F. Uddin, H. Akter, S. A.
Safizadeh Shabestari, O. Almidani, A. Islam, M. Gaudet, R. K. Kandasamy, T. Loney, A. A.
Tayoun, N. Nowotny, M. Woodbury-Smith, P. Rahman, W. M. Kuebler, M. Yaseen Hachim, J. L.
Casanova, B. K. Berdiev, A. Alsheikh-Ali and M. Uddin (2021). "Single-cell transcriptome
identifies FCGR3B upregulated subtype of alveolar macrophages in patients with critical
COVID-19." iScience **24**(9): 103030.
Nelson, R. K., A. Bush, J. Stokes, P. Nair and P. Akuthota (2020). "Eosinophilic Asthma." J
Allergy Clin Immunol Pract **8**(2): 465-473.
O'Bleness, M., V. B. Searles, A. Varki, P. Gagneux and J. M. Sikela (2012). "Evolution of
genetic and genomic features unique to the human lineage." Nat Rev Genet **13**(12): 853-866.
Oguchi, A., A. Suzuki, S. Komatsu, H. Yoshitomi, S. Bhagat, R. Son, R. J. P. Bonnal, S. Kojima,
M. Koido, K. Takeuchi, K. Myouzen, G. Inoue, T. Hirai, H. Sano, Y. Takegami, A. Kanemaru, I.
Yamaguchi, Y. Ishikawa, N. Tanaka, S. Hirabayashi, R. Konishi, S. Sekito, T. Inoue, J. Kere, S.
Takeda, A. Takaori-Kondo, I. Endo, S. Kawaoka, H. Kawaji, K. Ishigaki, H. Ueno, Y.
Hayashizaki, M. Pagani, P. Carninci, M. Yanagita, I. C. dagger, N. Parrish, C. Terao, K.
Yamamoto, Y. Murakawa and I. C. Members (2024). "An atlas of transcribed enhancers across
helper T cell diversity for decoding human diseases." Science **385**(6704): eadd8394.
Pollard, K. S., M. J. Hubisz, K. R. Rosenbloom and A. Siepel (2010). "Detection of nonneutral
substitution rates on mammalian phylogenies." Genome Res **20**(1): 110-121.
Pollen, A. A., U. Kilik, C. B. Lowe and J. G. Camp (2023). "Human-specific genetics: new tools
to explore the molecular and cellular basis of human evolution." Nat Rev Genet **24**(10): 687-711.
Radke, D. W., J. H. Sul, D. J. Balick, S. Akle, I. Alzheimer's Disease Neuroimaging, R. C. Green
and S. R. Sunyaev (2021). "Purifying selection on noncoding deletions of human regulatory loci
detected using their cellular pleiotropy." Genome Res **31**(6): 935-946.
Roadmap Epigenomics, C., A. Kundaje, W. Meuleman, J. Ernst, M. Bilenky, A. Yen, A. Heravi-
Moussavi, P. Kheradpour, Z. Zhang, J. Wang, M. J. Ziller, V. Amin, J. W. Whitaker, M. D.
Schultz, L. D. Ward, A. Sarkar, G. Quon, R. S. Sandstrom, M. L. Eaton, Y. C. Wu, A. R.
Pfenning, X. Wang, M. Claussnitzer, Y. Liu, C. Coarfa, R. A. Harris, N. Shoresh, C. B. Epstein,
E. Gjoneska, D. Leung, W. Xie, R. D. Hawkins, R. Lister, C. Hong, P. Gascard, A. J. Mungall, R.
Moore, E. Chuah, A. Tam, T. K. Canfield, R. S. Hansen, R. Kaul, P. J. Sabo, M. S. Bansal, A.
Carles, J. R. Dixon, K. H. Farh, S. Feizi, R. Karlic, A. R. Kim, A. Kulkarni, D. Li, R. Lowdon, G.
Elliott, T. R. Mercer, S. J. Neph, V. Onuchic, P. Polak, N. Rajagopal, P. Ray, R. C. Sallari, K. T.
Siebenthall, N. A. Sinnott-Armstrong, M. Stevens, R. E. Thurman, J. Wu, B. Zhang, X. Zhou, A.
E. Beaudet, L. A. Boyer, P. L. De Jager, P. J. Farnham, S. J. Fisher, D. Haussler, S. J. Jones,
W. Li, M. A. Marra, M. T. McManus, S. Sunyaev, J. A. Thomson, T. D. Tlsty, L. H. Tsai, W.
Wang, R. A. Waterland, M. Q. Zhang, L. H. Chadwick, B. E. Bernstein, J. F. Costello, J. R.
Ecker, M. Hirst, A. Meissner, A. Milosavljevic, B. Ren, J. A. Stamatoyannopoulos, T. Wang and
M. Kellis (2015). "Integrative analysis of 111 reference human epigenomes." Nature **518**(7539):
317-330.
Schaid, D. J., W. Chen and N. B. Larson (2018). "From genome-wide associations to candidate
causal variants by statistical fine-mapping." Nat Rev Genet **19**(8): 491-504.
Silvert, M., L. Quintana-Murci and M. Rotival (2019). "Impact and Evolutionary Determinants of
Neanderthal Introgression on Transcriptional and Post-Transcriptional Regulation." Am J Hum
Genet **104**(6): 1241-1250.
Soskic, B., E. Cano-Gamez, D. J. Smyth, W. C. Rowan, N. Nakic, J. Esparza-Gordillo, L.
Bossini-Castillo, D. F. Tough, C. G. C. Larminie, P. G. Bronson, D. Wille and G. Trynka (2019).

"Chromatin activity at GWAS loci identifies T cell states driving complex immune diseases." <u>Nat Genet</u> **51**(10): 1486-1493.

Stenson, P. D., E. V. Ball, M. Mort, A. D. Phillips, J. A. Shiel, N. S. Thomas, S. Abeysinghe, M. Krawczak and D. N. Cooper (2003). "Human Gene Mutation Database (HGMD): 2003 update." <u>Hum Mutat</u> **21**(6): 577-581.

Storey, J. D. (2002). "A Direct Approach to False Discovery Rates." <u>Journal of the Royal Statistical Society Series B: Statistical Methodology</u> **64**(3): 479-498.

Sullivan, P. F., J. R. S. Meadows, S. Gazal, B. N. Phan, X. Li, D. P. Genereux, M. X. Dong, M. Bianchi, G. Andrews, S. Sakthikumar, J. Nordin, A. Roy, M. J. Christmas, V. D. Marinescu, C. Wang, O. Wallerman, J. Xue, S. Yao, Q. Sun, J. Szatkiewicz, J. Wen, L. M. Huckins, A. Lawler, K. C. Keough, Z. Zheng, J. Zeng, N. R. Wray, Y. Li, J. Johnson, J. Chen, s. Zoonomia Consortium section, B. Paten, S. K. Reilly, G. M. Hughes, Z. Weng, K. S. Pollard, A. R. Pfenning, K. Forsberg-Nilsson, E. K. Karlsson and K. Lindblad-Toh (2023). "Leveraging base-pair mammalian constraint to understand genetic variation and human disease." <u>Science</u> **380**(6643): eabn2937.

Tanaka, N., M. Koido, A. Suzuki, N. Otomo, H. Suetsugu, Y. Kochi, K. Tomizuka, Y. Momozawa, Y. Kamatani, P. Biobank Japan, S. Ikegawa, K. Yamamoto and C. Terao (2021). "Eight novel susceptibility loci and putative causal variants in atopic dermatitis." <u>J Allergy Clin Immunol</u> **148**(5): 1293-1306.

Tashman, K. C., R. Cui, L. J. O'Connor, B. M. Neale and H. K. Finucane (2021). "Significance testing for small annotations in stratified LD-Score regression." <u>medRxiv</u>: 2021.003.2013.21249938.

Terao, C., H. Yoshifuji, T. Matsumura, T. K. Naruse, T. Ishii, Y. Nakaoka, Y. Kirino, K. Matsuo, T. Origuchi, M. Shimizu, Y. Maejima, E. Amiya, N. Tamura, T. Kawaguchi, M. Takahashi, K. Setoh, K. Ohmura, R. Watanabe, T. Horita, T. Atsumi, M. Matsukura, T. Miyata, Y. Kochi, T. Suda, K. Tanemoto, A. Meguro, Y. Okada, A. Ogimoto, M. Yamamoto, H. Takahashi, S. Nakayamada, K. Saito, M. Kuwana, N. Mizuki, Y. Tabara, A. Ueda, I. Komuro, A. Kimura, M. Isobe, T. Mimori and F. Matsuda (2018). "Genetic determinants and an epistasis of LILRA3 and HLA-B*52 in Takayasu arteritis." <u>Proc Natl Acad Sci U S A</u> **115**(51): 13045-13050.

Thibodeau, A., A. Uyar, S. Khetan, M. L. Stitzel and D. Ucar (2018). "A neural network based model effectively predicts enhancers from clinical ATAC-seq samples." <u>Sci Rep</u> **8**(1): 16048.

Tran, D. Q., J. Andersson, R. Wang, H. Ramsey, D. Unutmaz and E. M. Shevach (2009). "GARP (LRRC32) is essential for the surface expression of latent TGF-beta on platelets and activated FOXP3+ regulatory T cells." <u>Proc Natl Acad Sci U S A</u> **106**(32): 13445-13450.

Trynka, G., C. Sandor, B. Han, H. Xu, B. E. Stranger, X. S. Liu and S. Raychaudhuri (2013). "Chromatin marks identify critical cell types for fine mapping complex trait variants." <u>Nat Genet</u> **45**(2): 124-130.

Visscher, P. M., N. R. Wray, Q. Zhang, P. Sklar, M. I. McCarthy, M. A. Brown and J. Yang (2017). "10 Years of GWAS Discovery: Biology, Function, and Translation." <u>Am J Hum Genet</u> **101**(1): 5-22.

Ward, L. D. and M. Kellis (2012). "Evidence of abundant purifying selection in humans for recently acquired regulatory functions." <u>Science</u> **337**(6102): 1675-1678.

Yan, F., D. R. Powell, D. J. Curtis and N. C. Wong (2020). "From reads to insight: a hitchhiker's guide to ATAC-seq data analysis." <u>Genome Biol</u> **21**(1): 22.

Yin, X., K. Kim, H. Suetsugu, S. Y. Bang, L. Wen, M. Koido, E. Ha, L. Liu, Y. Sakamoto, S. Jo, R. X. Leng, N. Otomo, V. Laurynenka, Y. C. Kwon, Y. Sheng, N. Sugano, M. Y. Hwang, W. Li, M. Mukai, K. Yoon, M. Cai, K. Ishigaki, W. T. Chung, H. Huang, D. Takahashi, S. S. Lee, M. Wang, K. Karino, S. C. Shim, X. Zheng, T. Miyamura, Y. M. Kang, D. Ye, J. Nakamura, C. H. Suh, Y. Tang, G. Motomura, Y. B. Park, H. Ding, T. Kuroda, J. Y. Choe, C. Li, H. Niiro, Y. Park, C. Shen, T. Miyamoto, G. Y. Ahn, W. Fei, T. Takeuchi, J. M. Shin, K. Li, Y. Kawaguchi, Y. K. Lee, Y. Wang, K. Amano, D. J. Park, W. Yang, Y. Tada, K. Yamaji, M. Shimizu, T. Atsumi, A. Suzuki, T. Sumida, Y. Okada, K. Matsuda, K. Matsuo, Y. Kochi, H. Japanese Research Committee on Idiopathic Osteonecrosis of the Femoral, L. C. Kottyan, M. T. Weirauch, S. Parameswaran, S.

Eswar, H. Salim, X. Chen, K. Yamamoto, J. B. Harley, K. Ohmura, T. H. Kim, S. Yang, T. Yamamoto, B. J. Kim, N. Shen, S. Ikegawa, H. S. Lee, X. Zhang, C. Terao, Y. Cui and S. C. Bae (2021). "Meta-analysis of 208370 East Asians identifies 113 susceptibility loci for systemic lupus erythematosus." <u>Ann Rheum Dis</u> **80**(5): 632-640.

Zeng, B., J. Bendl, C. Deng, D. Lee, R. Misir, S. M. Reach, S. P. Kleopoulos, P. Auluck, S. Marenco, D. A. Lewis, V. Haroutunian, N. Ahituv, J. F. Fullard, G. E. Hoffman and P. Roussos (2024). "Genetic regulation of cell type-specific chromatin accessibility shapes brain disease etiology." <u>Science</u> **384**(6698): eadh4265.

Zhang, J. Y., X. M. Wang, X. Xing, Z. Xu, C. Zhang, J. W. Song, X. Fan, P. Xia, J. L. Fu, S. Y. Wang, R. N. Xu, X. P. Dai, L. Shi, L. Huang, T. J. Jiang, M. Shi, Y. Zhang, A. Zumla, M. Maeurer, F. Bai and F. S. Wang (2020). "Single-cell landscape of immunological responses in patients with COVID-19." <u>Nat Immunol</u> **21**(9): 1107-1118.

Zhang, K., J. D. Hocker, M. Miller, X. Hou, J. Chiou, O. B. Poirion, Y. Qiu, Y. E. Li, K. J. Gaulton, A. Wang, S. Preissl and B. Ren (2021). "A single-cell atlas of chromatin accessibility in the human genome." <u>Cell</u> **184**(24): 5985-6001 e5919.

Zhou, J., C. L. Theesfeld, K. Yao, K. M. Chen, A. K. Wong and O. G. Troyanskaya (2018). "Deep learning sequence-based ab initio prediction of variant effects on expression and disease risk." <u>Nat Genet</u> **50**(8): 1171-1179.

Zhou, J. and O. G. Troyanskaya (2015). "Predicting effects of noncoding variants with deep learning-based sequence model." <u>Nat Methods</u> **12**(10): 931-934.

Zoonomia, C. (2020). "A comparative genomics multitool for scientific discovery and conservation." <u>Nature</u> **587**(7833): 240-245.