COSMOS: COmmunity and Single Microbe Optimisation System

Lavanya Raajaraam[1,2] and Karthik Raman[1,2,3*]

[1]Bhupat and Jyoti Mehta School of Biosciences, Department of Biotechnology, Indian Institute of

Technology (IIT) Madras, Chennai – 600 036, INDIA

[2]Centre for Integrative Biology and Systems mEdicine (IBSE), Wadhwani School of Data Science

and AI, IIT Madras, Chennai – 600 036, INDIA

[3]Department of Data Science and AI, Wadhwani School of Data Science and AI, IIT Madras,

Chennai – 600 036, INDIA

*Corresponding author. E-mail: kraman@iitm.ac.in

ABSTRACT

Bioprocessing harnesses the potential of microbial communities and monocultures to convert

renewable resources like agricultural by-products and wastewater into valuable products.

While monocultures offer simplicity and control, microbial communities provide metabolic

diversity, enabling complex substrate breakdown and cooperative biosynthesis. However,

balancing efficiency with stability requires a nuanced approach, as each system presents unique

advantages and limitations, particularly under different environmental conditions. To

systematically explore the potential of these systems, we developed the COmmunity and Single

Microbe Optimisation System (COSMOS), a dynamic computational framework that compares

microbial monocultures and communities across diverse fermentation conditions. COSMOS

identifies optimal microbial systems tailored to specific carbon sources and environments. Our

findings highlight the impact of key factors such as environment, microbial interactions, and

carbon sources on the biosynthetic capabilities of both communities and monocultures. A key

result is the identification of the *Shewanella oneidensis* – *Klebsiella pneumoniae* community as

the most efficient producer of 1,3-propanediol under anaerobic conditions. Notably, this aligns

with previous experimental findings, with COSMOS accurately predicting the optimal carbon

source concentration and inoculum ratio used. Additional findings highlight the value of

communities for nutrient-limited processes and emphasise the importance of computational

screening to balance productivity with ease of control. The insights gained from this study offer

a roadmap toward optimising microbial systems for sustainable bioprocesses and circular bio-

economies.

Importance: The transition to sustainable biomanufacturing is vital for reducing reliance on

fossil-based resources in industries such as biofuels, wastewater treatment, and

pharmaceuticals. While monocultures are preferred for their predictability, microbial

communities offer greater metabolic versatility, particularly in nutrient-limited environments.

However, their complexity and the lack of systematic evaluation tools have hindered wider

adoption. Identifying the most efficient microbial system—whether monoculture or

community—and optimizing it for resource efficiency is key to sustainability and aligns with the

UN Sustainable Development Goals (SDG 9 and SDG 12). COSMOS addresses this gap by

providing a computational framework to assess microbial systems, minimizing the need for

extensive experimental testing. By enabling data-driven selection of scalable and efficient

bioprocessing strategies, COSMOS supports the transition to circular bio-economies and

sustainable manufacturing, ensuring resource-efficient production while reducing

environmental impact.

Keywords: metabolic modelling, genome-scale models, metabolic engineering, constraint-based

modelling, monoculture, sustainable bioprocess, circular bioeconomy, responsible consumption

and production, microbial systems, sustainability, SDGs

1   Introduction

Biomanufacturing supports sustainable development by converting renewable

resources, such as agricultural waste or wastewater, into valuable products like biofuels,

pharmaceuticals, and bioplastics (Süntar et al. 2021). Both microbial monocultures and

communities are used in these processes, each with distinct strengths and challenges.

Monocultures are easier to control, manipulate, and engineer for specific product yields, making

them ideal for simple, well-characterised bioprocesses. However, their productivity can reach a

plateau, even after genetic optimisation (Pachapur et al. 2015).

Microbial communities, on the other hand, leverage complementary metabolic

capabilities and interspecies cooperation (Ibrahim, Raajaraam, and Raman 2021). This has

sparked significant interest in engineering co-cultures tailored to achieve specific bioprocess

objectives (Rafieenia et al. 2024). Notably, even communities composed of strains from the

same organism can outperform individual strains. (H. Peng et al. 2024) demonstrated this by

developing a molecular toolkit of auxotrophic and overexpression yeast strains, enabling the

construction of diverse two- and three-member communities with distinct metabolic

capabilities. In contrast, heterologous communities, while more complex, offer unique

advantages. In these systems, organisms break down complex substrates—such as lignin or

cellulose—into simpler compounds, facilitating resource-sharing (X. "Nick" Peng, Gilmore, and

O'Malley 2016; Raajaraam and Raman 2024). Communities are also less prone to feedback

inhibition because one species may utilise by-products of another, leading to enhanced growth

and stability (Ghosh, Chowdhury, and Bhattacharya 2016). Furthermore, the 'division of labour'

across species helps distribute the metabolic burden, improving production efficiency for

complex products (Rafieenia, Atkinson, and Ledesma-Amaro 2022). However, communities are

inherently harder to manage and require careful optimisation to maintain stability and

maximise productivity.

Given these trade-offs, the choice between monocultures and communities is dependent

on the nature of the bioprocess. The cooperative behaviour of communities provides resilience

and efficiency (Ziganshin et al. 2013), whereas monocultures are well-studied and can be easier

to manipulate. Designing effective bioprocesses involves identifying the most suitable microbial

system for specific substrates or desired products. The decision is not always as straightforward

as selecting communities for lignocellulosic biomass conversion and monocultures for simpler

fermentations (Liang et al. 2020). Selecting the optimal microbial system for other processes

can be challenging, as it would require testing numerous combinations of organisms and

environmental conditions—an approach that is both labour-intensive and impractical (Fischer,

Klein-Marcuschamer, and Stephanopoulos 2008). Therefore, computational algorithms become

essential for managing this complexity and streamlining the selection process (García-Jiménez,

Torres-Bacete, and Nogales 2021).

While some algorithms optimize specific co-cultures, few can systematically evaluate

multiple microbial systems to identify the most suitable option for a given environment. For

instance, (Martinez et al. 2022) have developed an algorithm to optimise substrate-pulsing to

obtain stable communities. Other studies have explored strategies to induce microbial

communities for specialized product synthesis, with various computational approaches

developed to regulate these processes (Boruta 2023). There are also studies like (C. Foster et al.

2021) which focus on a single coculture and model growth kinetics to study how interspecies

cell fusion can influence bioproduction. Other algorithms like FLYCOP (García-Jiménez, García,

and Nogales 2018) optimises a specific microbial co-culture for synthesizing the desired

product. While (Wilken et al. 2018), use constraint-based modelling to find suitable bacterial

partners for anaerobic fungi in co-cultures. However, these tools are limited in scope—they

cannot comprehensively analyse multiple communities or compare them with monocultures to

determine the most effective system.

In this work, we present COSMOS, an algorithm that overcomes these limitations and

identifies the best microbial system by building pairwise communities from a set of individual

organisms. COSMOS integrates dynamic Flux Balance Analysis (FBA) (Henson and Hanly 2014;

Mahadevan, Edwards, and Francis J Doyle 2002) and Flux Variability Analysis (FVA) to simulate

the growth of communities and their constituent monocultures (Gudmundsson and Thiele

2010). Moreover, it compares the performance of each community against the highest-

performing monoculture, rather than relying on the average productivity of the monocultures,

ensuring that the best microbial system is chosen. It also enables users to make informed
decisions not only based on productivity but also on additional factors, such as the abundance
distribution of the community or any biological insights that might favour one community over
another.

In summary, this work contributes to the development of sustainable biomanufacturing
by identifying when communities or monocultures should be used and how bioresources can be
efficiently employed. It marks a step forward in creating economically viable, eco-friendly
processes, aligning with the global effort to promote sustainable industrial practices. This
framework aligns with the growing need for optimised bio-based processes that reduce waste
and reliance on fossil resources, promoting a circular economy.

2   Methods

2.1   Flux Balance Analysis (FBA)

The metabolic network of the organism is represented as a stoichiometric matrix A of size
$m \times n$ where m is the number of metabolites and n is the number of reactions. The entries in
each column represent the stoichiometric coefficients of the metabolites that constitute the
reaction. The linear programming (LP) problem is denoted by

$$max_v \qquad c^T v$$

$$\text{subject to} \quad A. v = 0$$

$$\text{while} \qquad v_l \leq v \leq v_u$$

where c is a vector of weights denoting the contribution of each reaction to the objective
function, $v \in R^n$ is the vector of metabolic fluxes, v_l and v_u are the lower and upper bounds,
respectively.

2.2   Dynamic FBA

Dynamic FBA is performed using the static optimisation approach where the entire batch time
is divided into intervals, and the LP is solved at each time interval as a standard FBA problem.
While FBA assumes that the cell maintains an intracellular steady state at a given time point, all
variables—both intracellular and extracellular—are subject to time variations. The growth rate
(μ) and intracellular fluxes (v), including product secretion rates (v_p) at any given time point
are determined by solving a standard FBA problem at the biomass concentration X. Instead of
using fixed substrate uptake rates as in the case of classic FBA, we utilise extracellular
concentrations of substrates (S) and products (P) to calculate dynamic substrate uptake rates
(v_s) based on specific kinetic parameters, as shown in Figure 1. These rates reflect the

maximum uptake capabilities at each time point and are applied as constraints in the
calculations.

Fig 1. Dynamic flux balance analysis model for a microbial community. The substrate flux
v_{s_j} for metabolite j is determined using substrate concentration S_j kinetic parameters V_{max} and
K_m at each timestep. The substrate fluxes, together with the lower and upper bounds, v_l and v_u
are used to solve the FBA problem. The FBA problem comprises the stoichiometric matrix A and
the vector of weights c which represents the contribution of each reaction to the objective
function, typically the maximization of the growth rate, μ. This optimization is performed
iteratively at each timestep dt for each species i using the current concentrations of biomass X_i,
substrate flux v_{s_j} and product flux v_{p_k} for product k.

2.2.1   Calculation of substrate uptake rate

The substrate uptake limit is limited by two factors – the amount of nutrient available in the
medium for each species i and the transport kinetics for the substrate. We have used a similar
approach as (Chiu, Levy, and Borenstein 2014; Varma and Palsson 1994; Wagner 2022) to
calculate the substrate uptake rate. The nutrient concentration in the medium is denoted by
$S_j(t)$, where the maximum amount of nutrient j that species i can import per gram biomass per
hour is $S_{j,conc}$

$$S_{j,conc} = \frac{S_j(t)}{X_i(t) * \Delta t}$$

154   The second limitation is transport, where the cell's transport mechanism may not be
155   able to import all the available nutrients. Nutrient transport often follows Michaelis–Menten
156   kinetics (Button 1991; Fiksen, Follows, and Aksnes 2013), and we therefore use the kinetic
157   parameters V_{max} and K_m to calculate the transport limit $S_{j,trans}$

$$S_{j,trans} = \frac{V_{max} * S_j(t)}{K_m + S_j(t)}$$

158 The substrate uptake rate v_s is the minimum of these two limits and is given by

$$v_s = min \left[\frac{S_j(t)}{(X_i(t) * \Delta t)}, \frac{V_{max} * S_j(t)}{K_m + S_j(t)} \right]$$

This substrate uptake rate is used as an additional constraint for the parsimonious flux balance
analysis to compute the growth rate μ_i of each species i and the medium concentration for each
substrate is updated.

2.2.2 Flux Variability Analysis (FVA)

To evaluate the production of each product under study we performed an FVA at each
time step. The maximum flux through the exchange reaction for k products v_{p_k} is evaluated as
follows

$$max_v \quad v_{p_k}$$

166 $$\text{subject to} \quad A. v = 0$$

167 $$\text{while} \quad v_l \leq v \leq v_u$$

168 The product concentration $P_{k,i}$ for product k in species i is given by

$$P_{k,i} = v_{p_k} * X_i(t) * \Delta t$$

2.3 Comparative analysis

High-quality GSMMs of more than 30 organisms were obtained from the BiGG models
database (King et al. 2016) and the BioModels database (Malik-Sheriff et al. 2020). These
models were filtered based on pathogenicity, model quality and annotation compatibility
(Lieven et al. 2020). We chose ten organisms for our final analysis, and though some of these are
pathogenic, bioproduction has been successfully demonstrated in all of them. These models
were used to form pairwise communities with a universal compartment shared by the
organisms, representing the external medium. The uptake kinetic parameters V_{max} and K_m were
assumed to be constant (V_{max} = 20 mmol/gDW/hr and K_m = 0.05 mmol) for all metabolites, as
the parameters for all the organisms were not readily available. This value of V_{max} and K_m lies
within the broad range reported in the BRENDA database (Scheer et al. 2011) and has been
utilised in several previous studies (Chiu, Levy, and Borenstein 2014; Roman and Wagner
2018). The pairwise community and the monoculture growth rates were simulated in the same
external medium, and the product concentrations in both cases were compared to find the best
system under each scenario and target product, as shown in Figure 2. The final biomass,
substrate, and product concentrations were averaged over the last five feasible solutions to

ensure robustness. A minimal abundance of 0.1 and at least a 10% increase in biomass was set
as a requirement for the community to be considered viable.

Fig 2. Workflow of COSMOS. The preprocessing step involves selecting target organisms,
ensuring model quality, and setting parameters like   ,   , initial biomass, and process time.
The algorithm uses these inputs to form pairwise communities and applies dynamic FBA and
FVA to simulate growth and productivity across mono- and co-cultures in the same
environment. The resulting growth, abundance, and productivity metrics help identify the
optimal microbial system for the given conditions, which can then be experimentally validated
for bioprocess implementation.

2.4   Statistical analysis

The distribution of the data was assessed using the Shapiro-Wilk test, which indicated a
non-normal distribution. Therefore, non-parametric statistical methods were applied. We
implemented a generalized additive model (GAM) regression in R (mgcv package) to model the
relationships between variables before hypothesis testing. This approach allowed us to account
for and exclude confounding factors before performing statistical comparisons. For group-wise
comparisons, we used the Kruskal-Wallis test, a non-parametric alternative to ANOVA. If a
statistically significant difference was detected ($p < 0.05$), we conducted post-hoc pairwise
comparisons using the Dunn test with Benjamini-Hochberg correction to adjust for multiple
comparisons. All statistical analyses were performed in R (version 4.4.2) using the mgcv and
stats packages.

3　Results

3.1　Navigating Trade-offs: Choosing Between Monocultures and Communities

The medium composition is one of the most substantial factors that can affect a bioprocess.
We come across both nutrient-dense or 'rich' feedstock like animal manure and sludge, which
have more nitrogen content and trace minerals, and 'less-dense' feedstock like wheat straw and
corn stover (Mohd Johari et al. 2020), which may lack essential micronutrients. The biosynthetic
capability of the organisms can also be affected as a result of this medium composition
(Williams et al. 2016). So, we need to either choose the target product based on the feedstock
availability or source the best-suited feedstock for our product of interest. Using COSMOS, we
analyse the effect of medium composition on the biosynthetic capability of a diverse set of
organisms.

To investigate the impact of medium composition, we designed two types of media: a
'minimal medium' and a 'rich medium.' The 'rich medium' contains over 60 metabolites
commonly found in growth media (Pacheco, Moel, and Segrè 2019). In contrast, the minimal
medium consists of around 40 essential metabolites required for the growth of all the individual
organisms under study. A complete list of organisms and media components is provided in
Supplementary Tables 1-3.

The initial concentration of each component was set to 10 mmol/L in both media. We
also investigated the impact of oxygen availability by growing communities under both
anaerobic and aerobic conditions, with the latter set to an oxygen concentration of 10 mmol/L.
Obligate anaerobes and aerobes were excluded from their respective simulations
(Supplementary Table 1), resulting in four distinct environments: aerobic-rich, aerobic-minimal,
anaerobic-rich, and anaerobic-minimal. An initial biomass of 0.01 g/L was chosen for both the
organisms of the co-culture to ensure sufficient growth. All simulations were run for 12 hours,

by which the carbon source was fully consumed, and the organisms reached the stationary
phase. We calculated the productivity of each product for both communities and the constituent
monocultures, as shown in Figure 3.

Figure 3. Productivity of communities and monocultures across varying
environmental conditions. The productivity of both communities and monocultures for
products across all four environments is represented. Five products, viz. adipic acid, xylitol,
catechol, butanol and butyrate, were excluded as they are not produced in any of the organisms
under study.

When comparing the four environments, we found that overall productivity is highest in
the aerobic-rich environment (Figure 3). However, we found the effect of the environment on
microbial systems to be more nuanced and product-dependent. While monocultures, on
average, performed better in the aerobic-rich environment, communities often achieved the
highest productivity for specific products. The list of best-performing microbial systems for the
aerobic-rich environment is summarized in Table 1. Our results highlight the importance of
conducting an analysis tailored to the specific product and environmental conditions to identify
the optimal microbial system.

To efficiently compare the productivity of communities and monocultures, we calculated
the productivity ratio for all products across the four environments. It should be noted that the
maximum productivity of the two monocultures was compared against the productivity of the
community. The productivity ratios across all four environments are represented in Figure 4.

In a community with organisms A and B, where

$$Mono_A + Mono_B = Comm_{AB}$$

252   the change in productivity is given as

$$\Delta Productivity = Productivity(Comm_{AB}) - \max\left(Productivity(Mono_A), Productivity(Mono_B)\right)$$

253   and the ratio of productivity of the community to that of the monoculture is given as

$$Productivity\ ratio\ = \frac{\Delta Productivity}{abs\left(\max\left(Productivity(Mono_A), Productivity(Mono_B)\right)\right)}$$

When comparing the productivity ratio of communities and monocultures, we find that the
anaerobic-rich environment predominantly favours community-based production, while
monocultures perform best in the aerobic-minimal medium ($p = 2.38 \times 10^{-10}$; Supplementary
Data 5). One possible explanation is that the incomplete nature of anaerobic fermentation can
lead to lower productivity in monocultures. In contrast, microbial communities may exchange
primary metabolites, enhancing both growth and productivity, particularly in anaerobic
conditions. To test this hypothesis, we analysed the abundance ratios of five communities across
all four environmental conditions, as shown in Table 2. Under anaerobic conditions, although
total biomass decreases, the abundance ratios of both organisms become more balanced. This
equilibrium may contribute to the improved performance of communities in anaerobic
environments. Additionally, the rich medium supports higher productivity by promoting better
growth and increased metabolite exchange. Based on these findings, simple monocultures may
be preferable for aerobic environments, while communities offer advantages for anaerobic
fermentation.

However, as discussed earlier, there are notable exceptions where communities excel in aerobic
environments. This underscores the importance of evaluating all candidate organisms as
monocultures and communities to determine the optimal microbial system for a given product
and environment. If the differences in productivity are minimal, monocultures may be
preferable due to easier process control, whereas co-cultures can be advantageous if they offer
superior biosynthetic capabilities.

For example, in the aerobic-rich medium, the sorbitol productivity of the *S. cerevisae – B. subtilis*
and the *S. cerevisae – L. lactis* communities are the highest and have 59% and 48%
improvement, respectively, when compared to monocultures. Likewise, in the anaerobic-
minimal medium*, Bacillus* subtilis outperforms all the other microbial communities in the
production of L-lactate, with the productivity of B. subtilis being 0.29 mmol/L/hr. It is to be
noted that the second-best alternative is *E. coli – P. aeruginosa* community, with a productivity
of 0.12 mmol/L/hr of L-lactate, which is less than half of the monoculture. These examples

demonstrate that while general trends provide useful insights, exceptions can be strategically
utilized to optimize bioprocesses.

Therefore, a systematic screening of both monocultures and communities is essential. In the
following sections, we explore the impact of additional factors, such as interaction type, carbon
sources and initial biomass ratio, on productivity and community dynamics.

Fig 3. Effect of medium composition and oxygen availability on the productivity of
communities vs monocultures. The average productivity ratio of communities across all
products and environments is represented. Positive values (blue) indicate increased
productivity in communities compared to monocultures, while negative values (red) indicate a
decline. In cases where monoculture productivity is extremely low, leading to an exceptionally
high productivity ratio, we classify it as 'High' and denote it with a navy blue colour.

Table 1. Optimal microbial for four products in the aerobic-rich environment

#	Fumarate			Ethanol			Spermidine			Glutathione		
	Microbial system	Productivity (mmol/L/hr)	Abundance	Microbial system	Productivity (mmol/L/hr)	Abundance	Microbial system	Productivity (mmol/L/hr)	Abundance	Microbial system	Productivity (mmol/L/hr)	Abundance
1	*P. aeruginosa - K. pneumoniae*	1.28	0.85,0.15	*S. oneidensis - K. pneumoniae*	2.04	0.19,0.81	*E. coli*	0.07	1	*S. oneidensis - K. pneumoniae*	0.26	0.19,0.81
2	*E. coli - K. pneumoniae*	1.22	0.60,0.40	*S. cerevisiae - K. pneumoniae*	1.42	0.13,0.87	*B. subtilis*	0.05	1	*E. coli - K. pneumoniae*	0.25	0.60,0.40
3	*S. oneidensis - K. pneumoniae*	1.10	0.19,0.81	*K. pneumoniae*	1.29	1	*S. cerevisiae - L. lactis*	0.03	0.84,0.16	*E. coli - P. putida*	0.19	0.72,0.28
4	*E. coli - P. putida*	1.07	0.72,0.28	*E. coli*	1.10	1	*S. cerevisiae*	0.02	1	*E. coli*	0.17	1
5	*B. subtilis - K. pneumoniae*	0.88	0.15,0.85	*P. aeruginosa - S. oneidensis*	0.63	0.89,0.11				*B. subtilis - K. pneumoniae*	0.13	0.15,0.85

295   **Table 2. Comparison of Microbial Communities Exhibiting Growth Across All Environments**

Community	Aerobic-rich			Aerobic-minimal			Anaerobic-rich			Anaerobic-minimal		
	Biomass	Total Biomass	Abundance	Biomass	Total Biomass	Abundance	Biomass	Total Biomass	Abundance	Biomass	Total Biomass	Abundance
E. coli - L. lactis	0.76,0.17	0.93	0.82,0.18	0.57,0.13	0.7	0.81,0.19	0.35,0.23	0.58	0.60,0.40	0.19,0.24	0.43	0.44,0.56
L. lactis - S. oneidensis	0.13,0.35	0.48	0.26,0.74	0.13,0.36	0.49	0.26,0.74	0.14,0.22	0.36	0.38,0.62	0.13,0.22	0.35	0.38,0.62
L. lactis - K. pneumoniae	0.12,0.80	0.92	0.13,0.87	0.12,0.35	0.47	0.15,0.85	0.13,0.48	0.61	0.21,0.79	0.14,0.29	0.43	0.32,0.68
B. subtilis - K. pneumoniae	0.14,0.80	0.94	0.15,0.85	0.14,0.66	0.8	0.18,0.82	0.17,0.46	0.63	0.27,0.73	0.17,0.29	0.46	0.37,0.63
S. oneidensis - K. pneumoniae	0.19,0.78	0.97	0.19,0.80	0.15,0.64	0.79	0.19,0.81	0.23,0.48	0.71	0.32,0.68	0.25,0.27	0.52	0.47,0.53

3.2 Synergy in Action: Positive Interactions Drive Higher Productivity

As we are dealing with pairwise communities, another interesting factor that we can analyse is the effect of interaction. To find the interaction type of a community, we compare the growth of the organism as a monoculture and in the community under the same environmental conditions. The communities were categorised into six interaction types, namely, competition, amensalism, parasitism, neutralism, commensalism and mutualism, based on a 10% difference in growth rates of the microbe in the co-culture and the monoculture (Heinken and Thiele 2015).

The average *Productivity ratio* was calculated for each product across the four environments, is shown in Figure 5. Mutualistic interactions generally lead to the highest productivity gains in communities, likely due to the increased biomass of the participating organisms. Parasitic interactions follow, with productivity varying depending on the product— some are better produced in communities, while others are more efficiently generated in monocultures. This variation may arise because, in some cases, the producer strain exhibits higher growth, whereas in others, it experiences reduced growth. The effect of interaction type on the productivity ratio was statistically significant ($p = 1.4 \times 10^{-10}$; Supplementary Data 6). However, no significant differences were observed among competition, commensalism, and amensalism, suggesting that these interaction types have minimal impact on productivity differences between communities and monocultures.

Fig 5. Effect of interaction type on the productivity of communities vs monocultures across products. Positive values (blue) indicate an increase in the productivity of a given

319 product in the community compared to the monocultures, while negative values (red) reflect a
320 decline in productivity. Neutralism was not observed.

3.3 Tailored Metabolism: Communities and Monocultures Show Distinct Carbon Preferences

Another key factor we examined is the impact of different carbon sources on the growth
and biosynthetic capabilities of microbial systems. In this analysis, we replaced the glucose in
the 'rich' and 'minimal' media with 10 mmol/L of the carbon source under investigation. We
tested seven different carbon sources and compared the productivity of communities and
monocultures across all four environments, as shown in Figure 6.

Our findings show that while there are some variations, lactose and sucrose consistently
enhance productivity in communities, whereas xylose and fructose favour monocultures.
Interestingly, the aerobic-rich medium deviates from the pattern observed in the other three
environments. While xylose and fructose promote higher productivity in communities in the
aerobic-rich environment, these carbon sources support higher productivity in monocultures in
other environments. Although the underlying cause remains unclear, this provides an additional
factor that could be manipulated to optimize the bioprocess. This insight can guide the design or
supplementation of fermentation media such that the choice of carbon source aligns with the
microbial system and environment. The effect of carbon source on the productivity ratio was
statistically significant ($p = 1.8 \times 10^{-4}$; Supplementary Data 7).

To explore the variation amongst communities, we calculated the productivity ratio for
four different products across communities in the aerobic-rich medium, as shown in
Supplementary Figure 1. Some communities, like *Shewanella oneidensis – Klebsiella pneumoniae*,
showed minimal sensitivity to various carbon sources. However, others, such as *S. cerevisiae – L.*
lactis, displayed varying productivity across carbon sources. In some cases, like *E. coli – S.*
cerevisiae, the community does not grow under specific carbon sources. Moreover, this
behaviour can also be product-dependent. Thus, product-specific analysis is crucial for
identifying the optimal combination of carbon source and microbial system.

Fig 6. Effect of carbon source on the productivity of communities vs monocultures. The effect of carbon sources on the *Productivity ratio* is compared across all four environments. Positive values (blue) indicate an increase in the productivity of a given product in the community compared to the monocultures, while negative values (red) reflect a decline in productivity.

3.4   Computational Screening Reveals Ideal Microbial Systems Across Fermentation Conditions

While communities generally perform better in challenging environments, exceptions may occur. Therefore, it is essential to analyse the specific product of interest before selecting the most suitable microbial system. Furthermore, variations between communities mean that choosing the optimal community for a given nutrient source is crucial. Since experimentally comparing multiple communities is time-consuming and labour-intensive, computational analysis becomes essential.

We compared the change in productivity between communities and monocultures across all products, as shown in Figure 8. The mean productivity ratio for 25 products was calculated across the four environments for each community. Positive values (blue) indicate higher productivity in communities compared to their constituent monocultures, while negative values (red) indicate lower productivity. Clear clustering patterns emerge both by product and community, highlighting preferences in microbial systems. The effect of community on the productivity ratio was found to be statistically significant, with a ($p = 2.2 \times 10^{-16}$;Supplementary Data 8).

Microbial communities such as *P. aeruginosa–S. oneidensis* and *E. coli–S. oneidensis* generally outperform their monocultures, particularly in the production of metabolites like

pyruvate, hydrogen, and glutathione. In contrast, communities like *Synechocystis spp.–K.* *pneumoniae* and *P. putida–K. pneumoniae* consistently exhibit lower productivity compared to their respective monocultures. Furthermore, metabolite production appears to be highly product- and organism-dependent. For instance, acetate and pyruvate are more efficiently synthesized in co-culture systems, whereas sorbitol and propionate yield higher levels in monocultures. Additionally, productivity patterns vary across different environmental conditions, underscoring the importance of evaluating microbial systems in a context-specific manner.

To investigate this behaviour further, we identified the best microbial systems for producing four key products from different classes under all four environmental conditions, as summarized in Supplementary Table 5. The observations reveal that productivity is highest in the aerobic-rich environment, as discussed in Sections 3.1 and 3.2. Notably, the anaerobic-rich environment also performs competitively and, in some cases, surpasses the aerobic minimal medium in productivity. The findings in Supplementary Table 5 show that some of the best-performing systems in aerobic environments are monocultures, while communities tend to outcompete monocultures in anaerobic environments. It also highlights variability within products—while fumarate is consistently better produced by communities across all environments, spermidine is predominantly produced by monocultures in most conditions. This highlights the necessity of selecting the optimal microbial consortium and environmental setting tailored to the specific product of interest.

Moreover, when choosing a community, the selection criteria need not be restricted to productivity but can also be extended to other factors like abundance ratios. When two systems exhibit similar productivity, choosing the one with a more balanced abundance may be preferable, as it can lead to a more stable community. Moreover, the final choice can also be guided by biological insights, such as ease of handling or specific biological characteristics desirable for the bioprocess of interest. The optimal systems in the other environments are provided in Supplementary Data 1-4.

Additionally, we compiled the top-performing systems for all 25 products across four environments in Supplementary Table 6. As expected from previous analyses, monocultures dominate in the aerobic-rich medium. Meanwhile, communities excel in the anaerobic-minimal medium, reinforcing the idea that harsher conditions favour biosynthesis in the communities. The detailed results for all 25 products are in Supplementary Data 1-4, with key findings aligning with experimental studies summarized in Table 3 alongside references.

Fig 8. Comparison of Productivity of Communities and Monocultures. This figure presents the mean productivity ratio of communities, averaged across four environments, for all the products under study. Positive values (blue) indicate an increase in the productivity of a given product in the community compared to the monocultures, while negative values (red) reflect a decline in productivity.

409   **Table 3. Results that align closely with experimental findings**

#	Product	System Name	Productivity in Community (mmol/L/hr)	Productivity in Monoculture A (mmol/L/hr)	Productivity in Monoculture B (mmol/L/hr)	Improvement in productivity (%)	Abundance	Environment	Reference
1	1,3 - Propanediol	*S. oneidensis - K. pneumoniae*	0.064	0	0.0519	23.31	0.32,0.68	Anaerobic-rich	(Wang et al. 2023)
2		*P. aeruginosa - K. pneumoniae*	0.053	0	0.018	196.11	0.58,0.42	Anaerobic-minimal	(Li, Zhu, and Ge 2019)
3	2,3 - Butanediol	*P. aeruginosa - K. pneumoniae*	0.073	0	0.036	103.06	0.58,0.42	Anaerobic-minimal	(Li, Zhu, and Ge 2019)
4	Hydrogen	*C. ljungdahlii - B. subtilis*	0.684	0.091	0	647.92	0.87,0.13	Anaerobic-rich	(Chang et al. 2008)
5	Ethanol	*S. cerevisiae – B. subtilis*	0.630	0.416	0.604	4.41	0.69,0.31	Aerobic-rich	(Tantipaibulvut et al. 2015)
		S. cerevisiae - L. lactis	0.584	0.416	0.009	40.22	0.84,0.16	Aerobic-rich	(Carvalho et al. 2015)
		S. cerevisiae - L. lactis	0.410	0.363	0.008	13.03	0.65,0.35	Aerobic-minimal	(Li, Zhu, and Ge 2019)

3.5   Optimising Biomass Ratios Enhances Productivity in Microbial Communities

The initial biomass ratio of organisms within a community can significantly influence
growth, abundance, and overall community dynamics. By adjusting the inoculum ratio, we can
further enhance the biosynthetic capabilities of the community. Tables 4 and 5 illustrate how
variations in inoculum ratios impact 1,3-propanediol (1,3-PDO) production in *P. aeruginosa - K.*
pneumoniae and *S. oneidensis - K. pneumoniae* communities. Product concentrations vary with
inoculum ratios, highlighting distinct responses between the two communities. While increasing
K. pneumoniae inoculum enhances product titer in both, the *S. oneidensis - K. pneumoniae*
community shows an increase in total biomass, whereas *P. aeruginosa - K. pneumoniae* displays
a growth reduction. Moreover, higher *K. pneumoniae* inoculum in *S. oneidensis - K. pneumoniae*
leads to a more skewed abundance ratio. These findings suggest that inoculum ratios can have
varied effects on community dynamics and biosynthetic output, emphasizing the need for
careful ratio optimization.

Table 4. Effect of inoculum ratio on the productivity of 1,3-PDO in *P. aeruginosa -*
***K. pneumoniae* community**

Biomass ratio	Biomass A (g/L)	Biomass B (g/L)	Total Biomass (g/L)	Abundance	Product concentration (mmol/L)
0.9,0.1	0.67	0.03	0.70	0.96,0.04	0.01
0.8,0.2	0.63	0.06	0.69	0.91,0.09	0.02
0.7,0.3	0.59	0.09	0.68	0.87,0.13	0.03
0.6,0.4	0.54	0.12	0.66	0.81,0.19	0.03
0.5,0.5	0.48	0.16	0.64	0.75,0.25	0.04
0.4,0.6	0.39	0.19	0.58	0.67,0.33	0.06
0.3,0.7	0.33	0.23	0.56	0.59,0.41	0.07
0.2,0.8	0.23	0.27	0.50	0.47,0.53	0.08
0.1,0.9	0.14	0.32	0.46	0.31,0.69	0.09

Table 5. Effect of inoculum ratio on the productivity of 1,3-PDO in *S. oneidensis - K. pneumoniae* community

Biomass ratio	Biomass A (g/L)	Biomass B (g/L)	Total biomass (g/L)	Abundance	Product concentration (mmol/L)
0.1,0.9	0.47	0.12	0.59	0.79,0.21	0.04
0.2,0.8	0.40	0.23	0.62	0.64,0.36	0.07
0.3,0.7	0.35	0.33	0.68	0.51,0.49	0.09
0.4,0.6	0.29	0.41	0.7	0.41,0.59	0.11
0.5,0.5	0.23	0.48	0.71	0.32,0.68	0.14
0.6,0.4	0.19	0.57	0.76	0.24,0.76	0.14
0.7,0.3	0.13	0.63	0.76	0.18,0.82	0.18
0.2,0.8	0.09	0.67	0.76	0.11,0.88	0.22
0.9,0.1	0.04	0.75	0.8	0.05,0.95	0.23

3.6   Validation

Though some of the results we obtain corroborate with experimental studies as listed in Table 3, these studies only talk about the biosynthetic capability of the community and not the monoculture. There are very few studies that compare the production capabilities of the monoculture and the community in the same environment. In one such study (Wang et al. 2023), they evaluated the production of 1,3-PDO in *Klebsiella pneumoniae* monoculture and the *Klebsiella pneumoniae – Shewanella oneidensis* community using glycerol as the carbon source under anaerobic conditions. They found that *Shewanella oneidensis* acts as an electron mediator in the co-culture to attain better production of 1,3-PDO. Notably, under anaerobic conditions, this community is the top producer of 1,3-PDO in our analysis. To demonstrate the reliability of COSMOS, we simulated the growth of the community and the monoculture in the medium used in the experimental study and compared the results. The first analysis is to identify the optimum glycerol concentration for the monoculture. We found 50 g/L glycerol to be the optimum concentration for 1,3-PDO production, which correlates with the experiment values, as shown in Table 6.

Table 6. Concentration of 1,3 propanediol in experiments and simulations under varying glycerol concentration

Glycerol concentration (g/L)	Concentration of 1,3 PDO in experiments (g/L)	Concentration of 1,3 PDO in COSMOS (g/L)
40	11.3	1.76
50	12.68	1.83
70	12.21	1.58

For the community analysis, the experimental group used a fed-batch reactor under anaerobic conditions where the initial glycerol concentration was 50 g/L. When the glycerol concentration falls below 10 g/L, additional glycerol is introduced to maintain a final concentration of 30 g/L. They found that the community produced 32.01 g/L of 1,3-PDO, which was significantly higher when compared to the monoculture. They also optimised the inoculum ratio and found that a 1:1 ratio of *Klebsiella pneumoniae*: *Shewanella oneidensis* resulted in the highest product concentration, though the exact values are not listed. We did a similar analysis using COSMOS and found that the community was, in fact, able to produce more product in the fed-batch system when compared to the monoculture. However, we found that both the inoculum ratios of 1:1 and 2:1 performed equally well, as shown in Table 7. This might be because we have used the wild-type models of both organisms and lack the strain-specific information that might be required.

Table 7. Concentration of 1,3 -propanediol with varying inoculum ratio according to simulations

Initial biomass ratio	1,3 Propanediol (g/L)	Abundance
2,1	2.42	0.04,0.96
1,1	2.20	0.07,0.93
1,2	0.31	0.1,0.9
1,3	0.28	0.12,0.88

The study reported monoculture and co-culture yields of 0.4 g/g and 0.44 g/g, respectively, reflecting a 10% improvement. COSMOS predicted yields of 1.50 g/g for monocultures and 1.33 g/g for co-cultures, indicating a 12.6% increase. Although the absolute values differ, the algorithm effectively captured the trend in performance, identifying optimal substrate concentrations and inoculum ratios. This demonstrates the potential of dynamic modelling in bioprocess design, offering a powerful tool for efficiently comparing multiple communities and monocultures to identify the most productive configurations.

4 Discussion

Microbial communities often exhibit superior biosynthetic capability compared to
monocultures (Rapp, Jenkins, and Betenbaugh 2020; Xu and Tschirner 2011). However, their
application in bioproduction remains limited to cases where their productivity is already known
or where their metabolic diversity is essential, such as when one community member can
metabolise a complex carbon source that the others cannot. Studies that directly compare the
biosynthetic potential of multiple communities remain scarce (Wilken et al. 2018), leaving a gap
in identifying the most effective microbial systems.

To address this, COSMOS provides a systematic approach to evaluating microbial systems
by assessing a wide range of co-cultures alongside their corresponding monocultures. Unlike
tools like FLYCOP (García-Jiménez, García, and Nogales 2018), which optimizes a single co-
culture, COSMOS evaluates a broad range of co-cultures alongside their constituent
monocultures to identify the most effective microbial system for a given product. It also offers
granular insights into the system's performance across a wider range of environmental
conditions. COSMOS is particularly valuable when working with low-quality feedstocks, such as
agricultural residues and wastewater, which are gaining importance due to global efforts to
transition toward second-generation feedstocks for improved food security and sustainability
(Russo et al. 2025). Given these challenges, optimizing the microbial system is just as crucial as
optimizing the environment to develop efficient and sustainable bioprocesses.

While productivity is often the primary factor in selecting a microbial system, the choice
between monocultures and communities depends on additional factors. If a community offers
substantially higher productivity, it is the better option for bioproduction. However, when
productivity is comparable, monocultures are preferred for their ease of control and simplicity.
In cases where operational simplicity takes precedence over maximum output, slightly less
productive monocultures can be selected. COSMOS streamlines this decision-making process by
assessing the biosynthetic potential of diverse microbial systems under diverse environments.

COSMOS employs parsimonious dynamic FBA (Henson and Hanly 2014; Mahadevan,
Edwards, and Francis J Doyle 2002) to model microbial growth in synthetic consortia, allowing
it to capture fluctuating growth rates. While traditional steady-state modelling is simpler to use,
it assumes equal growth rates and is better suited to natural communities (Chan, Simons, and
Maranas 2017). Therefore, we use dynamic FBA, which accounts for fluctuating growth and
provides promising results even with the lack of organism-specific parameters (Gomez, Höffner,
and Barton 2014). This makes COSMOS especially valuable for engineered or synthetic

consortia, where kinetic data is often unavailable, and the stability of the co-culture remains
uncertain.

By integrating Flux Variability Analysis (FVA), COSMOS minimizes redundancy in metabolic
predictions, providing more precise insights into productivity (Gudmundsson and Thiele 2010).
To ensure an unbiased comparison, COSMOS evaluates communities and monocultures under
identical conditions, benchmarking community productivity against the highest-performing
monoculture. This approach prioritizes final productivity or yield, making it a more effective
tool for selecting optimal bioproduction systems (Zaramela et al. 2021).

To understand the factors that influence bioproduction in cocultures and monocultures, we
tested COSMOS across four environments: aerobic-rich, aerobic-minimal, anaerobic-rich, and
anaerobic-minimal media. While overall productivity was highest in the aerobic nutrient-rich
conditions, communities outperformed monocultures in anaerobic environments. This
advantage is likely driven by cooperative interactions facilitated by metabolite exchange from
incomplete anaerobic fermentation. Additionally, the more balanced abundance ratios observed
in co-cultures under anaerobic conditions further support this hypothesis.

Interestingly, there were exceptions where communities achieved maximum productivity
in aerobic environments, while certain monocultures outperformed communities in anaerobic
conditions. This highlights the importance of systematically evaluating both communities and
monocultures before selecting the optimal microbial system for a specific product. COSMOS
enables us to maximize productivity in the chosen fermentation medium and provides insights
into the role of mutualistic interactions in community design. By leveraging these interactions,
COSMOS aids in optimizing both productivity and stability for biomanufacturing applications.

We also examined the effect of carbon sources on these systems. Certain carbon sources,
like lactose and sucrose, enable better productivity in the community, while fructose boosts
monoculture performance in most environments. This analysis aids in selecting or enriching
fermentation media with specific carbon sources. The algorithm can also be easily extended to
study the effect of two or more carbon sources. COSMOS also equips us to optimise the initial
inoculum ratio to further improve productivity.

To validate COSMOS, we applied it to the *Klebsiella pneumoniae – Shewanella oneidensis*
community and its corresponding monocultures, successfully capturing the effects of carbon
source concentration and inoculum ratio (Wang et al. 2023). The observed trends in yield and
productivity across different conditions aligned with experimental data, confirming the
reliability of the algorithm. Furthermore, several microbial communities identified through our
analysis have been experimentally validated as stable consortia, with some already proving
their biosynthetic potential (as listed in Table 3).

While COSMOS provides valuable results using standard kinetic parameters and wild-type

GSMMs, its accuracy can be significantly improved by using experimentally-determined kinetic

parameters and strain-specific GSMMs (C. J. Foster et al. 2021; Link, Christodoulou, and Sauer

2014). However, the availability of high-quality, manually curated models for diverse organisms

remains a limitation (Ponce-de-Leon et al. 2015). Improving the accessibility and accuracy of

metabolic models will strengthen computational predictions. Additionally, integrating multi-

omic data—such as transcriptomic and proteomic information—into GSMMs could further

refine the algorithm's performance, paving the way for more precise bioprocess optimization

(Sen and Orešič 2023).

Notably, the insights from this study have direct relevance to fields like biofuel production,

wastewater treatment, and pharmaceutical biosynthesis (Keasling et al. 2021). Moreover, by

optimising microbial systems for nutrient-limited and waste-derived feedstocks, our approach

aligns with the UN Sustainable Development Goals (SDGs), particularly SDG 9 (Industry,

Innovation, and Infrastructure) and SDG 12 (Responsible Consumption and Production)

(Keasling et al. 2021). In summary, COSMOS provides a robust computational framework to

evaluate and compare the productivity of monocultures and microbial communities, addressing

the challenges of experimental testing. It offers insights into how environmental factors, carbon

sources, and inoculum ratios affect performance, helping researchers make well-informed

decisions about which microbial system to use. Whether the goal is to work with nutrient-dense

or nutrient-limited media, our approach ensures the selection of the most effective system,

enhancing productivity, process efficiency, and community stability.

5   Conclusion

As industries pivot toward sustainable practices, bioresources like agricultural residues

and wastewater will play a vital role in producing specialty and bulk chemicals. Although

nutrient-rich media remain standard for high-value fermentation, optimizing bioprocesses with

nutrient-poor media is increasingly crucial for achieving net-zero carbon goals. However, the

limited economic feasibility and lack of robust tools have hindered their widespread adoption.

Our study demonstrates that *in silico* analyses can effectively optimise microbial systems to

enhance productivity, especially in challenging environments. Microbial communities are

generally better suited for anaerobic conditions due to their complementary metabolism and

cooperative interactions. However, certain communities can outperform monocultures even in

aerobic environments, demonstrating their enhanced biosynthetic capabilities. Likewise, some

monocultures may still surpass communities in anaerobic settings, underscoring the

importance of systematic computational screening to identify the most suitable candidates

before laboratory validation. Tools like COSMOS facilitate this process by helping us leverage

both productivity and ease of control, enabling informed decision-making. As more models and
kinetic data become available, the predictive accuracy of computational approaches will
continue to improve, further advancing bioprocess design.

Ultimately, this approach supports the rational design of microbial systems, advancing
synthetic biology and sustainable biomanufacturing. By understanding when to switch between
monocultures and communities and identifying key environmental and metabolic factors
influencing performance, we can develop more robust and efficient microbial systems. We
believe that our framework and such computational strategies will be key enablers in the
transition to circular bio-economies.

Acknowledgements

LR acknowledges the fellowship from the Ministry of Education, Government of India. KR
acknowledges support from the Science and Engineering Research Board (SERB) MATRICS
Grant MTR/2020/000490, IIT Madras, Centre for Integrative Biology and Systems mEdicine
(IBSE) and the Wadhwani School of Data Science and AI.

Data Availability

All models and analysis scripts used in this study are openly available at
https://github.com/RamanLab/COSMOS

Author Contributions

Lavanya Raajaraam: Conceptualization, Performing Experiments, Analysis of Results, Writing -
original draft. Karthik Raman: Conceptualisation, Analysis of Results.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal
relationships that could have appeared to influence the work reported in this paper.

References

Boruta, Tomasz. 2023. "Computation-Aided Studies Related to the Induction of Specialized
Metabolite Biosynthesis in Microbial Co-Cultures: An Introductory Overview."
Computational and Structural Biotechnology Journal 21: 4021–29.
doi:10.1016/j.csbj.2023.08.011.

Button, D. K. 1991. "Biochemical Basis for Whole-Cell Uptake Kinetics: Specific Affinity,
Oligotrophic Capacity, and the Meaning of the Michaelis Constant." *Applied and*
Environmental Microbiology 57(7): 2033–38. doi:10.1128/aem.57.7.2033-2038.1991.

Carvalho, Fernanda Paula, Whasley Ferreira Duarte, Disney Ribeiro Dias, Roberta Hilsdorf Piccoli, and Rosane Freitas Schwan. 2015. "Interaction of Saccharomyces Cerevisiae and Lactococcus Lactis in the Fermentation and Quality of Artisanal Cachaça." *Acta Scientiarum. Agronomy* 37: 51–60. doi:10.4025/actasciagron.v37i1.18397.

Chan, Siu Hung Joshua, Margaret N. Simons, and Costas D. Maranas. 2017. "SteadyCom: Predicting Microbial Abundances While Ensuring Community Stability" ed. Nathan D. Price. *PLOS Computational Biology* 13(5): e1005539. doi:10.1371/journal.pcbi.1005539.

Chang, Jui-Jen, Chia-Hung Chou, Cheng-Yu Ho, Wei-En Chen, Jiunn-Jyi Lay, and Chieh-Chen Huang. 2008. "Syntrophic Co-Culture of Aerobic *Bacillus* and Anaerobic *Clostridium* for Bio-Fuels and Bio-Hydrogen Production." *International Journal of Hydrogen Energy* 33(19): 5137–46. doi:10.1016/j.ijhydene.2008.05.021.

Chiu, Hsuan-Chao, Roie Levy, and Elhanan Borenstein. 2014. "Emergent Biosynthetic Capacity in Simple Microbial Communities." *PLOS Computational Biology* 10(7): e1003695. doi:10.1371/journal.pcbi.1003695.

Fiksen, Øyvind, Michael J. Follows, and Dag L. Aksnes. 2013. "Trait-Based Models of Nutrient Uptake in Microbes Extend the Michaelis-Menten Framework." *Limnology and Oceanography* 58(1): 193–202. doi:10.4319/lo.2013.58.1.0193.

Fischer, Curt R., Daniel Klein-Marcuschamer, and Gregory Stephanopoulos. 2008. "Selection and Optimization of Microbial Hosts for Biofuels Production." *Metabolic Engineering* 10(6): 295–304. doi:10.1016/j.ymben.2008.06.009.

Foster, Charles, Kamil Charubin, Eleftherios T. Papoutsakis, and Costas D. Maranas. 2021. "Modeling Growth Kinetics, Interspecies Cell Fusion, and Metabolism of a Clostridium Acetobutylicum/Clostridium Ljungdahlii Syntrophic Coculture." *mSystems* 6(1): 10.1128/msystems.01325-20. doi:10.1128/msystems.01325-20.

Foster, Charles J, Lin Wang, Hoang V Dinh, Patrick F Suthers, and Costas D Maranas. 2021. "Building Kinetic Models for Metabolic Engineering." *Current Opinion in Biotechnology* 67: 35–41. doi:10.1016/j.copbio.2020.11.010.

García-Jiménez, Beatriz, José Luis García, and Juan Nogales. 2018. "FLYCOP: Metabolic Modeling-Based Analysis and Engineering Microbial Communities." *Bioinformatics* 34(17): i954–63. doi:10.1093/bioinformatics/bty561.

García-Jiménez, Beatriz, Jesús Torres-Bacete, and Juan Nogales. 2021. "Metabolic Modelling Approaches for Describing and Engineering Microbial Communities." *Computational and Structural Biotechnology Journal* 19: 226–46. doi:10.1016/j.csbj.2020.12.003.

Ghosh, Shiladitya, Ranjana Chowdhury, and Pinaki Bhattacharya. 2016. "Mixed Consortia in Bioprocesses: Role of Microbial Interactions." *Applied Microbiology and Biotechnology* 100(10): 4283–95. doi:10.1007/s00253-016-7448-1.

Gomez, Jose A., Kai Höffner, and Paul I. Barton. 2014. "DFBAlab: A Fast and Reliable MATLAB Code for Dynamic Flux Balance Analysis." *BMC Bioinformatics* 15(1): 409. doi:10.1186/s12859-014-0409-8.

Gudmundsson, Steinn, and Ines Thiele. 2010. "Computationally Efficient Flux Variability Analysis." *BMC Bioinformatics* 11(1): 489. doi:10.1186/1471-2105-11-489.

Heinken, Almut, and Ines Thiele. 2015. "Anoxic Conditions Promote Species-Specific Mutualism
between Gut Microbes In Silico." *Applied and Environmental Microbiology* 81(12): 4049–
61. doi:10.1128/AEM.00101-15.

Henson, Michael A., and Timothy J. Hanly. 2014. "Dynamic Flux Balance Analysis for Synthetic
Microbial Communities." *IET Systems Biology* 8(5): 214–29. doi:10.1049/iet-
syb.2013.0021.

Ibrahim, Maziya, Lavanya Raajaraam, and Karthik Raman. 2021. "Modelling Microbial
Communities: Harnessing Consortia for Biotechnological Applications." *Computational
and Structural Biotechnology Journal* 19: 3892–3907. doi:10.1016/j.csbj.2021.06.048.

Keasling, Jay, Hector Garcia Martin, Taek Soon Lee, Aindrila Mukhopadhyay, Steven W. Singer,
and Eric Sundstrom. 2021. "Microbial Production of Advanced Biofuels." *Nature Reviews
Microbiology* 19(11): 701–15. doi:10.1038/s41579-021-00577-w.

King, Zachary A., Justin Lu, Andreas Dräger, Philip Miller, Stephen Federowicz, Joshua A.
Lerman, Ali Ebrahim, Bernhard O. Palsson, and Nathan E. Lewis. 2016. "BiGG Models: A
Platform for Integrating, Standardizing and Sharing Genome-Scale Models." *Nucleic Acids
Research* 44(D1): D515–22. doi:10.1093/nar/gkv1049.

Li, Ying, Siyu Zhu, and Xizhen Ge. 2019. "Co-Production of 1,3-Propanediol and 2,3-Butanediol
from Waste Lard by Co-Cultivation of Pseudomonas Alcaligenes and Klebsiella
Pneumoniae." *Current Microbiology* 76(4): 415–24. doi:10.1007/s00284-019-01628-5.

Liang, Jinsong, Mohammad Nabi, Panyue Zhang, Guangming Zhang, Yajing Cai, Qingyan Wang,
Zeyan Zhou, and Yiran Ding. 2020. "Promising Biological Conversion of Lignocellulosic
Biomass to Renewable Energy with Rumen Microorganisms: A Comprehensive Review."
Renewable and Sustainable Energy Reviews 134: 110335.
doi:10.1016/j.rser.2020.110335.

Lieven, Christian, Moritz E. Beber, Brett G. Olivier, Frank T. Bergmann, Meric Ataman, Parizad
Babaei, Jennifer A. Bartell, et al. 2020. "MEMOTE for Standardized Genome-Scale
Metabolic Model Testing." *Nature Biotechnology* 38(3): 272–76. doi:10.1038/s41587-
020-0446-y.

Link, Hannes, Dimitris Christodoulou, and Uwe Sauer. 2014. "Advancing Metabolic Models with
Kinetic Information." *Current Opinion in Biotechnology* 29: 8–14.
doi:10.1016/j.copbio.2014.01.015.

Mahadevan, Radhakrishnan, Jeremy S. Edwards, and 3rd Francis J Doyle. 2002. "Dynamic Flux
Balance Analysis of Diauxic Growth in Escherichia Coli." *Biophysical Journal* 83(3): 1331.
doi:10.1016/S0006-3495(02)73903-9.

Malik-Sheriff, Rahuman S., Mihai Glont, Tung V. N. Nguyen, Krishna Tiwari, Matthew G. Roberts,
Ashley Xavier, Manh T. Vu, et al. 2020. "BioModels-15 Years of Sharing Computational
Models in Life Science." *Nucleic Acids Research* 48(D1): D407–15.
doi:10.1093/nar/gkz1055.

Martinez, J. Andres, Matheo Delvenne, Lucas Henrion, Fabian Moreno, Samuel Telek, Christian
Dusny, and Frank Delvigne. 2022. "Controlling Microbial Co-Culture Based on Substrate
Pulsing Can Lead to Stability through Differential Fitness Advantages." *PLOS
Computational Biology* 18(10): e1010674. doi:10.1371/journal.pcbi.1010674.

Mohd Johari, Siti Aminah, Aqsha Aqsha, Noridah B. Osman, M. Rashid Shamsudin, Mariam Ameen, and Sharul Sham Dol. 2020. "Enhancing Biogas Production in Anaerobic Co-Digestion of Fresh Chicken Manure with Corn Stover at Laboratory Scale." *SN Applied Sciences* 2(7): 1260. doi:10.1007/s42452-020-3063-y.

Pachapur, Vinayak Laxman, Saurabh Jyoti Sarma, Satinder Kaur Brar, Yann Le Bihan, Gerardo Buelna, and Mausam Verma. 2015. "Biological Hydrogen Production Using Co-Culture versus Mono-Culture System." *Environmental Technology Reviews* 4(1): 55–70. doi:10.1080/21622515.2015.1068381.

Pacheco, Alan R., Mauricio Moel, and Daniel Segrè. 2019. "Costless Metabolic Secretions as Drivers of Interspecies Interactions in Microbial Ecosystems." *Nature Communications* 10(1): 103. doi:10.1038/s41467-018-07946-9.

Peng, Huadong, Alexander P. S. Darlington, Eric J. South, Hao-Hong Chen, Wei Jiang, and Rodrigo Ledesma-Amaro. 2024. "A Molecular Toolkit of Cross-Feeding Strains for Engineering Synthetic Yeast Communities." *Nature Microbiology* 9(3): 848–63. doi:10.1038/s41564-023-01596-4.

Peng, Xuefeng "Nick," Sean P Gilmore, and Michelle A O'Malley. 2016. "Microbial Communities for Bioprocessing: Lessons Learned from Nature." *Current Opinion in Chemical Engineering* 14: 103–9. doi:10.1016/j.coche.2016.09.003.

Raajaraam, Lavanya, and Karthik Raman. 2024. "Modeling Microbial Communities: Perspective and Challenges." *ACS Synthetic Biology* 13(8): 2260–70. doi:10.1021/acssynbio.4c00116.

Rafieenia, Razieh, Eliza Atkinson, and Rodrigo Ledesma-Amaro. 2022. "Division of Labor for Substrate Utilization in Natural and Synthetic Microbial Communities." *Current Opinion in Biotechnology* 75: 102706. doi:10.1016/j.copbio.2022.102706.

Rafieenia, Razieh, Cinzia Klemm, Piotr Hapeta, Jing Fu, María Gallego García, and Rodrigo Ledesma-Amaro. 2024. "Designing Synthetic Microbial Communities with the Capacity to Upcycle Fermentation Byproducts to Increase Production Yields." *Trends in Biotechnology* 0(0). doi:10.1016/j.tibtech.2024.10.015.

Rapp, Kent M, Jackson P Jenkins, and Michael J Betenbaugh. 2020. "Partners for Life: Building Microbial Consortia for the Future." *Current Opinion in Biotechnology* 66: 292–300. doi:10.1016/j.copbio.2020.10.001.

Roman, Magdalena San, and Andreas Wagner. 2018. "An Enormous Potential for Niche Construction through Bacterial Cross-Feeding in a Homogeneous Environment." *PLOS Computational Biology* 14(7): e1006340. doi:10.1371/journal.pcbi.1006340.

Russo, Claudio, Valerio Cirillo, Nausicaa Pollaro, Fabio Terribile, Andrea Chiodini, and Albino Maggio. 2025. "The Global Energy Challenge: Second-Generation Feedstocks on Marginal Lands for a Sustainable Biofuel Production." *Chemical and Biological Technologies in Agriculture* 12(1): 10. doi:10.1186/s40538-025-00729-7.

Scheer, Maurice, Andreas Grote, Antje Chang, Ida Schomburg, Cornelia Munaretto, Michael Rother, Carola Söhngen, et al. 2011. "BRENDA, the Enzyme Information System in 2011." *Nucleic Acids Research* 39(suppl_1): D670–76. doi:10.1093/nar/gkq1089.

739   Sen, Partho, and Matej Orešič. 2023. "Integrating Omics Data in Genome-Scale Metabolic
740   Modeling: A Methodological Perspective for Precision Medicine." *Metabolites* 13(7): 855.
741   doi:10.3390/metabo13070855.

742   Süntar, Ipek, Sümeyra Çetinkaya, Ülkü Selcen Haydaroğlu, and Solomon Habtemariam. 2021.
743   "Bioproduction Process of Natural Products and Biopharmaceuticals: Biotechnological
744   Aspects." *Biotechnology Advances* 50: 107768. doi:10.1016/j.biotechadv.2021.107768.

745   Tantipaibulvut, Sukon, Anawat Pinisakul, Phannee Rattanachaisit, Kadsarin Klatin, Benjaporn
746   Onsriprai, and Kanokwan Boonyaratsiri. 2015. "Ethanol Production from Desizing
747   Wastewater Using Co-Culture of *Bacillus Subtilis* and *Saccharomyces Cerevisiae*." *Energy*
748   *Procedia* 79: 1001–7. doi:10.1016/j.egypro.2015.11.600.

749   Varma, A, and B O Palsson. 1994. "Stoichiometric Flux Balance Models Quantitatively Predict
750   Growth and Metabolic By-Product Secretion in Wild-Type Escherichia Coli W3110."
751   *Applied and Environmental Microbiology* 60(10): 3724–31.
752   doi:10.1128/aem.60.10.3724-3731.1994.

753   Wagner, Andreas. 2022. "Competition for Nutrients Increases Invasion Resistance during
754   Assembly of Microbial Communities." *Molecular Ecology* 31(15): 4188–4203.
755   doi:10.1111/mec.16565.

756   Wang, Yanxia, Zijian Wan, Yueting Zhu, Haibo Hu, Yujia Jiang, Wankui Jiang, Wenming Zhang,
757   and Fengxue Xin. 2023. "Enhanced 1,3-Propanediol Production with High Yield from
758   Glycerol through a Novel Klebsiella–Shewanella Co-Culture." *Biotechnology for Biofuels*
759   *and Bioproducts* 16(1): 50. doi:10.1186/s13068-023-02304-4.

760   Wilken, St Elmo, Mohan Saxena, Linda R. Petzold, and Michelle A. O'Malley. 2018. "In Silico
761   Identification of Microbial Partners to Form Consortia with Anaerobic Fungi." *Processes*
762   6(1): 7. doi:10.3390/pr6010007.

763   Williams, C. Luke, Tyler L. Westover, Rachel M. Emerson, Jaya Shankar Tumuluru, and Chenlin
764   Li. 2016. "Sources of Biomass Feedstock Variability and the Potential Impact on Biofuels
765   Production." *BioEnergy Research* 9(1): 1–14. doi:10.1007/s12155-015-9694-y.

766   Xu, Lei, and Ulrike Tschirner. 2011. "Improved Ethanol Production from Various Carbohydrates
767   through Anaerobic Thermophilic Co-Culture." *Bioresource Technology* 102(21): 10065–
768   71. doi:10.1016/j.biortech.2011.08.067.

769   Zaramela, Livia S, Oriane Moyne, Manish Kumar, Cristal Zuniga, Juan D Tibocha-Bonilla, and
770   Karsten Zengler. 2021. "The Sum Is Greater than the Parts: Exploiting Microbial
771   Communities to Achieve Complex Functions." *Current Opinion in Biotechnology* 67: 149–
772   57. doi:10.1016/j.copbio.2021.01.013.

773   Ziganshin, Ayrat M., Jan Liebetrau, Jürgen Pröter, and Sabine Kleinsteuber. 2013. "Microbial
774   Community Structure and Dynamics during Anaerobic Digestion of Various Agricultural
775   Waste Materials." *Applied Microbiology and Biotechnology* 97(11): 5161–74.
776   doi:10.1007/s00253-013-4867-0.

BiGG

Choose organisms
and models

Check the model quality

Choose environments
and parameters

Monoculture simulation

Community simulation

$$v_{s_j} = \frac{V_{max_j} S_j}{K_{m_j} + S_j}$$

$$\max \mu = c^T v$$
$$A.v = 0$$
$$v_l \leq v \leq v_u$$

$$\frac{dX_i}{dt} = \mu_i X_i$$
$$\frac{dS_j}{dt} = \sum_i v_{s_j} X_i$$
$$\frac{dP_k}{dt} = \sum_i v_{p_k} X_i$$

v_3

v_1

v_2

Flux Variability Analysis

Dynamic Flux Balance Analysis

Concentration

Time

Analyse results

Concentration

Metabolites

Choose best system and environment

Experimental validation

Environments

Pyruvate
Ethanol
Hydrogen
Acetate
Putrescine
Glutathione
D-Lactate
Fumarate
Formate
Glycerol
Succinate
Gluconate
Spermidine
L-Lactate
Propane-1,2-diol
Propane-1,3-diol
2,3 Butanediol
Methanol
Xylitol
Catechol
Butyrate
Adipic acid
Butanol
Sorbitol
Propionate

Products

Anaer_Rich
Anaer_Min
Aer_Rich
Aer_Min

Product Class

- Diol
- Organic acid
- Carboxylic acid
- Alcohol
- SCFA
- Phenol
- Tripeptide
- Triol
- Gas
- Polyamine
- Sugar alcohol

Mean Productivity Ratio

High

3.89

-0.26