Awakening of the RuMP cycle for synthetic methylotrophy in the thermophile *Parageobacillus thermoglucosidasius*.

Miguel Paredes-Barrada[1], Annemieke S. Mathissen[1], Roland A. van der Molen[1], Pablo J. Jiménez-Huesa[1], Machiel Eduardo Polano[1], Stefano Donati[2], Miriam Abele[3], Christina Ludwig[3], Richard van Kranenburg[1] and Nico J. Claassens[1].

[1]Laboratory of Microbiology, Wageningen University, Stippeneng 4, 6708 WE, Wageningen, The Netherlands
[2]Novo Nordisk Foundation Center for Biosustainability, Søltofts Plads, 220, 212F, 2800 Kgs. Lyngby, Denmark.
[3]Technical University of Munich, Germany; TUM School of Life Sciences, Bayerisches Zentrum für Biomolekulare Massenspektrometrie (BayBioMS), Gregor-Mendel-Strasse 4, 85354 Freising, Germany.

Abstract

Given sustainability and scalability concerns of using sugar feedstocks for microbial bioproduction of bulk chemicals, widening the feedstock range for microbial cell factories is of high interest. Methanol is a one-carbon alcohol that stands out as an alternative feedstock for the bioproduction of chemicals, as it is electron-rich, water-miscible and can be produced from several renewable resources. Bioconversion of methanol into products under thermophilic conditions (>50°C) could be highly advantageous for industrial biotechnology. However, thermophilic natural methylotrophic microorganisms are typically not suitable for this purpose. Hence, we set out to implement synthetic methanol assimilation in the emerging thermophilic model organism *Parageobacillus thermoglucosidasius.* We engineered *P. thermoglucosidasius* to be strictly dependent for its growth on methanol assimilation via the core of the highly efficient ribulose monophosphate (RuMP) cycle, while co-assimilating ribose. Surprisingly, this did not require heterologous expression of RuMP enzymes. Instead, by laboratory evolution we awakened latent, native enzyme activities to form the core of the RuMP cycle. We obtained fast methylotrophic growth in which ~15% of biomass was strictly obtained from methanol. This work paves the way for developing a thermophilic bioproduction platform based on renewable methanol.

Keywords: *P. thermoglucosidasius,* thermophile, synthetic methylotrophy, RuMP cycle, methanol dehydrogenase.

Introduction

The use of microbial cell factories is a promising strategy to produce chemicals more sustainably. Several chemical products such as ethanol and lactate are already produced at a commercial scale by using this approach[1–5]. Sugar-based feedstocks are mostly used for the manufacturing of bio-based products, but there is an interest to move towards the use of alternative feedstocks to make bio-based manufacturing more sustainable and less dependent on agricultural land[6,7]. Alternative sustainable feedstocks should be generated from renewable resources and ideally not be dependent on agriculture, as this can potentially lead to competition with food production for production of bulk products, and a further reduction of the habitat for natural biodiversity.

Among alternative carbon feedstocks, one-carbon feedstocks such as formic acid, carbon monoxide and methanol are promising candidates, given that they can be produced from renewable resources. Among these feedstocks, methanol stands out because it is both water-miscible, electron rich, and its renewable production is already being scaled up. Methanol can be produced from biomass waste (bio-methanol) or from CO_2, water, and renewable energy (e-methanol)[6]. These characteristics make methanol suitable to sustainably produce the wide array of chemicals needed by society and appealing from a bioprocess and environmental perspective.

In nature there are several pathways known to enable growth on methanol as a sole carbon and energy source, such as the serine cycle, the reductive glycine pathway, or the ribulose monophosphate (RuMP) cycle. Among these pathways, the RuMP cycle is known to be one of the most efficient methanol assimilation pathways[6]. In the RuMP cycle, methanol is first oxidized to formaldehyde via methanol dehydrogenase (Mdh). Formaldehyde is assimilated by condensation with ribulose 5-phosphate (Ru5P) to form hexulose 6-phosphate (H6P), in a reaction catalyzed by hexulose 6-phosphate synthase (Hps). H6P is isomerized to fructose 6-phosphate (F6P), via 6-phospho-3-hexuloisomerase (Phi). F6P can be further converted into biomass via several routes dependent on the exact variant of the RuMP cycle[6]. In the most ATP-efficient versions of the RuMP cycle, F6P is subsequently phosphorylated to fructose 1,6-bisphosphate (F1,6P), which is split into the correspondent trioses phosphate, glyceraldehyde 3-phosphate (G3P) and dihydroxyacetone phosphate (DHAP). A part of these triose phosphate molecules will be further metabolized following the glycolytic route, and a part of the G3P will be redirected towards the RuMP cycle to replenish the precursor Ru5P (Figure 1).

Figure 1: Metabolic pathway representing the most ATP-efficient variant of the RuMP cycle[6]. The crucial enzymes for methylotrophy, Mdh, Hps and Phi are highlighted in bold. Mdh: methanol dehydrogenase, Hps: hexulose-6-phosphate synthase, Phi: 6-phospho 3-hexuloisomerase, RpiA: R5P isomerase, Rpe: Ru5P epimerase, Tkt: transketolase, Tal: transaldolase, Pfk: phosphofructokinase, Fba: F1,6P aldolase, TpiA: triose phosphate isomerase.

As methanol oxidation, which is the first step of microbial methanol assimilation, is a thermodynamically challenging reaction ($\Delta_rG'^m$=30.5 kJ/mol)[8], methanol oxidation represents a bottleneck for the implementation of methanol-based biotechnology[9,10]. However, it has been suggested that thermophilic organisms, which are known to have higher enzymatic rates, could be a solution to speed up the methanol oxidation bottleneck[11]. Moreover, the use of thermophilic organisms for biobased manufacturing can reduce the amount of energy needed to cool the bioreactors, as well as the risk of contamination by mesophilic organisms[12]. *Bacillus methanolicus* is the only well-known moderate thermophilic microorganism that can grow on methanol. It has an optimal growth temperature of 50°C and possesses the efficient RuMP cycle for growth on methanol as sole carbon and energy source (Figure 1)[13–15]. Even though *B. methanolicus* has been regarded as a promising thermophilic bacterial host for methanol-based manufacturing, it is notorious for being hard to cultivate. Despite many years of research, *B. methanolicus* lacks effective molecular tools, e.g., to modify its genome, which makes it a suboptimal microbial chassis for chemical production.

It is a major bottleneck for the realization of thermophilic methanol-based bioproduction that there is no
natural thermophilic methylotrophic organism that is suitable for industrial bioproduction. An alternative
approach to obtain a suitable thermophilic microbial chassis for chemical production from methanol, is
to engineer synthetic methylotrophy in a versatile, easier to handle thermophilic bacterium, with a
broad range of genetic tools available. *Parageobacillus thermoglucosidasius* (formerly known as
Geobacillus thermoglucosidasius) is a facultatively anaerobic thermophilic bacterium, which is
emerging as a promising thermophilic model-organism. During the last years, an increasing number of
genetic tools has been developed to effectively engineer its genome, which has harnessed its
potential as a microbial chassis for the production of bulk and fine chemicals[16–19]. Overall, this makes
P. thermoglucosidasius a promising thermophilic microbial chassis to engineer synthetic
methylotrophy[16]

In the last decades, scientific efforts towards implementing synthetic methylotrophy in non-
methylotrophic bacteria, yielded successful results in mesophilic model microorganisms. Full
methylotrophy has been implemented in *Escherichia coli* and *Saccharomyces cerevisiae* [20–26]*,* as well
as partial methylotrophy in *Pseudomonas putida* and *Bacillus subtilis*[27,28]*,* but such life-style change
has not been demonstrated yet for a thermophilic species. Implementing synthetic methylotrophy in a
non-methylotrophic microorganism is a challenging process, which implies a holistic rewiring of central
carbon and energy metabolism, as well as a readjustment of cellular homeostasis. Because of such
complexity, typically efforts to engineer methylotrophy relied on a mixed approach, including rational
engineering and adaptive laboratory evolution (ALE)[20–30]. These mixed approaches are often based
on rational engineering of methanol-dependent strains as a starting point for ALE. Methanol-
dependence can be achieved by one or more gene knockouts that make the cells unable to grow
solely on a certain auxiliary carbon substrate unless they co-assimilate methanol. Such deletion
strains are then normally complemented with heterologous enzymes that enable methanol and
formaldehyde assimilation, such as Mdh and other enzymes. After methanol-dependency and
heterologous enzymes are introduced, ALE is typically performed to enhance functional operation of
methanol assimilation. Examples of rationally engineered methanol-dependent bacterial strains co-
assimilating methanol and ribose (*rpe* deletion), xylose (*rpiA* deletion) or pyruvate (*tpiA or fba*
deletions) can be found in literature [21,22,24,27,30]. So far, in all cases that synthetic methylotrophy has
been engineered in a non-methylotrophic organism, a heterologous Mdh was overexpressed to enable
methanol oxidation in these strains [20,22,25,28,31] . The most used Mdh's in the field have been the ones
from *B. methanolicus* (BmMdh), *Geobacillus stearothermophilus* (GsMdh), *Corynebacterium*
glutamicum (CgMDH) and an engineered *Cupriavidus necator* N1 Mdh (CnMdh-CT4-1)[21,24,26,27,32,33]. In
this work, we present the first instance of synthetic partial methylotrophy via a native, promiscuous
alcohol dehydrogenase.

In the present study, we used a mixed rational and evolutionary approach to engineer partial synthetic
methylotrophy in a thermophilic bacterium for the first time, based on the methods and insights from
previous efforts in mesophiles. However, unlike previous efforts in mesophiles, we could harness via
laboratory evolution, the potential of the latent, native RuMP cycle enzymes without the expression of

any heterologous enzymes. In these evolved strains of *P. thermoglucosidasius*, we demonstrated

methanol-dependent growth with ribose co-assimilation. In addition, we discovered and benchmarked

via *in vivo* assays a novel, thermophilic methanol dehydrogenase native to *P. thermoglucosidasius*.

We believe this work is an important milestone in the path towards the realization of a thermophilic

microbial chassis to produce green chemicals from methanol.

Results

P. thermoglucosidasius harbors both native formaldehyde oxidation and assimilation enzymes.

To convert the non-methylotrophic *P. thermoglucosidasius* strain into a synthetic methylotroph, we

started by exploring the native formaldehyde conversion enzymes. Formaldehyde is not only a central

intermediate in the assimilation of methanol, but also a highly toxic metabolite as it can crosslink DNA

and proteins at low concentrations[20]. Hence most organisms, including non-methylotrophic bacteria

have developed different ways to convert formaldehyde into less toxic compounds.

A common way to detoxify formaldehyde found in microorganisms is its oxidation into formate and the

subsequent oxidation of formate into CO_2[34]. In *P. thermoglucosidasius* we identified a gene annotated

as *adhP* (ALF09200.1), which encodes a protein with 37 % homology to the enzyme AdhA from

Bacillus subtilis, a bacillithiol-dependent formaldehyde dehydrogenase with a detoxification role in this

organism [27,34]. We hypothesized that if AdhP was a formaldehyde detoxification system in *P.*

thermoglucosidasius, this protein would be upregulated upon addition of methanol as its

(promiscuous) oxidation could lead to elevated formaldehyde levels in the cell. Analysis of the wild-

type (WT) strain proteome revealed that AdhP was more than 18-fold overproduced when methanol

was added (8 g/L; Figure 2), hinting that AdhP could, indeed be involved in formaldehyde oxidation

and detoxification.

As observed in several previous studies that established synthetic methylotrophy via the RuMP cycle,

formaldehyde oxidation is usually needs to be eliminated[20,22–28,31]. Hence, we deleted the gene

encoding AdhP by homologous recombination and using a ThermoCas9 counterselection system [35].

We tested if there was an increased sensitivity to different concentrations of formaldehyde in the

Δ*adhP* strain in comparison to the WT, however no differences were found (Table S1).   We

hypothesized that this dissonance between the results obtained from the proteome analysis and the

similar formaldehyde sensitivity for the Δ*adhP* strain, could be explained by the existence of

complementary formaldehyde detoxification systems in *P. thermoglucosidasius.*

In *B. subtilis*, a second formaldehyde detoxification system is known. This second system involves the

enzymes HxlA and HxlB, which are homologues of the RuMP enzymes Hps and Phi [34,36,37] . These

enzymes can assimilate formaldehyde, by condensation with Ru5P and its further conversion into the

central metabolite F6P. *P. thermoglucosidasius* may also have this capacity as it natively encodes

HxlA and HxlB proteins (with 77.6 % and 76.7 % amino acid homology to the *B. methanolicus*

proteins). We observed that, while HxlA was not upregulated in the *P. thermoglucosidasius* WT

proteome during growth in the presence of methanol, HxlB was three-fold upregulated in the presence

of methanol (Figure 2). Therefore, we hypothesized that this assimilation route may also play a role in

formaldehyde detoxication in *P. thermoglucosidasius.* Based on this knowledge, we aimed to exploit
the native formaldehyde assimilation enzymes of *P. thermoglucosidasius* to harness synthetic
methanol assimilation. As AdhP may wastefully convert formaldehyde into formate and CO_2, which
would divert carbon from biomass synthesis, we decided to use the *P. thermoglucosidasius* Δ*adhP*
strain to further establish synthetic methylotrophy.

Awakening the latent RuMP cycle enzymes in *P. thermoglucosidasius* via the evolution of a
methanol-dependent strain*.*

To realize methanol assimilation via the RuMP in *P. thermoglucosidasius,* we developed a strain that
is strictly dependent on methanol assimilation during growth on ribose (Figure 3A). Hereto, we
disrupted a part of the native ribose assimilation route, by deleting the gene *rpe*, which encodes for the
enzyme ribulose-5-phosphate-3-epimerase*.* Rpe catalyzes the interconversion of ribulose to xylulose,
and its activity is essential for growth on ribose as the sole carbon source. The Δ*adhP* Δ*rpe* strain was
obtained with the ThermoCas9 counterselection tool, and we confirmed that this strain could not grow
with ribose as a sole carbon source. We could then use this strain to develop a methanol-assimilating
strain.

Methanol oxidation, the first step of methanol assimilation, is a crucial step to implement synthetic
methylotrophy. Typically, in synthetic methylotrophy a heterologous Mdh is introduced to enable
methanol oxidation [21,22,27,29,30]. *P. thermoglucosidasius* does not possess an annotated Mdh, however
it possesses many alcohol dehydrogenases that could potentially take this role. Alcohol
dehydrogenases are known to be promiscuous enzymes, which accept diverse types of alcohols as
substrates. An example of this is the methanol dehydrogenase from *B. methanolicu*s MGA3, which
has been demonstrated to oxidize propanol, butanol, and ethanol, even at higher rates than methanol
itself [38]. Thus, we hypothesized that via evolution of the Δ*adhP* Δ*rpe* methanol-dependent strain we
could potentially exploit a promiscuous side activity of a native alcohol dehydrogenase from *P.*
thermoglucosidasius, to take up the role of a methanol dehydrogenase.

Together with the natively present formaldehyde assimilation system encoded by the operon *hxlAB*,
we hypothesized that we could evolve *P. thermoglucosidasius* Δ*adhP* Δ*rpe* to oxidize and assimilate
methanol into biomass*.* To do this, we used the selective pressure dictated by the inability of the
Δ*adhP* Δ*rpe* strain to grow on ribose as sole substrate. The methanol dependent *P.*
thermoglucosidasius Δ*adhP* Δ*rpe* was cultivated in minimal TMM medium in the presence of ribose (4
g/L) with or without methanol (8 g/L). After three weeks, growth was observed in two independent
cultures containing methanol and ribose, but not in the ones containing only ribose. Two single
methanol-dependent isolates were isolated from these parallel cultures and designated PTM1 and
PTM2.

Figure 2: Average protein abundance ratios of the putative formaldehyde dissimilatory and assimilatory detoxification systems encoded by *adhP* and *hxlAB* in *P. thermoglucosidasius* WT or PTM1 cultures, supplemented with either mannose or ribose and methanol.

This result suggested that the native enzymatic machinery of the Δ*adhP* Δ*rpe* strain had evolved to co-assimilate ribose and methanol. To confirm the methanol dependency of the evolved PTM1 and PTM2 strains, we cultivated these strains together with the WT *P. thermoglucosidasius* in shake flasks and monitored their growth in the presence of either mannose, ribose, or a mixture of ribose and methanol as carbon sources (Figures 3B, S1). After an extended lag-phase, both PTM1 and PTM2 could grow in the presence of ribose and methanol, while not on ribose alone. These results suggested that the evolved *P. thermoglucosidasius* strains assimilated methanol into biomass via native enzymes.

Figure 3: A) Methanol-dependency selection scheme dictated by the *rpe* gene deletion and used to evolve synthetic methylotrophy with ribose co-assimilation. B) Growth curves of either the WT, or the

205 PTM1 partially methylotrophic strain of *P. thermoglucosidasius* cultivated in minimal media
206 supplemented with different carbon sources. These results confirmed the methanol dependency of the
207 evolved strain PTM1.

Validating methanol assimilation via [13]C labelling experiments

To validate methanol assimilation into biomass, we performed labelling experiments with [13]C-labelled
methanol. The evolved partially methylotrophic strains PTM1 and PTM2 were cultivated in minimal
TMM medium with [12]C-ribose (4 g/L) and [13]C-labelled methanol (8 g/L). Cell pellets were analyzed to
investigate the incorporation of the [13]C-labelled methanol in the proteogenic amino acids serine and
alanine. 3-phosphoglycerate (3PG) is the precursor of serine, while pyruvate is the precursor of
alanine (Figure 4A). 3PG is a glycolytic intermediate and pyruvate the product of glycolysis, therefore
these two metabolites are derived from the split of F6P into two triose phosphate molecules. We
hypothesized that if methanol was co-assimilated with ribose via the RuMP enzymes, one of the
carbon atoms of F6P should be labelled. Thus, 50% of the 3PG and pyruvate molecules should
contain one labelled carbon atom, and so should their corresponding amino acid products, serine, and
alanine (Figure 4A). The results of the analysis revealed that in the strain PTM1 47.5 % of the alanine
molecules contained one [13]C atom, while 46.7% of the serine molecules were labeled with one atom of
[13]C, matching well the expected values (Figure 4B). The labelling results for the strain PTM2 showed
slightly lower labelling, , being 39.5% and 38.3% for serine and alanine. The latter results on PTM2
suggested that this evolved strain possibly had leaky conversion of ribose into biomass independent
from methanol assimilation. Overall, we concluded that these results validated methanol assimilation
via the RuMP cycle enzymes in the partially methylotrophic evolved strains PTM1 and PTM2, and
according to these results, we calculated that methanol accounts for at least ~15% of the biomass in
the strain PTM1.

Figure 4: A) Metabolic scheme showing the expected labelling pattern of serine and alanine assuming
co-assimilation of [12]C-ribose and [13]C-methanol B) Relative abundance of [13]C-labelled carbons present

in the proteogenic amino acids alanine and serine, in cells of the partially methylotrophic evolved strain

PTM1 cultivated in the presence of a mixture of natural ^{12}C ribose and ^{13}C-labelled methanol. The WT

strain cultivated on natural ^{12}C ribose was used as control to account for naturally occurring ^{13}C atoms.

Whole genome sequencing of the evolved strains revealed crucial mutations for methanol assimilation.

The genomes of the evolved partially methylotrophic *P. thermoglucosidasius* strains PTM1 and PTM2,

as well as the parental strain Δ*adhP* Δ*rpe*, were sequenced and analyzed for mutations that occurred

during the evolution towards partial methylotrophy. A list with all mutations found in the evolved strains

can be found in the supplementary material (Tables S2, S3).

Among the observed mutations in both strains, a clear overlap was observed, as mutations were found

in the same specific genes in both evolved strains. We hypothesize that these overlapping mutations

were related to the methylotrophic phenotype observed in both evolved strains. The first two of these

overlapping mutated genes were related to methanol oxidation and assimilation via the latent RuMP

cycle enzymes in *P. thermoglucosidasius.*

In relation to methanol oxidation, both strains had different mutations upstream of the coding region of

adhT gene, an uncharacterized alcohol dehydrogenase from *P. thermoglucosidasius.* In PTM1

multiple substitutions occurred, while in PTM2 a large 369-bp deletion occurred in that region (Table

S3). We hypothesized that AdhT is a promiscuous alcohol dehydrogenase with methanol

dehydrogenase capabilities and these mutations upstream of the coding sequence increased

transcription and/or translation of this enzyme to allow faster methanol conversion via AdhT.

Moreover, both evolved strains had a mutation (G->A) in codon 26 of the *hxlR* gene, resulting in an

amino acid substitution (M26I). HxlR is a helix-turn-helix (HTH) transcriptional regulator controlling the

expression of the operon *hxlAB,* which encodes central enzymes of the RuMP cycle. We hypothesized

that the M26I mutation could be enhancing the expression of *hxlA* and *hxlB* and leading to an

upregulation of these crucial enzymes for methanol assimilation.

Furthermore, the strain PTM1 has a large 23 KB deletion involving the whole *rbsDACBK* ribose operon

and other genes. This seems a counterintuitive mutation as these strains was evolved while co-

utilizing ribose. In the genome of *P. thermoglucosidasius* there is a second copy of the ribose

transporter encoding the proteins *rbsABC*, but the genes *rbsD* and *rbsK,* which encode the proteins

activating ribose to R5P, are not duplicated. Ribose activation is a necessary step for ribose and

methanol co-assimilation, which indicates that this reaction can be possibly catalyzed by other

enzymes. However, this remains an open question. The strain PTM2 does not possess this large

deletion, but shows a truncation in the gene *rbsR,* the transcriptional regulator of the *rbsDACBK* ribose

operon.

The *ptsG* gene encoding a glucose phosphotransferase system, was truncated in both PTM1 and

PTM2 evolved strains. Although the mutations leading to the truncation of PtsG are not identical in

each of the strains, this mutation was found in both parallelly evolved strains, so we think it might be
important for the methylotrophic phenotype. However, the link between the found truncations and
methylotrophy remains unclear.

Another interesting mutation that may have a plausible link to the RuMP operation was found only in
PTM1. In this strain we found a SNP (1984003T>A), which caused an amino acid substitution in the
coding sequence of the *tkt* gene (T184A). *tkt* encodes the transketolase, an enzyme which catalyzes
two of the reactions of the RuMP cycle (G3P + F6P = E4P + Xu5P; G3P + S7P = X5P + R5P; Figure
1). The Tyr184 residue is highly conserved among different species of microorganisms, and given its
vicinity to the cofactor thiamine diphosphate binding site, this residue has been regarded as essential
for the transketolase activity[39–41]. The activity of Tkt is essential for methanol and ribose co-
assimilation via the RuMP, and therefore we believe this amino acid substitution did not result in a
death variant. However, what is the actual effect of this mutation remains an open question.

Proteomics analysis revealed that crucial enzymes for the RuMP cycle were upregulated.

To investigate the effect of some of the mutations observed in the *P. thermoglucosidasius* evolved
strains, we aimed to analyze their proteome and compared it with the one of the *P.*
thermoglucosidasius WT strain (Figure S2). Because the strain PTM2 showed lower methanol
assimilation in the [13]C-labelling, we decided to only continue analyzing the proteome of the PTM1
strain (Figure 2, 5). The PTM1 and WT strains were cultivated in the presence of either mannose or a
mixture of ribose and methanol as carbon sources.

In relation to the methanol oxidation and assimilation capabilities, we confirmed that the putative
methanol dehydrogenase AdhT, as well as HxlA and HxlB were significantly upregulated in the PTM1
strain in comparison to the WT (Figure 2). AdhT was 63-fold more abundant in the PTM1 strain than
in the WT, when a mixture of ribose and methanol were utilized as carbon source. This result
confirmed our hypothesis that the mutations observed in the upstream region of the *adhT* gene, in
PTM1, increased the expression of this alcohol dehydrogenase. The enzyme HxlA was more than 56-
fold overproduced in the PTM1 strain in comparison with the WT when a mixture of ribose and
methanol was used as carbon source, while it was more than 16-fold overproduced in the PTM1 when
grown on mannose. The enzyme HxlB was more than 32-fold overproduced in the evolved PTM1
strain in comparison with the WT when a mixture of ribose and methanol was used as carbon source,
while it was more than 15-fold overproduced when grown on mannose (Figure 2). These results
indicated that the mutation observed in the HxlR transcriptional regulator in the evolved partially
methylotrophic strains, increased the expression of the operon *hxlAB* without methanol present and
further boosted the expression with methanol, thus enabling efficient assimilation of formaldehyde into
fructose-6-phosphate. In conclusion, this strongly suggest that the native AdhT, HxlA and HxlB drive
methanol assimilation in *P. thermoglucosidasius*.

In *B. subtilis*, it was shown by X-ray crystallography that HxlR undergoes a conformational change
induced by the presence of formaldehyde, which promotes the formation of an intra-helical cysteine-
lysine bond, and allows DNA binding to activate the expression of *hxlAB*[36,42]. The domains of the HxlR

protein from *B. subtilis* and *P. thermoglucosidasius* responsible for this conformational change and
activation of the expression of *hxlAB* are conserved. When comparing the corresponding amino acid
residue 26 of the HxlR protein from both microorganisms, we have observed that this amino acid is
situated in the formaldehyde recognition pocket, exactly in front of the cysteine-lysine bridge which
activates HxlR. In light of this observation, we hypothesize that the M26I substitution found in the
evolved strains, could lead to a partial conformational change in the off state of HxlR, potentially
causing the constitutive increased expression of HxlAB observed in the evolved strain PTM1 (Figure
2). This hypothesis could explain why in the evolved strain PTM1, HxlAB are produced in higher
amounts even when methanol is not present in the media.

Figure 5: Average protein abundance ratios of AdhT, Zwf and Aldh when cultures of the WT and PTM1
strains were supplemented with a mixture of ribose and methanol as carbon source.

In addition, we found that the enzyme glucose-6-phosphate dehydrogenase (Zwf), was more than
325-fold overproduced in the evolved strain (Figure 5). This enzyme is part of the oxidative part of the
RuMP cycle that diverts carbon from biosynthesis to produce NADPH and CO_2, named the RuMP
dissimilatory cycle. This dissimilatory pathway includes the reactions catalyzed by the enzymes G6P
isomerase (encoded by *pgi*) which isomerizes F6P into G6P, G6P dehydrogenase (encoded by *zwf*)
which oxidizes G6P into glucono-1,5-lactone-6P, and 6-phosphogluconate dehydrogenase which
converts gluconate-6P back into the precursor Ru5P and CO_2 (Figure 6). This result could indicate that
the flux of carbon going through the RuMP dissimilatory pathway is considerable, causing that part of
the carbon assimilated via formaldehyde condensation with Ru5P gets lost in the form of CO_2, which is
undesirable to establish full methylotrophy in this bacterium, but could support extra NADPH
generation.

Another interesting highly downregulated enzyme in the PTM1 proteome compared to the WT, was an
enzyme potentially related to formaldehyde dissimilative detoxification. The enzyme annotated as

aldehyde dehydrogenase (AldH), was more than 581-fold downregulated in the evolved partially methylotrophic strain PTM1 relative to WT (Figure 5). In *Corynebacterium glutamicum*, a similar enzyme (CAF20817), which shares ~70% amino acid identity with AldH from *P. thermoglucosidasius*, was confirmed to be part of a dissimilative formaldehyde detoxification pathway[34,43,44]. In *C. glutamicum* this pathway involved the oxidation of formaldehyde to formic acid by the acetaldehyde dehydrogenase and the dissimilation of formate to CO_2. Thus, we hypothesize that the native AldH of *P. thermoglucosidasius* could be fulfilling a similar role, and the observed downregulation of AldH in the evolved PTM1 strain increases the pool of formaldehyde available for methylotrophic biosynthesis.

Figure 6: Metabolic pathway representing methanol and ribose co-assimilation via the RuMP cycle in the evolved partially methylotrophic strain PTM1 of *P. thermoglucosidasius*. Key native enzymes for methylotrophy, AdhT, Hps and Phi are in bold. AdhT: putative methanol dehydrogenase, Hps: hexulose-6-phosphate synthase encoded by *hxlA*, Phi: 6-phospho 3-hexuloisomerase encoded by *hxlB*, Zwf: G6P dehydrogenase, GapA: G3P dehydrogenase, RpiA: R5P isomerase, Rpe: R5P epimerase, Tkt: transketolase, Tal: transaldolase, Pfk: phosphofructokinase, Fba: fructose-1,6-bisphosphate aldolase, TpiA: triose phosphate isomerase, Pgk: phosphoglycerate kinase, Pgm: phosphoglycerate mutase, Eno: enolase, Pk: pyruvate kinase, Pdh: pyruvate dehydrogenase, Cs: citrate synthase, Pgi: phosphoglucose isomerase, Pgd: 6-phosphogluconate dehydrogenase.

In vivo assays confirmed that AdhT is a novel thermophilic methanol dehydrogenase.

To validate that AdhT, the putative native methanol dehydrogenase from *P. thermoglucosidasius*
indeed possessed methanol oxidation capabilities, we investigated the production of formaldehyde by
this enzyme. We utilized two different *in vivo* approaches to detect the activity of AdhT, as methanol
dehydrogenase activity is notoriously hard to measure *in vitro.*

In our first *in vivo* approach we confirmed the enzyme activity at thermophilic temperature (60°C) using
plasmid-based overexpression of AdhT in *P. thermoglucosidasius* WT. In the second approach we
further confirmed its activity and compared it to other commonly used Mdh's at mesophilic temperature
(37°C), in an *E. coli* formaldehyde biosensor strain[45].

For the first approach, *P. thermoglucosidasius* strains overexpressing *adhT* from plasmid under the
transcriptional control of a strong promoter (P1) and a medium strength promoter (P2) were cultivated
in the presence of methanol, and formaldehyde concentration was measured in the culture
supernatants by a colorimetric method[46,47]. The formaldehyde concentration detected in *P.*
thermoglucosidasius culture supernatants overexpressing the *adhT* gene under medium-strength P1
and strong P2 promoters, was respectively 15-fold and 19-fold higher. These results confirmed that
the overexpression of the native AdhT enzyme from *P. thermoglucosidasius* enabled methanol
oxidation, and hence AdhT can act as thermophilic methanol dehydrogenase. Formaldehyde was also
detected in low concentrations in the supernatants cultures of *P. thermoglucosidasius* harbouring an
empty plasmid (Figure 7). This observed modest native methanol oxidation activity in *P.*
thermoglucosidasius cultures, could be attributed to the presence of small amounts of AdhT expressed
from the wild-type genome of *P. thermoglucosidasius* or the promiscuous activity of other alcohol
dehydrogenases.

Figure 7: Average formaldehyde accumulation in culture supernatants of different *P.*
thermoglucosidasius strains cultivated in the presence of methanol and overexpressing *adhT* under

the control of the medium-strength promoter P1 (32.73 µM) and stronger promoter P2 (42.26 µM), or

harboring an empty plasmid (E; 2,14 µM).

The second *in vivo* approach to study the ability of AdhT to convert methanol into formaldehyde was

performed in the *E. coli* HOB formaldehyde biosensor, published by Schann et al. (2024). This

biosensor was engineered to be methionine and isoleucine auxotrophic. Growth of the biosensor can

be rescued by the production of formaldehyde, which together with pyruvate can be converted to

methionine and isoleucine via the combined activity of the 4-hydroxy-2-oxobutanoate (HOB) aldolase

(HAL) and HOB transaminase (HAT) reactions[45]. In this experiment, either AdhT or other reference

Mdh's (CgMdh and BsMdh) were overexpressed in the *E. coli* formaldehyde biosensor and cultivated

in varying concentrations of methanol. Growth of the biosensor implied that methanol was oxidized to

formaldehyde, concluding that the overproduced Mdh possessed methanol dehydrogenase activity.

When comparing AdhT performance to the other two tested Mdh's, AdhT performed similar to the

CgMdh, while BsMdh allowed better growth at lower methanol concentration (Figure 8). This likely

indicates that AdhT has a relatively low affinity for methanol and required higher methanol

concentrations to allow growth of the biosensor strain.

Overall the results from the mesophilic and thermophilic *in vivo* approaches demonstrated that the

enzyme AdhT of *P. thermoglucosidasius* can be regarded as a newly discovered, thermophilic

methanol dehydrogenase.

Figure 8: Average maximum OD_{600} of the *E. coli* formaldehyde biosensor strain, overexpressing AdhT,

BsMdh or CgMdh. The different strains were cultivated in M9 selective media supplemented with

methanol in different concentrations.

Discussion

This work represents a milestone towards the realization thermophilic synthetic methylotrophy in *P. thermoglucosidasius,* a promising thermophilic model organism for industrial use[16]. Our work presents the first example of achieving partial synthetic methylotrophy in a thermophilic organism, using a mixed rational and evolutionary approach. The present study also represents the first instance of establishing synthetic methylotrophy that was only reliant on activation of native latent RuMP cycle enzymes.

We analyzed which mutations led to methanol and ribose co-assimilation in the evolved partially methylotrophic strain PTM1, which included the modification of *adhT* upstream region and a point mutation in the *hxlR* regulator resulting in an amino acid substitution (M26I). We confirmed via proteomics that these mutations caused the overproduction of AdhT, HxlA and HxlB, which we believe are crucial enzymes for partial methylotrophy via the RuMP cycle. Other authors also demonstrated that during methylotrophic growth via the RuMP cycle, the Mdh, Hps and Phi enzymes were upregulated during evolution towards full methylotrophy and were key for methanol assimilation[22]. This was also demonstrated in the natural methylotroph *B. methanolicus*, in which adding extra copies of *hps* and *phi* (the homologues of *hxlAB*) increased the methanol tolerance and assimilation capacities of this bacterium [38,48].

We furthermore discovered a novel thermophilic methanol dehydrogenase native to *P. thermoglucosidasius,* the enzyme AdhT. In our work, we observed that AdhT was overproduced in the evolved strain PTM1, and hypothesized it was a promiscuous alcohol dehydrogenase with methanol oxidation capabilities. We confirmed that this enzyme is a methanol dehydrogenase via two different *in vivo* approaches based on the AdhT capacity to produce formaldehyde from methanol. As indicated by the growth of the *E. coli* formaldehyde biosensor overexpressing different Mdh's, AdhT was not as efficient as BsMdh at low methanol concentrations (40 mM), but had a comparable methanol oxidation capacity at 80 mM of methanol. This result could hint that AdhT has a higher K_m than BsMdh and a similar K_{cat}, however the exact kinetic properties of the enzyme remain unknown.

We also observed that the protein abundances of Zwf and AldH were, respectively 325 and 581-fold different in the PTM1 strain proteome in comparation with the WT, while no mutations occurred in the promoter or RBS regions of these genes during evolution of the PTM1 strain. We hypothesized that these differences in protein abundance existed due to allosteric regulation, or other types of regulation at the transcriptional, translational, or post-translational level, however this remains an open question.

Other studies have revealed that it is usual to find several formaldehyde detoxification systems in the same microorganism. For instance, *B. subtilis* possesses two different formaldehyde detoxification systems. One based on the HxlAB enzymes, and a second one based on a BSH-dependent aldehyde dehydrogenase encoded by *adhA*. In *P. thermoglucosidasius,* we hypothesize that there are three of these systems, HxlAB, AdhP and the uncharacterized aldehyde dehydrogenase encoded by *aldH*. We base this hypothesis on the fact that AldH was 581-fold less abundant in the evolved PTM1 strain in comparison to the WT. We think that AldH detoxifies formaldehyde by oxidation into formate, which

later gets dissimilated into CO_2, in a similar way as AdhA in *B, subtilis.* We hypothesize that this
drastic downregulation of AldH decreased the dissimilation of formaldehyde to CO_2 and thus increased
its availability for assimilation into biomass.

We believe that the evolved strain PTM1 is a promising thermophilic platform starting point to establish
full synthetic methylotrophy. To accomplish growth on methanol as a sole carbon source in this strain,
first the *rpe* gene should be restored into the genome of the strain PTM1 to allow the full RuMP cycle
to function. ALE is a common strategy to accomplish full methylotrophic growth after reconstitution of
the full RuMP cycle in a partially methylotrophic strain [22,24,30]. In this way, the strain is cultivated on a
mixed carbon source including methanol and an auxiliary carbon source, and during a series of
passages the auxiliary carbon source can be weaned out until growth on methanol as a sole carbon
source is realized. In the case of the *P. thermoglucosidasius* PTM1 strain, after the *rpe* gene is
reconstituted, it could be for example evolved with a mixture of methanol and ribose. The genes *zwf*
and *aldH* could be possible KO targets to make the PTM1 strain more suitable for full growth on
methanol. The *zwf* deletion would eliminate the carbon flux going through the RuMP dissimilatory
cycle, and the *aldH* deletion would decrease formaldehyde detoxification to CO_2, in both cases
increasing the carbon flux from methanol to biomass.

Materials and methods

General cultivation techniques, media composition, bacterial strains, and plasmids.
455

For the cultivation of *E. coli* strains, lysogeny broth (LB) medium was used containing 10 g/L tryptone
(Bacto™), 10 g/L NaCl, and 5 g/L yeast extract (Bacto™). Solid LB medium additionally contained 15
g/L agar. *E. coli* strains were generally cultivated in 10 mL liquid medium in falcon tubes at 37°C and
200 rpm overnight or on solid medium at 37°C overnight. *E. coli* HOB biosensor cultures were
supplemented with 0.25 mM of diaminopimelate acid (DAP).

For general cultivation of *P. thermoglucosidasius* strains, TGP medium was used containing 17 g/L
tryptone (Bacto™), 3 g/L peptone (Sigma Aldrich), 5 g/L NaCl, 2.5 g/L K_2HPO_4, 0.4% w/v glycerol, and
4 g/L sodium pyruvate. Solid TGP medium additionally contained 30 g/L agar. For the experimental
investigation of methanol-dependent growth, formaldehyde concentration determination via NASH
assay and the determination of growth curves of *P. thermoglucosidasius*, thermophilic minimal media
(TMM) was used. TMM contained: 494 mg/L NaCl, 145 mg/L $NaSO_4$, 24.7 mg/L KCl, 3.97 mg/L KBr,
184 mg/L $MgCl_2$·6 H2O, 89.2 mg/L $NaNO_3$, 8.37 g/L MOPS, 3.58 mg/L $FeSO_4$, 716 mg/L tricine, 229
mg/L K_2HPO_4, 0.51 g/L NH_4Cl, 73.5 mg/L $CaCl_2$, 2 µg/L biotin, 5 µg/L thiamine-HCl, 2 µg/L folic acid,
0.1ug/L vitamin B12, 0.127 µg/L spermidine, 174 µg/L spermine tetrahydrochloride, 0.7 mg/L hemin, 1
mg/L $FeCl_3$·6 H2O, 0.18 mg/L $ZnSO_4$·7 H2O, 0.12 mg/L $CuCl_2$·2 H2O, 0.12 mg/L, $MnSO_4$· H2O, and
0.18 mg/L $CoCl_2$·6 H2O, with a variable carbon source. Either D-ribose (4 g/L), glucose (5 g/L),
mannose (10 g/L), methanol (8 g/L) or a mixture of sugars and methanol were added as carbon
source. For the *in vivo* assays involving the *E. coli* HOB biosensor strain, M9 relaxing or selective
media were used (Table S3).

When needed for cloning or plasmid propagation, chloramphenicol (15 μg/L), ampicillin (100 μg/L) or
streptomycin (100 μg/L) was added to *E. coli* cultures, while either chloramphenicol (8 μg/L) or
thiamphenicol (6.5 μg/L) was added to *P. thermoglucosidasius* cultures .

Bacterial strain or plasmid	Description	Source or reference
P. thermoglucosidasius WT	Sporulation-deficient Type strain DSM 2542 with *sigF* gene deletion	49
P. thermoglucosidasius ΔadhP	*P. thermoglucosidasius WT* with *adhP* gene deletion	This work
P. thermoglucosidasius ΔadhP Δrpe	*P. thermoglucosidasius ΔadhP* with *rpe* gene deletion	This work
P. thermoglucosidasius PTM1	Methanol dependent isolate evolved from *P. thermoglucosidasius ΔadhP Δrpe*	This work
P thermoglucosidasius PTM2	Methanol dependent isolate evolved from *P. thermoglucosidasius ΔadhP Δrpe*	This work
P. thermoglucosidasius P1-*adhT*	*P. thermoglucosidasius WT* harboring pJS15A-pRpsl8-*adhT*	This work
P. thermoglucosidasius P2-*adhT*	*P. thermoglucosidasius WT* harboring pJS15A-pLDH-*adhT*	This work
P. thermoglucosidasius EV	*P. thermoglucosidasius WT* harboring pJS15A empty vector	This work
E. coli Neb5a	*E. coli* cloning host. *fhuA2Δ(argF-lacZ) U169 phoA glnV44 Φ80Δ(lacZ)M15 gyrA96 recA1 relA1 endA1 thi-1 hsdR17*	New England Biolabs
E. coli HOB biosensor	Formaldehyde biosensor *E. coli Δasd, ΔfrmRAB, ΔrhmA, rhmA.*	45
HOB –*Bsmdh*	*E. coli* formaldehyde biosensor harboring pZASS-*Bsmdh*	45
HOB –*Cgmdh*	*E. coli* formaldehyde biosensor harboring pZASS-*Cgmdh*	45
HOB –*adhT*	*E. coli* formaldehyde biosensor harboring pZASS-*adhT*	This work
pJS28	Shuttle vector *Geobacillus-E. coli. repB, ColE1, ChlR, oriT, truncated traJ*	50

pJS15A	Shuttle vector *Geobacillus-E. coli. repB, p15a, ChlR, oriT, truncated traJ*	This work
pJS15A-*adhP*KO	pJS15A containing ThermoCas9, a sgRNA targeting the *adhP* gene of DSM 2542	This work
pJS15A-*rpe*KO	pJS15A containing ThermoCas9, a sgRNA targeting the *rpe* gene of DSM 2542	This work
pJS15A-pRpsl8-*adhT*	pJS15A containing *adhT* from *P. thermoglucosidasius*, under the expression of pRpsl8	This work
pJS15A-pLDH-*adhT*	pJS15A containing *adhT* from *P. thermoglucosidasius*, under the expression of pLDH	This work
pZASS_*adhT*	pZASS containing *adhT* from *P. thermoglucosidasius.*	This work

479   Table 1: List of bacterial strains and plasmids used in this study, together with a brief description and
480   the source.

Plasmid assembly and genome modification.

Generally, primers were designed to either introduce BsaI sites and overhangs for Golden Gate
assembly or to provide homology for Gibson assembly (Table S5). All the desired DNA fragments
were amplified using Q5 polymerase (Q5 2x MasterMix, New England Biolabs), isolated from a 1%
agarose gel (Zymogen gel DNA recovery kit) and assembled by either Gibson (50 °C, 1 hour) or
Golden Gate ((37°C, 5 min; 16 °C, 5 min) x 30 ; 37°C, 20 min; 65°C, 20 min). 5 µL of assembly
mixture were transformed by heat shock in Neb5a commercial chemically competent cells (New
England Biolabs) and plasmid presence in bacterial cells was verified by colony PCR using OneTaq
polymerase (OneTaq 2x MM, New England Biolabs). Plasmids were isolated (QIAprep Spin Miniprep
Kit, Qiagen) and the sequence of all newly assembled plasmids were confirmed by Oxford Nanopore
whole plasmid sequencing (Plasmidsaurus).

pZASS_*adhT* was constructed by Gibson assembly by combining two PCR amplified fragments with
overlapping regions, one containing the *adhT* coding sequence, amplified from *P. thermoglucosidasius*
WT genome as a template, and the other containing the pZASS backbone[45]. To obtain the plasmid
pJS15A, a derivative of the Geobacilli:*E. coli* shuttle vector pJS28 (Table X), we replaced by Golden
Gate assembly the origin ColE1 from pJS28 by a lower copy number p15A ori [51]. To construct the
plasmids pJS15A-*adhP*KO and pJS15A-*rpe*KO, first *thermoCas9* under the expression of the *P.*
thermoglucosidasius native cellobiose inducible promoter *Pbgl*, and the sgRNA module including a
scrambled spacer were amplified by PCR and assembled into the backbone pJS15A by Golden Gate.
The resulting vector was later digested with SacI, and 1 KB homology arms (HA) flanking the

upstream and downstream regions of either the *adhP* or *rpe* gene of *P. thermoglucosidasius* were
included via Gibson assembly. The scrambled spacer was replaced by a targeting spacer via Gibson
assembly.

Confirmed plasmids were transformed into *P. thermoglucosidasius* by conjugation, following a slightly
adapted protocol from Hon Wai Wan (2013)[52]. Notable differences were that a 50 mL culture of *P.*
thermoglucosidasius was grown to mid-log phase (~0.7 OD_{600}) in a 250 mL baffled flask and then
centrifuged at 4500xg for 10 minutes before the actual conjugation.

To delete the genes *adhP* and *rpe*, we induced homologous recombination of HA from pJS15A-
*thermocas9-adhP*KO or pJS15A-*thermocas9-rpe*KO onto the genome of *P. thermoglucosidasius* WT
strain and selected against wild-type DNA by using ThermoCas9. One confirmed transformant of *P.*
thermoglucosidasius containing the editing plasmid was cultivated in 10 mL liquid TGP containing
chloramphenicol (8 µg/mL) at 60°C overnight, then transferred (1mL) to fresh 10 mL TGP containing
chloramphenicol (8 µg/mL) and cultivated at 68°C overnight. The resulting culture was then transferred
into fresh TGP containing chloramphenicol (8 µg/mL) and 1% w/v cellobiose and cultivated at 60°C
overnight to induce the *pBgl* promoter and enhance *thermocas9* transcription. Finally, the culture was
plated on a TGP plate with 1% w/v cellobiose using 1 mL, 100 µL, 10 µL, or 1 µL of liquid culture and
cultivated at 55° C. Gene deletions were confirmed by colony PCR.

Evolution of PTM partially methylotrophic strains of *P. thermoglucosidasius.*

To test if the *rpe* gene deletion indeed led to the inability to utilize ribose as a sole carbon source, we
inoculated *P. thermoglucosidasius* ΔadhP Δrpe in different falcon tubes containing 10 mL of TMM
supplemented with ribose in different concentrations (1, 2 and 4 g/L), and incubated them at 60°C and
180 rpm during at least two weeks. Growth was not observed in any of the cultures and therefore we
concluded that this strain was not able to utilize ribose as a sole carbon source.

P. thermoglucosidasius ΔadhP Δrpe was inoculated from glycerol stock in liquid TGP media and
incubated overnight at 60°C and 180 rpm. Having previously removed the preculture media by
centrifugation (5000 rpm, 3 minutes), 100 µL from the preculture were sub-cultured in a falcon tube
containing 10 mL of TMM supplemented with ribose (4 g/L) and methanol (8 g/L). The cells were
incubated at 60°C and 180 rpm for three weeks. After OD_{600} reached 0.6, methanol dependent growth
was confirmed by subculturing the cells in 250 mL baffled flasks containing 25 mL of TMM
supplemented with ribose (4 g/L) or with a mixture of ribose (4 g/L) and methanol (8 g/L). After 24h
only the cultures containing methanol had grown, confirming that the evolved population was methanol
dependent.

The evolved methanol-dependent *P. thermoglucosidasius* population was plated on TGP solid media
from glycerol stocks and incubated at 60°C overnight. Up to ten colonies per strain were picked and
screened for methanol-dependency in falcon tubes containing 10 mL of liquid TMM media containing
ribose (4 g/L) or a mixture of ribose (4 g/L) and methanol (8 g/L). The isolates PTM1 and PTM2 which
only grew when methanol was added, were selected for further analysis.

538   **Analysis of the genomic mutations occurring during evolution of partial methylotrophy.**

539   *P. thermoglucosidasius* genomic DNA was extracted from overnight cultures using the MasterPure™

540   Gram Positive DNA Purification Kit (Lucigen). The quality and concentration of the genomic DNA was

541   verified by Qubit (ThermoFisher) and the genome was sequenced (Illumina, Novogene). The reads

542   were mapped to a genome of reference and assembled in a contig, and the mutations differing

543   between the evolved strains PTM1 and PTM2, and the parental strain were analyzed.

544   **Validation of methanol assimilation via [13]C labelling experiments.**

545   To validate methanol assimilation into biomass, we performed steady-state labelling experiments with

546   [13]C labelled methanol. *P. thermoglucosidasius* WT, PTM1 and PTM2 strains were inoculated from

547   glycerol stocks in liquid TGP media and incubated overnight at 60°C 180 rpm. 100 µL from each

548   preculture were subcultured in 10 mL of TMM, containing natural labelled ribose (4 g/L) and

549   supplemented with [13]C-labelled methanol (8 g/L), having previously removed the preculture media by

550   centrifugation (5000 rpm, 3 minutes). Cells were incubated at 60°C and 180 rpm, grown to late

551   exponential phase (OD~1), subcultured a second time in fresh media with [13]C-labelled methanol and

552   incubated overnight in the same conditions. One mL of culture broth was harvested (OD~1),

553   centrifuged (13000 rpm, 1 minute) and cell pellets were stored at -20 until further processing. Pellets

554   were re-suspended in 200 µL of 6 M HCl and incubated at 110 °C for 16 hours to lyse the cells and

555   hydrolyze the respective proteomes. Hydrolyzed samples were then filtered on a filter plate (AcroPrep

556   Advance 96-well Filter, 0.2 µm membrane, product ID 8019) by positive pressure, and dried in a cold-

557   trap speedvac. Dried samples were then derivatized by adding 35 µL of pyridine (Sigma-Aldrich, cat

558   no. 270407) and 50 µL of MBTSFA + 1% (w/w) TBDMSCIS (Sigma-Aldrich, cat no. 375934), mixed by

559   pipetting and incubated at 60°C for 30 min. 75 µL of the derivatized samples were aliquoted to LC-MS

560   vials with glass inserts for GC-MS analysis within 12h from derivatization on an Agilent 5977 GC-MS

561   system with an Agilent DB-5ms capillary column (30m, inner diameter of 0.25 mm, film thickness of

562   0.25 µm, cat no. 122-5532). Samples were measured in full-scan mode, using a 1:100 split ratio, with

563   the following gradient: start at 160°C, hold for 2 min, ramp to 310 °C at 7 °C/min, hold for 1 min. We

564   considered for further analysis fragments listed in Long C.P. & Antoniewicz (2019) which included the

565   whole C backbone of the relative amino acid (Ala 260, Ser 390)[53]. Raw chromatographic data was

566   integrated using SmartPeak[54]. The relative data was corrected for the natural abundance of isotopes

567   in the derivatization agents used for GCMS analysis[55].

568   **Proteomic sample preparation**

569   *P. thermoglucosidasius* PTM1 and WT strains were inoculated from glycerol stocks in liquid TGP

570   media and incubated overnight at 60°C and 180 rpm. Having previously removed the preculture media

571   by centrifugation (5000 rpm, 3 minutes), 100 µL from each preculture were subcultured in triplicated

572   250 mL baffled flasks containing 25 mL of TMM supplemented with either mannose (10 g/L) or a

573   mixture of ribose (4 g/L) and methanol (8 g/L). Cells were incubated at 60°C and 180 rpm, grown to

574   mid exponential phase (OD_{600}=0.4-0.6), harvested by centrifugation (4700 rpm, 10 min), and stored at

575   -20 until further processing. Samples were processed for mass spectrometry according to the SPEED

protocol with slight modifications as described previously[56,57]. Shortly, cells were resuspended in 100

µL 100% TFA and transferred to clean plastic microtubes which were incubated in a thermocycler at

55 °C. A volume of 900 µL 2 M Tris solution (pH was not adjusted) was added and tubes were

vortexed. The lysed samples were quantified by Bradford assay and stored at -80 °C. A mass of

20 µg of protein per sample was reduced (10 mM TCEP) and carbamidomethylated (55 mM CAA)

for five minutes at 95 °C. The proteins were digested by adding trypsin (proteomics grade, Roche) at

a 1/50 enzyme/protein ratio (w/w) and incubated at 37°C overnight. Digests were acidified by the

addition of 3% (v/v) formic acid (FA) and desalted using self-packed StageTips (five disks per micro-

column, ø 1.5 mm, C18 material, 3 M Empore)[58]. The peptide eluates were dried to completeness

and stored at −80 °C. Before the LC-MS/MS measurement, all samples were freshly resuspended in

12 µl 0.1% FA in HPLC grade water and 5 µl were subjected to LC-ESI-MS/MS.

Proteomics data acquisition Exploris – microflow label-free DDA

Peptides were analyzed on a Vanquish Neo (microflow configuration) coupled to an Orbitrap Exploris

480 mass spectrometer (both Thermo Fisher Scientific). Around five micrograms of peptides were

applied onto a commercially available Acclaim PepMap 100 C18 column (2 µm particle size, 1 mm

ID × 150 mm, 100 Å pore size; Thermo Fisher Scientific) and separated using a stepped gradient with

acetonitrile concentration ranging from 3% to 24% to 31% solvent B (0.1% FA, 3% DMSO in ACN) in

solvent A (0.1% FA, 3% DMSO in HPLC grade water). A flow rate of 50 µl/min was applied. The mass

spectrometer was operated in data-dependent acquisition (DDA) and positive ionization mode. MS1

full scans (360–1300 m/z) were acquired with a resolution of 60,000, a normalized automatic gain

control target value of 100%, and a maximum injection time of 50 ms. Peptide precursor selection for

fragmentation was carried out at a 1.2 seconds cycle time. Only precursors with charge states from

two to six were selected, and dynamic exclusion of 30 s was enabled. Peptide fragmentation was

performed using higher energy collision-induced dissociation and normalized collision energy of 28%.

The precursor isolation window width of the quadrupole was set to 1.1 m/z. MS2 spectra were

acquired with a resolution of 15,000, a fixed first mass of 100 m/z, a normalized automatic gain control

target value of 100%, and a maximum injection time of 40 ms.

LC-MS/MS data analysis

Peptide identification and quantification was performed using the software MaxQuant (version

1.6.3.4)[59] with its built-in search engine Andromeda[60]. MS2 spectra were searched against the NCBI

P. thermoglucosidasius protein database (GCF_001295365.1, downloaded May 2024), supplemented

with common contaminants (built-in option in MaxQuant). Trypsin/P was specified as proteolytic

enzyme. Carbamidomethylated cysteine was set as fixed modification. Oxidation of methionine and

acetylation at the protein N-terminus was specified as variable modifications. Results were adjusted to

1% false discovery rate on peptide spectrum match (PSM) level and protein level employing a

target-decoy approach using reversed protein sequences. Label-Free Quantification (LFQ)[61] and

iBAQ[62] intensities were computed. The minimal peptide length was defined as seven amino acids and

the "match-between-runs" functionality was disabled. The ProteinGroups.txt result file from MaxQuant

was used as input for Omicsviewer 1.1.5[63] to perform t-tests of differential protein abundances of *P.*

thermoglucosidasius in the presence of mannose or ribose and methanol. Missing values were

imputed by a protein-specific constant value, which was defined as the lowest detected protein-

specific LFQ-value over all samples divided by two. Additionally, a maximal imputed LFQ value was

defined as 15% quantile of the protein distribution from the complete dataset.

Formaldehyde concentration estimation

P. thermoglucosidasius E, P1-*adhT* and P2-*adhT* strains were inoculated from glycerol stocks in liquid

TGP media and incubated overnight at 60°C and 180 rpm. 100 µL from each preculture were

subcultured in triplicated falcon tubes containing 10 mL of TMM supplemented with glucose (5 g/L),

methanol (8 g/L) and thiamphenicol (6.5 mg/L) and incubated at 57°C and 180 rpm. One mL samples

were taken at early stationary phase (OD_{600}~1) and formaldehyde concentration was estimated by

using a colorimetric method[46].

***adhT* complementation *in vivo* experiment in *E. coli* HOB biosensor**

E.coli HOB biosensor strains harboring the plasmids pZASS_*Bsmdh* and pZASS_*Cgmdh* were kindly

donated by Schann et al. (2024). pZASS_*adhT* was cloned and transformed by heat shock in the *E.*

coli HOB biosensor in house. Transformants of the HOB-*adhT* strain were confirmed via colony PCR

as explained above. The strains HOB-*Bsmdh*, HOB-*Cgmdh* and HOB-*adhT* were streaked from

glycerol stocks onto LB plates containing 0.25 mM DAP and streptomycin 100 mg/L, and grown

overnight at 37°C. Individual colonies from each strain were inoculated in 10 ml M9 relaxing liquid

medium (Table S 4) and incubated overnight at 37°C and 220 rpm. Cells were harvested and washed

3 times with M9 minimal medium at room temperature. The first washing step consisted of spinning

down the cultures at 4700 rpm for 10 minutes, followed by resuspension in 900 µl. The second and

third washing steps were performed at 1100 rpm for 1 minute. After washing was completed,

experimental strains were diluted to a final OD_{600} of 0.01 in M9 selective media containing either 0, 10,

40 or 80 mM of methanol (Table S3). OD_{600} was measured using a portable ImplenTM OD_{600}

DiluPhotometer. 150 µl of each culture was transferred to a transparent 96-well flat-bottomed plate

with 50 µl mineral oil pipetted on top. For each condition and strain, three replicates were used. The

growth of the different *E. coli* HOB strains was monitored for 70 hours in the BioTek synergy Neo2

multi-mode reader.

Data availability

644 The mass spectrometric raw files as well as the MaxQuant output files will be deposited to the

645 ProteomeXchange Consortium via the PRIDE[64] partner repository. gDNA raw reads corresponding to

646 the PTM1, PTM2 strains and the parental Δ*adhP* Δ*rpe* strain will be deposited in the European

647 Nucleotide Archive.

Acknowledgements

649 We acknowledge Sebastian Wenk and his co-authors for supplying and providing advice on the use of

650 the *E. coli* formaldehyde sensor strain. N.J.C. acknowledges support from his NWO Veni fellowship

651 (VI.Veni.192.156). S.D. acknowledges support by The Novo Nordisk Foundation (NNF20CC0035580).

References

1. Di Lorenzo, R. D., Serra, I., Porro, D., & Branduardi, P. (2022). State of the art on the microbial
production of industrially relevant organic acids. In *Catalysts* (Vol. 12, Issue 2). MDPI.
https://doi.org/10.3390/catal12020234

2. Blombach, B., Grünberger, A., Centler, F., Wierckx, N., & Schmid, J. (2021). Exploiting
unconventional prokaryotic hosts for industrial biotechnology. In *Trends in Biotechnology*.
Elsevier Ltd. https://doi.org/10.1016/j.tibtech.2021.08.003

3. Liu, J., Han, X., Tao, F., & Xu, P. (n.d.). Reprogramming a high robust *Geobacillus*
thermoglucosidasius 1 for efficient synthesis of polymer-grade lactic acid under 2 extremely high
temperature (60° C). https://doi.org/10.1101/2023.04.14.536835

4. Zhou, J., Wu, K., & Rao, C. V. (2016). Evolutionary engineering of *Geobacillus thermoglucosidasius*
for improved ethanol production. *Biotechnol. Bioeng*, *113*, 2156–2167.
https://doi.org/10.1002/bit.25983/abstract

5. Raita, M., Ibenegbu, C., Champreda, V., & Leak, D. J. (2016). Production of ethanol by thermophilic
oligosaccharide utilising *Geobacillus thermoglucosidasius* TM242 using palm kernel cake as a
renewable feedstock. *Biomass and Bioenergy*, *95*, 45–54.
https://doi.org/10.1016/j.biombioe.2016.08.015

6. Cotton, C. A., Claassens, N. J., Benito-Vaquerizo, S., & Bar-Even, A. (2020). Renewable methanol
and formate as microbial feedstocks. In *Current Opinion in Biotechnology* (Vol. 62, pp. 168–180).
Elsevier Ltd. https://doi.org/10.1016/j.copbio.2019.10.002

7. Liao, J. C., Mi, L., Pontrelli, S., & Luo, S. (2016). Fuelling the future: microbial engineering for the
production of sustainable biofuels. *Nature Reviews Microbiology*, *14*(5), 288–304.
https://doi.org/10.1038/nrmicro.2016.32

8. Beber, M. E., Gollub, M. G., Mozaffari, D., Shebek, K. M., Flamholz, A. I., Milo, R., & Noor, E.
(2022). eQuilibrator 3.0: a database solution for thermodynamic constant estimation. *Nucleic*
Acids Research, *50*(D1), D603–D609. https://doi.org/10.1093/nar/gkab1106

9. Heux, S., Brautaset, T., Vorholt, J. A., Wendisch, V. F., & Portais, J. C. (2018). Synthetic
Methylotrophy: Past, Present, and Future. In *Methane Biocatalysis: Paving the Way to*
Sustainability (pp. 133–151). Springer International Publishing. https://doi.org/10.1007/978-3-
319-74866-5_9

10. Krüsemann, J. L., Rainaldi, V., Cotton, C. A., Claassens, N. J., & Lindner, S. N. (2023). The
cofactor challenge in synthetic methylotrophy: bioengineering and industrial applications. *Current*
Opinion in Biotechnology, *82*, 102953. https://doi.org/10.1016/j.copbio.2023.102953

11. Whitaker, W. B., Sandoval, N. R., Bennett, R. K., Fast, A. G., & Papoutsakis, E. T. (2015). Synthetic methylotrophy: engineering the production of biofuels and chemicals based on the biology of aerobic methanol utilization. *Current Opinion in Biotechnology*, *33*, 165–175. https://doi.org/10.1016/j.copbio.2015.01.007

12. Komives, C. F., Cheung, L. Y.-Y., Pluschkell, S. B., & Flickinger, M. C. (2005). Growth of *Bacillus methanolicus* in seawater-based media. *Journal of Industrial Microbiology & Biotechnology*, *32*(2), 61–66. https://doi.org/10.1007/s10295-004-0195-9

13. Müller, J. E. N., Heggeset, T. M. B., Wendisch, V. F., Vorholt, J. A., & Brautaset, T. (2015). Methylotrophy in the thermophilic *Bacillus methanolicus*, basic insights and application for commodity production from methanol. In *Applied Microbiology and Biotechnology* (Vol. 99, Issue 2, pp. 535–551). Springer Verlag. https://doi.org/10.1007/s00253-014-6224-3

14. Müller, J. E. N., Meyer, F., Litsanov, B., Kiefer, P., & Vorholt, J. A. (2015). Core pathways operating during methylotrophy *of Bacillus methanolicus* MGA3 and induction of a bacillithiol-dependent detoxification pathway upon formaldehyde stress. *Molecular Microbiology*, *98*(6), 1089–1100. https://doi.org/10.1111/mmi.13200

15. Brautaset, T., Jakobsen, Ø. M., Josefsen, K. D., Flickinger, M. C., & Ellingsen, T. E. (2007). *Bacillus methanolicus*: A candidate for industrial production of amino acids from methanol at 50°C. In *Applied Microbiology and Biotechnology* (Vol. 74, Issue 1, pp. 22–34). https://doi.org/10.1007/s00253-006-0757-z

16. Paredes-Barrada, M., Kopsiaftis, P., Claassens, N. J., & van Kranenburg, R. (2024). *Parageobacillus thermoglucosidasius* as an emerging thermophilic cell factory. *Metabolic Engineering*, *83*, 39–51. https://doi.org/10.1016/j.ymben.2024.03.001

17. Millgaard, M., Bidart, G. N., Pogrebnyakov, I., Nielsen, A. T., & Welner, D. H. (2023). An improved integrative GFP-based vector for genetic engineering of *Parageobacillus thermoglucosidasius* facilitates the identification of a key sporulation regulator. *AMB Express*, *13*(1), 44. https://doi.org/10.1186/s13568-023-01544-9

18. Yang, Z., Li, Z., Li, B., Bu, R., Tan, G.-Y., Wang, Z., Yan, H., Xin, Z., Zhang, G., Li, M., Xiang, H., Zhang, L., & Wang, W. (2023). A thermostable type I-B CRISPR-Cas system for orthogonal and multiplexed genetic engineering. *Nature Communications*, *14*(1), 6193. https://doi.org/10.1038/s41467-023-41973-5

19. Simon, M., & Msci, H. L. (n.d.). The Expansion of Available Bioparts and the Development of CRISPR/Cas9 for Genetic Engineering of Parageobacillus thermoglucosidasius.

20. Chen, F. Y. H., Jung, H. W., Tsuei, C. Y., & Liao, J. C. (2020). Converting *Escherichia coli* to a synthetic methylotroph growing solely on methanol. *Cell*, *182*(4), 933-946.e14. https://doi.org/10.1016/j.cell.2020.07.010

21. Kim, S., Lindner, S. N., Aslan, S., Yishai, O., Wenk, S., Schann, K., & Bar-Even, A. (2020). Growth of *E. coli* on formate and methanol via the reductive glycine pathway. *Nature Chemical Biology*, *16*(5), 538–545. https://doi.org/10.1038/s41589-020-0473-5

22. Reiter, M. A., Bradley, T., Büchel, L. A., Keller, P., Hegedis, E., Gassler, T., & Vorholt, J. A. (2024). A synthetic methylotrophic *Escherichia coli* as a chassis for bioproduction from methanol. *Nature Catalysis*, *7*(5), 560–573. https://doi.org/10.1038/s41929-024-01137-0

23. Nieh, L.-Y., Chen, F. Y.-H., Jung, H.-W., Su, K.-Y., Tsuei, C.-Y., Lin, C.-T., Lee, Y.-Q., & Liao, J. C.
(2024). Evolutionary engineering of methylotrophic *E. coli* enables fast growth on methanol.
Nature Communications, *15*(1), 8840. https://doi.org/10.1038/s41467-024-53206-4

24. Keller, P., Reiter, M. A., Kiefer, P., Gassler, T., Hemmerle, L., Christen, P., Noor, E., & Vorholt, J.
A. (2022). Generation of an *Escherichia coli* strain growing on methanol via the ribulose
monophosphate cycle. *Nature Communications*, *13*(1), 5243. https://doi.org/10.1038/s41467-
022-32744-9

25. Whitaker, W. B., Jones, J. A., Bennett, K., Gonzalez, J., Vernacchio, V. R., Collins, S. M., Palmer,
M. A., Schmidt, S., Antoniewicz, M. R., Koffas, M. A., & Papoutsakis, E. T. (n.d.). Engineering the
Biological Conversion of Methanol to Specialty Chemicals in Escherichia coli.

26. Zhan, C., Li, X., Lan, G., Baidoo, E. E. K., Yang, Y., Liu, Y., Sun, Y., Wang, S., Wang, Y., Wang,
G., Nielsen, J., Keasling, J. D., Chen, Y., & Bai, Z. (2023). Reprogramming methanol utilization
pathways to convert *Saccharomyces cerevisiae* to a synthetic methylotroph. *Nature Catalysis*,
6(5), 435–450. https://doi.org/10.1038/s41929-023-00957-w

27. Gao, B., Zhao, N., Deng, J., Gu, Y., Jia, S., Hou, Y., Lv, X., & Liu, L. (2022). Constructing a
methanol-dependent *Bacillus subtilis* by engineering the methanol metabolism. *Journal of
Biotechnology*, *343*, 128–137. https://doi.org/10.1016/j.jbiotec.2021.12.005

28. Bruinsma, L., Wenk, S., Claassens, N. J., & Martins dos Santos, V. A. P. (2023). Paving the way
for synthetic C1 - Metabolism in *Pseudomonas putida* through the reductive glycine pathway.
Metabolic Engineering, *76*, 215–224. https://doi.org/10.1016/j.ymben.2023.02.004

29. Meyer, F., Keller, P., Hartl, J., Gröninger, O. G., Kiefer, P., & Vorholt, J. A. (2018). Methanol-
essential growth of *Escherichia coli*. *Nature Communications*, *9*(1).
https://doi.org/10.1038/s41467-018-03937-y

30. Keller, P., Noor, E., Meyer, F., Reiter, M. A., Anastassov, S., Kiefer, P., & Vorholt, J. A. (2020).
Methanol-dependent *Escherichia coli* strains with a complete ribulose monophosphate cycle.
Nature Communications, *11*(1). https://doi.org/10.1038/s41467-020-19235-5

31. Müller, J. E. N., Meyer, F., Litsanov, B., Kiefer, P., Potthoff, E., Heux, S., Quax, W. J., Wendisch,
V. F., Brautaset, T., Portais, J. C., & Vorholt, J. A. (2015). Engineering *Escherichia coli* for
methanol conversion. *Metabolic Engineering*, *28*, 190–201.
https://doi.org/10.1016/j.ymben.2014.12.008

32. Wu, T.-Y., Chen, C.-T., Liu, J. T.-J., Bogorad, I. W., Damoiseaux, R., & Liao, J. C. (2016).
Characterization and evolution of an activator-independent methanol dehydrogenase from
Cupriavidus necator N-1. *Applied Microbiology and Biotechnology*, *100*(11), 4969–4983.
https://doi.org/10.1007/s00253-016-7320-3

33. Lee, J. Y., Park, S. H., Oh, S. H., Lee, J. J., Kwon, K. K., Kim, S. J., Choi, M., Rha, E., Lee, H.,
Lee, D. H., Sung, B. H., Yeom, S. J., & Lee, S. G. (2020). Discovery and biochemical
characterization of a methanol dehydrogenase from *Lysinibacillus xylanilyticus*. *Frontiers in
Bioengineering and Biotechnology*, *8*. https://doi.org/10.3389/fbioe.2020.00067

34. Klein, V. J., Irla, M., Gil López, M., Brautaset, T., & Fernandes Brito, L. (2022). Unravelling
formaldehyde metabolism in bacteria: Road towards synthetic methylotrophy. *Microorganisms*,
10(2), 220. https://doi.org/10.3390/microorganisms10020220

35. Mougiakos, I., Mohanraju, P., Bosma, E. F., Vrouwe, V., Finger Bou, M., Naduthodi, M. I. S.,
Gussak, A., Brinkman, R. B. L., Van Kranenburg, R., & Van Der Oost, J. (2017). Characterizing a
thermostable Cas9 for bacterial genome editing and silencing. *Nature Communications*, *8*(1).
https://doi.org/10.1038/s41467-017-01591-4

36. Yurimoto, H., Hirai, R., Matsuno, N., Yasueda, H., Kato, N., & Sakai, Y. (2005). HxlR, a member of
the DUF24 protein family, is a DNA-binding protein that acts as a positive regulator of the
formaldehyde-inducible *hxlAB* operon in *Bacillus subtilis*. *Molecular Microbiology*, *57*(2), 511–
519. https://doi.org/10.1111/j.1365-2958.2005.04702.x

37. Yasueda, H., Kawahara, Y., & Sugimoto, S. (1999). *Bacillus subtilis yckG* and *yckF* Encode Two
Key Enzymes of the Ribulose Monophosphate Pathway Used by Methylotrophs, and *yckH* Is
Required for Their Expression. *Journal of Bacteriology*, *181*(23), 7154–7160.
https://doi.org/10.1128/JB.181.23.7154-7160.1999

38. Krog, A., Heggeset, T. M. B., Müller, J. E. N., Kupper, C. E., Schneider, O., Vorholt, J. A.,
Ellingsen, T. E., & Brautaset, T. (2013). Methylotrophic *Bacillus methanolicus* Encodes Two
Chromosomal and One Plasmid Born NAD+ Dependent Methanol Dehydrogenase Paralogs with
Different Catalytic and Biochemical Properties. *PLoS ONE*, *8*(3).
https://doi.org/10.1371/journal.pone.0059188

39. Kovina, M., Viryasov, M., Baratova, L., & Kochetov, G. (1996). Localization of reactive tyrosine
residues of baker's yeast transketolase. *FEBS Letters*, *392*(3), 293–294.
https://doi.org/10.1016/0014-5793(96)00836-8

40. Kovina, M. V, & Kochetov, G. A. (1998). Cooperativity and flexibility of active sites in homodimeric
transketolase. *FEBS Letters*, *440*(1–2), 81–84. https://doi.org/10.1016/S0014-5793(98)01423-9

41. GA, K., & IA, S. (2005). Binding of the coenzyme and formation of the transketolase active center.
IUBMB Life (International Union of Biochemistry and Molecular Biology: Life), *57*(7), 491–497.
https://doi.org/10.1080/15216540500167203

42. Zhu, R., Zhang, G., Jing, M., Han, Y., Li, J., Zhao, J., Li, Y., & Chen, P. R. (2021). Genetically
encoded formaldehyde sensors inspired by a protein intra-helical crosslinking reaction. *Nature*
Communications, *12*(1), 581. https://doi.org/10.1038/s41467-020-20754-4

43. Witthoff, S., Mühlroth, A., Marienhagen, J., & Bott, M. (2013). C$_1$ metabolism in *Corynebacterium*
glutamicum: an endogenous pathway for oxidation of methanol to carbon dioxide. *Applied and*
Environmental Microbiology, *79*(22), 6974–6983. https://doi.org/10.1128/AEM.02705-13

44. Lessmeier, L., Hoefener, M., & Wendisch, V. F. (2013). Formaldehyde degradation in
Corynebacterium glutamicum involves acetaldehyde dehydrogenase and mycothiol-dependent
formaldehyde dehydrogenase. *Microbiology*, *159*(Pt_12), 2651–2662.
https://doi.org/10.1099/mic.0.072413-0

45. Schann, K., Bakker, J., Boinot, M., Kuschel, P., He, H., Nattermann, M., Paczia, N., Erb, T.,
Bar-Even, A., & Wenk, S. (2024). Design, construction and optimization of formaldehyde growth
biosensors with broad application in biotechnology. *Microbial Biotechnology*, *17*(7).
https://doi.org/10.1111/1751-7915.14527

46. Nash, T. (1953). The colorimetric estimation of formaldehyde by means of the Hantzsch reaction.
Biochemical Journal, *55*(3), 416–421. https://doi.org/10.1042/bj0550416

47. Reeve, B., Martinez-Klimova, E., De Jonghe, J., Leak, D. J., & Ellis, T. (2016). The *Geobacillus*
Plasmid Set: A Modular Toolkit for Thermophile Engineering. *ACS Synthetic Biology*, *5*(12),
1342–1347. https://doi.org/10.1021/acssynbio.5b00298

48. Jakobsen, Ø. M., Benichou, A., Flickinger, M. C., Valla, S., Ellingsen, T. E., & Brautaset, T. (2006).
Upregulated transcription of plasmid and chromosomal ribulose monophosphate pathway genes
is critical for methanol assimilation rate and methanol tolerance in the methylotrophic bacterium
Bacillus methanolicus. *Journal of Bacteriology*, *188*(8), 3063–3072.
https://doi.org/10.1128/JB.188.8.3063-3072.2006

49. Marinus Petrus Machielsen, Mariska Van Hartskamp, & Richard Van Kranenburg. (2015).
Genetically modified (r)-lactic acid producing thermophilic bacteria (Patent WO2016012296A1).

50. Marinus Petrus Machielsen, Mariska Van Hartskamp & Richard Van Kranenburg. (2015).
Genetically modified (r)-lactic acid producing thermophilic bacteria. (Patent WO2016012296A1).

51. Richard Van Kranenburg, Mariska Van Hartskamp, Eelco Anthonius Johannes, Heintz Esther,
Johanna Geertruda Van Mullekom & Jurgen Snelders. (2013). *Genetic modification of*
homolactic thermophilic bacilli. (Patent WO2007085443A3)

52. Hon Wai Wan. (2013). Transformation of the thermophilic bacterium, *Geobacillus debilis,* by
conjugation with the mesophilic bacterium, *Escherichia coli.*

53. Long, C. P., & Antoniewicz, M. R. (2019). High-resolution 13C metabolic flux analysis. *Nature*
Protocols, *14*(10), 2856–2877. https://doi.org/10.1038/s41596-019-0204-0

54. Kutuzova, S., Colaianni, P., Röst, H., Sachsenberg, T., Alka, O., Kohlbacher, O., Burla, B., Torta,
F., Schrübbers, L., Kristensen, M., Nielsen, L., Herrgård, M. J., & McCloskey, D. (2020).
SmartPeak automates targeted and quantitative metabolomics data processing. *Analytical*
Chemistry, *92*(24), 15968–15974. https://doi.org/10.1021/acs.analchem.0c03421

55. Wahl, S. A., Dauner, M., & Wiechert, W. (2004). New tools for mass isotopomer data evaluation in
^{13}C flux analysis: Mass isotope correction, data consistency checking, and precursor
relationships. *Biotechnology and Bioengineering*, *85*(3), 259–268.
https://doi.org/10.1002/bit.10909

56. Abele, M., Doll, E., Bayer, F. P., Meng, C., Lomp, N., Neuhaus, K., Scherer, S., Kuster, B., &
Ludwig, C. (2023). Unified workflow for the rapid and in-depth characterization of bacterial
proteomes. *Molecular & Cellular Proteomics*, *22*(8), 100612.
https://doi.org/10.1016/j.mcpro.2023.100612

57. Doellinger, J., Schneider, A., Hoeller, M., & Lasch, P. (2020). sample preparation by easy
extraction and digestion (SPEED) - A universal, rapid, and detergent-free protocol for proteomics
based on acid extraction. *Molecular & Cellular Proteomics*, *19*(1), 209–222.
https://doi.org/10.1074/mcp.TIR119.001616

58. Rappsilber, J., Mann, M., & Ishihama, Y. (2007). Protocol for micro-purification, enrichment, pre-
fractionation and storage of peptides for proteomics using StageTips. *Nature Protocols*, *2*(8),
1896–1906. https://doi.org/10.1038/nprot.2007.261

59. Tyanova, S., Temu, T., & Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. *Nature Protocols*, *11*(12), 2301–2319.
https://doi.org/10.1038/nprot.2016.136

60. Cox, J., Neuhauser, N., Michalski, A., Scheltema, R. A., Olsen, J. V., & Mann, M. (2011).
Andromeda: A peptide search engine integrated into the MaxQuant environment. *Journal of*
Proteome Research, *10*(4), 1794–1805. https://doi.org/10.1021/pr101065j

61. Cox, J., Hein, M. Y., Luber, C. A., Paron, I., Nagaraj, N., & Mann, M. (2014). Accurate proteome-
wide label-free quantification by delayed normalization and maximal peptide ratio extraction,
termed MaxLFQ. *Molecular & Cellular Proteomics*, *13*(9), 2513–2526.
https://doi.org/10.1074/mcp.M113.031591

62. Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., & Selbach,
M. (2011). Global quantification of mammalian gene expression control. *Nature*, *473*(7347), 337–
342. https://doi.org/10.1038/nature10098

63. Chen Meng. (2022). Fast analyzing, interpreting and sharing quantitative omics data using
omicsViewer (1.1.3). *Zenodo*.

64. Perez-Riverol, Y., Bai, J., Bandla, C., García-Seisdedos, D., Hewapathirana, S., Kamatchinathan,
S., Kundu, D. J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., Walzer, M., Wang, S.,
Brazma, A., & Vizcaíno, J. A. (2022). The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Research*, *50*(D1), D543–D552.
https://doi.org/10.1093/nar/gkab1038

868