TFAP2C is a key regulator of intrauterine trophoblast cell invasion

and deep hemochorial placentation

Esteban M. Dominguez[1,*], Ayelen Moreno-Irusta[1], Regan L. Scott[1], Khursheed Iqbal[1,†],

and Michael J. Soares[1,2,3,*]

[1]Institute for Reproductive and Developmental Sciences, Department of Pathology &

Laboratory Medicine, University of Kansas Medical Center, Kansas City, Kansas 66160,

[2]Department of Obstetrics and Gynecology, University of Kansas Medical Center,

Kansas City, Kansas 66160, [3]Center for Perinatal Research, Children's Mercy Research

Institute, Children's Mercy, Kansas City, MO 64108

[†]Present address: Department of Animal and Food Sciences, Oklahoma State

University, Stillwater, Oklahoma

*Correspondence: edominguez@kumc.edu or msoares@kumc.edu

Conflict of interests

The authors have declared that no conflict of interest exists.

Abstract

The transcription factor AP-2 gamma (**TFAP2C**) has been identified as a key regulator of the trophoblast cell lineage and hemochorial placentation. The rat possesses deep placentation characterized by extensive intrauterine trophoblast cell invasion, which resembles human placentation. Initially, we examined *Tfap2c* expression in the rat placentation site and demonstrated its presence in multiple trophoblast cell lineages, including invasive trophoblast cells situated within the uterine-placental interface. Global genome-editing was used to explore the biology of *Tfap2c* in rat placentation. Homozygous global disruption of *Tfap2c* resulted in prenatal lethality between gestation days 8.5 and 9.5. Although, global disruption of one *Tfap2c* allele was compatible with survival, this heterozygous condition was surprisingly associated with diminished invasive trophoblast cell infiltration into the uterus. Since *Tfap2c* is expressed in multiple trophoblast cell lineages, we next explored the role of TFAP2C in the invasive trophoblast cell lineage using Cre-lox conditional mutagenesis. Invasive trophoblast cell-specific disruption of *Tfap2c* resulted in inhibition of intrauterine trophoblast cell invasion and intrauterine and postnatal growth restriction. In contrast to the global disruption of *Tfap2c*, the invasive trophoblast cell lineage was not impaired following conditional monoallelic disruption of *Tfap2c*. In summary, TFAP2C contributes to the progression of distinct stages of placental development. TFAP2C is a driver of early events in trophoblast cell development and reappears later in gestation as an essential regulator of the invasive trophoblast cell lineage. A subset of TFAP2C actions on trophoblast cells are dependent on gene dosage.

Introduction

The placenta is a transient organ that ensures viviparity and optimal fetal development (1–3). The human and rat possess a hemochorial placenta (4, 5), which is composed of specialized cellular constituents termed trophoblast cells (6). Among the trophoblast cell lineages contributing to the placentation site are invasive trophoblast cells, which in the human are referred to as extravillous trophoblast cells (7). Human and rat placentation sites are characterized by deep intrauterine trophoblast cell invasion (8–10). Invasive trophoblast cells facilitate the erosion of maternal uterine spiral arteries permitting a direct flow of maternal nutrients to trophoblast cells (8, 11–15). Uterine vasculature remodeling is a critical process ensuring a successful pregnancy outcome (13, 16, 17). There is a paucity of knowledge regarding regulatory mechanisms controlling development of the invasive trophoblast cell lineage and trophoblast cell-guided uterine spiral remodeling.

Members of the transcription factor AP-2 (**TFAP2**) family are key regulators of embryonic and extraembryonic development (18, 19). TFAP2C distinguishes itself as a regulator of the trophoblast cell lineage and hemochorial placentation (20–22). In the mouse, global disruption of the *Tfap2c* locus results in abnormalities within the trophoblast cell lineage and prenatal lethality prior to placentation (23, 24). Although, TFAP2C is an established regulator of the earliest stages of trophoblast cell lineage development (25–28), placental expression profiles and target gene occupancy also implicate TFAP2C in the regulation of multiple processes impacting mammalian placentation and placental function (29–40). Included in this broad spectrum of TFAP2C

actions on placentation is its potential involvement in invasive trophoblast cell biology

(38–41).

In this report, we present the results of in vivo experimentation directed toward

elucidating the physiological role of TFAP2C in the developing hemochorial placenta

with a focus on trophoblast cell-guided events within the uterine-placental interface. We

show that TFAP2C possesses essential roles during distinct stages of placental

development. TFAP2C is an essential regulator of early placental morphogenetic events

and reappears in a critical capacity for invasive trophoblast cell lineage development.

Finally, elements of TFAP2C actions on placentation are dose dependent.

Results

TFAP2C expression within the placentation site

The rat hemochorial placenta is comprised of multiple lineages of trophoblast cells

arranged into distinct compartments, including the labyrinth zone, junctional zone, and

uterine-placental interface (42, 43) (Figure 1). The labyrinth zone is the site of maternal-

fetal nutrient/waste transfer (44), whereas the junctional zone is composed of endocrine

cells targeting the maternal environment and the site where invasive trophoblast cells arise

(42, 43, 45). Cell composition of the uterine-placental interface is dynamic, including an

increase in the infiltration of invasive trophoblast cells as gestation proceeds (9, 38).

Intrauterine invasive trophoblast cells invade into the stroma located between uterine

blood vessels (interstitial) and within uterine blood vessels (endovascular). *Tfap2c*

transcripts were detected within the uterine-placental interface and increased during

pregnancy (Supplemental Figure 1A). In situ hybridization was used to determine the

distribution of *Tfap2c* transcripts within the hemochorial placenta. At gestation day (**gd**)

9.5, *Tfap2c* was prominently expressed in the ectoplacental cone (Figure 1A). *Tfap2c*

transcripts were also localized to the anti-mesometrial uterine deciduum (Figure 1A). As

gestation advanced (gd 11.5), *Tfap2c* expression was observed within trophoblast cells of

the developing placenta (Figure 1B). On gd 18.5, *Tfap2c* was abundantly expressed within

the junctional zone and in invasive trophoblast cells of the uterine-placental interface

(Figure 1C-E). The presence of *Tfap2c* transcripts in the trophoblast cell lineage was

supported by its colocalization with keratin 8 (**Krt8**), a pan-trophoblast cell marker (Figure

1A-C). Expression of *Tfap2c* transcripts in invasive trophoblast cells was further

demonstrated by co-localization with prolactin family 7, subfamily b, member 1 (*Prl7b1*)

transcripts (Figure 1D and E), an invasive trophoblast cell-specific transcript, and is

consistent with single cell RNA-sequencing of the of the uterine-placental interface

(Supplemental Figure 1B) (38). TFAP2C protein was distributed in invasive trophoblast

cells of the uterine-placental interface and throughout the junctional zone, replicating the

distribution of *Tfap2c* transcripts at gd 18.5 (Supplemental Figure 2). Within the invasive

trophoblast cell lineage, TFAP2C transcript and protein were localized to both

endovascular and interstitial trophoblast cells (Figure 1E**,** Supplemental Figure 3). *Tfap2c*

transcripts were localized to most trophoblast cell populations within the junctional zone,

including a smaller subset of invasive trophoblast progenitor cell populations (*Prl7b1*

positive; **Figure 1F**).

Global deletion of *Tfap2c* leads to prenatal lethality

Next, we investigated the in vivo role of *Tfap2c* in rat placenta development using

CRISPR/Cas9 genome editing. DNA-binding and dimerization domains encoded by Exon

4 of the *Tfap2c* gene were targeted (Supplemental Figure 4A). A germ line mutant rat was

generated possessing a 308-bp deletion within Exon 4 (Supplemental Figure 4B). The

deletion resulted in a frameshift and a premature stop codon (Supplemental Figure 4C).

Polymerase chain reaction (**PCR**) products corresponding to wild type and mutant alleles

were identified by genotyping (828 bp versus 520 bp, respectively, Supplemental Figure

4D). Null *Tfap2c* conceptuses (-/-) were obtained by mating heterozygous (+/-) *Tfap2*c

females and males. Breeding results collected throughout pregnancy and postnatally,

including Mendelian ratios, are presented in Supplemental Table 1. Null *Tfap2c*

conceptuses were viable until gd 8.5, which temporally coincides with the onset of

placental morphogenesis. This observation agrees with the timing of embryonic demise

observed following global *Tfap2c* mouse gene disruption (23, 24). Interestingly, we

observed an unexpected reduction in the number of surviving heterozygous pups

generated from *Tfap2c*$^{+/-}$ x *Tfap2c*$^{+/-}$, which prompted an investigation of placental

development in conceptuses possessing a single *Tfap2c* mutant allele.

Decreased *Tfap2*c gene dosage leads to a failure in intrauterine trophoblast cell

invasion

To explore possible *Tfap2c* gene dosage effects, *Tfap2c*$^{+/+}$ females and *Tfap2c*$^{+/-}$ males

were mated, and pregnancies assessed at gd 18.5. *Tfap2c*$^{+/+}$ and *Tfap2c*$^{+/-}$ placentation

sites exhibited a similar organizational structure, including a labyrinth zone, junctional

zone, and a uterine-placental interface (Figure 2A). We next assessed infiltration of

invasive trophoblast cells into the uterine-placental interfaces of *Tfap2c*$^{+/+}$ versus *Tfap2c*$^{+/-}$

placentation sites by monitoring the distribution of cytokeratin protein expression (9, 38)

(Figure 2B) and *Prl7b1* mRNA (46) (Figure 2C), both representing indices of invasive

trophoblast cells. The uterine-placental interface of *Tfap2c*$^{+/-}$ placentation sites were

depleted of invasive trophoblast cells (Figure 2B and C). Both interstitial and endovascular

invasive trophoblast cells were diminished in the uterine-placental interface of *Tfap2c*$^{+/-}$

placentation sites. This result was supported by quantification of the depth of invasion of

cytokeratin positive cells into the uterine parenchyma (Figure 2D). The depletion of

invasive trophoblast cells in the uterine-placental interface of $Tfap2c^{+/-}$ placentation sites

was further supported by diminished expression of several invasive trophoblast cell

associated transcripts (*Krt7, Krt8, Krt18, Prl7b1, Prl5a1, Cdkn1c, Cited2, Plac1, Peg3,*

Tfpi; (38) (Figure 2E). Similar findings were recorded when placentation sites were

interrogated from pregnancies generated from $Tfap2c^{+/-}$ females mated to $Tfap2c^{+/+}$ males

(Supplemental Figure 5 A-D).

However, breeding combinations did differentially affect postnatal heterozygote

offspring viability and heterozygote fetal and placental weights. Postnatal heterozygote

viability was prominently decreased in female $Tfap2c^{+/-}$ x male $Tfap2c^{+/-}$ but not in female

$Tfap2c^{+/+}$ x male $Tfap2c^{+/-}$ or female $Tfap2c^{+/-}$ x male $Tfap2c^{+/+}$ breeding combinations

(Supplemental Table 1-3). Genotypic differences in fetal, placenta, junctional zone, and

labyrinth zone weights at gd 18.5 were observed among breeding combinations. Mating

$Tfap2c^{+/+}$ females with $Tfap2c^{+/-}$ males was associated with significant decreases in

heterozygote placenta, junctional zone, and labyrinth zone weights (Figure 2F), whereas

only heterozygote fetal weights were significantly decreased by mating $Tfap2c^{+/-}$ females

with $Tfap2c^{+/+}$ males (Supplemental Figure 5 E). Some of these differences might be

associated with *Tfap2c* gene dosage effects on the maternal environment.

Invasive trophoblast cell conditional *Tfap2c* mutagenesis

The impact of TFAP2C dosage on invasive trophoblast cell development prompted a

direct analysis of the importance of TFAP2C on the invasive trophoblast cell lineage. We

generated an invasive trophoblast cell-specific *Tfap2c* gene disruption using Cre/loxP

system. CRISPR/Cas9 genome editing was used to insert loxP sites flanking Exon 4 of

the *Tfap2c* gene (Figure 3A). *Tfap2c* floxed (**Tfap2c^f/f**) rats were mated with *Prl7b1-Cre*

rats (47). Cre recombinase is specifically targeted to invasive trophoblast cells in *Prl7b1-*

Cre rats (47). TFAP2C protein was present in lysates of the *Tfap2c^f/f* uterine-placental

interface but not in lysates of the *Tfap2c^d/d* uterine-placental interface (Figure 3B).

Immunostaining supported depletion of TFAP2C from the gd 18.5 uterine-placental

interface but not from the JZ of *Tfap2c^d/d* placentation sites (Figure 3C). Our ability to

modulate TFAP2C in the invasive trophoblast cell lineage led us to evaluate a potential

role for TFAP2C in regulating the invasive trophoblast cell lineage.

TFAP2C-dependent intrauterine trophoblast cell invasion

Intrauterine trophoblast cell invasion was assessed in *Tfap2c^f/f* and *Tfap2c^d/d*

placentation sites. At gd 18.5 invasive trophoblast cells were present throughout the

Tfap2c^f/f uterine-placental interface but were not detected in the *Tfap2^d/d* uterine-placental

interface (Figure 4 A-C). Interstitial and endovascular invasive trophoblast cells were

absent from the uterine-placental interface of *Tfap2c^d/d* placentation sites. The depletion

of invasive trophoblast cells in the uterine-placental interface of *Tfap2c^d/d* placentation

sites was further supported by diminished expression of several invasive trophoblast cell-

associated transcripts (*Krt7, Krt8, Krt18, Prl7b1, Prl5a1, Cdkn1c, Cited2, Plac1, Peg3,*

Tfpi; Figure 4D). In contrast to the global disruption of *Tfap2c*, migration of invasive

trophoblast cells into the uterus was not impaired following conditional monoallelic

disruption of *Tfap2c* (Supplemental Figure 6). Single cell RNA-sequencing and single

nucleus assay for transposase-accessible chromatin-sequencing datasets (38, 41) reveal

a network of genes potentially regulated by TFAP2C in rat invasive trophoblast cells that,

include candidate contributors to remodeling extracellular matrix, cytoskeleton

restructuring, regulation of cell movement, and transcriptional regulators that target genes

encoding other proteins affecting the invasive trophoblast phenotype (Figure 4E;

Supplemental Dataset 1). Thus, TFAP2C has a fundamental role in regulating the invasive

trophoblast cell lineage, especially movement of invasive trophoblast cells into the uterus,

and a list of downstream effectors are available to investigate mechanisms underlying

TFAP2C actions.

Invasive trophoblast cells and natural killer (NK) cells exhibit a reciprocal

distribution within the uterine-placental interface

At midgestation, NK cells are an abundant constituent of the uterine-placental interface

(48). As gestation progresses, NK cells become depleted as trophoblast cells invade the

uterine-placental interface (48). Interestingly, we observed a retention of NK cells in

Tfap2c$^{d/d}$ uterine-placental interfaces depleted of invasive trophoblast cells (Figure 5A).

This observation was further supported by increased expression of NK cell-associated

transcripts (*Prf1, Klrb1c, Klrb1a, Klrb1, Ncr1*; Figure 5B) within *Tfap2c$^{d/d}$* uterine-placental

interfaces. These findings reinforce the observation that fundamental changes occur in

the *Tfap2c$^{d/d}$* uterine-placental interface and imply that invasive trophoblast cells affect

cellular dynamics within the uterine-placental interface.

Invasive trophoblast cells impact prenatal and postnatal growth

We next investigated whether the presence of invasive trophoblast cells within the

uterine-placental interface affected prenatal or postnatal growth. Depletion of invasive

trophoblast cells as observed in *Tfap2c$^{d/d}$* placentation sites was associated with

intrauterine placental growth restriction, including both junctional zone and labyrinth zone

compartments, and fetal growth restriction at gd 18.5 (Figure 6A). The adverse effects of

invasive trophoblast cell depletion on fetal weight extended to postnatal offspring growth.

Depletion of invasive trophoblast cells was associated with decreased postnatal body

weight (Figure 6B). The results are consistent with invasive trophoblast cells supporting

the nutrient demands required for optimal intrauterine growth. Deficits acquired prenatally

were sustained during the first three weeks of postnatal life.

Discussion

Placentation is a critical adaptation ensuring the success of viviparity (5, 49). Gene

manipulated animal models and trophoblast stem cells have provided insights into gene

regulatory networks controlling trophoblast cell lineage development and placentation

(21). TFAP2C figures prominently in these networks and represents a conserved

regulator of placentation (20–22). In the research reported here, we utilized genome-

edited rat models to elucidate roles for TFAP2C in placentation. The rat, like the human,

is a species exhibiting deep intrauterine trophoblast cell invasion (8–10). TFAP2C is

expressed in multiple trophoblast cell lineages of the developing rat placenta, including

invasive trophoblast cells residing in the uterine-placental interface. Consistent with

earlier reports in the mouse (23, 24), global disruption of *Tfap2c* in the rat led to prenatal

lethality (gd 8.5 to 9.5), prior to the initiation of placenta morphogenesis. Interestingly,

decreased gene dosage of *Tfap2c* (heterozygous) was compatible with pregnancy but

affected placentation, including diminished infiltration of invasive trophoblast cells into

the uterine-placental interface. A role for TFAP2C in the regulation of intrauterine

trophoblast cell invasion was further supported using cell lineage specific mutagenesis.

Tfap2c was conditionally disrupted in the invasive trophoblast cell lineage, which led to a

uterine-placental interface devoid of invasive trophoblast cells. Impairment of intrauterine

trophoblast cell invasion had consequences, including intrauterine and postnatal growth

restriction.

Involvement of TFAP2C in early events dictating trophoblast cell lineage development

were reinforced from phenotyping conceptuses possessing a global disruption of the

Tfap2c locus. TFAP2C is critical to activation of the trophectoderm transcriptome (36,

50) and development of trophectoderm (25, 26, 51, 52). Trophectoderm represents the

earliest stage of trophoblast cell lineage development (53, 54). The importance of

TFAP2C in early trophoblast cell lineage development is further exemplified by its

utilization in the direct reprogramming of fibroblasts to the trophoblast cell lineage (27,

28).

TFAP2C actions in the rat placentation site were affected by gene dosage. Monoallelic

disruption of *Tfap2c* resulted in the development of a rat uterine-placental interface

possessing a diminished number of invasive trophoblast cells. Observing a phenotype in

Tfap2c global heterozygotes was intriguing but not entirely new for *Tfap2c*.

Haploinsufficiency of *Tfap2c* is linked to an increased incidence of peripartum lethality

(55). Surviving *Tfap2c* heterozygotes exhibit postnatal growth restriction (24),

development of germ cell tumors (56), and hippocampal dysfunction (57). Placentation is

also affected by *Tfap2c* haploinsufficiency (55). The labyrinth zone compartment of the

mouse placenta is disorganized in *Tfp2c* heterozygotes (55). Interestingly, the invasive

trophoblast cell phenotype observed in global heterozygotes reported here was not a

feature of conceptuses with a monoallelic disruption of *Tfap2c* restricted to the invasive

trophoblast cell lineage. An explanation for this difference is likely connected to the wide-

spread involvement of TFAP2C in multiple stages of trophoblast cell differentiation (58).

This is supported by TFAP2C expression extending to many more trophoblast cells in

the junctional zone than express *Prl7b1*, the gene driving Cre recombinase expression in

our conditional mutagenesis rat model. Global *Tfap2c* haploinsufficiency at a key stage

of trophoblast cell lineage development distinct from *Prl7b1*-expressing trophoblast cells

must be relevant and responsive to *Tfap2c* gene dosage. Thus, *Tfap2c* gene dosage is

utilized in a differentiation stage specific manner to guide trophoblast cell development.

TFAP2C is utilized at different stages of trophoblast cell development. TFAP2C is

critical to the derivation of the trophoblast cell lineage and then repurposed for later

stages of placentation (58), including the regulation of invasive trophoblast cell

development. TFAP2C does not act alone. Context is critical for the actions of TFAP2C.

The presence of other transcription factor partners and stoichiometry influence TFAP2C

function during trophoblast cell development (21, 59). A short list of potential TFAP2C

partners in invasive trophoblast cell development is available (38, 41).

Lastly, we have taken transcript and open chromatin profiles (38, 41) and created a

path for elucidating gene involvement in the regulation of invasive trophoblast cell

lineage development and its impact on the uterine-placental interface. We provide

compelling evidence that TFAP2C is critical in establishing the invasive trophoblast cell

lineage. Furthermore, we effectively demonstrate that the presence of invasive

trophoblast cells in the uterus has consequences. Deficits in invasive trophoblast cell

guided uterine transformation resulted in intrauterine and postnatal growth restriction.

Most importantly, tools are in place to establish a gene regulatory network controlling

development and function of the invasive trophoblast cell lineage.

In summary, TFAP2C contributes to the progression of distinct stages of rat placental

development. TFAP2C is a driver of early events in trophoblast cell development and

reappears later in gestation as an essential regulator of the invasive trophoblast cell

lineage. A subset of TFAP2C actions on trophoblast cells are dependent on gene

dosage.

Methods

Animals and tissue collection

Holtzman Sprague–Dawley rats were acquired from Envigo. Animals were maintained

in an environmentally controlled facility (14 h light and 10 h dark cycle) with food and

water available ad libitum. Timed pregnancies were established using virgin female rats

(8 to 10 weeks of age) mated with adult male rats (>3 months of age). Mating was

confirmed by the presence of sperm in the vaginal lavage and was considered gd 0.5.

Pseudopregnant female rats were generated by mating with vasectomized male rats.

Detection of a seminal plug was considered day 0.5 of pseudopregnancy. Pregnant rats

were euthanized on specific days of gestation. Some placentation sites were frozen

intact in dry-ice cooled heptane and stored at -80 °C for subsequent histological

analyses, whereas others were dissected into uterine-placental interface, junctional

zone, and labyrinth zone compartments as previously described (38, 60). Dissected

placentation site compartments and fetuses were weighed, frozen in liquid nitrogen, and

stored at -80°C for subsequent biochemical analyses. All protocols using rats were

approved by the University of Kansas Medical Center Animal Care and Use Committee.

Generation of global and conditional *Tfap2c* mutant rat models

A global mutation at the rat *Tfap2c* locus was generated using CRISPR/Cas9 genome

editing technology (61). RNA guides targeting Exon 4 of the *Tfap2c* gene (Supplemental

Figure 4, Supplemental Table 4) were assembled with *CRISPR* RNA, trans-activating

CRISPR RNA, and Cas9 nuclease V3 (Integrated DNA Technologies). Conditional

Tfap2c disruption by the insertion of loxP sites targeted to 5' and 3' introns flanking Exon

4 (Figure 3, Supplemental Table 4, and Supplemental Table 5). Electroporation of

constructs were performed into one-cell rat embryos using a NEPA21 electroporator

(Nepa Gene Co Ltd). Electroporated embryos were transferred to pseudopregnant rats.

Offspring were screened for mutations by polymerase chain reaction (**PCR**), and the

deletion confirmed by DNA sequencing (GeneWiz). A founder mutant animal was

backcrossed with wild type rats to evaluate germline transmission. Genotyping and fetal

sex chromosome determination were performed as previously described (47, 61, 62).

DNA was extracted from tail-tip biopsies with Red Extract-N-Amp tissue PCR kit (XNAT-

1000RXN, Sigma-Aldrich) and used for genotyping. Primers used for genotyping of

global and conditional mutations and sex chromosome determinations are provided

(Supplemental Table 6). Body weight measurements were performed on offspring

generated from specific intercrosses.

Reverse transcription-quantitative PCR (RT-qPCR)

Total RNA was isolated from tissues with TRIzol (15596018, Thermo Fisher).

Extracted RNA (1 µg) was used to synthesize complementary DNA using a High-

Capacity Reverse Transcription kit (4368814, Applied Biosystems) and then diluted 1:10

in water. qPCR was performed using PowerSYBR Green PCR Master Mix (4367659,

Thermo Fisher) and transcript-specific primer sets (250 nM). Primer sequences are

provided (Supplemental Table 7). QuantStudio 5 Real-Time PCR system (Thermo

Fisher) cycling conditions were as follows: an initial step (95 °C for 10 min), preceded by

40 cycles of two-step PCR (95 °C for 15 s, 60 °C for 1 min), and then a dissociation step

(95 °C for 15 s, 60 °C for 1 min), and a sequential increase to (95 °C for 15 s). Relative

mRNA expression was calculated using the ΔΔCt method and glyceraldehyde 3-

phosphate dehydrogenase was used as a reference RNA.

In situ hybridization

In situ hybridization was performed using the RNAscope® Multiplex Fluorescent

Reagent kit version 2 (Advanced Cell Diagnostics) according to the manufacturer's

instructions. Images were captured on Nikon 90i upright microscopes (Nikon) with

Photometrics CoolSNAP-ES monochrome cameras (Roper). Probes were designed to

detect *Tfap2c* (860171, NM_201420.2, target region: 696-1733), *Krt8* (87304-C2,

NM_199370.1, target region: 134-1472) and *Prl7b1* (860181-C2, NM_153738.1, target

region: 28-900).

Immunohistochemistry

Frozen rat placentation sites were sectioned at 10 μm, mounted on slides, and fixed in

4% paraformaldehyde. Sections were blocked with 10% goat serum (50062Z, Thermo

Fisher) and incubated overnight at 4°C with a primary antibody for vimentin (1:300, sc-

6260, Santa Cruz Biotechnology), pan-cytokeratin antibody (1:100, F3418, Sigma-

Aldrich), perforin (1:300, TP251, Amsbio), or TFAP2C (1:150, 2320, Cell Signaling

Technology). Fluorescence-tagged secondary antibodies, Alexa Fluor 488-conjugated

goat anti mouse IgG (A11001, Thermo Fisher), and Alexa Fluor 568-conjugated goat

anti-rabbit IgG (A11011, Thermo Fisher) were used to visualize immunostaining.

Sections were counterstained with 4,6-diamidino-2-phenylindole (DAPI, 1:50,000,

D1306, Invitrogen). Slides were mounted in Fluoromount-G (0100-01, SouthernBiotech)

and imaged on Nikon 90i upright microscopes with Photometrics CoolSNAP-ES

monochrome cameras (Roper). Regions of the uterine-placental interface possessing

cytokeratin-positive cells were quantified using ImageJ software, as previously described

by our laboratory (63, 64).

Western blotting

Tissue lysates were processed using the radioimmunoprecipitation assay lysis

buffer (sc-24948A, Santa Cruz Biotechnology) and protein concentrations determined

using the DC Protein Assay Kit (5000112, Bio-Rad). Proteins (80 μg/lane) were

separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis. Separated

proteins were electrophoretically transferred to polyvinylidene difluoride membranes

(10600023, GE Healthcare) for 1 h at 25 V on a semi-dry transfer apparatus (Bio-Rad).

Membranes were subsequently blocked with 5% milk for 1 h at room temperature,

followed by incubation with antibodies to TFAP2C (1:100, sc-12762, Santa Cruz

Biotechnology) in 5% milk overnight at 4°C. After primary antibody incubation, the

membranes were washed in Tris-buffered saline with Tween 20 (TBS-T) three times (10

min/wash) at room temperature. The membranes were then incubated with anti-mouse

immunoglobulin G conjugated to horseradish peroxidase (HRP; 1:5000, Cell Signaling

Technology) in 5% milk for 1 h at room temperature, washed in TBS-T three times (10

min/wash) at room temperature, immersed in Immobilon Crescendo Western HRP

Substrate (WBLUR0500, Sigma-Aldrich), and luminescence detected using Chemi Doc

MP Imager (Bio-Rad).

Identification of TFAP2C and potential TFAP2C gene targets in rat invasive

trophoblast cells

Uniform manifold approximation and projection (**UMAP**) plots for *Tfap2c* and *Prl7b1*

were generated from rat gd 19.5 uterine-placental interface single cell RNA sequencing

(accession No. GSE206086, (38)). Gene expression profiles and open chromatin

containing TFAP2C DNA binding motifs identified in rat invasive trophoblast cells (38,

41) were integrated and ascribed function through manual annotation using the UniProt

(https://www.uniprot.org) and National Center for Biotechnology

(https://www.ncbi.nlm.nih.gov/) databases.

Statistical analysis

Statistical analyses were performed with GraphPad Prism 10.2.3 software. Statistical

comparisons were evaluated using Student's *t* test or one-way ANOVA with Tukey's or

Dunnett's post hoc tests as appropriate. Statistical significance was determined as

$P<0.05$.

Author Contributions

E.M.D., K. I., and M.J.S. conceived and designed the research; E.M.D. and A.M-I.,

performed experiments; K.I. provided experimental tools for analysis; E.M.D., A.M-I.,

R.L.S, and M.J.S. analyzed the data and interpreted results of experiments; E.M.D. and

M.J.S. prepared the manuscript. All authors read, contributed to editing, and approved

the final version of manuscript.

Acknowledgments

We thank Brandi Miller, Stacy Oxley, and Leslie Tracy for administrative assistance.

The work was supported by the Lalor Foundation (E.M.D., A.M.-I.), Kansas Idea Network

of Biomedical Research Excellence, P20 GM103418 (E.M.D, A.M.-I.), NIH grants:

HD115834 (A.M.-I.), HD104495 (R.L.S.), HD104071 (K.I.), HD020676 (M.J.S.),

HD105734 (M.J.S.), HD112559 (M.J.S.), and the Sosland Foundation (M.J.S.).

References

1. Georgiades P, Ferguson-Smith AC, Burton GJ. Comparative developmental anatomy of the

murine and human definitive placentae. *Placenta*. 2002;23(1):3–19.

2. Maltepe E, Fisher SJ. Placenta: the forgotten organ. *Annu Rev Cell Dev Biol*. 2015;31:523–552.

3. Burton GJ, Fowden AL, Thornburg KL. Placental Origins of Chronic Disease. *Physiol Rev*.

2016;96(4):1509–1565.

4. Enders AC, Welsh AO. Structural interactions of trophoblast and uterus during hemochorial

placenta formation. *J Exp Zool*. 1993;266(6):578–587.

5. Roberts RM, Green JA, Schulz LC. The evolution of the placenta. *Reproduction*.

2016;152(5):R179-189.

6. Hertig AT (Arthur T. *Human trophoblast*. Springfield, Ill. : C. C. Thomas; 1968.

7. Vićovac L, Aplin JD. Epithelial-mesenchymal transition during trophoblast differentiation. *Acta*

Anat (Basel). 1996;156(3):202–216.

8. Pijnenborg R, et al. Review article: trophoblast invasion and the establishment of

haemochorial placentation in man and laboratory animals. *Placenta*. 1981;2(1):71–91.

9. Ain R, Canham LN, Soares MJ. Gestation stage-dependent intrauterine trophoblast cell

invasion in the rat and mouse: novel endocrine phenotype and regulation. *Dev Biol*.

2003;260(1):176–190.

10. Velicky P, Knöfler M, Pollheimer J. Function and control of human invasive trophoblast

subtypes: Intrinsic vs. maternal control. *Cell Adh Migr*. 2016;10(1–2):154–162.

11. Kaufmann P, Black S, Huppertz B. Endovascular trophoblast invasion: implications for the

pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod*. 2003;69(1):1–7.

12. Red-Horse K, et al. Trophoblast differentiation during embryo implantation and formation of

the maternal-fetal interface. *J Clin Invest*. 2004;114(6):744–754.

13. Pijnenborg R, Vercruysse L, Hanssens M. The uterine spiral arteries in human pregnancy:

facts and controversies. *Placenta*. 2006;27(9–10):939–958.

14. Harris LK. Review: Trophoblast-vascular cell interactions in early pregnancy: how to remodel

a vessel. *Placenta*. 2010;31 Suppl:S93-98.

15. Whitley GSJ, Cartwright JE. Cellular and molecular regulation of spiral artery remodelling:

lessons from the cardiovascular field. *Placenta*. 2010;31(6):465–474.

16. Brosens I, Puttemans P, Benagiano G. Placental bed research: I. The placental bed:

from spiral arteries remodeling to the great obstetrical syndromes. *Am J Obstet Gynecol*.

2019;221(5):437–456.

17. Zhang L, et al. Unraveling the mysteries of spiral artery remodeling. *Placenta*. 2023;141:51–

56.

18. Hilger-Eversheim K, et al. Regulatory roles of AP-2 transcription factors in vertebrate

development, apoptosis and cell-cycle control. *Gene*. 2000;260(1–2):1–12.

19. Eckert D, et al. The AP-2 family of transcription factors. *Genome Biol*. 2005;6(13):246.

20. Kuckenberg P, Kubaczka C, Schorle H. The role of transcription factor Tcfap2c/TFAP2C in

trophectoderm development. *Reprod Biomed Online*. 2012;25(1):12–20.

21. Hemberger M, Hanna CW, Dean W. Mechanisms of early placental development in mouse

and humans. *Nat Rev Genet*. 2020;21(1):27–43.

22. Papuchova H, Latos PA. Transcription factor networks in trophoblast development. *Cell Mol*

Life Sci. 2022;79(6):337.

23. Auman HJ, et al. Transcription factor AP-2gamma is essential in the extra-embryonic lineages

for early postimplantation development. *Development*. 2002;129(11):2733–2747.

24. Werling U, Schorle H. Transcription factor gene AP-2 gamma essential for early murine

development. *Mol Cell Biol*. 2002;22(9):3149–3156.

25. Kuckenberg P, et al. The transcription factor TCFAP2C/AP-2gamma cooperates with CDX2 to

maintain trophectoderm formation. *Mol Cell Biol*. 2010;30(13):3310–3320.

26. Cao Z, et al. Transcription factor AP-2γ induces early Cdx2 expression and represses HIPPO

signaling to specify the trophectoderm lineage. *Development*. 2015;142(9):1606–1615.

27. Benchetrit H, et al. Extensive Nuclear Reprogramming Underlies Lineage Conversion into

Functional Trophoblast Stem-like Cells. *Cell Stem Cell*. 2015;17(5):543–556.

28. Kubaczka C, et al. Direct Induction of Trophoblast Stem Cells from Murine Fibroblasts. *Cell*

Stem Cell. 2015;17(5):557–568.

29. Shi D, Kellems RE. Transcription Factor AP-2γ Regulates Murine Adenosine Deaminase Gene

Expression during Placental Development*. *Journal of Biological Chemistry*.

1998;273(42):27331–27338.

30. LiCalsi C, et al. AP-2 family members regulate basal and cAMP-induced expression of human

chorionic gonadotropin. *Nucleic Acids Res*. 2000;28(4):1036–1043.

31. Sapin V, et al. Differential expression of retinoic acid-inducible (Stra) genes during mouse

placentation. *Mech Dev*. 2000;92(2):295–299.

32. Ben-Zimra M, et al. Uterine and placental expression of steroidogenic genes during rodent

pregnancy. *Molecular and Cellular Endocrinology*. 2002;187(1):223–231.

33. Peng L, Payne AH. AP-2 gamma and the homeodomain protein distal-less 3 are required for

placental-specific expression of the murine 3 beta-hydroxysteroid dehydrogenase VI gene,

Hsd3b6. *J Biol Chem*. 2002;277(10):7945–7954.

34. Oztürk A, et al. Proteomic identification of AP2 gamma as a rat placental lactogen II

trophoblast cell-specific enhancer binding protein. *Endocrinology*. 2006;147(9):4319–4329.

35. Ushizawa K, et al. Global gene expression analysis and regulation of the principal genes

expressed in bovine placenta in relation to the transcription factor AP-2 family. *Reprod Biol

Endocrinol*. 2007;5:17.

36. Kidder BL, Palmer S. Examination of transcriptional networks reveals an important role for

TCFAP2C, SMARCA4, and EOMES in trophoblast stem cell maintenance. *Genome Res*.

2010;20(4):458–472.

37. Oikawa M, et al. Generation of Tfap2c-T2A-tdTomato knock-in reporter rats via adeno-

associated virus-mediated efficient gene targeting. *Molecular Reproduction and Development*.

2022;89(3):129–132.

38. Scott RL, et al. Conservation at the uterine–placental interface. *Proceedings of the National*

Academy of Sciences. 2022;119(41):e2210633119.

39. Varberg KM, et al. Extravillous trophoblast cell lineage development is associated with active

remodeling of the chromatin landscape. *Nat Commun*. 2023;14(1):4826.

40. Kim M, et al. The transcriptional regulatory network modulating human trophoblast stem

cells to extravillous trophoblast differentiation. *Nat Commun*. 2024;15(1):1285.

41. Vu HTH, et al. Core conserved transcriptional regulatory networks define the invasive

trophoblast cell lineage. *Development*. 2023;150(15):dev201826.

42. Soares MJ, et al. RAT PLACENTATION: AN EXPERIMENTAL MODEL FOR INVESTIGATING THE

HEMOCHORIAL MATERNAL-FETAL INTERFACE. *Placenta*. 2012;33(4):233–243.

43. Shukla V, Soares MJ. Modeling Trophoblast Cell-Guided Uterine Spiral Artery Transformation

in the Rat. *Int J Mol Sci*. 2022;23(6):2947.

44. Knipp GT, Audus KL, Soares MJ. Nutrient transport across the placenta. *Advanced Drug*

Delivery Reviews. 1999;38(1):41–58.

45. Soares MJ, et al. Differentiation of trophoblast endocrine cells. *Placenta*. 1996;17(5–6):277–

289.

46. Wiemers DO, et al. The mouse prolactin gene family locus. *Endocrinology*. 2003;144(1):313–

325.

47. Iqbal K, et al. Conditionally mutant animal model for investigating the invasive trophoblast

cell lineage. *Development*. 2024;151(2):dev202239.

48. Chakraborty D, et al. Natural killer cells direct hemochorial placentation by regulating

hypoxia-inducible factor dependent trophoblast lineage decisions. *Proc Natl Acad Sci U S A*.

2011;108(39):16295–16300.

49. Amoroso EC. The evolution of viviparity. *Proc R Soc Med*. 1968;61(11 Pt 2):1188–1200.

50. Li L, et al. Lineage regulators TFAP2C and NR5A2 function as bipotency activators in

totipotent embryos. *Nat Struct Mol Biol*. 2024;31(6):950–963.

51. Zhu M, et al. Developmental clock and mechanism of de novo polarization of the mouse

embryo. *Science*. 2020;370(6522):eabd2703.

52. Zhu M, et al. Tead4 and Tfap2c generate bipotency and a bistable switch in totipotent

embryos to promote robust lineage diversification. *Nat Struct Mol Biol*. 2024;31(6):964–976.

53. Gardner RL, Beddington RS. Multi-lineage "stem" cells in the mammalian embryo. *J Cell Sci

Suppl*. 1988;10:11–27.

54. Rossant J. Stem cells from the Mammalian blastocyst. *Stem Cells*. 2001;19(6):477–482.

55. Kaiser S, et al. Reduced Gene Dosage of Tfap2c Impairs Trophoblast Lineage Differentiation

and Alters Maternal Blood Spaces in the Mouse Placenta1. *Biology of Reproduction*.

2015;93(2):31, 1–13.

56. Schemmer J, et al. Transcription factor TFAP2C regulates major programs required for

murine fetal germ cell maintenance and haploinsufficiency predisposes to teratomas in male

mice. *PLoS One*. 2013;8(8):e71113.

57. Loureiro-Campos E, et al. Constitutive deficiency of the neurogenic hippocampal modulator

AP2γ promotes anxiety-like behavior and cumulative memory deficits in mice from juvenile to

adult periods. *Elife*. 2021;10:e70685.

58. Sharma N, et al. Tpbpa-Cre-mediated deletion of TFAP2C leads to deregulation of Cdkn1a,

Akt1 and the ERK pathway, causing placental growth arrest. *Development*. 2016;143(5):787–

798.

59. Latos PA, et al. Elf5-centered transcription factor hub controls trophoblast stem cell self-

renewal and differentiation through stoichiometry-sensitive shifts in target gene networks.

Genes Dev. 2015;29(23):2435–2448.

60. Ain R, et al. Phenotypic analysis of the rat placenta. *Methods Mol Med*. 2006;121:295–313.

61. Iqbal K, et al. Catechol-O-methyltransferase and Pregnancy Outcome: an Appraisal in Rat.

Reprod Sci. 2021;28(2):462–469.

62. Dhakal P, Soares MJ. Single Step PCR-Based Genetic Sex Determination for Rat Tissues and

Cells. *Biotechniques*. 2017;62(5):232–233.

63. Rosario GX, Konno T, Soares MJ. Maternal hypoxia activates endovascular trophoblast cell

invasion. *Dev Biol*. 2008;314(2):362–375.

64. Nteeba J, et al. Poorly controlled diabetes mellitus alters placental structure, efficiency, and

plasticity. *BMJ Open Diabetes Res Care*. 2020;8(1):e001243.

Figures

Figure 1. *Tfap2c* transcript distribution during placenta development. **A-C**) *Tfap2c*
transcripts (magenta) were localized to the ectoplacental cone at gestation day (gd) 9.5
(**A**) and placentation sites at 11.5 (**B**), and 18.5 (**C**). *Tfap2c* transcripts were co-localized
with *Krt8* transcripts (cyan). Schematic diagrams representing gd 9.5, 11.5, and 18.5
placentation sites are initially presented for each panel. **D**) Co-localization of *Tfap2c*
(magenta) and *Prl7b1* (yellow) transcripts within the gd 18.5 placentation site. *Prl7b1* is
specifically expressed in the invasive trophoblast cell lineage. Scale bar: 500 μm. **E**)
High magnification images showing *Tfap2c* and *Prl7b1* transcript localization in
endovascular (arrowhead) and interstitial (arrow) invasive trophoblast cells within the
uterine-placental interface. Scale bar: 100 μm. (**F**) Co-localization of *Tfap2c* and *Prl7b1*
transcripts (arrows) within the junctional zone of the gd 15.5 placenta. Scale bar: 50 μm.
Co-localization is shown as whilte. Abbreviations: EPC, ectoplacental cone; AMD, anti-

561 mesometrial decidua; SpA, spiral artery; UPI, uterine-placental interface; P, placenta; JZ,
562 junctional zone; LZ, labyrinth zone.
563

Figure 2. *Tfap2c* gene dosage effects on placental development. Placentation sites were generated from *Tfap2c*$^{+/-}$ male and *Tfap2c*$^{+/+}$ female breeding. **A)** Identification of placentation site compartments *Tfap2c*$^{+/+}$ and *Tfap2c*$^{+/-}$ using vimentin immunostaining (magenta). **B**) Immunohistochemical localization of cytokeratin protein in gestation day (gd) 18.5 *Tfap2c*$^{+/+}$ and *Tfap2c*$^{+/-}$ placentation sites (cyan). **C**) Distribution of *Prl7b1* transcripts in gd 18.5 *Tfap2c*$^{+/+}$ and *Tfap2c*$^{+/-}$ placentation sites (yellow). **D)** Quantification of trophoblast cell invasion area (magenta dotted line) determined by cytokeratin immunostaining within the uterine-placental interface. Data are expressed as mean $\pm$ standard error of the mean (**SEM**). Each data point represents different uterine-placental interface tissues (n=4) obtained from four pregnancies. **E)** RT-qPCR of invasive trophoblast cell-specific transcripts in gd 18.5 *Tfap2c*$^{+/+}$ and *Tfap2c*$^{+/-}$ uterine-placental interface tissues. Data are expressed as mean $\pm$ SEM. Each data point represents a biological replicate obtained from six different pregnancies (n=6). **F)** Fetal, placenta, junctional zone and labyrinth zone weights for gd 18.5 *Tfap2c*$^{+/+}$ and *Tfap2c*$^{+/-}$ conceptuses. Data are expressed as mean $\pm$ SEM. Each data point represents a biological replicate obtained from six different pregnancies (*Tfap2*+$^{+/+}$, n=38; *Tfap2c* $^{+/-}$, n=32). Unpaired *t*-test: * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Abbreviations: UPI, uterine-placental interface; JZ, junctional zone; LZ, labyrinth zone; SpA, spiral artery. Scale bars: 500 μm.

Figure 3. Generation of an invasive trophoblast cell-specific *Tfap2c* deletion. A)
Schematic of *loxP* sites inserted into introns flanking Exon 4 of the *Tfap2c* gene. **B)**
TFAP2C western blot of the uterine-placental interface of *Tfap2c^{f/f}* and *Tfap2c^{d/d}* gd 18.5
placentation sites. **C)** TFAP2C immunostaining of *Tfap2c^{f/f}* and *Tfap2c^{d/d}* gd 18.5
placentation sites (left panels: TFAP2C immunostaining, white; right panels: TFAP2C
immunostaining, magenta, DAPI, white). Scale bar: 500 μm. Inserts to the right show
higher magnification images of uterine-placental interface tissues (top) and junctional
zone tissue (bottom) for each placentation site. Scale bar: 100 μm. Abbreviations: UPI,
uterine-placental interface; JZ, junctional zone; LZ, labyrinth zone; SpA, spiral artery.

Figure 4. Characterization of invasive trophoblast cell-specific *Tfap2c* deletion. A)
Immunohistochemical localization of cytokeratin (magenta) in gd 18.5 *Tfap2c^f/f* and
Tfap2c^d/d placentation sites. **B**) Distribution of *Prl7b1* transcripts (yellow) in gd 18.5
Tfap2c^f/f and *Tfap2c^d/d* placentation sites. Scale bar: 500 μm. Inserts to the right show
higher magnification images of the boxed areas within the uterine-placental interface
tissues for each placentation site. Scale bar: 100 μm **C)** Quantification of trophoblast cell
invasion area (green dotted line) determined by cytokeratin immunostaining within the
uterine-placental interface. Data are expressed as mean ± standard error of the mean
(**SEM**). Each data point represents different uterine-placental interface tissues obtained
from four pregnancies (n=4). **D)** RT-qPCR of invasive trophoblast cell-specific transcripts

in gd 18.5 *Tfap2c*^{f/f} and *Tfap2c*^{d/d} uterine-placental interface tissues. The left panel

corresponds to a schematic of the regions of the placentation site used for analysis.

Data are expressed as mean $\pm$ SEM. Each data point represents a biological replicate

from six different pregnancies (n=6). **E**) Table of cellular processes potentially regulated

by TFAP2C gene targets. Unpaired *t*-test: *$p < 0.05$, ***$p<0.001$, ****$p<0.0001$.).

Abbreviations: UPI, uterine-placental interface; JZ, junctional zone; LZ, labyrinth zone;

SpA, spiral artery.

Figure 5. *Tfap2c* conditional deletion in trophoblast invasive cells results in the retention of uterine natural killer (NK) cells within the uterine-placental interface.
A) Immunohistochemical localization of cytokeratin (cyan) and perforin (magenta) in gd 18.5 *Tfap2c^f/f^* and *Tfap2c^d/d^* placentation sites. **B)** RT-qPCR of NK cell-specific transcripts in gd 18.5 *Tfap2c^f/f^* and *Tfap2c^d/d^* uterine-placental interface tissues. The left panel corresponds to a schematic of the regions of the placentation site used for analysis. Data are expressed as mean ± standard error of the mean. Each data point represents a biological replicate from six different pregnancies (n=6). Unpaired *t*-test: *p< 0.05, **p<0.01, ***p<0.001, ****p<0.0001. Abbreviations: UPI, uterine-placental interface; JZ, junctional zone; LZ, labyrinth zone; SpA, spiral artery. Scale bar: 500 μm.

Figure 6. Impact of an invasive trophoblast cell-specific *Tfap2c* deletion on prenatal and postnatal development. A) Fetal, placenta, junctional zone, and labyrinth zone weights from gd 18.5 *Tfap2c^{f/f}* and *Tfap2c^{d/d}* gd 18.5 conceptuses. Data are expressed as mean ± standard error of the mean (**SEM**). Each data point represents a biological replicate from six pregnancies (*Tfap2c^{f/f}*, n=20; *Tfap2c^{d/d}*, n= 16). Unpaired *t*-test: *p< 0.05, **p<0.01, ***p<0.001, ****p<0.0001. **B**) Postnatal body weight for *Tfap2c^{f/f}* and *Tfap2c^{d/d}* pups. Data are expressed as mean ± SEM from six pregnancies (*Tfap2c^{f/f}*, n=24; *Tfap2c^{d/d}*, n= 29). Two-Way ANOVA and Tukey's Test: **p<0.01, ****p<0.0001. Abbreviations: JZ, junctional zone; LZ, labyrinth zone.

639 **Supplemental information**

640   Figures S1 to S6 and Table S1 to S7.

641   Dataset S1. Excel file.