High-throughput measurement of adipocyte size with open-source software using whole-slide adipose tissue images

Authors:

Ramalho, Alan[1,2]; Gauthier, Marie-Frédérique[1]; Maltais-Payette, Ina[1]; Ostinelli, Giada[1]; Hould, Frédéric[1]; Biertho, Laurent[1]; Tchernof, André[1,2]

Author's affiliations :

[1]Institut universitaire de Cardiologie et de Pneumologie de Québec-Université Laval, Québec, Québec, Canada

[2]École de Nutrition, Université Laval, Québec, Québec, Canada

Corresponding author

André Tchernof, PhD,

Institut Universitaire de Cardiologie et Pneumologie de Québec (IUCPQ), Université Laval,

2725 Chemin Sainte-Foy, Québec, QC, G1V 4G5, Canada.

Email : andre.tchernof@criucpq.ulaval.ca

ORCID

André Tchernof http://orcid.org/0000-0002-2587-1000

Keywords

Adipocytes, Adipose tissue, cell size, ImageJ, Histomorphometry

Abstract

The aim of this study was to create and validate a high-throughput method based on open-source

software for the measurement of adipocyte diameters in white adipose tissue histological sections.

Human omental and subcutaneous adipose tissue samples collected during bariatric surgery were

used to prepare hematoxylin and eosin-stained histological slides. Digital images were acquired.

Adipocyte diameters were measured both manually and with an automated procedure created using

ImageJ. Comparative analysis of our automated method with the manual measurement and

associations of the mean adipocyte diameters with cardiometabolic markers were used to validate

our method. A total of 377 adipose samples (190 participants) were included in the analysis.

Pearson correlation of mean adipocyte diameters shows a strong linear relationship between

methods (r=0.88, p<0.0001). The average diameter measured with the automated method was

significantly smaller (8.1±5.3µm difference, p<0.0001) compared to the manual method, likely

reflecting bias in selecting the cells measured with the manual approach. Pearson correlation

analyses between mean omental adipocyte diameters and markers of cardiometabolic risk show

that the diameters of both methods are significantly associated with the same parameters (fasting

concentrations of TG, HDL-Chol, homeostasis model assessment insulin resistance, and visceral

adiposity index values) with no significant differences between methods. There were also no

significant differences between the manual and automated method regarding the correlations

between mean subcutaneous adipocyte diameters and anthropometric or metabolic markers. In

conclusion, we have created and validated a rapid automated method based on open-source

software to measure adipocyte diameters from whole-slide adipose tissue images.

1- Introduction

The body mass index (BMI) is a poor indicator of individual cardiometabolic risk, and other metrics that better estimate this risk are increasingly favored in clinical practice and in research [1]. In this regard, adipose tissue dysfunction is a major driver of obesity-related cardiometabolic dysregulation [2–5]. Dysfunctional adipose tissue is characterized by reduced adipogenesis, low free fatty acid uptake, impaired triglyceride synthesis, impaired lipid storage, insulin resistance, fibrosis, immune cell infiltration, increased proinflammatory cytokine secretion, and adipocyte hypertrophy [2,4]. It is also strongly associated with visceral obesity, that is, excessive accumulation of fat in the abdominal depots surrounding the internal organs such as the omental (OM) and mesenteric fat depots [2,4], consistent with the notion that visceral obesity is an important risk factor for the development of cardiometabolic dysregulation and cardiometabolic diseases [2,4].

Adipose tissue expands via two main mechanisms: adipocyte hyperplasia where preadipocytes are recruited and develop into small new mature adipocytes through adipogenesis, and adipocyte hypertrophy where existing adipocytes increase in size to accommodate the storage of more lipids [2–5]. Studies have shown that expansion via hyperplasia, typical of subcutaneous (SC) depots, seems to favor the preservation of cardiometabolic health [2,3]. Conversely, expansion via hypertrophy is an important indicator of adipose tissue dysfunction and increased cardiometabolic risk [2,3]. Indeed, adipocyte hypertrophy is often implicated in visceral obesity and has been shown to be associated with an increased secretion of proinflammatory factors as well as with altered lipid metabolism, impaired glucose-insulin homeostasis, ectopic fat accumulation, and cardiovascular disorders, independently of BMI [2–4,6].

Consequently, measuring adipocyte size can provide indirect information on adipose tissue

function, fatty acid and glucose metabolism, and cardiometabolic pathophysiology [4,6,7].

Analysis of adipocyte size can also help better understand the pathophysiological processes of

obesity and cardiometabolic diseases as well as to characterize treatment responses to dietary

interventions, nutraceuticals, bioactive compounds, and medication [8,9].

Currently, there is no gold standard method to measure adipocyte size [4,6]. There are three main

methods utilized in research: imaging of collagenase digestion-isolated adipocytes, sizing of

osmium tetroxide-fixed cells, and histomorphometry [3,4]. Histomorphometry involves analyzing

hematoxylin and eosin (H&E) -stained histological slides of adipose tissue using an optical

microscope to measure the diameter or cross-sectional area of the adipocytes [4]. This method is

considered to be the most reliable, but it still has drawbacks [8]. First, there are several inherent

assumptions including a uniform cell size distribution and that cells are showing their largest

diameter or cross-sectional area [4]. Furthermore, fixation of the tissue reduces volume of the cells

by approximately 15-20%, directly measuring the diameter is not reliable for irregularly shaped

cells, distortion from poor fixation can occur, crush artifacts can be introduced during processing,

and the quality of the stain can vary between laboratories [4,6,8]. Differences can also exist

regarding specificities in cell sizing protocols which may influence the final results and the ability

to compare results between laboratories [10,11]. The most important drawback, however, is that

this method is very labor-intensive, time consuming, and has a high risk of measurement and

observer bias [7,8,12]. Due to the time required, only 100 to 300 cells per sample are typically

measured, decreasing measurement accuracy [12,13]. These are inherent issues of manual

histomorphometry as it requires the measurer to visually identify each adipocyte in the tissue

section and then measure their diameter or cross-sectional surface area. With the advent of

biological image analysis software, such as ImageJ (National Institutes of Health, USA), the

measurement of adipocyte surface area has become easier. ImageJ is a free easily accessible open-

source scientific image processing and analysis software platform originally created by the

National Institute of Health of the United States [14]. ImageJ is relatively easy to use and has been

widely adopted by the scientific community.

To address the main drawbacks of the histomorphometric method, several teams have developed

automated programs or protocols to detect and measure adipocyte size using histological slides

[7–9,11,12,15–18]. However, many of these automated solutions have issues such as adaptability,

cost, reliability, accessibility (both short- and long-term), and usability. Many tools cannot be

adapted sufficiently to account for considerable procedural differences such as sample quality

preservation, paraffin inclusion, microtomy, staining, and even imaging [10]. The cost of paid

software solutions as well at the costs of the equipment required can be prohibitory [12]. Some

software tools have been discontinued or are not accessible through secure channels [7,9,17,18].

Many of the current software solutions typically analyze digital photomicrographs of the tissue

sample, limiting the number of adipocytes measured [9,12,13]. Photomicrographs are used for

manual analysis as manually measuring each adipocyte in a complete section would require an

exorbitant amount of time. Cell sizing software tools should be capable of analyzing whole-slide

images of the tissue samples to maximize the number of adipocytes measured [12]. However,

many of the free software options were not designed to do this due to file size restrictions. Recent

advancements in both file size capacity of software tools and the affordability of slide scanners for

the acquisition of high-quality whole-slide images have opened the door for significant progress

in this regard [12].

Ultimately, there are important issues with the current histomorphometric methods of fat cell

sizing. The aim of this study was to develop and validate an easy-to-use method to measure

adipocyte size in whole tissue images. We hypothesized that measurements can be made

automatically and that average adipocyte diameter measurements obtained with the automated

method are consistent with those obtained with the manual method, while requiring significantly

less time and measuring a larger number of cells per sample. Additionally, we hypothesized that

the results obtained with the automated method are equivalent or superior to those of the manual

method for the prediction of cardiometabolic risk markers.

2- Results

2.1 Manual and automated cell size measurements

Regarding the initial elaboration phase of our study, the average adipocyte diameters from 16

samples obtained with the manual and automated methods were compared for the purpose of

establishing measurement parameters. Spearman correlation analysis showed a strong significant

association between the methods (r=0.90, p <0.0001). Paired t-test analysis indicated that the

average adipocyte diameters obtained with the two methods were not significantly different

(p=0.10) (data not shown).

Table 1: Participants' anthropometric characteristics and metabolic markers levels. Data are
expressed as the mean ± SD and minimum and maximum values.

	Mean ± SD	Min - Max
n	190	
Sex (men/women)	46:144	
Age (years)	38.0 ± 8.6	19 - 59
BMI (kg/m^2)	49.0 ± 7.3	35.6 – 74.8
Visceral adiposity index (VAI)	2.65 ± 1.62	0.69 – 11.26

Waist circumference (cm)	135 ± 14	106 – 184
Triglycerides (mmol/L)	1.64 ± 0.76	0.45 – 5.20
HDL-cholesterol (mmol/L)	1.18 ± 0.27	0.69 – 2.06
LDL-cholesterol (mmol/L)	2.73 ± 0.73	1.13 – 6.25
Total cholesterol (mmol/L)	4.63 ± 0.88	2.54 – 9.47
Glucose (mmol/L)	5.4 ± 0.4	4.2 – 7.7
HOMA-IR	6.2 ± 3.3	0.9 – 17.3
Apo B (g/L)	0.96 ± 0.21	0.52 – 1.96

For the validation study, a total of 370 and 374 samples (for the manual method and automated

method, respectively) from 190 participants were included in the analysis (**Table 1 and Table 2**).

As shown in **Table 2**, the average adipocyte diameter was $77.1 \pm 11.0\,\mu m$ and $69.1 \pm 10.5\,\mu m$ for

the manual and automated method, respectively. These average diameters were shown to be

significantly different by paired t-test ($p<0.0001$) (data not shown). The approximate amount of

time dedicated to the manual analysis was 10 minutes per tissue compared to 0.8 minute per tissue

for the automated method (based on a subgroup of 300 samples). The automated method detected

an average of 878 ± 511 cells compared to 130 ± 34 cells for the manual analysis ($p<0.0001$).

Table 2: Summary results of all adipocytes diameter (OM and SC) with the manual and automated
method. Data are expressed as the mean ± SD for the measured cell number and cell diameter. Time
required for analysis includes file processing and handling.

	Manual method	Automated method
N	370	374
Time required for analysis	~ 10 minutes/section	~ 0.8 minutes/section
Average number of cells measured ±SD	130 ± 34 cells	878 ± 511 cells
Average cell diameter ± SD	77.1 ± 11.0 µm	69.1 ± 10.5 µm

2.2 Comparisons between manual and automated cell size measurements

A Pearson correlation analysis was performed between the manual and automated cell size (for the

two depots) and is presented in **Figure 1a**. The Pearson coefficient of r=0.88 indicates that the two

methods are strongly consistent. We observed in the Bland Altman plot in **Figure 1b** an average

difference of -8.10 µm between the mean automated and manual cell diameters, as well as limits

of agreement of 2.22 µm and -18.43 µm. Most of the data points are within the limits of agreement,

indicating good agreement between methods. The magnitude of the difference does not appear to

vary based on the average diameter, suggesting the absence of systematic bias related to the

magnitude of the measurement [19]. **Figures 1c and 1d** show that the average cell size measured

with the automated method is significantly smaller compared to the manual method in both the

visceral and SC depots (-9.55 µm, $p < 0.0001$ and -6.61 µm, $p < 0.0001$, respectively) and this is also

true for the majority of individual samples.

Figure 1: Comparison of adipocyte diameters measured by the two methods. Pearson correlation in all cell diameter (OM and SC) between the manual and automated method **(a)** and Bland-Atlman plot with size differences **(b)**. Paired t-test between the two methods in omental **(c)** and subcutaneous **(d)** adipose tissues.

2.3 Associations with metabolic markers

The participants for this study were on average 38.0 ± 8.6 years old, including 46 men and 144 women with an average BMI of 49.0 ± 7.4 kg/m^2 and an average waist circumference of 135 ± 14 cm (**Table 1**). Pearson correlation analyses were performed to evaluate the association between cardiometabolic health markers and the mean adipocyte diameter obtained with each measurement

189 method (i.e. manual and automated) for each adipose tissue depot. The results are presented in

190 **Figure 2**. The automated cell size method was not significantly different from the manual method

191 for any of the cardiometabolic health markers for either depot, as determined using Meng's Z-test

192 (data not shown).

Figure 2: Comparison in association with anthropometrics and metabolic markers. Pearson correlation between mean adipocyte diameter with BMI, WC and cardiometabolic markers (TG, HDL-Chol, HOMA-IR and VAI) between the two methods in omental **(a)** and subcutaneous depot **(b)**.

Regarding the OM depot, we observed in **Figure 2a,** although not significantly different, a slightly

stronger correlation and lower p-value with the automated method with triglyceride (TG) levels

(r=0.28, p<0.0001 vs r=0.23, p=0.003), HDL-Cholesterol (HDL-Chol) (r=-0.34, p<0.0001 vs r=-

0.32, p<0.0001), the homeostatic model assessment of insulin resistance (HOMA-IR) (r=0.44,

p<0.0001 vs r=0.40, p<0.0001), and the visceral adiposity index (VAI) (r=0.28, p=0.0001 vs

r=0.24, p=0.001) compared to the manual method.

Regarding the SC depot (**Figure 2b**), although not significantly different, a slightly higher

correlation strength was observed with the automated method for the association with HDL-Chol

(r=-0.16, p=0.03 vs r=-0.09, p=0.24), HOMA-IR (r=0.21, p=0.01 vs r=0.17, p=0.05) and VAI

(r=0.18, p=0.02 vs r=0.13, p=0.09) compared to the manual method. Notably, the association with

HDL-Chol and VAI became significant with the automated method.

No significant associations were observed with total Cholesterol (Chol), LDL-Chol, and

apolipoprotein B (apoB) with either method in either depot.

3- Discussion

Our analysis shows that our automated method for measuring adipocyte size is rapid, valid, and

reliable. Pearson correlation analysis shows a strong agreement between both methods for the

determination of mean adipocyte diameters. Pearson correlation analyses and Meng's Z-test also

demonstrate that the associations between different cardiometabolic health markers and mean

adipocyte diameters measured using the automated method were equivalent to the associations

with the manual method. Adipocyte size is often used in research as a marker of adipose tissue

function and cardiometabolic health. As such, the significant associations between other

cardiometabolic health markers and the automated mean adipocyte diameter and their agreement

with the associations with the manual mean adipocyte diameters clearly demonstrate the validity

of our method. It should be noted that the weight status of the participants in this study likely

influenced these associations. Expansion through hypertrophy has its limitations as adipocytes can

only increase in volume up to a certain point [2,4]. Mean fat cell size eventually plateaus in

individuals with large adiposity excess as can be the case with candidates of bariatric surgery

[2,4,11]. This plateauing weakens the ability to detect significant associations with other variables

[4]. This may explain the lack of significant associations with some of the cardiometabolic risk

markers and why the associations observed with BMI and WC were weaker than expected based

on data from other studies [3,13].

Despite the strong association between methods, the paired t-test, grouped violin plots, and the

Bland-Altman plot demonstrate that the automated mean adipocyte diameters are on average

significantly smaller than those of the manual method. The average difference between the two

methods is 8.1 µm. Because there is no gold standard method for cell size measurement, it is

possible that this difference reflects better estimation of average cell size by our automated method

compared to the manual method. The manual method is dependent on the judgement of the user in

both the selection of the tissue section subregions to use and the identification of adipocytes to

measure. The selection of subregions is ideally completely randomized, but many factors such as

section edges, damaged subregions, and fibrotic subregions can influence this selection. This may

result in preferentially identifying better-looking subsections. Large adipocytes may be easier to

notice and identify visually than smaller ones, which could result in an overestimation of average

size with the manual approach. The automated method targets the entire section, thereby

eliminating such biases. Analyzing the entire histological section instead of a few subregions also

greatly increases the number of cells measured for each sample. Increasing the number of cells

measured increases the likelihood that the estimation of average cell size is accurate [12]. Other

teams have obtained similar differences in mean adipocyte size. Maguire et al (2020) and Palomäki

et al (2022) [11,12], both observed that mean cell size was smaller when using an automated

method with whole-slide images compared to methods that used photomicrographs of tissue

sections. Interestingly, Maguire et al (2020) [12] only observed this in high-fat diet mice, and the

opposite was true for chow-fed mice. Like us, Palomäki et al (2022) [11] used adipose tissue

samples from patients with obesity, but the difference they observed between sizing methods was

much smaller at -0.8 µm. This may be due to methodological differences between their study and

ours as they measured approximately 300 cells per sample using photomicrographs while also

using a smaller number of tissue samples [11]. Due to these reasons and the significant correlations

with cardiometabolic risk markers previously discussed, it is possible that our automated method

produces a more reliable estimation of mean adipocyte size compared to the manual method.

Our method has several advantages compared to the manual method. As discussed, our automated

method is significantly faster and uses whole-slide images of H&E-stained tissue sections,

significantly increasing the number of adipocytes measured. The automated detection of

adipocytes reduces the potential of observer and measurement bias. Our method is also very easy

to implement as its main component is a macro for ImageJ, that we provide here and can easily be

copied, making it very accessible. The only software needed is ImageJ, which is free, easily

accessible, and can be run on low-powered computers. The fact that ImageJ was developed by the

NIH and that it has a dedicated userbase provides good assurance that it should be maintained for

the foreseeable future. ImageJ is also easy to learn, especially for this application.

As mentioned, other teams have developed automated protocols using software, but many of these

solutions have issues including software cost, short- and long-term accessibility, and the inability

to use whole-slide images. Notable exceptions to most of these issues are Adiposoft, a plugin

created for ImageJ, and QuPath, another free open-source software for the analysis of digital

biological images [8,12,20]. Adiposoft was created and validated to analyze the number and size

of white adipocytes from adipose tissue histological sections [8,21]. Originally, it only analyzed

photomicrographs but is now capable of analyzing whole-slide images. Indeed, Adiposoft is a

capable program solution, but we encountered issues when attempting to use it with our samples.

While Adiposoft analyzed many of our samples without issue, there were significant problems

with some samples such as adipocytes not being detected and the identification of large empty

areas as adipocytes. Our experience suggests that it may be difficult to adapt Adiposoft, using its

current settings options, to certain experimental conditions. Adiposoft also outputs a unique

spreadsheet for each image, complicating data management when analyzing many samples.

QuPath is a powerful tool for the analysis of whole-slide images and large 2D data [20]. Maguire

et al (2020) developed an interactive plugin for QuPath for the measurement of adipocytes [12].

Their results show that their method reliably measures adipocyte number and size from whole-

slide images [12]. However, their method involves several steps that require both QuPath and

ImageJ. The adaptability of their method to different experimental conditions is also uncertain.

The QuPath method developed by Palomäki et al (2022) only uses tools that are already built into

QuPath and no other analysis software is required. Their analysis shows that their method is valid

for good and average quality whole-slide images and photomicrographs [11]. However, the

method requires training a pixel classifier with good quality histology sections of adipose tissue

[11]. This step increases the complexity and requires technical knowledge, which may hinder other

teams from adopting the method. Ultimately, even with free software solutions capable of

analyzing whole-slide images, there are issues that hinder their adoption. In contrast, our method

is more straightforward and simpler to implement. It requires only ImageJ and our macro, which

is arguably easier. Our macro uses standard image processing methods, which should allow other

teams to easily understand it and adapt it to their experimental conditions. Furthermore, the macro

does not need to be downloaded, maximizing its accessibility.

Our study has several strengths. First, to validate our automated method, we used both correlation

analyses and Bland-Altman plots to evaluate the strength of the linear relationship and agreement

between the methods, respectively. While a strong correlation between methods of measurement

of the same variable is expected, it does not indicate that there is good agreement between methods

[19]. Bland-Altman plots evaluate the differences between the methods, allowing for a better

determination of comparability between methods [19]. Second, 367 white adipose tissue samples

obtained from male and female human participants were analyzed by both methods. Such a large

sample size greatly increases the validity of our analysis and is much larger than samples used by

other teams in the validation of their cell-sizing tools [12,15,16]. Furthermore, we used both SC

and OM adipose tissue samples from patients from two different cohorts, which indicates how our

automated method can be applied to different scenarios. Third, we evaluated how the methods

influence the associations between cardiometabolic health markers and mean cell diameters to

establish the validity and usefulness of our automated method.

This study does have some limitations. First, all the samples came from participants with obesity

undergoing bariatric surgery. As such, it cannot be stated for certain that our method would

adequately perform in populations without obesity. This is unlikely to be an issue due to obesity

being a very heterogeneous disease and that we used a very large sample size which included

varied types of cellular morphologies. Rodent white adipose tissue is similar to that of humans

regarding morphology and gross anatomical regions, indicating rodent tissues could likely be

analyzed with our method [17]. Second, all the histological slides were prepared by the histology

department at our Institute by highly trained technicians using high-quality equipment. The slides

were typically of high quality. As such, our method may not function well with lower quality

slides. However, poor quality slides are an impediment to all sizing methods, and, as mentioned,

our macro can be adapted to optimize its performance to different experimental conditions. Our

method also requires the use of whole-slide images with uniform illumination, but the recent

decrease in the cost of slide scanners alleviates this issue [12]. Third, only mean adipocyte diameter

was considered and not adipocyte size distribution in each sample. As only approximately 100

adipocytes were measured per sample, it is possible that some size subgroups would be

underrepresented with the manual method, resulting in the comparison of the two methods in this

regard to be of questionable relevance.

In conclusion, we have successfully developed a rapid, automated method for the measurement of

adipocytes in H&E-stained histological sections of white adipose tissue. This automated method

consists of a straightforward macro for ImageJ that analyzes whole-slide images of histological

white adipose tissue sections. We have shown that this method is valid and reliable. Compared to

the usual manual method, our automated method allows for the measurement of a much greater

number of cells while requiring considerably less time. Furthermore, our method can easily be

adopted and adapted by other teams, which could increase the uniformity between laboratories.

4- Methods

4.1 Participants characteristics and tissue sampling

Participants are patients who underwent bariatric surgery and were recruited through the Biobank

infrastructure at the *Insititut universitaire de cardiologie et de pneumologie de Québec –*

Université Laval. All participants provided written, informed consent. This study was approved

by the medical ethics committees of Laval University (approval numbers 2017-2710, 2021-3398

and 2019-3218) and adheres to the principles of the Declaration of Helsinki. Clinical and

anthropometric characteristics such as age, sex, height, weight, BMI, and waist circumference

were recorded during the pre-surgery consultation or on the morning of surgery. Fasting blood

samples were obtained and lipid profiles (total Chol, HDL-Chol, LDL-Chol, TG, and apoB levels)

and hematological parameters (fasting insulin, fasting glucose, and HbA1c) were measured by the

biochemistry department of the Institute. OM and SC adipose tissue samples were collected from

the greater omentum and the abdominal SC adipose tissue compartment, respectively, during

surgery and immediately frozen at -80°C. Approximately 60 to 80mg of frozen adipose tissue was

fixed in 10% formalin (Sigma-Aldrich®, HT501128-4L) overnight at 4°C and embedded in

paraffin the following day in accordance with our validated protocol [10]. Five µm sections were

mounted on slides, stained with H&E and used for the manual and automated measurements of

cell size. HOMA-IR was calculated according to the equation: fasting insulin (µU/mL) * fasting

glucose (mmol/L) /22.5 [22]. VAI was calculated using validated sex-specific equations [23]. The

equations are as follows: males: $VAI = \left(\frac{Waist\ Circumference}{39.68+(1.88 X BMI)}\right) * \left(\frac{TG}{1.03}\right) * \left(\frac{1.31}{HDL}\right)$ females: $VAI =$

$\left(\frac{Waist\ Circumference}{36.58+(1.89 X BMI)}\right) * \left(\frac{TG}{0.81}\right) * \left(\frac{1.52}{HDL}\right)$[23].

Initially, a small group of 16 samples were selected for the elaboration of the automated method

and an initial validation. The samples were obtained from four participants (two men and two

women, 45 ± 9.7 years old, BMI 48 ± 6.6 kg/m^2 at the time of bariatric surgery). This group

allowed us to determine the parameters for the automated measurements in a variety of cell size

ranges using adipose tissue samples from two depots.

Subsequently, validation of the method was performed with a larger group composed of 190

participants who had undergone bariatric surgery and were included in this study.

4.2 Manual adipose cell size measurements

Briefly, five subregions with minimal damage and artifacts were randomly acquired from each

histological adipose tissue section using a Leica DMLB microscope (Leica, Wetzlar, Germany)

coupled with a MotiCam580 (Motic, British Columbia, Canada) at 5x magnification for each

sample. Manual histomorphometric analyses were performed blinded to patient identity or type of

depot according to the method described by Laforest et al (2018) [10]. The digital

photomicrographs were converted to 8-bit, the background was subtracted, background noise was

reduced using a median filter, and a manual threshold was applied to convert the images to binary

in order to segment adipocytes using ImageJ (National Institutes of Health, USA). The cross-

sectional area of each adipocyte was manually measured using the wand tool on each image.

Incomplete cells on the edge of the images were excluded. The paintbrush tool was used to smooth

the cell surface or fill-in membranes when needed. A minimum of 100 cells per tissue were

measured. The cross-sectional areas were used to calculate diameters, expressed in µm.

4.3 Automated adipose cell size measurements

Whole-slide images were acquired with an Axio Scan Z1 slide scanner (Zeiss, Oberkochen,

Germany) at 20x magnification. Whole-slide digital images were first converted to 8-bit images

and then into binary using ImageJ. The thresholding method was selected based on the best

segmentation of the adipocytes. Multiple thresholding methods were tested and triangle

thresholding produced the best result for our images. The automatic thresholding compensates for

differences in light intensity during acquisition of H&E staining variation within samples. The

threshold is adjusted and optimized automatically on each image. Other parameters such as

removing outliers, eroding, and despeckling were tested to reduce background noise.

Area analysis was performed through a particle analysis component in ImageJ. The particle

analysis is based on the size and the circularity of objects. Minimum and maximum size cut-offs

of 315 and 40000 µm² (equivalent to a diameter of 20 and 225 µm), respectively, were applied so

all objects inferior to 20 µm and superior to 225 µm in diameter were excluded. The minimum

cut-off enabled the exclusion of objects unlikely to be adipocytes. The maximum cut-off is

important to control for potentially damaged tissues. Objects larger than 225 µm in diameter likely

represent two or more cells where the cell membrane collapsed or was damaged during tissue

processing. This would result in an overestimation of cell size. The roundness of mature adipocytes

allows us to use circularity as an important component of this analysis. Different circularity value

cut-offs were tested during the initial elaboration and validation phase. The circularity cut-off

values have been selected to maximize the number of detected cells while having the best fit with

the manual cell sizes without including false detections due to poor tissue quality (cell damage).

Once the steps of the procedure were established, the different parameters were incorporated into

a macro file in ImageJ to perform automatic fast-paced analysis on all images. The macro is as

follows:

`run("8-bit");`

`setAutoThreshold("Triangle dark");`

`//run("Threshold...");`

`setOption("BlackBackground", false);`

`run("Convert to Mask");`

`run("Erode");`

`run("Remove Outliers...", "radius=2.5 threshold=50 which=Bright");`

`run("Erode");`

`run("Remove Outliers...", "radius=5.5 threshold=50 which=Bright");`

419  run("Analyze   Particles...",   "size=315-40000   circularity=0.63-1.00   display   exclude

420   summarize add");

421 Mean surface areas were extracted and converted to diameters expressed in µm. For quality

422 assurance, when the number of detected cells was less than 200, the whole-slide image was visually

423 inspected to determine why the count was low. If there was an issue in cell detection, the threshold

424 was manually adjusted to improve segmentation and the image was reanalyzed. If the detection of

425 cells appeared accurate, the initial results were kept if the total number of cells detected was greater

426 than 100. If less than 100 cells were detected, this was considered insufficient, and the results were

427 omitted from subsequent analyses. A simplified schematic overview of the two methods is

428 presented in **Figure 3**.

Figure 3: Schematic overview of the manual **(a)** and the automated **(b)** adipocyte diameter measurements in image J. Descriptive steps with disadvantages and advantages of each method are provided.

4.4 Statistical analyses

Spearman correlations and paired t-tests were used for the initial validation with 16 adipose tissue samples with GraphPad Prism 10.0.3 (GraphPad Software, LLC, San Diego, Calif, USA).

465 For the final validation of the automated method using 377 adipose tissue samples from 190

466 patients, the strength of association between the mean adipocyte diameters from each method was

467 analyzed using paired t-tests and Pearson correlations. Disagreement between the mean adipocyte

468 diameters from each method was evaluated using Bland-Altman plots. Pearson correlations were

469 used to analyze the association between mean adipocyte diameters from each method with

470 different markers of cardiometabolic health (BMI, WC, total Chol, HDL-Chol, LDL-Chol, TG,

471 apoB, HOMA-IR, and VAI). Meng's Z-test for correlated correlation coefficients was used to

472 determine if the correlations differed significantly based on the method used. Statistical analysis

473 was performed using RStudio® 2024.04.2 Build 764 (Posit Software, PBC, Boston, Mass. USA)

474 with R version 4.3.3 and with GraphPad Prism® Version 10.3.1 (509) (GraphPad Software, LLC,

475 San Diego, Calif, USA).

Authors' contribution

Conceptualization: MFG, AR, and AT. Methodology: MFG and AR. Software: MFG. Validation: AR, IMP, MFG, and AT. Formal analysis and Investigation: AR, IMP, and MFG. Resources: AT, FH, LB, and MFG. Data Curation: IMP, GO, AR, and MFG. Supervision: AT. Writing - Original Draft: AR, MFG, and AT. Writing - Review & Editing: AR, MFG, IMP, GO, FH, LB, and AT. Visualization: AR and IMP. Supervision: AT. Project Administration: AT, MFG, and AT Funding acquisition: AT. All authors have read and approved of the final work.

Funding information and sponsorship

This study was funded by the Natural Sciences and Engineering Research Council of Canada, Grant/Award Numbers: 2017-05825. AR is a recipient of the Doctoral Training Scholarship from the *Fonds de Recherche du Québec (FRQ)*. IMP and GO are recipients of postdoctoral scholarships from the FRQ.

Disclosures

AT receives research funding from Johnson & Johnson, Medtronic, GI Windows and BioTwin for studies on obesity and bariatric surgery, and acted as a consultant for Bausch Health, Novo Nordisk and BioTwin.

Data availability statement

The participants of this study did not give written consent for their data to be shared publicly, so due to the sensitive nature of the research supporting data is not available.

5- References

1. Wharton S, Lau DCW, Vallis M, Sharma AM, Biertho L, Campbell-Scherer D, Adamo K, Alberga A, Bell R, Boulé N, Boyling E, Brown J, Calam B, Clarke C, Crowshoe L, Divalentino D, Forhan M, Freedhoff Y, Gagner M, Glazer S, Grand C, Green M, Hahn M, Hawa R, Henderson R, Hong D, Hung P, Janssen I, Jacklin K, Johnson-Stoklossa C, Kemp A, Kirk S, Kuk J, Langlois MF, Lear S, McInnes A, Macklin D, Naji L, Manjoo P, Morin MP, Nerenberg K, Patton I, Pedersen S, Pereira L, Piccinini-Vallis H, Poddar M, Poirier P, Prud'homme D, Salas XR, Rueda-Clausen C, Russell-Mayhew S, Shiau J, Sherifali D, Sievenpiper J, Sockalingam S, Taylor V, Toth E, Twells L, Tytus R, Walji S, Walker L, Wicklum S. Obesity in adults: a clinical practice guideline. CMAJ. 2020 Aug 4;192(31):E875–91.

2. Tchernof A, Després JP. Pathophysiology of human visceral obesity: an update. Physiol Rev. 2013 Jan;93(1):359–404.

3. Ye RZ, Richard G, Gévry N, Tchernof A, Carpentier AC. Fat Cell Size: Measurement Methods, Pathophysiological Origins, and Relationships With Metabolic Dysregulations. Endocr Rev. 2022 Jan 12;43(1):35–60.

4. Laforest S, Labrecque J, Michaud A, Cianflone K, Tchernof A. Adipocyte size as a determinant of metabolic disease and adipose tissue dysfunction. Crit Rev Clin Lab Sci. 2015;52(6):301–13.

5. Lessard J, Laforest S, Pelletier M, Leboeuf M, Blackburn L, Tchernof A. Low abdominal subcutaneous preadipocyte adipogenesis is associated with visceral obesity, visceral adipocyte hypertrophy, and a dysmetabolic state. Adipocyte. 2014 Jul 1;3(3):197–205.

6. Chen HC, Farese RV. Determination of adipocyte size by computer image analysis. J Lipid Res. 2002 Jun;43(6):986–9.

7. Okamoto Y, Higashiyama H, Inoue H, Kanematsu M, Kinoshita M, Asano S. Quantitative image analysis in adipose tissue using an automated image analysis system: differential effects of peroxisome proliferator-activated receptor-alpha and -gamma agonist on white and brown adipose tissue morphology in AKR obese and db/db diabetic mice. Pathol Int. 2007 Jun;57(6):369–77.

8. Galarraga M, Campión J, Muñoz-Barrutia A, Boqué N, Moreno H, Martínez JA, Milagro F, Ortiz-de-Solórzano C. Adiposoft: automated software for the analysis of white adipose tissue cellularity in histological sections. J Lipid Res. 2012 Dec;53(12):2791–6.

9. Yosofvand M, Liyanage S, Kalupahana NS, Scoggin S, Moustaid-Moussa N, Moussa H. AdipoGauge software for analysis of biological microscopic images. Adipocyte. 2020 Dec;9(1):360–73.

10. Laforest S, Pelletier M, Michaud A, Daris M, Descamps J, Soulet D, Jensen MD, Tchernof A. Histomorphometric analyses of human adipose tissues using intact, flash-frozen samples. Histochem Cell Biol. 2018 Mar;149(3):209–18.

11. Palomäki VA, Koivukangas V, Meriläinen S, Lehenkari P, Karttunen TJ. A Straightforward Method for Adipocyte Size and Count Analysis Using Open-source Software QuPath. Adipocyte. 2022 Dec;11(1):99–107.

12.   Maguire AS, Woodie LN, Judd RL, Martin DR, Greene MW, Graff EC. Whole-slide image
analysis outperforms micrograph acquisition for adipocyte size quantification. Adipocyte.
2020 Dec;9(1):567–75.

13.   Laforest S, Michaud A, Paris G, Pelletier M, Vidal H, Géloën A, Tchernof A. Comparative
analysis of three human adipocyte size measurement methods and their relevance for
cardiometabolic risk. Obes Silver Spring Md. 2017 Jan;25(1):122–31.

14.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods. 2012 Jul;9(7):671–5.

15.   Hu Y, Yu J, Cui X, Zhang Z, Li Q, Guo W, Zhao C, Chen X, Meng M, Li Y, Guo M, Qiu J,
Shen F, Wang D, Ma X, Xu L, Shen F, Gu X. Combination Usage of AdipoCount and
Image-Pro Plus/ImageJ Software for Quantification of Adipocyte Sizes. Front Endocrinol.
2021;12:642000.

16.   Osman OS, Selway JL, Kępczyńska MA, Stocker CJ, O'Dowd JF, Cawthorne MA, Arch
JR, Jassim S, Langlands K. A novel automated image analysis method for accurate
adipocyte quantification. Adipocyte. 2013 Jul 1;2(3):160–4.

17.   Parlee SD, Lentz SI, Mori H, MacDougald OA. Quantifying size and number of adipocytes
in adipose tissue. Methods Enzymol. 2014;537:93–122.

18.   Zhi X, Wang J, Lu P, Jia J, Shen HB, Ning G. AdipoCount: A New Software for Automatic
Adipocyte Counting. Front Physiol. 2018;9:85.

19.   Giavarina D. Understanding Bland Altman analysis. Biochem Medica. 2015 Jun
5;25(2):141.

20.   Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD,
McQuaid S, Gray RT, Murray LJ, Coleman HG, James JA, Salto-Tellez M, Hamilton PW.
QuPath: Open source software for digital pathology image analysis. Sci Rep. 2017 Dec
4;7(1):16878.

21.   Adiposoft [Internet]. ImageJ Wiki. [cited 2024 Oct 1]. Available from:
https://imagej.github.io/plugins/adiposoft

22.   Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis
model assessment: insulin resistance and beta-cell function from fasting plasma glucose and
insulin concentrations in man. Diabetologia. 1985 Jul;28(7):412–9.

23.   Amato MC, Giordano C, Galia M, Criscimanna A, Vitabile S, Midiri M, Galluzzo A,
AlkaMeSy Study Group. Visceral Adiposity Index: a reliable indicator of visceral fat
function associated with cardiometabolic risk. Diabetes Care. 2010 Apr;33(4):920–2.

a)

Manual Measurement of Cell Size

Method
Five photomicrographs of each H&E-stained slide are acquired in bright-field at 5x magnification.

The photomicrographs are imported into ImageJ.

Disadvantages
-Difficult to analyze low-quality images
-Cannot use whole slide images due to file size and time constraints

Method
Manual threshold is applied to segment counterstained adipocytes.

Images are manually processed to reduce background signal.

Disadvantages
- Residual background signal

Method
At least 100 cells are counted using the magic wand tool.

Surface areas are extracted from each counted adipocyte.

Disadvantages
-Time consuming
- Poor representation of whole tissue
-Measurement and observer bias

b)

Automated Measurement of Cell Size

Analyze Particles	×
Size (µm^2):	315-40000
	☐ Pixel units
Circularity:	0.63-1.00
Show:	Overlay ▼

☑ Display results ☐ Exclude on edges
☐ Clear results ☐ Include holes
☑ Summarize ☐ Record starts
☐ Add to Manager ☐ In situ Show

OK   Cancel   Help

Method
Whole slide images of the H&E-stained tissue sections acquired at 20x magnification.

Whole slide images are imported into ImageJ.

Advantages
-High-quality images of each section
-Automated acquisition of whole slide images

Method
Automatic thresholding is applied.

Automatic adjustment to reduce background signal.

Advantages
-Possibility of batch processing
-Reproducible results

Method
Automated analysis of particles with cut-offs for size (20-225µm) and circularity (0.63-1.00).

Advantages
-Rapid and high-capacity measurement of adipocytes