Article
Electroconvulsive therapy generates a postictal wave of spreading depolarization in mice and humans

Zachary P Rosenthal[1,*], Joseph B. Majeski[2], Ala Somarowthu[3], Davin K Quinn[4], Britta E. Lindquist[5], Mary E. Putt[6], Antoneta Karaj[6], Chris G Favilla[7], Wesley B. Baker[3], Golkoo Hosseini[8], Jenny P Rodriguez[8], Mario A Cristancho[8,9], Yvette I Sheline[7,8,9,10], C. William Shuttleworth[11], Christopher C. Abbott[4], Arjun G Yodh[2], Ethan M Goldberg[3,7,12]

1. Psychiatry Residency Physician-Scientist Research Track, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
2. Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA, USA
3. Division of Neurology, Department of Pediatrics, The Children's Hospital of Philadelphia, PA, USA
4. Department of Psychiatry, University of New Mexico School of Medicine, Albuquerque, NM, USA
5. Department of Neurology, University of California San Francisco School of Medicine, San Francisco, CA, USA
6. Department of Biostatistics, Epidemiology & Informatics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
7. Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
8. Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
9. Center for Neuromodulation in Depression and Stress, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
10. Department of Radiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
11. Department of Neurosciences, University of New Mexico School of Medicine, Albuquerque, NM, USA
12. Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA
* Corresponding author, Zachary.Rosenthal@pennmedicine.upenn.edu

Abstract

Electroconvulsive therapy (ECT) is a fast-acting, highly effective, and safe treatment for medication-resistant depression. Historically, the clinical benefits of ECT have been attributed to generating a controlled seizure; however, the underlying neurobiology is understudied and unresolved. Using optical neuroimaging of neural activity and hemodynamics in a mouse model of ECT, we demonstrated that a second brain event follows seizure: cortical spreading depolarization (CSD). We found that ECT pulse parameters and electrode configuration directly shaped the wave dynamics of seizure and subsequent CSD. To translate these findings to human patients, we used non-invasive diffuse optical monitoring of cerebral blood flow and oxygenation during routine ECT treatments. We observed that human brains reliably generate hyperemic waves after ECT seizure which are highly consistent with CSD. These results challenge a long-held assumption that seizure is the primary outcome of ECT and point to new opportunities for optimizing ECT stimulation parameters and treatment outcomes.

Introduction

Electroconvulsive therapy (ECT) is a life-saving intervention for medication-resistant depression. Treatment consists of electrical pulses delivered to the brain to elicit a controlled (~30-90 second) electrographic seizure while under general anesthesia and muscle relaxants to minimize physical movement. Nearly a century since it was discovered, ECT remains the most clinically effective treatment for severe depression[1]. A typical index course of six to twelve ECT treatments achieves rapid symptom improvement in 60-80% of patients, reducing suicide risk by 50% compared to matched controls who do not receive ECT[2,3]. ECT is also highly effective in mania, psychosis, Parkinson's disease, catatonia, and even status epilepticus – particularly for those with the most severe, medication-resistant symptoms. The procedure is safe and generally well-tolerated, including in pediatric, geriatric, and pregnant populations. Potential risks, such as cognitive slowing or memory impairment, are typically modest compared to improvement in neuropsychiatric symptoms, and resolve when treatment is discontinued. Unfortunately, the proven efficacy and safety of ECT have been overshadowed by stigma and negative portrayal in popular media. As a result, contemporary research on ECT is limited, and the treatment is underutilized[4]. Much could be learned from ECT and the basic physiology underlying its rapid-acting efficacy across a varied range of brain disorders, particularly when modern pharmacology has failed.

For the last eight decades, it has been assumed that seizure is necessary for ECT to elicit clinical improvement[3]. However, not all ECT-induced seizures are therapeutic, and conversely, electrical stimulation below the seizure threshold may also be an effective

treatment[5,6]. The role of seizure in the therapeutic mechanism of ECT is thus uncertain.

An important clue to this mechanism may be that ECT induces lasting inhibitory plasticity

in brain activity, dampening cortical response to stimulation and reliably reducing

propensity for future seizures (i.e., raising the seizure threshold)[7-11]. It has been

hypothesized that ECT may elicit inhibitory plasticity through the process of seizure

suppression[12], but few studies have explored the cellular, circuit, or network level

mechanisms of brain inhibition after ECT.

During ECT, brain activity is typically monitored with a minimal scalp

electroencephalography (EEG) montage consisting of two leads on the forehead to

measure seizure quality, duration and post-ictal suppression in each hemisphere.

However, quantitative seizure metrics from EEG monitoring have shown limited predictive

value for ECT clinical outcomes[5-7,13-16]. A great need exists for more robust brain activity

biomarkers that can predict treatment efficacy, detect side effects, or guide selection of

stimulation parameters. Because ECT electrical pulses saturate the scalp EEG signal, no

study has ever measured brain activity during pulse delivery in human patients or animal

models. Computational models have helped predict the brain electric field evoked by a

single electrical pulse[5,17-19], but there are likely opportunities to further optimize

stimulation parameters and improve clinical outcomes using empirical measurements of

brain activity during treatment. For example, a seminal randomized controlled trial

showed that the choice of two electrode spatial configurations (right unilateral vs.

bitemporal), as well as two pulse durations (0.3 vs. 0.5 ms), significantly modulates the

clinical efficacy and side effects of ECT[20,21]. The remaining stimulation parameter space

is vast, mostly untested, and with unknown effects on brain activity and clinical

outcomes[20]. This includes infinite combinations of pulse properties (duration, frequency,

total count, current, polarity, waveform) and electrode spatial configurations. Only three

electrode placements have been tested clinically – right unilateral, bitemporal, and

bifrontal. There are likely alternative stimulation parameters that would provide superior

symptom reduction and lower risk of side effects compared to the current standard-of-

care configurations and titration algorithms.

To bridge these knowledge gaps, we employed optical neuroimaging to record brain

activity during ECT in both rodent models and human patients. Our data demonstrate that

a second brain event follows ECT-induced seizure: cortical spreading depolarization

(CSD). In mice, we show that clinically relevant choices of electrode placements and

pulse parameters directly modulate the spatial and temporal properties of both seizure

and subsequent CSD. We then translate these findings from mice to humans. Using non-

invasive diffuse optical monitoring of brain tissue oxygen saturation and blood flow in

patients receiving ECT, we find evidence for post-ictal hyperemic waves that are highly

consistent with CSD. These discoveries have important implications for ECT's

mechanism of action and for optimizing stimulation to target specific brain outcomes.

Methods

Mouse model

All procedures described below were approved by the Institutional Animal Care and Use

Committee at the Children's Hospital of Philadelphia in compliance with AAALAC

guidelines. Mice were raised in standard cages in a double barrier mouse facility with a

12-hr/12-hr light/dark cycle and *ad libitum* access to food and water. All experiments used

n=10 eight-week-old mice (male and female) hemizygous for *Thy1-jRGECO1a* (JAX

030525) on a C57BL/6J background, to enable optical imaging of the fluorescent calcium

sensor protein jRGECO1a. Pups were genotyped by PCR prior to experiments to confirm

the presence of the *Thy1-jRGECO1a* transgene, using the forward primer 5'-

ACAGAATCCAAGTCGGAACTC-3'   and   reverse   primer   5'-

CCTATAGCTCTGACTGCGTGAC-3'.

Cranial window and electrode implantation

Mice were treated preoperatively with subcutaneous buprenorphine-SR (1.0 mg/kg),

meloxicam (5.0 mg/kg) and cefazolin (500 mg/kg). Mice were anesthetized with isoflurane

(3% induction, 1.5% maintenance). Body temperature was maintained via heating pad.

The scalp was shaved, sterilized with alcohol and betadine, incised at midline, and

retracted to expose the dorsal skull. Five brass electrode pins with 1.78mm diameter

flanges (DigiKey 4443-0-00-15-00-00-03-0) were attached to the intact skull surface using

a thin layer of silver conductive epoxy (MG Chemicals 8331D). Electrodes were

stereotactically centered relative to bregma (see **Fig. 1a**; #1, frontal: X = 0.00 mm, Y =

4.09 mm; #2 and #3, temporo-parietal: X = ±3.40 mm, Y = 0.47 mm; #4 and #5, occipital:

X = ±4.69 mm, Y = -4.89 mm). A custom steel headbar was attached to the posterior skull

and a cranial window was formed with optically clear dental cement (C&B-Metabond,

Parkell Inc., Edgewood, NY). The window was rendered transparent and hardened with

clear UV-cure gel nail polish. Animals were allowed to recover from surgery for at least

one week prior to imaging.

Widefield imaging system

Widefield imaging of neuronal dynamics (jRGECO1a fluorescence) and cerebral

hemodynamics (optical intrinsic signal, OIS) was performed using a modified version of

a previous method[22], on a Leica M205FA stereoscope fitted with a 0.63× objective to

capture the dorsal convexity of the cortex (a ~1 cm^2 area, binned to 128×128 pixels, 0.078

mm pixel-width). System optical spectra are depicted in **Extended Data Fig. 1**. White

light LED illumination (X-Cite XYLIS) was bandpass filtered to green (525/50 nm,

Chroma) and directed through the objective at the cortex. Light reflected and emitted from

the brain was then separated into two channels using a 560 nm dichroic and image-

splitting optics (Hamamatsu W-VIEW GEMINI). The OIS channel was filtered with a

512/25 nm bandpass and attenuated with a 5% transmission neutral density filter

(Chroma), optimized to capture light reflectance at the isosbestic point of hemoglobin to

approximate total hemoglobin concentration. The jRGECO1a red fluorescence channel

was filtered with a 630/92 nm bandpass filter. Both channels were detected side-by-side

and spatially co-registered on a single CMOS camera (Hamamatsu ORCA-Flash4.0 V3).

An exposure time of 50 ms was used, achieving a sampling rate of 15.7 Hz.

Combined imaging and electroconvulsive therapy in mice

Mice were monitored with head-fixed widefield imaging while under dexmedetomidine

anesthesia (0.5 mg/kg, IP), which reliably achieved a plane of sedation in ~10 minutes

with loss of response to toe-pinch and emergence of ~3 Hz anteroposterior global slow

waves that are characteristic of anesthesia[23,24]. ECT pulses were delivered to pairs of

cranial electrodes using constant current pulse trains (described below) with an isolated

pulse stimulator (A-M Systems 4100). At the end of each recording session, anesthesia

was reversed with atipamezole (0.5 mg/kg). Similar to prior reports, seizures were noted

to elicit ~1-60 seconds of high-amplitude, aperiodic neuronal fluorescence activity globally

across the cortex, as well as tonic-clonic limb and tail movements. Several pilot titrations

were also conducted using etomidate as well as ketamine/xylazine anesthesia, which

likewise demonstrated CSD after ECT seizure (data not shown); dexmedetomidine was

ultimately selected because of its hemodynamic safety[25], reversibility, and favorable

pharmacodynamic profile as an alpha agonist, avoiding acting directly on GABAergic or

glutamatergic currents.

To broadly survey the stimulation parameter space, n=38 ECT stimulus titrations were

performed, each testing a different combination of electrode spatial configuration (5

electrodes, 10 possible pairs), frequency (5, 10, 25, 50, 100 Hz) and pulse counts (25,

50, 100 pulses); as shown in **Extended Data Table 1**. All stimulation trials used constant-

current, bipolar square waves of 0.5 ms pulse width, similar to routine brief pulse ECT in

humans. Within each titration, stimulation was delivered first at 1 mA current and then

successively up-titrated to 2 mA, 5 mA, 10 mA, and 25 mA, akin to the individualized low

amplitude seizure therapy (ILAST) strategy of ECT[26]. At each current step, brain activity

was monitored at baseline for 5 seconds, during stimulation, and then for at least 90s, or

until the sustained return of baseline anteroposterior slow waves from anesthesia, before

being restimulated at the next highest current step. At the current threshold when seizure

elicited CSD, recordings were continued for 10 minutes, and the titration was terminated.

Titration sessions within individual mice were spaced at least 1 week apart to facilitate

washout of acute effects of stimulation.

To control for effects of resampling the same individuals across titrations with different

conditions, mixed effects modeling adjusted for mouse ID. Stimulation conditions were

partially randomized and balanced across n=10 mice such that each mouse was treated

with multiple frequencies and electrode configurations in varying session order. For any

given electrode spatial configuration, multiple frequencies were tested; for each

frequency, a mixture of unilateral and bilateral electrode configurations were tested. In

n=4 titrations, animals were restimulated one more time to elicit a second CSD – these

four secondary CSDs exhibited similar intrinsic amplitude and duration to initial CSDs,

and were thus pooled into **Figs. 2a**, **2g**, and **2h** (n=42). Secondary CSDs were excluded

from analysis in **Figs. 2f** and **3**. One titration was excluded from analysis because of

inadequate head-fixation leading to motion artifact that precluded accurate quantification;

it was however visually scored as a right-sided CSD and pooled into **Fig. 2f** for

completeness (n=39).

Mouse widefield optical imaging signal processing

Data were processed using an open-source widefield optical imaging toolkit[27]. Imaging

data were converted into two-channel tiff files. A binary brain mask was manually drawn

in MATLAB for each recording; all subsequent analyses were performed on pixels labeled

as brain. Image sequences from each mouse (as well as the brain mask for each mouse)

were affine-transformed to Paxinos atlas space using the positions of bregma and

lambda[28]. A standardized mask of the stimulation electrodes was approximated using

Paxinos coordinates (see seizure to electrode distance permutation analysis below).

Right and left hemispheric divisions were segmented using a straight line drawn through

midline of each image. A dark image with no illumination was subtracted from all frames,

and then data were spatially smoothed with a Gaussian filter. The jRGECO1a calcium-

sensor fluorescence signal (%dF/F) and total hemoglobin OIS signal (%dA/A) were

calculated at each pixel by subtracting and dividing the 20^{th} percentile value from the first

4 seconds of baseline (pre-stimulation) raw signal from each recording. The total

hemoglobin OIS signal was multiplied by -1 (absorbance) so that positive sign changes

indicate increased blood volume/total hemoglobin concentration. Red-shifted

fluorophores have significantly reduced hemoglobin absorption artifact compared to

green fluorophores[22]. Given that fluorescence %dF/F event peaks were more than an

order of magnitude larger than hemoglobin %dA/A changes, we opted not to regress the

hemodynamic signal out of the fluorescence signal.

Event detection and quantification in neuronal fluorescence data

Optical detection of cortical hemodynamics and calcium dynamics has been extensively

cross-validated as a surrogate for routine electrophysiology for detecting CSD, offering

rich spatial information about traveling waves.[29-32]. The temporal bounds of seizure and

CSD events were identified by taking the derivative of the root mean square of the

neuronal fluorescence signal at each pixel. We then used the MATLAB function

'findchangepts' to find abrupt signal changes at the start and end of seizure and CSD

events. We used an event detection threshold of %dF/F signal rising greater than 20% of

baseline. Note, typical widefield %dF/F (fluorescence change) values are +/- 5-10%

during routine physiological fluctuations in a mouse. In contrast, seizure and CSD are

such large events that %dF/F fluorescence typically rises by 50-200%. The event

detection threshold of 20% was determined through trial and error to accurately identify

the bounds of seizure and CSD. This approach was cross-validated against the current

gold standard for electrographic CSD detection - visual scoring by a trained clinician[33].

Recordings (and pixels therein) not crossing this threshold were excluded from analysis.

For each event (seizure or CSD) in each pixel, duration was calculated as the difference

between start and end times, and amplitude was calculated as the peak %dF/F value

between the start and end time. Then, average event amplitude was computed within

right hemisphere, left hemisphere, and total brain space by averaging across pixels.

Overall event durations were computed by comparing the event start times in the first and

last pixel with right hemisphere, left hemisphere, and total brain space. For CSDs, the

first pixel within the brain mask to have an event start time was used as a global reference

point for computing time to peak at other pixels for spatial maps of CSD trajectory. For

CSDs, the percentage of pixels from each hemisphere recruited into the event was

calculated. Average within-hemisphere CSD areal propagation speed was computed by

dividing the area recruited into the CSD by the event duration within each hemisphere.

Average linear propagation speed was computed by identifying the minimum and

maximum points within each hemisphere of the CSD trajectory maps and drawing a line

between them. This line profile of the spatial gradient was fit with a linear regression from

which the slope was computed and converted into mm/min. Of note, clinical seizure

duration in human ECT does not consistently include the 0.5-8 second stimulation period,

because EEG activity cannot be measured during electrical pulse delivery. Our mouse

optical neuroimaging paradigm allows us to observe that high-amplitude cortical

discharges occur during stimulus delivery; we thus included this activity as part of the

seizure.

Both seizure and CSD are well known to trigger a period of suppressed brain activity. To

calculate post-event suppression time, a short time Fourier transform (STFT) power

spectrum was determined for each pixel and for average right/left hemisphere

fluorescence. Hemispheric suppression endpoints were detected using the findchangepts

function for the fluorescence signal in each hemisphere. The suppression end point for

each pixel was computed by finding the inflection point of up-trending 1-3Hz power

(associated with return of slow waves from anesthesia) that is nearest to the respective

global suppression endpoint. Then, the suppression duration was computed by taking the

difference between the suppression end time and the event (seizure or SD) end time.

Suppression times were then averaged within each hemisphere.

The maximum intensity seizure pixel was identified for all end-titration (i.e., CSD-

generating) seizures (n=42) by identifying the median value within the top 99% of pixel

fluorescence amplitude in seizure maps. We then calculated the shortest distance

between the seizure peak pixel and the nearest stimulation electrode (**Fig. 2i**), as well as

the shortest distance between the seizure peak pixel and the CSD initiation reference

pixel (**Fig. 2j**). Electrode-to-seizure and seizure-to-CSD distances were averaged across

all 42 events. To assess the significance of these spatial relationships, a shuffle analysis

was conducted by randomly permuting the data and recalculating the average distances

for 10,000 iterations. The distribution of shuffled distances was compared to the actual

distances, and statistically verified by t-test with Bonferroni correction for multiple

comparisons.

Statistical comparisons of seizure and CSD phenotypes

For analyses below summarizing all recorded seizure and CSD events, distribution

normality testing was performed using the D'Agostino & Pearson test to determine use of

parametric or non-parametric testing. We used non-parametric Kruskal-Wallis ANOVA

with Dunn's multiple comparison test for comparison of seizure laterality index between

stimulation types (**Fig. 2c**), as well as comparison of seizure duration (**Extended Data**

Fig. 3a) and amplitude (**Extended Data Fig. 3b**) across CSD outcomes. Post-event

suppression times (**Extended Data Fig. 3c,d**) were compared using the parametric

Brown-Forsythe ANOVA with Dunnett's multiple comparison test. The relative

contribution of the three types of stimulation configurations (right, left, bilateral) for right,

left and bilateral CSDs were compared using Fisher's exact test (**Fig. 2f**). Average CSD

fluorescence amplitude (**Fig. 2g**) and duration (**Fig. 2h**) were compared using the non-

parametric Mann-Whitney test.

Mixed effects model of stimulation parameter titration

For analyses of the effect of titrating pulse parameters (frequency, current level) on

seizure properties (overall duration, average mean fluorescence amplitude, **Fig. 3**), we

excluded cases that did not cross the seizure threshold in any pixels, thus excluding all

evoked responses at 1 mA. A mixed effects model was fit using the outcome of either

seizure duration or average peak fluorescence amplitude (%dF/F), transformed on a log

scale to achieve approximate normality. Frequency and current level were the

independent variables. A model including an interaction between current level and

frequency was fit and the interaction evaluated using an F-test with a Satterthwaite

correction for the degrees of freedom. If the interaction was significant ($p < 0.05$) the

model was refit for each current level and the effect of increasing frequency evaluated.

Specific values of frequency were replaced with integer values (levels 1,2,3,4,5

correspond to 5, 10, 25, 50, 100 Hz) to succinctly evaluate trends in the outcome as a

function of frequency. If the interaction was not significant, the procedure was repeated

and the main effects of frequency and amplitude were evaluated marginally, followed by

Wald tests of specific contrasts. Models of current and frequency effects included

electrode placement, animal ID, and pulse counts as covariates to adjust for imbalances

across conditions. Electrode placement and pulse count were tested in separate

sensitivity analyses to preclude convergence problems with the mixed effects models due

to the small number of animals. Similarly, frequency was included as an ordered variable

in the model with pulse count.

Time-to-event Model of CSD

To assess how modulating pulse parameters impacts the probability of CSD after seizure,

we fit a Cox model substituting current level for time and including frequency as the

independent variable and stratification by animal ID. Current is a surrogate for time in this

model because an animal receives increasing current levels at each frequency until the

event, CSD, is observed. Once a CSD was observed, no further increases in current were

assessed. Frequency was considered both as an ordered variable, to assess the overall

trend with increasing frequency, and as a categorical variable to describe the results at

specific frequencies. Electrode placement is included in the models reported here and in

sensitivity analyses substituting pulse count for electrode frequency. Hypothesis tests of

variables with individual terms are based on Wald tests, and for those with multiple

categories (electrode placement, frequency) on likelihood ratio tests (LRT). Hypothesis

tests are two-sided with a Type I error rate of 0.05, uncorrected for multiple comparisons.

Diffuse Optics and Vitals Monitoring in ECT Patients

All procedures described below were approved by the University of Pennsylvania

Institutional Review Board for the observational cohort study, SWEET COMBO: Studying

Waves Evoked by Electroconvulsive Therapy with Combined Optical Monitoring of Blood

flow and Oxygenation. Patients included in the study (n=18) were recruited from the pool

of patients already being treated with ECT at Pennsylvania Hospital (Penn Medicine

Health System) between February and July 2024. All participants provided written

informed consent to participate. Of note, participation in this cross-sectional study had no

impact on which patients were treated with ECT, nor in what manner. All patients were

treated with standard of care protocols for Pennsylvania Hospital using a Σigma device

(SigmaStim). Study enrollees include 9 women and 9 men, ages ranging 23-82 (median

age 60). Enrollees self-identified as white (n=14), black (n=3), Asian (n=1) and Hispanic

(n=1).

During ECT treatment, in addition to routine clinical monitoring (EKG, EEG, vitals), two

diffuse optical monitoring probes were placed on the forehead surface in the regions

overlying right (F4) and left (F3) prefrontal cortex. The probes and optical system provide

simultaneous measurement of blood flow index using diffuse correlation spectroscopy[34,35]

(DCS) and tissue oxygen saturation using frequency-domain diffuse optical

spectroscopy[35,36] (FD-DOS) from scalp, skull, and cortex underlying the probes (see

Supplemental Methods on instrumentation)[35]. To improve accuracy of the recovered

tissue oxygen saturation, FD-DOS combined data from four separations (1.5, 2, 2.5, and

3 cm) on the tissue surface to extract a single datapoint. For DCS flow monitoring, each

optical probe utilized a long and short source-detector pair that were fit independently.

The light from the short separation pair (1 cm) penetrates scalp and skull, while light from

the long separation pair (2.5 cm) probes deeper to the cortex. The optical property data

collected concurrently with FD-DOS was input to the DCS fitting to account for changes

in absorption and scattering that may otherwise confound the recovered blood flow index.

Briefly, FD-DOS and DCS data were fit using the semi-infinite, homogenous solutions to

the frequency domain-photon diffusion equation and the correlation diffusion equation,

respectively. The interested reader is directed to reference[34] for greater detail on the

theoretical models used. In the main text, we only show blood flow obtained from the long

separation pair that is sensitive to cortex (example short separation data is shown in

Extended Data Fig. 5 and **Supplemental Discussion**).

Diffuse optical recordings started from a pre-stimulation baseline period and lasted for

several minutes after treatment; the treatment period had variable duration depending on

the procedure room schedule, effects of anesthesia, and patient response. Routine

intermittent vitals monitoring (collected at 2-minute intervals) was supplemented with

continuous vitals monitoring (20 second intervals) using a noninvasive finger monitor for

cardiac output (Acumen IQ cuff; HemoSphere monitor, Edwards)[37]. A total of n=27

recordings were performed on n=17 patients **(Extended Data Table 2** and **3)**. Fifteen of

these recordings were excluded because the recording signal did not meet quality control

standards (see **Supplemental Discussion/Methods**). The remaining n=12 recordings

are presented in **Fig. 4** and **Extended Data Fig. 4**. Based on prior measurements in

cross-validated CSDs[38-42], we defined CSD as a post-ictal wave of at least 2-minute

duration before return to 100% of baseline, and with peak >200% of baseline blood flow

and/or with >±5% fluctuation around baseline brain oxygen saturation. For recordings

where bilateral sensor data was available, if these criteria were met in both hemispheres,

the CSDs were classified as bilateral; if criteria were met in only one hemisphere, CSD

was classified as unilateral.

Results

ECT seizure is followed by cortical spreading depolarization in mice

We created a new mouse ECT model to facilitate optical neuroimaging of large-scale

brain activity during ECT stimulation. We used transgenic *Thy1-jRGECO1a* mice which

express the red fluorescent calcium sensor jRGECO1a in excitatory neurons[43]. Mice were

implanted with a transparent polymer window overlying the intact skull to enable optical

access to activity-dependent dynamics across the cortical surface. We measured

neuronal activity with jRGECO1a fluorescence as well as hemodynamic activity with

optical intrinsic signal (OIS) imaging, using green light illumination and a 1-photon

widefield fluorescence mesoscope (optical set-up shown in **Fig. 1b,** spectra shown in

Extended Data Fig. 1). To facilitate concurrent electrical stimulation within the imaging

field, each mouse cranial window was implanted with five brass electrodes stereotactically

affixed to the skull surface overlying frontal, temporo-parietal, and occipital cortices (**Fig.**

1a). Brain activity was recorded with widefield imaging under dexmedetomidine

anesthesia during 153 ECT stimulation blocks. To broadly survey the stimulation

parameter space (see Methods and **Extended Data Table 1**), we controlled the

stimulation pulse parameters using an isolated pulse stimulator, with spatial configuration

varied by changing which pair of electrodes were stimulated. For each set of tested

conditions, stimulation was titrated by delivering successive rounds of pulses with

increasing current (1, 2, 5, 10, 25 mA, **Fig. 1c**) followed by a monitoring period before

restimulating.

383
Fig. 1. Optical neuroimaging in a mouse model of ECT demonstrates a current threshold
where seizure is followed by cortical spreading depolarization.
a, Schematic of mouse cranial window with optical access to the dorsal cortex. Within each
window, five brass electrodes were attached to the intact skull overlying frontal (#1), temporo-
parietal (#2, 3), and occipital (#4, 5) cortices.
b, Green-filtered illumination (525/50 nm) was projected through the microscope objective onto
the brain; reflected and emitted light from the brain was separated by a 560 nm dichroic mirror
into two spatially co-registered channels that were bandpass filtered for isosbestic point
hemoglobin reflectance of green illumination (512/25 nm) or jRGECO1a red fluorescence
emission (630/92 nm). Electrical stimulation was delivered via pairs of brass electrode pins using
an isolated pulse stimulator. See **Extended Data Fig. 1** and **Supplemental Methods**.
c, Example video frames (from **Extended Data Video 1**) of neuronal activity (jRGECO1a relative
fluorescence %dF/F) in an individual mouse being stimulated with 5 Hz pulses to electrodes 1
and 4. Stimulation was titrated in successive rounds with increasing current (**c'**, note rainbow
color map corresponds to current titration); in each round we monitored brain activity for 5 seconds

of baseline, during stimulation, and then for 90 seconds (or the resumption of baseline slow wave
dynamics.). Data from each round of stimulation in the current titration are depicted in still-frame
maps of cortical activity in (**c**) and time series averaged within a right hemisphere ROI (**c"**).
402
Representative neuronal calcium dynamics during ECT titration are shown in **Fig. 1**. Low

amplitude current steps (1 and 2 mA) elicited a small <15% dF/F evoked response and

then an immediate return of slow wave activity characteristic of anesthesia (**Fig. 1c**). At

higher current (5, 10 mA), ECT elicited seizure activity (high-amplitude, irregular

discharges generalized across the cortex), followed by 5-10 seconds of post-ictal

suppression, and then return of slow wave activity. At 25 mA, seizure was followed by a

qualitatively distinct electrical event: cortical spreading depolarization (CSD, see

Extended Data Video 1, depicted as still frames in **Fig. 1c** and time series in **Fig. 1c"**).

CSD is a slow-traveling, high-amplitude wave of electrochemical depolarization that can

be detected with high spatiotemporal fidelity using all-optical detection of neural dynamics

or hemodynamics[29-32]. Post-ictal CSD was observed as a slowly propagating, high-

amplitude wavefront of neuronal calcium elevation that concentrically expanded across

the whole brain over the course of ~160 seconds, followed by a prolonged period of

suppressed neuronal calcium dynamics. All 38/38 ECT titration sessions reached the

threshold to trigger post-ictal CSD. Direct current electrocorticography (DC-ECoG) in a

different rodent ECT model likewise demonstrates post-ictal CSD in a current-thresholded

manner (see **Extended Data Fig. 2, Supplemental Discussion**).

Note, despite their magnitude, CSDs are virtually invisible on routine alternating current

(AC)-amplified EEG, as they are physically obscured by volume conduction and spatial

blurring through the scalp and skull, and are digitally hidden with high-pass filtering (>0.5

Hz)[33,44,45]. Notably, subjecting mouse brain fluorescence data to the same >0.5 Hz high-

pass filtering rendered CSD waves invisible on widefield imaging, and at point

measurements throughout the brain appeared as a flat line that is otherwise

indistinguishable from post-ictal suppression (see **Extended Data Video 1**).

Electrode configuration shapes the spatial topography of seizure and CSD

We then considered whether systematically varying electrode configuration would impact

the spatial topography of both seizure and post-ictal CSD waves. In our mouse model

with five electrodes, ten electrode pairs were possible for left unilateral (L, green), right

unilateral (R, magenta), or bilateral (BL, black) stimulation. Intuitively, we observed that

electrode configuration shaped the spatial topography of both the evoked seizure (**Fig.**

2a, mapping seizure fluorescence amplitude at each pixel) as well as the trajectory of the

subsequent CSD traveling wave (**Fig. 2b**, mapping time-to-peak of CSD at each pixel).

Bilateral electrode placement produced relatively symmetric seizures, while R or L

unilateral electrode placement recruited asymmetric seizure with higher amplitude in the

stimulated hemisphere than the contralateral hemisphere (**Fig. 2c,d,** L vs R $p^{****} <$

0.0001, L vs BL $p^{****} < 0.0001$, R vs BL $p^{**} = 0.0026$).

442
Fig. 2. Electrode configuration shapes the spatial topography of seizure and CSD.
a, Group average maps of seizure amplitude at each pixel (%dF/F). Each panel represents one
of 10 possible pairs of 5 electrodes, averaged across multiple stimulation trials (electrode pair and
number of trials in each average below, from n=42 end-titration seizures in n=10 mice).
Fluorescence colormap is normalized to each plot's min and max; absolute values of fluorescence
are shown in **c** and **d**. Stimulated electrode pairs are depicted as colored dots. Note seizures
tended to localize near stimulation electrodes.
b, Example individual map of post-ictal CSD trajectory for each electrode configuration, depicting
time-to-CSD-peak at each pixel. Note that some CSDs were recruited bilaterally to the entirety of
both hemispheres, while others were recruited unilaterally to one hemisphere. Bilateral CSDs
typically had some degree of temporal lag between the two hemispheres. Colormaps are
normalized to the earliest and latest pixel, with absolute values of overall event durations shown
in h.
c, Asymmetry index for all detected seizures across all titrations (see Methods) evoked by right
(R, n=25), left (L, n=34) and bilateral (BL, n=38) electrode configurations. Asymmetry index was
calculated as the ratio of the left:right hemisphere maximum seizure amplitude averaged across
pixels within each hemisphere. Kruskal-Wallis test with Dunn's multiple comparison correction L
vs. R (p**** < 0.0001), L vs BL (p****< 0.0001), R vs BL (p**=0.0026); Kruskal-Wallis statistic
61.83.
d, Seizure amplitude (fluorescence, (%dF/F) values averaged within the right (x-axis) and left (y-
axis) hemispheres for each event. A range of frequency and current conditions were tested (see
Fig. 3), producing a wide range of fluorescence intensities. Electrode configuration for each
seizure is color coded (L green, R magenta, BL Black).

e, Percentage of pixels within the left hemisphere and right hemisphere where CSD was detected,
from n=42 recorded events. Icons depict right-sided, left-sided, and bilateral CSD event types.
f, Distribution of n=39 CSD outcomes resulting from right (magenta), left (green) or bilateral
(black) stimulation. Fisher's exact test with null hypothesis that the distributions of electrode
configurations are equal in all three CSD outcomes (p < 0.0001 for 3x3 interaction).
g, Average amplitude (%dF/F) for bilateral and unilateral CSDs (selecting whichever hemisphere
had the largest activation). CSD amplitude was similar whether it occurred in one hemisphere or
both, via two-tailed Mann-Whitney test, p=0.3306 (U=171, n=42).
h, Overall duration of bilateral CSDs was slightly longer than that of unilateral CSDs, two-tailed
Mann-Whitney test p**=0.0011 (U= 94, n=42). This was due to asymmetry in the start times for
each hemisphere, illustrated in bilateral CSDs shown in b.
i, Average distance (millimeters) of seizure peak pixel to nearest stimulation electrode (n=42 end-
titration seizures), compared to 10,000 permutations of shuffled data, using a two-tailed t-test with
Bonferroni corrected p****< 0.0001.
j, Average distance (millimeters) of seizure peak pixel to CSD initiation pixel. Same statistical
approach as i.
482
In contrast to preceding seizures, post-ictal CSDs occurred in an all-or-none fashion

anatomically, recruiting either ~0% or ~100% of cortical pixels in a given hemisphere (**Fig.**

2b,e). Trials that evoked bilateral CSD waves were disproportionately triggered by

bilateral stimulation, while trials that evoked unilateral CSD waves were disproportionately

triggered by same-side unilateral electrodes (**Fig. 2f**, p < 0.0001). Bilateral and unilateral

CSDs tended to achieve similar fluorescence amplitude (**Fig. 2g,** p= 0.331), but bilateral

CSD events tended to be longer in duration (**Fig. 2h,** p = 0.0011), primarily because of

cases of asynchronous onset of multifocal CSD between hemispheres (e.g., **Fig. 2b**

frontotemporal configuration 1,2 in panel 1, where CSD in L hemisphere peaked ~1

minute before the R hemisphere). CSD propagation was noted to originate variably from

both singleton and multiple foci, with all foci inevitably expanding to invade the entire

hemisphere in which they started, but never traveling across midline. On average within

any given hemisphere, we observed that CSD wavefronts spread at 4.1 mm/min (±

standard deviation of 0.9 mm/min), consistent with prior electrophysiology. CSDs

expanded in 2D at an average rate of 13.2 mm^2/minute (± standard deviation of 3.6

mm^2/min). Seizure, while always generalized bilaterally throughout cortex, tended to

exhibit highest amplitude at a short distance from the stimulation electrodes (**Fig. 2a,i,**

p**** < 0.0001), consistent with prior predictions from brain electric field models for

various stimulation geometries[5]. Likewise, CSD initial foci tended to occur in close spatial

proximity to pixels exhibiting peak fluorescence values during the preceding seizure (**Fig.**

2b,j, p**** < 0.0001).

Pulse parameters modulate seizure and post-ictal CSD

Next, we examined titration data to test how pulse parameters, current (mA) and

frequency (Hz), influence seizure and CSD. For each treatment session, individual mice

were assigned to a pulse frequency and spatial configuration, and then serially stimulated

with increasing current steps, monitoring the evoked seizures at each step with optical

imaging, and ending at the current threshold where seizure led to CSD. For each seizure,

we measured the maximum amplitude (peak fluorescence intensity averaged across all

brain pixels, **Fig. 3a**) and overall event duration across the brain (**Fig. 3b**). Current

titrations in individual mice at a fixed frequency and electrode configuration are

represented as individual lines in **Fig. 3a,b**, with titration curves ending at the current

threshold where the evoked seizure triggered a CSD. Some combinations of frequency

and current were not tested (e.g., the combination of 100 Hz stimulation at 25 mA was

not reached in any titration because all mice experienced CSD at lower current steps).

Trials that did not meet our amplitude threshold for defining a seizure (see Methods) were

excluded from statistical analysis. To assess how pulse current and frequency modify

brain outcomes, we fit a mixed effects model (see Methods), using a random term to

account for repeated measures from each animal and fixed effects for the assigned

electrode configuration and pulse count.

524
Fig. 3. Pulse current and frequency modulate evoked seizure amplitude and duration, as
well as subsequent cortical spreading depolarization. Each line in **a** and **b** represents an
individual mouse being serially stimulated at a fixed pulse frequency (5, 10, 25, 50, or 100 Hz,
yellow-orange-red-black color coding). Mice were first stimulated at 1 mA current, seizure was
monitored until return to baseline, and then mice were restimulated in increasing in steps to 2, 5,
10, and 25 mA (rainbow color coding) until the evoked seizure was sufficient to trigger CSD.
Individual lines for each mouse titration in **a** and **b** thus end at the current step that elicited CSD.
The x-axis in **a-c** corresponds to the current step in the titration, while the y-axes measure three
different brain event outcomes.
a, Seizure amplitude (y-axis, fluorescence intensity %dF/F) averaged across the whole cortex as
a function of current step (x-axis, mA) for each mouse titration (individual lines) at one of five
frequencies (yellow-orange-red-black color coding). Data analyzed using mixed effects modeling
to account for repeated trials on individual animals (see Methods). n is the number of individual
mouse titration trials tested at each frequency.
b, Same seizure titrations as in a) but with y-axis measuring seizure duration (seconds). Note
inversion of the order of frequency groups relative to a.
c, Cumulative probability of experiencing CSD after seizure at each current step within each
frequency condition. Each line represents a group summary for the fixed frequency conditions 5,
10, 25, 50 and 100 Hz. See main text for results of statistical analysis.
544
For every mouse titration, across all tested frequencies, seizure amplitude went up with

increasing current (note positive slope to each titration line shown in **Fig. 3a**). The effect

of current differed by frequency (p< 0.001 F-test, with LRT interaction vs. main effects

model). Increasing pulse frequency significantly increased the amplitude of the resultant

seizure, producing steeper slopes in titrations for higher frequency conditions of 50 and

100 Hz shown in **Fig. 3a**. Within individual current levels, frequency did not significantly

modulate seizure amplitude at low current steps (2 mA, p=0.63), but each level increase

in pulse frequency increased seizure amplitude by a factor of 1.2 fold, significantly so at

5 mA (p< 0.001) and 10 mA (p=0.001), but not at 25 mA (p=0.11), see **Fig. 3a**. Electrode

placement did not significantly influence seizure amplitude, either in the interaction model

(p=0.70) or the individual models (p>0.05). In sensitivity analyses, higher pulse count did

not significantly contribute to current and frequency effects (p=0.45 LRT). Higher

frequency pulses thus elicit higher amplitude seizures independent of other variables.

In the same seizure titrations, increasing current led to increased seizure duration (**Fig.**

3b, p< 0.001), in a manner that did not vary significantly by pulse frequency condition

(p=0.39). Compared to seizures at the 2 mA current step, seizure duration increased 2.8-

fold at 5 mA (p< 0.001), 3.5-fold at 10 mA (p< 0.001) and 2.8-fold at 25 mA (p=0.005)

when controlling for individual frequency level and electrode placement. Somewhat

surprisingly, though increasing frequency led to larger seizure amplitude, seizures

became briefer. Compared to 5 Hz stimulation, seizure duration decreased by a factor of

0.6 for 10 Hz (p=0.08), by 0.4 at 10 Hz (p=0.004), and by a factor of 0.3 at 50 and 100 Hz

(p< 0.001). Seizures generally increased in duration with increasing current, but seizures

at CSD threshold often had reduced duration, apparently because the expanding CSD

terminated the ictal state. Frequency (p< 0.001) and pulse count (p=0.006) were

associated with longer seizure duration. Delivery of 50 pulses yielded 1.9-fold longer

seizure duration than 25 pulses (p=0.006) and 100 pulses yielded 2.4 fold longer seizure

duration than 25 pulses (p=0.012). Duration at pulse counts of 50 and 100 did not differ

significantly from each other. (p=0.49)

These parameters also impact the probability of generating CSD. Seizures that were

sufficient to elicit CSD had significantly higher seizure amplitudes **(Extended Data Fig.**

3, p < 0.0001 for both unilateral and bilateral CSD), but no difference in seizure duration

compared to seizures that do not elicit CSD. We thus asked if higher frequency stimulation

(which elicits higher amplitude seizures) was more like to result in a CSD. To this end, we

built a Cox model with current level as an analog of time (sequentially increased from 1,

2, 5, 10, 25 mA until CSD is achieved) and adjusting for electrode placement and pulse

count. We find that increasing the stimulation frequency by one level (e.g., 5 to 10 Hz)

increased the cumulative probability of CSD 2.1-fold at any point in the titration (95% CI

1.6-2.7, p< 0.001, **Fig. 3c**). Neither electrode placement (p=0.199) nor pulse count

(p=0.64) was associated with the probability of CSD in this model. Compared to the lowest

intensity 5 Hz stimulation, the probability of CSD was 16.5-fold greater at 100 Hz (p <

0.001) and 3.5-fold greater, but not significantly so, at 50 Hz (p=0.23); 10 Hz (p=0.87)

and 25 Hz (p=0.18) did not differ from 5 Hz. Thus, at any current level, higher stimulation

frequency statistically increased the likelihood of eliciting CSD, effectively leading CSD to

occur earlier in a titration at lower current. Lastly, we quantified post-event suppression

time by measuring the time between the terminal event (seizure or CSD) and subsequent

return of baseline slow wave 1-3 Hz spectral power from anesthesia (**Extended Data Fig.**

3c,d). We observed that post-ictal suppression times were relatively brief (20-70 seconds

before return of slow waves), but suppression times were significantly longer in
hemispheres affected by CSD (110-270s; see Figure for pairwise comparisons.)

Using concurrent hemodynamic imaging, we observed that seizure elicited a blush of
hyperemia. During CSD, this blush was followed by subsequent traveling waves of hypo-
and hyperperfusion, which propagated slowly in the wake of the neuronal depolarization
wave, consistent with prior reports[31,46]. Intuitively, we observe that bilateral CSD
consistently triggers bilateral hemodynamic traveling waves (**Extended Data Video 2**),
unilateral CSD triggers hemodynamic waves in only one hemisphere (**Extended Data**
Video 3), and seizure without CSD generates the initial blush, but no post-ictal traveling
hemodynamic wave (**Extended Data Video 4**). In each of these videos, two ROI pixel
time series are presented to illustrate that CSD hemodynamic waves can occur with a
variable post-ictal time delays and morphology, depending on where in the brain one
takes a point measurement in the path of a multifocal traveling wave. Optical detection of
this hemodynamic wave is a reliable biomarker of CSD in both rodents and humans[29,46-
50].

Cerebral hemodynamics in human ECT patients show expected features of post-
ictal CSD
Given that CSDs were reliably induced in mice across a wide range of stimulation
parameters, we hypothesized that CSDs could also occur in human patients receiving
ECT. We further hypothesized that the hemodynamic waves of post-ictal CSD could be
translationally detected using non-invasive diffuse optical monitoring of cerebral blood

flow and oxygen saturation (**Fig. 4a**). To this end, we measured local cerebral

hemodynamics using multimodal optical sensors on the forehead during routine human

ECT treatments in a single-center observational cohort study. We define four criteria for

detecting CSD in these measurements. First, we expect to see an initial hemodynamic

wave during seizure; subsequently during the post-ictal period, we expect hemodynamics

will either return to baseline in cases/hemispheres without CSD, or exhibit a second wave

(consistent with Extended Data Videos 1, 3, 4 and prior reports[38]). Second, based on prior

invasive regional measurements in human patients, if this post-ictal wave is a CSD, it will

exhibit >2-minute long duration of 200-600% increase in cerebral blood flow relative to

baseline[39]. Third, based on prior oximetry during CSD, we expect post-ictal waves will

exhibit variable polyphasic waves of either hypoxic, hyperoxic, or mixed response,[40,41]

with amplitude modulation of ±5% from baseline[42]. Because flow data was not available

in all subjects, we consider meeting either criterion two (blood flow) or three (oximetry)

sufficient. Fourth, given the impact of electrode spatial configuration on the hemispheric

laterality of mouse CSDs (**Fig. 2**), we predict that right unilateral ECT will elicit post-ictal

hemodynamic waves that are primarily right-sided (and occasionally bilateral), while

bilateral stimulation will elicit primarily bilateral waves.

Fig. 4. Cerebral hemodynamics during four ECT sessions from a single patient exhibit expected features of post-ictal CSD.

a, Diagram of diffuse optical probes (magenta on right at F4, green on left at F3), frontal EEG (grey circles at FP1 and FP2), and right unilateral electrodes (yellow).

b, Simplified schematic of non-invasive diffuse optics sensors, using paired red-light sources and detectors to regionally measure both deep (brain) and superficial (skin, scalp) blood flow (diffuse correlation spectroscopy, DCS) from two source-detector separations of 1 cm and 2.5 cm and oxygen saturation (frequency-domain diffuse optical spectroscopy, FD-DOS) from four source-detector separations of 1.5, 2, 2.5, and 3 cm. For clarity, the co-located DCS and FD-DOS probes are separated in the schematic. See **Methods** and **Supplemental Discussion and Methods** on diffuse optics.

c, Series of four recordings on four different treatment days from the same subject receiving right unilateral ECT. Note, the blood flow data shown are derived from the source-detector pair with largest separation and thus largest depth penetration into the brain (Data at shorter separations are given in **Extended Data Fig. 5**). The oxygen saturation data is the result of combining data from all FD-DOS source-detector pairs for improved signal quality.

In this pilot study, we obtained n=12 recordings of bilateral cerebral hemodynamics during

right unilateral ECT (n=5 patients) and bitemporal ECT (n=5 patients), summarized in

Extended Data Table 2 and **3**. We first present recordings from four treatments on

different days from an index course of right unilateral ECT for a 50-year-old woman with treatment-resistant depression (**Fig. 4b)**. All four treatments resulted in ~60s, bilateral seizures on frontal EEG (grey shaded region). During seizure, relative cerebral blood flow (measured with DCS long source-detector separation) increased by 200-500%. This change was observed predominantly in the right hemisphere, with an associated 5-10% *decrease* in brain $\%O_2$ saturation (derived from the FD-DOS signal).

After the seizure ended electrographically (i.e., scored by a clinician as a flat line on EEG), all four recordings exhibited an inflection point and then a distinct post-ictal wave of 200-600% increase in relative blood flow that lasted for over 2-minutes. Interestingly, unlike the preceding seizure, this second hyperemic wave was associated with a 5-10% *increase* in brain oxygen saturation, suggesting a hyperoxic surge of blood flow that kept pace with the metabolic demand of the brain[40,47]. These regional measurements of post-ictal hemodynamic waves exhibited variable waveform morphology and time delays relative to the initial electrographic seizure. In the first three recordings, post-ictal hyperemic waves were restricted to the right hemisphere (magenta lines), and in the last recording the hemodynamic changes occurred bilaterally, indicating three right hemispheric post-ictal events and one bilateral event. These hemispheric comparisons provide a within-subject control to demonstrate that hemodynamic changes are due to local brain metabolism and not to global autonomic changes. To further corroborate that these 200-600% changes in blood flow index are not due to systemic circulation, we implemented non-invasive cardiac output monitoring throughout the procedure. We observe that systemic blood flow from the heart was relatively constant at 5 ± 2 L/min

throughout the procedure. Taken together, these waves meet all four of our criteria for

defining post-ictal CSD. Of note, over this interval, the patient exhibited a relatively typical

clinical response to ECT. At her pre-treatment evaluation she reported intermittent

suicidal ideation, anhedonia, excess sleep, feelings of worthlessness, hopelessness, and

low energy. After 2 weeks (6 treatments) of ECT, she reported her mood, energy, and

sleep had improved, and that her suicidal thoughts had resolved. She denied any side

effects of treatment.

We collected eight more recordings from n=8 patients (**Extended Data Fig. 4**) and

similarly observed post-ictal hyperemic waves meeting criteria for CSD in an additional

3/3 recordings of right unilateral ECT as well as 4/5 recordings of bitemporal ECT. In all

recordings where sensor data was available bilaterally, putative CSDs occurred

bilaterally. This post-ictal surge in cerebral blood flow was also evident in the superficial

short separation DCS detectors, with variable amplitude relative to the long separation

DCS detectors **(**see **Extended Data Fig. 5** and **Supplemental Discussion**). Finally, we

provide additional replication cases obtained independently at another institution using a

commercial continuous-wave functional near infrared spectroscopy (fNIRS) instrument to

measure cerebral oximetry in human ECT patients (see **Extended Data Fig. 6** and

Supplemental Discussion). Thus, the hemodynamic signatures of post-ictal CSD were

reliably observed in three different modalities of non-invasive optical monitoring.

Discussion

In this study, we show that electroconvulsive therapy (ECT) reliably elicits post-ictal

cortical spreading depolarization (CSD) in both mouse models and in human patients.

This discovery unifies several previously disconnected observations: (1) that ECT induces

lasting inhibitory plasticity in brain activity[7], (2) that therapeutic response to ECT requires

high energy seizures of sufficient magnitude[51], and (3) that stimulation parameters

modulate seizure and clinical outcomes from ECT[20]. Our findings suggest that strong

stimulation elicits seizure of sufficient magnitude to generate post-ictal CSD.

Spreading depolarization may engage mechanisms of inhibitory plasticity that contribute

to the brain's clinical response to ECT. Post-ictal CSD has primarily been observed with

intracranial electrocorticography (ECoG) in both human patients[52-54] and in animal

models, dating back to Leão's original demonstration that CSD can be induced by direct

electrical or mechanical stimulation of cortex[55-57]. Because of the requirement for invasive

neurosurgery for electrophysiologic detection of CSD, virtually all reports of post-ictal

CSD in human patients have been obtained in individuals with brain lesions that already

required surgery (e.g., stroke and traumatic brain injury). In the context of injury and

metabolically fragile brain tissue, CSD is known to exacerbate excitotoxicity and worsen

clinical outcomes. However, in the context of seizure in an uninjured brain, preclinical

evidence suggests that CSD may act as an intrinsic protective mechanism, terminating

the seizure and inducing inhibitory plasticity that prevents future seizures[38,58]. CSD is not

known to be harmful in brains without underlying injury; in this context, CSD has been

observed to paradoxically induce growth and plasticity genes, downregulate cell death

genes, and protect against subsequent injury[54,59-62]. It remains unknown whether ECT-

induced CSDs are harmful or potentially part of the therapeutic mechanism. This cannot

be answered with this pilot study's limited sample of 12 recordings over single treatments

(nearly all of which exhibited evidence of CSD). This will be tested in future large-scale

clinical studies. One potential clue is the observation that post-ictal blood flow waves were

associated with a 5-10% *increase* in cerebral O_2 saturation – a feature associated with

CSDs with preserved neurovascular coupling and adequate perfusion, in contrast to the

decreased O_2 saturation seen in CSDs from ischemic injuries[47]. If CSD proves to be part

of the therapeutic mechanism of ECT, this would invite the exciting possibility of

bypassing seizure to directly trigger CSD. Indeed, we find that certain stimulation

conditions (i.e., brief trains of high frequency pulses) more efficiently generate CSD with

minimal seizure duration, while low frequency pulses require higher levels of current to

elicit high-amplitude seizure or CSD.

The inhibitory, antiseizure effects of CSD might help explain the long-known inhibitory

effects of ECT. ECT raises the seizure threshold in over 90% of patients[7], requiring

clinicians to progressively increase stimulation intensity between treatments to maintain

adequate seizure[11]. Ironically, this inhibitory effect makes ECT highly effective at treating

status epilepticus, particularly in super-refractory cases when maximal anti-seizure

pharmacotherapy has failed[63-65]. Furthermore, in those treated with ECT, EEG

demonstrates sustained increases in aperiodic power spectral density (an index of cortical

inhibitory tone)[66,67], as well as suppressed evoked measures of excitability (transcranial

magnetic stimulation evoked potentials)[10]. Human PET imaging has likewise shown that

ECT increases cortical GABA concentrations[68], and rodent studies have demonstrated

corresponding changes in GABA release[69] and GABA receptor mRNA levels[70]. In the

immediate aftermath of CSD, synaptic activity is potently suppressed, initially by

depolarization block of axonal action potentials, then by inhibition of presynaptic

neurotransmitter release due to activation of adenosine A1 receptors[71,72]. Adenosine

accumulation is also observed with seizures[73] and ECT[74]. Within inhibitory circuits, ECT-

induced plasticity may require cell-type specific molecular regulators of circuit

excitability[75] and act on excitability of single neurons[76]. We hypothesize that post-ictal

CSD may contribute to these multi-scale forms of inhibitory plasticity after ECT.

To our knowledge, this is the first study to record brain activity during ECT pulse delivery.

We used optical recordings to circumvent electrical signal artifacts that confound routine

EEG monitoring during stimulation. Previous computational predictions have shown that

larger evoked electric fields correlate with improved clinical response[15,77,78], which we

speculate may be due to increased probability of a large electric field triggering CSD.

Real-time recording of brain activity during ECT may prove valuable for future studies on

stimulus parameter optimization, just as it has for transcranial magnetic stimulation

(TMS)[79-81], deep brain stimulation (DBS)[82,83], and ultrasound neuromodulation[84]. Of note,

in our mouse model, we observed that high frequency stimulation at 100 Hz sometimes

evoked only a single high-amplitude spike during pulse delivery, without persistent post-

stimulation ictal activity. We opted to include evoked activity during stimulation as part of

the seizure, though the physiology governing persistent post-stimulation seizure may

differ from acute evoked discharges during stimulation. Both brief high-amplitude

discharges during stimulation as well as persistent post-stimulation seizure were sufficient

to generate CSD in our mouse model.

Post-ictal CSD has the potential to explain many epiphenomena seen in other

measurements of brain activity from ECT patients. We observed that larger seizures were

more likely to generate CSD in mice, which could theoretically explain why higher EEG

ictal spectral power is associated with improved clinical outcomes in humans[12,14,15]. Our

rodent data also showed longer post-event suppression time after CSD compared to

seizure alone, which could explain why longer suppression times are associated with

improved patient outcomes[12,85] and longer latency to reorientation[86]. Extended EEG

recording after treatment may be necessary to validate if post-ictal suppression is

associated with CSD[45,87]. Looking beyond EEG, CSD is known to produce lasting

changes in brain hemodynamics, which could account for why effective ECT seizures

elicit lasting perfusion changes 1-hour after treatment[88] as well as on longer

timescales[89,90]. Furthermore, we observed that right unilateral stimulation primarily

generated unilateral post-ictal waves. Unilateral CSD could play a role in ECT-induced

volumetric changes that occur preferentially in the stimulated hemisphere and correlate

with clinical outcomes and brain network functional connectivity changes[91-96]. CSDs are

so named because neocortex is most accessible for recording brain activity. Indeed, this

is a key limitation of our optical methods for recording cortical dynamics - spreading

depolarization can also occur in subcortical structures, which might have important

implications for both the clinical benefits and side effects of ECT.

We speculate that post-ictal CSD has been occurring undetected during the millions of

treatments that have been performed over the last 86 years. In clinical practice, EEG is

typically turned off after the seizure terminates, and furthermore, EEG cannot reliably

detect CSDs [33,44] unless the skull has been penetrated[45,97]. This is thought to be due to

digital high-pass filtering of slow waveforms, as well as spatial smearing from volume

conduction of CSD dynamics through the brain, skull, and scalp. Very focal CSDs with

narrow-width wavefronts may be particularly challenging to detect[45]. Optical recordings in

rodent models have long proven a valuable surrogate for CSD detection, providing richer

spatial detail on traveling waves compared to electrographic recordings[29,46-50]. Here we

replicate these findings in both optical imaging of neural and blood dynamics, and provide

supplemental verification of CSD with DC-ECoG (Extended Data Figure 2). However,

non-invasive optical methods have not yet been extended to human subjects prior to this

work.

Indeed, Dreier et al. have proposed that non-invasive optical hemodynamic measures

should prove sufficient to detect CSD in human patients[97]. Here we find that optical

features of CSD can be detected non-invasively at patient bedside, and meet all of our

criteria for defining CSD based on prior literature on the amplitude, duration, and

morphology of CSD hemodynamic waves[38-42]. This approach to CSD detection may have

valuable clinical applications in neurocritical care for individuals without intracranial ECoG

implants. Previous studies have characterized the hemodynamic features of CSD using

invasive probes in the brain[39-41,98] in those with brain injuries. In migraine patients, CSD

has been detected non-invasively using functional MRI[42,99] and radiotracer/PET

imaging[100,101], revealing that CSD is the physiologic basis of migraine aura (and may

perhaps explain headache as an occasional side effect of ECT). Note, PET and MRI are

incompatible with ECT due to the need for mechanical ventilation during treatment. In this

study we implemented a non-invasive investigational device that combined frequency-

domain diffuse optical spectroscopy (FD-DOS) to measure tissue oxygenation with

diffuse correlation spectroscopy (DCS) to capture blood flow rate changes.

Our human optical recordings have several limitations. First, it is not possible to obtain

electrographic verification of post-ictal CSD in these recordings – which would require

invasive neurosurgery for ECoG implantation, and would not be clinically appropriate in

ECT patients. As expected from prior reports[33,44,45,97], we were unable to detect CSD

using scalp DC-EEG (see **Supplement**). Second, our approach relied on regional point

measurements over non-hair bearing skin. Future investigations using broader spatial

coverage at multiple points would be needed to detect CSD propagation across a brain

hemisphere. Third, signal quality is significantly influenced by patient motion, skin

temperature, and variation in skull and skin microanatomy that can influence the detected

optical path distribution (see **Supplemental Discussion**/**Methods**). These factors can

introduce systematic errors that influence absolute tissue optical properties. We have

ameliorated these effects by computing relative change in blood flow and oxygenation

indices within each individual.

This study implemented a new mouse model of ECT that enabled experimental control

over stimulation combined with real-time monitoring of brain activity with widefield

imaging. To model the three standard ECT electrode placements (right unilateral,

bitemporal, bifrontal), we attached five electrodes to the intact skull with silver-conductive

epoxy to test ten configurations of electrode pairs. This builds on prior innovation using

screw electrodes in rats, which have been shown to have superior translational validity to

conventional rodent ECT with auricular (ear clip) electrodes.[102] Skull electrodes help

control for the confounding effects of current shunting through the low-resistance skin[103].

Indeed, we observed the use of ear clip electrodes required 5-10 times more current to

elicit seizure or CSD compared to skull electrodes (see **Supplement**). In addition, we

substituted a conventional rodent ECT device for a broadly customizable isolated pulse

stimulator, enabling the first systematic evaluation of ECT pulse parameters in an animal

model. We found that increasing pulse frequency increased seizure amplitude and

probability of CSD, while low frequency pulses elicited long duration seizures without

increased probability of CSD. This builds on prior studies showing that high frequency

electrical pulses can elicit CSD in hippocampal slices[104], and that low intensity stimulation

does not elicit CSD[105]. One limitation of this study is its inability to exhaustively survey the

virtually infinite space of pulse parameters and electrode configurations[20]. We opted to

survey multiple electrode and pulse parameters simultaneously, using

pseudorandomization to sample stimulation conditions in a balanced fashion, and linear

mixed effects modeling to isolate pulse parameter-specific effects in titration data. This is

an imperfect solution, and future investigations would benefit from systematically

modulating individual stimulation parameters, as well as testing for specific effects in

translational models of neuropsychiatric disease. Lastly, using a current titration strategy,

we identified that increasing pulse frequency lowered the current threshold required to

elicit a seizure of a given magnitude. This has important ramifications for novel ECT

protocols that modulate current amplitude[26,78,106]. Future investigations should explore

whether high frequency stimulation is a more current-sparing approach of achieving

therapeutic efficacy. This study thus provides a translational framework for measuring

brain activity to optimize ECT stimulation parameters and precisely control seizure and

post-ictal CSD. These findings further highlight opportunities to modernize ECT and

directly target treatment-induced mechanisms of brain plasticity.

Author Contributions:
All authors helped with data interpretation and editorial feedback on the paper.
ZPR conceptualized overall project; designed, performed, and analyzed results of mouse experiments; generated Penn IRB protocol, collected and analyzed human data; wrote original manuscript, produced figures; contributed funding.
JBM contributed to diffuse optics method development, human data collection and analysis, and text in the original manuscript
AS contributed to Penn mouse imaging data analysis and text in the original manuscript
DKQ designed and performed UNM human recordings
BEL contributed UNM mouse recordings and text in the original manuscript.
MEP conceptualized and implemented time-to-event and linear mixed effects modeling and contributed text in the original manuscript.
AK contributed to the statistical analysis presented in the original manuscript.
CGF contributed to methodological development of human studies and IRB protocol.
WBB contributed to conceptualization and methods development for diffuse optics.
GH is an ECT clinician who helped supervise human data collection.
JPR is an ECT clinician who helped supervise human data collection.
MAC is an ECT clinician who helped supervise human data collection and advise on project conceptualization.
YIS helped supervise development of human studies and funding acquisition.
CWS conceptualized and supervised UNM mouse experiments.
CCA is an ECT clinician and conceptualized and supervised UNM human case series.
AGY contributed diffuse optics resources for human recordings, method development, data analysis, postdoctoral supervision and funding support of JBM.
EMG contributed to method development of mouse ECT model and widefield imaging, editorial supervision of original manuscript, postdoctoral supervision of ZPR.

Acknowledgements: We would like to thank the patients who participated in the study for their generosity and openness. This work was supported by the Penn Psychiatry Residency Research Track award NIH R25MH119043 (ZPR); Institute for Translational Medicine and Therapeutics of the University of Pennsylvania NIH UL1TR001878 (ZPR); NIH P41-EB029460 (AGY); NIH NS078805, NS051288, NS070680 (BEL); NIH NS106901 (CWS); NIH P50HD105354 (MEP, AK). The authors would like to thank Emily Hoddeson for assistance with mouse husbandry and genotyping, Dr. Adam Bauer for advice on mouse optical methods, Dr. Jin-Moo Lee for sharing breeders from the *Thy1-jRGECO1a* mouse line, ANT Neuro for loaning DC-EEG equipment, Katy Brown for facilitating human continuous vitals monitoring, Dr. Holly Lisanby for advice on project conceptualization, Jurre Blom for assistance with manuscript illustrations, Dr. Brian White for help launching the collaboration between the diffuse optics and ECT research teams, and Dr. Rodrigo M. Forti for help with data interpretation and diffuse optical instrumentation.

Competing Interests: No financial conflicts or completing interests to disclose.

Data Availability Source Data are provided in the Goldberg Lab gnode repository (https://gin.g-node.org/GoldbergNeuroLab).

References:

1. Espinoza, R. T. & Kellner, C. H. Electroconvulsive Therapy. *N Engl J Med* **386**, 667-672, (2022).
2. Kaster, T. S., Blumberger, D. M., Gomes, T., Sutradhar, R., Wijeysundera, D. N. & Vigod, S. N. Risk of suicide death following electroconvulsive therapy treatment for depression: a propensity score-weighted, retrospective cohort study in Canada. *Lancet Psychiatry* **9**, 435-446, (2022).
3. Fink, M. What was learned: studies by the consortium for research in ECT (CORE) 1997-2011. *Acta Psychiatr Scand* **129**, 417-426, (2014).
4. Sackeim, H. A. Modern Electroconvulsive Therapy: Vastly Improved yet Greatly Underused. *JAMA Psychiatry* **74**, 779-780, (2017).
5. Deng, Z. D., Robins, P. L., Regenold, W., Rohde, P., Dannhauer, M. & Lisanby, S. H. How electroconvulsive therapy works in the treatment of depression: is it the seizure, the electricity, or both? *Neuropsychopharmacology* **49**, 150-162, (2024).
6. Regenold, W. T., Noorani, R. J., Piez, D. & Patel, P. Nonconvulsive Electrotherapy for Treatment Resistant Unipolar and Bipolar Major Depressive Disorder: A Proof-of-concept Trial. *Brain Stimul* **8**, 855-861, (2015).
7. Duthie, A. C., Perrin, J. S., Bennett, D. M., Currie, J. & Reid, I. C. Anticonvulsant Mechanisms of Electroconvulsive Therapy and Relation to Therapeutic Efficacy. *J ECT* **31**, 173-178, (2015).
8. Lloyd, R. L. & Sattin, A. The Behavioral Physiology and Antidepressant Mechanisms of Electroconvulsive Shock. *J ECT* **31**, 159-166, (2015).
9. Voineskos, D., Levinson, A. J., Sun, Y., Barr, M. S., Farzan, F., Rajji, T. K., Fitzgerald, P. B., Blumberger, D. M. & Daskalakis, Z. J. The Relationship Between Cortical Inhibition and Electroconvulsive Therapy in the Treatment of Major Depressive Disorder. *Sci Rep* **6**, 37461, (2016).
10. Bajbouj, M., Lang, U. E., Niehaus, L., Hellen, F. E., Heuser, I. & Neu, P. Effects of right unilateral electroconvulsive therapy on motor cortical excitability in depressive patients. *J Psychiatr Res* **40**, 322-327, (2006).
11. Krystal, A. D., Coffey, C. E., Weiner, R. D. & Holsinger, T. Changes in seizure threshold over the course of electroconvulsive therapy affect therapeutic response and are detected by ictal EEG ratings. *J Neuropsychiatry Clin Neurosci* **10**, 178-186, (1998).
12. Perera, T. D., Luber, B., Nobler, M. S., Prudic, J., Anderson, C. & Sackeim, H. A. Seizure expression during electroconvulsive therapy: relationships with clinical outcome and cognitive side effects. *Neuropsychopharmacology* **29**, 813-825, (2004).
13. Zhang, J. Y., Xu, S. X., Zeng, L., Chen, L. C., Li, J., Jiang, Z. Y., Tan, B. J., Gu, C. L., Lai, W. T., Kong, X. M., Wang, J., Rong, H. & Xie, X. H. Improved Safety of Hybrid Electroconvulsive Therapy Compared With Standard Electroconvulsive Therapy in Patients With Major Depressive Disorder: A Randomized, Double-Blind, Parallel-Group Pilot Trial. *Front Psychiatry* **13**, 896018, (2022).
14. Miller, J., Jones, T., Upston, J., Deng, Z. D., McClintock, S. M., Ryman, S., Quinn, D. & Abbott, C. C. Ictal Theta Power as an Electroconvulsive Therapy Safety Biomarker: A Pilot Study. *J ECT* **38**, 88-94, (2022).
15. Miller, J., Jones, T., Upston, J., Deng, Z. D., McClintock, S. M., Erhardt, E., Farrar, D. & Abbott, C. C. Electric Field, Ictal Theta Power, and Clinical Outcomes in

Electroconvulsive Therapy. *Biol Psychiatry Cogn Neurosci Neuroimaging* **8**, 760-767,
(2023).

16. Gillving, C., Ekman, C. J., Hammar, A., Landen, M., Lundberg, J., Movahed Rad, P.,
Nordanskog, P., von Knorring, L. & Nordenskjold, A. Seizure Duration and
Electroconvulsive Therapy in Major Depressive Disorder. *JAMA Netw Open* **7**,
e2422738, (2024).

17. Khadka, N., Deng, Z. D., Lisanby, S. H., Bikson, M. & Camprodon, J. A. Computational
Models of High-Definition Electroconvulsive Therapy for Focal or Multitargeting
Treatment. *J ECT*, (2024).

18. Argyelan, M. *et al.* Electroconvulsive therapy-induced volumetric brain changes converge on
a common causal circuit in depression. *Mol Psychiatry* **29**, 229-237, (2024).

19. Deng, Z. D., Argyelan, M., Miller, J., Quinn, D. K., Lloyd, M., Jones, T. R., Upston, J.,
Erhardt, E., McClintock, S. M. & Abbott, C. C. Electroconvulsive therapy, electric field,
neuroplasticity, and clinical outcomes. *Mol Psychiatry* **27**, 1676-1682, (2022).

20. Peterchev, A. V., Rosa, M. A., Deng, Z. D., Prudic, J. & Lisanby, S. H. Electroconvulsive
therapy stimulus parameters: rethinking dosage. *J ECT* **26**, 159-174, (2010).

21. Sackeim, H. A., Prudic, J., Nobler, M. S., Fitzsimons, L., Lisanby, S. H., Payne, N., Berman,
R. M., Brakemeier, E. L., Perera, T. & Devanand, D. P. Effects of pulse width and
electrode placement on the efficacy and cognitive effects of electroconvulsive therapy.
Brain Stimul **1**, 71-83, (2008).

22. Wang, X., Padawer-Curry, J. A., Bice, A. R., Kim, B., Rosenthal, Z. P., Lee, J.-M., Goyal,
M. S., Macauley, S. L. & Bauer, A. Q. Spatiotemporal relationships between neuronal,
metabolic, and hemodynamic signals in the awake and anesthetized mouse brain. *Cell*
Reports **43**, (2024).

23. Brier, L. M., Landsness, E. C., Snyder, A. Z., Wright, P. W., Baxter, G. A., Bauer, A. Q.,
Lee, J. M. & Culver, J. P. Separability of calcium slow waves and functional connectivity
during wake, sleep, and anesthesia. *Neurophotonics* **6**, 035002, (2019).

24. Mitra, A., Kraft, A., Wright, P., Acland, B., Snyder, A. Z., Rosenthal, Z., Czerniewski, L.,
Bauer, A., Snyder, L., Culver, J., Lee, J. M. & Raichle, M. E. Spontaneous Infra-slow
Brain Activity Has Unique Spatiotemporal Dynamics and Laminar Structure. *Neuron* **98**,
297-305 e296, (2018).

25. Mukhtar, F., Regenold, W. & Lisanby, S. H. Recent advances in electroconvulsive therapy in
clinical practice and research. *Faculty Reviews* **12**, (2023).

26. Peterchev, A. V., Krystal, A. D., Rosa, M. A. & Lisanby, S. H. Individualized Low-
Amplitude Seizure Therapy: Minimizing Current for Electroconvulsive Therapy and
Magnetic Seizure Therapy. *Neuropsychopharmacology* **40**, 2076-2084, (2015).

27. Brier, L. M. & Culver, J. P. Open-source statistical and data processing tools for wide-field
optical imaging data in mice. *Neurophotonics* **10**, 016601, (2023).

28. Franklin, K. B. J. & Paxinos, G. *The Mouse Brain in Stereotactic Coordinates*. (Academic
Press, 2012).

29. Chung, D. Y., Sugimoto, K., Fischer, P., Bohm, M., Takizawa, T., Sadeghian, H., Morais, A.,
Harriott, A., Oka, F., Qin, T., Henninger, N., Yaseen, M. A., Sakadzic, S. & Ayata, C.
Real-time non-invasive in vivo visible light detection of cortical spreading
depolarizations in mice. *J Neurosci Methods* **309**, 143-146, (2018).

30. Balbi, M., Vanni, M. P., Silasi, G., Sekino, Y., Bolanos, L., LeDue, J. M. & Murphy, T. H.
Targeted ischemic stroke induction and mesoscopic imaging assessment of blood flow
and ischemic depolarization in awake mice. *Neurophotonics* **4**, 035001, (2017).
31. Zhao, H. T., Tuohy, M. C., Chow, D., Kozberg, M. G., Kim, S. H., Shaik, M. A. & Hillman,
E. M. C. Neurovascular dynamics of repeated cortical spreading depolarizations after
acute brain injury. *Cell Rep* **37**, 109794, (2021).
32. Smith, S. E. *et al.* Astrocyte deletion of alpha2-Na/K ATPase triggers episodic motor
paralysis in mice via a metabolic pathway. *Nat Commun* **11**, 6164, (2020).
33. Hofmeijer, J., van Kaam, C. R., van de Werff, B., Vermeer, S. E., Tjepkema-Cloostermans,
M. C. & van Putten, M. Detecting Cortical Spreading Depolarization with Full Band
Scalp Electroencephalography: An Illusion? *Front Neurol* **9**, 17, (2018).
34. Carp, S. A., Robinson, M. B. & Franceschini, M. A. Diffuse correlation spectroscopy:
current status and future outlook. *Neurophotonics* **10**, 013509, (2023).
35. Durduran, T., Choe, R., Baker, W. B. & Yodh, A. G. Diffuse Optics for Tissue Monitoring
and Tomography. *Rep Prog Phys* **73**, (2010).
36. Zhou, X., Xia, Y., Uchitel, J., Collins-Jones, L., Yang, S., Loureiro, R., Cooper, R. J. &
Zhao, H. Review of recent advances in frequency-domain near-infrared spectroscopy
technologies [Invited]. *Biomed Opt Express* **14**, 3234-3258, (2023).
37. Kouz, K. *et al.* Intraoperative hypotension when using hypotension prediction index software
during major noncardiac surgery: a European multicentre prospective observational
registry (EU HYPROTECT). *BJA Open* **6**, 100140, (2023).
38. Tamim, I., Chung, D. Y., de Morais, A. L., Loonen, I. C. M., Qin, T., Misra, A., Schlunk, F.,
Endres, M., Schiff, S. J. & Ayata, C. Spreading depression as an innate antiseizure
mechanism. *Nat Commun* **12**, 2206, (2021).
39. Lemale, C. L., Luckl, J., Horst, V., Reiffurth, C., Major, S., Hecht, N., Woitzik, J. & Dreier,
J. P. Migraine Aura, Transient Ischemic Attacks, Stroke, and Dying of the Brain Share
the Same Key Pathophysiological Process in Neurons Driven by Gibbs-Donnan Forces,
Namely Spreading Depolarization. *Front Cell Neurosci* **16**, 837650, (2022).
40. Winkler, M. K., Dengler, N., Hecht, N., Hartings, J. A., Kang, E. J., Major, S., Martus, P.,
Vajkoczy, P., Woitzik, J. & Dreier, J. P. Oxygen availability and spreading
depolarizations provide complementary prognostic information in neuromonitoring of
aneurysmal subarachnoid hemorrhage patients. *J Cereb Blood Flow Metab* **37**, 1841-
1856, (2017).
41. Hecht, N., Haddad, D., Neumann, K., Schumm, L., Dengler, N. F., Wessels, L., Domer, P.,
Helgers, S., Meinert, F., Major, S., Lemale, C. L., Dreier, J. P., Vajkoczy, P. & Woitzik,
J. Reduced brain oxygen response to spreading depolarization predicts worse outcome in
ischaemic stroke. *Brain*, (2024).
42. Hadjikhani, N., Sanchez Del Rio, M., Wu, O., Schwartz, D., Bakker, D., Fischl, B., Kwong,
K. K., Cutrer, F. M., Rosen, B. R., Tootell, R. B., Sorensen, A. G. & Moskowitz, M. A.
Mechanisms of migraine aura revealed by functional MRI in human visual cortex. *Proc*
Natl Acad Sci U S A **98**, 4687-4692, (2001).
43. Dana, H., Novak, O., Guardado-Montesino, M., Fransen, J. W., Hu, A., Borghuis, B. G.,
Guo, C., Kim, D. S. & Svoboda, K. Thy1 transgenic mice expressing the red fluorescent
calcium indicator jRGECO1a for neuronal population imaging in vivo. *PLoS One* **13**,
e0205444, (2018).

44. Dreier, J. P., Major, S., Lemale, C. L., Kola, V., Reiffurth, C., Schoknecht, K., Hecht, N.,
Hartings, J. A. & Woitzik, J. Correlates of Spreading Depolarization, Spreading
Depression, and Negative Ultraslow Potential in Epidural Versus Subdural
Electrocorticography. *Front Neurosci* **13**, 373, (2019).
45. Chamanzar, A., George, S., Venkatesh, P., Chamanzar, M., Shutter, L., Elmer, J. & Grover,
P. An Algorithm for Automated, Noninvasive Detection of Cortical Spreading
Depolarizations Based on EEG Simulations. *IEEE Trans Biomed Eng* **66**, 1115-1126,
(2019).
46. Ayata, C., Shin, H. K., Salomone, S., Ozdemir-Gursoy, Y., Boas, D. A., Dunn, A. K. &
Moskowitz, M. A. Pronounced hypoperfusion during spreading depression in mouse
cortex. *J Cereb Blood Flow Metab* **24**, 1172-1182, (2004).
47. Dreier, J. P. The role of spreading depression, spreading depolarization and spreading
ischemia in neurological disease. *Nat Med* **17**, 439-447, (2011).
48. de Crespigny, A., Rother, J., van Bruggen, N., Beaulieu, C. & Moseley, M. E. Magnetic
resonance imaging assessment of cerebral hemodynamics during spreading depression in
rats. *J Cereb Blood Flow Metab* **18**, 1008-1017, (1998).
49. Zhou, C., Yu, G., Furuya, D., Greenberg, J., Yodh, A. & Durduran, T. Diffuse optical
correlation tomography of cerebral blood flow during cortical spreading depression in rat
brain. *Opt Express* **14**, 1125-1144, (2006).
50. Yamato, H., Jin, T. & Nomura, Y. Near infrared imaging of intrinsic signals in cortical
spreading depression observed through the intact scalp in hairless mice. *Neurosci Lett*
701, 213-217, (2019).
51. Sackeim, H. A., Prudic, J., Devanand, D. P., Kiersky, J. E., Fitzsimons, L., Moody, B. J.,
McElhiney, M. C., Coleman, E. A. & Settembrino, J. M. Effects of stimulus intensity and
electrode placement on the efficacy and cognitive effects of electroconvulsive therapy. *N
Engl J Med* **328**, 839-846, (1993).
52. Fabricius, M., Fuhr, S., Willumsen, L., Dreier, J. P., Bhatia, R., Boutelle, M. G., Hartings, J.
A., Bullock, R., Strong, A. J. & Lauritzen, M. Association of seizures with cortical
spreading depression and peri-infarct depolarisations in the acutely injured human brain.
Clin Neurophysiol **119**, 1973-1984, (2008).
53. Dreier, J. P., Major, S., Pannek, H. W., Woitzik, J., Scheel, M., Wiesenthal, D., Martus, P.,
Winkler, M. K., Hartings, J. A., Fabricius, M., Speckmann, E. J., Gorji, A. & group, C. s.
Spreading convulsions, spreading depolarization and epileptogenesis in human cerebral
cortex. *Brain* **135**, 259-275, (2012).
54. Dreier, J. P. *et al.* Recording, analysis, and interpretation of spreading depolarizations in
neurointensive care: Review and recommendations of the COSBID research group. *J
Cereb Blood Flow Metab* **37**, 1595-1625, (2017).
55. Leao, A. A. Further observations on the spreading depression of activity in the cerebral
cortex. *J Neurophysiol* **10**, 409-414, (1947).
56. Leao, A. A. P. Spreading Depression of Activity in the Cerebral Cortex. *Journal of
Neurophysiology* **7**, 359-390, (1944).
57. Bragin, A., Penttonen, M. & Buzsaki, G. Termination of epileptic afterdischarge in the
hippocampus. *J Neurosci* **17**, 2567-2579, (1997).
58. Zakharov, A., Chernova, K., Burkhanova, G., Holmes, G. L. & Khazipov, R. Segregation of
seizures and spreading depolarization across cortical layers. *Epilepsia* **60**, 2386-2397,
(2019).

59. Dell'Orco, M., Weisend, J. E., Perrone-Bizzozero, N. I., Carlson, A. P., Morton, R. A.,
Linsenbardt, D. N. & Shuttleworth, C. W. Repetitive spreading depolarization induces
gene expression changes related to synaptic plasticity and neuroprotective pathways.
Front Cell Neurosci **17**, 1292661, (2023).
60. Nedergaard, M. & Hansen, A. J. Spreading depression is not associated with neuronal injury
in the normal brain. *Brain Res* **449**, 395-398, (1988).
61. Kobayashi, S., Harris, V. A. & Welsh, F. A. Spreading depression induces tolerance of
cortical neurons to ischemia in rat brain. *J Cereb Blood Flow Metab* **15**, 721-727, (1995).
62. Matsushima, K., Hogan, M. J. & Hakim, A. M. Cortical spreading depression protects
against subsequent focal cerebral ischemia in rats. *J Cereb Blood Flow Metab* **16**, 221-
226, (1996).
63. Lambrecq, V., Villega, F., Marchal, C., Michel, V., Guehl, D., Rotge, J. Y. & Burbaud, P.
Refractory status epilepticus: electroconvulsive therapy as a possible therapeutic strategy.
Seizure **21**, 661-664, (2012).
64. Garcia-Lopez, B. *et al.* Electroconvulsive Therapy in Super Refractory Status Epilepticus:
Case Series with a Defined Protocol. *Int J Environ Res Public Health* **17**, (2020).
65. Ahmed, J., Metrick, M., Gilbert, A., Glasson, A., Singh, R., Ambrous, W., Brown, L.,
Aykroyd, L. & Bobel, K. Electroconvulsive Therapy for Super Refractory Status
Epilepticus. *J ECT* **34**, e5-e9, (2018).
66. Smith, S. E., Ma, V., Gonzalez, C., Chapman, A., Printz, D., Voytek, B. & Soltani, M.
Clinical EEG slowing induced by electroconvulsive therapy is better described by
increased frontal aperiodic activity. *Transl Psychiatry* **13**, 348, (2023).
67. Smith, S. E., Kosik, E. L., van Engen, Q., Kohn, J., Hill, A. T., Zomorrodi, R., Blumberger,
D. M., Daskalakis, Z. J., Hadas, I. & Voytek, B. Magnetic seizure therapy and
electroconvulsive therapy increase aperiodic activity. *Transl Psychiatry* **13**, 347, (2023).
68. Sanacora, G., Mason, G. F., Rothman, D. L., Hyder, F., Ciarcia, J. J., Ostroff, R. B., Berman,
R. M. & Krystal, J. H. Increased cortical GABA concentrations in depressed patients
receiving ECT. *Am J Psychiatry* **160**, 577-579, (2003).
69. Green, A. R. & Vincent, N. D. The effect of repeated electroconvulsive shock on GABA
synthesis and release in regions of rat brain. *Br J Pharmacol* **92**, 19-24, (1987).
70. Kang, I., Miller, L. G., Moises, J. & Bazan, N. G. GABAA receptor mRNAs are increased
after electroconvulsive shock. *Psychopharmacol Bull* **27**, 359-363, (1991).
71. Lindquist, B. E. & Shuttleworth, C. W. Adenosine receptor activation is responsible for
prolonged depression of synaptic transmission after spreading depolarization in brain
slices. *Neuroscience* **223**, 365-376, (2012).
72. Lindquist, B. E. & Shuttleworth, C. W. Evidence that adenosine contributes to Leao's
spreading depression in vivo. *J Cereb Blood Flow Metab* **37**, 1656-1669, (2017).
73. Boison, D. Adenosinergic signaling in epilepsy. *Neuropharmacology* **104**, 131-139, (2016).
74. Lewin, E. & Bleck, V. Electroshock seizures in mice: effect on brain adenosine and its
metabolites. *Epilepsia* **22**, 577-581, (1981).
75. Chang, A. D. *et al.* Narp Mediates Antidepressant-Like Effects of Electroconvulsive
Seizures. *Neuropsychopharmacology* **43**, 1088-1098, (2018).
76. Ueta, Y., Yamamoto, R. & Kato, N. Layer-specific modulation of pyramidal cell excitability
by electroconvulsive shock. *Neurosci Lett* **709**, 134383, (2019).

77. Lee, W. H., Lisanby, S. H., Laine, A. F. & Peterchev, A. V. Comparison of electric field
strength and spatial distribution of electroconvulsive therapy and magnetic seizure
therapy in a realistic human head model. *Eur Psychiatry* **36**, 55-64, (2016).
78. Abbott, C. C., Quinn, D., Miller, J., Ye, E., Iqbal, S., Lloyd, M., Jones, T. R., Upston, J.,
Deng, Z., Erhardt, E. & McClintock, S. M. Electroconvulsive Therapy Pulse Amplitude
and Clinical Outcomes. *Am J Geriatr Psychiatry* **29**, 166-178, (2021).
79. Gogulski, J., Ross, J. M., Talbot, A., Cline, C. C., Donati, F. L., Munot, S., Kim, N., Gibbs,
C., Bastin, N., Yang, J., Minasi, C., Sarkar, M., Truong, J. & Keller, C. J. Personalized
Repetitive Transcranial Magnetic Stimulation for Depression. *Biol Psychiatry Cogn
Neurosci Neuroimaging* **8**, 351-360, (2023).
80. Wang, J. B., Hassan, U., Bruss, J. E., Oya, H., Uitermarkt, B. D., Trapp, N. T., Gander, P. E.,
Howard, M. A., 3rd, Keller, C. J. & Boes, A. D. Effects of transcranial magnetic
stimulation on the human brain recorded with intracranial electrocorticography. *Mol
Psychiatry* **29**, 1228-1240, (2024).
81. Peterchev, A. V., Goetz, S. M., Westin, G. G., Luber, B. & Lisanby, S. H. Pulse width
dependence of motor threshold and input-output curve characterized with controllable
pulse parameter transcranial magnetic stimulation. *Clin Neurophysiol* **124**, 1364-1372,
(2013).
82. Paulk, A. C., Zelmann, R., Crocker, B., Widge, A. S., Dougherty, D. D., Eskandar, E. N.,
Weisholtz, D. S., Richardson, R. M., Cosgrove, G. R., Williams, Z. M. & Cash, S. S.
Local and distant cortical responses to single pulse intracranial stimulation in the human
brain are differentially modulated by specific stimulation parameters. *Brain Stimul* **15**,
491-508, (2022).
83. Huang, Y., Zelmann, R., Hadar, P., Dezha-Peralta, J., Richardson, R. M., Williams, Z. M.,
Cash, S. S., Keller, C. J. & Paulk, A. C. Theta-burst direct electrical stimulation remodels
human brain networks. *Nat Commun* **15**, 6982, (2024).
84. Murphy, K. R. *et al.* Optimized ultrasound neuromodulation for non-invasive control of
behavior and physiology. *Neuron*, (2024).
85. Moulier, V., Guehl, J., Eveque-Mourroux, E., Quesada, P. & Rotharmel, M. A Retrospective
Study of Postictal Suppression during Electroconvulsive Therapy. *J Clin Med* **11**, (2022).
86. J, C. M. P., J, P. A. J. V., Stuiver, S., Aalbregt, E., Schmettow, M., Hofmeijer, J., van
Waarde, J. A. & M, J. A. M. v. P. Seizure duration predicts postictal
electroencephalographic recovery after electroconvulsive therapy-induced seizures. *Clin
Neurophysiol* **148**, 1-8, (2023).
87. Hickman, L. B., Kafashan, M., Labonte, A. K., Chan, C. W., Huels, E. R., Guay, C. S., Guan,
M. J., Ching, S., Lenze, E. J., Farber, N. B., Avidan, M. S., Hogan, R. E. & Palanca, B. J.
A. Postictal generalized electroencephalographic suppression following electroconvulsive
therapy: Temporal characteristics and impact of anesthetic regimen. *Clin Neurophysiol*
132, 977-983, (2021).
88. Pottkamper, J. C. M., Verdijk, J., Aalbregt, E., Stuiver, S., van de Mortel, L., Norris, D. G.,
van Putten, M., Hofmeijer, J., van Wingen, G. A. & van Waarde, J. A. Changes in
postictal cerebral perfusion are related to the duration of electroconvulsive therapy-
induced seizures. *Epilepsia* **65**, 177-189, (2024).
89. Hirano, J., Takamiya, A., Yamagata, B., Hotta, S., Miyasaka, Y., Pu, S., Iwanami, A.,
Uchida, H. & Mimura, M. Frontal and temporal cortical functional recovery after

electroconvulsive therapy for depression: A longitudinal functional near-infrared spectroscopy study. *J Psychiatr Res* **91**, 26-35, (2017).

90. Downey, D., Brigadoi, S., Trevithick, L., Elliott, R., Elwell, C., McAllister-Williams, R. H. & Anderson, I. M. Frontal haemodynamic responses in depression and the effect of electroconvulsive therapy. *J Psychopharmacol* **33**, 1003-1014, (2019).

91. Mulders, P. C. R. *et al.* Structural changes induced by electroconvulsive therapy are associated with clinical outcome. *Brain Stimul* **13**, 696-704, (2020).

92. Verdijk, J. *et al.* Longitudinal resting-state network connectivity changes in electroconvulsive therapy patients compared to healthy controls. *Brain Stimul* **17**, 140-147, (2024).

93. Abbott, C. C., Jones, T., Lemke, N. T., Gallegos, P., McClintock, S. M., Mayer, A. R., Bustillo, J. & Calhoun, V. D. Hippocampal structural and functional changes associated with electroconvulsive therapy response. *Transl Psychiatry* **4**, e483, (2014).

94. Argyelan, M. *et al.* Electric field causes volumetric changes in the human brain. *Elife* **8**, (2019).

95. Cano, M., Lee, E., Cardoner, N., Martinez-Zalacain, I., Pujol, J., Makris, N., Henry, M., Via, E., Hernandez-Ribas, R., Contreras-Rodriguez, O., Menchon, J. M., Urretavizcaya, M., Soriano-Mas, C. & Camprodon, J. A. Brain Volumetric Correlates of Right Unilateral Versus Bitemporal Electroconvulsive Therapy for Treatment-Resistant Depression. *J Neuropsychiatry Clin Neurosci* **31**, 152-158, (2019).

96. Leaver, A. M., Vasavada, M., Kubicki, A., Wade, B., Loureiro, J., Hellemann, G., Joshi, S. H., Woods, R. P., Espinoza, R. & Narr, K. L. Hippocampal subregions and networks linked with antidepressant response to electroconvulsive therapy. *Mol Psychiatry* **26**, 4288-4299, (2021).

97. Drenckhahn, C., Winkler, M. K., Major, S., Scheel, M., Kang, E. J., Pinczolits, A., Grozea, C., Hartings, J. A., Woitzik, J., Dreier, J. P. & group, C. s. Correlates of spreading depolarization in human scalp electroencephalography. *Brain* **135**, 853-868, (2012).

98. Thomas, R., Shin, S. S. & Balu, R. Applications of near-infrared spectroscopy in neurocritical care. *Neurophotonics* **10**, 023522, (2023).

99. Gollion, C., Christensen, R. H., Ashina, H., Al-Khazali, H. M., Fisher, P. M., Amin, F. M., Lauritzen, M. & Ashina, M. Somatosensory migraine auras evoked by bihemispheric cortical spreading depression events in human parietal cortex. *J Cereb Blood Flow Metab*, 271678X241290606, (2024).

100. Woods, R. P., Iacoboni, M. & Mazziotta, J. C. Brief report: bilateral spreading cerebral hypoperfusion during spontaneous migraine headache. *N Engl J Med* **331**, 1689-1692, (1994).

101. Olesen, J., Larsen, B. & Lauritzen, M. Focal hyperemia followed by spreading oligemia and impaired activation of rCBF in classic migraine. *Ann Neurol* **9**, 344-352, (1981).

102. Theilmann, W., Loscher, W., Socala, K., Frieling, H., Bleich, S. & Brandt, C. A new method to model electroconvulsive therapy in rats with increased construct validity and enhanced translational value. *J Psychiatr Res* **53**, 94-98, (2014).

103. Vöröslakos, M., Takeuchi, Y., Brinyiczki, K., Zombori, T., Oliva, A., Fernández-Ruiz, A., Kozák, G., Kincses, Z. T., Iványi, B., Buzsáki, G. & Berényi, A. Direct effects of transcranial electric stimulation on brain circuits in rats and humans. *Nature Communications* **9**, (2018).

104. Su, Y., Radman, T., Vaynshteyn, J., Parra, L. C. & Bikson, M. Effects of high-frequency
stimulation on epileptiform activity in vitro: ON/OFF control paradigm. *Epilepsia* **49**,
1586-1593, (2008).
105. Turner, D. A., Degan, S., Galeffi, F., Schmidt, S. & Peterchev, A. V. Rapid, Dose-
Dependent Enhancement of Cerebral Blood Flow by transcranial AC Stimulation in
Mouse. *Brain Stimul* **14**, 80-87, (2021).
106. Abbott, C. C. *et al.* Amplitude-determined seizure-threshold, electric field modeling, and
electroconvulsive therapy antidepressant and cognitive outcomes.
Neuropsychopharmacology **49**, 640-648, (2024).
1256