Microbial transcriptional responses to host diet maintain gut microbiome homeostasis in the

American cockroach

Amber N. DePoy[a], Helen E. Wall[a]*, Kara A. Tinker[a]*, Elizabeth A. Ottesen [a]#

[a] Department of Microbiology, University of Georgia, Athens, Georgia, USA

Running title: Cockroach hindgut microbial response to host diet

#Address correspondence to Elizabeth A. Ottesen, ottesen@uga.edu.

*Present address: Helen E. Wall, Center for Disease Control and Prevention, Atlanta, Georgia,

USA; Kara A. Tinker, National Energy Technology Laboratory (NETL), Pittsburgh, PA, USA

Helen E. Wall and Amber N. DePoy contributed equally to this work. Author order was

determined based on alphabetical order.

Abstract

Diet is considered a key determinant of gut microbiome composition and function.

However, studies in the American cockroach have revealed surprising stability in hindgut

microbiome taxonomic composition following shifts in host diet. To discover microbial activities

underlying this stability, we analyzed microbial community transcriptomes from hindguts of

cockroaches fed diverse diets. We used a taxon-centric approach in which we clustered genomes

based on taxonomic relatedness and functional similarity and examined the transcriptional

profiles of each cluster independently. In total, we analyzed a set of 18 such "genome clusters",

including key taxa within *Bacteroidota*, *Bacillota*, *Desulfobacterota,* and *Euryarcheaeota* phyla.

We found that microbial transcriptional responses to diet varied across diets and microbial

functional profiles, with the strongest transcriptional shifts seen in taxa predicted to be primarily

focused on degradation of complex dietary polysaccharides. These groups upregulated genes

associated with utilization of diet-sourced polysaccharides in response to bran and dog food

diets, while they upregulated genes for degradation of potentially host-derived polysaccharides in

response to tuna, butter, and starvation diets. In contrast, chemolithotrophic taxa, such as

Desulfobacterota and *Methanimicrococcus,* exhibited stable transcriptional profiles, suggesting

that compensatory changes in the metabolism of other microbial community members are

sufficient to support their activities across major dietary shifts. These results provide new insight

into microbial activities supporting gut microbiome stability in the face of variable diets in

omnivores.

Importance

The gut microbiome of omnivores must adjust to constantly changing dietary nutrient

compositions. This study describes gut microbial transcriptional responses to dietary shifts that

support gut microbiome homeostasis in omnivorous cockroaches. Our results provide insight into

the complex web of metabolic interactions mediating gut microbiome composition and stability.

We found a key role for polysaccharide-degrading microbes (including *Bacteroides* and

Dysgonomonas) in supporting the metabolic activity of other gut microbes through utilization of

diverse dietary polysaccharides or, in the absence of dietary polysaccharides, host-origin

polysaccharides. Further, the results show remarkable stability in gene expression across diets for

many gut microbes including secondary degraders of simple sugars, sulfate-reducing bacteria,

and methanogens, showing that the metabolism of these microbes is more dependent on the

metabolic products of primary degraders rather than dietary nutrients. This research provides

new insights into the complexity of microbe-microbe and host-microbe interactions underlying

gut microbiome homeostasis and dysbiosis.

Introduction

The gut microbiome shares an intimate relationship with its host, impacting digestion,

nutrition, immunity, and more (1). Given the influence of microbes on their host, gut microbiome

dysbiosis has been implicated in a wide range of diseases (2-7). In turn, the host can modify the

gut microbiome through a variety of pathways (8-12). Of these, diet has long been considered

one of the most direct drivers of gut microbiome composition (13-17). The fiber content of diets

is thought to play a particularly important role in shaping gut microbiome composition, with

many (18) studies reporting differences in the taxonomic and functional community profile of

mice (19-21), humans (22, 23), and other mammals (13, 24-26) fed Western diets vs. high-fiber

diets.

Despite the importance of diet in shaping the gut microbiome (23, 27, 28), short-term

dietary interventions in humans often fail to elicit shifts in gut microbiome composition greater

than pre-existing, individual-to-individual differences (27, 29, 30). Multiple long-term studies of

individual humans have shown stability over time (23, 27, 28, 31). Even dietary interventions

that have been shown to induce short-term responses in the gut microbiome have shown longer-

term reversion towards the native composition before or after cessation of the intervention (32-

34). Together, these results emphasize the importance of understanding microbe-microbe or

host-microbe interactions that may assist in the maintenance of gut microbiome homeostasis.

We are exploring gut microbiome homeostasis in response to host dietary shifts using an

omnivorous cockroach (*Periplaneta americana*) as a model system. The American cockroach

hindgut microbiome is highly diverse and includes many taxa from clades found in mammalian

and human guts (35-38). This gut microbiome also shows broad taxonomic and functional

similarities to mammalian omnivores, including abundant *Bacteroidota* and *Bacillota* with

taxonomic and functional profiles similar to their sister clades found in mammalian guts (39).

However, the ability of *Periplaneta americana* to survive for long periods of time on diverse and

nutrient-poor diets allows us to design studies using more extreme dietary shifts than allowable

in mammalian studies. Studies have found that, like humans, the cockroach gut microbiome

exhibits stability across highly divergent diets (35, 37, 40, 41). Therefore, we hypothesize that a

combination of host-microbe and microbe-microbe interactions support gut microbiome

homeostasis.

Here, we examined hindgut microbial community activity in cockroaches fed a control

diet of dog food as well as high-fiber, high-protein, high-fat, and starvation diets. A previous

study showed that these diets did not induce substantial, global changes in gut microbiome

composition (35). Using metagenomic and metatranscriptomic data, we identified taxonomic and

phylogenetic clusters of related microbes showing evidence of similar functional roles. We then

generated pangenomes for each cluster and analyzed their transcriptional responses to diet

independently. These results provide new insight into microbial activities underlying observed

compositional stability in the cockroach hindgut microbiome.

Results

We sequenced shotgun community DNA and shotgun community mRNA reads from 45

individual insects fed one of five distinct dietary conditions: bran, dog food (laboratory

maintenance diet), tuna, butter, and starvation. We obtained 25 metagenomes (5 from each diet)

and 20 metatranscriptomes (4 from each diet). Following quality-filtering, removal of host

DNA/RNA reads, and removal of remaining rRNA reads, we obtained 1,021,038,314

metagenome (870,360-533,070,442 per sample) and 78,022,090 metatranscriptome reads

(891,624-5,551,790 per sample). After read translation, 35.14% of metagenome and 62.44% of

metatranscriptome reads were mapped by homology search to protein sequences in RefSeq or

our single cell genomes from the cockroach gut (39).

Taxonomic profiles and overall community composition

Overall, the metagenomes and metatranscriptomes were dominated by *Bacteroidota*,

Bacillota (formerly *Firmicutes*), and *Desulfobacterota*. Alpha diversity (Shannon index) did not

vary significantly across the diets for the metagenomes ($p = 0.28$, $p_{adj} \geq 0.39$) nor

metatranscriptomes ($p = 0.04$, $p_{adj} > 0.08$) (Figure S1). However, we did observe large variability

in metagenome alpha diversity (Shannon index) within the butter diet (Figure S1).

PERMANOVA analysis of Bray-Curtis dissimilarities based on taxonomic profiles showed that

while diet exerted little effect over metagenome community composition ($p = 0.20$), it was a

significant driver of metatranscriptome composition ($p = 0.01$) (Figure 1A). Bran-fed

cockroaches had lower dissimilarity compared to other diets for each of the datasets, while

butter-fed cockroaches had the highest dissimilarity (Figure 1A). The high variability seen in

butter-fed cockroaches was likely associated with "blooms" in the abundance of individual taxa

among a subset of samples in both metagenome and metatranscriptome datasets (Figure 1).

Changes in microbial abundance and activity levels across diets

To evaluate how key taxa within the cockroach gut community respond to host dietary

shifts, we employed a taxon-centric strategy for metatranscriptomic analyses. Pangenomes were

constructed for eight genome clusters in *Bacteroidota*, seven in *Bacillota*, two in

Desulfobacterota, and one in *Euryarchaeota*. *Bacteroidota* genome clusters represent roughly

genus-level clusters of *Bacteroides, Parabacteroides*, *Dysgonomonas*, *Alistipes*, and

Odoribacter, as well as unclassified *Paludibacteraceae*, *Azobacteroidaceae*, and *Bacteroidales*.

Bacillota genome clusters often could not be built at genus level (42) but included three family-

level clusters (*Clostridiaceae*, *Lactobacillaceae*, and *Entercoccaceae*) where co-occurrence data

suggested a single population, while *Oscillospiraceae* and *Lachnospiraceae* were each split into

two sub-populations that showed distinct co-occurrence and transcriptional profiles (Figure S2-

S3); Other sub-populations identified within *Oscillospiraceae* and *Lachnospiraceae* did not have

enough sequence coverage in the datasets to be analyzed further. For *Desulfobacterota,* clusters

include *Frigididesulfovibrio* and *Desulfosarcina*. *Methanimicrococcus* was the only methanogen

with transcriptional abundance high enough to generate a genome cluster. See Supplemental

Table 1 for a full list of genome references used to build genome clusters.

Many individual genome clusters showed little change in average metagenome relative

abundance between diet groups (Figure 2). However, *Dysgonomonas*, *Lactobacillaceae*, and

Enterococcaceae, exhibited higher average metagenome relative abundance in the butter diet,

much of which is attributable to "blooms" of different taxa in a subset of samples (Figures 1B,

2). In contrast, the fraction of metatranscriptome reads assigned to individual genome clusters

was more variable. *Bacteroides* had increases in relative abundance of transcripts for the bran

diet while *Lachnospiraceae_2* and *Clostridiaceae* had more transcription in the dog food diet.

Odoribacter, unclassified *Bacteroidales*, and *Desulfosarcina* had more transcripts in the butter

diet, while *Desulfosarcina* was very transcriptionally active in the tuna diet. Some groups, such

as *Parabacteroides* and *Frigididesulfovibrio*, showed little to no change in abundance in the

metagenome and metatranscriptome between diets (Figure 2).

High-level trends in gene expression of key metabolic pathways

Figure 3 shows expression levels of genes assigned to key functional categories in

different diets for individual taxonomic clusters. These functional profiles underscore established

patterns of gene expression for these groups, with carbohydrate-active enzymes (CAZymes)

expressed by nearly all groups within *Bacteroidota* and *Bacillota* phyla, with exception of the

Bacteroidales taxonomic cluster (Figure 3). *Bacteroides*, *Dysgonomonas*, and *Parabacteroides*

showed some of the highest transcription of CAZymes (Figure 3) and are hypothesized to be

primary degraders of polysaccharides (39). Beyond the primary degraders, other taxa transcribed

CAZymes ($\geq$ 0.1% average relative abundance) and the difference in transcription varied by diet

(Figure S4). Many of these taxa had more transcription of CAZymes in the bran and dog food

diets, but the CAZyme type and specificity varied among taxa (Figure S4). Even in diets without

abundant carbohydrates, there were CAZymes transcribed at high levels. For many taxa,

particularly–*Bacteroidota*—CAZymes targeting eukaryotic components (GH109, GT41,

CBM51) and mannose (GH92, GH38) were transcribed more during host starvation (Figure S4).

Not all *Bacteroidota* appeared to specialize in fiber degradation. *Odoribacter* and a group

of unclassified *Bacteroidales* both transcribed genes related to carbon fixation at high levels

(Figure 3) but showed differences in the specific carbon fixation genes utilized. *Odoribacter*

primarily transcribed genes involved in converting acetyl-CoA and CO_2 to pyruvate (K14534

and K03737) (Figure S5A). Due to higher transcription of both genes, this implicates a role in

the dicarboxylate-hydroxybutyrate cycle of carbon fixation. *Bacteroidales* primarily transcribed

genes for the conversion of acetyl-CoA and CO_2 to phosphoenol-pyruvate (K03737, K01006)

but also highly transcribed genes to convert succinyl-CoA and CO_2 to 2-oxoglutarate (K00174,

K00175, and K00177) (Figure S5B). These are metabolic steps in both the dicarboxylate-

hydroxybutyrate cycle and the reductive citrate cycle. However, *Bacteroidales* was not

transcribing several genes in the dicarboxylate-hydroxybutyrate pathway that *Odoribacter*

transcribed, suggesting that *Bacteroidales* utilizes these genes in the reductive citrate cycle

(Figure S5).

168   While *Bacteroidota* and *Bacillota* groups transcribed CAZymes at high levels, they

differed in nutrient acquisition strategies; *Bacteroidota* typically transcribed TonB/Sus type

transporter systems and *Bacillota* primarily transcribed ABC-type transporters (Figure 3). In

Bacillota, we observed diet- and taxon-driven differences in ABC transporter transcription

(Figure 3). For example, *Lactobacillaceae* transcribed ABC transporters to a greater extent in the

butter diet and less so in the tuna diet (Figure 3). For this group, only a single ATP-binding

cassette (K20344, a subfamily C bacteriocin exporter), was upregulated in the tuna diet while the

butter diet induced upregulation of diverse ABC transporters (Figure S6). Finally,

Lactobacillaceae responded to host starvation by upregulating ABC transporters for nucleosides,

ribose/D-Xylose, lincosamide, and streptogramin A (Figure S6).

178   Chemolithotrophic groups, including sulfate reducers and methanogens, were also highly

transcriptionally active (Figure 3). Although transcription of sulfate reduction genes was largely

not significantly changing in response to host diet, *Desulfosarcina* and *Frigididesulfovibrio*

showed trends in transcription of sulfur metabolism genes. For the butter diet, *Desulfosarcina*

had reduced transcription of some sulfur metabolism genes compared to the other diets (Figure

S7). Conversely, *Frigididesulfovibrio* increased transcription of some sulfur metabolism genes in

cockroaches fed butter, particularly genes involved in the conversion of PAPS to APS (K01082

and K06881), an anaerobic sulfite reductase (K16951), and an anaerobic dimethyl sulfoxide

reductase (K07306) (Figure S7B). Methane metabolism genes for *Methanimicrococcus* did not

change gene expression significantly in response to host diet shift. Although not significant,

some methane metabolism genes were transcribed more in the bran and tuna diets (Figure S8).

Additionally, there were transcriptional differences that suggest changes in preferred substrates

between methanol and methylated amines depending on host diet (Figure S8).

Nearly all taxa increased transcription of genes associated with stress responses in the

butter diet, evident by increased transcription of chaperones and folding catalysts (Figure 3).

Specifically, DnaJK chaperones, heat-shock proteins, peroxiredoxin, and thioredoxin were all

differentially expressed in the butter diet relative to all other diets (Figure S9). Taxa that showed

significant differential expression of these stress transcripts included *Bacteroides*,

Dysgonomonas, *Parabacteroides*, *Alistipes*, *Paludibacteraceae*, *Lactobacillaceae*, and

*Lachnospiraceae*_1. The most consistent trend of increased transcription in these stress response

genes was for DnaK (K04043) and chaperonin GroEL (K04077) in the butter diet (Figure S9).

Despite a strong stress response seen in many taxonomic clusters, there was no decrease in the

relative abundance of these groups in the metagenomes (Figure 2). However, some taxa

exhibited increased transcriptional activity in the butter diet (as measured by relative abundance

in the metatranscriptome), suggesting that they were more active under this condition and/or

were less impacted by this stress than other taxa (Figure 2).

Some taxa also exhibited high transcription of stress-related genes in diets other than

butter. For example, *Dysgonomonas* transcribed thioredoxin (K03671) and NADH-dependent

peroxiredoxin subunit C (K24119) to a greater extent in starvation (Figure S9).

*Lachnospiraceae*_1 exhibited greater transcription of NADH-dependent peroxiredoxin subunit C

(K24119) in the tuna diet than in any of the other diets (Figure S9). *Clostridiaceae* and

*Lachnospiraceae*_2 transcribed chaperoin GroEL (K04077) and DnaK (K04043) at higher levels

than other taxa in all diets, not only the butter diet.

Significant gene expression changes in response to host diet shift

For each genome cluster, we analyzed genes that displayed significantly altered

expression across diets using DESeq2 (43). Across all taxa and all pairwise comparisons of diets,

a total of 1215 genes were identified as significantly changing with respect to host diet (Figure

4). However, there were striking differences in the number of significantly differentially

expressed genes across taxa. Those previously identified as potential primary degraders (39)

showed stronger transcriptional responses to diet than other organisms. Additionally, taxa varied

in which diets elucidated the strongest responses, with some organisms responding most strongly

to bran, while others responded most to tuna or butter diets (Figure 4).

Significant gene expression for Bacteroidota taxa

Members of the *Bacteroidota* phylum had the greatest number of significantly changing

genes, with *Bacteroides* (520 significant genes) and *Dysgonomonas* (198 significant genes)

having far more than other taxa (Figure S10). *Bacteroides* had the most changes in gene

expression when comparing bran to tuna, butter, and starvation (Figure 4). In contrast to

Bacteroides, *Dysgonomonas*, and *Parabacteroides*, other *Bacteroidota* taxa (*Alistipes*,

Azobacteroidaceae, *Odoribacter*, *Paludibacteraceae*, and *Bacteroidales*) show substantially

fewer differences in gene expression across diets despite having high read counts (Figure S10,

Figure 4).

Bacteroides showed a particularly strong response in gene expression between the bran

and other diets (Figure 4). When comparing gene expression between the tuna and bran diet,

Bacteroides differentially expressed 127 genes, 16 of which were upregulated in the bran diet

(Figure 4). Bran-responsive genes included four CAZyme genes: two that degrade diverse

substrates (GH3, GH32) and two that degrade xylan (CE1, GH10) (Table S2). There were four

transporters upregulated in the bran diet, including a putative fructose porter, SusD starch

binding proteins, and SusD homologs. Other bran upregulated genes included a fructokinase,

glycan binding proteins, TolC family proteins, and RagB/SusD family nutrient uptake proteins.

The 20 CAZymes upregulated in the tuna (vs. bran) diet included CAZymes specific for pectin

(CE8, PL1_2, PL1, GH28, GH105), chitin or peptidoglycan (CBM50), rhamnogalacturonan

(PL11_1, GH106), and diverse substrates (GH43_4, GH43_10, GH2, CE12). Other genes

upregulated in cockroaches fed tuna (vs. bran) included FAD-dependent oxidoreductase, NADH-

dependent peroxiredoxin subunit C, fimbrillin family protein, and Mfa1 family fimbria major

subunit (Table S2). When comparing bran and butter gene expression patterns, *Bacteroides*

differentially expressed 119 genes, 40 of which were upregulated in the bran diet (Figure 4). The

same four CAZymes upregulated in bran (vs. tuna) were also upregulated in response to butter

(vs. bran), with the addition of three CAZymes that degrade diverse substrates (CE20, GH43_1,

and GH67). Other genes upregulated in the butter diet (vs. bran) included chaperonin GroEL

(*groL*), a putative lipoprotein, a possible iron receptor (RagA), the chloroplast envelope TIC

translocase complex, and NADH-dependent peroxiredoxin subunit C. During host starvation,

Bacteroides differentially expressed 122 genes compared to the bran diet. CAZymes that were

upregulated in the bran diet (vs. starvation) were similar to those seen for other diets, including

GH32 (diverse substrates), CE1 (xylan), GH10 (xylan), GH67 (alpha-glucuronidase), and

GH43_24 (Exo-β-1,3-galactanase). Some upregulated genes in starvation (vs. bran) included

FAD-dependent oxidoreductase, possible iron receptor (RagA), and NADH-dependent

peroxiredoxin subunit C. There were no CAZyme genes that were significantly upregulated

during host starvation (vs. bran). However, CAZymes that showed increased, if not significantly

so, expression included those potentially involved in breakdown of host mucins, including

GH109 and mannosidases (Figure S4) (44). In addition, a single gene, xylulokinase was

upregulated in starvation when compared to the butter diet.

Dysgonomonas and *Parabacteroides* showed fewer significantly differentially expressed

genes, but overall exhibited similar responses (Figure 4). Like *Bacteroides*, *Dysgonomonas*

upregulated CAZymes targeting xylan, pectin, and diverse substrates in the bran diet (Table S3).

Dysgonomonas also upregulated SusC and SusD transporters, L-arabinose isomerase, xylose

isomerase, and L-ribulokinase in the bran diet (vs. butter). During starvation (vs. bran),

Dysgonomonas upregulated NADH-dependent peroxiredoxin subunit F, DNA-binding ferritin-

like protein (oxidative damage protectant), and a fatty acid repression mutant protein (predicted

oxidoreductase). *Dysgonomonas* responded to the tuna diet (vs. bran) with pectin CAZymes

(GH28, PL1_2, CE8). *Parabacteroides* responded to the bran diet with upregulation of

CAZymes that target diverse substrates (Table S4). Other genes that were upregulated by

Parabacteroides in the bran diet included SusC and SusD transporters (Table S4). Unlike

Bacteroides and *Dysgonomonas*, *Parabacteroides* did not upregulate pectin CAZymes in

response to the tuna diet (Table S4). In response to the tuna diet, *Parabacteroides* instead

upregulated SusC and SusD transporters and other related starch binding outer membrane

protein, SusD/RagB family. The butter diet triggered upregulation of GroEL (*groL*) and DnaK

genes (Table S4). In addition, *Parabacteroides* only upregulated three genes during host

starvation (vs. bran): NADH-dependent peroxiredoxin subunit F, cytochrome bd ubiquinol

oxidase subunit I, and cytochrome bd ubiquinol oxidase subunit II (Table S4).

Alistipes had far fewer differentially expressed genes as compared to other members of

Bacteroidota, largely changing gene expression in response to the butter diet (Figure 4). Many of

the genes that were upregulated in the butter diet were stress response genes, including

chaperonins and heat-shock related proteins. Six genes were upregulated in the dog food diet (vs.

butter). Functions of these genes included outer membrane protein OmpA, Mfa1 family fimbria

major subunit, and thioredoxin-dependent peroxiredoxin. Two other taxa of *Bacteroidota*,

Azobacteroidaceae and *Paludibacteraceae*, differentially expressed genes in the bran to butter

comparison (Figure 4). *Azobacteroidaceae* upregulated a putative arabinose porter and putative

outer membrane porin in the bran diet (vs. butter) (Table S5). Some of the other genes

upregulated in the bran diet included c-type cytochrome, IS982 family transposase, and L-

arabinose isomerase. As was observed in many organisms, DnaK genes were upregulated in the

butter diet compared to bran (Table S5). In addition, *Azobacteroidaceae* upregulated a

hypothetical protein, T9SS type A sorting domain-containing protein, and an arabinose

isomerase in the dog food diet (vs. butter). *Paludibacteraceae* differentially expressed 15 genes

in the comparison of bran and butter diets (Figure 4), and 10 of these genes were upregulated in

the bran diet (vs. butter). Some bran upregulated genes included elongation factor Tu, 50S

ribosomal protein L11 (rplK), and 30S ribosomal protein S11 (rpsK) (Table S6). There was also

upregulation of xylulokinase, RagB/SusD family nutrient uptake outer membrane protein, and

ribulokinase in the bran diet (vs. butter). Butter upregulated genes included chloroplast envelope

TIC translocase (TIC) complex, molecular chaperone protein (Hsp90 homologue), and HSP20

family protein, presented in highest to lowest transcription (Table S6). *Paludibacteraceae*

upregulated a single gene for a glucoronate isomerase in the dog food diet (vs. butter).

Odoribacter upregulated fewer genes, and only in the bran vs. butter and tuna vs. butter

comparisons (Figure 4). Of these 19 upregulated genes, four were upregulated in the butter diet

(vs. tuna), all of which were transporters; these transporters included an outer membrane protein

transport protein, multidrug efflux pump, Triclosan resistance efflux pump, and TolC family

protein. All genes that *Odoribacter* upregulated in the tuna and bran diet (vs. butter) included

H^+-translocating ATPases (Table S7). *Bacteroidales* only had a single gene, a two-sector V-type

ATPase or ATP synthase, that was upregulated in the butter diet (vs. tuna).

Significant gene expression for Bacillota taxa

Among the *Bacillota*, *Oscillospriaceae_*2 also showed a similar pattern of gene

expression changes when comparing bran to tuna, butter, and starvation, suggesting a strong

response to diets with a high polysaccharide content (Figure 4). However, the types of genes

upregulated by *Oscillospiraceae_*2 in response to these diets were different from those observed

in *Bacteroides*. When comparing tuna to bran, *Oscillospiraceae_*2 differentially expressed 26

genes, of which 15 genes were upregulated in the bran diet. None of these bran upregulated

genes were CAZymes but instead included ABC transporters for arabinose, arabinose isomerase

genes, and electron transporters involved in sulfur metabolism with anaerobic dimethyl sulfoxide

reductase (Pyrogallol hydroxytransferase) (Table S8). Genes that were upregulated in the tuna

diet included ferrous iron transporter B, amidohydrolase, and aminobenzoyl-glutamate utilization

protein B. The latter two genes are involved in lysine biosynthesis.

Lactobacillaceae (*Bacillota*) had 154 genes that were significantly changing with diet.

Interestingly, many of these differences in gene expression were identified in comparisons

between tuna and all other diets, suggesting a strong response to a high-protein diet (Figure 3).

Lactobacillaceae upregulated four CAZymes consistently in all diets compared to tuna,

including CBM50 (chitin or peptidoglycan), CE9 (N-acetylglucosamine-6-phosphate

deacetylase), GH1 (beta-glucosidases and beta-galactosidases), and GT2 (diverse substrates).

Only seven genes were upregulated in the tuna diet (vs. starvation), including

maltose/maltotriose/maltodextrin (up to 7 glucose units) transporters, NADP-dependent

phosphogluconate dehydrogenase, septation ring formation regulator, 3-phenylpropionate/trans-

cinnamate dioxygenase ferredoxin reductase component, Lipoprotein-anchoring transpeptidase,

and 6-phospho-beta-glucosidase. Many of the same genes were upregulated in the bran, butter,

and starvation diet when compared to tuna. Genes that were consistently upregulated in diets

other than tuna included formate C-acetyltransferase, phosphoenolpyruvate-protein

phosphotransferase (PTS system enzyme I), pyruvate kinase, and putative sigma-54 modulation

protein (Table S9).

*Lachnospiraceae*_1 (*Bacillota*) also had a strong response to the tuna diet, but only when

compared to the bran diet (Figure 4). There were 34 differentially expressed genes in this

comparison, 12 of which were upregulated in the bran diet. Bran upregulated genes included a

putative multiple sugar transport system, flagellin, and glycine/betaine/sarcosine/D-proline

family reductase selenoprotein B. Tuna upregulated genes (vs. bran) included those for

peptide/nickel transport system substrate-binding protein, IS21-like element helper ATPase and

IS21 family transposase (Table S10). *Lachnospiraceae*_1 only upregulated a single CAZyme,

which has N-acetylglucosamine-6-phophate deacetylase activity (CE9), for the tuna diet (vs.

bran).

Significant gene expression of chemolithotrophs

The two sulfate reducing taxa, *Frigididesulfovibrio* and *Desulfosarcina*, did not respond

strongly to host diet. *Desulfosarcina* did not differentially express any genes across the diets.

Frigididesulfovibrio differentially expressed only 17 genes when comparing bran to butter

(Figure 4), 15 of which were upregulated in the bran diet (vs. butter). Four of these bran-

upregulated genes were transporters including a putative porin, a dicarboxylate (succinate,

fumarate, malate) transporter, Na^+-dependent C4-dicarboxylate (fumarate, succinate) uptake

transporter, and a 2,3-diketo-L-gulonate (2,3-DKG) transporter (Table S11). Other bran-

upregulated genes included several related to choline metabolism, including five BCCT family

transporters and choline TMA-lyase-activating enzyme. The two genes that were upregulated in

the butter diet (vs. bran) were for a multidrug efflux pump subunit AcrA (membrane-fusion

protein) and opacity protein LomR and related surface antigens. In the tuna diet (vs. butter),

several genes related to histidine degradation were upregulated, including putative histidine

porter and histidine ammonia-lyase. Further, *Fridgididesulfovibrio* upregulated two subunits of

the 50S ribosome in both bran and tuna diets compared to butter.

Methanimicrococcus differentially expressed very few genes, with most gene expression

changes identified in the bran vs. dog food comparison (Figure 4). Many of these differentially

expressed genes were hypothetical proteins. *Methanimicrococcus* also showed a response

between bran and butter diet with four differentially expressed genes. Three of these genes were

upregulated in the butter diet (vs. bran) including methanol:5-hydroxybenzimidazolylcobamide

Co-methyltransferase, ethanolamine permease, and peroxiredoxin (Table S12).

Discussion

The aim of this study was to utilize metagenomic and metatranscriptomic techniques to

better understand how microbial transcriptional responses support the strong stability in

taxonomic composition of the cockroach gut microbiome following dietary shifts. Although we

observed somewhat higher variability in taxonomic composition across samples in the

metagenome dataset (Figure 1B), the metagenome results support the key observation of the

previous study that diet was not a strong driver of gut microbiome community composition (35).

In contrast, metatranscriptome composition did exhibit changes in microbial abundance and

transcript abundance in responses to dietary shifts. As a result, we focused primarily on analysis

of transcriptional responses to diet that may support this stability.

We used a taxon-centric, pangenome approach to generate a composite transcriptome for

clusters of closely related genomes. This approach was implemented to take maximal advantage

of available reference genomes—including highly fragmented, host-relevant, single-cell

sequenced genomes (39)—while avoiding having to select a single reference genome that may

not perfectly match the unclassified dominant organisms in the sample. A similar approach has

previously been used for analysis of transcriptional patterns among marine microorganisms (45-

47). The set of organisms selected for our analysis included *Bacteroides*, *Dysgonomonas*,

Parabacteroides, *Alistipes*, unclassified *Azobacteroidaceae*, *Paludibacteraceae*, *Bacteroidales*,

Clostridiaceae, *Lactobacillaceae*, *Enterococcaceae*, *Lachnospiraceae*, *Oscillospiraceae*;

Frigididesulfovibrio; *Desulfosarcina*; and *Methanimicrococcus*. These organisms were selected

because their high relative abundance in the metatranscriptome. They represent a variety of

putative trophic groups (48), including primary polysaccharide degraders, secondary degraders

of polysaccharolytic byproducts, and chemolithotrophs including acetogens, sulfate reducers, and

methanogens.

Through this work we have expanded what is known about a wide range of

microorganisms' transcriptional response to host dietary shift. In general, changes in the relative

abundance of individual taxa in the metatranscriptome, which is assumed to represent their

overall transcriptional activity, broadly reflected hypothesized substrate specializations. Taxa

that have previously been hypothesized to act as primary degraders, including *Bacteroides*,

Dysgonomonas, and *Parabacteroides*, showed higher transcriptional activity in cockroaches fed

bran and dog food diets (Figure 2) (49-52). Conversely, other members of *Bacteroidota* had

higher transcriptional activity in the butter and/or tuna diets (Figure 2). Members of *Bacillota*

varied in which diets they were most active in, with some more active in cockroaches fed butter,

tuna, or dog food. Both *Desulfosarcina* and *Frigididesulfovibrio* exhibited reduced activity in the

bran diet, whereas *Methanimicrococcus* were more active in the bran fed cockroaches (Figure 2).

Microbial responses to high-polysaccharide diets (bran and dog food)

The microbial community of bran-fed cockroaches had the most stable taxonomic

community structure in both metagenome and metatranscriptome datasets (Figure 1A).

Bacteroides had the highest transcriptional activity in the bran diet followed by *Dysgonomonas*

and *Parabacteroides* (Figure 2). All three of these taxa transcribed a wide range of genes

responsible for the degradation of complex carbohydrates, including genes containing inulin- and

xylan-targeting CAZyme domains that are upregulated in cockroaches eating bran or dog food.

This activity supports findings from *Bacteroides* spp. from other environments that release large

amounts of polysaccharide degradation products extracellularly, providing the potential for these

primary degraders to create an "everybody wins" scenario in terms of cross-feeding relationships

(53, 54). Previous studies have implicated *Bacteroides* in cross-feeding of fermentation products

with secondary degraders, sulfate reducers, and methanogens (54-58). Given the potential for

cross-feeding relationships, gains in growth and carbohydrate degradation among the primary

degraders may simultaneously promote growth of a wide range of other organisms, thereby

minimizing compositional changes in gut microbiome composition, as observed here. Further

studies are needed to assess specific cross-feeding interactions that occur between these primary

degraders and other microbiome members in the cockroach hindgut.

Some *Bacillota* members also showed strong transcriptional responses to bran. Rather

than breaking down primary carbohydrates, their transcriptional patterns suggest that these

organisms are making use of by-products from primary polysaccharide degradation. *Bacillota*

using simple sugars released from primary degraders has been shown in *Roseburia*

(*Lachnospiraceae*) (59), while aldouronate (hemicellulose depolymerization product) is

commonly metabolized by Gram-positive members of *Clostridiaceae* and *Oscillospiraceae* (60).

In our data (Figure 3), *Bacillota* members such as *Lactobacillaceae*, *Lachnospiraceae*_1,

*Oscillospiraceae*_1, and *Oscillospiraceae*_2 responded to a bran diet by upregulating ABC

transporters that target mono- and short chain oligosaccharides such as maltose, oligosaccharide

(glucuronate-linked to a xylo-oligosaccharide), ribose, and arabinose, suggesting that they are

primarily acting as secondary degraders. Other *Bacillota* show evidence of amino acid utilization

in the bran diet. *Lachnospiraceae*_1 and *Oscillospiraceae*_2 upregulated glycine

betaine/sarcosine reductase and transporters associated with Stickland metabolism. Related

organism *Closridium sporogenes* has been shown to reduce amino acids using the sugar

fermentation by-product H_2 (61-63).

Methanimicrococcus transcribed various genes related to methyl-reducing

methanogenesis at higher levels in the bran diet (Figure S6), although many of these genes were

not significantly upregulated. Previous studies have shown that *Methanimicrococcus* can utilize

diverse methyl-containing organic compounds such as methylamines and methanol (39, 63, 64).

Degradation of plant-sourced phosphatidylcholine by *Bacteroidota* and *Desulfobacterota* has

been shown to release methylamines (65, 66), and methanol is associated with degradation of

pectin (65, 66). In rumen isolates, consumption of these byproducts increased energy yield and

fiber degradation of other fiber degraders in the gut (67, 68). Although not significantly

differentially expressed, genes for the coenzyme 420 hydrogenanse (Frh) are transcribed more in

the bran diet (Figure S8), suggesting that the activity of methanogens is closely tied to the

activity of fiber degraders in the cockroach. *M. blatticola* lacks many enzymes of the methyl-

branch of the Wood-Ljungdahl pathway. Therefore, any electrons transferred to Frh would be

used for anabolism (69). This suggests a crucial role for methanogens in maintaining the low

partial pressure of H_2 and utilizing fermentation byproducts that may impact fermentation

activity in the gut. It also demonstrates how cross-feeding relationships may act to stabilize gut

taxonomic composition.

446   In many cases, organisms that respond strongly to bran also respond to dog food, likely

because it has the second highest carbohydrate content of the tested diets. Taxa that responded

strongly to the dog food diet included *Dysgonomonas*, *Bacteroides*, *Lactobacillaceae*, and

Parabacteroides, listed in highest to lowest number of dog food upregulated genes. Other

organisms responding to dog food appear to be interacting with the amino acid content of this

food including *Paludibacteraceae* and *Azobacteroidaceae*.

Microbial responses to a high-protein (tuna) diet

In general, the tuna diet elicited a strong response to protein components by *Bacillota,*

and *Frigididesulfovibrio*. Genes involved in amino acid degradation were upregulated in the tuna

diet by *Lactobacillaceae*, *Lachnospiraceae*_1, *Oscillospiraceae*_1, *Oscillospiraceae*_2, and

Frigididesulfovibrio. Proteolysis in the human gut is also largely attributed to *Bacillota*

members, including activity on aromatic amino acids (70-72). *Lactobacillaceae* upregulated

several genes involved in aromatic amino acid metabolism. *Frigididesulfovibrio* also responded

to the tuna diet with upregulation of several genes for histidine metabolism. Although their

primary metabolism remains unchanged by host diet, this may reflect an ability by this sulfate

reducer to take advantage of additional electron donor substrates if present.

In contrast to the tuna diet response by *Bacillota* and *Frigididesulfovibrio*, *Bacteroides*

and *Dysgonomonas* responded with transcriptional patterns suggesting utilization of

polysaccharides in tuna. We did not expect polysaccharide degraders to engage extensively with

this diet. However, there was strong upregulation of genes associated with pectin degradation by

various primary and secondary degraders. We hypothesize this CAZyme activity is related to

utilization of galacturonan, which—in addition to being a key component of pectin–is known to

be present in tuna and tuna by-products (73). In addition, most canned tuna products contain

vegetable broth as a flavor enhancer, which may also contribute to the pectin activity that we

observed. Even in a diet very low in polysaccharides, such as tuna, these primary degraders

identify and scavenge relevant substrates for their specialized metabolisms.

Microbial responses to a high-fat (butter) diet

Microbial responses to the butter diet represent one of the major surprises of this work.

There was wide-spread transcriptional upregulation of stress response proteins including RND

efflux pumps, oxidative stress, and heat shock proteins (Figure 3, Figure S7). Across many

organisms, this stress response was associated with upregulation of *groEL* and *dnaK* (Figure 3,

Figure S7). These two genes are known to allow bacteria to survive better in adverse conditions

and are involved in adhesion and invasion processes (74, 75). GroEL is the bacterial counterpart

of mammalian heat shock protein 60 and plays an essential role in bacterial survival during

chemical, heat, or oxidative stress. DnaK is a chaperone that is part of the heat shock protein

family and involved in stress response (76). Previous studies have shown that a high-fat diet

induces oxidative stress in the host, which may be the reason we see a wide-spread upregulation

of stress genes in the microbiome (77). A similar stress response to changes in redox state that

has been noted in gut epithelial cells in mice on high-fat diets (78, 79). The butter diet in the

cockroach also appears to induce lower gut microbiome stability (Figure 1). Instability has not

been explicitly connected to high-fat diets, but studies in mice eating high-fat diets have

indicated blooms of organisms (19, 20, 80). A study of human diet and microbiome composition

also indicated greater variability in community profiles of individuals eating diets high in fats

(81). Together, these results support findings that high fat diets may have profound impacts on

the gut microbiome.

In addition to the observed stress response, *Bacteroides* and *Dysgonomonas* respond to

the butter diet by transcribing CAZyme domains associated with degradation of GlcNAc and

mannose in host tissue (Figure S2). These transcripts are also upregulated in response to

starvation, which may reflect degradation of host mucins from the peritrophic membrane; this

starvation response is discussed in more detail below. In addition, *Lactobacillaceae* upregulate

genes with CAZyme domains that target diverse substrates. Despite primary polysaccharide

degraders attempting to scavenge appropriate substrates here, much of their transcriptional

activity is reduced, with remaining activity redirected towards survival in suboptimal conditions.

Despite this reduced transcriptional activity in diets low in polysaccharides, primary degraders

may avoid dropping in relative abundance in the metagenome because other taxa are

experiencing similar stresses (82, 83).

In contrast, *Enterococcaceae* and *Lactobacillaceae* exhibited blooms in a subset of

cockroaches eating butter (Figure 1), and both taxa had increased relative abundance in the

transcriptome (Figure 2). *Enterococcaceae* and *Lactobacillaceae* spp. have been found in

association with high-fat diets in mice (84-86). Although *Enterococcaceae* did not significantly

upregulate any genes across the diets, *Enterococcus faecalis* is known for efficient glycerol

metabolism (87, 88). This ability may provide *Enterococcaceae* an advantage in cockroaches

eating butter. Furthermore, *Enterococcus* species are known to produce reactive oxygen species

that damage host epithelial cells (89, 90). Oxygenation in the gut produces a colonization benefit

for *Lactobacillus* in humans (91), as they are equipped to handle oxidative stress (92).

Enterococcus activities may cause the observed stress response from microbes throughout the

hindgut community, allowing *Lactobacillaceae* to take advantage of changing environmental

conditions in the gut (89, 90).

Microbial responses to host starvation

Bacteroides, *Parabacteroides*, and *Dysgonomonas* responded to host starvation (and to a

lesser extent butter) by upregulating GH109, a CAZyme that targets GlcNAc (93). *Bacteroides*

and *Dysgonomonas* also transcribe gene clusters with mannose-targeting CAZyme domains

(GH92, GH76). Although the hindgut of insects is surrounded by a chitinous ectoderm and lacks

a mammalian-like mucus layer, the midgut of insects produces a peritrophic membrane that is

continually extruded into the hindgut and may serve as a source of polysaccharides and glycans

(94). From work in the Colorado potato beetle, glycosylation of the peritrophic membrane

includes mannose and GlcNAc, especially the structures $GlcNAc_2Man_3$, $GlcNAc_2Man_3Fuc$ and

$GlcNAc_2Man_5$ (44). *Dysgonomonas* also highly expresses gene clusters with mannose-specific

CAZyme domains in low-carbohydrate diets and indicates lower expression of these genes in

carbohydrate-rich diets (Table S3). This transcriptional adjustment likely allows them to use

host-sourced polysaccharides in a diet short on complex carbohydrates. Studies in mammals

have widely identified host mucins as an important substrate for gut fiber degraders in the

absence of microbially accessible polysaccharides (95). These results suggests that the

peritrophic membrane in cockroaches may serve a similar role in the ecology of the cockroach

gut microbiome.

Microbial transcriptional stability across diets

While we focus on transcripts whose expression changed significantly across the diets, it

is worth noting the overall stability of gene expression in many of these microbes. While primary

degraders such as *Bacteroides*, *Dysgonomonas*, and *Parabacteroides* showed large number of

significantly changing genes, many key taxa had few or even no transcripts that were found to be

significantly different across diets or were responsive to only one diet (Figure 4). While the

ability to detect significantly changing transcripts is somewhat dependent on transcriptome

coverage, multiple groups with few or no diet-responsive transcripts, such as *Desulfosarcina* and

Frigididesulfovibrio, did so despite transcriptome coverage similar to or greater than groups with

larger numbers of significantly changing transcripts. These results suggest that they are not

directly dependent on dietary components to support their metabolism and are supported by other

host or microbial activities. Given the extensive shifts in metabolic activities observed in key

primary degraders, and the likely position of less diet-responsive microbes at the 'bottom' of the

microbial food web, this points towards a scenario in which the actions of these primary

degraders are sufficient to compensate for dietary changes and in turn support the metabolism of

microbes further down the food chain. This ability is likely not infinite, which could in turn

explain why polysaccharide-rich diets are associated with the greatest overall compositional and

transcriptional stability. Together, these findings support a 'keystone taxa' role for fiber

degraders in structuring gut microbiome metabolism.

Conclusions

The primary goal of this work was to gain insight into microbial transcriptional activities that

may explain the remarkable compositional stability of the cockroach gut microbiome following

substantial dietary shifts. A key finding of this work suggests the activity of polysaccharide

degraders—including *Bacteroides* and *Dysgonomonas*—is highly responsive to diet and plays a

key role in supporting gut microbiome homeostasis. These organisms are highly versatile,

scavenging polysaccharides from even low-fiber foods such as tuna, and using components of

the peritrophic matrix under starvation conditions and on low-carbohydrate butter diets. Higher

fiber diets result in lower sample-to-sample variability in gut microbiome composition and

transcriptional activity. Secondary degraders like *Bacillota* appear to adapt to available polymer

degradation products—such as oligosaccharides and oligopeptides—depending on the level of

fiber and protein in the host's diet. In contrast, a high-fat diet is associated with upregulation of

stress proteins across many gut microbes. This stress was associated with reduced gut

microbiome stability, with individual samples showing large-scale blooms of different taxa. This

finding aligns well with literature concluding that the 'Western diet' rich in fats and simple

sugars can lead to gut microbiome dysbiosis in many organisms (18, 24). Overall, these results

suggest that the omnivorous cockroach is a promising model for disentangling microbial

responses to diet. In addition, it suggests that taxon-centric analysis of metatranscriptomes is

highly promising for understanding the complex web of microbe-microbe and host-microbe

interactions underlying gut microbiome homeostasis and dysbiosis.

Materials and Methods

Diet manipulation and sample collection

Laboratory colonies of *Periplaneta americana* are maintained in mixed age, mixed sex

colonies in aquarium tanks at room temperature on a diet of dog food *ad libitum*. Each tank is

provided with corn cob bedding, cardboard tubes for nesting, and a cellulose sponge saturated

with water. Each dietary treatment group was housed in a single plastic tank that contained

pebbles for bedding, a large weigh boat for shelter, and food and water in weigh boats. Food,

water, and weigh boats were changed as needed, and any ootheca or deceased cockroaches were

removed daily. Treatments included dog food (Kroger Nutritionally Complete Bite Size Adult

Dog Food; composed of 21% protein, 8% fat, and 6% fiber), bran (Bob's Red Mill Organic High

Fiber Oat Bran Hot Cereal), butter (Kroger Unsalted Butter Sticks), tuna (StarKist Selects Low

Sodium Chunk Light Tuna in Water), and starvation (Table 1).

After 14 days on the assigned diet, all cockroaches were sacrificed and their hindgut

microbiota preserved for DNA or RNA extraction. Individual cockroaches were removed from

tanks, weighed, and placed on ice in sterile culture plates. Once sufficiently torpid, cockroaches

were dissected, and the entire gut was removed. Any visible debris, including fat bodies or

exoskeleton, was removed with forceps. The hindgut was then separated from the rest of the gut

using a scalpel and submerged in either 100 μL of 1XTE buffer or RNALater (Ambion, Austin,

TX, USA). Hindgut samples placed in 1XTE buffer were immediately stored at -80°C. Hindgut

samples placed in RNALater were crushed with a sterile glass stirring rod. The suspended gut

lumen was then removed and stored at -80°C.

DNA and RNA extraction, sequencing library preparation

Microbial DNA and RNA was extracted from hindgut samples as outlined previously in

(39). Briefly, DNA was extracted using a modified version of the EZNA Bacteria Kit (Omega

Biotek, Norcross, GA). Microbial RNA was extracted using a modified version of the HP Total

RNA Kit (Omega Biotek, Norcross, GA). Detailed protocols for DNA and RNA extractions can

be found here: dx.doi.org/10.17504/protocols.io.jz5cp86,

https://dx.doi.org/10.17504/protocols.io.jz6cp9e. Sequencing libraries were prepared using the

NEBNext Ultra Directional RNA Library Prep Kit for Illumina (NEB) following product

guidelines with an RNA fragmentation time of 10 min. The prepared library was submitted to the

Georgia Genomics and Bioinformatics Core facility for normalization, pooling, and sequencing

(Illumina HiSeq; Illumina, Inc., San Diego, CA).

Sheared DNA samples were end-repaired, dA-tailed, and adaptor ligated as instructed

with the NEBNext Ultra II DNA Library Prep Kit for Illumina (New England BioLabs Inc).

Adaptor-ligated fragments were size-selected with a magnetic bead protocol (Glenn et al., 2019)

derived from (Rohland and Reich, 2012). Size-selected and adapter-ligated DNA was enriched

via PCR using the NEBNext Ultra II DNA Library Prep Kit for Illumina with the NEBNext

Multiplex Oligos for Illumina as dual index primers (New England BioLabs Inc.). A final

metagenome pool was sent for sequencing at Novogene Corporation Inc. (Sacramento, CA)

following their specifications.

Upon receiving sequences, FastQC and MultiQC were used to check for low-quality

regions. Adapter sequences were trimmed using BBDuk from BBTools. Reads were then filtered

613 to remove rRNA sequences using SortMeRNA (96). Sequences that aligned to the *Periplaneta*

614 *americana* (97), *Blattabacterium* genome or plasmid (98) were removed using BBSplit. Paired-

615 end reads were merged using BBMerge and reads that did not successfully merge were rescued

616 by pseudo-merging using a custom laboratory script. For this, the forward read was joined with

617 the reverse complement of the reverse read using a string 10 Ns as an internal spacer to denote a

618 gap of unknown length.

619 *Construction of reference genome clusters for key taxonomic clusters*

620  An obstacle to RNA-seq analysis of gut microbiota is the relative lack of available

621 reference genomes. To surmount this problem, we utilized a pangenome mapping approach

622 based loosely on that employed in (45). This approach takes the available reference genomes

623 associated with taxonomic clusters of interest to construct a genome cluster for analysis of

624 group-level transcriptional activity.

625  We identified 18 genome clusters for analysis of the metatranscriptome data (see Table

626 S1 for a full list of reference genomes used for constructing each cluster). Genome sequences for

627 inclusion in each genome cluster were identified using a variety of metrics, including taxonomic

628 annotation, phylogenetic information from 16S rRNA gene and genome trees, abundance in

629 metagenome and metatranscriptome based on RefSeq homology searches as described below,

630 and co-correlation analyses.

631  Pangenomes of orthologous gene clusters for each taxon were built with Anvi'o (99). We

632 used *anvi-pan-genome* with DIAMOND (100) to calculate genome similarity within the taxon

633 and bin individual genes into gene clusters using the MCL algorithm (101, 102). We used *anvi-*

634 *pan-genome* with DIAMOND (100) to calculate genome similarity within the taxon and bin

635 individual genes into gene clusters using the MCL algorithm (103, 104). Functional annotations

for gene clusters were found using DIAMOND (100) to match functions and pathways in KEGG

(105), COG (103, 104, 106), and CAZy (107, 108). Additional annotations from a DIAMOND

search of each taxon's reference genomes to the Transporter Classification Database (TCDB)

were added to gene clusters (109).

Metatranscriptome and metagenome reads were translated and mapped to protein

sequences in RefSeq Prokaryotic Genomes (accessed 11/2021) (110) and to our dataset of 96

single-cell genomes from cockroach gut microbes (39, 111) using DIAMOND (100). The top-

scoring hit (based on bit score) and all equal hits were kept. Custom scripts were used to map

reads to genome clusters and build a per-sample gene cluster hit count table.

Analysis of genome cluster abundance and transcriptional activity

To calculate genome cluster abundance, the number of metatranscriptome and

metagenome read hits to each taxon was divided by the number of reads in that sample with hits

to the combined RefSeq and SAG reference database.

Weighted Bray-Curtis dissimilarities were calculated in R on whole-community

taxonomic tables using *vegdist()* from the vegan package (112). Differences in dissimilarity

values between diets were tested for significance in R with the Kruskal-Wallace method

followed by Dunn's test with Bonferroni correction (*dunnTest()* from FSA package and

kruskal.test() from stats package, respectively). Beta-dispersion in dissimilarity values were

tested using the betadisper() function from the vegan package (112). Permutational multivariate

analysis of variance (PERMANOVA) was carried out on dissimilarities using *adonis2()* from the

vegan R package with host diet as the factor of variation.

For all genome clusters, DESeq2 (43, 113) was used for pairwise comparisons between

diet treatments to find significantly differentially expressed gene clusters between diets. Relative

transcriptional abundances of gene clusters found to be significantly differentially expressed

between any pairwise diet comparison were visualized with heatmaps to determine their diet

specificity.

Acknowledgements:

This work was supported by the National Institute of General Medical Sciences of the National

Institutes of Health (NIH) under award number R35GM133789.

Data Availability:

Data associated with this study are available from NCBI SRA under BioProject accession

number PRJNA994902. Scripts and workflow for the metatranscriptomic analyses can be found

here: https://github.com/adepoy1/cockroach_metatranscriptome_manuscript.

References

1. Wong JMW, de Souza R, Kendall CWC, Emam A, Jenkins DJA. 2006. Colonic Health: Fermentation and Short Chain Fatty Acids. Journal of Clinical Gastroenterology 40.

2. Evrensel A, Ceylan ME. 2015. The Gut-Brain Axis: The Missing Link in Depression. Clin Psychopharmacol & Neurosci 13:239-244.

3. Kang D-W, Park JG, Ilhan ZE, Wallstrom G, Labaer J, Adams JB, Krajmalnik-Brown R. 2013. Reduced Incidence of *Prevotella* and Other Fermenters in Intestinal Microflora of Autistic Children. PLoS ONE 8:e68322.

4. Le Roy T, Llopis M, Lepage P, Bruneau A, Rabot S, Bevilacqua C, Martin P, Philippe C, Walker F, Bado A, Perlemuter G, Cassard-Doulcier AM, Gerard P. 2013. Intestinal microbiota determines development of non-alcoholic fatty liver disease in mice. 62:1787-1794.

5. Ferrer M, Ruiz A, Lanza F, Haange S-B, Oberbach A, Till H, Bargiela R, Campoy C, Segura MT, Richter M, Von Bergen M, Seifert J, Suarez A. 2013. Microbiota from the distal guts of lean and obese adolescents exhibit partial functional redundancy besides clear differences in community structure. Environ Microbiol 15:211-226.

6. Ley RE, Turnbaugh PJ, Klein S, Gordon JI. 2006. Human gut microbes associated with obesity. Nature 444:1022-1023.

7. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. 2006. An obesity-associated gut microbiome with increased capacity for energy harvest. Nature 444:1027-1031.

8. Kurilshikov A, Medina-Gomez C, Bacigalupe R, Radjabzadeh D, Wang J, Demirkan A, Le Roy CI, Raygoza Garay JA, Finnicum CT, Liu X, Zhernakova DV, Bonder MJ, Hansen TH, Frost F, Rühlemann MC, Turpin W, Moon J-Y, Kim H-N, Lüll K, Barkan E, Shah SA, Fornage M, Szopinska-Tokov J, Wallen ZD, Borisevich D, Agreus L, Andreasson A, Bang C, Bedrani L, Bell JT, Bisgaard H, Boehnke M, Boomsma DI, Burk RD, Claringbould A, Croitoru K, Davies GE, Van Duijn CM, Duijts L, Falony G, Fu J, Van Der Graaf A, Hansen T, Homuth G, Hughes DA, Ijzerman RG, Jackson MA, Jaddoe VWV, Joossens M, Jørgensen T, et al. 2021. Large-scale association analyses identify host factors influencing human gut microbiome composition. Nature Genetics 53:156-165.

9. Boccuto L, Tack J, Ianiro G, Abenavoli L, Scarpellini E. 2023. Human Genes Involved in the Interaction between Host and Gut Microbiome: Regulation and Pathogenic Mechanisms. Genes 14:857.

10. Sutherland DB, Suzuki K, Fagarasan S. 2016. Fostering of advanced mutualism with gut microbiota by Immunoglobulin A. Immunological Reviews 270:20-31.

11. Peterson DA, McNulty NP, Guruge JL, Gordon JI. 2007. IgA Response to Symbiotic Bacteria as a Mediator of Gut Homeostasis. Cell Host & Microbe 2:328-339.

12. Meyer-Hoffert U, Hornef MW, Henriques-Normark B, Axelsson LG, Midtvedt T, Pütsep K, Andersson M. 2008. Secreted enteric antimicrobial activity localises to the mucus surface layer. Gut 57:764.

13. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin AS, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ. 2014. Diet rapidly and reproducibly alters the human gut microbiome. Nature 505:559-563.

715   14.   Cotillard A, Kennedy SP, Kong LC, Prifti E, Pons N, Le Chatelier E, Almeida M,
716   Quinquis B, Levenez F, Galleron N, Gougis S, Rizkalla S, Batto J-M, Renault P, Doré J,
717   Zucker J-D, Clément K, Ehrlich SD, Blottière H, Leclerc M, Juste C, De Wouters T,
718   Lepage P, Fouqueray C, Basdevant A, Henegar C, Godard C, Fondacci M, Rohia A,
719   Hajduch F, Weissenbach J, Pelletier E, Le Paslier D, Gauchi J-P, Gibrat J-F, Loux V,
720   Carré W, Maguin E, Van De Guchte M, Jamet A, Boumezbeur F, Layec S. 2013. Dietary
721   intervention impact on gut microbial gene richness. Nature 500:585-588.
722   15.   Walker AW, Ince J, Duncan SH, Webster LM, Holtrop G, Ze X, Brown D, Stares MD,
723   Scott P, Bergerat A, Louis P, McIntosh F, Johnstone AM, Lobley GE, Parkhill J, Flint
724   HJ. 2011. Dominant and diet-responsive groups of bacteria within the human colonic
725   microbiota. The ISME Journal 5:220-230.
726   16.   Drasar BS, Crowther JS, Goddard P, Hawksworth G, Hill MJ, Peach S, Williams REO,
727   Renwich A. 1973. The relation between diet and the gut microflora in man. Proceedings
728   of the Nutrition Society 32:49-52.
729   17.   Fava F, Gitau R, Griffin BA, Gibson GR, Tuohy KM, Lovegrove JA. 2013. The type and
730   quantity of dietary fat and carbohydrate alter faecal microbiome and short-chain fatty
731   acid excretion in a metabolic syndrome 'at-risk' population. International Journal of
732   Obesity 37:216-223.
733   18.   Statovci D, Aguilera M, MacSharry J, Melgar S. 2017. The Impact of Western Diet and
734   Nutrients on the Microbiota and Immune Response at Mucosal Interfaces. Frontiers in
735   Immunology 8.
736   19.   Hildebrandt MA, Hoffmann C, Sherrill–Mix SA, Keilbaugh SA, Hamady M, Chen YY,
737   Knight R, Ahima RS, Bushman F, Wu GD. 2009. High-Fat Diet Determines the
738   Composition of the Murine Gut Microbiome Independently of Obesity. Gastroenterology
739   137:1716-1724.e2.
740   20.   Turnbaugh PJ, Bäckhed F, Fulton L, Gordon JI. 2008. Diet-Induced Obesity Is Linked to
741   Marked but Reversible Alterations in the Mouse Distal Gut Microbiome. Cell Host &
742   Microbe 3:213-223.
743   21.   Turnbaugh PJ, Ridaura VK, Faith JJ, Rey FE, Knight R, Gordon JI. 2009. The Effect of
744   Diet on the Human Gut Microbiome: A Metagenomic Analysis in Humanized
745   Gnotobiotic Mice. Science Translational Medicine 1:6ra14-6ra14.
746   22.   De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S,
747   Pieraccini G, Lionetti P. 2010. Impact of diet in shaping gut microbiota revealed by a
748   comparative study in children from Europe and rural Africa. Proc Natl Acad Sci U S A
749   107:14691-6.
750   23.   Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL,
751   Clemente JC, Knight R, Heath AC, Leibel RL, Rosenbaum M, Gordon JI. 2013. The
752   long-term stability of the human gut microbiota. Science 341:1237439.
753   24.   Bortolin RC, Vargas AR, Gasparotto J, Chaves PR, Schnorr CE, Martinello KB, Silveira
754   AK, Rabelo TK, Gelain DP, Moreira JCF. 2018. A new animal diet based on human
755   Western diet is a robust diet-induced obesity model: comparison to high-fat and cafeteria
756   diets in term of metabolic and gut microbiota disruption. International Journal of Obesity
757   42:525-534.
758   25.   Panasevich MR, Meers GM, Linden MA, Booth FW, II JWP, Fritsche KL, Wankhade
759   UD, Chintapalli SV, Shankar K, Ibdah JA, Rector RS. 2018. High-fat, high-fructose,
760   high-cholesterol feeding causes severe NASH and cecal microbiota dysbiosis in juvenile

Ossabaw swine. American Journal of Physiology-Endocrinology and Metabolism
314:E78-E92.

26. Amato KR, Yeoman CJ, Cerda G, A. Schmitt C, Cramer JD, Miller MEB, Gomez A, R.
Turner T, Wilson BA, Stumpf RM, Nelson KE, White BA, Knight R, Leigh SR. 2015.
Variable responses of human and non-human primate gut microbiomes to a Western diet.
Microbiome 3.

27. Chen L, Wang D, Garmaeva S, Kurilshikov A, Vich Vila A, Gacesa R, Sinha T, Segal E,
Weersma RK, Wijmenga C, Zhernakova A, Fu J. 2021. The long-term genetic stability
and individual specificity of the human gut microbiome. Cell 184:2302-2315.e12.

28. Mehta RS, Abu-Ali GS, Drew DA, Lloyd-Price J, Subramanian A, Lochhead P, Joshi
AD, Ivey KL, Khalili H, Brown GT, Dulong C, Song M, Nguyen LH, Mallick H, Rimm
EB, Izard J, Huttenhower C, Chan AT. 2018. Stability of the human faecal microbiome in
a cohort of adult men. Nature Microbiology 3:347-355.

29. Schloissnig S, Arumugam M, Sunagawa S, Mitreva M, Tap J, Zhu A, Waller A, Mende
DR, Kultima JR, Martin J, Kota K, Sunyaev SR, Weinstock GM, Bork P. 2013. Genomic
variation landscape of the human gut microbiome. Nature 493:45-50.

30. Garud NR, Good BH, Hallatschek O, Pollard KS. 2019. Evolutionary dynamics of
bacteria in the gut microbiome within and across hosts. PLOS Biology 17:e3000102.

31. Wu GD, Chen J, Hoffmann C, Bittinger K, Chen Y, Keilbaugh SA, Bewtra M, Knights
D, Walters WA, Knight R, Sinha R, Gilroy E, Gupta K, Baldassano R, Nessel L, Li H,
Bushman FD, Lewis JD. 2011. Linking Long-Term Dietary Patterns with Gut Microbial
Enterotypes. Science 334:105-108.

32. Leeming ER, Johnson AJ, Spector TD, Le Roy CI. 2019. Effect of Diet on the Gut
Microbiota: Rethinking Intervention Duration. Nutrients 11.

33. Fragiadakis GK, Wastyk HC, Robinson JL, Sonnenburg ED, Sonnenburg JL, Gardner
CD. 2020. Long-term dietary intervention reveals resilience of the gut microbiota despite
changes in diet and weight. Am J Clin Nutr 111:1127-1136.

34. Willis HJ, Slavin JL. 2020. The Influence of Diet Interventions Using Whole, Plant Food
on the Gut Microbiome: A Narrative Review. J Acad Nutr Diet 120:608-623.

35. Tinker KA, Ottesen EA. 2016. The Core Gut Microbiome of the American Cockroach,
Periplaneta americana, Is Stable and Resilient to Dietary Shifts. Appl Environ Microbiol
82:6603-6610.

36. Mikaelyan A, Thompson CL, Hofer MJ, Brune A. 2016. Deterministic Assembly of
Complex Bacterial Communities in Guts of Germ-Free Cockroaches. Appl Environ
Microbiol 82:1256-63.

37. Schauer C, Thompson C, Brune A. 2014. Pyrotag sequencing of the gut microbiota of the
cockroach *Shelfordella lateralis* reveals a highly dynamic core but only limited effects of
diet on community structure. PLoS One 9:e85861.

38. Schauer C, Thompson CL, Brune A. 2012. The Bacterial Community in the Gut of the
Cockroach *Shelfordella lateralis* Reflects the Close Evolutionary Relatedness of
Cockroaches and Termites. 78:2758-2767.

39. Dukes HE, Tinker KA, Ottesen EA. 2023. Disentangling hindgut metabolism in the
American cockroach through single-cell genomics and metatranscriptomics. Frontiers in
Microbiology 14.

40. Lampert N, Mikaelyan A, Brune A. 2019. Diet is not the primary driver of bacterial
community structure in the gut of litter-feeding cockroaches. BMC Microbiology 19.

807 41. Pérez-Cobas AE, Maiques E, Angelova A, Carrasco P, Moya A, Latorre A. 2015. Diet
808   shapes the gut microbiota of the omnivorous cockroach *Blattella germanica*. FEMS
809   Microbiol Ecol 91.
810 42. Yutin N, Galperin MY. 2013. A genomic update on clostridial phylogeny: Gram-negative
811   spore formers and other misplaced clostridia. Environmental Microbiology
812   doi:10.1111/1462-2920.12173:n/a-n/a.
813 43. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
814   for RNA-seq data with DESeq2. Genome Biology 15.
815 44. Liu D, De Schutter K, Smargiasso N, De Pauw E, Van Damme EJM, Smagghe G. 2019.
816   The N-glycan profile of the peritrophic membrane in the Colorado potato beetle larva
817   (*Leptinotarsa decemlineata*). Journal of Insect Physiology 115:27-32.
818 45. Ottesen EA, Young CR, Eppley JM, Ryan JP, Chavez FP, Scholin CA, DeLong EF.
819   2013. Pattern and synchrony of gene expression among sympatric marine microbial
820   populations. Proceedings of the National Academy of Sciences 110:E488-E497.
821 46. Ottesen EA, Young CR, Gifford SM, Eppley JM, Marin R, Schuster SC, Scholin CA,
822   DeLong EF. 2014. Multispecies diel transcriptional oscillations in open ocean
823   heterotrophic bacterial assemblages. Science 345:207-212.
824 47. Sharma AK, Becker JW, Ottesen EA, Bryant JA, Duhamel S, Karl DM, Cordero OX,
825   Repeta DJ, Delong EF. 2014. Distinct dissolved organic matter sources induce rapid
826   transcriptional responses in coexisting populations of *Prochlorococcus*, *Pelagibacter* and
827   the OM60 clade. Environmental Microbiology 16:2815-2830.
828 48. Wang T, Goyal A, Dubinkina V, Maslov S. 2019. Evidence for a multi-level trophic
829   organization of the human gut microbiome. PLOS Computational Biology 15:e1007524.
830 49. Vera-Ponce De León A, Jahnes BC, Duan J, Camuy-Vélez LA, Sabree ZL. 2020.
831   Cultivable, Host-Specific Bacteroidetes Symbionts Exhibit Diverse Polysaccharolytic
832   Strategies. Applied and Environmental Microbiology 86.
833 50. Kmezik C, Mazurkewich S, Meents T, McKee LS, Idström A, Armeni M, Savolainen O,
834   Brändén G, Larsbrink J. 2021. A polysaccharide utilization locus from the gut bacterium
835   *Dysgonomonas mossii* encodes functionally distinct carbohydrate esterases. Journal of
836   Biological Chemistry 296:100500.
837 51. Ravcheev DA, Godzik A, Osterman AL, Rodionov DA. 2013. Polysaccharides utilization
838   in human gut bacterium *Bacteroides thetaiotaomicron*: comparative genomics
839   reconstruction of metabolic and regulatory networks. BMC Genomics 14:873.
840 52. Sonnenburg ED, Zheng H, Joglekar P, Higginbottom SK, Firbank SJ, Bolam DN,
841   Sonnenburg JL. 2010. Specificity of Polysaccharide Use in Intestinal *Bacteroides* Species
842   Determines Diet-Induced Microbiota Alterations. Cell 141:1241-1252.
843 53. Rakoff-Nahoum S, Foster KR, Comstock LE. 2016. The evolution of cooperation within
844   the gut microbiota. Nature 533:255-259.
845 54. Rakoff-Nahoum S, Coyne J, Michael, Comstock E, Laurie. 2014. An Ecological Network
846   of Polysaccharide Utilization among Human Intestinal Symbionts. Current Biology
847   24:40-49.
848 55. Catlett JL, Carr S, Cashman M, Smith MD, Walter M, Sakkaff Z, Kelley C, Pierobon M,
849   Cohen MB, Buan NR. 2022. Metabolic Synergy between Human Symbionts *Bacteroides*
850   and *Methanobrevibacter*. Microbiology Spectrum 10:e01067-22.

56. Shetty SA, Kuipers B, Atashgahi S, Aalvink S, Smidt H, De Vos WM. 2022. Inter-
species Metabolic Interactions in an In-vitro Minimal Human Gut Microbiome of Core
Bacteria. npj Biofilms and Microbiomes 8.
57. Fernandez-Julia P, Commane DM, Van Sinderen D, Munoz-Munoz J. 2022. Cross-
feeding interactions between human gut commensals belonging to the Bacteroides and
Bifidobacterium genera when grown on dietary glycans. Microbiome Research Reports
doi:10.20517/mrr.2021.05.
58. Rey FE, Gonzalez MD, Cheng J, Wu M, Ahern PP, Gordon JI. 2013. Metabolic niche of
a prominent sulfate-reducing human gut bacterium. Proceedings of the National
Academy of Sciences 110:13582-13587.
59. Belenguer A, Duncan SH, Calder AG, Holtrop G, Louis P, Lobley GE, Flint HJ. 2006.
Two Routes of Metabolic Cross-Feeding between *Bifidobacterium adolescentis* and
Butyrate-Producing Anaerobes from the Human Gut. Applied and Environmental
Microbiology 72:3593-3599.
60. Chow V, Nong G, Preston JF. 2007. Structure, function, and regulation of the
aldouronate utilization gene cluster from *Paenibacillus sp.* strain JDR-2. Journal of
Bacteriology 189:8863-8870.
61. Hoogerheide JC, Kocholaty W. 1938. Metabolism of the strict anaerobes (genus:
Clostridium): Reduction of amino-acids with gaseous hydrogen by suspensions of *Cl.
sporogenes*. Biochemical Journal 32:949-957.
62. Doelle HW. 1969. 10 - NITROGEN METABOLISM AS AN ENERGY SOURCE FOR
ANAEROBIC MICROORGANISMS (CLOSTRIDIUM), p 402-422. *In* Doelle HW (ed),
Bacterial Metabolism doi:https://doi.org/10.1016/B978-1-4832-3135-8.50013-8.
Academic Press.
63. Kurth JM, Op den Camp HJM, Welte CU. 2020. Several ways one goal—
methanogenesis from unconventional substrates. Applied Microbiology and
Biotechnology 104:6839-6854.
64. Thomas CM, Taib N, Gribaldo S, Borrel G. 2021. Comparative genomic analysis of
Methanimicrococcus blatticola provides insights into host adaptation in archaea and the
evolution of methanogenesis. ISME Communications 1.
65. Ametaj BN, Zebeli Q, Saleem F, Psychogios N, Lewis MJ, Dunn SM, Xia J, Wishart DS.
2010. Metabolomics reveals unhealthy alterations in rumen metabolism with increased
proportion of cereal grain in the diet of dairy cows. Metabolomics 6:583-594.
66. Kelly WJ, Leahy SC, Kamke J, Soni P, Koike S, Mackie R, Seshadri R, Cook GM,
Morales SE, Greening C, Attwood GT. 2019. Occurrence and expression of genes
encoding methyl-compound production in rumen bacteria. Animal Microbiome 1.
67. McAllister TA, Okine EK, Mathison GW, Cheng KJ. 1996. Dietary, environmental and
microbiological aspects of methane production in ruminants. Canadian Journal of Animal
Science:231-243.
68. Wolin MJ, Miller TL, Stewart CS. 1997. Microbe-microbe interactions. *In* Hobson PN,
Stewart CS (ed), The Rumen Microbial Ecosystem. Blackie Academic & Professional,
New York.
69. Thomas CM, Taib N, Gribaldo S, Borrel G. 2021. Comparative genomic analysis of
Methanimicrococcus blatticola provides insights into host adaptation in archaea and the
evolution of methanogenesis. ISME Communications 1:1-11.

70. Macfarlane GT, Allison C, Gibson SAW, Cummings JH. 1988. Contribution of the
microflora to proteolysis in the human large intestine. Journal of Applied Bacteriology
64:37-46.
71. Barker HA. 1981. Amino Acid Degradation by Anaerobic Bacteria. Annual Review of
Biochemistry 50:23-40.
72. Sugiyama Y, Mori Y, Nara M, Kotani Y, Nagai E, Kawada H, Kitamura M, Hirano R,
Shimokawa H, Nakagawa A, Minami H, Gotoh A, Sakanaka M, Iida N, Koyanagi T,
Katayama T, Okamoto S, Kurihara S. 2022. Gut bacterial aromatic amine production:
aromatic amino acid decarboxylase and its effects on peripheral serotonin production.
Gut Microbes 14.
73. Naghdi S, Rezaei M, Tabarsa M, Abdollahi M. 2023. Ultrasonic-assisted enzymatic
extraction of sulfated polysaccharide from Skipjack tuna by-products. Ultrasonics
Sonochemistry 95:106385.
74. Hennequin C, Porcheray F, Waligora-Dupriet A, Collignon A, Barc M, Bourlioux P,
Karjalainen T. 2001. GroEL (Hsp60) of *Clostridium difficile* is involved in cell
adherance. Microbiology 147:87-96.
75. Gomes SL, Simão RCG. 2014. Stress Responses: Heat, Reference Module in Biomedical
Sciences doi:10.1016/b978-0-12-801238-3.02459-4.
76. Calloni G, Chen T, Schermann SM, Chang HC, Genevaux P, Agostini F, Tartaglia GG,
Hayer-Hartl M, Hartl FU. 2012. DnaK functions as a central hub in the *E. coli* chaperone
network. Cell Rep 1:251-64.
77. Wypych A, Ożgo M, Bernaciak M, Herosimczyk A, Barszcz M, Gawin K, Ciechanowicz
AK, Kucia M, Pierzchała M, Poławska E, Lepczyński A. 2024. Effect of feeding high fat
diets differing in fatty acid composition on oxidative stress markers and protein
expression in mouse kidney. Journal of Animal and Feed Sciences 33:170-184.
78. Enriquez JR, McCauley HA, Zhang KX, Sanchez JG, Kalin GT, Lang RA, Wells JM.
2022. A dietary change to a high-fat diet initiates a rapid adaptation of the intestine. Cell
Reports 41:111641.
79. Qiao Y, Sun J, Ding Y, Le G, Shi Y. 2013. Alterations of the gut microbiota in high-fat
diet mice is strongly linked to oxidative stress. Applied Microbiology and Biotechnology
97:1689-1697.
80. Devkota S, Chang EB. 2015. Interactions between Diet, Bile Acid Metabolism, Gut
Microbiota, and Inflammatory Bowel Diseases. Digestive Diseases 33:351-356.
81. Wang B, Kong Q, Li X, Zhao J, Zhang H, Chen W, Wang G. 2020. A High-Fat Diet
Increases Gut Microbiota Biodiversity and Energy Expenditure Due to Nutrient
Difference. Nutrients 12:3197.
82. Metzl-Raz E, Kafri M, Yaakov G, Soifer I, Gurvich Y, Barkai N. 2017. Principles of
cellular resource allocation revealed by condition-dependent proteome profiling. eLife
6:e28034.
83. Wang Q, Lin J. 2022. Environment-specificity and universality of the microbial growth
law. Communications Biology 5:891.
84. Heisel T, Montassier E, Johnson A, Al-Ghalith G, Lin Y-W, Wei L-N, Knights D, Gale
CA. 2017. High-Fat Diet Changes Fungal Microbiomes and Interkingdom Relationships
in the Murine Gut. mSphere 2:10.1128/msphere.00351-17.
85. Xiao L, Sonne SB, Feng Q, Chen N, Xia Z, Li X, Fang Z, Zhang D, Fjære E, Midtbø LK,
Derrien M, Hugenholtz F, Tang L, Li J, Zhang J, Liu C, Hao Q, Vogel UB, Mortensen A,

Kleerebezem M, Licht TR, Yang H, Wang J, Li Y, Arumugam M, Wang J, Madsen L,
Kristiansen K. 2017. High-fat feeding rather than obesity drives taxonomical and
functional changes in the gut microbiota in mice. Microbiome 5:43.
86. Lin H, An Y, Hao F, Wang Y, Tang H. 2016. Correlations of Fecal Metabonomic and
Microbiomic Changes Induced by High-fat Diet in the Pre-Obesity State. Scientific
Reports 6:21618.
87. Bizzini A, Zhao C, Budin-Verneuil A, Sauvageot N, Giard JC, Auffray Y, Hartke A.
2010. Glycerol is metabolized in a complex and strain-dependent manner in
Enterococcus faecalis. J Bacteriol 192:779-85.
88. Doi Y. 2018. Lactic acid fermentation is the main aerobic metabolic pathway in
Enterococcus faecalis metabolizing a high concentration of glycerol. Appl Microbiol
Biotechnol 102:10183-10192.
89. Zeng N, Wu F, Lu J, Li X, Lin S, Zhou L, Wang Z, Wu G, Huang Q, Zheng D, Gao J,
Wu S, Chen X, Chen M, Meng F, Shang H, He Y, Chen P, Wei H, Li Z, Zhou H. 2023.
High-fat diet impairs gut barrier through intestinal microbiota-derived reactive oxygen
species. Science China Life Sciences doi:10.1007/s11427-022-2283-4.
90. Strickertsson JAB, Desler C, Martin-Bertelsen T, Machado AMD, Wadstrøm T, Winther
O, Rasmussen LJ, Friis-Hansen L. 2013. *Enterococcus faecalis* Infection Causes
Inflammation, Intracellular Oxphos-Independent ROS Production, and DNA Damage in
Human Gastric Cancer Cells. PLoS ONE 8:e63147.
91. Lee J-Y, Bae E, Kim HY, Lee K-M, Yoon SS, Lee D-C. 2021. High-Fat-Diet–Induced
Oxidative Stress Linked to the Increased Colonization of Lactobacillus sakei in an Obese
Population. Microbiology Spectrum 9.
92. Wolf G, Strahl A, Meisel J, Hammes WP. 1991. Heme-dependent catalase activity of
lactobacilli. International Journal of Food Microbiology 12:133-140.
93. Liu QP, Sulzenbacher G, Yuan H, Bennett EP, Pietz G, Saunders K, Spence J, Nudelman
E, Levery SB, White T, Neveu JM, Lane WS, Bourne Y, Olsson ML, Henrissat B,
Clausen H. 2007. Bacterial glycosidases for the production of universal red blood cells.
Nature Biotechnology 25:454-464.
94. Bell WJ, Adiyodi KG (ed). 1982. The American Cockroach. Chapman and Hall, London.
95. Desai MS, Seekatz AM, Koropatkin NM, Kamada N, Hickey CA, Wolter M, Pudlo NA,
Kitamoto S, Terrapon N, Muller A, Young VB, Henrissat B, Wilmes P, Stappenbeck TS,
Nunez G, Martens EC. 2016. A Dietary Fiber-Deprived Gut Microbiota Degrades the
Colonic Mucus Barrier and Enhances Pathogen Susceptibility. Cell 167:1339-1353 e21.
96. Kopylova E, Noé L, Touzet H. 2012. SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics 28:3211-3217.
97. Li S, Zhu S, Jia Q, Yuan D, Ren C, Li K, Liu S, Cui Y, Zhao H, Cao Y, Fang G, Li D,
Zhao X, Zhang J, Yue Q, Fan Y, Yu X, Feng Q, Zhan S. 2018. The genomic and
functional landscapes of developmental plasticity in the American cockroach. Nature
Communications 9.
98. Sabree ZL, Kambhampati S, Moran NA. 2009. Nitrogen recycling and nutritional
provisioning by Blattabacterium, the cockroach endosymbiont. Proceedings of the
National Academy of Sciences 106:19521-19526.
99. Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef
M, Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD,
Kivenson V, Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin

A, Scott JJ, Vázquez-Campos X, Mckay LJ, Mcdaniel EA, Stevens SLR, Anderson RE,
Fuessel J, Fernandez-Guerra A, Maignien L, Delmont TO, Willis AD. 2021. Community-
led, integrated, reproducible multi-omics with anvi'o. Nature Microbiology 6:3-6.

100. Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using
DIAMOND. Nature Methods 12:59-60.

101. Van Dongen S. 2008. Graph Clustering Via a Discrete Uncoupling Process. SIAM
Journal on Matrix Analysis and Applications 30:121-141.

102. Enright AJ, Van Dongen S, Ouzounis CA. 2002. An efficient algorithm for large-scale
detection of protein families. Nuc Acids Res 30:1575-1584.

103. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for
genome-scale analysis of protein functions and evolution. Nucleic Acids Research 28:33-
36.

104. Galperin MY, Wolf YI, Makarova KS, Vera Alvarez R, Landsman D, Koonin EV. 2020.
COG database update: focus on microbial diversity, model organisms, and widespread
pathogens. Nucleic Acids Research 49:D274-D281.

105. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020.
KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score
threshold. Bioinformatics 36:2251-2252.

106. Agarwala R, Barrett T, Beck J, Benson DA, Bollin C, Bolton E, Bourexis D, Brister JR,
Bryant SH, Canese K, Cavanaugh M, Charowhas C, Clark K, Dondoshansky I, Feolo M,
Fitzpatrick L, Funk K, Geer LY, Gorelenkov V, Graeff A, Hlavina W, Holmes B,
Johnson M, Kattman B, Khotomlianski V, Kimchi A, Kimelman M, Kimura M, Kitts P,
Klimke W, Kotliarov A, Krasnov S, Kuznetsov A, Landrum MJ, Landsman D, Lathrop S,
Lee JM, Leubsdorf C, Lu Z, Madden TL, Marchler-Bauer A, Malheiro A, Meric P,
Karsch-Mizrachi I, Mnev A, Murphy T, Orris R, Ostell J, O'Sullivan C, Palanigobu V, et
al. 2018. Database resources of the National Center for Biotechnology Information.
Nucleic Acids Research 46:D8-D13.

107. Drula E, Garron M-L, Dogan S, Lombard V, Henrissat B, Terrapon N. 2021. The
carbohydrate-active enzyme database: functions and literature. Nucleic Acids Research
50:D571-D577.

108. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. 2014. The
carbohydrate-active enzymes database (CAZy) in 2013. Nuc Acids Res 42:D490-D495.

109. Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, Lam
K, Jing, Katie, Tian N, Russum S, Wang J, Medrano-Soto A. 2021. The Transporter
Classification Database (TCDB): 2021 update. Nucleic Acids Research 49:D461-D467.

110. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, Mcveigh R, Rajput B,
Robbertse B, Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y, Blinkova
O, Brover V, Chetvernin V, Choi J, Cox E, Ermolaeva O, Farrell CM, Goldfarb T, Gupta
T, Haft D, Hatcher E, Hlavina W, Joardar VS, Kodali VK, Li W, Maglott D, Masterson
P, Mcgarvey KM, Murphy MR, O'Neill K, Pujar S, Rangwala SH, Rausch D, Riddick
LD, Schoch C, Shkeda A, Storz SS, Sun H, Thibaud-Nissen F, Tolstoy I, Tully RE,
Vatsan AR, Wallin C, Webb D, Wu W, Landrum MJ, Kimchi A, et al. 2016. Reference
sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional
annotation. Nucleic Acids Research 44:D733-D745.

111. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L,
Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome
annotation pipeline. Nucleic Acids Research 44:6614-6624.
112. Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos
P, Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D,
Carvalho G, Chirico M, De Caceres M, Durand S, Evangelista H, FitzJohn R, Friendly
M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlinn D, Ouellette M, Ribeiro CE,
Smith T, Stier A, Ter Braak C, Weedon J. 2022. vegan: Community Ecology Package,
vR package version 2.6-4. https://CRAN.R-project.org/package=vegan.
113. Zhu A, Ibrahim JG, Love MI. 2018. Heavy-tailed prior distributions for sequence count
data: removing the noise and preserving large differences. Bioinformatics 35:2084-2092.

1043

1044

1045 Figure Legends and tables

1046 **Table 1.** Nutrient information for 100g serving of diet treatment. All cockroaches were raised on

1047 our laboratory control diet of dog food, composed of 21% protein, 8% fat, and 6% fiber.

1048 Carbohydrate contents of dog food are estimated based on levels of other macronutrients, ash,

1049 and moisture.

Diet Treatment	Calories	Protein (g)	Carbohydrate (g)	Fat (g)	Fiber (g)
Bran	375	17.5	67.5	5	17.5
Butter	714	0	0	79	0
Tuna	107	27	0	1	0
Starvation	N/A	N/A	N/A	N/A	N/A
Dog Food	350-399	21	45	8	6

1050

Figure 1. Taxonomy and beta diversity metrics for the metagenome and metatranscriptome

datasets (A)Within diet and all-diet comparison of weighted Bray-Curtis dissimilarities were

calculated using taxonomic community profiles from the metagenome and metatranscriptome

datasets. Tests of significance among the diets were done with a Kruskal test and Dunn's test.

Beta dispersion was tested using the betadisper() function in vegan. (B) Relative abundance of

families in metagenome samples. (C) Relative abundance of families in metatranscriptome

samples. Each column represents a hindgut sample from a single cockroach. Families with a

maximum relative abundance of at least 5% in either the metatranscriptome or metagenome

datasets are shown.

Figure 2. Average relative abundance of genome clusters in metatranscriptome and metagenome

datasets across diet treatments plotted using ggradar package for R.

Figure 3. Within-genome cluster transcriptional abundance of a manually curated list of KEGG

pathways (binned at third hierarchy), RND efflux pumps (gene clusters classified as belonging to

TCDB families 2.A.6 or 8.A.50), TonB-dependent transporters (gene clusters classified as

belonging to TCDB families 1.B.14 or 2.C.1), and total summed CAZymes. Relative abundance

was averaged within diet treatment.

Figure 4. Number of differentially expressed genes identified by DESeq2. Each number within

each square is the number of differentially expressed genes when those two diets were compared.

1069

A

legend:
- metagenome
- metatranscriptome

x-axis: all samples, bran, dog food, tuna, butter, starvation
y-axis: weighted Bray-Curtis distance

B

y-axis: Relative abundance of families in metagenomes
x-axis: bran, dog food, tuna, butter, starvation

C

y-axis: Relative abundance of families in metatranscriptomes
x-axis: bran, dog food, tuna, butter, starvation

Legend:
- others below 5%
- f_Holophaga group
- f_Eggerthellaceae
- f_Treponemataceae
- f_Synergistaceae
- f_Nitrobacteraceae
- f_Hafniaceae
- f_Caulobacteraceae
- f_Desulfobacteraceae
- f_Desulfovibrionaceae
- f_Erysipelotrichaceae
- f_Enterococcaceae
- f_Lactobacillaceae
- f_Clostridiaceae
- f_Christensenellaceae
- f_Lachnospiraceae
- f_Ruminococcaceae
- f_Odoribacteraceae
- f_Dysgonomonadaceae
- unclassified Bacteroidales
- f_Tannerellaceae
- f_Bacteroidaceae