Metabolic flexibility among fiber degraders supports gut microbiome stability across host diets

in the American cockroach

Amber N. DePoy[a], Helen E. Wall[a]*, Kara A. Tinker[a]*, Ian M. Black[b], Elizabeth A. Ottesen [a]#

[a] Department of Microbiology, University of Georgia, Athens, Georgia, USA

[b] Complex Carbohydrate Research Center, University of Georgia, Athens, Georgia, USA

Running title: Cockroach hindgut microbial response to host diet

Keywords: cockroach (*Periplaneta americana*), gut microbiome, metatranscriptome, diet

#Address correspondence to Elizabeth A. Ottesen, ottesen@uga.edu.

*Present address: Helen E. Wall, Center for Disease Control and Prevention, Atlanta, Georgia,

USA; Kara A. Tinker, National Energy Technology Laboratory (NETL), Pittsburgh, PA, USA

Amber N. DePoy and Helen E. Wall contributed equally to this work. Author order was

determined by alphabetical order.

Abstract

Scavenging omnivores face challenges in maintaining gut microbiome homeostasis due

to a highly variable diet. Work in mammalian models has highlighted the importance of fiber-

degrading taxa and their utilization of dietary polysaccharides and host glycans in structuring gut

microbiome composition and function. However, less is known about biological mechanisms

supporting gut microbiome homeostasis in insects. Here, we present a metatranscriptomic

analysis of hindgut microbial transcriptional responses to diverse host diets in the American

cockroach *(Periplaneta americana)*, a hindgut-fermenting and omnivorous insect. We analyzed

microbial community transcriptomes from hindguts of cockroaches fed high-fiber, high-protein,

high-fat, and starvation diets. Microbial taxa predicted to be primary fiber degraders increased

transcription of genes associated with utilization of dietary polysaccharides in insects fed a high-

fiber diet. In contrast, these taxa responded to starvation and low fiber diets with upregulation of

transcripts associated with degradation of host glycans. Analysis of N- and O-glycans in the

cockroach midgut confirm the presence of predicted substrates for these pathways. As a result,

we hypothesize that peritrophic matrix glycans serve as a critical carbohydrate subsidy

supporting insect gut microbial metabolism in the absence of dietary fiber. Observed stability in

gut microbiome composition and the transcriptional activity of many secondary fermenters and

hydrogenotrophic taxa in the gut suggests that this strategy can be highly successful. Together,

this work supports a novel physiological role for the peritrophic matrix of insects and other

invertebrates in supporting gut microbiome homeostasis across dietary shifts.

Introduction

The gut microbiome shares an intimate relationship with its host, impacting digestion, nutrition, immunity, and more [1-5]. The host can modify the gut microbiome through various pathways [6-10], but diet is considered one of the most direct drivers of gut microbiome composition [11-15]. Much of the work studying the effect of diet on the gut microbiome has been done in mammalian hosts, such as mice [16-18], humans [11,19-21], and other mammals [22-24]. These mammalian studies have shown that dietary fiber and plant polysaccharides play a particularly important role shaping gut microbiome composition [25-27]. Despite the importance of diet in shaping the gut microbiome, multiple long-term studies of individual humans have shown stability over time [20,28-30]. Short-term dietary interventions in humans often fail to elicit shifts in the gut microbiome composition more than pre-existing individual-to-individual differences [28,31,32]. These results emphasize the need for an understanding of microbe-microbe or host-microbe interactions that may assist in the maintenance of gut microbiome homeostasis.

The gut microbiome of many insects is less diverse and more variable than that of mammals [2], however there are numerous exceptions, including dung beetles, termites, ants, and cockroaches [33-38]. Insect models such as these provide an opportunity to explore diverse and evolutionarily ancient biological mechanisms supporting host-gut microbe symbioses. In this work, we explore gut microbiome homeostasis in response to host dietary shifts in the omnivorous American cockroach (*Periplaneta americana*). The American cockroach hindgut microbial community is highly diverse and includes many taxa from clades found in mammalian and human guts [33,39-41]. These taxa include those in the abundant Bacteroidota and Bacillota phyla that have similar taxonomic and functional profiles to their sister clades found in

mammalian guts [42]. Multiple studies have demonstrated that, like humans, the cockroach gut

microbiome exhibits stability across highly divergent diets [33,40,43,44]. This stability is likely

advantageous to their lifestyle as scavenging omnivores, but host and microbial activities

supporting this stability are poorly understood.

Our goal of this study was to explore the microbial activities underlying compositional

stability with host diet changes, with a particular focus on understanding metabolic shifts

supporting the ability of key gut taxa to persist in the absence of a dietary source with preferred

growth substrates. We examined hindgut microbial community activity using metagenomic and

metatranscriptomic approaches in cockroaches fed a high-fiber, high-protein, high-fat, and

starvation diet. Metagenomic analyses confirmed taxonomic stability, while metatranscriptomic

profiles showed plasticity in transcriptional activity among key taxa, particularly primary fiber

degraders. Analysis of carbohydrate active enzyme expression highlights a role for host glycans

in supporting fiber degraders in low fiber diets. Together, these results provide new insight into

microbial responses to dietary shifts that support gut microbiome homeostasis.

Materials and Methods

Diet manipulation and sample collection

Cockroaches (*Periplaneta americana*) are maintained in mixed age, mixed sex colonies

in aquarium tanks at room temperature on a diet of dog food *ad libitum*. Each tank is provided

with corn cob bedding, cardboard tubes for nesting, and a cellulose sponge saturated with water.

For experiments using controlled diets, each treatment group was housed in a single plastic tank

that contained pebbles for bedding, a large weigh boat for shelter, and food and water in weigh

boats. Food, water, and weigh boats were changed as needed, and any ootheca or deceased

cockroaches were removed daily. Treatments included bran (Bob's Red Mill Organic High Fiber

Oat Bran Hot Cereal), butter (Kroger Unsalted Butter Sticks), tuna (StarKist Selects Low

Sodium Chunk Light Tuna in Water), and starvation (Table 1).

After 14 days on the assigned diet, all cockroaches were sacrificed and their hindgut

microbiota preserved for DNA or RNA extraction. Individual cockroaches were removed from

tanks, weighed, and placed on ice in sterile culture plates. Once sufficiently torpid, cockroaches

were dissected, and the entire gut was removed. Any visible debris, including fat bodies or

exoskeleton, was removed with forceps. The hindgut was then separated from the rest of the gut

using a scalpel and submerged in either 100 μL of 1XTE buffer or RNALater (Ambion, Austin,

TX, USA). Hindgut samples placed in 1XTE buffer were immediately stored at -80°C. Hindgut

samples placed in RNALater were crushed with a sterile glass stirring rod. The suspended gut

lumen was then aspirated and stored at -80°C.

DNA and RNA extraction, sequencing library preparation

Microbial DNA and RNA were extracted from hindgut samples as outlined previously in

[42]. Briefly, DNA and RNA were extracted using a modified version of the EZNA Bacteria Kit

and HP Total RNA Kit (Omega Biotek, Norcross, GA), respectively. For metatranscriptome

sequencing, RNA libraries were prepared using the NEBNext Ultra Directional RNA Library

Prep Kit for Illumina (NEB) following product guidelines with an RNA fragmentation time of 10

min. For metagenome sequencing, DNA libraries were prepared using NEBNext Ultra II DNA

Library Prep Kit for Illumina (New England BioLabs Inc) and dual index primers from the

NEBNext Multiplex Oligos for Illumina (New England BioLabs Inc.). Size selection was done

using a protocol adapted from [45]. The prepared RNA library was sequenced (Illumina HiSeq)

by the Georgia Genomics and Bioinformatics Core (Athens, GA), while the metagenome pool

was sequenced (Illumina NovSeq) by Novogene Corporation Inc. (Sacramento, CA).

Raw sequences were checked for low-quality regions using FastQC and MultiQC.

Adapter sequences were trimmed using BBDuk from BBTools. Ribosomal RNA sequences were

removed using SortMeRNA [46]. *Periplaneta americana* [47] and *Blattabacterium* [48]

sequences were removed using BBSplit. Paired-end reads were merged using BBMerge and

reads that did not successfully merge were rescued by pseudo-merging the forward read with an

internal spacer of 10 Ns and the reverse complement of the reverse read. This approach allowed

the processing of pseudo-merged reads through downstream scripts in the same manner as

merged reads.

Construction of reference genome clusters for key taxonomic clusters

An obstacle to RNA-seq analysis of gut microbiota is the relative lack of available

reference genomes. To surmount this problem, we utilized a pangenome mapping approach

based loosely on that employed in [49]. This approach uses available reference genomes

associated with taxonomic clusters of interest to construct genome clusters for analysis of group-

level transcriptional activity. Reads were mapped to pangenomes constructed from clusters of

closely related genomes that show high levels of co-occurrence in the metagenome and

metatranscriptomes (e.g., Figures S1-S2). These analyses allowed us to compile genetic

information from multiple closely related reference genomes while minimizing ambiguity of

read mapping in conserved regions. Pangenomes were constructed for 18 genome clusters (see

Table S1 for a full list of reference genomes used for constructing each cluster). Genome

sequences for inclusion in each genome cluster were identified using a variety of metrics,

including taxonomic annotation, phylogenetic information from 16S rRNA gene and genome

trees, abundance in metagenome and metatranscriptome based on RefSeq homology searches as

described below, and co-correlation analyses.

Pangenomes of orthologous gene clusters for each taxon were built with Anvi'o [50]. We

used *anvi-pan-genome* with DIAMOND [51] to calculate genome similarity within the taxon and

bin individual genes into gene clusters using the MCL algorithm [52,53]. Functional annotations

for gene clusters were found using DIAMOND [51] to match functions and pathways in KEGG

[54], COG [55,56], and CAZy [57]. Additional annotations from a DIAMOND search of each

taxon's reference genomes to the Transporter Classification Database (TCDB) were added to

gene clusters [58].

Metatranscriptome and metagenome reads were translated and mapped to protein

sequences in RefSeq Prokaryotic Genomes (accessed 11/2021) [59] and to our dataset of 96

single-cell genomes from cockroach gut microbes [42,60] using DIAMOND [51]. The top-

scoring hit (based on bit score) and all equal hits were kept. Custom scripts were used to map

reads to genome clusters and build a per-sample gene cluster hit count table. See supplemental

Table S1-S3 for read tracking through the analysis steps.

Analysis of genome cluster abundance and transcriptional activity

To calculate genome cluster abundance, the number of metatranscriptome and

metagenome read hits to each taxon was divided by the number of reads in that sample with hits

to the combined RefSeq and SAG reference database. Weighted Bray-Curtis dissimilarities were

calculated in R on whole-community taxonomic tables using *vegdist()* from the vegan package

[61]. Differences in dissimilarity values between diets were tested for significance in R with the

Kruskal-Wallis method followed by Dunn's test with Bonferroni correction (*dunnTest()* from

FSA package and *kruskal.test()* from stats package, respectively). Permutational multivariate

analysis of variance (PERMANOVA) was carried out on dissimilarities using *adonis2()* from the

vegan R package with host diet as the factor of variation. Redundancy analyses (RDA) were

done with variance stabilized data normalized with DESeq2 [62,63]. For all taxa, DESeq2

[62,63] was used for pairwise comparisons between diet treatments to find significantly

differentially expressed gene clusters between diets.

Results

Taxonomic profiles and overall community composition

To address gut microbiome stability and the functions underlying that stability, we

sequenced metagenomes and metatranscriptomes from cockroaches fed four distinct dietary

conditions: bran (high-fiber), tuna (high-protein), butter (high-fat), and starvation. Metagenomic

reads were assigned to a diverse set of families, including Bacteroidaceae, Dysgonomonadaceae,

Tannerellaceae, Lachnospiraceae, and Desulfovibrionaceae (Figure 1). Relative abundances of

these families changed very little between diet treatments. Alpha diversity (Shannon index) did

not vary significantly across the diets ($p = 0.43$) (Figure S3). PERMANOVA analysis of beta

diversity using weighted Bray-Curtis dissimilarities based on taxonomic profiles confirmed that

diet was not significantly correlated with overall metagenome community composition ($r^2 =$

$0.18, p = 0.24$) (Figure 1), although we did observe significant enrichment/depletion of

individual taxa (Figure S4). Much of this taxa enrichment/depletion appears to be associated with

increased intra-treatment variability and associated 'blooms' of multiple taxa in individual

samples within the butter diet (Figure S1, Figure S5).

In contrast, diet was a significant driver of metatranscriptome composition ($r^2 = 0.30, p =$

0.01) and alpha diversity (Shannon index) ($p = 0.005$; bran vs. starve, $p = 0.02$) (Figure 1, Figure

S3). DESeq2 analysis of taxonomic composition showed significant enrichment of diverse

Bacteroidaceae in the bran diet compared to the other diets, as well as enrichment of

Lachnospiraceae, Leuconostocaceae, and Clostridiaceae (Figure S4). Few taxa were enriched in

the other diets (Figure S4).

Response of individual taxa to host diet treatment

Transcriptional profiles of each taxon were analyzed using redundancy analysis (RDA),

which demonstrated that diet was significantly associated with overall transcriptional activity in

most taxa, excluding Bacteroidales, Enterococcaceae, Lactobacillaceae, and *Desulfosarcina*

(Figure S6). When 'diet' was modeled as a factor, it explained 8.3% (Bacteroidales) to 21.9%

(Lactobacillaceae) of the variance (Figure S6). For many organisms, the presence/absence of

fiber alone (bran vs. all other diets) explained a comparable amount of transcriptional variance to

'diet' as a factor. This emphasizes the effect and importance of dietary fiber in shaping microbial

transcriptional profiles.

To characterize microbial responses and the extent of responses to host diet changes, we

used DESeq2 to identify genes that displayed significantly altered expression in pairwise

comparisons of the diet treatments. A combined total of 2,684 gene clusters were changed

significantly between host diets (Figure 2). Members of the Bacteroidota phylum altered

expression of the greatest number of genes, with *Bacteroides*, *Dysgonomonas,* and

Parabacteroides having far more than every other group (Figure 2, Figure S7, Table S4). Many

genes significantly changed expression between the high-fiber (bran) diet compared to each of

the other diets (Figure 2). Two Bacillota taxa, Oscillospiraceae_2 and Lactobacillaceae, had high

numbers of significantly responding genes (Figure S7, Table S5). However, other Bacillota taxa

had fewer differentially transcribed genes. Chemolithotrophic groups (sulfate reducers and

methanogens) had a range of significantly changing genes but showed less of a response with

gene expression changes than other taxa (Figure S7, Table S6).

Carbohydrate active enzymes (CAZyme) utilization between the host diets

Given the importance of fiber in shaping transcription, we analyzed genes active in the

breakdown of dietary polysaccharides (CAZymes). CAZyme gene expression (% of all mapped

transcripts) was highest in the bran diet (2.1% of all transcription in bran) and decreased with the

amount of fiber present in the diet (Figure 3B). Despite starvation having no dietary fiber input,

CAZyme gene expression remained high under starvation, still accounting for 0.99% of all

transcription (Figure 3B). *Bacteroides*, *Dysgonomonas*, and *Parabacteroides* produced the

highest number of CAZyme transcripts, emphasizing their role as primary degraders (Figure 3).

We used dbCAN-seq [64] to associate CAZyme family annotations with potential

substrates (Figure 3). These annotations can be somewhat ambiguous as fiber substrate

breakdown can overlap between CAZyme family classifications, but it remains a valuable way to

identify changes in putative substrate profiles across diets. CAZymes targeting three major

substrate types were expressed at high levels: xylan, pectin, and host glycans (Figure 3A).

Xylan-targeting CAZymes were expressed 2.4-fold more in the bran diet compared to the other

diets (Figure 3C). Pectin breakdown genes were expressed 2-fold more in tuna diet than butter

and starve but only slightly more than in bran (Figure 3D). Host glycan-targeting CAZymes were

among the most highly expressed across all conditions (Figure 3E). While all CAZyme groups

showed decreased expression in starvation (and butter), host glycan targeting CAZymes showed

smaller diet-dependent fluctuations than other CAZyme groups.

Patterns of CAZyme and carbohydrate utilization gene expression in Bacteroides

As one of the most active genome clusters in the metatranscriptome and the largest

source of CAZyme transcripts, we performed a more in-depth analysis of CAZyme expression

and carbohydrate utilization gene expression in *Bacteroides*. For these analyses, we expressed

transcription of each fiber substrate metabolism genes as a fraction of *Bacteroides* transcripts.

This standardization method allows us to examine changes in transcriptional activity within

Bacteroides (equivalent to single-organism RNA-seq) in isolation from changes in the relative

abundance of this organism within the metatranscriptome.

For *Bacteroides*, overall transcription of xylan targeting CAZymes was highest in the

bran diet (Figure 4A). Several CAZymes that target the xylan backbone (GH10 and GH3) were

significantly upregulated in bran (Figure 4A), as were CAZyme families targeting arabinose

(GH51) and glucuronate (GH115) side chains of xylan. Although not significant, CE1 and GH67

(required for removal of acetyl groups and glucuronic acid side chains) also had greater

transcription in the bran diet (Figure 4A). While xylan gene expression remains high in

starvation, much of this signal is associated with transcription of broad specificity CAZyme

families (i.e., GH2, GH3) that are associated with utilization of a wide range of substrates. We

also examined KEGG orthologs associated with processing of sugars released during xylan

breakdown, focusing on early processing steps prior to entry into shared metabolic pathways.

Bacteroides expressed two genes for xylose processing into xylulose-5P (for entry into the

pentose phosphate pathway) at high levels the bran diet (Figure 4D). *Bacteroides* also expressed

L-arabinose isomerase (K01804) at high levels in both bran and tuna diets (Figure 4G).

Arabinose can be released from either arabinoxylan or pectin processing.

Bacteroides responded to the tuna diet with increased expression of pectin targeting

CAZymes (Figure 4B). There were ten CAZyme families that were significantly upregulated in

tuna, six of which were specific for pectin (Figure 4B), including CE8 demethylases, GH105 and

GH106 for depolymerization of the pectin backbone, and polysaccharide lyases PL1, PL11, and

PL9 (Figure 4B). Although not specifically associated as a pectin CAZyme family, GH51 (L-

arabinfuranosidase) was also significantly upregulated in the tuna diet (Figure 4A). After initial

pectin breakdown, the main monosaccharides released for further metabolism include rhamnose,

galactose, galacturonic acid, fucose, and arabinose. *Bacteroides* expressed rhamnose and

galacturonic acid metabolism genes more in the tuna diet compared to the bran diet (8.6 and 4.3-

fold more, respectively) (Figure 4E, 4H). Galactose metabolism genes did not show the same

increase in the tuna diet but were highly expressed in most diets (Figure 4K).

Relative to total organismal transcriptional activity, *Bacteroides* showed 1.91-fold higher

expression of host glycan degradation CAZymes in starvation than in the bran diet (Figure 4C).

Eight CAZyme families associated with host glycan utilization were significantly upregulated in

starvation vs. bran. Most of these CAZymes were associated with N-glycan degradation (Figure

4C). *Bacteroides* increased expression of GH38 (α-mannosidases) 6.3-fold more in host

starvation compared to bran, suggesting target of mannoses (Figure 4C). In addition to GH38,

other CAZyme families with mannosidase activity (GH130, GH125, GH92) were significantly

upregulated in starvation compared to bran. *Bacteroides* also targeted fucose with two CAZymes,

one of which (GH29) was significantly upregulated in starvation. CAZyme families targeting O-

glycans were expressed to a lesser extent. Cleavage of terminal α-GalNAc residues from O-

glycans by GH109 was significantly upregulated in starvation. Monosaccharides from N-glycan

degradation include mannose, fucose, and GlcNAc. Genes associated with mannose breakdown

were transcribed under all conditions, although they were not significantly upregulated under

starvation (Figure 4F). Although not significant, fucose metabolism genes were transcribed more

in starvation and less in the butter diet (Figure 4L).

Taxa showing functional stability among the host diets

While primary degraders showed changes in CAZyme gene expression and downstream

utilization of the monosaccharides to each of the diets, these significant changes underscore the

extent to which other taxa were less affected by host diet changes. Secondary degraders,

including many organisms within the Bacillota, showed fewer significant differences in gene

expression across diets, and the presence/absence of dietary fiber explained a smaller fraction of

their overall transcriptional profile (Figure 2, Figure S6-S7). While the major fiber sources in

each diet require unique carbohydrate active enzymes to degrade, this degradation produces a

limited set of monosaccharides, which are ultimately metabolized via conserved core metabolic

pathways. Summed gene expression among core metabolic pathways, such as glycolysis, pentose

phosphate, and citrate pathways showed high and stable expression across the diets in most

organisms (Figure S8).

Other groups that are not directly involved in polysaccharide degradation, including

sulfate reducers (*Desulfosarcina*, *Frigididesulfovibrio*) and methanogens (*Methanimicrococcus*),

expressed very few genes differentially between the diets (Figure 2). Despite an overall

transcriptional abundance like that of key fiber degraders, *Desulfosarcina* did not differentially

express any genes across the diets, while *Frigididesulfovibrio* differentially expressed only 60

genes. None of the significantly enriched genes for *Frigididesulfovibrio* were related to sulfate

reduction. Of the differentially expressed genes, most were enriched in the tuna diet (28 out of 60

genes) compared to each of the other diets, including three genes associated with the taurine

metabolism pathway (K03851, K03852). *Methanimicrococcus* differentially expressed few

genes, with most gene expression changes identified in the bran vs. butter comparison (17 genes)

(Figure 2). Ten of these genes were transcribed more in the bran diet including four hypothetical

proteins, two coenzyme F420 hydrogenase subunits (frhA and frhG), cobaltochelatase subunit

CobN, oxidoreductase hydrogenase small subunit, and respiratory nitrate reductase subunit

gamma. Conversely, there were seven genes that were transcribed more in the butter diet

compared to bran, including methanol:5-hydroxybenzimidazolylcobamide Co-methyltransferase,

monomethylamine corrinoid protein, ethanolamine permease, acetate uptake transporter, and

peroxiredoxin (Table S6).

High-fat diet induces community stress response

Second to the high fiber diet, many of cockroach gut bacteria exhibited strong

transcriptional responses to the butter (high-fat) diet, including some taxa that showed minimal

responses to other dietary treatments (Figure 2). Across all taxa, there were 1130 genes that were

significantly differentially expressed between butter and other diets. Of these genes, 38% (410

genes) were enriched in the butter diet compared to the other diets. Many of these genes were

annotated as chaperones and folding catalysts including DnaJK chaperones, heat-shock proteins,

peroxiredoxin, and thioredoxin (Figure S9). 11 out of 18 taxa exhibited significantly increased

expression in at least one of these transcripts, and 6 of the remaining taxa showed increases

below the significance threshold.

Discussion

In this study, we examined cockroach gut microbiome transcriptional activity under

diverse host diets to better understand metabolic responses to changing dietary inputs and how

they may support gut microbiome homeostasis. Our previous work showed that the American

cockroach gut microbiome demonstrates high taxonomic stability following dietary shifts [33],

even following such extreme treatments as starvation for up to 60 days. Much of the stability

observed in the American cockroach bears similarities to the long-term taxonomic stability seen

in humans [20,28-30]. However, factors contributing to this stability are poorly understood.

Metagenomic analysis supported 16S rRNA-based findings that the cockroach gut

microbiome showed high taxonomic stability across diets (Figure 1). However, we observed

diet-driven changes in both the relative abundance of different taxonomic groups in the

metatranscriptome and the transcriptional profiles of individual organisms. *Bacteroides*,

Dysgonomonas, and *Parabacteroides* have the greatest plasticity in gene expression across host

diets, suggesting that they play an important role in shaping gut microbiome responses to host

diet (Figure 2). These taxa have previously been hypothesized to be primary fiber degraders

based on the number of CAZymes (carbohydrate active enzymes) present in their genomes and

the level at which these CAZymes are transcribed (Figure 3) [42].

Analysis of transcripts annotated as CAZymes showed diet-associated differences in

polysaccharide utilization profiles. The oat bran diet had the most dietary fiber content (Table 1)

and contains arabinoxylan along with beta-glucans [65]. We found that this bran diet led to high

expression of CAZyme families associated with arabinoxylan utilization. Highly transcribed

CAZymes including enzymes associated with side chain removal and degradation of main chain

xylose-xylose bonds, including GH2, GH3, GH10 and CE1. Several *Bacteroides* isolates from

the human gut are also known to highly express these xylanase CAZyme families when grown

on xylan [66-68].

The tuna diet was associated with unexpectedly high transcription of CAZymes

associated with pectin utilization, including CE8 (pectin methyl-esterase), which has been shown

to be particularly important in pectin metabolism by *Bacteroides thetaiotaomicron* [69]. We note

that the canned tuna used includes vegetable broth as a flavor enhancer and hypothesize that gut

bacteria may be responding to pectin present in this additive. In addition, there have been reports

of rhamnose, xylose, mannose, glucuronic acid, and galacturonic acid in processing Skipjack

tuna by-products, which suggests that glycans naturally present in tuna may be inducing

expression of CAZymes typically associated with pectin degradation [70].

The butter diet was associated with a decrease of CAZyme transcripts (Figure 3) as well

as widespread transcriptional upregulation of stress response genes including RND efflux pumps,

oxidative stress, and heat shock proteins (Figure S9). Two notable proteins that showed increased

expression in the butter diet, GroEL and dnaK, are known to play an important role in bacterial

survival during chemical, heat, or oxidative stress [71-73]. While additional work is needed to

determine if high fat diets induce host oxidative stress in the cockroach, high-fat diets in mouse

models have been associated with oxidative stress [74] and changes in redox state in gut

epithelial cells [75,76]. The butter diet in the cockroach also appears to reduce gut microbiome

stability, with blooms of facultative anaerobes such as *Enterococcaceae* and *Lactobacillaceae*

(*Leuconostocaceae*) observed in metagenomic analyses of a subset of cockroaches eating butter

(Figure S4-S5). These results are notably similar to studies that have found blooms of organisms

[18,77,78] and growth of facultative anaerobes, including *Enterococcaceae* and *Lactobacillaceae*

spp. in mice eating high-fat diets [79-81]. Studies in humans have also indicated greater

variability in community profiles of individuals eating high-fat diets [82]. Together, these results

are consistent with a pattern of high fat diet causing oxidative stress in the gut as well as blooms

of oxygen-tolerant microbiota and a reduction in the abundance and activity of fiber degrading

bacteria. The remarkable similarity in responses to high fat diet between cockroaches and

mammalian models suggests that the cockroach gut may represent a useful model for exploring

these linkages further.

A major goal of this work was to identify microbial metabolic activities that support the

observed persistence of diverse gut microbes under starvation conditions as well as other diets

lacking preferred growth substrates. A key finding was that overall CAZyme transcription

remained relatively high during host starvation, including strong and stable transcription of

CAZymes for utilizing host glycans (Figure 3). This signal was particularly notable in

Bacteroides, with eight CAZyme families associated with host glycan utilization showing

significantly higher expression under starvation compared to bran and/or butter (Figure 4C). The

utilization of host glycans has been widely reported in the gut symbionts of mammals such as

humans and mice, which helped identify host mucins as an important substrate for gut fiber

degraders in the absence of microbially accessible polysaccharides [83-85]. More broadly, host

mucins have been shown to play a role in shaping symbiotic microbiota across the animal

kingdom, from invertebrates such as corals, squid light organs, and insects to higher animals

including zebra fish, mice, and humans [86-90]. However, it was not immediately clear that this

strategy would be applicable to the cockroach hindgut microbiome, as the insect hindgut is

surrounded by a chitinous ectoderm and lacks a mammalian-like mucus layer [91,92].

A possible source of host glycans for the insect hindgut microbiota is the peritrophic

matrix, which acts as a barrier between the midgut absorptive tissues and the microbiota,

analogous to a mammalian mucus layer [93]. In cockroaches, the peritrophic matrix is

continually produced in the dorsal midgut and extruded along with digested food into the hindgut

[91,92]. While peritrophic matrices can contain both O- and N-glycans [94], our data suggests a

greater role of N-glycans in supporting the insect hindgut microbiota. We observed high

expression of GH38 alpha-mannosidases, which have previously been shown to act on N-glycans

[95,96]. Additionally, we identified other CAZymes associated with fucosidase, glucosidase, and

galactosidase activity. Analysis of N-glycans in cockroach midgut preparations confirmed the

presence of glycans containing these residues (Figure S10). These structures are also consistent

with a published analysis of N-glycans in the peritrophic matrix of the Colorado potato beetle

[94]. While expression of CAZymes active on O-glycans was much lower, *Bacteroides* did

express genes to break down bonds present in O-glycan analyses of cockroach midguts (Figure

S11). Overall, these results suggest that mannosylated N-glycans in the peritrophic matrix play a

critical role in shaping the hindgut microbiota of cockroaches. Our hypothesis that the insect

peritrophic matrix is a mechanism by which the host can shape the hindgut (as well as midgut)

microbiota has implications for understanding hindgut microbial communities in other insects.

As an example, honeybees represent another well-studied insect system with both a peritrophic

matrix and a hindgut microbiota critical to host health [97-99]. An examination of data from the

insect hindgut metagenome also suggests the presence of glycan-targeting CAZymes in the

honeybee gut microbiota [100].

Finally, we note that for organisms further down the 'microbial food web', transcriptional

activity remained remarkably stable across diets. While primary degraders exhibit large

transcriptional shifts in response to changing diets, these efforts release a limited set of sugar

monomers that, in turn, are processed via central metabolic pathways into a limited suite of

fermentation end products. Despite having a comparable level of coverage in the

metatranscriptome to that seen in some fiber degrading bacteria, sulfate reducing bacteria and

methanogens, in particular, showed remarkable stability in transcriptional activity across diets.

These organisms primarily survive on hydrogen and other fermentation products (short chain

fatty acids and/or methylamines) released during the degradation of fiber by primary degraders.

Our results suggest that the transcriptional plasticity exhibited by the fiber degraders is sufficient

to maintain production of these end products at levels that support the continued persistence and

activity of the rest of the gut microbiome, leading to the observed overall stability in gut

microbiome composition seen overall.

While these results may seem contradictory to the many studies showing dietary impacts

on gut microbiome composition, we would like to emphasize two points. First, we note that

overall stability in gut microbiome seems to vary among organisms, with laboratory mice in

particular showing strong responses to diet while the human gut microbiota frequently does not

exhibit large structural changes that are greater than pre-existing inter-individual variability

[28,31,32]. Further, many measures of microbial community composition and responses to

dietary shifts are primarily aimed at identifying differences, while ignoring similarities. The goal

of this study was to emphasize instead mechanisms that support gut microbiome homeostasis in

the face of dietary differences.

Conclusions

This work emphasized the importance of fiber degraders in supporting overall gut

microbiome homeostasis. In our model, transcriptional plasticity in fiber degrading bacteria and

– to a lesser extent– secondary fermenters was sufficient to maintain key hydrogenotrophic taxa

at stable relative abundance and minimal alterations to their overall transcriptional profiles. Our

results also suggest a role for host glycans present in the peritrophic matrix and passed into the

hindgut as part of normal digestive function in providing a critical nutritional subsidy when the

host diet lacks sufficient fiber sources. Together, these results provide new insight into host and

microbial mechanisms supporting gut microbiome homeostasis in omnivorous cockroaches.

Acknowledgements:

This work was supported by the National Institute of General Medical Sciences of the National

Institutes of Health (NIH) under award number R35GM133789. Glycan analyses were supported

425 under the NIH award R24GH137782. We would like to thank Lauren Hudgens for help with the

426 glycan analyses.

427 **Data Availability:**

428 Data associated with this study are available from NCBI SRA under BioProject accession

429 number PRJNA994902. Scripts and workflow for the metatranscriptomic analyses can be found

430 here: https://github.com/adepoy1/cockroach_metatranscriptome_manuscript.

431 **Tables:**

432 Table 1. Nutrient information for 100g serving of diet treatment. All cockroaches were raised on
433 our laboratory control diet of dog food, composed of 21% protein, 8% fat, and 6% fiber.
434

Diet Treatment	Calories	Protein (g)	Carbohydrate (g)	Fat (g)	Fiber (g)
Bran (high-fiber)	375	17.5	67.5	5	17.5
Butter (high-fat)	714	0	0	79	0
Tuna (high-protein)	107	27	0	1	0
Starvation (starve)	N/A	N/A	N/A	N/A	N/A

435
436 **Figure Captions:**
437 **Figure 1.** Taxonomic composition of metagenomes and metatranscriptomes. A) Bar plot of
438 relative abundance (%) of families in the metagenome and metatranscriptome datasets. Each bar
439 represents an average relative abundance of 5 replicates (metagenome) or 4 replicates
440 (metatranscriptome) per diet. Families with a maximum relative abundance of at least 5% in any
441 sample of either the metatranscriptome or metagenome dataset are included in the taxonomy. B)
442 Non-metric multidimensional scaling (NMDS) using a weighted Bray-Curtis dissimilarity metric
443 for the metagenome and metatranscriptome datasets, labeled accordingly.
444
445 **Figure 2.** Number of differentially expressed genes identified by DESeq2. Each number within
446 each square is the number of differentially expressed genes when those two diets were compared.
447
448 **Figure 3.** CAZyme gene expression across the gut microbiome. A) Sankey diagram of the
449 summed relative abundance of CAZyme genes (> 0.01% relative abundance) associated with the
450 breakdown of substrates. Relative abundance was calculated as a proportion of all transcripts in
451 each sample to allow for comparisons between taxa. Lines of the Sankey diagram represent the
452 summed relative abundance of the average across all diets, while the stacked bar plots represent
453 summed relative abundance of transcripts in each fiber substrate averaged for each diet. B) Bar
454 plot of transcription of B) all, C) xylan, D) pectin, and E) host glycan CAZymes across all diets.
455 Note that different scales were used on the y-axes of the bar plots to account for differences in
456 expression levels of each of the substrate CAZyme families.
457

Figure 4. Xylan (A), pectin (B), and host glycan (C) utilization by *Bacteroides*. Bar plots of average (n = 4) relative abundance (%) of CAZyme families and KEGG orthologs associated with polysaccharide degradation. Each bar plot is labelled with its associated substrate of degradation: xylose (D), rhamnose (E), mannose (F), arabinose (G), galacturonate (H), GlcNAc (I), galactose (K), and fucose (L). Note each bar plot has a different y-axis scale to emphasize the difference between expression levels between diets. Monosaccharide degradation following its respective polysaccharide are together in a box, while overlapping monosaccharides are separately boxed and colored. Abbreviations, galacturonate: KDG = 2-Dehydro-3-deoxy-D-gluconate, KDPG = 2-Dehydro-3-deoxy-6-phospho-D-glyconate, G3P = D-Glyceraldehyde 3-phosphate. All monosaccharide degradation pathways feed into central metabolism pathways (J) shown in lower left bar plot.

References

1. Sommer F, Backhed F. The gut microbiota--masters of host development and physiology. *Nat Rev Microbiol*. 2013;11(4):227-38. doi:10.1038/nrmicro2974

2. Engel P, Moran NA. The gut microbiota of insects - diversity in structure and function. *FEMS Microbiol Rev*. 2013;37(5):699-735. doi:10.1111/1574-6976.12025

3. Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI. Worlds within worlds: evolution of the vertebrate gut microbiota. *Nat Rev Microbiol*. 2008;6:776-788.

4. Hooper LV, Gordon JI. Commensal Host-Bacterial Relationships in the Gut. *Science*. 2001;292(5519):1115-1118.

5. Dillon RJ, Dillon VM. The gut bacteria of insects: nonpathogenic interactions. *Annu Rev Entomol*. 2004;49:71-92. doi:10.1146/annurev.ento.49.061802.123416

6. Kurilshikov A, Medina-Gomez C, Bacigalupe R, et al. Large-scale association analyses identify host factors influencing human gut microbiome composition. *Nat Genet*. 2021;53(2):156-165. doi:10.1038/s41588-020-00763-1

7. Boccuto L, Tack J, Ianiro G, Abenavoli L, Scarpellini E. Human Genes Involved in the Interaction between Host and Gut Microbiome: Regulation and Pathogenic Mechanisms. *Genes*. 2023;14(4):857. doi:10.3390/genes14040857

8. Sutherland DB, Suzuki K, Fagarasan S. Fostering of advanced mutualism with gut microbiota by Immunoglobulin A. *Immunological Reviews*. 2016;270(1):20-31. doi:10.1111/imr.12384

9. Peterson DA, McNulty NP, Guruge JL, Gordon JI. IgA Response to Symbiotic Bacteria as a Mediator of Gut Homeostasis. *Cell Host & Microbe*. 2007;2(5):328-339. doi:10.1016/j.chom.2007.09.013

10. Meyer-Hoffert U, Hornef MW, Henriques-Normark B, et al. Secreted enteric antimicrobial activity localises to the mucus surface layer. *Gut*. 2008;57(6):764. doi:10.1136/gut.2007.141481

11. David LA, Maurice CF, Carmody RN, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature*. 2014;505(7484):559-563. doi:10.1038/nature12820

12. Cotillard A, Kennedy SP, Kong LC, et al. Dietary intervention impact on gut microbial gene richness. *Nature*. 2013;500(7464):585-588. doi:10.1038/nature12480

13. Walker AW, Ince J, Duncan SH, et al. Dominant and diet-responsive groups of bacteria within the human colonic microbiota. *The ISME Journal*. 2011;5(2):220-230. doi:10.1038/ismej.2010.118

14. Drasar BS, Crowther JS, Goddard P, et al. The relation between diet and the gut microflora in man. *Proc Nutr Soc*. 1973;32(2):49-52. doi:10.1079/pns19730014

15. Fava F, Gitau R, Griffin BA, Gibson GR, Tuohy KM, Lovegrove JA. The type and quantity of dietary fat and carbohydrate alter faecal microbiome and short-chain fatty acid excretion in a metabolic syndrome 'at-risk' population. *International Journal of Obesity*. 2013;37(2):216-223. doi:10.1038/ijo.2012.33

16. Hildebrandt MA, Hoffmann C, Sherrill-Mix SA, et al. High-fat diet determines the composition of the murine gut microbiome independently of obesity. *Gastroenterology*. 2009;137(5):1716-24 e1-2. doi:10.1053/j.gastro.2009.08.042

17. Turnbaugh PJ, Ridaura VK, Faith JJ, Rey FE, Knight R, Gordon JI. The Effect of Diet on the Human Gut Microbiome: A Metagenomic Analysis in Humanized Gnotobiotic Mice. *Science Translational Medicine*. 2009;1(6):6ra14-6ra14. doi:10.1126/scitranslmed.3000322

18. Turnbaugh PJ, Bäckhed F, Fulton L, Gordon JI. Diet-Induced Obesity Is Linked to
Marked but Reversible Alterations in the Mouse Distal Gut Microbiome. *Cell Host & Microbe*.
2008;3(4):213-223. doi:10.1016/j.chom.2008.02.015
19. De Filippo C, Cavalieri D, Di Paola M, et al. Impact of diet in shaping gut microbiota
revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci*
USA. 2010;107(33):14691-6. doi:10.1073/pnas.1005963107
20. Faith JJ, Guruge JL, Charbonneau M, et al. The long-term stability of the human gut
microbiota. *Science*. 2013;341(6141):1237439. doi:10.1126/science.1237439
21. Statovci D, Aguilera M, MacSharry J, Melgar S. The Impact of Western Diet and
Nutrients on the Microbiota and Immune Response at Mucosal Interfaces. Review. *Front*
Immunol. 2017;8doi:10.3389/fimmu.2017.00838
22. Bortolin RC, Vargas AR, Gasparotto J, et al. A new animal diet based on human Western
diet is a robust diet-induced obesity model: comparison to high-fat and cafeteria diets in term of
metabolic and gut microbiota disruption. *International Journal of Obesity*. 2018;42(3):525-534.
doi:10.1038/ijo.2017.225
23. Panasevich MR, Meers GM, Linden MA, et al. High-fat, high-fructose, high-cholesterol
feeding causes severe NASH and cecal microbiota dysbiosis in juvenile Ossabaw swine. *Am J*
Physiol Endocrinol Metab. 2018;314(1):E78-E92. doi:10.1152/ajpendo.00015.2017
24. Amato KR, Yeoman CJ, Cerda G, et al. Variable responses of human and non-human
primate gut microbiomes to a Western diet. *Microbiome*. 2015;3(1)doi:10.1186/s40168-015-
0120-7
25. Jensen N, Maldonado-Gomez M, Krishnakumar N, et al. Dietary fiber monosaccharide
content alters gut microbiome composition and fermentation. *Appl Environ Microbiol*.
2024;90(8)
26. Tanes C, Bittinger K, Gao Y, et al. Role of dietary fiber in the recovery of the human gut
microbiome and its metabolome. *Cell Host & Microbe*. 2021;29(3):394-407 e5.
doi:10.1016/j.chom.2020.12.012
27. Vinelli V, Biscotti P, Martini D, et al. Effects of Dietary Fibers on Short-Chain Fatty
Acids and Gut Microbiota Composition in Healthy Adults: A Systematic Review. *Nutrients*.
2022;14(13)doi:10.3390/nu14132559
28. Chen L, Wang D, Garmaeva S, et al. The long-term genetic stability and individual
specificity of the human gut microbiome. *Cell*. 2021;184(9):2302-2315.e12.
doi:10.1016/j.cell.2021.03.024
29. Mehta RS, Abu-Ali GS, Drew DA, et al. Stability of the human faecal microbiome in a
cohort of adult men. *Nat Microbiol*. 2018;3(3):347-355. doi:10.1038/s41564-017-0096-0
30. Wu GD, Chen J, Hoffmann C, et al. Linking Long-Term Dietary Patterns with Gut
Microbial Enterotypes. *Science*. 2011;334:105-108.
31. Schloissnig S, Arumugam M, Sunagawa S, et al. Genomic variation landscape of the
human gut microbiome. *Nature*. 2013;493(7430):45-50. doi:10.1038/nature11711
32. Garud NR, Good BH, Hallatschek O, Pollard KS. Evolutionary dynamics of bacteria in
the gut microbiome within and across hosts. *PLOS Biology*. 2019;17(1):e3000102.
doi:10.1371/journal.pbio.3000102
33. Tinker KA, Ottesen EA. The Core Gut Microbiome of the American Cockroach,
Periplaneta americana, Is Stable and Resilient to Dietary Shifts. *Appl Environ Microbiol*. Nov
15 2016;82(22):6603-6610. doi:10.1128/AEM.01837-16

34.   Moreau CS, Rubin BER. Diversity and Persistence of the Gut Microbiome of the Giant
Neotropical Bullet Ant. *Integr Comp Biol*. 2017;57(4):682-689. doi:10.1093/icb/icx037
35.   Súarez-Moo P, Cruz-Rosales M, Ibarra-Laclette E, Desgarennes D, Huerta C, Lamelas A.
Diversity and Composition of the Gut Microbiota in the Developmental Stages of the Dung
Beetle *Copris incertus* Say (Coleoptera, Scarabaeidae). *Front Microbiol*. 2020;11:1698.
doi:10.3389/fmicb.2020.01698
36.   Vikram S, Arneodo JD, Calcagno J, et al. Diversity structure of the microbial
communities in the guts of four neotropical termite species. *PeerJ*. 2021;9:e10959.
doi:10.7717/peerj.10959
37.   Ohkuma M, Brune A. Diveristy, Structure, and Evolution of the Termite Gut Microbial
Community. In: Bignell DE, Roisin Y, Lo N, eds. *Biology of Termites: A modern Synthesis*.
Springer; 2010:413-438.
38.   Tinker KA, Ottesen EA. Differences in Gut Microbiome Composition Between
Sympatric Wild and Allopatric Laboratory Populations of Omnivorous Cockroaches. *Front
Microbiol*. 2021;12:703785. doi:10.3389/fmicb.2021.703785
39.   Mikaelyan A, Thompson CL, Hofer MJ, Brune A. Deterministic Assembly of Complex
Bacterial Communities in Guts of Germ-Free Cockroaches. *Appl Environ Microbiol*.
2016;82(4):1256-63. doi:10.1128/AEM.03700-15
40.   Schauer C, Thompson C, Brune A. Pyrotag sequencing of the gut microbiota of the
cockroach *Shelfordella lateralis* reveals a highly dynamic core but only limited effects of diet on
community structure. *PLoS One*. 2014;9(1):e85861. doi:10.1371/journal.pone.0085861
41.   Schauer C, Thompson CL, Brune A. The bacterial community in the gut of the Cockroach
Shelfordella lateralis reflects the close evolutionary relatedness of cockroaches and termites.
Appl Environ Microbiol. 2012;78(8):2758-67. doi:10.1128/AEM.07788-11
42.   Dukes HE, Tinker KA, Ottesen EA. Disentangling hindgut metabolism in the American
cockroach through single-cell genomics and metatranscriptomics. Original Research. *Front
Microbiol*. 2023;14doi:10.3389/fmicb.2023.1156809
43.   Lampert N, Mikaelyan A, Brune A. Diet is not the primary driver of bacterial community
structure in the gut of litter-feeding cockroaches. *BMC Microbiology*.
2019;19(1)doi:10.1186/s12866-019-1601-9
44.   Pérez-Cobas AE, Maiques E, Angelova A, Carrasco P, Moya A, Latorre A. Diet shapes
the gut microbiota of the omnivorous cockroach *Blattella germanica*. *FEMS Microbiol Ecol*.
2015;91(4)doi:10.1093/femsec/fiv022
45.   Glenn TC, Nilsen RA, Kieran TJ, et al. Adapterama I: universal stubs and primers for 384
unique dual-indexed or 147,456 combinatorially-indexed Illumina libraries (iTru & iNext).
PeerJ. 2019;7:e7755. doi:10.7717/peerj.7755
46.   Kopylova E, Noé L, Touzet H. SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. *Bioinformatics*. 2012;28(24):3211-3217.
doi:10.1093/bioinformatics/bts611
47.   Li S, Zhu S, Jia Q, et al. The genomic and functional landscapes of developmental
plasticity in the American cockroach. *Nat Commun*. 2018;9(1)doi:10.1038/s41467-018-03281-1
48.   Sabree ZL, Kambhampati S, Moran NA. Nitrogen recycling and nutritional provisioning
by *Blattabacterium*, the cockroach endosymbiont. *Proc Natl Acad Sci USA*.
2009;106(46):19521-19526. doi:10.1073/pnas.0907504106

49.   Ottesen EA, Young CR, Eppley JM, et al. Pattern and synchrony of gene expression
among sympatric marine microbial populations. *Proc Natl Acad Sci USA*. 2013;110(6):E488-
E497. doi:doi:10.1073/pnas.1222099110
50.   Eren AM, Kiefl E, Shaiber A, et al. Community-led, integrated, reproducible multi-omics
with anvi'o. *Nat Microbiol*. 2021;6(1):3-6. doi:10.1038/s41564-020-00834-3
51.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.
Nat Methods. 2015;12(1):59-60. doi:10.1038/nmeth.3176
52.   Van Dongen S. Graph Clustering Via a Discrete Uncoupling Process. *SIAM Journal on
Matrix Analysis and Applications*. 2008;30(1):121-141. doi:10.1137/040608635
53.   Enright AJ, Van Dongen S, Ouzounis CA. An efficient algorithm for large-scale detection
of protein families. *Nucleic Acids Res*. 2002;30(7):1575-1584.
54.   Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids
Res*. 2000;28(1):27-30. doi:10.1093/nar/28.1.27
55.   Tatusov RL, Galperin MY, Natale DA, Koonin EV. The COG database: a tool for
genome-scale analysis of protein functions and evolution. *Nucleic Acids Res*. 2000;28(1):33-36.
doi:10.1093/nar/28.1.33
56.   Galperin MY, Wolf YI, Makarova KS, Vera Alvarez R, Landsman D, Koonin EV. COG
database update: focus on microbial diversity, model organisms, and widespread pathogens.
Nucleic Acids Res. 2020;49(D1):D274-D281. doi:10.1093/nar/gkaa1018
57.   Drula E, Garron M-L, Dogan S, Lombard V, Henrissat B, Terrapon N. The carbohydrate-
active enzyme database: functions and literature. *Nucleic Acids Res*. 2021;50(D1):D571-D577.
doi:10.1093/nar/gkab1045
58.   Saier MH, Reddy VS, Moreno-Hagelsieb G, et al. The Transporter Classification
Database (TCDB): 2021 update. *Nucleic Acids Res*. 2021;49(D1):D461-D467.
doi:10.1093/nar/gkaa1004
59.   O'Leary NA, Wright MW, Brister JR, et al. Reference sequence (RefSeq) database at
NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res*.
2016;44(D1):D733-D745. doi:10.1093/nar/gkv1189
60.   Tatusova T, DiCuccio M, Badretdin A, et al. NCBI prokaryotic genome annotation
pipeline. *Nucleic Acids Res*. 2016;44(14):6614-6624. doi:10.1093/nar/gkw569
61.   *vegan: Community Ecology Package*. Version R package version 2.6-4. 2022.
https://CRAN.R-project.org/package=vegan
62.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology*. 2014;15(12)doi:10.1186/s13059-014-0550-8
63.   Zhu A, Ibrahim JG, Love MI. Heavy-tailed prior distributions for sequence count data:
removing the noise and preserving large differences. *Bioinformatics*. 2018;35(12):2084-2092.
doi:10.1093/bioinformatics/bty895
64.   Zheng J, Hu B, Zhang X, et al. dbCAN-seq update: CAZyme gene clusters and substrates
in microbiomes. *Nucleic Acids Res*. 2023;51(D1):D557-D563. doi:10.1093/nar/gkac1068
65.   Westerlund E, Andersson R, Åman P. Isolation and chemical characterization of water
soluble mixed-linked ß-glucans and arabinoxylans in oat milling fractions. *Carbohydr Polym*.
1993;20:115-123.
66.   Zhang M, Cheken JR, Dodd D, et al. Xylan utilization in human gut commensal bacteria
is orchestrated by unique modular organization of polysaccharide-degrading enzymes. *Proc Natl
Acad Sci USA*. 2014;111(35):E3708-E3717. doi:10.1073/pnas.1406156111

67. Wang K, Pereira GV, Cavalcante JJ, Zhang M, Mackie R, Cann I. *Bacteroides intestinalis*
DSM 17393, a member of the human colonic microbiome, upregulates multiple endoxylanases
during growth on xylan. *Scientific Reports*. 2016;6:34360. doi:10.1038/srep34360
68. Despres J, Forano E, Lepercq P, et al. Xylan degradation by the human gut *Bacteroides*
xylanisolvens XB1A(T) involves two distinct gene clusters that are linked at the transcriptional
level. *BMC Genomics*. 2016;17:326. doi:10.1186/s12864-016-2680-8
69. Duan CJ, Basle A, Liberato MV, et al. Ascertaining the biochemical function of an
essential pectin methylesterase in the gut microbe *Bacteroides thetaiotaomicron. J Biol Chem*.
2020;295(52):18625-18637. doi:10.1074/jbc.RA120.014974
70. Naghdi S, Rezaei M, Tabarsa M, Abdollahi M. Ultrasonic-assisted enzymatic extraction
of sulfated polysaccharide from Skipjack tuna by-products. *Ultrasonics Sonochemistry*.
2023;95:106385. doi:10.1016/j.ultsonch.2023.106385
71. Hennequin C, Porcheray F, Waligora-Dupriet A, et al. GroEL (Hsp60) of *Clostridium*
difficile is involved in cell adherance. *Microbiology*. 2001;147:87-96.
72. Gomes SL, Simão RCG. Stress Responses: Heat. *Reference Module in Biomedical*
Sciences. 2014.
73. Calloni G, Chen T, Schermann SM, et al. DnaK functions as a central hub in the *E. coli*
chaperone network. *Cell Reports*. 2012;1(3):251-64. doi:10.1016/j.celrep.2011.12.007
74. Wypych A, Ożgo M, Bernaciak M, et al. Effect of feeding high fat diets differing in fatty
acid composition on oxidative stress markers and protein expression in mouse kidney. *J Anim*
Feed Sci. 2024;33(2):170-184. doi:10.22358/jafs/175920/2024
75. Enriquez JR, McCauley HA, Zhang KX, et al. A dietary change to a high-fat diet initiates
a rapid adaptation of the intestine. *Cell Reports*. 2022;41(7):111641.
doi:10.1016/j.celrep.2022.111641
76. Qiao Y, Sun J, Ding Y, Le G, Shi Y. Alterations of the gut microbiota in high-fat diet mice
is strongly linked to oxidative stress. *Applied Microbiology and Biotechnology*.
2013;97(4):1689-1697. doi:10.1007/s00253-012-4323-6
77. Hildebrandt MA, Hoffmann C, Sherrill–Mix SA, et al. High-Fat Diet Determines the
Composition of the Murine Gut Microbiome Independently of Obesity. *Gastroenterology*.
2009;137(5):1716-1724.e2. doi:10.1053/j.gastro.2009.08.042
78. Devkota S, Chang EB. Interactions between Diet, Bile Acid Metabolism, Gut Microbiota,
and Inflammatory Bowel Diseases. *Digestive Diseases*. 2015;33(3):351-356.
doi:10.1159/000371687
79. Heisel T, Montassier E, Johnson A, et al. High-Fat Diet Changes Fungal Microbiomes
and Interkingdom Relationships in the Murine Gut. *mSphere*. 2017;2(5):10.1128/msphere.00351-
17. doi:doi:10.1128/msphere.00351-17
80. Xiao L, Sonne SB, Feng Q, et al. High-fat feeding rather than obesity drives taxonomical
and functional changes in the gut microbiota in mice. *Microbiome*. 2017;5(1):43.
doi:10.1186/s40168-017-0258-6
81. Lin H, An Y, Hao F, Wang Y, Tang H. Correlations of Fecal Metabonomic and
Microbiomic Changes Induced by High-fat Diet in the Pre-Obesity State. *Scientific Reports*.
2016;6(1):21618. doi:10.1038/srep21618
82. Wang B, Kong Q, Li X, et al. A High-Fat Diet Increases Gut Microbiota Biodiversity and
Energy Expenditure Due to Nutrient Difference. *Nutrients*. 2020;12(10):3197.
doi:10.3390/nu12103197

83.   Desai MS, Seekatz AM, Koropatkin NM, et al. A Dietary Fiber-Deprived Gut Microbiota
Degrades the Colonic Mucus Barrier and Enhances Pathogen Susceptibility. *Cell*.
2016;167(5):1339-1353 e21. doi:10.1016/j.cell.2016.10.043
84.   Sonnenburg JL, Xu J, Leip DD, et al. Glycan Foraging in Vivo by an Intestine-Adapted
Bacterial Symbiont. *Science*. 2005;307:1955-1959.
85.   Martens EC, Chiang HC, Gordon JI. Mucosal glycan foraging enhances fitness and
transmission of a saccharolytic human gut bacterial symbiont. *Cell Host & Microbe*.
2008;4(5):447-57. doi:10.1016/j.chom.2008.09.007
86.   Nyholm SV, Deplancke B, Gaskins HR, Apicella MA, McFall-Ngai MJ. Roles of *Vibrio
fischeri* and nonsymbiotic bacteria in the dynamics of mucus secretion during symbiont
colonization of the *Euprymna scolopes* light organ. *Appl Environ Microbiol*. 2002;68(10):5113-
22. doi:10.1128/AEM.68.10.5113-5122.2002
87.   Smith TJ, Sundarraman D, Melancon E, Desban L, Parthasarathy R, Guillemin K. A
mucin-regulated adhesin determines the spatial organization and inflammatory character of a
bacterial symbiont in the vertebrate gut. *Cell Host & Microbe*. 2023;31(8):1371-1385 e6.
doi:10.1016/j.chom.2023.07.003
88.   Berkhout MD, Ioannou A, de Ram C, Boeren S, Plugge CM, Belzer C. Mucin-driven
ecological interactions in an in vitro synthetic community of human gut microbes. *Glycobiology*.
2024;34(12)doi:10.1093/glycob/cwae085
89.   Moriyama M, Hayashi T, Fukatsu T. A mucin protein predominantly expressed in the
female-specific symbiotic organ of the stinkbug *Plautia stali*. *Scientific Reports*.
2022;12(1):7782. doi:10.1038/s41598-022-11895-1
90.   Krediet CJ, Ritchie KB, Paul VJ, Teplitski M. Coral-associated micro-organisms and
their roles in promoting coral health and thwarting diseases. *Proc R Soc B*.
2013;280(1755):20122328. doi:10.1098/rspb.2012.2328
91.   Bell WJ, Adiyodi KG, eds. *The American Cockroach*. Chapman and Hall; 1982.
92.   Terra WR. The origin and functions of the insect peritrophic membrane and peritrophic
gel. *Arch Insect Biochem Physiol*. 2001;47(2):47-61. doi:10.1002/arch.1036
93.   Tellam RL, Wigffels G, Willadsen P. Peritrophic matrix proteins. *Insect Biochem Mol
Biol*. 1999;29:87-101.
94.   Liu D, De Schutter K, Smargiasso N, De Pauw E, Van Damme EJM, Smagghe G. The N-
glycan profile of the peritrophic membrane in the Colorado potato beetle larva (*Leptinotarsa
decemlineata*). *J Insect Physiol*. 2019;115:27-32. doi:10.1016/j.jinsphys.2019.03.009
95.   Suits MD, Zhu Y, Taylor EJ, et al. Structure and kinetic investigation of *Streptococcus
pyogenes* family GH38 alpha-mannosidase. *PLoS One*. 2010;5(2):e9006.
doi:10.1371/journal.pone.0009006
96.   Cordeiro RL, Santos CR, Domingues MN, et al. Mechanism of high-mannose N-glycan
breakdown and metabolism by *Bifidobacterium longum*. *Nat Chem Biol*. 2023;19(2):218-229.
doi:10.1038/s41589-022-01202-4
97.   Teixeira ADD, Marques-Araujo S, Zanuncio JC, Serrao JE. Peritrophic membrane origin
in adult bees (Hymenoptera): immunolocalization. *Micron*. 2015;68:91-97.
doi:10.1016/j.micron.2014.09.009
98.   Raymann K, Moran NA. The role of the gut microbiome in health and disease of adult
honey bee workers. *Curr Opin Insect Sci*. 2018;26:97-104. doi:10.1016/j.cois.2018.02.012

99.   Zheng H, Powell JE, Steele MI, Dietrich C, Moran NA. Honeybee gut microbiota
promotes host weight gain via bacterial metabolism and hormonal signaling. *Proc Natl Acad Sci*
USA. 2017;114(18):4775-4780. doi:10.1073/pnas.1701819114
100.   Engel P, Martinson VG, Moran MA. Functional diversity within the simple gut
microbiota of the honey bee. *Proc Natl Acad Sci USA*. 2012;109(27):11002-11007.
doi:10.1073/pnas.1202970109
743