Navigating the Fitness Landscapes of Plasmodium falciparum Dihydrofolate Reductase: Evolutionary Insights into Antifolate Resistance

Danny Muzata[1,2#], Dwipanjan Sanyal[3#], Shivram A[1,2], Deeptanshu Pandey[4], Soumyananda Chakraborti[4], Aneesh S. Chivukula[2], Ravi Jasuja[5], Vladimir N. Uversky[6], Daniel Zielinski[7], Prajna D. Upadhyay[2], Mark Albers[8], and Sourav Chowdhury[4*]

Affiliations:
[1] One Health Trust, Bangalore, India
[2] Department of Computer Science and Information Systems, Birla Institute of Technology and Science-Pilani, Hyderabad, India.
[3] Structural Biology and Bioinformatics Division, CSIR-Indian Institute of Chemical Biology, Kolkata, India
[4] Department of Biological Sciences, Birla Institute of Technology and Science-Pilani, Hyderabad, India.
[5] Research Program in Men's Health: Aging and Metabolism, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA
[6] USF Health Byrd Alzheimer's Research Institute, Morsani College of Medicine, University of South Florida, Tampa, FL 33612, USA
[7] Department of Bioengineering, University of California, San Diego, La Jolla, CA, USA
[8] Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, USA

[*]**Corresponding Author:**

Sourav Chowdhury
Department of Biological Sciences,
Office: J Block-109,
Birla Institute of Technology and Science, Hyderabad Campus,
Hyderabad, Telangana, India, 500078
Email: sourav.chowdhury@hyderabad.bits-pilani.ac.in

[#]*These authors contributed equally to the work.*

Abstract

The rapid emergence of drug resistance in malaria parasites poses a significant challenge to the efficacy of antifolate treatments. Traditional drug development approaches, which often rely on empirical screening with limited mechanistic insights, tend to overlook the complex evolutionary mechanisms that enable *Plasmodium falciparum* to evade drug inhibition while preserving enzyme functionality. In this study, we employed computational techniques to investigate the mutational landscape of dihydrofolate reductase (DHFR), focusing on regions essential for enzyme stability and resistance. Our analysis uncovered conserved residues essential for stability, mutation hotspots that enhance adaptability under drug pressure and co-evolving clusters revealing critical functional interdependencies. Through integrated approaches including mutational scanning, epistatic interaction modeling, and fitness trajectory mapping, we elucidated distinct evolutionary pathways that drive resistance. We were able to track the adaptive paths taken by wild-type residues upon mutation, revealing the steps required to reach high-fitness peaks within the rugged fitness landscape. These findings provide valuable insights into the molecular mechanisms of antifolate resistance. We suggest that future drug design should target co-evolving networks and conserved regions to support the development of next-generation therapies to overcome resistance.

Keywords: *Malaria*, *Plasmodium falciparum*, *Sequence space analysis*, *Protein fitness, Dihydrofolate reductase, Protein evolution, Antifolate resistance, Folate pathway, Drug design*

Introduction

To date, malaria remains one of the most challenging infectious diseases to control partly due to the parasite's ability to rapidly evolve resistance to frontline drugs[1,2]. Among the most challenging cases of resistance are mutations in *P. falciparum* dihydrofolate reductase (*pf*DHFR), which reduce the effectiveness of antifolate therapies[3,4]. Antifolate drugs have long been used in treating not only malaria but other diseases including bacterial infections and cancer[5,6]. These drugs exert their effects by targeting and inhibiting key enzymes in folate metabolism, a critical pathway for the synthesis of nucleotides and certain amino acids. In *P. falciparum*, several metabolic pathways are targeted by antimalarial drugs. One good example is the hemoglobin degradation pathway and the mitochondrial electron transport chain that is targeted by drugs such as chloroquine and artemisinin[2]. Heat shock proteins also present promising drug targets due to their essential roles in parasite survival and virulence across the various life cycle stages[1]. These alternative drug targets highlight the complexity of *P. falciparum* biology and emphasize the importance of targeting multiple pathways to combat resistance. However, the folate pathway, particularly DHFR, remains one of the most critical targets in antimalarial therapy due to its central role in DNA synthesis and repair[7–9].

In *P. falciparum*, DHFR catalyzes the NADPH-dependent reduction of dihydrofolate to tetrahydrofolate, an essential cofactor required for the synthesis of purines, thymidylate, and certain amino acids[8,10]. This reduction reaction makes DHFR an indispensable enzyme in the folate pathway and a prime target for antifolate drugs such as pyrimethamine and cycloguanil[8,11]. Single and multiple mutations at residue positions 16, 50, 51, 59, 108, and 164 of *pf*DHFR have been linked to antifolate resistance in malaria. Mutations at these sites confer resistance to pyrimethamine and cycloguanil by reducing its binding affinity to the protein rendering the drug less effective[3,12,13].

Despite significant research into DHFR mutations and their role in antifolate resistance, gaps remain in understanding how these mutations affect protein fitness, structure, and function at the molecular level. The concept of sequence space provides a theoretical framework for mapping all possible amino acid combinations in a protein and exploring the evolutionary constraints that shape functional regions. Sequence space analysis has proven valuable in uncovering

evolutionarily constrained regions as well as identifying conserved and flexible sites within
protein structures[14]. Thus, in this study, we carried out a thorough sequence space analysis of
DHFR protein to explore the evolution of antifolate resistance and its implications for drug
design. We developed an evolutionary dynamics simulation model to map adaptive fitness
trajectories. This approach integrates both deterministic and stochastic elements, simulating how
mutations traverse the fitness landscape while balancing short-term gains and long-term
evolutionary potential.

Results

Sequence space analysis

In this study, we explored the sequence space of *pf*DHFR to gain insights into how mutations
influence protein fitness and stability. This sequence-based study provided important insights
into the protein's evolutionary dynamics. It highlighted how mutations at critical positions
influence its stability, function, and mutational tolerance. Through careful analysis of sequence
conservation, co-evolution and network-based interactions, we inferred how sequence variability
could lead to significant functional consequences in the enzyme, especially in the context of
resistance-related mutations.

Figure 1: Sequence Space Analysis of DHFR Reveals Key Regions Under Evolutionary Constraints and Functional Adaptation. (A) Shannon Entropy plot shows sequence variability across the DHFR domain. Residues with higher Shannon Entropy scores exhibit greater sequence variability (high spikes), while residues with lower scores (close to zero) are highly conserved. The mutation sites *C41*, *N42*, *C50*, and *S99* exhibit high entropy, indicating their propensity for evolutionary changes. **(B)** Per-residue disorder propensity of pfDHFR evaluated by six commonly used disorder predictors: PONDR® FIT, PONDR® VSL2, PONDR® VL3, PONDR® VLXT, IUPred Short, and IUPred Long. MDP corresponds to the mean disorder profile calculated by averaging the outputs of individual predictors. Residues with disorder scores ≥ 0.5 are considered disordered, while those between 0.15 and 0.5 are flexible. **(C)** The Mutual Information (MI) plot illustrates strong MI values (darker blue) indicating residues that

123 co-vary. **(D)** 3D structure of the DHFR protein, color-coded by regions of variability and

124 highlighting mutation and active sites. Active site residues are highlighted in yellow sticks, while

125 mutation sites are highlighted in magenta sticks. High-entropy regions in active site are shown in

126 white indicating areas prone to evolutionary change. Highly conserved regions are shown in

127 blue.

Shannon entropy analysis revealed critical positions essential for protein function and mutation-tolerant positions

128
129

We performed Shannon Entropy (SE) calculation for each residue position to identify conserved

and variable regions within the protein sequence which play a crucial role in the proteins

structure and function. Our analysis revealed six conserved residue positions, that is, positions

30, 32, 38, 39, 91 and 104 which exhibited zero entropy values (**Figure 1A**). These regions are

likely to be evolutionarily constrained, as mutations in these positions could disrupt protein

stability or function. Certain active site positions were also found to be relatively conserved

(close to zero) implying that mutations in these positions could have disruptive effects on the

pyrimethamine drug binding. Conversely, regions with higher entropy values were identified

mostly in the flexible loop regions between residue positions 9–25 and 69–89. Other regions

towards the C-terminus (190–219) also displayed high spikes (**Figure 1A**). These positions

displayed greater variability, suggesting they may be more tolerant to mutations. Positions C41,

N42, C50, and S99 also exhibited high variability and have mutations linked to cycloguanil and

pyrimethamine resistance.

Intrinsically disordered prediction revealed that the DHFR is predominantly ordered

We further explored the structural flexibility of the DHFR using six commonly used disorder

predictors[15]: PONDR® FIT[16], PONDR® VSL2[17], PONDR® VL3[17], PONDR® VLXT[18], IUPred

Short[19], and IUPred Long[19] and generated its disorder profile. The outputs of these predictors

were averaged into a mean disorder profile (MDP), providing a consensus view of the protein's

disorder propensity. Intrinsic disorder often correlates with regions capable of tolerating

mutations contributing to evolutionary flexibility.

Our findings revealed that the DHFR protein is predominantly ordered, with most residues
displaying low disorder scores (< 0.5), indicating structural stability. A few flexible regions
(disorder scores between 0.15 and 0.5) were identified at positions 1–24, 60–125, 199–203, and
214–219. However, as per the outputs of PONDR® VSL2[17], which is one of the more accurate
disorder predictors, several key regions of elevated disorder were identified, that is, residues
spanning position 13-21, 61-65, 76-90, 103-107, 123-124, and 217-219. These regions suggest
points of potential mutational tolerance, contributing to the protein's adaptability (**Figure 1B**).

Mutual information calculation revealed tight clusters of inter-dependent residues in the N-
terminus and C-terminus

While SE calculation offered insights into evolutionary conservation rates, it did not account for
interactions between individual residues, particularly how certain positions may evolve in
tandem due to structural or functional dependencies. To explore these interactions, we employed
Mutual Information (MI) analysis. This measures the co-evolution of residue pairs and identifies
compensatory mutations that help maintain protein structure and function. The MI analysis
revealed key co-evolutionary relationships between specific residues, providing critical insights
into the structural and functional constraints imposed by selective pressures. A high MI value
indicates significant positional interdependence, reflecting strong co-evolutionary relationships
between residues. In the MI heatmap (**Figure 1C**), dense clusters, such as those seen between
residues 1–11 and 181–216, highlight regions with notable co-variation.

MI interaction networks highlight co-evolution of mutational sites in DHFR sequence space

Building on the MI calculations, the graph-based analysis was performed to vividly reveal the
complex co-evolutionary relationships between residues in the active site. These MI networks
provide a clear visual representation of how different regions of the protein interact. The residue-
residue networks provided detailed evidence of epistatic interactions, where mutations at one site
could influence compensatory changes at another, preserving enzyme function. The interaction
networks for I5 and C6 (**Figure 2A**) demonstrated a tightly knit structural cluster potentially
critical for drug-binding and resistance mechanisms. They exhibited strong interactions (MI >
0.8) with residues positions C8, C9, K10, V11, K19, and E12. Residues F49 and I103 also

displayed strong and moderate interactions with M95, I103 and K124, S200, F107, and L110

respectively, suggesting their central role in maintaining the active site configuration.

We extended this analysis to the mutational sites which also revealed intriguing co-evolutionary

relationships. Specifically, residues like A7 showed significant interactions with K10, V11 and

E12 while C41 interacted strongly with Y61 and K67. Similarly, N42 and C50 formed

interactions with Y61 and K67.

Figure 2: Mutual Information Networks Reveal Co-evolving Residues Critical for DHFR

Function and Adaptation. (**A**) Interaction network for active site **I5**, showing strong MI values

with residues like I2, C8, and D1. (**B**) This shows a network centered on active site **C6** with

strong interactions extending to K10, V11, and C9, implying that these residues may co-evolve

to maintain catalytic function under selective pressures. (**C**) Interaction network for **F49** showing

significant interactions with residues such as I103 and T53. (**D**) Interaction network around

active site **I103** displayed extensive interactions with several residues such as N115 and K124

highlighting the interconnectedness of residues involved in DHFR's active site dynamics. (**E**)

Interaction network centered on mutation site **A7** showing significant correlations with residues

such as V11, A4, and K10. (**F**) Network around mutation site C41, displaying strong MI

interactions with residues like K55 and C9. (**G**) Mutation site N42 shows strong interactions with

W103 and K67 (**H**) Network focused on mutation site C50 highlights strong interactions with

residues K55 and V11.

Maximal clique analysis and community detection identified densely connected cliques

In our study, we performed Maximal Clique Analysis (MCA)[20,21] and Community Detection[22] to

identify densely connected sub-networks (cliques) and partition the protein into distinct

interaction groups. These analyses provide a deeper layer of insight compared to MI analysis, as

they go beyond pairwise interactions by revealing how clusters of residues interact within a

broader network. Important cliques were identified in regions containing known active and

mutation sites, including residues 6, 7, 42, 49, 155 (**Figure 3A**). These residues form part of

highly interconnected clusters, indicating strong structural and functional interdependencies.

Residues 49 and 155, which form part of pyrimethamine-binding pocket, were observed in

multiple cliques alongside other critical residues highlighting their importance in stabilizing the

surrounding structural framework of the protein. Mutation sites such as residues 7 and 42 also

formed part of distinct cliques where they interacted with other residues. It is worth noting that

residue 6, a drug-binding pocket, was identified within the same clique as residue 7, a known

mutation site. The analysis revealed that the active sites and mutation sites often reside in highly

interconnected cliques, suggesting that mutations at these sites might affect the stability and

function of nearby residues through their cooperative interactions. Community detection (**Figure**

3B) further uncovered broader groupings of residues that frequently interact within the same sub-

network. It highlighted functional domains or regions that evolve together.

Figure 3: **Maximal Clique and Mutational Sensitivity Analyses Reveal Inter-Residue**

Networks and Evolutionarily Constrained Regions in DHFR. (**A**) Maximal clique detection

analysis visualizes cliques or complete subgraphs within the DHFR residue network. Different

cliques represent tightly connected regions of residues which indicate areas of high inter-residue

coordination and co-evolution. (**B**) Community detection analysis using a modularity-based

algorithm identified 11 distinct communities within the DHFR protein. These communities

represent clusters of residues that are likely to have high intra-group connectivity. (**C**) Deep

mutational scan heatmap showing the fitness effects of single amino acid substitutions across the

DHFR protein. The heatmap reveals positions that are highly sensitive to mutations, with dark

blue indicating deleterious effects (negative fitness scores) and light shades or orange

representing beneficial or neutral mutations.

Deep mutational scanning yielded mutational hotspots with beneficial fitness effects that may

facilitate evolutionary adaptation under drug pressure.

The Deep Mutational Scanning (DMS) analysis was performed to gain insights into the

functional consequences of amino acid substitutions across the protein sequence. By quantifying

the fitness effects of each mutation, this analysis allowed us to map the sequence space of

DHFR, identifying regions that are intolerant to mutations, as well as those that can tolerate or

even benefit from substitutions. A significant number of deleterious mutations were observed in

key regions of the DHFR enzyme (**Figure 3C**), particularly around the active site. Positions such

as I155, which is a critical residue within the active site, show dramatic reductions in fitness for

nearly all substitutions. This suggests that I155 plays an essential role in maintaining the

enzyme's catalytic function and structural stability. Similarly, D45, F49 and I155, another set of

residues located in the active site, also shows a strong deleterious signal upon mutation with

almost all substitutions. The positions G96, G156, G157 and Y217 strongly emerged as key sites

where mutations lead to significant fitness losses. These residues are situated near the catalytic

core of the enzyme and substitutions here likely disrupt the enzyme's stability or interfere with

essential catalytic processes. The DMS analysis also uncovered key mutation hotspots with

beneficial fitness effects, particularly in the regions spanning residues 120 to 155 and 190 to 211.

The fitness trajectories for active site residues illustrates distinct adaptive paths for various

mutations, highlighting the complexity of the fitness landscape for pfDHFR

Our fitness trajectory analysis provides critical insights into the distinct adaptive pathways of key

active site residues in *pf*DHFR. This reveals how the enzyme navigates complex fitness

landscapes under selective pressure. To simulate these evolutionary paths, we developed an

epistasis-based fitness prediction model using Dijkstra's shortest path algorithm. In this model,

mutations are represented as nodes, and the edges between them are weighted by the absolute

fitness differences between the mutations. This approach identifies the most efficient mutational

paths—those requiring the fewest steps—from the wild-type to both local and global fitness
peaks. The model captures both stochastic and deterministic evolutionary dynamics, offering a
realistic framework for simulating biological evolution. Stochastic sampling introduces
randomness, reflecting genetic drift and the exploration of multiple paths. Deterministically, the
algorithm favors mutations with immediate fitness benefits, emulating natural selection. This
dual approach balances short-term incremental adaptation with long-distance evolutionary leaps**,**
mirroring the unpredictable yet selective nature of real-world evolution.

Using deep mutational scan data to guide these pathways, the model reveals how near-neutral
and compensatory mutations expand the genotype-phenotype space. This allows the enzyme to
smooth rugged fitness landscapes and adapt under selective pressure. For instance, mutations at
position 49 exhibited quick transitions to fitness peaks such as A51S, highlighting the mutational
tolerance at this site. However, other substitutions at the same site, such as F49K and F49W,
required longer pathways to reach comparable peaks, indicating variability in their fitness effects
(**Figure 4A**). Similarly, mutations S99C and S99T demonstrated shorter evolutionary paths to
local fitness peaks like C50S, suggesting faster adaptation, while mutations like S99F followed
more extended trajectories (**Figure 4C**). This trend was also observed at positions 102 and 103,
with certain mutations achieving rapid adaptation, while others required more gradual, complex
walks. Conversely, position 155 presented one of the most challenging landscapes, with most
mutations demanding over 100 evolutionary steps to achieve fitness peaks (**Figure 4F**). A
comparable trend was seen at position 161 (**Figure 4H**), where mutations such as Y161L and
Y161K required long evolutionary walks toward global peaks, highlighting the ruggedness of the
landscape at these critical sites.

These findings underscore that evolutionary adaptation in DHFR is not uniform. While certain
residues, such as 49 and 99, serve as stepping stones toward resistance by facilitating quick
fitness gains, others like 155 demand more extensive evolutionary navigation. The interplay
between stochastic exploration and deterministic selection captured by our model reflects the
intricate dynamics of DHFR evolution, offering a deeper understanding of how the enzyme
balances adaptability and functional constraints under selective pressure.

Figure 4: **Fitness Trajectories Highlight Adaptive Paths of Key Active Site Residues in**

DHFR. (A) At position 49, mutations such as F49L and F49Y achieve quicker routes to fitness

peaks, while others like F49K, F49W, and F49H follow longer, more gradual trajectories. **(B)**

Position 45 shows gradual fitness increases across all mutations, except D45E which achieved a

relatively quicker fitness. **(C)** Mutations at position 99 exhibit variability in trajectory length.

Positions S99T and S99C reach fitness peaks faster, while S99G requires the longest path. **(D)**

Position 102 shows variation in mutation paths, with S102A and S102P achieving fitness

improvements more quickly than S102E and S102F. **(E)** Mutations at positions 103 generally

follow longer trajectories to fitness peaks. **(F)** At position 155, most mutations require extended

walks of over 100 steps to reach fitness peaks. **(G)** Shows similar patterns at position 161, where

Y161L and Y161K mutations follow longer trajectories. **(H)** Position 176 demonstrates both

short and long trajectories, with some mutations reaching fitness peaks more rapidly, while

others require extended adaptive steps.

Discussion

The folate pathway plays a crucial role in the survival of *P. falciparum* by supplying essential

cofactors for nucleotide and amino acid synthesis[7]. This makes DHFR a prime target for

antifolate drugs. However, the emergence of resistance mutations in this essential protein has led

to significant challenges in the effective treatment of malaria. These mutations have enabled the

parasite to evade drug inhibition while preserving enzyme functionality[23]. Further complexity

arises from the genomic organization of the *pfdhfr-ts* gene. This is a bifunctional gene located on

chromosome 4 that encodes both DHFR and thymidylate synthase within the same polypeptide

chain[24,25]. This dual functionality introduces potential pleiotropic effects, where mutations

selected for resistance in the DHFR domain could unintentionally affect the function of the TS

domain, or vice versa. Similar to the observations of Raff and Kaufman[26], such

interdependencies highlight the complexity of interconnected biological systems, where targeted

changes can yield large-scale, unintended consequences across multiple functions. The

interdependence between these two enzymatic activities creates challenges in predicting the

evolutionary consequences of mutations, as adaptations beneficial for one activity might impair

the other. While the direct impact of promoter mutations on this gene remains largely

unexplored[27], these mutations could potentially influence gene dosage and expression levels,

thereby affecting drug resistance mechanisms. This genomic context implies that resistance

arises not only from individual enzyme mutations but also from the complex regulation of gene

expression. Thus, understanding the evolutionary dynamics driving these mutational escapes in

DHFR is essential for designing novel antifolate therapies.

Mutational landscapes are inherently complex[28], reflecting the diverse roles amino acid

substitutions play in shaping protein function and evolution. Proteins, including DHFR, are

subjected to a range of evolutionary pressures that necessitate both structural stability and

adaptability[29,30]. However, mutations that confer drug resistance often introduce trade-offs that

impact their fitness[31]. This balance is complicated by the fact that near-neutral mutations, while

individually minor, can act as stepping stones toward significant adaptive shifts[32]. The study of

mutational landscapes using tools such as deep mutational scanning has proven critical in

identifying mutations that influence protein adaptability without drastic functional

compromises[33,34]. Some mutations improve fitness only marginally but, over time, accumulate to

confer drug resistance[35]. These insights highlight how proteins like DHFR navigate a rugged

landscape, with regions of both mutational tolerance and structural constraint. This

understanding is essential for predicting the evolution of resistance and identifying critical points

for therapeutic intervention.

The evolutionary landscape of proteins is defined by peaks of high fitness separated by valleys of

low fitness. This ruggedness reflects the complex interplay between mutations, structural

integrity, and functional performance. Adaptive walks along this landscape often involve both

stochastic exploration and deterministic processes that select for advantageous traits[36]. Near-

neutral mutations facilitate exploration by expanding the search space which allows enzymes to

navigate fitness valleys more efficiently[32]. Over time, these mutations can accumulate to provide

a selective advantage, especially under environmental pressures. Evolutionary trajectories are

further shaped by epistasis, where interactions between multiple mutations influence fitness

outcomes[37]. These complexities emphasize the importance of studying not just individual

mutations but also their collective impact on protein function.

The present study provides critical insights into the mutational and evolutionary landscapes of *P.*
falciparum DHFR with emphasis on both structural constraints and adaptive strategies employed
by the protein. Conserved residues were identified and highlighted as structurally indispensable
elements that are less tolerant to mutations. These regions offer promising targets for novel
antifolate therapies, as disrupting them could impair the parasite's ability to maintain
functionality and evolve resistance. Network-based analyses further revealed key hubs and
cliques of interacting residues. The presence of such interconnected, co-evolving clusters
highlights that resistance arises not through isolated mutations but via cooperative networks that
preserve stability despite structural alterations. Several mutation hotspots were also identified
with beneficial fitness effects, particularly in regions spanning residues 120–155 and 190–211.
While many mutations, especially at key active site residues like I155, D45, and F49, caused
significant fitness losses due to their critical role in protein stability, certain regions exhibited
greater tolerance to substitutions. These mutation-tolerant regions suggest potential adaptive
flexibility which enable the enzyme to navigate selective pressures by accommodating beneficial
changes without compromising overall functionality. The integration of conserved elements,
mutation hotspots, co-evolving clusters, and interconnected networks underscores the
multifaceted nature of resistance. Targeting these networks and conserved regions provides a
promising approach for the development of future-proof antifolate therapies, disrupting the
parasite's ability to adapt and evolve in response to drug pressure.

To simulate adaptive evolutionary pathways and explore how mutations influence fitness
landscapes, we developed an epistatic fitness prediction model. Our model provides valuable
insights into how *P. falciparum* DHFR navigates complex fitness landscapes under selective
pressure. This dual framework captures the unpredictable aspects of evolution while also
reflecting selective pressures that favor beneficial mutations. Through this approach, we modeled
both rapid adaptive walks and gradual evolutionary trajectories, mirroring the diverse paths
proteins take toward higher fitness in dynamic environments. The use of deep mutational scan
data enabled us to assign fitness values to individual mutations. This provided a quantitative lens
to evaluate how various mutations affect enzyme stability and function. By mapping these
trajectories, we demonstrated that near-neutral mutations and compensatory interactions play
pivotal roles in expanding the genotype-phenotype space, thereby smoothing the rugged fitness

landscape. This adaptive flexibility allows DHFR to navigate resistance pathways efficiently,
overcoming mutational barriers while retaining functionality.

The insights from this model emphasize the importance of epistatic interactions—where
mutations at one site influence the fitness outcomes of others—in shaping evolutionary
pathways. It underscores that resistance in DHFR is not merely a product of isolated mutations
but rather emerges from complex mutational networks that facilitate both short-term gains and
long-term adaptability. This predictive framework enhances our understanding of antifolate
resistance mechanisms and lays the groundwork for designing novel antifolate therapies that can
effectively disrupt these evolutionary strategies.

While our study offers significant insights into the evolution of antifolate resistance, it is not
without limitations. This research was solely based on protein sequence analysis and did not
delve into the structural dynamics of DHFR, which could provide additional insights into
enzyme stability, conformational changes, and protein-drug interactions. Despite incorporating
stochastic sampling to simulate randomness in evolutionary choices, the model may still miss
certain complexities of real-world adaptation, such as temporary fitness losses or parallel
exploration of multiple paths. These limitations highlight the challenges of representing rugged
fitness landscapes, where local optima may be revisited and evolutionary trajectories can follow
non-linear patterns over time. Moreover, the limited availability of fitness data constrains the
scope of the fitness landscape explored. Expanding the dataset or integrating complementary
experimental data would provide a more comprehensive view of mutational effects across a
broader range of residues. Future experimental validation of our computational findings is
crucial. Molecular docking simulations and fitness assays in biological systems will confirm
whether the identified adaptive pathways correspond to real-world resistance mechanisms.
Integrating advanced computational tools, such as machine learning models, could enhance
predictive accuracy and uncover novel mutational pathways that are difficult to identify using
conventional algorithms. These efforts will contribute to the development of future-proof
antifolate therapies capable of addressing the challenges posed by evolving resistance in *P.*
falciparum.

Materials and Methods

Sequences preprocessing

For this study, we used the crystal structure of the wild-type *P. falciparum* DHFR-TS protein

(PDB ID: 3QGT) as the reference for our analyses. The protein structure consists of two

domains: the DHFR domain (first 231 residues) and the TS domain (last 288 residues), separated

by an 89-residue interjunction region (Supplemental Figure 1). Given the focus on DHFR, we

truncated the DHFR domain, which stretches from amino acid position 10 to 228 in the full-

length protein as annotated in UNIPROT[38]. This truncation was performed using the PYMOL

Molecular Graphics System (version 3.0.3)[39]. For consistency, the truncated sequence was re-

indexed from 1 to 219, a scheme used throughout the study.

Homology Modeling and Structure Validation

Upon examination, the DHFR protein contained ten missing residues in its sequence from

position 86 to 95. To fill in these gaps, homology modeling was performed using Modeller

(Version 10.5)[40,41]. This software generated several structural models and ranked them using the

Discrete Optimized Protein Energy (DOPE) scoring function. The model with the lowest DOPE

score was selected as the most accurate representation of the full DHFR structure (**Supplemental**

Figure 2). To ensure the accuracy of the modeled structure, we calculated the Root Mean Square

Deviation (RMSD) between the modeled residues and the crystal structure (**Supplemental**

Figure 3). An RMSD value of 0.306 Å indicated high structural similarity. After validation, we

extracted the amino acid sequence (219 residues) from the modeled DHFR structure for further

analyses.

Sequence Space Analysis

To explore the mutational landscape of DHFR, we performed a sequence space analysis using

the multiple sequence alignment (MSA) of homologous DHFR proteins. First, we retrieved 5000

non-redundant protein sequences homologous to DHFR using the NCBI BlastP tool[42]. The

sequences were cleaned to ensure high quality, non-redundancy and biological relevance. We

then aligned the cleaned sequences using the ClustalW algorithm[43], and the resulting MSA

served as the basis for the subsequent analyses.

Occupancy Calculation

Occupancy calculations were performed to assess how frequently specific amino acids occupy
each residue position in homologous DHFR sequences. Positions with high occupancy values
represent conserved residues, likely essential for the protein's structural stability. By comparing
occupancy data with mutation sites, we identified residues where mutations may have significant
impacts on the protein's structure and function.

Shannon Entropy Calculation

Shannon entropy $(S(i))$[44] was calculated to quantify the variability of amino acids at each
sequence position. Positions with low entropy values represent conserved residues, critical for
structural and functional integrity, while high entropy values indicate positions prone to
mutations.

$$S(i) = -\sum_{a_i=1}^{20} P(a_i) \, log P(a_i)$$

Where *i* represents the sequence position, and *P(a_i)* is the probability of amino acid *a* occurring
at position *i* in the MSA. Lower values of *S(i)* correspond to highly conserved residues, while
higher values suggest increased variability.

Mutual Information Calculation

To investigate co-evolving residues, we calculated mutual information (MI)[45,46] for each pair of
positions in the MSA. MI quantifies the degree of correlation between two sequence positions,
revealing residues that might be functionally or structurally interdependent. These co-varying
residues could indicate regions where mutational compensation occurs, maintaining overall
protein function despite mutations in specific positions.

$$MI(i,j) = \sum_{a_i=1}^{21} \sum_{b_j=1}^{21} P(a_i, b_j) \, log \frac{P(a_i, b_j)}{P(a_i) \, P(b_j)}$$

In this formula, *P(ai, bj)* represents the probability of amino acids a and b occurring at positions *i*

and *j*, respectively, while $P(a_i)$ and $P(b_j)$ are the individual probabilities at each position . The

MI(i,j) values range from 0 (uncorrelated residues) to MI_{max} (the most interdependent residue

pairs). In this study, a cutoff MI value of 0.5 was chosen to focus on highly co-varying positions.

The MI values were visualized as a heatmap and a co-evolutionary matrix, generated using

python programming.

Intrinsic Disorder Prediction

Intrinsic disorder prediction analysis was conducted to explore the presence of disordered

regions within the DHFR protein. Disordered regions, known for their inherent flexibility, can

tolerate higher levels of mutation and often play significant roles in protein-protein interactions.

For a comprehensive evaluation, we employed six widely used disorder predictors: PONDR®

FIT[16], PONDR® VSL2[17], PONDR® VL3[17], PONDR® VLXT[18], IUPred Short[19], and IUPred

Long[19]. These predictors leverage distinct predictive approaches to identify disordered regions

within the protein. PONDR® VSL2 is particularly optimized for proteins containing both

structured and disordered regions[47]. By integrating the outputs from all six predictors, we

generated a mean disorder profile (MDP), providing a consensus disorder score for each residue.

Residues with scores above 0.5 were classified as disordered, while those between 0.15 and 0.5

were considered flexible but ordered. This disorder profile was further analyzed to examine

potential correlations between disordered regions, known mutation sites, and active site residues,

helping to elucidate the role of structural flexibility in resistance mutations and the maintenance

of protein function.

Graph-Based Analysis

Following the MI calculations, we performed a graph-based analysis of the identified co-

evolving residues. Nodes in the graph represent individual residues, while edges denote strong

mutual information-based connections between them. The MI threshold was used to define the

strength of these interactions, with different ranges highlighting strong (MI > 0.8), moderate (0.6

< MI < 0.8), and weak (0.4 < MI < 0.6) interactions. This graph-based analysis allowed us to

visualize key mutation sites (A16, N42, C50, S99, S108, and I155) and their network of

interactions, especially with active site residues. The resulting network graphs provided insight

into how mutations at specific residues might influence structural stability and drug resistance

through interdependencies with nearby residues.

To investigate densely connected sub-networks within the DHFR protein, we employed Maximal

Clique Analysis (MCA)[20,21] and Community Detection[22] methods. In MCA, we aimed to identify

groups of residues that form maximal cliques, where every residue (node) within a clique is

directly connected to every other residue. This method focuses on identifying highly

interconnected sets of residues that are structurally and functionally interdependent. Each residue

in the protein is represented as a node and the edges between these nodes represent significant

interactions, which are based on proximity in the protein structure or known co-evolutionary

relationships. The primary goal was to identify the largest cliques within the overall interaction

network. Community detection was also performed to divide the protein's residue interaction

network into distinct subgroups or communities. Unlike MCA, which focuses on fully connected

sub-networks, community detection identifies groups of residues that are more likely to interact

within their community than with residues outside it.

Fitness Trajectories Analysis Using Epistatic Prediction Model

To explore the epistatic interactions between multiple mutations, we developed an epistasis-

based fitness prediction model to simulate adaptive walks within the *P. falciparum* DHFR

protein. The model integrates both stochastic and deterministic elements to account for the

complexity of the fitness landscape. The epistatic data used were derived from deep mutation

scan which predict the fitness effects of individual mutations. The goal of this analysis was to

map the evolutionary trajectories of key active site residues as they navigate through the fitness

landscape.

We used Dijkstra's shortest path algorithm[48] to construct the adaptive walks, representing the

protein's potential mutational paths in search of fitness improvement. In this graph-based

approach, mutations were treated as nodes, and fitness differences between mutations as

weighted edges. The shortest path corresponds to the fewest evolutionary steps required to

maximize fitness. The fitness values for each mutation were calculated from the epistatic

prediction data, where a positive value indicates a beneficial mutation and a negative value

represents a deleterious one. Given that a set of mutations $M = \{m_1, m_2,\ldots, m_n\}$ and their fitness

values $f(m_i)$, our goal was to identify the shortest adaptive walk that maximizes fitness. The

distance between two mutations is defined as the absolute difference between their fitness

values;

$$d(m_i, m_j) = |f(m_i) - f(m_j)|$$

511   We sought to find the mutation path P from an initial wild-type m_{wt} to a final high-fitness peak

512   mutation m_{peak}, such that the sum of fitness differences along the path is minimized;

$$P = \arg\min \sum_{i=1}^{k-1} d(m_i, m_{i+1})$$

A combination of deterministic and stochastic methods was employed. The algorithm

deterministically follows the mutation path with the highest fitness improvement, i.e., aiming

directly for the global peak unless the local peak provides a substantially higher immediate

fitness gain. When multiple peaks are available, the algorithm stochastically samples between the

local and global peaks based on fitness differences. The stochastic component is controlled by a

probability threshold, which adjusts based on the fitness difference between the local peak and

the global peak:

$$\text{probability threshold} = \begin{cases} 0.3 & \text{if } \Delta f > 2 \\ 0.5 & \text{if } 1 < \Delta f \leq 2 \\ 0.7 & \text{if } \Delta f \leq 1 \end{cases}$$

522   This threshold governs the likelihood of choosing a local versus a global peak, simulating a

523   situation where evolutionary pressures might favor local exploration or long-distance leaps to

524   distant fitness peaks.

AUTHOR CONTRIBUTIONS

SC and DS conceived and planned the work. SC motivated and guided the work. DM performed all the formal analysis, model development, manuscript composition, and critical review. VNU performed the disorder analysis. SC, DS, VNU, SA, ASC, DZ, RJ and PDU provided critical feedback and assisted in refining the manuscript.

ACKNOWLEDGMENT

DM acknowledges One Health Trust, Bangalore, India. DM and SC acknowledge the Director, Birla Institute of Technology and Science-Pilani, Hyderabad, India.

DECLARATION

The authors declare that they have no conflicts of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

1.   Ahmad T, Alhammadi BA, Almaazmi SY, Arafa S, Blatch GL, Dutta T, et al. Plasmodium falciparum heat shock proteins as antimalarial drug targets: An update. Cell Stress and Chaperones. 2024 Apr 1;29(2):326–37.

2.   Huthmacher C, Hoppe A, Bulik S, Holzhütter HG. Antimalarial drug targets in Plasmodium falciparum predicted by stage-specific metabolic network analysis. BMC Systems Biology. 2010 Aug 31;4(1):120.

3.   Sirawaraporn W, Prapunwattana P, Sirawaraporn R, Yuthavong Y, Santi DV. The dihydrofolate reductase domain of Plasmodium falciparum thymidylate synthase-dihydrofolate reductase. Gene synthesis, expression, and anti-folate-resistant mutants. Journal of Biological Chemistry. 1993 Oct 15;268(29):21637–44.

4. Kreutzfeld O, Tumwebaze PK, Byaruhanga O, Katairo T, Okitwi M, Orena S, et al. Decreased Susceptibility to Dihydrofolate Reductase Inhibitors Associated With Genetic Polymorphisms in Ugandan Plasmodium falciparum Isolates. J Infect Dis. 2021 Aug 30;225(4):696–704.

5. Wróbel A, Arciszewska K, Maliszewski D, Drozdowska D. Trimethoprim and other nonclassical antifolates an excellent template for searching modifications of dihydrofolate reductase enzyme inhibitors. J Antibiot. 2020 Jan;73(1):5–27.

6. Fernández-Villa D, Aguilar MR, Rojo L. Folic Acid Antagonists: Antimicrobial and Immunomodulating Mechanisms and Applications. Int J Mol Sci. 2019 Oct 9;20(20):4996.

7. Hyde JE. Exploring the folate pathway in Plasmodium falciparum. Acta Trop. 2005 Jun;94(3):191–206.

8. Bilsland E, Van Vliet L, Williams K, Feltham J, Carrasco MP, Fotoran WL, et al. Plasmodium dihydrofolate reductase is a second enzyme target for the antimalarial action of triclosan. Sci Rep. 2018 Jan 18;8(1):1038.

9. Nzila A, Ward SA, Marsh K, Sims PF, Hyde JE. Comparative folate metabolism in humans and malaria parasites (part I): pointers for malaria treatment from cancer chemotherapy. Trends in parasitology. 2005 Jun;21(6):292.

10. Chusacultanachai S, Thiensathit P, Tarnchompoo B, Sirawaraporn W, Yuthavong Y. Novel antifolate resistant mutations of *Plasmodium falciparum* dihydrofolate reductase selected in *Escherichia coli*. Molecular and Biochemical Parasitology. 2002 Mar 1;120(1):61–72.

11. Yuthavong Y, Tarnchompoo B, Vilaivan T, Chitnumsub P, Kamchonwongpaisan S, Charman SA, et al. Malarial dihydrofolate reductase as a paradigm for drug development against a resistance-compromised target. Proc Natl Acad Sci U S A. 2012 Oct 16;109(42):16823–8.

12. Kayode AT, Ajogbasile FV, Akano K, Uwanibe JN, Oluniyi PE, Eromon PJ, et al. Polymorphisms in Plasmodium falciparum dihydropteroate synthetase and dihydrofolate reductase genes in Nigerian children with uncomplicated malaria using high-resolution melting technique. Sci Rep. 2021 Jan 12;11(1):1–8.

13. Amusengeri A, Tata RB, Tastan Bishop Ö. Understanding the Pyrimethamine Drug Resistance Mechanism via Combined Molecular Dynamics and Dynamic Residue Network Analysis. Molecules. 2020 Feb 18;25(4):904.

14. Sanyal D, Banerjee S, Bej A, Chowdhury VR, Uversky VN, Chowdhury S, et al. An integrated understanding of the evolutionary and structural features of the SARS-CoV-2 spike receptor binding domain (RBD). International Journal of Biological Macromolecules. 2022 Sep;217:492–505.

15. Dayhoff II GW, Uversky VN. Rapid prediction and analysis of protein intrinsic disorder. Protein Science. 2022;31(12):e4496.

16. Xue B, Dunbrack RL, Williams RW, Dunker AK, Uversky VN. PONDR-FIT: A Meta-Predictor of Intrinsically Disordered Amino Acids. Biochimica et biophysica acta. 2010 Jan 25;1804(4):996.

17. Peng K, Radivojac P, Vucetic S, Dunker AK, Obradovic Z. Length-dependent prediction of protein intrinsic disorder. BMC Bioinformatics. 2006 Apr 17;7(1):208.

18. Romero P, Obradovic Z, Li X, Garner EC, Brown CJ, Dunker AK. Sequence complexity of disordered protein. Proteins. 2001 Jan 1;42(1):38–48.

19. Dosztányi Z, Csizmók V, Tompa P, Simon I. The pairwise energy content estimated from amino acid composition discriminates between folded and intrinsically unstructured proteins. J Mol Biol. 2005 Apr 8;347(4):827–39.

20. Pardalos PM, Rodgers GP. A branch and bound algorithm for the maximum clique problem. Computers & Operations Research. 1992 Jul 1;19(5):363–75.

21. Konc J, Janezic D. An improved branch and bound algorithm for the maximum clique problem. MATCH Commun Math Comput Chem. 2007;58, 569-590.

22. Grant WP, Ahnert SE. Modular decomposition of protein structure using community detection. Journal of Complex Networks. 2019 Feb 1;7(1):101–13.

23. Heinberg A, Kirkman L. The molecular basis of antifolate resistance in Plasmodium falciparum: looking beyond point mutations. Ann N Y Acad Sci. 2015 Apr;1342(1):10–8.

24. Dj B, Wb L, T H, J I. Molecular cloning and sequence analysis of the Plasmodium falciparum dihydrofolate reductase-thymidylate synthase gene. Proceedings of the National Academy of Sciences of the United States of America [Internet]. 1987 Dec [cited 2024 Oct 7];84(23). Available from: https://pubmed.ncbi.nlm.nih.gov/2825189/

25. Mita T, Tanabe K, Kita K. Spread and evolution of Plasmodium falciparum drug resistance. Parasitology International. 2009 Sep;58(3):201–9.

26. Garfield DA, Wray GA. The Evolution of Gene Regulatory Interactions. BioScience. 2010 Jan 1;60(1):15–23.

27. Ferreira de Azevedo M, del Portillo HA. Control of Gene Expression in Plasmodium. 2006 Jul 17 [Updated 2007 May 11]. In: Gruber A, Durham AM, Huynh C, et al., editors. Bioinformatics in Tropical Disease Research: A Practical and Case-Study Approach [Internet]. Bethesda (MD): National Center for Biotechnology Information (US); 2008. Chapter B01.

28. Wagner A. Evolvability-enhancing mutations in the fitness landscapes of an RNA and a protein. Nat Commun. 2023 Jun 19;14(1):3624.

29. Siltberg-Liberles J, Grahnen JA, Liberles DA. The Evolution of Protein Structures and Structural Ensembles Under Functional Constraint. Genes. 2011 Oct 28;2(4):748.

626   30.   Halabi N, Rivoire O, Leibler S, Ranganathan R. Protein Sectors: Evolutionary Units of
627   Three-Dimensional Structure. Cell. 2009 Aug 21;138(4):774–86.

628   31.   Matange N, Bodkhe S, Patel M, Shah P. Trade-offs with stability modulate innate and
629   mutationally acquired drug resistance in bacterial dihydrofolate reductase enzymes.
630   Biochemical Journal. 2018 Jun 29;475(12):2107–25.

631   32.   Rendel MD. Adaptive evolutionary walks require neutral intermediates in RNA fitness
632   landscapes. Theoretical Population Biology. 2011 Feb 1;79(1):12–8.

633   33.   Maes S, Deploey N, Peelman F, Eyckerman S. Deep mutational scanning of proteins in
634   mammalian cells. Cell Reports Methods. 2023 Nov 20;3(11):100641.

635   34.   Dunham AS, Beltrao P. Exploring amino acid functions in a deep mutational landscape.
636   Molecular Systems Biology. 2021 Jul 22;17(7):e10305.

637   35.   Saralamba N, Nakeesathit S, Mayxay M, Newton PN, Osorio L, Kim JR, et al. Geographic
638   distribution of amino acid mutations in DHFR and DHPS in Plasmodium vivax isolates
639   from Lao PDR, India and Colombia. Malaria Journal. 2016 Sep 21;15(1):484.

640   36.   Park SC, Neidhart J, Krug J. Greedy adaptive walks on a correlated fitness landscape.
641   Journal of Theoretical Biology. 2016 May 21;397:89–102.

642   37.   Starr TN, Thornton JW. Epistasis in protein evolution. Protein Science : A Publication of
643   the Protein Society. 2016 Feb 28;25(7):1204.

644   38.   The UniProt Consortium. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic
645   Acids Research. 2023 Jan 6;51(D1):D523–31.

646   39.   Yuan S, Chan HCS, Hu Z. Using PyMOL as a platform for computational drug design.
647   WIREs Computational Molecular Science. 2017;7(2):e1298.

648   40.   Fiser A, Do RK, Sali A. Modeling of loops in protein structures. Protein Sci. 2000
649   Sep;9(9):1753–73.

650   41.   Eswar N, Webb B, Marti-Renom MA, Madhusudhan MS, Eramian D, Shen MY, et al.
651   Comparative protein structure modeling using Modeller. Curr Protoc Bioinformatics. 2006
652   Oct;Chapter 5:Unit-5.6.

653   42.   McGinnis S, Madden TL. BLAST: at the core of a powerful and diverse set of sequence
654   analysis tools. Nucleic Acids Research. 2004 Jul 1;32(Web Server issue):W20.

655   43.   Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of
656   progressive multiple sequence alignment through sequence weighting, position-specific gap
657   penalties and weight matrix choice. Nucleic Acids Res. 1994 Nov 11;22(22):4673–80.

658   44.   Strait BJ, Dewey TG. The Shannon information entropy of protein sequences. Biophysical
659   Journal. 1996 Jul 1;71(1):148–55.

45. Gloor GB, Martin LC, Wahl LM, Dunn SD. Mutual Information in Protein Multiple
Sequence Alignments Reveals Two Classes of Coevolving Positions. Biochemistry. 2005
May 1;44(19):7156–65.

46. Simonetti FL, Teppa E, Chernomoretz A, Nielsen M, Marino Buslje C. MISTIC: mutual
information server to infer coevolution. Nucleic Acids Research. 2013 Jul 1;41(W1):W8–
14.

47. Johnson DE, Xue B, Sickmeier MD, Meng J, Cortese MS, Oldfield CJ, et al. High-
Throughput Characterization of Intrinsic Disorder in Proteins from the Protein Structure
Initiative. Journal of structural biology. 2012 May 29;180(1):201.

48. Cui Q, Purisima EO, Wang E. Protein evolution on a human signaling network. BMC Syst
Biol. 2009 Feb 18;3(1):21.

Supplemental Information

Navigating the Fitness Landscapes of Plasmodium falciparum Dihydrofolate Reductase: Evolutionary Insights into Antifolate Resistance

Danny Muzata[1,2#], Dwipanjan Sanyal[3#], Shivram A[1,2], Deeptanshu Pandey[4], Soumyananda Chakraborti[4], Aneesh S. Chivukula[2], Ravi Jasuja[5], Vladimir N. Uversky[6], Daniel Zielinski[7], Prajna D. Upadhyay[2], Mark Albers[8], and Sourav Chowdhury[5*]

Affiliations:
[1] One Health Trust, Bangalore, India
[2] Department of Computer Science and Information Systems, Birla Institute of Technology and Science-Pilani, Hyderabad, India.
[3] Structural Biology and Bioinformatics Division, CSIR-Indian Institute of Chemical Biology, Kolkata, India
[4] Department of Biological Sciences, Birla Institute of Technology and Science-Pilani, Hyderabad, India.
[5] Research Program in Men's Health: Aging and Metabolism, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA
[6] USF Health Byrd Alzheimer's Research Institute, Morsani College of Medicine, University of South Florida, Tampa, FL 33612, USA
[7] Department of Bioengineering, University of California, San Diego, La Jolla, CA, USA
[8] Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts, USA

*Corresponding Author:

Sourav Chowdhury
Department of Biological Sciences,
Office: J Block-109,
Birla Institute of Technology and Science, Hyderabad Campus,
Hyderabad, Telangana, India, 500078
Email: sourav.chowdhury@hyderabad.bits-pilani.ac.in

#*These authors contributed equally to the work.*

728   **Supplemental Figures**

731   **Supplemental Figure S1:** This illustrates the 608 long *pf*DHFR-TS (PDB ID: 3QGT) dimeric

732   assembly. In cyan are the DHFR domains with DHFR-TS junction in magenta and yellow region

733   as TS domains. The DHFR domains are complexed with the drug pyrimethamine and NADPH

734   cofactor

Supplemental Figure S2: This plot presents the validation results of homology modeling performed using Modeller. The DOPE scores for the model and template structures are compared across the alignment positions. The plot shows how well the model (red) aligns with the template (green). However, the misalignment observed between positions 0 to 96 is expected as it is due to the existing gap in the template structure.

745 **Supplemental Figure S3:** Shows the structural alignment of the modeled DHFR domain (cyan)

746 with the template structure (green) using PyMOL. The RMSD was calculated to assess the

747 structural similarity between the model and the template. The RMSD value of 0.306 Å indicates

748 a good alignment, validating the accuracy of the homology model. This alignment helps confirm

749 that the modeled structure retains the key features and overall topology of the template.