Analysis of sparse animal social networks

Helen K Mylne[1,2*], Jackie Abell[3], Colin M Beale[1,2,4], Lauren JN Brent[5], Jakob Bro-Jørgensen[6], Kate E Evans[7,8], Jordan DA Hart[5], Dabwiso Sakala[9,10], Twakundine Simpamba[11], David Youldon[10], and Daniel W Franks[1‡].

1 Department of Biology, University of York, York, UK

2 Leverhulme Centre for Anthropocene Biodiversity, University of York, York, UK

3 Centre for Agroecology, Water and Resilience, Coventry University, Coventry, UK

4 York Environmental Sustainability Institute, University of York, York, UK

5 Centre for Research in Animal Behaviour, University of Exeter, Exeter, UK

6 Mammalian Behaviour and Evolution Group, Department of Evolution, Ecology and Behaviour, University of Liverpool, Liverpool, UK

7 Elephants for Africa, London, UK

8 Gothenburg Global Biodiversity Centre, Gothenburg, Sweden

9 Department of Geography, University of Zurich, Zürich, Switzerland

10 African Lion and Environmental Research Trust, Livingstone, Zambia

11 Department of National Parks & Wildlife, Zambia

‡Senior author

* helen.mylne@york.ac.uk

Abstract

Low-density social networks can be common in animal societies, even among species generally considered to be highly social. Social network analysis is commonly used to analyse animal societal structure, but edge weight (strength of association between two individuals) estimation methods designed for dense networks can produce biased measures when applied to low-density networks. Frequentist methods suffer when data availability is low, because they contain an inherent flat prior that will accept any

possible edge weight value, and contain no uncertainty in their output. Bayesian methods can accept alternative priors, so can provide more reliable edge weights that include a measure of uncertainty, but they can only reduce bias when sensible prior values are selected. Currently, neither accounts for zero-inflation, so they produce edge weight estimates biased towards stronger associations than the true social network, which can be seen through diagnostic plots of data quality against output estimate. We address this by adding zero-inflation to the model, and demonstrate the process using group-based data from a population of male African savannah elephants. We show that the Bayesian approach performs better than the frequentist to reduce the bias caused by these problems, though the Bayesian requires careful consideration of the priors. We recommend the use of a Bayesian framework, but with a conditional prior that allows the modelling of zero-inflation. This reflects the fact that edge weight derivation is a two-step process: i) probability of ever interacting, and ii) frequency of interaction for those who do. Additional conditional priors could be added where the biology requires it, for example in a society with strong community structure, such as female elephants in which kin structure would create additional levels of social clustering. Although this approach was inspired by reducing bias observed in sparse networks, it could have value for networks of all densities.

1 Introduction

Social networks underlie all animal social behaviour, and analysing the factors that affect their structure can improve our understanding of how individuals interact. Social network density forms an important part of social structure, defined by the proportion of dyads (all of the possible pairs of individuals) who could potentially associate within a population that do associate. Some animal social networks are dense, characterised by frequent interactions and connections among a high proportion of its members. Others are sparse, with a high proportion of dyads never associating with one another. This difference in network density can influence an individual's rate of social learning [1–3] or their risk of pathogenic infection [4–7], which can cascade to population-level impacts. Differences in network density may be due to inherent differences in gregariousness between species, or inter-population variation in resource availability or predation risk

imposing different limits on the propensity of different populations to aggregate. ₁₃

Studies of animal social networks tend to address gregarious species, typically ₁₄
characterised by dense social networks. However, being gregarious and living in groups ₁₅
is not the same as having strong social preferences: two species can be equally ₁₆
gregarious, but one has individuals associating randomly, while in the other they only ₁₇
associate with a few select partners. As of 2019, 78% of published studies focused on ₁₈
the social structure of either birds (e.g., [8–10]) or group-living mammals [11], such as ₁₉
cetaceans (e.g., [12–15]), primates (e.g., [16–18]) and social carnivores (e.g., [19–21]). ₂₀
Understandably, researchers generally focus on highly gregarious species because their ₂₁
associations are more frequent and they are more likely to be co-located so they are ₂₂
easier to sample for social network studies, and there can be clearer applications for the ₂₃
work concerning conservation (e.g., [19]) or captive animal welfare [22]. However, this ₂₄
means that we know only relatively little about other taxonomic groups and species ₂₅
that may have sparser networks, such as solitary mammals (e.g., [23]) or reptiles ₂₆
(e.g., [24]). Less gregarious animals can still have strong friendships or specific ₂₇
preferences regarding who they should associate with, which might impact their ₂₈
behaviour and survival, including species such as African savannah elephants ₂₉
(*Loxodonta africana*), or mountain and western gorillas (*Gorilla beringei beringei* and ₃₀
Gorilla gorilla gorilla) that are generally considered highly social but can show ₃₁
relatively low average rates of social association [25–29]. ₃₂

The two core components of a social network are the individuals within the ₃₃
population, called nodes, and the associations between them, known as the edges. In ₃₄
some networks, we only need to know if a connection exists between a dyad, but the ₃₅
strength of that association (henceforth "edge weight") is not important. In others, we ₃₆
need to know the probability of a dyad being observed together. For many studies, edge ₃₇
weights are the basis of further network structural analyses, making it critical to ₃₈
calculate them reliably. ₃₉

Edge weight estimation methods are sensitive to the density of a network, so may be ₄₀
problematic when networks are sparse, as we will demonstrate in this paper. Methods ₄₁
designed for dense networks may produce biased measures of edge weight, which can ₄₂
lead to further bias in downstream calculations of node centrality or network density, ₄₃
but this is not something that has previously been examined. Similarly, connections ₄₄

between dyads that do not associate need to be modelled as zero-strength edges, but   45
these methods may be either poor at detecting non-association or falsely identify   46
zero-strength edges. In turn, these errors lead to over- and underestimation of edge   47
weight, which will affect all network studies, regardless of whether they are interested in   48
edge presence or weight. The pitfalls of different methods, which we will identify in this   49
paper, should be considered to decide if zero-inflation is likely to influence edge weight   50
estimation and if edges are more likely to be over- or underestimated.   51

Animal social network studies often struggle with limited data quality, exacerbating   52
the difficulties of analysing sparse networks using common methods. Frequentist models   53
— such as the commonly-used simple ratio index (SRI [30]) — are biased towards   54
extreme edge weights (0 and 1, respectively indicating that it is impossible for a pair to   55
ever be together or apart), especially when sample size is low [31,32], and so accept zero   56
edge weight values too readily. They are therefore liable to underestimating network   57
density by overinflating the number of dyads showing non-association. Furthermore,   58
they contain no measures of uncertainty so cannot highlight the dyads whose edge   59
weight is most likely to have suffered from low sample size. In contrast, Bayesian edge   60
weight models can, as with any Bayesian analysis, accept a non-flat prior, making them   61
more reliable when performed correctly. However, they introduce bias if the analyst   62
chooses to use inappropriate priors designed for the analysis of dense networks. These   63
edge weight priors can imply that non-association is highly implausible, producing   64
posterior distributions biased towards stronger associations. This, in turn, can lead to   65
errors in estimating network density. It is, therefore, inappropriate to use either the SRI   66
or a Bayesian model with a prior designed for a dense network when analysing networks   67
that are not fully connected, as we will later demonstrate. So far, this does not appear   68
to have been an issue in publications, as Bayesian edge weight models are a relatively   69
recent innovation and have only been used for populations with dense networks   70
(e.g., [33–40]). However, as Bayesian models generally become more popular, it is likely   71
that researchers will start to use them to calculate the edge weights of less gregarious   72
species, for which a more robust analysis method needs to be established. We can   73
overcome some of the issues with methods designed for dense networks by having very   74
large quantities of data [32,41], but this is rarely a plausible solution in animal social   75
network studies, and it does not solve all of the issues associated with edge weight   76

estimation, as we will here discuss.

We need a model that reflects how the edge weights are generated in the natural system and incorporates the likelihood of obtaining a true zero, to avoid modelling a connection where one does not exist or biasing our model by supplying inappropriate information regarding the probability of extreme edge weights. All common methods, whether they are frequentist such as SRI (e.g. [10, 21, 24, 42, 43]), half-weight index (HWI) (e.g., [44–47]; or HWIG when correcting for individual gregariousness [48]) or twice-weight index (TWI) (e.g., [49, 50]), or Bayesian such as the Bayesian framework for Inference of Social Networks (BISoN [41, 51]; e.g., [34, 38–40]) or STRAND [52, 53], model a network based on the assumption that the social processes that produce the edge weight contain only a single step: they implicitly assume a fully-connected network where every individual forms a connection with every other, and the only step in deriving the edge weight is determining how strong that connection will be. Edge weight may be correlated with dyad characteristics such as age (e.g., [49]), kinship status (e.g., [25]), phenotypic similarity (e.g., [54]), or anything else that has the potential to induce a preference (or not) for associating, but the decision over how much time to spend together is only a single step.

In contrast to these modelling assumptions, a network that is not fully connected has an additional data-generating step: before defining the weight of the connection, there is a precursory step that determines whether or not a dyad connects at all. This could be driven by a choice to avoid another individual [24, 55] or by spatial or temporal barriers that preclude individuals ever having the opportunity to associate (e.g., one member of the population dies before another is born, or no movement corridor exists between two subpopulations so their home ranges cannot overlap). This creates an additional step that defines the level of zero-inflation in the edge weights. Only once it has been defined that a dyad associates can the edge weight be determined. The sparser a network is, the more important it is to capture the first step of the underlying functional process. It is important to note that while this initial step of edge weight calculation has a Bernoulli outcome, the final network density is not a binary result of dense or sparse but a continuum composed of many dyadic edge weights that lie between zero and one. Models that assume a fully connected network become progressively worse as network density declines, but there is no intrinsic cost to always

using a method that captures both steps of the underlying social process. ₁₀₉

In this paper we examine two different network modelling approaches for their ₁₁₀ applicability to sparse networks: the SRI [30], which is the approach that has ₁₁₁ historically been the most commonly used to estimate the association between ₁₁₂ individuals, and BISoN [41, 51], which is a newer method of using Bayesian models to ₁₁₃ estimate the association index with uncertainty and incorporate more appropriate prior ₁₁₄ information. We demonstrate the application of both of these methods on a sparse ₁₁₅ network of male savannah elephants. We show how Bayesian models overcome many of ₁₁₆ the limitations of the SRI but can suffer problems if supplied with priors that are not ₁₁₇ appropriate for the particular use case. Finally, we introduce a modification of the ₁₁₈ Bayesian framework that follows a two-step data-generating process, which can deal ₁₁₉ with the pitfalls of sparse network analysis. We propose that this modification be ₁₂₀ applied to networks of any density and would benefit future animal network studies. ₁₂₁

2 Challenges of sparse social networks for current common methods _{122 123}

2.1 Common network analysis methods ₁₂₄

With gambit-of-the-group data [31], the SRI and Bayesian methods estimate the edge ₁₂₅ weight between individuals based on the total number of observations in which a dyad ₁₂₆ was together versus apart. However, despite using the same input data to estimate the ₁₂₇ same target measure, they suffer from different problems. As with any frequentist ₁₂₈ measure, SRI estimates are drawn based on the assumption that all possible edge weight ₁₂₉ values are equally likely — the equivalent of assuming a flat prior — so can produce any ₁₃₀ output edge weight, even when data are insufficient to support extreme values. To ₁₃₁ combat this, researchers often only include individuals with total sightings exceeding a ₁₃₂ threshold value, but taking a subset of the data creates its own issues, discussed below. ₁₃₃ As a Bayesian alternative, BISoN [41, 51] and other Bayesian approaches such as ₁₃₄ STRAND [52, 53] can use non-flat priors to inform the model of the prior probability of ₁₃₅ extreme values, so we can set up the model to be more uncertain about extreme edge ₁₃₆ values when the dyad observation count is low. Bayesian approaches, therefore, work ₁₃₇

better with lower sample sizes and uneven sampling than the SRI approach (see ₁₃₈ supporting information: The SRI is too trusting of extreme values). However, as with ₁₃₉ the SRI, a single-step prior is appropriate only for fully connected, or close to fully ₁₄₀ connected, networks. For more sparse networks, we should appropriately modify the ₁₄₁ prior to reflect the biological processes: here, we consider the underlying two-step social ₁₄₂ process that generates connections (or not) and then determines the weight of that ₁₄₃ connection. ₁₄₄

The SRI is typically calculated as the number of times a pair of individuals were ₁₄₅ observed associating divided by the total number of sightings that either individual was ₁₄₆ observed: ₁₄₇

$$SRI_{ij} = together_{ij}/sightings_{ij} \tag{1}$$

$$sightings_{ij} = sightings_i + sightings_j + together_{ij}$$

Where $together_{ij}$ is the number of observations in which individuals i and j were in ₁₄₈ the same group; $sightings_i$ and $sightings_j$ indicate the respective number of ₁₄₉ observations of individuals i and j in the absence of the other; and $sightings_{ij}$ is the ₁₅₀ total number of times they were seen as a whole. As a simple proportion, this measure ₁₅₁ expects extreme values such as 0 and 1 to be as likely as anything in between. It is a ₁₅₂ common misconception that frequentist models are less biased than Bayesian models ₁₅₃ because prior information is not incorporated. This has the opposite effect: a ₁₅₄ frequentist model still has a prior, but it is forced to be flat and uninformative, so it ₁₅₅ biases the results towards extreme values, making outliers just as plausible as the ₁₅₆ average value [56]. The SRI also contains no measure of uncertainty: with this method, ₁₅₇ a value of SRI = 0 occurs when a dyad is never observed in the same group, with no ₁₅₈ greater uncertainty around edge weight estimates for poorly sampled dyads than those ₁₅₉ with many observations, but we should be more certain when more data are available ₁₆₀ (Fig S1.1). ₁₆₁

These issues are the case regardless of network density, but their potential impact on ₁₆₂ analyses is exacerbated when dealing with extreme edge weight scores. In the real world, ₁₆₃ a lack of connection between individuals has important implications for social structure, ₁₆₄ information and disease transfer, and individual centrality. As the true network density ₁₆₅

declines, the likelihood of obtaining an edge weight of zero (spurious or correct)
increases, and the SRI's lack of uncertainty and treatment of all possible edge weights
as equally plausible become a greater issue. We must, therefore, be extremely careful
about modelling zero-inflation, but the flat prior and lack of uncertainty seen in the SRI
combine to make it very trusting of extreme edge weight values. It is important to note
that while we are focussing on the SRI for this paper, other frequentist edge weight
measures, including the half-weight and twice-weight indexes, all suffer the same pitfalls.
Furthermore, to avoid confusion, it is also worth noting that pre-network permutation
tests, which have known problems [57–59], do nothing to deal with the issues faced by
frequentist measures.

With Bayesian approaches, we still estimate the edge weight based on the proportion
of total dyad sightings in which they were together, but now as a model parameter
rather than a simple ratio calculation, for example:

$$together_{ij} \sim binomial(sightings_{ij}, edgeweight_{ij}) \tag{2}$$
$$logit(edgeweight_{ij}) \sim normal(\mu, \sigma)$$

Where $together_{ij}$ and $sightings_{ij}$ are the respective number of observations of a
dyad in the same group and in total, as above; and μ and σ are the prior mean and
standard deviation. Estimates of edge weight are now described by posterior
distributions, which narrow as $sightings_{ij}$ increases, capturing uncertainty in the edge
weight. Incorporating a user-defined prior can limit misleading outputs when the sample
sizes are small, as the range of likely edge weight values is reduced. Through this use of
sensible priors and quantification of uncertainty, Bayesian approaches are inherently
more capable of dealing with the small sample sizes that are common in studies of
animal social systems. While BISoN is not the only option available for the Bayesian
estimation of edge weight (see also STRAND [52,53]), we will focus on it in this paper.

2.2   SRI requires deletion of data

With a flat prior and no implicit uncertainty measure, limited data availability about a
dyad can produce unreliable SRI values, so researchers will usually set an arbitrary
minimum threshold number of observations per individual to include that node in the

analysis (e.g., [25, 29, 42, 44, 46, 49]). While this does prevent inaccurate edge weights ₁₉₃
created by the most poorly sampled dyads, it simultaneously introduces new ₁₉₄
issues [60, 61] that are often overlooked. Firstly, data are often hard-won and already ₁₉₅
limited, yet this practice requires the discarding of data, sometimes in large quantities. ₁₉₆
Secondly, data removal may not be completely-at-random [62], which is known to ₁₉₇
introduce bias [61, 63]. For example, individuals that are less neophobic, more ₁₉₈
philopatric, or possess distinctive identifying features are more likely to be included in ₁₉₉
the analysis, because the probability of their having repeat observations is greater. If a ₂₀₀
study next performs a regression analysis on calculated network measures, this ₂₀₁
non-random data deletion can directly bias the results, especially if these factors are ₂₀₂
linked to the exposure or outcome variables in question. Thirdly, the specific value of ₂₀₃
these observation thresholds often varies between studies, as the actual choice of how ₂₀₄
many sightings is considered "enough" usually depends on the number of individuals ₂₀₅
that will remain after filtering out those poorly sampled. For example, Chiyo *et al.* [25] ₂₀₆
included all male elephants with at least 15 observations in their three-year study ₂₀₇
period, while Murphy *et al.* Murphy2020-yk used an inclusion threshold of just five ₂₀₈
observations per four year sampling period. Therefore, comparing the results of different ₂₀₉
studies and identifying how many sightings are truly "enough" is difficult. Even with ₂₁₀
several sightings per individual, scores of 0 and 1 can be overinflated relative to their ₂₁₁
true underlying network, because there are only a certain number of possible edge ₂₁₂
weight values when sighting counts are low (Fig S1.1). ₂₁₃

In contrast, employing a Bayesian framework prevents the requirement for data ₂₁₄
inclusion thresholds, which improves data quality and reduces biases caused by the ₂₁₅
possibility of data missing not-completely-at-random. ₂₁₆

2.3   Bayesian models facilitate the use of more appropriate prior ₂₁₇ distributions ₂₁₈

A model prior should always be carefully considered, which is not possible under the ₂₁₉
frequentist framework. Given how few papers have thus far used a Bayesian method for ₂₂₀
edge weight estimation, most social network analyses are based on flat frequentist priors, ₂₂₁
which are generally inappropriate (if nothing is known at all of the population in ₂₂₂

question, then a flat prior may still be a reasonable choice, though this would be ²²³ extremely unusual). The *bisonR* [51] package for running BISoN models supplies very ²²⁴ broad default priors, but the package authors strongly recommend tailoring them to ²²⁵ each specific scenario. Most social networks contain at least some true zeros. Any ²²⁶ model, frequentist or Bayesian, that uses a flat or very broad prior will struggle to ²²⁷ model these zero values because they are modelling only the second step of what is ²²⁸ inherently a two-step process. ²²⁹

Unlike the SRI, with BISoN we can change our model structure to incorporate two ²³⁰ steps — first to determine if a dyad will associate at all, and second to define the ²³¹ strength of that connection — if we supply it with an appropriate prior, which package ²³² default priors are not. Here, we have presented a prior allowing BISoN to model both ²³³ steps without changing the social network analysis workflow. While we could instead ²³⁴ use a mixture model with zero-inflation to model a two-step process, using either a ²³⁵ frequentist (see supporting information: Standard zero-inflated model) or Bayesian ²³⁶ framework, we can keep the overall method relatively simple by changing only the prior ²³⁷ structure. All network models should include some option to account for zero-inflation, ²³⁸ and selecting the best method for measuring edge weights requires careful consideration: ²³⁹ Bayesian techniques require integration of current understanding of the system, and a ²⁴⁰ prior predictive check, which can lead to biases if carried out using inappropriate priors, ²⁴¹ but their ability to estimate uncertainty and accept non-flat priors makes them more ²⁴² appropriate than frequentist methods. ²⁴³

2.4   Prior selection in a Bayesian edge weight model ²⁴⁴

Bayesian methods have been successfully applied to dense networks, using both ²⁴⁵ simulated and empirical data [34,64,65], and their use in primate research has been ²⁴⁶ discussed in subsequent publications [66,67]. However, the best way to use them for ²⁴⁷ sparse networks has not yet been evaluated. Prior choices require careful consideration ²⁴⁸ for any analysis, not just social networks. Fig 1a shows an improper uniform prior, in ²⁴⁹ which all edge weights are considered equally likely before exposure to the data. By ²⁵⁰ relaxing the assumption that all edge weights are equally probable, we can make the ²⁵¹ prior more informative. When we demonstrate the use of BISoN on an example dataset, ²⁵²

we first show the outcome of this uniform prior, followed by the default prior supplied in the *bisonR* package [51]: a wide symmetrical prior (logit(*edge weight$_{ij}$*) $\sim normal(0, 2.5)$) that presumes very little about the average edge weight (Fig 1b). Third, we show the result of using a strong asymmetrical prior (*edge weight$_{ij}$* $\sim beta(1, 5)$) that indicates weaker edges are the most likely but still allows for some strong associations (Fig 1c). How we define the prior depends on our knowledge and assumptions of the system: the sparser the network, the more right-skewed the prior distribution should be.

Fig 1. Prior assumptions for single-step BISoN models. Prior assumptions for a) SRI or a BISoN model with a uniform prior, b) a BISoN model with a weak, symmetric prior (logit normal, $\mu = 0$, $\sigma = 2.5$) and c) a BISoN model with a strong asymmetric prior (beta, $a = 1, b = 5$).

Biases can appear in the edge posterior when the prior distribution does not reflect the underlying social process: knowing that the network is of a lower density can allow us to adapt our prior to a more suitable distribution, and the assumptions we can make based on that knowledge are critical to dealing with poorly sampled dyads. When the prior average is a long way from that of most dyads, it induces a systematic shift in the direction the data pushes the posterior from the prior. In a sparse network, many of the dyads will never associate, so a prior designed for a dense network will usually be shifted towards peaking at zero by the data. The data for poorly sampled dyads will, by definition, not be able to shift the posterior as far from the prior as the data from the well-sampled dyads, so if they all shift in the same direction, the level of sampling could potentially have a greater effect on the final edge weight than the actual raw data values. When this shift is consistently towards zero, poorly sampled dyads, on average, receive edge weight distributions with higher average values than well-sampled dyads. It is difficult to imagine a scenario in which the number of observations of a dyad would

be strongly dependent on the strength of their social bond or vice versa, therefore this correlation highlights a problem with the edge weight data. We show this happening in the example analysis. Again, without well-sampled data, the choice of prior (including the SRI's uniform prior) becomes critical to the success of the edge weight estimation and all subsequent analyses.

3 Introducing a two-step prior for sparse networks

By varying the prior distribution, Bayesian methods can be tailored to a sparser or denser network. The modification that we propose simply extends this ability: to alter the BISoN prior structure to reflect a two-step social process producing the edge weights, in which step one defines the extent of zero-inflation and step two the weight of the non-zero social bonds. We structure our model to use two separate edge weight priors, selecting which to use depending on whether a dyad has ever been observed together (Fig 2), such as:

$$together_{ij} \sim binomial(sightings_{ij}, edgeweight_{ij}) \qquad (3)$$

$$edgeweight_{ij} \sim beta(a, b)$$

$$\text{If } together_{ij} = 0: \ a < 1 \text{ and } b = \text{large}$$

$$\text{If } together_{ij} > 0: \ a = 1 \text{ and } b = \text{small}$$

Where again $together_{ij}$ refers to the number of times a pair was observed in the same group, and $sightings_{ij}$ is the number of times they were observed in total. This is not a major change to the method itself, as BISoN has always been capable of using a conditional prior. The change is that by capturing both parts of the underlying social process in a conditional prior, we only need to change one part of our overall method.

The new prior is simple to apply: if a dyad is never observed in the same social group, then the model will use a prior that increases the probability of an edge weight of zero; if a dyad has been observed interacting on at least one occasion, it will receive an edge weight calculated using a prior with less or no zero-inflation. This, therefore, allows the model to identify the most appropriate prior and incorporate the probability of non-association as an initial step before identifying the best edge weight distribution per dyad. As before, poorly sampled dyads will return an edge weight more similar to

Fig 2. Conditional edge weight prior for a BISoN model of a sparse network.
The blue line indicates the lower prior that will be selected when a dyad has never been
observed in the same group (in this case $beta(a = 0.7, b = 10)$). The purple line shows
the upper prior choice for when a dyad has been observed grouping together on at least
one occasion (in this case $beta(a = 1, b = 5)$). While in this model we have chosen to
still have this prior peak at zero, this reflects that we still expect very low edge weights
amongst those that do associate, and the data immediately overwhelm the possibility of
zero being an option. Note that the upper prior is the same as the strongly right-skewed
prior shown above.

their respective prior, while well-sampled dyads will shift further. The model becomes

less uncertain about a zero edge as the total sighting count increases: the uncertainty in

these values indicates that zero is the most likely edge weight but still allows stronger

relationships as a possibility when sampling is poor. In contrast, for a dyad observed

together on at least one occasion, we use a wider prior that allows higher edge weights

to be observed, again becoming more confident in the results as total sampling increases.

The most logical step here may appear to be to make our regular prior more

right-skewed so it is closer to zero. However, if we were to use only an extremely

right-skewed prior, such as the lower prior of this combination, to encapsulate the

zero-inflation and the true associations, a prohibitively large amount of association data

would be required for any truly strong edge weights to register.

The values a researcher might choose for the prior depend on their study species and

expected edge weights. In the following example, we have opted to have the prior for

associated dyads peaking at zero, but this is immediately overwhelmed by the data.

This produces edge weight values that are still far more likely to be close to zero than to

one, based on the evidence that male elephants do not appear to form strong

relationships with one another [25, 26, 29]. If we were expecting to see a strongly ³¹⁹
bimodal distribution in the final edge weights, with some dyads never associating and ³²⁰
others showing consistent association, we would shift our upper prior further from zero. ³²¹

4   Example dataset

For this example, we used data on sparse networks based on associations between male ³²³
elephants of the Mosi-Oa-Tunya National Park (MOTNP), Zambia, collected from May ³²⁴
2016 until December 2017 (all permissions for data collection were supplied by ³²⁵
Department of National Parks and Wildlife, Zambia: permit number ³²⁶
TJ/DNPW/101/13/18). No additional ethics permission was required alongside the ³²⁷
research permit, but to avoid causing any stress to the elephants, we monitored them ³²⁸
from a vehicle from a safe distance, and kept all noise to a minimum while in the ³²⁹
National Park. ³³⁰

This population is relatively small, totalling approximately 500 individuals, with ³³¹
similar numbers of males and females. The park is well connected to large areas of ³³²
suitable habitat, and the elephants are free-ranging, so individuals may leave the area ³³³
for long periods. Data were collected during daily drives through the park, marking ³³⁴
down the identities of all individuals. From these observations, we created 504 days of ³³⁵
gambit-of-the-group [31] data comprising 213 males sighted in 481 groups. Finally, we ³³⁶
converted this to a data frame containing every possible dyad with counts of the total ³³⁷
number of times that dyad had been observed and how many of those occasions they ³³⁸
were in the same group. Data analyses were run by calling Stan (version 2.26.1 [68]) ³³⁹
from R (version 4.2.1 [69]) using the *cmdstanr* [70] package. ³⁴⁰

5   Results from empirical data

5.1   Simple Ratio Index (SRI)

When we calculate the SRI values for this population (Fig 3a), we get an extremely high ³⁴³
proportion of zeros (17385, equating to 77.00% of dyads). While a much lower number ³⁴⁴
of dyads receive an SRI of one (12 dyads, equating to just 0.05% of the total ³⁴⁵
population), the likelihood of these apparent permanent alliances would be exceedingly ³⁴⁶

small given the population average.

Fig 3. Edge weights for MOTNP elephants using the SRI. a) The vast majority of dyads have an SRI score (green points) of zero, indicating that the output is certain that these pairs never associate with one another, regardless of the number of sightings. b) This translates to a large split between the edge weights calculated for poorly sampled dyads that have (purple line) or have not (blue line) ever been observed grouping together. We see a very strong effect of dyad sighting count on the edge weight for dyads observed together on at least one occasion, until we reach around 12 sightings per dyad.

Given this, the standard next step with the SRI would be to set a threshold for the minimum number of sightings per node required to be included in the analysis. If we set this threshold to five sightings per node, the proportion of dyads with an SRI score of zero shifts down to 46.13%, and all dyads scoring a 100% association rate are removed. A threshold of 10 sightings per individual drops the 0% association rate down to 26.25%, showing high instability in the SRI outputs. This is highlighted by Fig 3b, which shows the association between the total number of observations per dyad, and the calculated edge weight for that dyad, stemming from the extreme number of zeros calculated by the SRI. The overall results would strongly depend on the threshold value we choose, which, as previously stated, is an arbitrary choice.

We could potentially use Fig 3b to determine a less arbitrary inclusion threshold by selecting the point at which the patterns in the edge weights weaken. However, this example shows that this is somewhere between 10 and 20 sightings per dyad. This would require removing at least all individuals with fewer than 10 observations. In this case, that would take our total sample size down from 213 to just 74 elephants. To remove 65% of the sample could severely bias our results if the remaining portion is not entirely representative of the total population. Therefore, we cannot rely on objective means to determine the threshold level.

5.2 Bayesian model with a single-step prior

When we use the BISoN framework with a single-step prior, we can see a limited effect of prior shape on the posterior distribution, but the inclusion of zero as a possible edge value is very important. When using a fully flat or highly right-skewed prior, in which zero is a possible option, we see very similar posterior distributions (Fig 4a and c), whereas for a normal prior peaking at 0.5 for which extreme edge weights are unlikely (and actual values of 0 and 1 are impossible) we see an overall much broader and less skewed posterior (Fig 4b).

Fig 4. Edge weights for MOTNP elephants using BISoN models with single-step priors. Results using BISoN with uniform (left), default (middle) and right-skewed (right) priors (insets in top row show the respective prior for the column). a-c) The top row shows the posterior distribution for 100 randomly sampled dyads, with green curves indicating pairs that have never been observed in the same group and purple curves showing dyads that were observed in the same group on at least one occasion. The right-skewed and uniform priors produce posterior distributions that initially appear very similar because they still allow an edge weight of zero to be a plausible draw, whereas the default prior draws all values together. d-f) The bottom row shows the effect of total sighting count per dyad on their median (green points) and full (yellow points) posterior distribution. The blue line indicates dyads never observed together, while the purple line shows dyads observed together at least once. As the prior becomes increasingly right skewed, the effect of sighting counts on median edge weight declines.

We now also have a situation in which there is a negative correlation between mean edge weight and total observations per dyad at low sighting counts, especially when using very broad priors. Looking at Figs 4d and 4e, we can see that poorly sampled dyads appear to show higher edge weights than well-sampled ones (an effect the authors have also seen in social network data for macaques). Again, this is most severe for the logit normal prior, which draws all of the values towards the middle instead of creating variation between dyads. The effect is also stronger for the flat prior than for the right-skewed, for which the trend is almost entirely removed because those dyads with the fewest sightings do not have sufficient data to discredit high edge weight model draws, so the posterior is very wide, with high median values. Similar reasoning also explains the differences between BISoN with a flat prior and the SRI: with the SRI, low sighting counts largely score exactly zero, whereas with BISoN low sighting counts are given a very wide posterior (shown in yellow) so their median (green) is substantially above zero.

Based on these graphs, the best solution would appear to be to use BISoN with a strongly right-skewed prior, which can model the edge weights without creating strong patterns of social association based on data quality. However, this prior may not allow sufficient movement towards a higher edge value for dyads showing a strong social bond. Therefore, we need a prior that better reflects the underlying social process, such that it allows for some dyads to attain higher edge weights, but without increasing the prior average and causing poorly sampled dyads to consistently obtain a higher edge weight than well-sampled ones. Our modification to BISoN to use a prior that reflects the two-step underlying social process facilitates edge weight estimation in a manner robust to zero inflation and which allows stronger edges to be estimated.

5.3 Bayesian model with a two-step prior

When using the two-step prior, the posterior distributions of the edge weights (Fig 5a) appear very similar to those with single-step priors, though with greater confidence in some of the weakest edges, as we would expect. The median edge weight is associated much more weakly with the total number of sightings than with the broader priors (Fig 5b), even at low dyad sighting counts. While the shape of the curve for pairs seen

grouping together is very similar to the single-step right-skewed prior, the overall effect   404
of low sighting count has been removed by the model's increased ability to obtain   405
median estimates very close to zero. We now have edge estimates that are more reliable   406
than the previous BISoN models, as they use a prior that allows the modelling of   407
zero-inflation.   408

**Fig 5. Edge weights for MOTNP elephants using a BISoN model with a
conditional prior.** a) posterior density lines for every dyad (blue indicates density
lines drawn for pairs never observed in the same group, purple for those seen together
on at least one occasion); and b) edge weight as a response to number of sightings per
dyad (full posterior distribution in yellow, median per dyad in green). While some effect
of sighting count on the median edge weight for individual dyads remains, the impact is
much reduced compared with the previous BISoN plots, and it lacks the values of 100%
association observed with the SRI.

5.4 Using downstream network measures to assess edge weight   409
   quality   410

In general, we should be able to see directly in our edge weight values if there is a   411
problem in our estimation method. Sometimes however, the impacts of the method are   412
masked until subsequent analyses of the outputs. For example, in an analysis to identify   413
the effects of individual trait values on network centrality, we may find that our   414
centrality values contain spurious patterns, such as in the number of sightings or   415
number of different partners that an individual is observed with compared to network   416
centrality. Here, we will show that using an inappropriate single-step prior for edge   417
weight can severely impact downstream analyses of network structure. We explore this   418
using eigenvector centrality, a measure of how connected an individual is to other   419

well-connected individuals, as it is commonly used and is the centrality measure most ₄₂₀ robust to causal analysis [71]. As with the edge weights, we can plot measures of ₄₂₁ centrality calculated per model against a) the number of other individuals in the ₄₂₂ population with whom they were never observed associating, and b) their individual ₄₂₃ sighting count, and look for any unexpected trends that may indicate problems with the ₄₂₄ outputs. ₄₂₅

When we plot the SRI-based eigenvector data against the number of non-associated ₄₂₆ elephants, we see a negative correlation (Fig 6a), exactly as we would expect. If, ₄₂₇ however, we compare the SRI-based centrality to the number of sightings per individual, ₄₂₈ we see a positive correlation for poorly sampled dyads (Fig 6b), which indicates that ₄₂₉ there is a problem with the estimation. This relationship is created by the extreme edge ₄₃₀ values caused by limited numbers of samples, especially for elephants observed fewer ₄₃₁ than 10 times. This could be used as an objective method of determining an inclusion ₄₃₂ threshold, but as before, this would require the discarding of more than half the data. ₄₃₃ We obtain similar trends when calculating the edge weight using a zero-inflated mixture ₄₃₄ model (Fig S2.2). ₄₃₅

Fig 6. Eigenvector centrality estimates based on the SRI edge weight calculations. a) There is a negative relationship between SRI-based centrality and the total number of potential partners with whom a focal was never observed associating, which is as expected. However, b) indicates an issue with the SRI-based centrality in that there is a strongly positive effect of the number of observations per individual on their social position at low observation counts.

We see something very different when we look at the eigenvector centrality plots ₄₃₆ based on the single-step BISoN models. Startlingly, there is an apparent reversal of the ₄₃₇ effect of the number of non-associated individuals in the population, such that elephants ₄₃₈

observed with the most partners receive the lowest centrality scores, while those 439
observed with the fewest partners receive the highest (Fig 7a-c). This is, of course, 440
nonsense. Eigenvector centrality as a measure combines the number of association 441
partners an individual has with the number of association partners their partners have. 442
The most central elephants therefore cannot be those with the fewest association 443
partners. The edge weight estimation process must have an underlying error to create 444
this trend. 445

Fig 7. Eigenvector centrality estimates based on BISoN edge weight models with single-step priors. Priors per column are shown as insets on the top row: left to right uniform, default and right-skewed. In all graphs, yellow points indicate the uncertainty around the median, displayed in blue. a-c) The top row indicates the impact of number of dyad partners with whom an individual was never observed associating, where we see a spurious positive effect. The more right-skewed prior suffers more weakly from this than the uniform-based or default-based centrality, but still sufficiently to create serious cause for concern. There is no biologically or statistically meaningful explanation for this to be a correct trend. The pattern is explained by d-f) in which the elephants with the most observations occupy the social positions of lowest centrality.

This seemingly impossible pattern occurs because of an underlying trend in the 446
number of observations per elephant (Fig 7d-f). Individuals with fewer overall 447
observations contribute fewer sightings to dyad observations. When the edge weight 448
prior is designed for dense networks, the posterior for most dyads is shifted closer to 449

zero from the prior. As the data for poorly sampled dyads are less influential on the ⁤450
distance the posterior can move, dyads with fewer observations consistently appear to ⁤451
have a stronger edge weight than well-sampled dyads. When this translates through to ⁤452
individual effects, and poorly sampled individuals end up in more poorly sampled dyads, ⁤453
it is exaggerated so individuals observed rarely have the strongest average edge weights ⁤454
and, therefore, the highest eigenvector centrality. ⁤455

Finally, we can now consider eigenvector centrality calculated from the edge weights ⁤456
produced using BISoN with a two-step prior and see that we successfully avoid all ⁤457
downfalls from the other methods. Changing to a conditional prior has successfully ⁤458
reversed the spurious correlation between number of associates and node centrality ⁤459
(Fig 8a) by removing the effect of sighting count on eigenvector (Fig 8b). We have ⁤460
avoided creating estimates biased by the data quality, so any subsequent analyses using ⁤461
these data will be more reliable than those calculated using an improper prior while also ⁤462
avoiding the need to remove poorly sampled dyads as we would with the SRI. ⁤463

Fig 8. Eigenvector centrality estimates based on BISoN edge weight model with a two-step prior. a) Using our modified prior (inset), to reflect a two-step social process, we have successfully restored the negative trend in node centrality such that those seen with the most different dyad partners on average have the highest centrality, and those with the fewest have the lowest. b) Individual sighting count now has extremely limited effect on eigenvector centrality.

This quick sanity check of the eigenvector against the number of associates can ⁤464
instantly highlight if there is a problem with the edge weights, and show how sighting ⁤465
count could affect centrality. Our modification does not entirely eliminate this, and care ⁤466
should be taken to look for any anomalies. For example, in the MOTNP data, there is a ⁤467
single sighting where 89 elephants crossed the river and came up into the surrounding ⁤468

mopane forest together, of which 31 were males with known identities. For two males, this was their only observation. Therefore, these two individuals were "always" seen with one another and are in the "seen grouping together at least once" group for 30 dyad pairs. This means that, despite having very wide uncertainty margins for all dyads due to low total counts, they will still have an above-average median edge weight due to the nature of their single observation. When carried forward to a centrality analysis, these elephants will display a slightly elevated eigenvector centrality. Using the modified BISoN framework means this elevation is much lower than when using an unconditional prior or the SRI, and the number of individuals for which this could be an issue is reduced, limiting the impact on the regression. However, we still need to be aware of the potential for data artefacts that an occurrence such as this may leave. It is worth noting that while none of these methods is perfect, by using a method that incorporates current knowledge, quantifies uncertainty, and does not induce issues in downstream analytical data frames, we can rest assured that we are using the best possible option.

6   Discussion

We have shown that current animal social network analysis methods do not cope well with sparse network data. Frequentist measures such as the SRI suffer from an innate and unavoidable reliance on a flat prior, which overestimates the confidence that can be placed on extreme edge weight values. Combine this prior with an inability to incorporate uncertainty, and researchers must rely on setting arbitrary inclusion thresholds when analysing poorly sampled dyads, leading to data loss and potentially biased results. While Bayesian models can account for these problems in dense networks, they can still produce edge weights overly influenced by the frequency of sightings per dyad when inappropriate priors are applied to sparse networks. By modifying the BISoN framework to use a conditional prior, reflecting a two-step underlying social process, we can create a model where the edge weights are more reliable than those from the SRI, a zero-inflated mixture model, or an uninformed Bayesian model. As such, they are no longer overly dependent on data quality, which is commonly poor in animal social network studies.

When checking our models, we used several diagnostics to compare output quality

and determine if our BISoN modification improved the current methods. The first question is whether a result fits with what we know of the population: the SRI produces far too many dyads with zero probability of ever associating, even when we set a high inclusion threshold. At just 66 km^2 [72], the MOTNP is not sufficiently large to preclude the interaction of elephants by simply never meeting, with a bull easily able to cross it in a single day. It is, therefore, extremely unlikely that this many dyads are truly never together, as there would not be enough space for them all to move without encountering other elephants. This allows us to make two informed assumptions: i) the proportion of zeros produced by the SRI is overinflated; and ii) that well-sampled dyads that are never recorded associating have elected not to do so, rather than happening to never encounter one another. When using BISoN, the conditional prior can help to alleviate the effect of sighting count on median edge weight by using the broader prior with a higher average only when we have evidence that the dyad has elected to sometimes associate. While this solution does not entirely remove the impact of limited sightings on edge weight, it does minimise it compared to a single-step prior.

At first glance at the edge weights alone, it seems that the conditional prior is not particularly advantageous over the right-skewed single-step prior: the basic outputs for the edge weight look fairly similar. However, we have modelled the zero-inflation, which, while it appears to have only had a limited effect on the edge weights themselves, has a very noticeable and important impact on our subsequent analyses. The node centrality values produced can act as a suitable sanity check. While there appears to be little difference between the right-skewed BISoN outputs and the two-step BISoN outputs when we consider only the edge weights, the eigenvector centrality from a single-step prior in a sparse network creates an impossible situation in which individuals with more partners receive a lower centrality score. This trend is corrected by the use of a two-step prior. We can use such checks to ensure our outputs make sense in downstream analyses.

Selecting the two distributions remains important for successful edge weight estimation when applying a two-step prior to a sparse network model. Making the zero-centred prior too strong will not allow enough variation when data are limited, but if too weak some of the possible edge weights may still present as too high and create trends in edge weight and centrality based on sighting counts. Similarly, if the probability mass for the non-zero prior is not shifted far enough from zero, any stronger

relationships will be missed; shift it too much, and the same trends of increased edge ²³¹
weight with reduced sightings will be observed within the group who have been ⁵³²
observed together. In this instance, we have opted for a large overlap between the priors ⁵³³
such that dyads only seen once or twice will receive relatively similar posterior ⁵³⁴
distributions regardless of which prior they are assigned to. By giving both priors a ⁵³⁵
peak at zero, a dyad observed just once but who were together at that time will still ⁵³⁶
have a low average edge weight (though the zero-peak is always immediately overcome ⁵³⁷
by the data), matching our assumptions of a sparse network. This particular conditional ⁵³⁸
prior is the one that works for our specific elephant population and is not a silver bullet ⁵³⁹
prior that will work for all network analyses. ⁵⁴⁰

Every author should identify their own priors to create a set that works for their ⁵⁴¹
own population, which does not have to require that both priors peak at zero: unlike ⁵⁴²
with the male elephants, a network can be sparse but still contain some strong ⁵⁴³
associations, in which case an upper prior that allows higher edge weights may be ⁵⁴⁴
preferable. For example, if we were to repeat this analysis with the female elephants of ⁵⁴⁵
MOTNP, we would expect a more bimodal distribution in our output because of the ⁵⁴⁶
fission-fusion structure of female elephant society [73–75]. In this case, we could shift ⁵⁴⁷
our non-zero prior to be more distinct from the lower, producing a bimodal distribution ⁵⁴⁸
of edge weights, indicating strong social bonds within certain dyads. It could even be ⁵⁴⁹
taken a step further to add more prior distributions as necessary. Again, using the ⁵⁵⁰
example of the female elephants, we may, in this scenario, want the prior to reflect a ⁵⁵¹
three-step social process: a zero-centred prior for those never seen in the same social ⁵⁵²
group, a right-skewed or more central prior for those who have been seen grouping ⁵⁵³
together but who are unrelated; and a left-skewed prior for related females. Hierarchical ⁵⁵⁴
networks in which we see kinship groups within subgroups, which themselves join ⁵⁵⁵
together to form larger association groups are very common in animal societies. This ⁵⁵⁶
method could be easily expanded to allow for the use of different priors depending on ⁵⁵⁷
which level of association a dyad is expected to fall into. ⁵⁵⁸

It should be noted that the assessment of the edge weight is rarely the end of the ⁵⁵⁹
analysis. Generally, we calculate the edge weights so that we can later assess trends in ⁵⁶⁰
node-, dyad- or network-level characteristics. Therefore, the edge weight calculation ⁵⁶¹
only creates the data ready for a future regression model. There is no issue with ⁵⁶²

checking the edge weights produced — just as we would check for errors in any other raw data — and running the model again using an improved prior if there is a problem. If edge weight outputs are dependent on the dyad sighting count, or there are issues with downstream calculations such as the centrality, there is no issue with adjusting the prior, so long as it still reflects our original understanding of the population.

In conclusion, the shape of the prior is, as with all analyses, the key to estimating edge weight, and using a prior structure that reflects the underlying social process is especially important for sparse networks, though the concept applies to dense networks too. With a frequentist model, this means considering the implications of a flat prior and including in our discussion any impacts this may have on the results. For a Bayesian model, it means tailoring the prior to our individual situation rather than using the broad default priors supplied by statistical packages, which will usually allow too much probability density over highly improbable edge weight values. In the case of sparse networks, whether the model is Bayesian or frequentist, we need to model the zero-inflation induced by the first step of the underlying social process. While a zero-inflated mixture model can do this better than the SRI, we still recommend that a Bayesian model with a conditional prior is a better option. This means allowing edge weights to sample down to zero using a lower prior and the possibility for stronger relationships with an upper prior. When investigating denser networks, our prior distributions may differ, but we can still consider the underlying social process as containing more than one step, even if the reduced proportion of zero relationships masks the effect. We can therefore use a conditional prior in a Bayesian framework to identify greater nuance in social network data, for networks of any density.

7   Acknowledgements

We would like to thank the African Lion and Environmental Research Trust (ALERT) for providing the example dataset and the Zambian Department of National Parks and Wildlife (DNPW) for allowing its collection and analysis. Analyses for this paper were run on the Viking cluster, a high-performance computing facility provided by the University of York. We are grateful for computational support from the University of York, IT Services and the Research IT team.

References

1. Singh J. Collaborative Networks as Determinants of Knowledge Diffusion Patterns. Manage Sci. 2005;51(5):756–770.

2. Franz M, Nunn CL. Network-based diffusion analysis: a new method for detecting social learning. Proc Biol Sci. 2009;276(1663):1829–1836.

3. Hobaiter C, Poisot T, Zuberbühler K, Hoppitt W, Gruber T. Social network analysis shows direct evidence for social transmission of tool use in wild chimpanzees. PLoS Biol. 2014;12(9):e1001960.

4. Kretzschmar M, Morris M. Measures of concurrency in networks and the spread of infectious disease. Math Biosci. 1996;133(2):165–195.

5. Cross PC, Lloyd-Smith JO, Bowers JA, Hay CT, Hofmeyr M, Getz WM. Integrating association data and disease dynamics in a social ungulate: bovine tuberculosis in African buffalo in the Kruger National Park. Ann Zool Fennici. 2004;41(6):879–892.

6. Naug D. Structure of the social network and its influence on transmission dynamics in a honeybee colony. Behav Ecol Sociobiol. 2008;62(11):1719–1725.

7. Silk MJ, Croft DP, Delahay RJ, Hodgson DJ, Boots M, Weber N, et al. Using Social Network Measures in Wildlife Disease Ecology, Epidemiology, and Management. Bioscience. 2017;67(3):245–257.

8. Silk MJ, Croft DP, Tregenza T, Bearhop S. The importance of fission-fusion social group dynamics in birds. Ibis. 2014;156(4):701–715.

9. Farine DR, Aplin LM, Sheldon BC, Hoppitt W. Interspecific social networks promote information transmission in wild songbirds. Proc Biol Sci. 2015;282(1803):20142804.

10. Beck KB, Farine DR, Firth JA, Sheldon BC. Variation in local population size predicts social network structure in wild songbirds. J Anim Ecol. 2023;92(12):2348–2362.

11. Webber QMR, Vander Wal E. Trends and perspectives on the use of animal social network analysis in behavioural ecology: a bibliometric approach. Anim Behav. 2019;149:77–87.

12. Foster EA, Franks DW, Morrell LJ, Balcomb KC, Parsons KM, van Ginneken A, et al. Social network correlates of food availability in an endangered population of killer whales, *Orcinus orca*. Anim Behav. 2012;83(3):731–736.

13. Weiss M. Developing methods and applications for the analysis of cetacean social networks. University of Exeter; 2021.

14. Gerber L, Connor RC, Allen SJ, Horlacher K, King SL, Sherwin WB, et al. Social integration influences fitness in allied male dolphins. Curr Biol. 2022;32(7):1664–1669.e3.

15. Marfurt SM, Allen SJ, Bizzozzero MR, Willems EP, King SL, Connor RC, et al. Association patterns and community structure among female bottlenose dolphins: environmental, genetic and cultural factors. Mamm Biol. 2022;102(4):1373–1387.

16. Kasper C, Voelkl B. A social network analysis of primate groups. Primates. 2009;50(4):343–356.

17. Brent LJN, Lehmann J, Ramos-Fernández G. Social network analysis in the study of nonhuman primates: a historical perspective. Am J Primatol. 2011;73(8):720–730.

18. Sueur C, Jacobs A, Amblard F, Petit O, King AJ. How can social network analysis improve the study of primate behavior? Am J Primatol. 2011;73(8):703–719.

19. Abell J, Kirzinger MWB, Gordon Y, Kirk J, Kokeŝ R, Lynas K, et al. A social network analysis of social cohesion in a constructed pride: implications for ex situ reintroduction of the African lion (*Panthera leo*). PLoS One. 2013;8(12):e82541.

20. Dorning J, Harris S. Individual and seasonal variation in contact rate, connectivity and centrality in red fox (*Vulpes vulpes*) social groups. Sci Rep. 2019;9(1):20095.

21. Stratford K, Stratford S, Périquet S. Dyadic associations reveal clan size and social network structure in the fission–fusion society of spotted hyaenas. Afr J Ecol. 2020;58(2):182–192.

22. Kleinhappel TK, John EA, Pike TW, Wilkinson A, Burman OHP. Animal welfare: a social networks perspective. Sci Prog. 2016;99(Pt 1):68–82.

23. Connor T, Frank K, Qiao M, Scribner K, Hou J, Zhang J, et al. Social network analysis uncovers hidden social complexity in giant pandas. Ursus. 2023;2023(34e9):1–13.

24. Leu ST, Farine DR, Wey TW, Sih A, Bull CM. Environment modulates population social structure: experimental evidence from replicated social networks of wild lizards. Anim Behav. 2016;111:23–31.

25. Chiyo PI, Archie EA, Hollister-Smith JA, Lee PC, Poole JH, Moss CJ, et al. Association patterns of African elephants in all-male groups: the role of age and genetic relatedness. Anim Behav. 2011;81(6):1093–1099.

26. Goldenberg SZ, de Silva S, Rasmussen HB, Douglas-Hamilton I, Wittemyer G. Controlling for behavioural state reveals social dynamics among male African elephants, *Loxodonta africana*. Anim Behav. 2014;95:111–119.

27. Rosenbaum S, Maldonado-Chaparro AA, Stoinski TS. Group structure predicts variation in proximity relationships between male–female and male–infant pairs of mountain gorillas (Gorilla beringei beringei). Primates. 2016;57(1):17–28.

28. Morrison RE, Eckardt W, Stoinski TS, Brent LJN. Comparing measures of social complexity: larger mountain gorilla groups do not have a greater diversity of relationships. Proc Biol Sci. 2020;287(1931):20201026.

29. Murphy D, Mumby HS, Henley MD. Age differences in the temporal stability of a male African elephant (*Loxodonta africana*) social network. Behav Ecol. 2020;31(1):21–31.

30. Cairns SJ, Schwager SJ. A comparison of association indices. Anim Behav. 1987;35(5):1454–1469.

31. Franks DW, Ruxton GD, James R. Sampling animal association networks with the gambit of the group. Behav Ecol Sociobiol. 2010;64(3):493–503.

32. Hoppitt WJE, Farine DR. Association indices for quantifying social relationships: how to deal with missing observations of individuals or groups. Anim Behav. 2018;136:227–238.

33. Gettler LT, Redhead D, Dzabatou A, Lew-Levy S. BaYaka forager food sharing networks in the Congo Basin: The roles of gender homophily and kin sharing. Am J Biol Anthropol. 2023;181(1):59–69.

34. Motes-Rodrigo A, Albery GF, Negron-Del Valle JE, Philips D, Cayo Biobank Research Unit, Platt ML, et al. Social responses to a hurricane drive greater epidemic risk. bioRxiv. 2023;.

35. Redhead D, Maliti E, Andrews JB, Borgerhoff Mulder M. The interdependence of relational and material wealth inequality in Pemba, Zanzibar. Philos Trans R Soc Lond B Biol Sci. 2023;378(1883):20220288.

36. Redhead D, Ragione AD, Ross CT. Friendship and partner choice in rural Colombia. Evol Hum Behav. 2023;44(5):430–441.

37. Debetencourt B, Barry MM, Arandjelovic M, Stephens C, Maldonado N, Boesch C. Camera traps unveil demography, social structure, and home range of six unhabituated Western chimpanzee groups in the Moyen Bafing National Park, Guinea. Am J Primatol. 2024;86(2):e23578.

38. Sadoughi B, Mundry R, Schülke O, Ostner J. Social network shrinking is explained by active and passive effects but not increasing selectivity with age in wild macaques. Proc Biol Sci. 2024;291(2018):20232736.

39. Siracusa ER, Pavez-Fox MA, Negron-Del Valle JE, Phillips D, Platt ML, Snyder-Mackler N, et al. Social ageing can protect against infectious disease in a group-living primate. bioRxiv. 2024;.

40. Testard C, Shergold C, Acevedo-Ithier A, Hart J, Bernau A, Negron-Del Valle JE, et al. Natural disaster alters the adaptive benefits of sociality in a primate. bioRxiv. 2024;.

41. Hart J, Weiss MN, Franks D, Brent L. BISoN: A Bayesian framework for inference of social networks. Methods Ecol Evol. 2023;14(9):2411–2420.

42. Croft DP, Krause J, Darden SK, Ramnarine IW, Faria JJ, James R. Behavioural trait assortment in a social network: patterns and implications. Behav Ecol Sociobiol. 2009;63(10):1495–1503.

43. de Silva S, Ranjeewa ADG, Kryazhimskiy S. The dynamics of social networks among female Asian elephants. BMC Ecol. 2011;11:17.

44. Bejder L, Fletcher D, BrÄger S. A method for testing association patterns of social animals. Anim Behav. 1998;56(3):719–725.

45. Lusseau D, Schneider K, Boisseau OJ, Haase P, Slooten E, Dawson SM. The bottlenose dolphin community of Doubtful Sound features a large proportion of long-lasting associations. Behav Ecol Sociobiol. 2003;54(4):396–405.

46. Napper CJ, Hatchwell BJ. Social dynamics in nonbreeding flocks of a cooperatively breeding bird: causes and consequences of kin associations. Anim Behav. 2016;122:23–35.

47. Fang G, Jiao HT, Wang MY, Huang PZ, Liu XM, Qi XG, et al. Female demographic changes contribute to the maintenance of social stability within a primate multilevel society. Anim Behav. 2022;192:101–108.

48. Godde S, Humbert L, Côté SD, Réale D, Whitehead H. Correcting for the impact of gregariousness in social network analyses. Anim Behav. 2013;85(3):553–558.

49. Turner JW, Bills PS, Holekamp KE. Ontogenetic change in determinants of social network position in the spotted hyena. Behav Ecol Sociobiol. 2017;72(1):10.

50. van Leeuwen EJC, Cronin KA, Haun DBM. Population-specific social dynamics in chimpanzees. Proceedings of the National Academy of Sciences. 2018;115(45):11393–11400.

51. Hart JDA, Franks DW, Brent L, Weiss MN. BisonR - Bayesian inference of social networks with R; 2022.

52. Ross CT, McElreath R, Redhead D. Modelling human and non-human animal network data in R using STRAND; 2022.

53. Redhead D, McElreath R, Ross CT. Reliable network inference from unreliable data: A tutorial on latent network modeling using STRAND. Psychol Methods. 2023;.

54. Zonana DM, Gee JM, Bridge ES, Breed MD, Doak DF. Assessing Behavioral Associations in a Hybrid Zone through Social Network Analysis: Complex Assortative Behaviors Structure Associations in a Hybrid Quail Population. Am Nat. 2019;193(6):852–865.

55. Farine DR, Firth JA, Aplin LM, Crates RA, Culina A, Garroway CJ, et al. The role of social and ecological processes in structuring animal populations: a case study from automated tracking of wild birds. R Soc Open Sci. 2015;2(4):150057.

56. van Zwet E. A default prior for regression coefficients. Stat Methods Med Res. 2019;28(12):3799–3807.

57. Franks DW, Weiss MN, Silk MJ, Perryman RJY, Croft DP. Calculating effect sizes in animal social network analysis. Methods Ecol Evol. 2020;12(1):33–41.

58. Weiss MN, Franks DW, Brent LJN, Ellis S, Silk MJ, Croft DP. Common datastream permutations of animal social network data are not appropriate for hypothesis testing using regression models. Methods Ecol Evol. 2021;12(2):255–265.

59. Hart JDA, Weiss MN, Brent LJN, Franks DW. Common permutation methods in animal social network analysis do not control for non-independence. Behav Ecol Sociobiol. 2022;76(11):151.

60. Kossinets G. Effects of missing data in social networks. Soc Networks. 2006;28(3):247–268.

61. Huisman M. Imputation of missing network data: Some simple procedures. J Soc Biol Struct. 2009;10(1):1–29.

62. Rubin DB. Inference and missing data. Biometrika. 1976;63(3):581–592.

63. Krause RW, Huisman M, Steglich C, Snijders TAB. Missing Network Data A Comparison of Different Imputation Methods. In: 2018 IEEE/ACM International Conference on Advances in Social Networks Analysis and Mining (ASONAM). IEEE; 2018. p. 159–163.

64. Pavez-Fox MA, Escabi-Ruiz CM, Hart JDA, Negron-Del Valle JE, Phillips D, Montague MJ, et al. Cost and benefits of being social: examining the influence of sociality on faecal parasites in free-ranging rhesus macaques. bioRxiv. 2022;.

65. Hart J. Bayesian Methods for Animal Social Network Analysis. University of Exeter; 2023.

66. Bonnell TR, Vilette C, Henzi SP, Barrett L. Network reaction norms: taking account of network position and plasticity in response to environmental change. Behav Ecol Sociobiol. 2023;77(3):35.

67. Torfs JRR, Stevens JMG, Verspeek J, Laméris DW, Guéry JP, Eens M, et al. Multi-group analysis of grooming network position in a highly social primate. PLoS One. 2023;18(4):e0284361.

68. Stan Development Team. Stan Modeling Language Users Guide and Reference Manual, 2.26.1; 2023; https://mc-stan.org

69. R Core Team. R: A language and environment for statistical computing; 2022.

70. Gabry J, Češnovar R. cmdstanr: R Interface to 'CmdStan'; 2021.

71. Dablander F, Hinne M. Node centrality measures are a poor substitute for causal inference. Sci Rep. 2019;9(1):6846.

72. Youldon DA, Abell J, Briffitt JS, Chama L, Channings MD, Kilundo A, et al. Patch-occupancy survey of elephant (*Loxodonta africana*) surrounding Livingstone, Zambia. Koedoe. 2017;59(1).

73. Douglas-Hamilton I. On the ecology and behaviour of the African elephant: "the elephants of Lake Manyara". University of Oxford; 1972.

74. Moss CJ, Poole JH. Relationships and social structure of African elephants. Primate social relationships: an integrated approach. 1983;315:325.

75. Wittemyer G, Douglas-Hamilton I, Getz WM. The socioecology of elephants: analysis of the processes creating multitiered social structures. Anim Behav. 2005;69(6):1357–1371.

S1 File. The SRI is too trusting of extreme values. Simulation plot to show the difference in potential outputs from SRI versus a Bayesian estimator given true values and the number of times a pair were observed.

S2 File. Standard zero-inflated model. Frequentist two-step model outputs as an alternative option to BISoN.