Analysis of sparse animal social networks

Helen K Mylne[1,2*], Jackie Abell[3], Colin M Beale[1,2,4], Lauren JN Brent[5], Jakob Bro-Jørgensen[6], Kate E Evans[7,8], Jordan DA Hart[5], Dabwiso Sakala[9,10], Twakundine Simpamba[11], David Youldon[10], and Daniel W Franks[1‡].

1 Department of Biology, University of York, York, UK

2 Leverhulme Centre for Anthropocene Biodiversity, University of York, York, UK

3 Centre for Agroecology, Water and Resilience, Coventry University, Coventry, UK

4 York Environmental Sustainability Institute, University of York, York, UK

5 Centre for Research in Animal Behaviour, University of Exeter, Exeter, UK

6 Mammalian Behaviour and Evolution Group, Department of Evolution, Ecology and Behaviour, University of Liverpool, Liverpool, UK

7 Elephants for Africa, London, UK

8 Gothenburg Global Biodiversity Centre, Gothenburg, Sweden

9 Department of Geography, University of Zurich, Zürich, Switzerland

10 African Lion and Environmental Research Trust, Livingstone, Zambia

11 Department of National Parks & Wildlife, Zambia

‡Senior author

* helen.mylne@york.ac.uk

Abstract

Low-density social networks can be common in animal societies, even among species generally considered to be highly social. Social network analysis is commonly used to analyse animal societal structure, but edge weight (strength of association between two individuals) estimation methods designed for dense networks can produce biased measures when applied to low-density networks. Frequentist methods suffer when data availability is low, because they contain an inherent flat prior that will accept any

possible edge weight value, and contain no uncertainty in their output. Bayesian methods can accept alternative priors, so can provide more reliable edge weights that include a measure of uncertainty, but they can only reduce bias when sensible prior values are selected. Currently, neither accounts for zero-inflation, so they produce edge weight estimates biased towards stronger associations than the true social network, which can be seen through diagnostic plots of data quality against output estimate. We address this by adding zero-inflation to the model, and demonstrate the process using group-based data from a population of male African savannah elephants. We show that the Bayesian approach performs better than the frequentist to reduce the bias caused by these problems, though the Bayesian requires careful consideration of the priors. We recommend the use of a Bayesian framework, but with a conditional prior that allows the modelling of zero-inflation. This reflects the fact that edge weight derivation is a two-step process: i) probability of ever interacting, and ii) frequency of interaction for those who do. Additional conditional priors could be added where the biology requires it, for example in a society with strong community structure, such as female elephants in which kin structure would create additional levels of social clustering. Although this approach was inspired by reducing bias observed in sparse networks, it could have value for networks of all densities.

1 Introduction

Social networks underlie all animal social behaviour, and analysing the factors that affect their structure can improve our understanding of how individuals interact. Social network density forms an important part of social structure, defined by the proportion of dyads (all of the possible pairs of individuals) who could potentially associate within a population that do associate. Some animal social networks are dense, characterised by frequent interactions and connections among a high proportion of its members. Others are sparse, with a high proportion of dyads never associating with one another. This difference in network density can influence an individual's rate of social learning [1–3] or their risk of pathogenic infection [4–7], which can cascade to population-level impacts. Differences in network density may be due to inherent differences in gregariousness between species, or inter-population variation in resource availability or predation risk

imposing different limits on the propensity of different populations to aggregate. 13

Studies of animal social networks tend to address gregarious species, typically 14 characterised by dense social networks. However, being gregarious and living in groups 15 is not the same as having strong social preferences: two species can be equally 16 gregarious, but one has individuals associating randomly, while in the other they only 17 associate with a few select partners. As of 2019, 78% of published studies focused on 18 the social structure of either birds (e.g., [8–10]) or group-living mammals [11], such as 19 cetaceans (e.g., [12–15]), primates (e.g., [16–18]) and social carnivores (e.g., [19–21]). 20 Understandably, researchers generally focus on highly gregarious species because their 21 associations are more frequent and they are more likely to be co-located so they are 22 easier to sample for social network studies, and there can be clearer applications for the 23 work concerning conservation (e.g., [19]) or captive animal welfare [22]. However, this 24 means that we know only relatively little about other taxonomic groups and species 25 that may have sparser networks, such as solitary mammals (e.g., [23]) or reptiles 26 (e.g., [24]). Less gregarious animals can still have strong friendships or specific 27 preferences regarding who they should associate with, which might impact their 28 behaviour and survival, including species such as African savannah elephants 29 (*Loxodonta africana*), or mountain and western gorillas (*Gorilla beringei beringei* and 30 *Gorilla gorilla gorilla*) that are generally considered highly social but can show 31 relatively low average rates of social association [25–29]. 32

The two core components of a social network are the individuals within the 33 population, called nodes, and the associations between them, known as the edges. In 34 some networks, we only need to know if a connection exists between a dyad, but the 35 strength of that association (henceforth "edge weight") is not important. In others, we 36 need to know the probability of a dyad being observed together. For many studies, edge 37 weights are the basis of further network structural analyses, making it critical to 38 calculate them reliably. 39

Edge weight estimation methods are sensitive to the density of a network, so may be 40 problematic when networks are sparse, as we will demonstrate in this paper. Methods 41 designed for dense networks may produce biased measures of edge weight, which can 42 lead to further bias in downstream calculations of node centrality or network density, 43 but this is not something that has previously been examined. Similarly, connections 44

between dyads that do not associate need to be modelled as zero-strength edges, but $\quad$ 45
these methods may be either poor at detecting non-association or falsely identify $\quad$ 46
zero-strength edges. In turn, these errors lead to over- and underestimation of edge $\quad$ 47
weight, which will affect all network studies, regardless of whether they are interested in $\quad$ 48
edge presence or weight. The pitfalls of different methods, which we will identify in this $\quad$ 49
paper, should be considered to decide if zero-inflation is likely to influence edge weight $\quad$ 50
estimation and if edges are more likely to be over- or underestimated. $\quad$ 51

Animal social network studies often struggle with limited data quality, exacerbating $\quad$ 52
the difficulties of analysing sparse networks using common methods. Frequentist models $\quad$ 53
— such as the commonly-used simple ratio index (SRI [30]) — are biased towards $\quad$ 54
extreme edge weights (0 and 1, respectively indicating that it is impossible for a pair to $\quad$ 55
ever be together or apart), especially when sample size is low [31, 32], and so accept zero $\quad$ 56
edge weight values too readily. They are therefore liable to underestimating network $\quad$ 57
density by overinflating the number of dyads showing non-association. Furthermore, $\quad$ 58
they contain no measures of uncertainty so cannot highlight the dyads whose edge $\quad$ 59
weight is most likely to have suffered from low sample size. In contrast, Bayesian edge $\quad$ 60
weight models can, as with any Bayesian analysis, accept a non-flat prior, making them $\quad$ 61
more reliable when performed correctly. However, they introduce bias if the analyst $\quad$ 62
chooses to use inappropriate priors designed for the analysis of dense networks. These $\quad$ 63
edge weight priors can imply that non-association is highly implausible, producing $\quad$ 64
posterior distributions biased towards stronger associations. This, in turn, can lead to $\quad$ 65
errors in estimating network density. It is, therefore, inappropriate to use either the SRI $\quad$ 66
or a Bayesian model with a prior designed for a dense network when analysing networks $\quad$ 67
that are not fully connected, as we will later demonstrate. So far, this does not appear $\quad$ 68
to have been an issue in publications, as Bayesian edge weight models are a relatively $\quad$ 69
recent innovation and have only been used for populations with dense networks $\quad$ 70
(e.g., [33–40]). However, as Bayesian models generally become more popular, it is likely $\quad$ 71
that researchers will start to use them to calculate the edge weights of less gregarious $\quad$ 72
species, for which a more robust analysis method needs to be established. We can $\quad$ 73
overcome some of the issues with methods designed for dense networks by having very $\quad$ 74
large quantities of data [32, 41], but this is rarely a plausible solution in animal social $\quad$ 75
network studies, and it does not solve all of the issues associated with edge weight $\quad$ 76

estimation, as we will here discuss.   77

We need a model that reflects how the edge weights are generated in the natural   78
system and incorporates the likelihood of obtaining a true zero, to avoid modelling a   79
connection where one does not exist or biasing our model by supplying inappropriate   80
information regarding the probability of extreme edge weights. All common methods,   81
whether they are frequentist such as SRI (e.g. [10, 21, 24, 42, 43]), half-weight index 82
(HWI) (e.g., [44–47]; or HWIG when correcting for individual gregariousness [48]) or   83
twice-weight index (TWI) (e.g., [49, 50]), or Bayesian such as the Bayesian framework   84
for Inference of Social Networks (BISoN [41, 51]; e.g., [34, 38–40]) or STRAND [52, 53], 85
model a network based on the assumption that the social processes that produce the   86
edge weight contain only a single step: they implicitly assume a fully-connected network 87
where every individual forms a connection with every other, and the only step in   88
deriving the edge weight is determining how strong that connection will be. Edge   89
weight may be correlated with dyad characteristics such as age (e.g., [49]), kinship   90
status (e.g., [25]), phenotypic similarity (e.g., [54]), or anything else that has the 91
potential to induce a preference (or not) for associating, but the decision over how much 92
time to spend together is only a single step.   93

In contrast to these modelling assumptions, a network that is not fully connected   94
has an additional data-generating step: before defining the weight of the connection,   95
there is a precursory step that determines whether or not a dyad connects at all. This 96
could be driven by a choice to avoid another individual [24, 55] or by spatial or temporal 97
barriers that preclude individuals ever having the opportunity to associate (e.g., one 98
member of the population dies before another is born, or no movement corridor exists   99
between two subpopulations so their home ranges cannot overlap). This creates an   100
additional step that defines the level of zero-inflation in the edge weights. Only once it 101
has been defined that a dyad associates can the edge weight be determined. The sparser   102
a network is, the more important it is to capture the first step of the underlying   103
functional process. It is important to note that while this initial step of edge weight 104
calculation has a Bernoulli outcome, the final network density is not a binary result of 105
dense or sparse but a continuum composed of many dyadic edge weights that lie   106
between zero and one. Models that assume a fully connected network become   107
progressively worse as network density declines, but there is no intrinsic cost to always 108

using a method that captures both steps of the underlying social process. ₁₀₉

In this paper we examine two different network modelling approaches for their ₁₁₀ applicability to sparse networks: the SRI [30], which is the approach that has ₁₁₁ historically been the most commonly used to estimate the association between ₁₁₂ individuals, and BISoN [41,51], which is a newer method of using Bayesian models to ₁₁₃ estimate the association index with uncertainty and incorporate more appropriate prior ₁₁₄ information. We demonstrate the application of both of these methods on a sparse ₁₁₅ network of male savannah elephants. We show how Bayesian models overcome many of ₁₁₆ the limitations of the SRI but can suffer problems if supplied with priors that are not ₁₁₇ appropriate for the particular use case. Finally, we introduce a modification of the ₁₁₈ Bayesian framework that follows a two-step data-generating process, which can deal ₁₁₉ with the pitfalls of sparse network analysis. We propose that this modification be ₁₂₀ applied to networks of any density and would benefit future animal network studies. ₁₂₁

2 Challenges of sparse social networks for current common methods

₁₂₂
₁₂₃

2.1 Common network analysis methods ₁₂₄

With gambit-of-the-group data [31], the SRI and Bayesian methods estimate the edge ₁₂₅ weight between individuals based on the total number of observations in which a dyad ₁₂₆ was together versus apart. However, despite using the same input data to estimate the ₁₂₇ same target measure, they suffer from different problems. As with any frequentist ₁₂₈ measure, SRI estimates are drawn based on the assumption that all possible edge weight ₁₂₉ values are equally likely — the equivalent of assuming a flat prior — so can produce any ₁₃₀ output edge weight, even when data are insufficient to support extreme values. To ₁₃₁ combat this, researchers often only include individuals with total sightings exceeding a ₁₃₂ threshold value, but taking a subset of the data creates its own issues, discussed below. ₁₃₃ As a Bayesian alternative, BISoN [41,51] and other Bayesian approaches such as ₁₃₄ STRAND [52,53] can use non-flat priors to inform the model of the prior probability of ₁₃₅ extreme values, so we can set up the model to be more uncertain about extreme edge ₁₃₆ values when the dyad observation count is low. Bayesian approaches, therefore, work ₁₃₇

better with lower sample sizes and uneven sampling than the SRI approach (see ^{138} supporting information: The SRI is too trusting of extreme values). However, as with ^{139} the SRI, a single-step prior is appropriate only for fully connected, or close to fully ^{140} connected, networks. For more sparse networks, we should appropriately modify the ^{141} prior to reflect the biological processes: here, we consider the underlying two-step social ^{142} process that generates connections (or not) and then determines the weight of that ^{143} connection. ^{144}

The SRI is typically calculated as the number of times a pair of individuals were ^{145} observed associating divided by the total number of sightings that either individual was ^{146} observed: ^{147}

$$SRI_{ij} = together_{ij}/sightings_{ij} \qquad (1)$$

$$sightings_{ij} = sightings_i + sightings_j + together_{ij}$$

Where $together_{ij}$ is the number of observations in which individuals i and j were in ^{148} the same group; $sightings_i$ and $sightings_j$ indicate the respective number of ^{149} observations of individuals i and j in the absence of the other; and $sightings_{ij}$ is the ^{150} total number of times they were seen as a whole. As a simple proportion, this measure ^{151} expects extreme values such as 0 and 1 to be as likely as anything in between. It is a ^{152} common misconception that frequentist models are less biased than Bayesian models ^{153} because prior information is not incorporated. This has the opposite effect: a ^{154} frequentist model still has a prior, but it is forced to be flat and uninformative, so it ^{155} biases the results towards extreme values, making outliers just as plausible as the ^{156} average value [56]. The SRI also contains no measure of uncertainty: with this method, ^{157} a value of SRI = 0 occurs when a dyad is never observed in the same group, with no ^{158} greater uncertainty around edge weight estimates for poorly sampled dyads than those ^{159} with many observations, but we should be more certain when more data are available ^{160} (Fig S1.1). ^{161}

These issues are the case regardless of network density, but their potential impact on ^{162} analyses is exacerbated when dealing with extreme edge weight scores. In the real world, ^{163} a lack of connection between individuals has important implications for social structure, ^{164} information and disease transfer, and individual centrality. As the true network density ^{165}

declines, the likelihood of obtaining an edge weight of zero (spurious or correct) 166
increases, and the SRI's lack of uncertainty and treatment of all possible edge weights 167
as equally plausible become a greater issue. We must, therefore, be extremely careful 168
about modelling zero-inflation, but the flat prior and lack of uncertainty seen in the SRI 169
combine to make it very trusting of extreme edge weight values. It is important to note 170
that while we are focussing on the SRI for this paper, other frequentist edge weight 171
measures, including the half-weight and twice-weight indexes, all suffer the same pitfalls. 172
Furthermore, to avoid confusion, it is also worth noting that pre-network permutation 173
tests, which have known problems [57–59], do nothing to deal with the issues faced by 174
frequentist measures. 175

With Bayesian approaches, we still estimate the edge weight based on the proportion 176
of total dyad sightings in which they were together, but now as a model parameter 177
rather than a simple ratio calculation, for example: 178

$$together_{ij} \sim binomial(sightings_{ij}, edgeweight_{ij}) \tag{2}$$
$$logit(edgeweight_{ij}) \sim normal(\mu, \sigma)$$

Where $together_{ij}$ and $sightings_{ij}$ are the respective number of observations of a 179
dyad in the same group and in total, as above; and μ and σ are the prior mean and 180
standard deviation. Estimates of edge weight are now described by posterior 181
distributions, which narrow as $sightings_{ij}$ increases, capturing uncertainty in the edge 182
weight. Incorporating a user-defined prior can limit misleading outputs when the sample 183
sizes are small, as the range of likely edge weight values is reduced. Through this use of 184
sensible priors and quantification of uncertainty, Bayesian approaches are inherently 185
more capable of dealing with the small sample sizes that are common in studies of 186
animal social systems. While BISoN is not the only option available for the Bayesian 187
estimation of edge weight (see also STRAND [52,53]), we will focus on it in this paper. 188

2.2 SRI requires deletion of data 189

With a flat prior and no implicit uncertainty measure, limited data availability about a 190
dyad can produce unreliable SRI values, so researchers will usually set an arbitrary 191
minimum threshold number of observations per individual to include that node in the 192

analysis (e.g., [25, 29, 42, 44, 46, 49]). While this does prevent inaccurate edge weights ⁠₁₉₃ created by the most poorly sampled dyads, it simultaneously introduces new ⁠₁₉₄ issues [60, 61] that are often overlooked. Firstly, data are often hard-won and already ⁠₁₉₅ limited, yet this practice requires the discarding of data, sometimes in large quantities. ⁠₁₉₆ Secondly, data removal may not be completely-at-random [62], which is known to ⁠₁₉₇ introduce bias [61, 63]. For example, individuals that are less neophobic, more ⁠₁₉₈ philopatric, or possess distinctive identifying features are more likely to be included in ⁠₁₉₉ the analysis, because the probability of their having repeat observations is greater. If a ⁠₂₀₀ study next performs a regression analysis on calculated network measures, this ⁠₂₀₁ non-random data deletion can directly bias the results, especially if these factors are ⁠₂₀₂ linked to the exposure or outcome variables in question. Thirdly, the specific value of ⁠₂₀₃ these observation thresholds often varies between studies, as the actual choice of how ⁠₂₀₄ many sightings is considered "enough" usually depends on the number of individuals ⁠₂₀₅ that will remain after filtering out those poorly sampled. For example, Chiyo *et al.* [25] ⁠₂₀₆ included all male elephants with at least 15 observations in their three-year study ⁠₂₀₇ period, while Murphy *et al.* Murphy2020-yk used an inclusion threshold of just five ⁠₂₀₈ observations per four year sampling period. Therefore, comparing the results of different ⁠₂₀₉ studies and identifying how many sightings are truly "enough" is difficult. Even with ⁠₂₁₀ several sightings per individual, scores of 0 and 1 can be overinflated relative to their ⁠₂₁₁ true underlying network, because there are only a certain number of possible edge ⁠₂₁₂ weight values when sighting counts are low (Fig S1.1). ⁠₂₁₃

In contrast, employing a Bayesian framework prevents the requirement for data ⁠₂₁₄ inclusion thresholds, which improves data quality and reduces biases caused by the ⁠₂₁₅ possibility of data missing not-completely-at-random. ⁠₂₁₆

2.3 Bayesian models facilitate the use of more appropriate prior ⁠₂₁₇ distributions ⁠₂₁₈

A model prior should always be carefully considered, which is not possible under the ⁠₂₁₉ frequentist framework. Given how few papers have thus far used a Bayesian method for ⁠₂₂₀ edge weight estimation, most social network analyses are based on flat frequentist priors, ⁠₂₂₁ which are generally inappropriate (if nothing is known at all of the population in ⁠₂₂₂

question, then a flat prior may still be a reasonable choice, though this would be ²²³
extremely unusual). The *bisonR* [51] package for running BISoN models supplies very ²²⁴
broad default priors, but the package authors strongly recommend tailoring them to ²²⁵
each specific scenario. Most social networks contain at least some true zeros. Any ²²⁶
model, frequentist or Bayesian, that uses a flat or very broad prior will struggle to ²²⁷
model these zero values because they are modelling only the second step of what is ²²⁸
inherently a two-step process. ²²⁹

Unlike the SRI, with BISoN we can change our model structure to incorporate two ²³⁰
steps — first to determine if a dyad will associate at all, and second to define the ²³¹
strength of that connection — if we supply it with an appropriate prior, which package ²³²
default priors are not. Here, we have presented a prior allowing BISoN to model both ²³³
steps without changing the social network analysis workflow. While we could instead ²³⁴
use a mixture model with zero-inflation to model a two-step process, using either a ²³⁵
frequentist (see supporting information: Standard zero-inflated model) or Bayesian ²³⁶
framework, we can keep the overall method relatively simple by changing only the prior ²³⁷
structure. All network models should include some option to account for zero-inflation, ²³⁸
and selecting the best method for measuring edge weights requires careful consideration: ²³⁹
Bayesian techniques require integration of current understanding of the system, and a ²⁴⁰
prior predictive check, which can lead to biases if carried out using inappropriate priors, ²⁴¹
but their ability to estimate uncertainty and accept non-flat priors makes them more ²⁴²
appropriate than frequentist methods. ²⁴³

2.4 Prior selection in a Bayesian edge weight model ²⁴⁴

Bayesian methods have been successfully applied to dense networks, using both ²⁴⁵
simulated and empirical data [34, 64, 65], and their use in primate research has been ²⁴⁶
discussed in subsequent publications [66, 67]. However, the best way to use them for ²⁴⁷
sparse networks has not yet been evaluated. Prior choices require careful consideration ²⁴⁸
for any analysis, not just social networks. Fig 1a shows an improper uniform prior, in ²⁴⁹
which all edge weights are considered equally likely before exposure to the data. By ²⁵⁰
relaxing the assumption that all edge weights are equally probable, we can make the ²⁵¹
prior more informative. When we demonstrate the use of BISoN on an example dataset, ²⁵²

we first show the outcome of this uniform prior, followed by the default prior supplied in the $bisonR$ package [51]: a wide symmetrical prior (logit($edge$ $weight_{ij}) \sim normal(0, 2.5)$) that presumes very little about the average edge weight (Fig 1b). Third, we show the result of using a strong asymmetrical prior ($edge$ $weight_{ij} \sim beta(1, 5)$) that indicates weaker edges are the most likely but still allows for some strong associations (Fig 1c). How we define the prior depends on our knowledge and assumptions of the system: the sparser the network, the more right-skewed the prior distribution should be.

Fig 1. Prior assumptions for single-step BISoN models. Prior assumptions for a) SRI or a BISoN model with a uniform prior, b) a BISoN model with a weak, symmetric prior (logit normal, $\mu = 0$, $\sigma = 2.5$) and c) a BISoN model with a strong asymmetric prior (beta, $a = 1, b = 5$).

Biases can appear in the edge posterior when the prior distribution does not reflect the underlying social process: knowing that the network is of a lower density can allow us to adapt our prior to a more suitable distribution, and the assumptions we can make based on that knowledge are critical to dealing with poorly sampled dyads. When the prior average is a long way from that of most dyads, it induces a systematic shift in the direction the data pushes the posterior from the prior. In a sparse network, many of the dyads will never associate, so a prior designed for a dense network will usually be shifted towards peaking at zero by the data. The data for poorly sampled dyads will, by definition, not be able to shift the posterior as far from the prior as the data from the well-sampled dyads, so if they all shift in the same direction, the level of sampling could potentially have a greater effect on the final edge weight than the actual raw data values. When this shift is consistently towards zero, poorly sampled dyads, on average, receive edge weight distributions with higher average values than well-sampled dyads. It is difficult to imagine a scenario in which the number of observations of a dyad would be strongly dependent on the strength of their social bond or vice versa, therefore this correlation highlights a problem with the edge weight data. We show this happening in the example analysis. Again, without well-sampled data, the choice of prior (including the SRI's uniform prior) becomes critical to the success of the edge weight estimation and all subsequent analyses.

3 Introducing a two-step prior for sparse networks

By varying the prior distribution, Bayesian methods can be tailored to a sparser or
denser network. The modification that we propose simply extends this ability: to alter
the BISoN prior structure to reflect a two-step social process producing the edge
weights, in which step one defines the extent of zero-inflation and step two the weight of
the non-zero social bonds. We structure our model to use two separate edge weight
priors, selecting which to use depending on whether a dyad has ever been observed
together (Fig 2), such as:

$$together_{ij} \sim binomial(sightings_{ij}, edgeweight_{ij}) \tag{3}$$

$$edgeweight_{ij} \sim beta(a, b)$$

$$\text{If } together_{ij} = 0: \ a < 1 \text{ and } b = \text{large}$$

$$\text{If } together_{ij} > 0: \ a = 1 \text{ and } b = \text{small}$$

Where again $together_{ij}$ refers to the number of times a pair was observed in the same
group, and $sightings_{ij}$ is the number of times they were observed in total. This is not a
major change to the method itself, as BISoN has always been capable of using a
conditional prior. The change is that by capturing both parts of the underlying social
process in a conditional prior, we only need to change one part of our overall method.

Fig 2. Conditional edge weight prior for a BISoN model of a sparse network.
The blue line indicates the lower prior that will be selected when a dyad has never been
observed in the same group (in this case $beta(a = 0.7, b = 10)$). The purple line shows
the upper prior choice for when a dyad has been observed grouping together on at least
one occasion (in this case $beta(a = 1, b = 5)$). While in this model we have chosen to
still have this prior peak at zero, this reflects that we still expect very low edge weights
amongst those that do associate, and the data immediately overwhelm the possibility of
zero being an option. Note that the upper prior is the same as the strongly right-skewed
prior shown above.

The new prior is simple to apply: if a dyad is never observed in the same social
group, then the model will use a prior that increases the probability of an edge weight
of zero; if a dyad has been observed interacting on at least one occasion, it will receive
an edge weight calculated using a prior with less or no zero-inflation. This, therefore,
allows the model to identify the most appropriate prior and incorporate the probability
of non-association as an initial step before identifying the best edge weight distribution

per dyad. As before, poorly sampled dyads will return an edge weight more similar to their respective prior, while well-sampled dyads will shift further. The model becomes less uncertain about a zero edge as the total sighting count increases: the uncertainty in these values indicates that zero is the most likely edge weight but still allows stronger relationships as a possibility when sampling is poor. In contrast, for a dyad observed together on at least one occasion, we use a wider prior that allows higher edge weights to be observed, again becoming more confident in the results as total sampling increases. The most logical step here may appear to be to make our regular prior more right-skewed so it is closer to zero. However, if we were to use only an extremely right-skewed prior, such as the lower prior of this combination, to encapsulate the zero-inflation and the true associations, a prohibitively large amount of association data would be required for any truly strong edge weights to register.

The values a researcher might choose for the prior depend on their study species and expected edge weights. In the following example, we have opted to have the prior for associated dyads peaking at zero, but this is immediately overwhelmed by the data. This produces edge weight values that are still far more likely to be close to zero than to one, based on the evidence that male elephants do not appear to form strong relationships with one another [25, 26, 29]. If we were expecting to see a strongly bimodal distribution in the final edge weights, with some dyads never associating and others showing consistent association, we would shift our upper prior further from zero.

4   Example dataset

For this example, we used data on sparse networks based on associations between male elephants of the Mosi-Oa-Tunya National Park (MOTNP), Zambia, collected from May 2016 until December 2017 (all permissions for data collection were supplied by Department of National Parks and Wildlife, Zambia: permit number TJ/DNPW/101/13/18). No additional ethics permission was required alongside the research permit, but to avoid causing any stress to the elephants, we monitored them from a vehicle from a safe distance, and kept all noise to a minimum while in the National Park.

This population is relatively small, totalling approximately 500 individuals, with

similar numbers of males and females. The park is well connected to large areas of ³³²
suitable habitat, and the elephants are free-ranging, so individuals may leave the area ³³³
for long periods. Data were collected during daily drives through the park, marking ³³⁴
down the identities of all individuals. From these observations, we created 504 days of ³³⁵
gambit-of-the-group [31] data comprising 213 males sighted in 481 groups. Finally, we ³³⁶
converted this to a data frame containing every possible dyad with counts of the total ³³⁷
number of times that dyad had been observed and how many of those occasions they ³³⁸
were in the same group. Data analyses were run by calling Stan (version 2.26.1 [68]) ³³⁹
from R (version 4.2.1 [69]) using the *cmdstanr* [70] package. ³⁴⁰

5   Results from empirical data ³⁴¹

5.1   Simple Ratio Index (SRI) ³⁴²

When we calculate the SRI values for this population (Fig 3a), we get an extremely high ³⁴³
proportion of zeros (17385, equating to 77.00% of dyads). While a much lower number ³⁴⁴
of dyads receive an SRI of one (12 dyads, equating to just 0.05% of the total ³⁴⁵
population), the likelihood of these apparent permanent alliances would be exceedingly ³⁴⁶
small given the population average. ³⁴⁷

Fig 3. Edge weights for MOTNP elephants using the SRI. a) The vast majority
of dyads have an SRI score (green points) of zero, indicating that the output is certain
that these pairs never associate with one another, regardless of the number of sightings.
b) This translates to a large split between the edge weights calculated for poorly
sampled dyads that have (purple line) or have not (blue line) ever been observed
grouping together. We see a very strong effect of dyad sighting count on the edge weight
for dyads observed together on at least one occasion, until we reach around 12 sightings
per dyad.

Given this, the standard next step with the SRI would be to set a threshold for the ³⁴⁸
minimum number of sightings per node required to be included in the analysis. If we set ³⁴⁹
this threshold to five sightings per node, the proportion of dyads with an SRI score of ³⁵⁰
zero shifts down to 46.13%, and all dyads scoring a 100% association rate are removed. ³⁵¹
A threshold of 10 sightings per individual drops the 0% association rate down to 26.25%, ³⁵²
showing high instability in the SRI outputs. This is highlighted by Fig 3b, which shows ³⁵³
the association between the total number of observations per dyad, and the calculated ³⁵⁴
edge weight for that dyad, stemming from the extreme number of zeros calculated by ³⁵⁵

the SRI. The overall results would strongly depend on the threshold value we choose, which, as previously stated, is an arbitrary choice.

We could potentially use Fig 3b to determine a less arbitrary inclusion threshold by selecting the point at which the patterns in the edge weights weaken. However, this example shows that this is somewhere between 10 and 20 sightings per dyad. This would require removing at least all individuals with fewer than 10 observations. In this case, that would take our total sample size down from 213 to just 74 elephants. To remove 65% of the sample could severely bias our results if the remaining portion is not entirely representative of the total population. Therefore, we cannot rely on objective means to determine the threshold level.

5.2 Bayesian model with a single-step prior

When we use the BISoN framework with a single-step prior, we can see a limited effect of prior shape on the posterior distribution, but the inclusion of zero as a possible edge value is very important. When using a fully flat or highly right-skewed prior, in which zero is a possible option, we see very similar posterior distributions (Fig 4a and c), whereas for a normal prior peaking at 0.5 for which extreme edge weights are unlikely (and actual values of 0 and 1 are impossible) we see an overall much broader and less skewed posterior (Fig 4b).

Fig 4. Edge weights for MOTNP elephants using BISoN models with single-step priors. Results using BISoN with uniform (left), default (middle) and right-skewed (right) priors (insets in top row show the respective prior for the column). a-c) The top row shows the posterior distribution for 100 randomly sampled dyads, with green curves indicating pairs that have never been observed in the same group and purple curves showing dyads that were observed in the same group on at least one occasion. The right-skewed and uniform priors produce posterior distributions that initially appear very similar because they still allow an edge weight of zero to be a plausible draw, whereas the default prior draws all values together. d-f) The bottom row shows the effect of total sighting count per dyad on their median (green points) and full (yellow points) posterior distribution. The blue line indicates dyads never observed together, while the purple line shows dyads observed together at least once. As the prior becomes increasingly right skewed, the effect of sighting counts on median edge weight declines.

We now also have a situation in which there is a negative correlation between mean edge weight and total observations per dyad at low sighting counts, especially when using very broad priors. Looking at Figs 4d and 4e, we can see that poorly sampled

dyads appear to show higher edge weights than well-sampled ones (an effect the authors have also seen in social network data for macaques). Again, this is most severe for the logit normal prior, which draws all of the values towards the middle instead of creating variation between dyads. The effect is also stronger for the flat prior than for the right-skewed, for which the trend is almost entirely removed because those dyads with the fewest sightings do not have sufficient data to discredit high edge weight model draws, so the posterior is very wide, with high median values. Similar reasoning also explains the differences between BISoN with a flat prior and the SRI: with the SRI, low sighting counts largely score exactly zero, whereas with BISoN low sighting counts are given a very wide posterior (shown in yellow) so their median (green) is substantially above zero.

Based on these graphs, the best solution would appear to be to use BISoN with a strongly right-skewed prior, which can model the edge weights without creating strong patterns of social association based on data quality. However, this prior may not allow sufficient movement towards a higher edge value for dyads showing a strong social bond. Therefore, we need a prior that better reflects the underlying social process, such that it allows for some dyads to attain higher edge weights, but without increasing the prior average and causing poorly sampled dyads to consistently obtain a higher edge weight than well-sampled ones. Our modification to BISoN to use a prior that reflects the two-step underlying social process facilitates edge weight estimation in a manner robust to zero inflation and which allows stronger edges to be estimated.

5.3 Bayesian model with a two-step prior

When using the two-step prior, the posterior distributions of the edge weights (Fig 5a) appear very similar to those with single-step priors, though with greater confidence in some of the weakest edges, as we would expect. The median edge weight is associated much more weakly with the total number of sightings than with the broader priors (Fig 5b), even at low dyad sighting counts. While the shape of the curve for pairs seen grouping together is very similar to the single-step right-skewed prior, the overall effect of low sighting count has been removed by the model's increased ability to obtain median estimates very close to zero. We now have edge estimates that are more reliable

than the previous BISoN models, as they use a prior that allows the modelling of zero-inflation.

Fig 5. Edge weights for MOTNP elephants using a BISoN model with a conditional prior. a) posterior density lines for every dyad (blue indicates density lines drawn for pairs never observed in the same group, purple for those seen together on at least one occasion); and b) edge weight as a response to number of sightings per dyad (full posterior distribution in yellow, median per dyad in green). While some effect of sighting count on the median edge weight for individual dyads remains, the impact is much reduced compared with the previous BISoN plots, and it lacks the values of 100% association observed with the SRI.

5.4 Using downstream network measures to assess edge weight quality

In general, we should be able to see directly in our edge weight values if there is a problem in our estimation method. Sometimes however, the impacts of the method are masked until subsequent analyses of the outputs. For example, in an analysis to identify the effects of individual trait values on network centrality, we may find that our centrality values contain spurious patterns, such as in the number of sightings or number of different partners that an individual is observed with compared to network centrality. Here, we will show that using an inappropriate single-step prior for edge weight can severely impact downstream analyses of network structure. We explore this using eigenvector centrality, a measure of how connected an individual is to other well-connected individuals, as it is commonly used and is the centrality measure most robust to causal analysis [71]. As with the edge weights, we can plot measures of centrality calculated per model against a) the number of other individuals in the population with whom they were never observed associating, and b) their individual sighting count, and look for any unexpected trends that may indicate problems with the outputs.

When we plot the SRI-based eigenvector data against the number of non-associated elephants, we see a negative correlation (Fig 6a), exactly as we would expect. If, however, we compare the SRI-based centrality to the number of sightings per individual, we see a positive correlation for poorly sampled dyads (Fig 6b), which indicates that there is a problem with the estimation. This relationship is created by the extreme edge

values caused by limited numbers of samples, especially for elephants observed fewer ⁠431
than 10 times. This could be used as an objective method of determining an inclusion ⁠432
threshold, but as before, this would require the discarding of more than half the data. ⁠433
We obtain similar trends when calculating the edge weight using a zero-inflated mixture ⁠434
model (Fig S2.2). ⁠435

Fig 6. Eigenvector centrality estimates based on the SRI edge weight calculations. a) There is a negative relationship between SRI-based centrality and the total number of potential partners with whom a focal was never observed associating, which is as expected. However, b) indicates an issue with the SRI-based centrality in that there is a strongly positive effect of the number of observations per individual on their social position at low observation counts.

We see something very different when we look at the eigenvector centrality plots ⁠436
based on the single-step BISoN models. Startlingly, there is an apparent reversal of the ⁠437
effect of the number of non-associated individuals in the population, such that elephants ⁠438
observed with the most partners receive the lowest centrality scores, while those ⁠439
observed with the fewest partners receive the highest (Fig 7a-c). This is, of course, ⁠440
nonsense. Eigenvector centrality as a measure combines the number of association ⁠441
partners an individual has with the number of association partners their partners have. ⁠442
The most central elephants therefore cannot be those with the fewest association ⁠443
partners. The edge weight estimation process must have an underlying error to create ⁠444
this trend. ⁠445

Fig 7. Eigenvector centrality estimates based on BISoN edge weight models with single-step priors. Priors per column are shown as insets on the top row: left to right uniform, default and right-skewed. In all graphs, yellow points indicate the uncertainty around the median, displayed in blue. a-c) The top row indicates the impact of number of dyad partners with whom an individual was never observed associating, where we see a spurious positive effect. The more right-skewed prior suffers more weakly from this than the uniform-based or default-based centrality, but still sufficiently to create serious cause for concern. There is no biologically or statistically meaningful explanation for this to be a correct trend. The pattern is explained by d-f) in which the elephants with the most observations occupy the social positions of lowest centrality.

This seemingly impossible pattern occurs because of an underlying trend in the ⁠446
number of observations per elephant (Fig 7d-f). Individuals with fewer overall ⁠447
observations contribute fewer sightings to dyad observations. When the edge weight ⁠448
prior is designed for dense networks, the posterior for most dyads is shifted closer to ⁠449
zero from the prior. As the data for poorly sampled dyads are less influential on the ⁠450

distance the posterior can move, dyads with fewer observations consistently appear to ₄₅₁ have a stronger edge weight than well-sampled dyads. When this translates through to ₄₅₂ individual effects, and poorly sampled individuals end up in more poorly sampled dyads, ₄₅₃ it is exaggerated so individuals observed rarely have the strongest average edge weights ₄₅₄ and, therefore, the highest eigenvector centrality. ₄₅₅

Finally, we can now consider eigenvector centrality calculated from the edge weights ₄₅₆ produced using BISoN with a two-step prior and see that we successfully avoid all ₄₅₇ downfalls from the other methods. Changing to a conditional prior has successfully ₄₅₈ reversed the spurious correlation between number of associates and node centrality ₄₅₉ (Fig 8a) by removing the effect of sighting count on eigenvector (Fig 8b). We have ₄₆₀ avoided creating estimates biased by the data quality, so any subsequent analyses using ₄₆₁ these data will be more reliable than those calculated using an improper prior while also ₄₆₂ avoiding the need to remove poorly sampled dyads as we would with the SRI. ₄₆₃

Fig 8. Eigenvector centrality estimates based on BISoN edge weight model with a two-step prior. a) Using our modified prior (inset), to reflect a two-step social process, we have successfully restored the negative trend in node centrality such that those seen with the most different dyad partners on average have the highest centrality, and those with the fewest have the lowest. b) Individual sighting count now has extremely limited effect on eigenvector centrality.

This quick sanity check of the eigenvector against the number of associates can ₄₆₄ instantly highlight if there is a problem with the edge weights, and show how sighting ₄₆₅ count could affect centrality. Our modification does not entirely eliminate this, and care ₄₆₆ should be taken to look for any anomalies. For example, in the MOTNP data, there is a ₄₆₇ single sighting where 89 elephants crossed the river and came up into the surrounding ₄₆₈ mopane forest together, of which 31 were males with known identities. For two males, ₄₆₉ this was their only observation. Therefore, these two individuals were "always" seen ₄₇₀ with one another and are in the "seen grouping together at least once" group for 30 ₄₇₁ dyad pairs. This means that, despite having very wide uncertainty margins for all dyads ₄₇₂ due to low total counts, they will still have an above-average median edge weight due to ₄₇₃ the nature of their single observation. When carried forward to a centrality analysis, ₄₇₄ these elephants will display a slightly elevated eigenvector centrality. Using the modified ₄₇₅ BISoN framework means this elevation is much lower than when using an unconditional ₄₇₆ prior or the SRI, and the number of individuals for which this could be an issue is ₄₇₇

reduced, limiting the impact on the regression. However, we still need to be aware of   478
the potential for data artefacts that an occurrence such as this may leave. It is worth   479
noting that while none of these methods is perfect, by using a method that incorporates   480
current knowledge, quantifies uncertainty, and does not induce issues in downstream   481
analytical data frames, we can rest assured that we are using the best possible option.   482

6   Discussion   483

We have shown that current animal social network analysis methods do not cope well   484
with sparse network data. Frequentist measures such as the SRI suffer from an innate   485
and unavoidable reliance on a flat prior, which overestimates the confidence that can be   486
placed on extreme edge weight values. Combine this prior with an inability to   487
incorporate uncertainty, and researchers must rely on setting arbitrary inclusion   488
thresholds when analysing poorly sampled dyads, leading to data loss and potentially   489
biased results. While Bayesian models can account for these problems in dense   490
networks, they can still produce edge weights overly influenced by the frequency of   491
sightings per dyad when inappropriate priors are applied to sparse networks. By   492
modifying the BISoN framework to use a conditional prior, reflecting a two-step   493
underlying social process, we can create a model where the edge weights are more   494
reliable than those from the SRI, a zero-inflated mixture model, or an uninformed   495
Bayesian model. As such, they are no longer overly dependent on data quality, which is   496
commonly poor in animal social network studies.   497

When checking our models, we used several diagnostics to compare output quality   498
and determine if our BISoN modification improved the current methods. The first   499
question is whether a result fits with what we know of the population: the SRI produces   500
far too many dyads with zero probability of ever associating, even when we set a high   501
inclusion threshold. At just 66 km^2 [72], the MOTNP is not sufficiently large to   502
preclude the interaction of elephants by simply never meeting, with a bull easily able to   503
cross it in a single day. It is, therefore, extremely unlikely that this many dyads are   504
truly never together, as there would not be enough space for them all to move without   505
encountering other elephants. This allows us to make two informed assumptions: i) the   506
proportion of zeros produced by the SRI is overinflated; and ii) that well-sampled dyads   507

that are never recorded associating have elected not to do so, rather than happening to   508
never encounter one another. When using BISoN, the conditional prior can help to   509
alleviate the effect of sighting count on median edge weight by using the broader prior   510
with a higher average only when we have evidence that the dyad has elected to   511
sometimes associate. While this solution does not entirely remove the impact of limited   512
sightings on edge weight, it does minimise it compared to a single-step prior.   513

At first glance at the edge weights alone, it seems that the conditional prior is not   514
particularly advantageous over the right-skewed single-step prior: the basic outputs for   515
the edge weight look fairly similar. However, we have modelled the zero-inflation, which,   516
while it appears to have only had a limited effect on the edge weights themselves, has a   517
very noticeable and important impact on our subsequent analyses. The node centrality   518
values produced can act as a suitable sanity check. While there appears to be little   519
difference between the right-skewed BISoN outputs and the two-step BISoN outputs   520
when we consider only the edge weights, the eigenvector centrality from a single-step   521
prior in a sparse network creates an impossible situation in which individuals with more   522
partners receive a lower centrality score. This trend is corrected by the use of a two-step   523
prior. We can use such checks to ensure our outputs make sense in downstream analyses.   524

Selecting the two distributions remains important for successful edge weight   525
estimation when applying a two-step prior to a sparse network model. Making the   526
zero-centred prior too strong will not allow enough variation when data are limited, but   527
if too weak some of the possible edge weights may still present as too high and create   528
trends in edge weight and centrality based on sighting counts. Similarly, if the   529
probability mass for the non-zero prior is not shifted far enough from zero, any stronger   530
relationships will be missed; shift it too much, and the same trends of increased edge   531
weight with reduced sightings will be observed within the group who have been   532
observed together. In this instance, we have opted for a large overlap between the priors   533
such that dyads only seen once or twice will receive relatively similar posterior   534
distributions regardless of which prior they are assigned to. By giving both priors a   535
peak at zero, a dyad observed just once but who were together at that time will still   536
have a low average edge weight (though the zero-peak is always immediately overcome   537
by the data), matching our assumptions of a sparse network. This particular conditional   538
prior is the one that works for our specific elephant population and is not a silver bullet   539

prior that will work for all network analyses. ₅₄₀

Every author should identify their own priors to create a set that works for their ₅₄₁ own population, which does not have to require that both priors peak at zero: unlike ₅₄₂ with the male elephants, a network can be sparse but still contain some strong ₅₄₃ associations, in which case an upper prior that allows higher edge weights may be ₅₄₄ preferable. For example, if we were to repeat this analysis with the female elephants of ₅₄₅ MOTNP, we would expect a more bimodal distribution in our output because of the ₅₄₆ fission-fusion structure of female elephant society [73–75]. In this case, we could shift ₅₄₇ our non-zero prior to be more distinct from the lower, producing a bimodal distribution ₅₄₈ of edge weights, indicating strong social bonds within certain dyads. It could even be ₅₄₉ taken a step further to add more prior distributions as necessary. Again, using the ₅₅₀ example of the female elephants, we may, in this scenario, want the prior to reflect a ₅₅₁ three-step social process: a zero-centred prior for those never seen in the same social ₅₅₂ group, a right-skewed or more central prior for those who have been seen grouping ₅₅₃ together but who are unrelated; and a left-skewed prior for related females. Hierarchical ₅₅₄ networks in which we see kinship groups within subgroups, which themselves join ₅₅₅ together to form larger association groups are very common in animal societies. This ₅₅₆ method could be easily expanded to allow for the use of different priors depending on ₅₅₇ which level of association a dyad is expected to fall into. ₅₅₈

It should be noted that the assessment of the edge weight is rarely the end of the ₅₅₉ analysis. Generally, we calculate the edge weights so that we can later assess trends in ₅₆₀ node-, dyad- or network-level characteristics. Therefore, the edge weight calculation ₅₆₁ only creates the data ready for a future regression model. There is no issue with ₅₆₂ checking the edge weights produced — just as we would check for errors in any other ₅₆₃ raw data — and running the model again using an improved prior if there is a problem. ₅₆₄ If edge weight outputs are dependent on the dyad sighting count, or there are issues ₅₆₅ with downstream calculations such as the centrality, there is no issue with adjusting the ₅₆₆ prior, so long as it still reflects our original understanding of the population. ₅₆₇

In conclusion, the shape of the prior is, as with all analyses, the key to estimating ₅₆₈ edge weight, and using a prior structure that reflects the underlying social process is ₅₆₉ especially important for sparse networks, though the concept applies to dense networks ₅₇₀ too. With a frequentist model, this means considering the implications of a flat prior ₅₇₁

and including in our discussion any impacts this may have on the results. For a $\quad$ 572
Bayesian model, it means tailoring the prior to our individual situation rather than $\quad$ 573
using the broad default priors supplied by statistical packages, which will usually allow $\quad$ 574
too much probability density over highly improbable edge weight values. In the case of $\quad$ 575
sparse networks, whether the model is Bayesian or frequentist, we need to model the $\quad$ 576
zero-inflation induced by the first step of the underlying social process. While a $\quad$ 577
zero-inflated mixture model can do this better than the SRI, we still recommend that a $\quad$ 578
Bayesian model with a conditional prior is a better option. This means allowing edge $\quad$ 579
weights to sample down to zero using a lower prior and the possibility for stronger $\quad$ 580
relationships with an upper prior. When investigating denser networks, our prior $\quad$ 581
distributions may differ, but we can still consider the underlying social process as $\quad$ 582
containing more than one step, even if the reduced proportion of zero relationships $\quad$ 583
masks the effect. We can therefore use a conditional prior in a Bayesian framework to $\quad$ 584
identify greater nuance in social network data, for networks of any density. $\quad$ 585

7 Acknowledgements $\quad$ 586

We would like to thank the African Lion and Environmental Research Trust (ALERT) $\quad$ 587
for providing the example dataset and the Zambian Department of National Parks and $\quad$ 588
Wildlife (DNPW) for allowing its collection and analysis. Analyses for this paper were $\quad$ 589
run on the Viking cluster, a high-performance computing facility provided by the $\quad$ 590
University of York. We are grateful for computational support from the University of $\quad$ 591
York, IT Services and the Research IT team. $\quad$ 592

References

1. Singh J. Collaborative Networks as Determinants of Knowledge Diffusion
   Patterns. Manage Sci. 2005;51(5):756–770.

2. Franz M, Nunn CL. Network-based diffusion analysis: a new method for
   detecting social learning. Proc Biol Sci. 2009;276(1663):1829–1836.

3. Hobaiter C, Poisot T, Zuberbühler K, Hoppitt W, Gruber T. Social network analysis shows direct evidence for social transmission of tool use in wild chimpanzees. PLoS Biol. 2014;12(9):e1001960.

4. Kretzschmar M, Morris M. Measures of concurrency in networks and the spread of infectious disease. Math Biosci. 1996;133(2):165–195.

5. Cross PC, Lloyd-Smith JO, Bowers JA, Hay CT, Hofmeyr M, Getz WM. Integrating association data and disease dynamics in a social ungulate: bovine tuberculosis in African buffalo in the Kruger National Park. Ann Zool Fennici. 2004;41(6):879–892.

6. Naug D. Structure of the social network and its influence on transmission dynamics in a honeybee colony. Behav Ecol Sociobiol. 2008;62(11):1719–1725.

7. Silk MJ, Croft DP, Delahay RJ, Hodgson DJ, Boots M, Weber N, et al. Using Social Network Measures in Wildlife Disease Ecology, Epidemiology, and Management. Bioscience. 2017;67(3):245–257.

8. Silk MJ, Croft DP, Tregenza T, Bearhop S. The importance of fission-fusion social group dynamics in birds. Ibis. 2014;156(4):701–715.

9. Farine DR, Aplin LM, Sheldon BC, Hoppitt W. Interspecific social networks promote information transmission in wild songbirds. Proc Biol Sci. 2015;282(1803):20142804.

10. Beck KB, Farine DR, Firth JA, Sheldon BC. Variation in local population size predicts social network structure in wild songbirds. J Anim Ecol. 2023;92(12):2348–2362.

11. Webber QMR, Vander Wal E. Trends and perspectives on the use of animal social network analysis in behavioural ecology: a bibliometric approach. Anim Behav. 2019;149:77–87.

12. Foster EA, Franks DW, Morrell LJ, Balcomb KC, Parsons KM, van Ginneken A, et al. Social network correlates of food availability in an endangered population of killer whales, *Orcinus orca*. Anim Behav. 2012;83(3):731–736.

13. Weiss M. Developing methods and applications for the analysis of cetacean social networks. University of Exeter; 2021.

14. Gerber L, Connor RC, Allen SJ, Horlacher K, King SL, Sherwin WB, et al. Social integration influences fitness in allied male dolphins. Curr Biol. 2022;32(7):1664–1669.e3.

15. Marfurt SM, Allen SJ, Bizzozzero MR, Willems EP, King SL, Connor RC, et al. Association patterns and community structure among female bottlenose dolphins: environmental, genetic and cultural factors. Mamm Biol. 2022;102(4):1373–1387.

16. Kasper C, Voelkl B. A social network analysis of primate groups. Primates. 2009;50(4):343–356.

17. Brent LJN, Lehmann J, Ramos-Fernández G. Social network analysis in the study of nonhuman primates: a historical perspective. Am J Primatol. 2011;73(8):720–730.

18. Sueur C, Jacobs A, Amblard F, Petit O, King AJ. How can social network analysis improve the study of primate behavior? Am J Primatol. 2011;73(8):703–719.

19. Abell J, Kirzinger MWB, Gordon Y, Kirk J, Kokeŝ R, Lynas K, et al. A social network analysis of social cohesion in a constructed pride: implications for ex situ reintroduction of the African lion (*Panthera leo*). PLoS One. 2013;8(12):e82541.

20. Dorning J, Harris S. Individual and seasonal variation in contact rate, connectivity and centrality in red fox (*Vulpes vulpes*) social groups. Sci Rep. 2019;9(1):20095.

21. Stratford K, Stratford S, Périquet S. Dyadic associations reveal clan size and social network structure in the fission–fusion society of spotted hyaenas. Afr J Ecol. 2020;58(2):182–192.

22. Kleinhappel TK, John EA, Pike TW, Wilkinson A, Burman OHP. Animal welfare: a social networks perspective. Sci Prog. 2016;99(Pt 1):68–82.

23. Connor T, Frank K, Qiao M, Scribner K, Hou J, Zhang J, et al. Social network analysis uncovers hidden social complexity in giant pandas. Ursus. 2023;2023(34e9):1–13.

24. Leu ST, Farine DR, Wey TW, Sih A, Bull CM. Environment modulates population social structure: experimental evidence from replicated social networks of wild lizards. Anim Behav. 2016;111:23–31.

25. Chiyo PI, Archie EA, Hollister-Smith JA, Lee PC, Poole JH, Moss CJ, et al. Association patterns of African elephants in all-male groups: the role of age and genetic relatedness. Anim Behav. 2011;81(6):1093–1099.

26. Goldenberg SZ, de Silva S, Rasmussen HB, Douglas-Hamilton I, Wittemyer G. Controlling for behavioural state reveals social dynamics among male African elephants, *Loxodonta africana*. Anim Behav. 2014;95:111–119.

27. Rosenbaum S, Maldonado-Chaparro AA, Stoinski TS. Group structure predicts variation in proximity relationships between male–female and male–infant pairs of mountain gorillas (Gorilla beringei beringei). Primates. 2016;57(1):17–28.

28. Morrison RE, Eckardt W, Stoinski TS, Brent LJN. Comparing measures of social complexity: larger mountain gorilla groups do not have a greater diversity of relationships. Proc Biol Sci. 2020;287(1931):20201026.

29. Murphy D, Mumby HS, Henley MD. Age differences in the temporal stability of a male African elephant (*Loxodonta africana*) social network. Behav Ecol. 2020;31(1):21–31.

30. Cairns SJ, Schwager SJ. A comparison of association indices. Anim Behav. 1987;35(5):1454–1469.

31. Franks DW, Ruxton GD, James R. Sampling animal association networks with the gambit of the group. Behav Ecol Sociobiol. 2010;64(3):493–503.

32. Hoppitt WJE, Farine DR. Association indices for quantifying social relationships: how to deal with missing observations of individuals or groups. Anim Behav. 2018;136:227–238.

33. Gettler LT, Redhead D, Dzabatou A, Lew-Levy S. BaYaka forager food sharing networks in the Congo Basin: The roles of gender homophily and kin sharing. Am J Biol Anthropol. 2023;181(1):59–69.

34. Motes-Rodrigo A, Albery GF, Negron-Del Valle JE, Philips D, Cayo Biobank Research Unit, Platt ML, et al. Social responses to a hurricane drive greater epidemic risk. bioRxiv. 2023;.

35. Redhead D, Maliti E, Andrews JB, Borgerhoff Mulder M. The interdependence of relational and material wealth inequality in Pemba, Zanzibar. Philos Trans R Soc Lond B Biol Sci. 2023;378(1883):20220288.

36. Redhead D, Ragione AD, Ross CT. Friendship and partner choice in rural Colombia. Evol Hum Behav. 2023;44(5):430–441.

37. Debetencourt B, Barry MM, Arandjelovic M, Stephens C, Maldonado N, Boesch C. Camera traps unveil demography, social structure, and home range of six unhabituated Western chimpanzee groups in the Moyen Bafing National Park, Guinea. Am J Primatol. 2024;86(2):e23578.

38. Sadoughi B, Mundry R, Schülke O, Ostner J. Social network shrinking is explained by active and passive effects but not increasing selectivity with age in wild macaques. Proc Biol Sci. 2024;291(2018):20232736.

39. Siracusa ER, Pavez-Fox MA, Negron-Del Valle JE, Phillips D, Platt ML, Snyder-Mackler N, et al. Social ageing can protect against infectious disease in a group-living primate. bioRxiv. 2024;.

40. Testard C, Shergold C, Acevedo-Ithier A, Hart J, Bernau A, Negron-Del Valle JE, et al. Natural disaster alters the adaptive benefits of sociality in a primate. bioRxiv. 2024;.

41. Hart J, Weiss MN, Franks D, Brent L. BISoN: A Bayesian framework for inference of social networks. Methods Ecol Evol. 2023;14(9):2411–2420.

42. Croft DP, Krause J, Darden SK, Ramnarine IW, Faria JJ, James R. Behavioural trait assortment in a social network: patterns and implications. Behav Ecol Sociobiol. 2009;63(10):1495–1503.

43. de Silva S, Ranjeewa ADG, Kryazhimskiy S. The dynamics of social networks among female Asian elephants. BMC Ecol. 2011;11:17.

44. Bejder L, Fletcher D, BrÄger S. A method for testing association patterns of social animals. Anim Behav. 1998;56(3):719–725.

45. Lusseau D, Schneider K, Boisseau OJ, Haase P, Slooten E, Dawson SM. The bottlenose dolphin community of Doubtful Sound features a large proportion of long-lasting associations. Behav Ecol Sociobiol. 2003;54(4):396–405.

46. Napper CJ, Hatchwell BJ. Social dynamics in nonbreeding flocks of a cooperatively breeding bird: causes and consequences of kin associations. Anim Behav. 2016;122:23–35.

47. Fang G, Jiao HT, Wang MY, Huang PZ, Liu XM, Qi XG, et al. Female demographic changes contribute to the maintenance of social stability within a primate multilevel society. Anim Behav. 2022;192:101–108.

48. Godde S, Humbert L, Côté SD, Réale D, Whitehead H. Correcting for the impact of gregariousness in social network analyses. Anim Behav. 2013;85(3):553–558.

49. Turner JW, Bills PS, Holekamp KE. Ontogenetic change in determinants of social network position in the spotted hyena. Behav Ecol Sociobiol. 2017;72(1):10.

50. van Leeuwen EJC, Cronin KA, Haun DBM. Population-specific social dynamics in chimpanzees. Proceedings of the National Academy of Sciences. 2018;115(45):11393–11400.

51. Hart JDA, Franks DW, Brent L, Weiss MN. BisonR - Bayesian inference of social networks with R; 2022.

52. Ross CT, McElreath R, Redhead D. Modelling human and non-human animal network data in R using STRAND; 2022.

53. Redhead D, McElreath R, Ross CT. Reliable network inference from unreliable data: A tutorial on latent network modeling using STRAND. Psychol Methods. 2023;.

54. Zonana DM, Gee JM, Bridge ES, Breed MD, Doak DF. Assessing Behavioral Associations in a Hybrid Zone through Social Network Analysis: Complex Assortative Behaviors Structure Associations in a Hybrid Quail Population. Am Nat. 2019;193(6):852–865.

55. Farine DR, Firth JA, Aplin LM, Crates RA, Culina A, Garroway CJ, et al. The role of social and ecological processes in structuring animal populations: a case study from automated tracking of wild birds. R Soc Open Sci. 2015;2(4):150057.

56. van Zwet E. A default prior for regression coefficients. Stat Methods Med Res. 2019;28(12):3799–3807.

57. Franks DW, Weiss MN, Silk MJ, Perryman RJY, Croft DP. Calculating effect sizes in animal social network analysis. Methods Ecol Evol. 2020;12(1):33–41.

58. Weiss MN, Franks DW, Brent LJN, Ellis S, Silk MJ, Croft DP. Common datastream permutations of animal social network data are not appropriate for hypothesis testing using regression models. Methods Ecol Evol. 2021;12(2):255–265.

59. Hart JDA, Weiss MN, Brent LJN, Franks DW. Common permutation methods in animal social network analysis do not control for non-independence. Behav Ecol Sociobiol. 2022;76(11):151.

60. Kossinets G. Effects of missing data in social networks. Soc Networks. 2006;28(3):247–268.

61. Huisman M. Imputation of missing network data: Some simple procedures. J Soc Biol Struct. 2009;10(1):1–29.

62. Rubin DB. Inference and missing data. Biometrika. 1976;63(3):581–592.

63. Krause RW, Huisman M, Steglich C, Snijders TAB. Missing Network Data A Comparison of Different Imputation Methods. In: 2018 IEEE/ACM International Conference on Advances in Social Networks Analysis and Mining (ASONAM). IEEE; 2018. p. 159–163.

64. Pavez-Fox MA, Escabi-Ruiz CM, Hart JDA, Negron-Del Valle JE, Phillips D, Montague MJ, et al. Cost and benefits of being social: examining the influence of sociality on faecal parasites in free-ranging rhesus macaques. bioRxiv. 2022;.

65. Hart J. Bayesian Methods for Animal Social Network Analysis. University of Exeter; 2023.

66. Bonnell TR, Vilette C, Henzi SP, Barrett L. Network reaction norms: taking account of network position and plasticity in response to environmental change. Behav Ecol Sociobiol. 2023;77(3):35.

67. Torfs JRR, Stevens JMG, Verspeek J, Laméris DW, Guéry JP, Eens M, et al. Multi-group analysis of grooming network position in a highly social primate. PLoS One. 2023;18(4):e0284361.

68. Stan Development Team. Stan Modeling Language Users Guide and Reference Manual, 2.26.1; 2023; https://mc-stan.org

69. R Core Team. R: A language and environment for statistical computing; 2022.

70. Gabry J, Češnovar R. cmdstanr: R Interface to 'CmdStan'; 2021.

71. Dablander F, Hinne M. Node centrality measures are a poor substitute for causal inference. Sci Rep. 2019;9(1):6846.

72. Youldon DA, Abell J, Briffitt JS, Chama L, Channings MD, Kilundo A, et al. Patch-occupancy survey of elephant (*Loxodonta africana*) surrounding Livingstone, Zambia. Koedoe. 2017;59(1).

73. Douglas-Hamilton I. On the ecology and behaviour of the African elephant: "the elephants of Lake Manyara". University of Oxford; 1972.

74. Moss CJ, Poole JH. Relationships and social structure of African elephants. Primate social relationships: an integrated approach. 1983;315:325.

75. Wittemyer G, Douglas-Hamilton I, Getz WM. The socioecology of elephants: analysis of the processes creating multitiered social structures. Anim Behav. 2005;69(6):1357–1371.

S1 File. The SRI is too trusting of extreme values. Simulation plot to show the difference in potential outputs from SRI versus a Bayesian estimator given true values and the number of times a pair were observed.

S2 File. Standard zero-inflated model. Frequentist two-step model outputs as an alternative option to BISoN.

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure

Figure