Rare species do not disproportionately contribute to phylogenetic diversity in a subalpine

plant community

Leah N Veldhuisen[1,2,*] (https://orcid.org/0000-0002-7094-0108), Verónica Zepeda[3]

(https://orcid.org/0000-0001-8559-2882), Brian J Enquist[1,2] (https://orcid.org/0000-0002-6124-

7096), Katrina M Dlugosch[1] (https://orcid.org/0000-0002-7302-6637)

[1]Department of Ecology & Evolutionary Biology, University of Arizona, Tucson, AZ, 85721

[2]Rocky Mountain Biological Laboratory, 8000 County Rd. 317, Box 519, Crested Butte, CO,

81224

[3] Laboratorio Nacional de Ciencias de la Sostenibilidad, Instituto de Ecología, Universidad

Nacional Autónoma de México, Mexico City, Mexico, 04510

[*]author for correspondence: Leah N Veldhuisen, leah.veldhuisen@gmail.com

ABSTRACT

• **Premise of the study**: Worldwide, 36% of all plant species are exceedingly rare, and are at the

highest risk of extinction. Extinction risk has also been observed to be phylogenetically

clustered, threatening the loss of lineages that contribute unique evolutionary diversity to

communities. Rare plants' contributions to phylogenetic diversity are largely unknown, however.

We investigate whether rare species (as measured by local abundance and range size) contribute

disproportionately to phylogenetic diversity in a subalpine plant community.

• **Methods:** We collected abundance data at three sites in Washington Gulch near the Rocky

Mountain Biological Laboratory (RMBL, Gothic, Colorado, USA) in 2021 and 2022. We then

calculated range size for each species. We calculated phylogenetic signal for abundance and

range size, compared community phylogenetic metrics weighted by range size and abundance to

unweighted metrics, and quantified the change in phylogenetic diversity when removing single

species and groups of species ranked by rarity.

• **Key results**: We found phylogenetic signal for abundance, but not range size. There was no

difference between rarity-weighted and unweighted phylogenetic diversity metrics. Finally,

phylogenetic diversity did not decline more when we removed single rare species and groups of

rare species than when we removed single common species and groups of common species.

• **Conclusions**: Overall, we found that rare species do not disproportionately contribute to

phylogenetic diversity. This suggests that rare species likely provide phylogenetic redundancy

with common species, and losing rare species may not cause disproportionate drops in

phylogenetic diversity in this system.

KEY WORDS: local abundance; phylogenetic diversity; rare species; range size; Rocky

Mountain Biological Laboratory

INTRODUCTION

Worldwide, 36% of all plant species are exceedingly rare (observed only five or fewer

times globally; (Enquist et al., 2019), and are at the highest risk of extinction (Yessoufou and

Davies, 2016; Enquist et al., 2019). At the same time, extinction risk has been observed to be

phylogenetically clustered (Willis et al., 2008; Eiserhardt et al., 2015; González-Orozco et al.,

2016), threatening the loss of lineages that contribute unique evolutionary diversity to

communities. The exact ecological contributions of rare species need more clarification in many

systems (Violle et al., 2017; Dee et al., 2019), but they are known to provide key ecosystem

functions ranging from providing pollinator resources to fire resilience to diversity across trophic

levels (Bracken and Low, 2012; Mouillot et al., 2013; Leitão et al., 2016). However, their

contributions to phylogenetic diversity have been explored in very few systems (but see

(Anderson et al., 2004; Mi et al., 2012; Herzog and Latvis, 2022).

Understanding factors impacting phylogenetic diversity is key, as it is more positively

correlated with biomass than species richness or functional diversity (Cadotte et al., 2008).

Phylogenetic diversity also promotes ecosystem stability and productivity (Cadotte et al., 2012;

Cadotte, 2013; Davies et al., 2016), diversity in other taxonomic groups (Dinnage et al., 2012;

Milcu et al., 2013), and can increase resistance to invasion (Galland et al., 2019; Delavaux et al.,

2023). More broadly, phylogenetic diversity can capture variation between species that is missed

when only studying one or a few traits (Cadotte et al., 2009; E-Vojtkó et al., 2023) and more

accurately reflects niche differences (Huang et al., 2020). It is also an increasingly important axis

of diversity to consider in plant conservation (Tucker et al., 2012; Molina-Venegas et al., 2021).

Given the benefits of phylogenetic diversity, the contributions of rare species are important to

clarify in light of ongoing species losses (White et al., 2023).

Rarity can be categorized in multiple ways in terms of small population size, small range

size, narrow habitat specificity, or combinations thereof (Rabinowitz, 1981). Here we

investigate two major axes of rarity: local abundance and range size. Species can be categorized

as rare using these metrics individually or in combination, and they do not always correlate

(Sexton et al., 2009), so it is unclear whether range size might follow the same phylogenetic

diversity patterns as abundance. For example, a species may have very high local abundance but

have a very restricted distribution or vice versa. Species' range sizes and limits are shifting,

shrinking or expanding with global change (Hardie and Hutchings, 2010; Chen et al., 2011;

Sheth and Angert, 2018). Like low-abundance species, species with small ranges are also at

higher risk of extinction (Harnik et al., 2012; Manes et al., 2021). To our knowledge, there has

never been a test of rare species' contributions to phylogenetic diversity based on range size.

We might expect both low abundance and small-ranged species to be more

phylogenetically diverse than random (overdispersed) in relation to each other and their larger

community (Mi et al., 2012). The niche differentiation hypothesis suggests that each species

within a community occupies its own niche, that rare species use distinct resources, and that rare

species will be distinct from each other and more abundant species (Hanski, 1982; Gaston, 1994;

Grime, 1998). We now know that phylogenetic diversity is rarely a suitable proxy for functional

diversity (Gerhold et al., 2015; Cadotte et al., 2019; E-Vojtkó et al., 2023), though the

association of phylogenetic diversity with ecosystem properties suggests that it can capture

aspects of community-level function (Cadotte et al. 2009; E-Vojtkó et al. 2023; Cadotte et al.

2008; Davies et al. 2016).

Previous work supports rare species' distinctiveness in some systems and by some

metrics. A study by Jiao et al. (2017) found that rare microbes have higher alpha and beta

diversity than their abundant counterparts, but this study did not test phylogenetic diversity. In

desert annual plants, rare species do not closely exhibit a well-established functional tradeoff

between water use efficiency and relative growth rate seen in the dominant species, suggesting

niche differentiation in physiological traits (Ge et al., 2019). Like Jiao et al. (2017), Ge and

colleagues (2019) did not test the relatedness of species evading the tradeoff, leaving open an

important question about how phylogenetic relatedness may explain differences in rare species'

ecological strategies. In tropical tree plots worldwide, rare species were found to be

phylogenetically more diverse than abundant species, although this only held true in less

disturbed plots (Mi et al., 2012). In yeasts growing in decaying cactus tissue, researchers found

that abundance was linked to phylogenetic relatedness, but it varied depending which subfamily

of cactus the yeast was growing within (Anderson et al., 2004).

Alternatively, both low abundance and small-ranged species may be randomly distributed

across the phylogeny. In this case, we would not expect low abundance and small-ranged species

to disproportionately contribute to overall phylogenetic diversity, and instead they would provide

phylogenetic redundancy with more abundant species. In tall grass prairies, rare species indeed

did not contribute disproportionate phylogenetic diversity, and provided phylogenetic

redundancy within the community (Herzog and Latvis, 2022).

Here, we investigate whether low abundance and small-ranged species contribute

disproportionately to phylogenetic diversity in a subalpine plant community. Consistent with the

idea that rare species are unique from more common ones, we hypothesize that rare species

disproportionately contribute to phylogenetic diversity. We test this idea across an elevational

gradient in the Colorado Rocky Mountains near the Rocky Mountain Biological Laboratory

(RMBL, Gothic, Colorado, USA). We first predict (1) that range size and local abundance will

both show phylogenetic signal, indicating that common and rare species occupy different

phylogenetic lineages. Then, we predict (2) that weighting phylogenetic diversity metrics by high

abundance or large range size will decrease phylogenetic diversity, consistent with the prediction

that rare species contribute more to diversity than common species. Finally, we predict (3) that

removing species according to rarity will decrease phylogenetic diversity more when the

removed species are rare than when they are common.

MATERIALS & METHODS

Study system & abundance data

We collected abundance data for an angiosperm community at three sites in Washington

Gulch near the Rocky Mountain Biological Laboratory (RMBL, Gothic, Colorado, USA) from

June to August in 2021 and 2022. RMBL is located in the East River valley of the West Elk

mountains, approximately 10 kilometers from Crested Butte, Colorado. Study sites were located

at 2815 m (38°53'50"N, 106°58'43"W), 3165 m (38°57'38"N, 107°01'53"W), and 3380 m

(38°58'10"N, 107°01'53"W) in elevation. At each of the three sites, we had five 1.2 m x 1.2 m

plots that were randomly placed within each site each summer (see more detailed description of

sites and plots in (Veldhuisen et al., 2023) and (Sloat et al., 2015)).

Within each 1.2 m x 1.2 m plot at the three sites, we identified all vascular plants to

species level and confirmed our identifications against RMBL Herbarium specimens (herbarium

code RMBL) and iNaturalist observations. Each plot was sampled once per year near the peak of

the growing season (approximately mid-July, depending on the year and elevation). Plant

sampling dates were staggered to ensure all communities had the same relative phenological time

point. In each plot we counted all individuals of every species. We only counted plants rooted

within the plot, and unknown and non-native species were excluded from all analyses. To

quantify abundance, we averaged local abundance across all five plots at each site and the two

data collection years. We did not include invasive species or species that were only identified to

genus in any of our analyses.

Range size

We calculated range size as Extent of Occurrence (EOO) and Area of Occupancy (AOO)

(Gaston and Fuller, 2009; Staude et al., 2020). To obtain these values, we downloaded

occurrence data for each species using the 'RGBIF' package (Appendix S1; Chamberlain and

Boettiger, 2017) and calculated AOO and EOO with the 'red' package (Cardoso, 2017). We

restricted GBIF downloads to occurrences in North America with <1000m coordinate

uncertainty and observations that have been made since 1990 to ensure coordinate accuracy. We

used only the AOO values to calculate phylogenetic signal (Prediction 1) and rank species for

testing the impact of their removal (Prediction 2), because AOO is less sensitive to extreme and

outlier observations (Gaston and Fuller, 2009; Staude et al., 2020).

Phylogenetic metrics

To calculate phylogenetic signal and community phylogenetic diversity, we used the

'ALLMB.tre' angiosperm phylogeny from (Smith and Brown, 2018). This phylogeny includes

all genera in our dataset and 54 of 68 species. If a species was missing from the phylogeny but

was the only member of its genus in the plots, we replaced it with a closely related congener in

the Smith and Brown phylogeny (Appendix S2). Only the genus *Senecio* had multiple species in

the plots missing from the phylogeny. We replaced *Senecio integerrimus* Nutt. with sister species

S. triangularis Hook (Pelser et al., 2007). We removed *S. crassulus* A. Gray and *S. bigelovii* A.

Gray from all phylogenetic analyses because they have no available genetic or phylogenetic

information. We could not find an up-to-date genus-level phylogeny for *Erigeron* to inform

replacements, so we removed *Erigeron elatior* and *Erigeon glacialis* from all analyses.

We calculated phylogenetic signal for both abundance and range size with Blomberg's K

(Blomberg et al., 2003) and Pagel's λ (Pagel, 1999). We calculated Blomberg's K with the

"phylosignal()" function in the 'picante' R package (Kembel et al., 2010) with 5000

randomizations, and Pagel's λ with the function "phylosig()" in the 'phytools' R package

(Revell, 2011) with 5000 iterations and model set to "lambda." Blomberg's K or Pagel's λ values

near 1 indicate trait evolution consistent with Brownian motion, while K less than 1 or Pagel's λ

near zero suggest less phylogenetic signal than expected by Brownian motion.

We calculated Faith's phylogenetic diversity (PD; (Faith, 1992), mean phylogenetic

distance (MPD), and mean nearest taxon distance (MNTD) and their associated standard effect

sizes (SES) in 'picante' (Kembel et al., 2010). Faith's Phylogenetic Diversity (Faith, 1992) is the

total evolutionary distance (branch length) across a phylogenetic tree describing the relationships

among the taxa in a dataset. This metric will increase if relationships are more distant across the

tree (species are more evolutionarily divergent). Mean Phylogenetic Distance is the mean

distance between pairs of species, and so it also increases as evolutionary relationships are more

distant across the tree, but will be less sensitive to rare cases of very distantly-related species that

would otherwise elevate Faith's Phylogenetic Diversity. Mean Nearest Taxon Distance is the

mean of the distance to the closest relative of each taxon, and so it reflects the typical distance to

any species' closest relative in the dataset and increases as closest relatives are more distant.

Therefore, MNTD is more reflective of the tendency of species to have closer relatives than it is

of the overall evolutionary diversity across the entire tree (community of species) that is captured

by the former metrics (for further discussion, see review by (Tucker et al., 2017)).

To calculate SES, we used each metric's "sample.pool" null model to draw species from

the RMBL community with 5000 iterations randomly. Positive SES values indicate that the

species group in question is more phylogenetically diverse than a random draw of the same

number of species from the regional species pool (overdispersion). SES values above 1.96 will

have P values above 0.975, and indicate statistical significance of the SES. Negative SES values

indicate that the species group in question is less phylogenetically diverse than a random draw of

the same number of species from the regional species pool (underdispersion). SES values below

-1.96 will generate P values below 0.025, and indicate statistical significance of the SES.

Weighting by abundance and range size

To test prediction 2, we calculated MPD and MNTD weighted by abundance and range

size. We then compared the weighted versus unweighted standard effect sizes for each

community. To weight these metrics by abundance, we used the "ses.mpd" function in 'picante'

(Kembel et al., 2010), but switched the "abundance.weighted=" parameter from *false* to *true*. To

weight by range size, we weighted by the log of the AOO value. PD cannot be weighted by

abundance, since it is directly correlated with species richness. We defined a significant change

in diversity in response to weighting as a sign change in the SES or if SES absolute values

became greater than 1.96.

Removing species

To test prediction 3, we removed individual species and groups of species and

recalculated phylogenetic diversity after the removal. For each site, we ranked species in order of

decreasing rarity in terms of both local abundance and range size, such that the most abundant

and largest range size species were ranked first and the lowest abundance and smallest range size

species were ranked last. This resulted in two separate rankings of rarity, one for local abundance

and one for range size. Then, we removed one species at a time in order of increasing rarity

(removing most abundant and largest range sizes first) and calculated the SES for PD, MPD, and

MNTD of the resulting community. We also tested for autocorrelation across species of adjacent

ranks within each site using the Durbin-Wallis test with 1, 2 and 3 steps to see if removal of

similarly-ranked species had similar effects.

To test the impact of removing groups of species at a time, we removed groups of ten

species, also in order of abundance and range size. Using a sliding window approach, we created

groups of ten species starting with the top 10 most abundant species, then shifting to species

ranked 2-11, 3-12, and so on until the 10 least abundant species. We then calculated the SES of

phylogenetic diversity for the community with each group of 10 removed. We analyzed all data

in R version 4.3.1 (R Core Team, 2023).

RESULTS

Community composition & phylogenetic signal

The low elevation site contained 39 species across 17 families, the middle site had 27

species across 14 families, and the high site had 36 species across 20 families. We found that

local abundance ranged from 0.5 to 573.5 individuals per site when pooled across plots and

averaged across years, with a median average abundance of 10.5 individuals per site. At all three

sites, the least abundant species had only one individual in one year with an average of 0.5

individuals per site across years (*Noccaea fendleri, Primula pauciflora* and *Rumex densiflorus* at

the low elevation site, *Eucephalus breweri* at the middle and *Arctostaphylos uva-ursi, Erigeron*

speciosus, Hydrophyllum capitatum, Senecio integerrimus and *Erigeron eatonii* at the high

elevation site). At the low elevation site, the most abundant species was *Collomia linearis* with

an average abundance of 573.5. At the middle elevation site, *Heliomeris multiflora* was the most

abundant with an average of 253.5 individuals in the site, and at the high elevation site

Polygonum douglasii was the most abundant with an average of 130 individuals in the site.

Range size measured as AOO spanned from 108 km^2 for *Rumex densiflorus* to 469,436 km^2 for

Achillea millefolium, with a median range size 2316 km^2. Both abundance and range size

followed standard rank abundance curves (Fig. 1A-B).

We found phylogenetic signal for abundance with only Blomberg's K (Blomberg's K =

0.304, P-value = 0.04; Pagel's λ = 0.926, P-value =0.17; Fig. 2A), but not for range size with

either metric (Blomberg's K = 0.09, P-value = 0.7; Pagel's λ = 7.3x10^{-5}, P-value =1; Fig. 2B).

Weighting community phylogenetic metrics

Weighting MPD and MNTD by abundance (Fig. 3A) and range size (Fig. 3B) did not

significantly change phylogenetic diversity at any of our sites. At the low elevation site, MPD

SES was positive when unweighted, and negative when weighted by range size but both SES

values were still very near zero and nonsignificant (Fig. 3B). Also at the low elevation site,

weighting by abundance increased SES values for MPD and MNTD, but SES values stayed

nonsignificant. At other sites, impacts of weighting were variable and depended on the metric in

question (Fig. 3A-B), but never caused the SES values to be statistically significant.

Removing species

Because range size did not show phylogenetic signal, we only show results for abundance

(for single species range size removal results see Appendix S3). For abundance, removing

species in order of decreasing abundance impacted SES values inconsistently across metrics and

sites (Fig. 4). Removing the rarest species did not consistently decrease phylogenetic diversity by

any metric. At the high elevation site, removing the most abundant species did not change the

SES for PD, slightly lowered SES for MPD, and increased the SES for MNTD (Fig. 4).

Removing the least abundant species increased the SES for PD, MPD and MNTD, and the

middle abundance species both increased and decreased SES values for all metrics depending on

the species (Fig. 4).

At the middle elevation site, removal of the most and least abundant species increased

SES across all metrics (Fig. 4). Removing other low-abundance species (ranked 20-26 out of 29)

both increased and decreased SES values depending on metric and individual species (Fig. 4).

Like the high elevation site, removing the middle-ranked abundance species had mixed impacts

on community phylogenetic diversity by all metrics (Fig. 4).

Finally, at the low elevation site removal of the most and least abundant species lowered

SES values by all metrics (Fig. 4). For both PD and MTD, SES values decreased with the

removal of each of the top five most abundant species (Fig. 4). This decrease in SES was also

mostly true for MPD, except that the removal of the third most abundant species did not change

the SES value. While the removal of the least abundant species decreased SES values for all

metrics, removal of the next three least abundant species either did not change or increased SES

values depending on metric and species (Fig. 4). Like the middle and high elevation sites,

removing the middle ranked species at the low elevation site had variable impacts on SES values

for all metrics (Fig. 4). We tested for autocorrelation between points within each site at one, two

and three steps, and found no significant autocorrelation (P>0.05 for all; Appendix S4).

Removing groups of ten species in order of abundance also did not produce consistent

patterns (Fig. 5). Removing the groups with the most rare species did not decrease SES for PD,

and instead increased SES at all sites (Fig. 5). At the low elevation site, removing groups of

common species had the most negative impact on SES values, while removing the most common

species did not impact SES at the high elevation site, and increased the SES at the middle

elevation site (Fig. 5). We only show results for PD here, as results for MPD and MNTD were

similarly inconsistent (Appendix S5). Like the removal of individual species, we only removed

groups of species based on abundance, because we found no phylogenetic signal in range size

(Fig. 2B).

DISCUSSION

We hypothesized that rare species (quantified by local abundance and range size) would

contribute disproportionately to phylogenetic diversity, but our results do not support our

associated predictions. We found that abundance did show phylogenetic signal, but range size

did not. We predicted that weighting MPD and MNTD by higher abundance and larger range

size would decrease phylogenetic diversity (consistent with disproportionately high contributions

by rare species), but it did not. Finally, we predicted that removing rare species (individually or

in groups) would decrease phylogenetic diversity more than removing common species, but this

was not consistently the case. Our results are strikingly consistent in suggesting that rare species

do not have disproportionately high contributions to phylogenetic diversity in this system.

Our results are consistent with a similar recent study that found that rare and common

species contribute similarly to phylogenetic diversity in tallgrass prairies (Herzog and Latvis,

2022). The authors sampled 21 tallgrass prairie communities, and calculated PD each time they

added increasingly rare species, which were quantified by percent cover. They then calculated

breakpoints where phylogenetic diversity changed significantly from one species' addition to the

next, but found that the location of statistically significant breakpoints was not consistently

associated with more rare species (Herzog and Latvis, 2022). While we did not calculate

breakpoints in our data, Herzog and Latvis's results are similar in that species' phylogenetic

contributions are not consistently predictable based on rarity.

Based on our results, we find no evidence that rare species are phylogenetically distant

from the rest of the community. However, we did not investigate their functional uniqueness,

which would likely produce a different pattern. Phylogenetic diversity often correlates poorly

with variation in individual traits (CaraDonna and Inouye, 2015; Gerhold et al., 2015; Maitner et

al., 2022; E-Vojtkó et al., 2023), indicating that phylogenetic and functional diversity patterns

are not interchangeable. Similarly, prioritizing phylogenetic diversity for conservation efforts

does not always maximize functional diversity, even when traits show phylogenetic signal

(Mazel et al., 2017). Work in other systems has suggested that rare species provide the most

unique functional trait combinations (Mouillot et al., 2013) and contribute significantly to

functional diversity (Leitão et al., 2016; Coulon et al., 2023). Whether our rare RMBL plant

community members are functionally unique or redundant is unknown, but our results suggest at

least that low abundance and small range size do provide redundancy in terms of phylogenetic

diversity.

In contrast, Mi at al. (2012) tested for niche differentiation by comparing the

phylogenetic diversity of rare and common species in temperate and tropical tree plots, and

suggested that the niche differentiation hypothesis is supported in some of their plots because of

rare species' phylogenetic distinctiveness. While we now know we cannot assume relatedness

predicts niche space in a straightforward way (Mason and Pavoine, 2013; Davies et al., 2016;

Blonder, 2018), Mi and colleague's results are informative. In particular, they found that rare

species contribute more phylogenetic diversity only in plots with lower levels of disturbance (Mi

et al., 2012). If disturbance has caused local extinctions of rare species before the time frame of

our study, the rarest species will no longer be present, and we would instead be observing the

less rare species that are still present. If this were the case, it may still be true that rare species

historically contributed the most phylogenetic diversity, but we would not know. We also

measured the impact of the removal of one species at a time and in groups of ten, but patterns of

actual loss in this system are unknown and are likely to be highly system-specific (Zavaleta and

Hulvey, 2004; Hooper et al., 2012; Sanders et al., 2018; Chase et al., 2020; Carmona et al.,

2021). We did not evaluate disturbance or historical species loss at our sites, and this would be

an important avenue for future research in this system.

The addition of new species via invasions or range shifts may also drive declines in

phylogenetic diversity (Selvi et al., 2016; Fitt and Lancaster, 2017; Daru et al., 2021), regardless

of whether native species are declining. We know that range shifting and invasive species are

impacting this system at the same time others are declining (Renwick et al., 2016; Zorio et al.,

2016; Shepard et al., 2022). We did not include non-native species in our analyses since

quantifying their range size is more complex, and only four non-native species occur at low

abundance in our plots to begin with. Three of the four non-native species are in Asteraceae, and

as such would likely not change our results. Evidence does suggest plants are shifting their

elevational ranges in this system (Zorio et al., 2016), however, so investigating the phylogenetic

impact of newcomers may be important for future work.

349   Finally, it is notable that we did find phylogenetic signal in abundance but not in range

size. Herzog & Latvis (2022) tested for phylogenetic signal in abundance in their tall grass

prairie communities, and found signal at the regional scale and in 5 of their 21 communities.

Previous work has tested for phylogenetic signal in range size with mixed results (Morin and

Lechowicz, 2013; Zacaï et al., 2017; Pigot et al., 2018). Pigot and colleagues (2018) found that

introduced bird species have strong phylogenetic signal in their native range size, but random

samples of birds do not show phylogenetic signal in range size, so this finding is unique to

introduced species. Morin and Lechowicz (2013) found weak but significant signal in range size

in tree species in forest plots worldwide. Zacaï (Morin and Lechowicz, 2013; Zacaï et al., 2017;

Pigot et al., 2018) and colleagues (2017) found that phylogenetic signal in range size varies

regionally, and is likely influenced by spatio-temporal environmental stability over geological

time. Each of these studies address range size across very broad spatial and temporal scales

(Morin and Lechowicz, 2013; Zacaï et al., 2017; Pigot et al., 2018), while our study only

includes present-day species native to the North American Rocky Mountains. This indicates that

the scale tested likely influences whether or not range size shows phylogenetic signal, which is

consistent with other scale-dependent community phylogenetic patterns (Cavender-Bares et al.,

2009; Willis et al., 2009; Park et al., 2020).

CONCLUSIONS

We found that rare species do not contribute disproportionately to phylogenetic diversity,

and both raise and lower phylogenetic diversity relative to richness depending on the identity of

the species, elevation and phylogenetic metric. This finding is consistent with previous work (Mi

et al., 2012; Herzog and Latvis, 2022), and suggests that rare species may add phylogenetic

redundancy rather than diversity in this community. To further clarify the impact of rare species

on biodiversity, future work should assess which species are disappearing, which species may

have already gone locally extinct, the functional redundancy of rare species, and the impact of

incoming species to the system.

ACKNOWLEDGEMENTS

This study was conducted on the traditional and unceded land of the Ute People. The

authors thank RMBL and private landowners for land access, RMBL for securing permits,

Dlugosch lab members for manuscript feedback, and Dan Park for discussion of analyses and

results. This work was funded by Rocky Mountain Biological Laboratory, Botanical Society of

America, and University of Arizona Graduate & Professional Student Council awards to LNV

and United States National Science Foundation grant #1750280 to KMD. BJE acknowledges that

this material is based upon work supported by the U.S. Department of Energy, Office of Science,

Office of Biological and Environmental Research, Earth and Environmental Systems Sciences

Division and Data Management Program, under Award Number DEAC02- 05CH11231, as part

of the Watershed Function Scientific Focus Area and the ExaShed project. BJE was also

supported by NSF awards 2225078 and 2225076.

AUTHOR CONTRIBUTIONS

391 L.N.V. and K.M.D. conceived of the study. V.Z. collected field data. L.N.V. generated range

392 size data and analyzed data. L.N.V wrote the original draft with input from K.M.D. and V.Z. All

393 authors provided feedback. L.N.V., B.J.E and K.M.D acquired funding. All authors approved the

394 final version of the manuscript.

DATA AVAILABILITY STATEMENT

Range size and abundance data can be accessed at

https://doi.org/10.6073/pasta/320111b8e067c3b443fa25a719aac7ae. Code is available at

https://github.com/lveldhuisen/phylogenetics-and-ranges.

Additional supporting information may be found online in the Supporting Information

section at the end of the article.

Appendix S1: Citations for GBIF downloads used to calculate range size.

Appendix S2: Species in our research sites listed with replacement species used from

Smith & Brown's ALLMB.tre phylogeny (see also Methods).

Appendix S3: Results of phylogenetic diversity (PD, MPD & MNTD) from removing

single species based on rage size.

Appendix S4: P-values for autocorrelation tests at each site for both abundance and

range size, all three phylogenetic metrics and 1, 2 and 3 steps.

Appendix S5: Results of additional phylogenetic diversity metrics (MPD & MNTD)

from removing groups of species based on abundance.

LITERATURE CITED

Anderson, T. M., M. Lachance, and W. T. Starmer. 2004. The Relationship of Phylogeny to Community Structure: The Cactus Yeast Community. *The American Naturalist* 164: 709–721.

Blomberg, S. P., T. Garland, and A. R. Ives. 2003. Testing for Phylogenetic Signal in Comparative Data: Behavioral Traits Are More Labile. *Evolution* 57: 717–745.

Blonder, B. 2018. Hypervolume concepts in niche- and trait-based ecology. *Ecography* 41: 1441–1455.

Bracken, M. E. S., and N. H. N. Low. 2012. Realistic losses of rare species disproportionately impact higher trophic levels: Loss of rare 'cornerstone' species. *Ecology letters* 15: 461–467.

Cadotte, M. W. 2013. Experimental evidence that evolutionarily diverse assemblages result in higher productivity. *Proceedings of the National Academy of Sciences of the United States of America* 110: 8996–9000.

Cadotte, M. W., M. Carboni, X. Si, and S. Tatsumi. 2019. Do traits and phylogeny support congruent community diversity patterns and assembly inferences? *The Journal of ecology* 107: 2065–2077.

Cadotte, M. W., B. J. Cardinale, and T. H. Oakley. 2008. Evolutionary history and the effect of biodiversity on plant productivity. *Proceedings of the National Academy of Sciences of the United States of America* 105: 17012–17017.

Cadotte, M. W., J. Cavender-Bares, D. Tilman, and T. H. Oakley. 2009. Using phylogenetic, functional and trait diversity to understand patterns of plant community productivity. *PloS one* 4: e5695.

Cadotte, M. W., R. Dinnage, and D. Tilman. 2012. Phylogenetic diversity promotes ecosystem stability. *Ecology* 93: S223–S233.

CaraDonna, P. J., and D. W. Inouye. 2015. Phenological responses to climate change do not exhibit phylogenetic signal in a subalpine plant community. *Ecology* 96: 355–361.

Cardoso, P. 2017. red - an R package to facilitate species red list assessments according to the IUCN criteria. *Biodiversity data journal*: e20530.

Carmona, C. P., R. Tamme, M. Pärtel, F. de Bello, S. Brosse, P. Capdevila, R. González-M, et al. 2021. Erosion of global functional diversity across the tree of life. *Science advances* 7: eabf2675.

Cavender-Bares, J., K. H. Kozak, P. V. A. Fine, and S. W. Kembel. 2009. The merging of community ecology and phylogenetic biology. *Ecology letters* 12: 693–715.

Chamberlain, S. A., and C. Boettiger. 2017. R Python, and Ruby clients for GBIF species occurrence data. *PeerJ Preprints*.

Chase, J. M., S. A. Blowes, T. M. Knight, K. Gerstner, and F. May. 2020. Ecosystem decay exacerbates biodiversity loss with habitat loss. *Nature* 584: 238–243.

Chen, I.-C., J. K. Hill, R. Ohlemüller, D. B. Roy, and C. D. Thomas. 2011. Rapid range shifts of species associated with high levels of climate warming. *Science* 333: 1024–1026.

Coulon, N., M. Lindegren, E. Goberville, A. Toussaint, A. Receveur, and A. Auber. 2023. Threatened fish species in the Northeast Atlantic are functionally rare. *Global ecology and biogeography: a journal of macroecology* 32: 1827–1845.

Daru, B. H., T. J. Davies, C. G. Willis, E. K. Meineke, A. Ronk, M. Zobel, M. Pärtel, et al. 2021. Widespread homogenization of plant communities in the Anthropocene. *Nature communications* 12: 6983.

Davies, T. J., M. C. Urban, B. Rayfield, M. W. Cadotte, and P. R. Peres-Neto. 2016. Deconstructing the relationships between phylogenetic diversity and ecology: a case study on ecosystem functioning. *Ecology* 97: 2212–2222.

Dee, L. E., J. Cowles, F. Isbell, S. Pau, S. D. Gaines, and P. B. Reich. 2019. When Do Ecosystem Services Depend on Rare Species? *Trends in ecology & evolution* 34: 746–758.

Delavaux, C. S., T. W. Crowther, C. M. Zohner, N. M. Robmann, T. Lauber, J. van den Hoogen, S. Kuebbing, et al. 2023. Native diversity buffers against severity of non-native tree invasions. *Nature* 621: 773–781.

Dinnage, R., M. W. Cadotte, N. M. Haddad, G. M. Crutsinger, and D. Tilman. 2012. Diversity of plant evolutionary lineages promotes arthropod diversity. *Ecology letters* 15: 1308–1317.

Eiserhardt, W. L., F. Borchsenius, C. M. Plum, A. Ordonez, and J.-C. Svenning. 2015. Climate-driven extinctions shape the phylogenetic structure of temperate tree floras. *Ecology letters* 18: 263–272.

Enquist, B. J., X. Feng, B. Boyle, B. Maitner, E. A. Newman, P. M. Jørgensen, P. R. Roehrdanz, et al. 2019. The commonness of rarity: Global and future distribution of rarity across land plants. *Science Advances* 5: eaaz0414.

E-Vojtkó, A., F. de Bello, Z. Lososová, and L. Götzenberger. 2023. Phylogenetic diversity is a weak proxy for functional diversity but they are complementary in explaining community assembly patterns in temperate vegetation. *The Journal of ecology*.

Faith, D. P. 1992. Conservation evaluation and phylogenetic diversity. *Biological conservation* 61: 1–10.

Fitt, R. N. L., and L. T. Lancaster. 2017. Range shifting species reduce phylogenetic diversity in high latitude communities via competition. *The Journal of animal ecology* 86: 543–555.

Galland, T., G. Adeux, H. Dvořáková, A. E-Vojtkó, I. Orbán, M. Lussu, J. Puy, et al. 2019.
Colonization resistance and establishment success along gradients of functional and
phylogenetic diversity in experimental plant communities. *The Journal of ecology* 107:
2090–2104.

Gaston, K. J. 1994. Biodiversity - measurement. *Progress in Physical Geography: Earth and*
Environment 18: 565–574.

Gaston, K. J., and R. A. Fuller. 2009. The sizes of species' geographic ranges. *The Journal of*
applied ecology 46: 1–9.

Gerhold, P., J. F. Cahill, M. Winter, I. V. Bartish, and A. Prinzing. 2015. Phylogenetic patterns
are not proxies of community assembly mechanisms (they are far better). *Functional*
ecology 29: 600–614.

Ge, X.-Y. M., J. P. Scholl, U. Basinger, T. E. Huxman, and D. L. Venable. 2019. Functional trait
trade-off and species abundance: insights from a multi-decadal study. *Ecology letters* 22:
583–592.

González-Orozco, C. E., L. J. Pollock, A. H. Thornhill, B. D. Mishler, N. Knerr, S. W. Laffan, J.
T. Miller, et al. 2016. Phylogenetic approaches reveal biodiversity threats under climate
change. *Nature climate change* 6: 1110–1114.

Grime, J. P. 1998. Benefits of plant diversity to ecosystems: immediate, filter and founder
effects. *The Journal of ecology* 86: 902–910.

Hanski, I. 1982. Dynamics of Regional Distribution: The Core and Satellite Species Hypothesis.
Oikos 38: 210–221.

Hardie, D. C., and J. A. Hutchings. 2010. Evolutionary ecology at the extremes of species'
ranges. *Environmental Review* 18: 1–20.

Harnik, P. G., C. Simpson, and J. L. Payne. 2012. Long-term differences in extinction risk
among the seven forms of rarity. *Proceedings. Biological sciences / The Royal Society* 279:
4969–4976.

Herzog, S. A., and M. Latvis. 2022. Community-level phylogenetic diversity does not differ
between rare and common lineages across tallgrass prairies in the northern Great Plains.
Ecology and evolution 12: e9453.

Hooper, D. U., E. C. Adair, B. J. Cardinale, J. E. K. Byrnes, B. A. Hungate, K. L. Matulich, A.
Gonzalez, et al. 2012. A global synthesis reveals biodiversity loss as a major driver of
ecosystem change. *Nature* 486: 105–108.

Huang, M., X. Liu, M. W. Cadotte, and S. Zhou. 2020. Functional and phylogenetic diversity
explain different components of diversity effects on biomass production. *Oikos* 129: 1185–
1195.

Jiao, S., W. Chen, and G. Wei. 2017. Biogeography and ecological diversity patterns of rare and
abundant bacteria in oil-contaminated soils. *Molecular ecology* 26: 5305–5317.

Kembel, S. W., P. D. Cowan, M. R. Helmus, W. K. Cornwell, H. Morlon, D. D. Ackerly, S. P.
Blomberg, and C. O. Webb. 2010. Picante: R tools for integrating phylogenies and ecology.
Bioinformatics 26: 1463–1464.

Leitão, R. P., J. Zuanon, S. Villéger, S. E. Williams, C. Baraloto, C. Fortunel, F. P. Mendonça,
and D. Mouillot. 2016. Rare species contribute disproportionately to the functional structure
of species assemblages. *Proceedings. Biological sciences / The Royal Society* 283.

Maitner, B. S., D. S. Park, B. J. Enquist, and K. M. Dlugosch. 2022. Where we've been and
where we're going: the importance of source communities in predicting establishment
success from phylogenetic relationships. *Ecography* 2022.

Manes, S., M. J. Costello, H. Beckett, A. Debnath, E. Devenish-Nelson, K.-A. Grey, R. Jenkins,
et al. 2021. Endemism increases species' climate change risk in areas of global biodiversity
importance. *Biological conservation* 257: 109070.

Mason, N. W. H., and S. Pavoine. 2013. Does trait conservatism guarantee that indicators of
phylogenetic community structure will reveal niche-based assembly processes along stress
gradients? *Journal of vegetation science: official organ of the International Association for
Vegetation Science* 24: 820–833.

Mazel, F., A. O. Mooers, G. V. D. Riva, and M. W. Pennell. 2017. Conserving Phylogenetic
Diversity Can Be a Poor Strategy for Conserving Functional Diversity. *Systematic biology*
66: 1019–1027.

Milcu, A., E. Allan, C. Roscher, T. Jenkins, S. T. Meyer, D. Flynn, H. Bessler, et al. 2013.
Functionally and phylogenetically diverse plant communities key to soil biota. *Ecology* 94:
1878–1885.

Mi, X., N. G. Swenson, R. Valencia, W. J. Kress, D. L. Erickson, Á. J. Pérez, H. Ren, et al.
2012. The contribution of rare species to community phylogenetic diversity across a global
network of forest plots. *The American naturalist* 180: E17–30.

Molina-Venegas, R., M. Á. Rodríguez, M. Pardo-de-Santayana, C. Ronquillo, and D. J.
Mabberley. 2021. Maximum levels of global phylogenetic diversity efficiently capture plant
services for humankind. *Nature ecology & evolution* 5: 583–588.

Morin, X., and M. J. Lechowicz. 2013. Niche breadth and range area in North American trees.
Ecography 36: 300–312.

Mouillot, D., D. R. Bellwood, C. Baraloto, J. Chave, R. Galzin, M. Harmelin-Vivien, M.
Kulbicki, et al. 2013. Rare species support vulnerable functions in high-diversity
ecosystems. *PLoS biology* 11: e1001569.

Pagel, M. 1999. Inferring the historical patterns of biological evolution. *Nature* 401: 877–884.

Park, D. S., X. Feng, B. S. Maitner, K. C. Ernst, and B. J. Enquist. 2020. Darwin's naturalization
conundrum can be explained by spatial scale. *Proceedings of the National Academy of*
Sciences of the United States of America 117: 10904–10910.

Pelser, P. B., B. Nordenstam, J. W. Kadereit, and L. E. Watson. 2007. An ITS phylogeny of tribe
Senecioneae (Asteraceae) and a new delimitation of Senecio L. *Taxon* 56: 1077–1104.

Pigot, A. L., E. E. Dyer, D. W. Redding, P. Cassey, G. H. Thomas, and T. M. Blackburn. 2018.
Species invasions and the phylogenetic signal in geographical range size. *Global ecology*
and biogeography: a journal of macroecology 27: 1080–1092.

Rabinowitz, D. 1981. Seven forms of rarity. *Biological aspects of rare plant conservation.*

R Core Team. 2023. R: A language and environment for statistical computing.

Renwick, K. M., M. E. Rocca, and T. J. Stohlgren. 2016. Biotic disturbance facilitates range shift
at the trailing but not the leading edge of lodgepole pine's altitudinal distribution. *Journal of*
vegetation science: official organ of the International Association for Vegetation Science
27: 780–788.

Revell, L. J. 2011. phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology & Evolution* 3: 217–223.

Sanders, D., E. Thébault, R. Kehoe, and F. J. Frank van Veen. 2018. Trophic redundancy reduces
vulnerability to extinction cascades. *Proceedings of the National Academy of Sciences of*
the United States of America 115: 2419–2424.

Selvi, F., E. Carrari, and A. Coppi. 2016. Impact of pine invasion on the taxonomic and
phylogenetic diversity of a relict Mediterranean forest ecosystem. *Forest ecology and*
management 367: 1–11.

Sexton, J. P., P. J. McIntyre, A. L. Angert, and K. J. Rice. 2009. Evolution and Ecology of
Species Range Limits. *Annual review of ecology, evolution, and systematics* 40: 415–436.

Shepard, I. D., S. A. Wissinger, Z. T. Wood, and H. S. Greig. 2022. Predators balance
consequences of climate-change-induced habitat shifts for range-shifting and resident
species. *The Journal of animal ecology* 91: 334–344.

Sheth, S. N., and A. L. Angert. 2018. Demographic compensation does not rescue populations at
a trailing range edge. *Proceedings of the National Academy of Sciences of the United States*
of America 115: 2413–2418.

Sloat, L. L., A. N. Henderson, C. Lamanna, and B. J. Enquist. 2015. The Effect of the
Foresummer Drought on Carbon Exchange in Subalpine Meadows. *Ecosystems* 18: 533–
545.

Smith, S. A., and J. W. Brown. 2018. Constructing a broadly inclusive seed plant phylogeny.
American journal of botany 105: 302–314.

Staude, I. R., D. M. Waller, M. Bernhardt-Römermann, A. D. Bjorkman, J. Brunet, P. De Frenne, R. Hédl, et al. 2020. Replacements of small- by large-ranged species scale up to diversity loss in Europe's temperate forest biome. *Nature ecology & evolution* 4: 802–808.

Tucker, C. M., M. W. Cadotte, S. B. Carvalho, T. J. Davies, S. Ferrier, S. A. Fritz, R. Grenyer, et al. 2017. A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biological reviews of the Cambridge Philosophical Society* 92: 698–715.

Tucker, C. M., M. W. Cadotte, T. J. Davies, and T. G. Rebelo. 2012. Incorporating geographical and evolutionary rarity into conservation prioritization. *Conservation biology: the journal of the Society for Conservation Biology* 26: 593–601.

Veldhuisen, L. N., B. J. Enquist, and K. M. Dlugosch. 2023. Phylogenetic diversity of flowering plants declines throughout the growing season across a subalpine elevational gradient. *bioRxiv*: 2023.11.06.565878.

Violle, C., W. Thuiller, N. Mouquet, F. Munoz, N. J. B. Kraft, M. W. Cadotte, S. W. Livingstone, and D. Mouillot. 2017. Functional rarity: The ecology of outliers. *Trends in ecology & evolution* 32: 356–367.

White, H. J., C. M. McKeon, R. J. Pakeman, and Y. M. Buckley. 2023. The contribution of geographically common and rare species to the spatial distribution of biodiversity. *Global ecology and biogeography: a journal of macroecology* 32: 1730–1747.

Willis, C. G., M. Halina, C. Lehman, P. B. Reich, A. Keen, S. McCarthy, and J. Cavender-Bares. 2009. Phylogenetic community structure in Minnesota oak savanna is influenced by spatial extent and environmental variation. *Ecography* 33: no–no.

Willis, C. G., B. Ruhfel, R. B. Primack, A. J. Miller-Rushing, and C. C. Davis. 2008. Phylogenetic patterns of species loss in Thoreau's woods are driven by climate change. *Proceedings of the National Academy of Sciences* 105: 17029–17033.

Yessoufou, K., and T. J. Davies. 2016. Reconsidering the Loss of Evolutionary History: How Does Non-random Extinction Prune the Tree-of-Life? *In* R. Pellens, and P. Grandcolas [eds.], Biodiversity Conservation and Phylogenetic Systematics: Preserving our evolutionary heritage in an extinction crisis, Topics in Biodiversity and Conservation, 57–80. Springer International Publishing, Cham.

Zacaï, A., E. Fara, A. Brayard, R. Laffont, J.-L. Dommergues, and C. Meister. 2017. Phylogenetic conservatism of species range size is the combined outcome of phylogeny and environmental stability. *Journal of biogeography* 44: 2451–2462.

Zavaleta, E. S., and K. B. Hulvey. 2004. Realistic species losses disproportionately reduce grassland resistance to biological invaders. *Science (New York, N.Y.)* 306: 1175–1177.

Zorio, S. D., C. F. Williams, and K. A. Aho. 2016. Sixty-Five Years of Change in Montane Plant Communities in Western Colorado, U.S.A. *Arctic, Antarctic, and Alpine Research* 48: 703–722.

FIGURES & CAPTIONS

Figure. 1: Rank abundance (A) and range size (B) curves for RMBL species at each site.

Figure 2: Phylogenies with log values of abundance (A) and range size (B) indicated by branch

colors. A) Abundance shows significant phylogenetic signal for Blomberg's K only (Blomberg's

K = 0.304, P-value = 0.04, Pagel's λ = 0.926, P-value =0.168), while B) range size does not

show any significant signal (Blomberg's K = 0.09, P-value = 0.7, Pagel's λ = 7.3×10^{-5}, P-value

=1).

Figure 3: Standard effect sizes to compare weighted and unweighted MPD and MNTD.

Unweighted values are in blue and weighted are in green. A) shows weighting by abundance. B)

shows weighting by range size. Each of the three panels in A) and B) show the different sites.

Weighting by abundance and range size does not significantly change MPD or MNTD in any of

the sites.

Figure 4: Change in SES values when individual species are removed in order of decreasing

abundance. The horizontal black line in each figure shows the standard effect size of the metric

for the overall community at each site, and dots show how the SES changes when the species at

the rank indicated on the x-axis is removed. MNTD values are shown in the top row, MPD

values are in the middle row, and PD are on the bottom. The left column shows the low elevation

site, middle column is the middle elevation site, and right is the high elevation site. There is no

clear pattern suggesting that removing rarer species (right side of each panel) decreases MNTD,

MPD or PD more than removing more abundant species (left side of each panel).

Figure 5: Change in SES values for PD when groups of ten species are removed in order of

decreasing abundance. The horizontal black line in each figure shows the standard effect size of

the metric for the overall community at each site, and dots show how the SES changes when the

group is removed. Numbers on the x-axis are the ranking of the most abundant species in the

group that was removed. For example, for the point at x=5, the species that were removed were

ranked 5-14. The left panel shows the low elevation site, middle panel is the middle elevation

site, and right is the high elevation site. There is no clear pattern suggesting that removing rarer

species (right side of each panel) decreases PD more than removing more abundant species (left

side of each panel).

A)

log(0.5)   trait value   log(573.5)

length=67.879

B)

log(108)   trait value   log(469,436)

length=67.879