Dominant-negative *TP53* mutations potentiated by the HSF1-regulated proteostasis network

Stephanie Halim[1,2], Rebecca M. Sebastian[1,2], Kristi E. Liivak[2], Jessica E. Patrick[2], Tiffani Hui[3], David R. Amici[4,5,6,7], Andrew O. Giacomelli[8], Paulina Rios[9,10], Vincent L. Butty[9,10,11], William C. Hahn[12,13], Francisco J. Sánchez-Rivera[9,10], Marc L. Mendillo[4,5,6], Yu-Shan Lin[3], Matthew D. Shoulders[2,9,13,14]

[1]Contributed equally to this work
[2]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA, USA
[3]Department of Chemistry, Tufts University, Medford, MA, USA
[4]Department of Biochemistry and Molecular Genetics, Northwestern University Feinberg School of Medicine, Chicago, IL, USA
[5]Simpson Querrey Institute for Epigenetics, Northwestern University Feinberg School of Medicine, Chicago, IL, USA
[6]Robert H. Lurie Comprehensive Cancer Center, Northwestern University Feinberg School of Medicine, Chicago, IL, USA
[7]Medical Scientist Training Program, Northwestern University Feinberg School of Medicine, Chicago, IL
[8]Humber Polytechnic, Toronto, ON, Canada
[9]Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology, Cambridge, MA, USA
[10]Department of Biology, Massachusetts Institute of Technology, Cambridge, MA, USA
[11]BioMicro Center, Massachusetts Institute of Technology, Cambridge, MA, USA
[12]Dana-Farber Cancer Institute, Boston, MA, USA
[13]Broad Institute of MIT and Harvard, Cambridge, MA, USA
[14]To whom correspondence may be addressed:
    Matthew D. Shoulders
    Department of Chemistry
    Massachusetts Institute of Technology
    77 Massachusetts Avenue, 16-573A
    Cambridge, MA 02139
    Phone: (617)452-3525
    Email: mshoulde@mit.edu

ABSTRACT

Protein mutational landscapes are sculpted by the impacts of the resulting amino acid substitutions on protein stability and folding or aggregation kinetics. These properties can, in turn, be modulated by the composition and activities of the cellular proteostasis network. Heat shock factor 1 (HSF1) is the master regulator of the cytosolic and nuclear proteostasis networks, dynamically tuning the expression chaperones and quality control factors in these cellular compartments to meet demand. Chronic increases in HSF1 levels and activity are prominent hallmarks of cancer cells, for reasons that remain poorly understood. One plausible explanation is that the consequent upregulation of proteostasis factors could biophysically facilitate the acquisition, selection, and maintenance of oncogenic mutations. Here, we experimentally evaluate the impacts of chronic HSF1 activation on the mutational landscape accessible to p53, the protein encoded by the most mutated gene in human cancers. Specifically, we apply quantitative deep mutational scanning of p53 to assess how HSF1 activation shapes the mutational trajectories by which p53 can escape cytotoxic pressure conferred by the small molecule nutlin-3, a potent antagonist of the p53 negative regulator MDM2. We find that HSF1 activation broadly increases the selection of dominant-negative substitutions within p53. This effect of HSF1 activation was particularly notable for non-conservative, biophysically unfavorable amino acid substitutions within buried regions of the p53 DNA-binding domain. These results indicate that chronic HSF1 activation shapes the oncogenic mutational landscape, preferentially supporting the acquisition and maintenance of cancer-associated substitutions that are biophysically disruptive.

INTRODUCTION

Cancers typically arise by multi-step acquisition of genetic alterations that dysregulate cell growth and survival programs, leading to malignant phenotypes[1-3]. In addition to mutated genes, cancer cells also co-opt non-mutated genes within stress pathways to aid cell growth, in a process known as non-oncogene addiction[4, 5]. Particularly noteworthy in this regard, pathways associated with maintenance of proteostasis are often upregulated in cancer, likely owing to challenges arising from dysregulated protein synthesis, nutrient starvation, subunit imbalance within protein complexes, and perhaps, expression of oncoproteins with destabilizing amino acid substitutions[6-11]. Most commonly, the heat shock response (HSR), which is controlled by the master transcription factor HSF1, is frequently upregulated in cancer[12, 13]. HSF1 can also modulate additional cell remodeling programs in tumors[14-16]. High basal expression of heat shock proteins regulated by HSF1 is widely observed in malignant cells[13], as are overexpression and constitutive activity of HSF1 itself[13-15, 17]. HSF1 supports the emergence of tumors in mice following exposure to mutagens, and high levels of HSF1 expression are associated with increased mortality rates in breast cancer[16, 17]. Additionally, HSF1 and the HSF1-regulated chaperone heat shock protein 90 (HSP90) can facilitate the development of resistance to chemotherapeutic agents, although molecular-level mechanisms of this phenomenon are unclear[18, 19]. Accordingly, much attention has been drawn not just to HSP90[20] and other chaperones but also to HSF1 as a potential chemotherapeutic target, motivating the development of potent small molecules that target HSR components either specifically through inhibition of individual chaperones or broadly through targeting HSF1[21, 22].

One intriguing hypothesis to explain how these phenomena benefit cancers is that an enhanced proteostasis environment amplifies the accessibility of novel mutations in cancer-associated proteins via its influence on the folding and degradation of the resulting protein variants. Successful emergence of gain-of-function mutations in all proteins is constrained by the biophysical properties of the evolving protein, particularly as most functionally important amino acid substitutions are biochemically non-conservative, and therefore, often biophysically deleterious[23, 24]. A growing body of literature has investigated the impact of proteostasis network components, such as HSP90 and HSP70, on protein evolution[23, 25-32]. Among other advances, such studies have shown that proteostasis network remodeling mediated by stress-responsive transcription factors can have major impacts on the mutational space accessible to viral pathogens that parasitize host chaperones[28-31], and that these effects are often mediated directly by the influence of proteostasis network composition on client protein folding and stability[23, 26, 30, 33].

Building on these studies, one compelling possibility regarding the roles of HSF1 in cancer is that HSF1 overexpression supports malignancy by creating a permissive protein folding environment that directly facilitates the emergence of proliferation-promoting oncogenic mutations. However, a role for HSF1 in defining accessible oncogene mutational space has never been experimentally explored.

The tumor suppressor p53 is a key transcription factor and regulator of the response to DNA damage and other oncogenic stimuli. When activated, p53 induces cell-cycle arrest, senescence, or apoptosis[34]. *TP53*, which encodes the p53 protein, is the most frequently mutated gene in cancer[35]. The majority of these mutations are missense mutations within p53's DNA-binding domain[36]. While some *TP53* missense mutations confer a loss-of-function effect typical of tumor suppressors, mutations in *TP53* can also function via a dominant-negative effect. In such cases, the mutated *TP53* allele encodes a protein variant that inhibits the function of residual wild-type p53, possibly through either heterotetramerization via a C-terminal oligomerization domain or by inducing co-aggregation and consequent loss of function[37-39].

p53 interacts extensively with the HSF1-regulated chaperones HSP90 and HSP70, which in turn help to regulate the stability and activity of p53[40-44]. Notably, while wild-type p53 engages transiently with chaperones, particularly during folding and maturation, several p53 variants display more stable interactions with HSP70 and HSP90, contributing to increased variant p53 levels[45, 46]. Considering the large number of destabilizing p53 substitutions and the extensive interactions between p53 and HSF1-regulated chaperones, p53 represents a compelling model system to explore the impacts of HSF1 activation on the evolution of the cancer-associated proteome.

Here, we apply deep mutational scanning (DMS) of p53 in concert with chemical genetic regulation of HSF1 to examine the impacts of proteostasis modulation on the fitness of dominant-negative p53 variants. We observed that constitutive activation of HSF1 broadly increases the fitness of diverse dominant-negative p53 variants, including several substitutions within hot-spot sites frequently associated with cancer. The impact of HSF1 activation on p53's mutational spectrum was most evident in substitutions of hydrophobic amino acids from nonpolar to polar amino acids within buried regions of the DNA-binding domain of p53. These substitutions are also most likely to be destabilizing, aggregation-inducing, or folding-disruptive, consistent with the notion that HSF1 activity increases the fitness of mutations that perturb p53 stability or folding. Altogether, these results indicate that HSF1 can directly potentiate oncogene evolution by opening access to otherwise biophysically problematic amino acid sequences. Further, these results implicate proteostasis network inhibition as a potential therapeutic strategy to prevent acquisition of resistance mutations during chemotherapy.

RESULTS

TP53 mutational library integrated with chemical genetic regulation of the HSR

To assess the impact of HSF1 on the mutational landscape of dominant-negative p53, we first needed to establish a system in which HSF1 activity could be robustly regulated in an appropriate cell line. We chose to use A549 cells, which are an epithelial tumor model derived from a human alveolar basal cell adenocarcinoma[47]. A549 cells exclusively express wild-type p53, critically enabling our experimental workflow[48]. Pharmacologic activation of HSF1 is traditionally achieved via treatment with HSP90 or HSP70 inhibitors or toxins like arsenite[49]. Such approaches are not useful here, as they activate HSF1 indirectly by causing massive cellular protein misfolding, and ultimately drive apoptosis via proteostatic overload. Instead, we employed regulated expression of a constitutively active HSF1 variant, termed cHSF1. We constructed a stable, single-colony A549 cell line in which cHSF1 expression was placed under the control of a doxycycline (dox)-responsive promoter[50-53]. In these cells, termed A549[cHSF1] cells, treatment with dox activates expression of cHSF1, which then proceeds to upregulate HSR-controlled gene expression independent of protein misfolding stress. Thus, A549[cHSF1] cells couple inducible expression of cHSF1, via dox treatment, to induction of the HSR.

To test the activation of HSF1, we treated A549[cHSF1] cells with dox or vehicle for 24 h and then evaluated the transcript levels of established HSF1 target genes using qPCR. In our optimized cell line, we observed modest upregulation of *DNAJB1* and *HSPA1A* transcripts during HSF1 activation as compared to vehicle treatment, indicating that our cHSF1 construct was functionally modulating the HSR as intended (**Figure 1A**). Critically, we obtained cells in which HSF1 induction upregulated HSR genes to levels that are still well within the physiologically accessible regime, a feature that is known to be critical to avoid off-target induction of genes not normally targeted by HSF1[52, 54]. As evidence, treatment with STA-9090[55], an HSP90 inhibitor and robust activator of endogenous HSF1, had considerably stronger effects than the dox treatment (**Figure 1A**). Moreover, a resazurin metabolic activity assay[56] indicated that cell growth and viability were not altered by dox-mediated HSF1 induction (**Figure S1A**).

We next aimed to comprehensively assess how HSF1 activation remodeled the proteostasis network. We treated A549[cHSF1] cells with dox for 24 h and quantified differentially transcribed genes using RNA-seq. 159 transcripts were significantly and differentially expressed with >1.5 $\log_2$fold-change upon HSF1 activation as compared to the vehicle-treated control, highlighting that HSF1 activation did not massively perturb the global transcriptome (**Table S1**). Instead, known components of the HSR, including *HSP90AA1* and *HSPA1A*, were highly enriched among the upregulated transcripts (**Figure 1B**). Gene set enrichment analysis using the MSigDB c5 collection[57] further confirmed that genes related to

the HSR were highly enriched following activation of HSF1 (**Table S2**). Notably, known p53-induced genes, such as *BAX* and *CDKN1A*, were not substantially impacted by HSF1 activation (**Table S1**).

We next asked to what extent HSF1-regulated chaperones might directly alter p53 proteostasis. To address this question, we identified the subset of transcripts encoding HSF1-regulated chaperones that are known to interact with p53. Of the 159 transcripts significantly upregulated by HSF1 activation, 22 (13.8%) encode proteins classified by the APID (Agile Protein Interaction DataAnalyzer) database as interacting with p53 (**Table S3**)[58]. Included within these p53-interacting proteins, 17 are either chaperones or co-chaperones, including the well-validated and HSF1-regulated p53 chaperones HSP90 and HSP70 (**Figure 1C**).

We transduced these A549[cHSF1] cells with a high-quality, lentiviral-based *TP53* mutational library in which all possible single amino acid substitutions across the p53 protein are present[37, 59]. We performed the transduction at a low multiplicity of infection (MOI = 0.25) to ensure that a single p53 variant was expressed in each cell, _alongside_ endogenous wild-type p53 (**Figure 1D**). To evaluate the diversity of the resulting A549[cHSF1] p53 variant library, which we termed A549[cHSF1](p53-Lib) cells, we harvested genomic DNA from the A549[cHSF1](p53-Lib) cells, then amplified the library-encoded *TP*53 gene using PCR before deep-sequencing. Out of 7467 potential missense mutations generated from 19 possible amino acid substitutions at 393 sites, 7465 (99.9%) of the possible substitutions were observed with a read depth >10 counts per amino acid substitution (**Figure S1B** and **Table S4**).

HSF1 activation modulates the p53 mutational landscape during MDM2 inhibition

We next sought to apply DMS to test whether HSF1 activity alters tolerance for p53 mutations that promote escape from nutlin-3 selection. Our approach was to perform selections in the HSF1-enhanced or the basal proteostasis environment in the presence of nutlin-3, a p53-activating agent[59]. Nutlin-3 inhibits the interaction between p53 and mouse double minute 2 homolog (MDM2), an E3 ubiquitin ligase and negative regulator of p53[60]. In unstressed cells, MDM2 binds to p53 to trigger ubiquitination and degradation. Binding of nutlin-3 to MDM2 releases p53 from the MDM2 complex, allowing p53 to accumulate and induce a transcriptional program that drives apoptosis and cell cycle arrest. Cells that express a dominant-negative p53 variant alongside endogenous wild-type p53 are strongly positively selected in the presence of nutlin-3, because dominant-negative p53 attenuates the wild-type p53-mediated activation of cell cycle arrest and thereby allows cells to continue to proliferate (**Figure 2A**)[59].

We treated A549[cHSF1](p53-Lib) cells with nutlin-3 (or vehicle) in the context of either a basal or an HSF1-activated proteostasis environment, allowing the selection to proceed under these conditions over a 12-day period[59]. After selection, we followed a published protocol to deep-sequence the library-

encoded *TP53* amplicons[59]. We next calculated the resulting changes in p53 variant frequency as the log$_2$ fold-change in normalized read counts of amino acid substitutions between our various selection versus control conditions. The resulting mutational log$_2$ fold-change and site-level log$_2$ fold-change values are available in **Tables S5** and **S6**.

We expected that nutlin-3 would be the dominant force affecting the fitness of p53 variants, as cells that fail to express a dominant-negative p53 variant cannot survive nutlin-3 selection. Consistent with this expectation, we observed a very strong and positive enrichment of missense mutations within the DNA-binding domain of p53, located from residues 100–300, upon nutlin-3 selection in both the basal (**Figure S2A**) and the HSF1-activated (**Figure S2B**) proteostasis environments relative to the corresponding vehicle-treated (no nutlin-3) controls. Importantly, the sites enriched during this nutlin-3 selection in both the basal and HSF1-activated environments overlapped with previous selections for dominant-negative p53 using saturation mutagenesis in both the A549 and AML cell lines[37, 59]. Moreover, the fitness of p53 variants identified as somatic mutations within the *TP53* database[61, 62] was higher as compared to all missense mutations observed in the DMS experiment, confirming that nutlin-3 was specifically selecting for cancer-associated, dominant-negative *TP53* mutations (**Figures S2C** and **S2D**). We also observed an exceptionally high degree of correlation for the cumulative site log$_2$ fold-change values between biological replicates of nutlin-3 treatment versus vehicle under both basal and HSF1 activated proteostasis environments (**Figures S3A** and **S3B**). This observation further indicates that nutlin-3 imposes a very strong selection pressure.

Next, to isolate the impact of HSF1 activation on dominant-negative p53 mutational tolerance, we evaluated whether and how HSF1 activation affected p53 variant fitness specifically during MDM2 inhibition. We observed an increase in p53 variant fitness across much of the *TP53* gene during HSF1 activation (**Figure 2B**). The correlation between individual replicates, shown in **Figure S4A**, was positive and highly significant, indicating the results were reproducible. That said, while reasonable for a DMS experiment[29-31, 63], the correlation was not as strong as the correlation observed for nutlin-3 treatment versus control (**Figures S3A** and **S3B**), an observation that is consistent with the exceptionally strong selection pressure imposed by MDM2 inhibition itself.

The observation that chronic HSF1 activation, which is commonly observed across diverse cancers[12-17], broadly increased p53 mutational tolerance (**Figure 2B**) motivated us to analyze the underlying effect in greater depth. We first subdivided variants by domain and examined the distribution of the net site log$_2$ fold-change and mutational log$_2$ fold-change for mutations that encoded amino acid substitutions within each domain. In all functional domains examined, we observed a significant and often

substantial increase in net site fitness (**Figure 2C**). Across these domains we observed an HSF1-dependent increase in mutational fitness for missense mutations only (**Figure 2D**). HSF1 had no significant impact on synonymous mutations (**Figure 2D**), suggesting that the effects of HSF1 are not arising from a general increase in cell fitness. Also noteworthy, HSF1 had minimal effects on p53 variant enrichment in the absence of nutlin-3 treatment, which provides the underlying need for p53 variant selection with MDM2 inhibition (**Figure S4B**). In sum, HSF1 activation specifically enhanced the fitness of non-synonymous dominant-negative *TP53* mutations under MDM2 inhibition, while not impacting synonymous mutations in all domains of p53.

HSF1 activation supports the accumulation of biophysically destabilizing, non-conservative amino acid substitutions within buried regions of the p53 DNA binding domain

We next examined the impacts of HSF1 activation during MDM2 inhibition on variant fitness within the DNA-binding domain of p53 (**Figures 3** and **S5**), because the majority of known oncogenic *TP53* mutations are localized to this region[61]. Moreover, the DNA-binding domain is the only structurally characterized region of p53, enabling further analyses. The DNA-binding domain of p53 is composed of a core β-sandwich scaffold supporting a DNA-binding surface comprising two loops (Loop 2 and Loop 3) stabilized by Zn^{2+} coordination, as well as a loop-sheet-helix motif that contains Loop 1[64]. The impact of HSF1 was seen most in sites 238–249 within Loop 3, as well as at sites V173 and H179 within Loop 2 (**Figure S5**). Several cancer hot-spots are localized in this region, including sites G245, R248, and R249, all of which displayed a net increase in fitness upon HSF1 activation. These observations indicate that HSF1 can directly support the acquisition of hot-spot mutations in p53 that lead to malignant transformation.

Relatively fewer p53 sites displayed an overall net negative fitness upon HSF1 activation during MDM2 inhibition. Interestingly, several of the few sites with net negative fitness (V172, R174, and T211; **Figures 3B** and **3C**) are localized at the surface of a pocket that interacts with the N-terminal tail of the DNA-binding domain. Interaction of the N-terminal tail of the DNA-binding domain with residues in this pocket has been shown to both increase p53 thermodynamic stability as well as decrease aggregation propensity[65]. A possible explanation for the decrease in fitness at these specific sites is that variants within this region increase the propensity for mutant p53 to co-aggregate with wild-type p53, with resultant dominant-negative consequences favorable in nutlin-3 selection. Such destabilization and aggregation may be attenuated in the supportive proteostasis environment created by HSF1 activation, leading to their relative decrease in fitness during simultaneous HSF1 activation and MDM2 inhibition.

Prototypical oncogenic p53 amino acid substitutions are classed into either DNA-contact variants with minimal impact on thermodynamic stability or structural mutations that significantly perturb the

stability of the DNA-binding domain[66]. Mutations within this second class are frequently localized to either the Zn^{2+}-binding site, or within the β-sandwich motif at the hydrophobic core of the DNA-binding domain, prompting us to analyze these sites in more detail. In particular, Zn^{2+} coordination is critical for p53's ability to bind DNA, and loss of Zn^{2+} binding is associated with destabilization and aggregation of p53[66-68]. We examined the fitness of amino acid substitutions within the Zn^{2+} coordination sites (C176 and H179 of Loop 2 and C238 and C242 of Loop 3; **Figures 3B** and **3C**). As with other cancer-associated mutations, we observed that the net site fitness for all four coordinating residues increased following HSF1 activation (**Figure 3C**).

We then turned our attention to sites within the β-sandwich motif and at the hydrophobic core of the DNA-binding domain where substitutions could be particularly biophysically disruptive. We first tested whether there was a correlation between relative solvent accessibility (RSA; **Table S7**) and net site fitness within the p53 DNA binding domain as a consequence of nutlin-3-mediated p53 selection regardless of the proteostasis environment. Indeed, when we examined the RSA of sites during nutlin-3 selection versus vehicle treatment in either the basal or in the HSF1-activated proteostasis environment, we observed a negative correlation between net site fitness and RSA (**Figures S6A–D**). This observation coincides with the expectation that substitutions to buried residues within the p53 DNA binding domain are likely to elicit dominant-negative behavior and merit further scrutiny.

Accordingly, we asked whether HSF1 activation during MDM2 inhibition preferentially impacted the fitness of p53 amino acid substitutions within buried regions of the p53 DNA-binding domain. We noted that variants classified as buried (RSA < 0.2) displayed marginally higher net fitness as compared to sites that were exposed (RSA > 0.2; **Figure S6E**). Motivated by this intriguing result, we clustered variants into three classes based on their biochemical properties:

(a) Conservative substitutions involving replacement of a nonpolar amino acid for another nonpolar amino acid or a polar amino acid for another polar amino acid.

(b) Non-conservative substitutions involving a switch from a nonpolar to a polar amino acid.

(c) Non-conservative substitutions involving a switch from a polar to a nonpolar amino acid.

Then, we analyzed whether HSF1 activation impacted non-conservative substitutions more strongly than conservative substitutions. In exposed regions of p53 DNA-binding domain, there was no significant effect in any mutation class upon HSF1 activation. Strikingly, however, non-conservative substitutions replacing a nonpolar amino acid with a charged or polar amino acid within buried p53 sites

displayed substantially and significantly higher fitness upon HSF1 activation relative to either conservative substitutions or non-conservative substitutions from polar to nonpolar amino acids in buried regions (**Figure 4A**). These results suggest that HSF1 activation particularly enables the p53 protein to more robustly access non-conservative, biophysically disruptive substitutions — especially the introduction of polar residues in hydrophobic, buried domains.

To further explore this phenomenon, we examined 42 missense mutations in the p53 DNA-binding domain for which thermodynamic stability has been experimentally determined[66, 69-71]. We grouped the variants into those that were stabilizing ($\Delta\Delta G$ < 0.125 kcal/mol) or destabilizing ($\Delta\Delta G$ > 0.125 kcal/mol). We observed a significant increase in fitness for mutations that were thermodynamically destabilizing as compared to stabilizing mutations (**Figure S7A**).

Since thermodynamic stability measurements have been experimentally performed for only a very limited number of p53 DNA-binding domain variants, we built on the result observed in **Figure S7A** by using Rosetta to estimate the change in thermodynamic stability for all possible amino acid substitutions within the DNA-binding domain (**Table S8**)[72]. To assess the accuracy of the calculated $\Delta\Delta G$ values, we first compared the computed to the experimentally determined $\Delta\Delta G$ values for the aforementioned 42 amino acid substitutions. We observed a strong correlation between the two approaches (**Figure S7B**), validating the computational approach.

As expected, our computations predicted that substitutions from nonpolar to polar amino acids in buried regions have by far the greatest impact on p53 stability in buried regions of the DNA-binding domain (**Figure 4B**). The trend observed in **Figure 4B** mirrored the observed trend discussed in **Figure 4A**, namely that HSF1 activation impacted non-conservative nonpolar to polar substitutions in buried regions of the DNA-binding domain of p53 the most. We next assessed whether and how HSF1 activation preferentially impacted thermodynamically destabilizing amino acid substitutions in the p53 DNA-binding domain. First, to evaluate whether there was a difference in estimated thermodynamic stability depending on the direction and magnitude of their change in fitness with HSF1 activation in a nutlin-3 environment we binned variants by their mutational fold-change HSF1 activation and nutlin-3 versus nutlin-3 alone selection, and then plotted the predicted $\Delta\Delta G$. In buried regions of the DNA-binding domain of p53, we observed that substitutions with enhanced fitness upon HSF1 activation were more likely to display a higher estimated $\Delta\Delta G$ than variants that displayed decreased fitness (**Figure 4C**). In contrast, we observed little difference in the effect of HSF1 activation on destabilized versus stabilized variants for exposed sites of the DNA-binding domain of p53 (**Figure 4C**). In sum, HSF1 activation specifically supports the emergence of destabilizing p53 substitutions during MDM2 inhibition.

HSF1-potentiation of destabilizing and oncogenic dominant-negative p53 variants is reproducible in head-to-head competition assays and generalizable across cancer cell lines

With the key conclusion from our large-scale DMS data established, we pursued detailed studies on a few specific p53 variants to better understand and explore these phenomena. In particular, we studied two p53 variants (V173Y and F113K) in buried regions of the p53 DNA-binding domain that were predicted to be biophysically destabilizing ($\Delta\Delta G$ > 5 kcal/mol) and representing the nonpolar-to-polar substitutions we found to be most potentiated by HSF1. F113K in particular is also charged in addition to being polar (**Figure 5A**). We also studied a prototypical oncogenic, dominant-negative p53 substitution commonly found in cancer patients (R273H). p53[R273H] displayed enhanced fitness upon HSF1 activation in our DMS experiments, albeit with a smaller effect size than the F113K or V173Y substitutions.

We first conducted qPCR assays to assess the dominant-negative activity of these p53 variants in an HSF1-activated environment. We observed strong suppression of the transcript levels of p53 target genes, *BAX* and *CDKN1A* (p21), when these variants were co-expressed with wild-type p53 in an HSF1-activated environment alongside MDM2 inhibition via nutlin-3 treatment (**Figure 5B**).

With dominant-negative behavior demonstrated, to test our results in a focused biochemical experiment we performed a pairwise competition experiment for cells expressing only wild-type p53 versus cells expressing both wild-type p53 and each individual p53 variant under the relevant conditions used in the DMS experiment. Briefly, we transduced A549[cHSF1] cells with lentivirus encoding either wild-type or variant p53 along with a corresponding fluorescent marker: BFP for wild-type p53 or GFP for p53 variants (**Figure 5C**). These markers enabled fluorescent readout of the abundance of each variant in a pre-competition mixture of the two cell lines, as well as 72–96 h post-treatment with either nutlin-3 alone or nutlin-3 and dox to induce HSF1. We observed enrichment of the p53 variants relative to wild-type upon nutlin-3 treatment, confirming that the variants are indeed exerting a dominant-negative effect (**Figure 5D**)[59, 60]. Upon HSF1 activation, we observed an additional, highly significant enrichment of these same variants (**Figure 5E**), validating the DMS results in A549[cHSF1] cells for individual variants. Notably, the common oncogenic p53 variant R273H showed an even larger effect size than the other variants in the head-to-head competition, suggesting that our A549[cHSF1] DMS results may underestimate the importance of HSF1 activity in potentiating common oncogenic p53 substitutions.

To assess whether the fitness-enhancing consequences of HSF1 activity for dominant-negative p53 substitutions are generalizable beyond A549 cells, we pursued a similar set of competition experiments in U2OS cells, an osteosarcoma cell line[73]. Importantly, U2OS cells, like A549 cells, are a cancer line that genomically encodes wild-type p53[74]. As with our A549 cells, we engineered U2OS[cHSF1] cells

with dox-regulated induction of HSF1 activity (**Figure S8**). We then introduced the same p53 variants (**Figure 5A**) and repeated the competition experiments shown in **Figure 5C** using the resulting U2OS[cHSF1] cell lines. Once again, we observed enrichment of the p53 variants relative to wild-type p53 upon nutlin-3 treatment, consistent with these variants exerting a dominant-negative effect (**Figure 5F**) in U2OS cells. This enrichment was strongly potentiated upon HSF1 activation, with even larger effect sizes than in A549[cHSF1] cells (**Figure 5G**). Thus, the effects of HSF1 activity on p53 variant fitness are generalizable beyond just one cancer model.

HSF1-potentiation of dominant-negative p53 substitutions is associated with aggregation modulation and likely involves the activities of HSF1-induced chaperones

Amino acid substitutions that induce mild structural defects within the p53 DNA-binding domain are likely to function through tetramerization with wild-type p53, where the presence of one or more non-functional p53 subunits reduces the ability of the overall complex to bind to DNA and induce transcription[38, 75]. For such non-functional p53 subunits to retain dominant-negative function, they must be soluble and not aggregated in cells. Interestingly along these lines, confocal microscopy analysis of A549[cHSF1] cells co-expressing wild-type p53 and the V173Y variant revealed visible p53 aggregates in the nucleus under basal conditions (**Figure 6**, top row). HSF1 activation resulted in reduced aggregation and even total clearance of aggregation in some cells, suggesting that HSF1 may act to prevent aggregation-inducing conformations of destabilized p53 variants in a manner that promotes dominant-negative function via tetramerization (**Figure 6**, middle row).

We sought to better understand the underlying forces shaping HSF1 potentiation of dominant-negative p53 variants. Specifically, we hypothesized that inhibition of key p53-interacting chaperones induced by HSF1 might strongly reduce the fitness of HSF1-potentiated p53 variants in a nutlin-3 environment. We performed pairwise competition assays with the V173Y, F113K, and R273H p53 variants identified from the DMS screen in the presence of either an HSP90 inhibitor (STA-9090) or an HSP70 inhibitor (VER-155008) in an MDM2-inhibited environment[55, 76]. Based on the potentiation of these variants by HSF1 activation, we hypothesized that they would perform more poorly upon chaperone inhibition. Indeed, we observed a decrease in p53 mutant fitness in the presence of these chaperone inhibitors (**Figures 7A** and **7B**). We next asked if the reason for the loss in fitness could be related to excessive mutant p53 protein aggregation. After treating V173Y p53-expressing cells for 24 h with STA-9090 to inhibit HSP90, we again visualized p53 state using confocal microscopy. We observed increased V173Y p53 aggregation upon HSP90 inhibition (**Figure 6**, bottom row), the opposite phenotype of the

reduced aggregation observed for V173Y p53 upon HSF1 activation. Taken together, these observations support the idea that chaperone networks modulate p53 variant fitness by tuning protein stability and aggregation propensity.

HSF1-potentiated p53 variants are enriched in patients

We questioned whether HSF1 activation impacts p53 mutations observed in cancer patients. We first analyzed distributions of the effects of HSF1, represented by the mutational $\log_2$ fold-change of HSF1 and nutlin-3 versus nutlin-3 alone, for all possible missense mutations from our DMS data to variants observed in cancer patients in the *TP53* database[61, 62]. We compared the mutational fold-change of nutlin-3 and HSF1 versus nutlin-3 alone of all missense mutations in the DMS and missense mutations with and without the dominant-negative loss of function (DNE_LOF) annotation in the TP53 database (**Figure 8A**). We observed that missense mutations annotated as DNE_LOF were significantly impacted by HSF1 activation in a nutlin-3 environment, while missense mutations without the DNE_LOF annotation were not potentiated by HSF1. This comparison suggests that HSF1-potentiated dominant-negative p53 variants are indeed enriched in the patient population.

We next asked whether HSF1 activity levels might be higher in patients who carry an HSF1-potentiated p53 substitution. We defined "HSF1-potentiated variants" as p53 variants with increased fitness in an HSF1 activated and nutlin-3-treated environment, specifically where the mutational $\log_2$ fold-change of HSF1 and nutlin-3 versus nutlin-3 alone was >0.05. Using the TCGA PanCancer Atlas of lung adenocarcinoma available in cBioPortal[77-79], we found that expression of sentinel genes that report on HSF1 activity in patients with wild-type p53 was significantly lower than HSF1 sentinel gene activity observed in patients carrying a p53 variant potentiated by HSF1 activity in our DMS studies (**Figure 8B**). Together, these observations support the notion that HSF1 is a fundamental force shaping p53 mutational landscapes in human cancers.

DISCUSSION

This study shows that HSF1 activity increases the fitness of p53 dominant-negative variants that can drive cancer. Moreover, our analyses of p53's structured DNA-binding domain show that the p53 mutational fitness-enhancing impact of HSF1 activation is strongly biased towards supporting the emergence of thermodynamically destabilizing substitutions located within buried protein regions. In particular, HSF1 upregulation improves the fitness of non-conservative substitutions where buried nonpolar amino acids are replaced by polar or charged amino acids. This effect is observed across multiple cancer models. RNA-seq analysis shows that HSF1-activated genes are enriched in p53-interacting chaperones, supporting the hypothesis that HSF1 directly impacts the fitness of p53 variants through modulation of folding, aggregation, degradation, or stability of mutant p53 via HSF1-regulated chaperones. Visualization of a destabilizing p53 variant within cells under basal conditions, HSF1 activation, and HSP90 inhibition is consistent with the notion that HSF1 activation reduces aggregation of otherwise unstable p53 variants. Thus, our data are consistent with the notion that HSF1 over-activation may overcome wild-type p53 suppression to restore the dominant-negative function of mutant p53, potentially amplifying the feed-forward circuit of HSF1 activation and continued oncogenic p53 stabilization. Finally, using human cancer patient data, we find that these phenomena may be relevant in a clinical setting.

The work presented here is, to our knowledge, the first to show that HSF1 can directly shape the mutational space accessible for malignant transformation. These results have several interesting implications. First, while the literature has clearly highlighted HSF1 activation as an oncogenic helper[7, 80, 81], it is yet unclear whether the commonly observed increase in HSF1 expression and activity occurs prior to the onset of tumorigenesis, or whether constitutive HSF1 activation is a consequence of proteome instability arising from accumulated mutations and genomic damage. Our results suggest that activation of HSF1 can facilitate early tumorigenic events by increasing the fitness of some oncogenic driver mutations. Second, resistance to chemo-therapeutic agents is frequently driven by mutations within the targeted proteins. While both HSF1 upregulation as well as downstream chaperones have been identified as facilitators of chemo-resistance, the mechanism has primarily been associated with alterations in metabolic or autophagy pathways. Further studies on HSF1 targets that interact with p53 should be pursued to better understand the interplay between chaperone networks and oncogenesis [18, 19, 82, 83]. Our results suggest that HSF1 activity may facilitate chemo-resistance by tuning the accessibility of resistance mutations. Given the recent therapeutic interest in inhibitors of both HSF1 itself and

HSF1-regulated chaperones such as HSP90 and HSP70[21, 22, 51, 84], these results indicate that such inhibitors may be particularly effective in combination therapy to prolong effective outcomes by reducing the emergence of resistant tumor populations.

The capacity of HSF1 or other types of proteostasis network upregulation to tune the stability, folding, and/or aggregation of oncoproteins could have tremendous implications for cancer immunotherapy. Variants with enhanced stability may not be able to be degraded, and thus, unable to be efficiently presented to the immune system by MHC-1 proteins. In accordance with our findings and published work in this field, we reason that downregulation of proteostasis networks may enable a more robust immune response to tumor presenting antigens[85].

We emphasize that the impacts of proteostasis network modulation on oncoprotein mutational spectra are likely to extend far beyond just p53. This study should motivate efforts to more fully understand how HSF1 shapes the mutational spectra of additional oncoproteins and chemotherapy targets, and the potential of modulating chaperone networks or even HSF1 itself as a chemotherapeutic target.

LIMITATIONS OF STUDY

While we observed a loss of fitness of HSF1-potentiated p53 variants with chaperone inhibition, HSP90 and HSP70 inhibition can also induce compensatory HSF1 activation[51], and untangling the relative effects of chaperone activity versus HSF1 activity is challenging. Considering these competing phenomena would likely require a comprehensive understanding of the effects of individual chaperones on individual p53 substitutions. Related to this point, while we focused our follow-up experiments on sentinel HSF1-targeted genes, namely HSP70 and HSP90, other HSF1-induced p53 interactors may also play a role in the relative fitness of certain p53 variants. These effects could include direct interactions with p53 or secondary effects due to remodeling of the proteostasis environment and would also require targeted mechanistic work on individual p53 variants.

A second caveat to consider is that the DMS library only contains single amino acid substitutions in the p53 protein. Large deletions, multiple missense mutations, or copy number variations, which can all impact p53 biology and play a role in tumorigenesis, were not considered in our work. There is reason to believe that HSF1 may play a role in such cases due to the proteostasis imbalances caused by such p53 variants, and this phenomenon should be further studied.

Finally, while our study provides evidence for roles of HSF1 activity in oncogenic mutational spectra across multiple cell lines, we are still limited by the model systems employed. The work here should motivate related *in vivo* studies in the future.

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health (1R35GM136354 to M.D.S.), (1R01AI168166 to Y.-S.L), MIT HEALS (to M.D.S.), and by an American Cancer Society–Ellison Foundation Research Scholar Award (to M.D.S.). Additional support was provided by Koch Institute Support (core) under NIH (core) grant P30-CA14051 from the NIH/NCI and by the MIT CEHS core via the NIH/NIEHS (Grant P30-ES002109). Work in the Sánchez-Rivera laboratory is supported by the Howard Hughes Medical Institute (Hanna Gray Fellowship, GT15656), V Foundation for Cancer Research (V2022-028), NCI 1P01CA291694-01A1, Virginia and D.K. Ludwig Fund for Cancer Research, MIT HEALS Initiative, Koch Institute Frontier Research Program, Casey and Family Foundation Cancer Research Fund, Michael (1957) and Inara Erdei Fund, MIT Research Support Committee, Upstage Lung Cancer Foundation, and a Traditional Project Award from the Bridge Project, a partnership between the Koch Institute for Integrative Cancer Research at MIT and the Dana-Farber/Harvard Cancer Center. P.R. was supported by a National Science Foundation GRFP Award.

DECLARATION OF INTERESTS

F.J.S.R. has consulted for Repare Therapeutics, Ono Pharma, and Merck. W.C.H. is a consultant for Thermo Fischer Scientific, Solasta Ventures, KSQ Therapeutics, Frontier Medicines, Jubilant Therapeutics, RAPPTA Therapeutics, Serinus Biosciences, Kestral Therapeutics, Function Oncology, Crane Biotherapeutics and Perceptive. A.O.G. is a consultant for Atlas Venture.

AUTHOR CONTRIBUTIONS

S.H, R.M.S., K.E.L, T.H., A.O.G., Y.-S.L., and M.D.S. designed experiments. S.H, K.E.L, R.M.S., and T.H. conducted experiments. R.M.S. and M.D.S. conceived the project. A.O.G. and W.C.H. provided key reagents. All authors analyzed results and contributed to writing and/or editing the manuscript. Y.-S.L. and M.D.S. supervised the research.

MATERIALS AVAILABILITY

Sequencing data have been deposited at SRA as SRA: PRJNA1178283 and are publicly available as of the date of publication. RNA-seq data have been deposited at GEO as GEO:GSE304320 and are publicly available as of the date of publication.

STAR Methods

METHOD DETAILS

Cell culture. A549 cells were a kind gift from the Prof. William Hahn Lab at Harvard Medical School. U2OS cells were purchased from ATCC. Cells were grown in DMEM medium (Corning), supplemented with 10% heat-inactivated fetal bovine serum (FBS, Cellgro) and 1% penicillin/streptomycin/glutamine (Cellgro) at 37 °C with 5% CO_2(g).

Plasmids. To create stable A549[cHSF1] and U2OS[cHSF1] cell lines, cHSF1[51] was cloned into the pIN-DUCER20 lentiviral vector (AddGene #44012) using Gateway cloning. pINDUCER20 expresses both a gene of interest under a tetracycline responsive TRE2 promoter as well as a tetracycline activator (rtTA3) under a constitutive promoter, enabling inducible regulation of a gene of interest following a single lentiviral transduction. The *TP53* library was expressed in a modified lentiviral pMT_BRD025 vector (AddGene #113569)[59]. To generate cell lines co-expressing p53 variants, fluorescent markers downstream of p53 and a t2a ribosome skip sequence were cloned into LX313-TP53-WT (Addgene #118014) plasmid using Gibson assembly (NEB E2621S). Site-directed mutagenesis (NEB E0554S) was used to generate point mutations in the p53 gene.

Lentivirus production. LentiX cells (Takara Bio), cultured as described above, were co-transfected with the structural plasmids necessary for virus production (psPAX2 and pMDM2.G from AddGene) along with the lentiviral vectors for either pINDUCER20.cHSF1, LX313-TP53-WT or variants, or the *TP53* mutational library. Cells were transfected using TransIT-Lenti (Mirus) for 24 h, after which the media was removed and replaced with fresh media. Media containing viral particles was collected at 48 h and cell debris was removed by centrifugation at 500 × *g* for 10 min. Viral supernatant was either concentrated using lentivirus precipitation solution (PEG-IT SBI) or aliquoted and stored at −80 °C until use. To measure the titer of the *TP53* library lentivirus, A549 cells were infected with serially diluted virus in 96-well plates. The infected cells were then selected in puromycin (Gibco) for 48 h and surviving cells were quantified using resazurin (Sigma).

Lentiviral titer assay. A549[cHSF1] cells were seeded in 96-well plates (Corning) at a density of 3×10^5 cells/well in DMEM medium. The following day media was removed and replaced with viral media containing polybrene at a final concentration of 8 μ/mL. After a 96-h incubation, media was removed and replaced with 100 μL of DMEM containing 0.01 mg/mL resazurin sodium salt (Sigma). After 2 h of incubation, resorufin fluorescence (excitation 530 nm; emission 590 nm) was quantified using a Take-3 plate reader (BioTeK). Experiments were conducted in biological triplicate. Viral titer in transducing

units per mL (TU/mL) was calculated as: [(number of cells plated) x (fraction of surviving cells)] / (volume of virus). The average of the calculated TU/mL over the linear range of the assay was used for subsequent calculation of appropriate multiplicity of infection.

Stable cell line engineering. For the construction of A549^{cHSF1} cells, A549 cells were transduced with lentivirus co-encoding a G418-resistance gene and rtTA3 alongside cHSF1 in the presence of 2 µg/mL polybrene (Sigma-Aldrich). Heterostable cell lines were then selected using 1 mg/mL G418 (Enzo Life Sciences). Clonal populations were screened based on functional testing of the cHSF1 construct using real-time polymerase chain reaction (RT-PCR; described below) with or without 1 µg/mL dox (Alfa Aesar). For the construction of U2OS^{cHSF1} cells, U2OS cells were transduced with lentivirus co-encoding a G418-resistance gene and rtTA3 alongside cHSF1 in the presence of 2 µg/mL polybrene (Sigma-Aldrich). Heterostable cell lines were then selected using 150 µg/mL G418 (Enzo Life Sciences) and functional testing of the cHSF1 construct was conducted using RT-PCR with or without 0.01 µg/mL dox.

Resazurin viability assay. A549^{cHSF1} cells were seeded in 96-well plates (Corning) at a density of 3 × 10^5 cells/well in DMEM medium and then treated with 0.1% DMSO, 1 µg/mL dox, 2.5 µM nutlin-3 (Cayman Chemical Company), or 1 µg/mL dox and 2.5 µM nutlin-3. 48 h post-treatment, media was removed and replaced with 100 µL of DMEM containing 0.01 mg/mL resazurin sodium salt (Sigma). After 2 h of incubation, resorufin fluorescence (excitation 530 nm; emission 590 nm) was quantified using a Take-3 plate reader (BioTeK). Experiments were conducted in biological triplicate.

RT-PCR. A549^{cHSF1} cells were treated with 1 µg/mL dox for 24 h for assessment of cHSF1 construct function, while a 6 h treatment with 500 nM STA-9090 (MedChem Express) was used as a positive control for HSR activation. RNA was extracted using the EZNA Total RNA Kit I (Omega). qRT-PCR reactions were performed on cDNA prepared from 1000 ng of total cellular RNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems). The Fast Start Universal SYBR Green Master Mix (Roche) and appropriate primers purchased Sigma were used for amplifications (6 min at 95 °C then 45 cycles of 10 s at 95 °C, 30 s at 60 °C) in a Light Cycler 480 II Real-Time PCR machine. The primers used for *DNAJB1* were 5′-TGTGTGGCTGCACAGTGAAC-3′ (forward) and 5′-ACGTTTCTCGGGTGTTTTGG-3′ (reverse), primers for *HSPA1A* were 5′-GGAGGCGGAGAAGTACA-3′ (forward) and 5′- GCTGATGATGGGGTTACA-3′ (reverse), primers for *RPLP2* were 5′-CCATTCAGCTCACTGATAACCTTG-3′ (forward) and 5′-CGTCGCCTCCTACCTGCT-3′ (reverse), primers for *CDKN1A* were 5′-CGACTGTGATGCGCTAATGG-3′ (forward) and 5′-CCGTGGGAAGGTAGAGCTTG-3′ (reverse), and primers for *BAX* were 5′-CAGCAAACTGGTGCTCAAGG-3′ (forward) and 5′-TCCTGGAGACAGGACATCA-3′ (reverse). Transcripts were normalized to the housekeeping genes *RPLP2*. All measurements were performed in technical triplicate. Data were analyzed

using the LightCycler® 480 Software, Version 1.5 (Roche) and data are reported as the mean ±95% confidence intervals.

RNA-Seq. A549^{cHSF1} cells were seeded at 7.5 x 10^4 cells/well in a 12-well plate in DMEM media. Cells were then treated with either 0.01 % DMSO or 1 µg/mL dox for 24 h. Cellular RNA was harvested using the RNeasy Plus Mini Kit with QIAshredder homogenization columns (Qiagen). RNA samples were quantified using an Advanced Analytical Fragment Analyzer. The initial steps were performed on a Tecan EVO150.10 ng of total RNA was used for library preparation. 3′DGE-custom primers 3V6NEXT-bmc#1-24 were added to a final concentration of 1 µM. (5'-/5Biosg/ACACTCTTTCCCTACACGAC-GCTCTTCCGATCT [BC6]N10T30VN-3′ where 5Biosg = 5′ biotin, [BC6] = 6bp barcode specific to each sample/well, N10 = Unique Molecular Identifiers, Integrated DNA technologies) were used to generate two subpools of 24 samples each[86, 87]. After addition of the oligonucleotides, Maxima H Minus RT was added per the manufacturer's recommendations with the template-switching oligo 5V6NEXT (10 µM, [5V6NEXT: 5′-iCiGiCACACTCTTTCCCTACACGACGCrGrGrG-3′ where iC: iso-dC, iG: iso-dG, rG: RNA G]), followed by incubation at 42 °C for 90 min and inactivation at 80 °C for 10 min. Following the template switching reaction, cDNA from 24 wells containing unique well identifiers were pooled together and cleaned using RNA Ampure beads at 1.0×. cDNA was eluted with 17 µL of water followed by digestion with Exonuclease I at 37 °C for 30 min, and inactivation at 80 °C for 20 min. Second strand synthesis and PCR amplification was done by adding the Advantage 2 Polymerase Mix (Clontech) and the SINGV6 primer (10 pM, Integrated DNA Technologies 5′-/5Biosg/ACACTCTTTCCCTACACGACGC-3′) directly to half of the exonuclease reaction volume. Eight cycles of PCR were performed, followed by clean-up using regular SPRI beads at 0.6×, and elution with 20 µL of Resuspension Buffer (Illumina). Successful amplification of cDNA was confirmed using the Fragment Analyzer. Illumina libraries were then produced using Nextera FLEX tagmentation substituting P5NEXTPT5-bmc primer (25 µM), Integrated DNA Technologies, (5′-AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGAC-GCTCTTCCG*A*T*C*T*-3′ where * = phosphorothioate bonds) in place of the normal N500 primer. Final libraries were cleaned using SPRI beads at 0.7× and quantified using the Fragment Analyzer and qPCR before being loaded for paired-end sequencing using the Illumina NextSeq500 in paired-end mode (26/50 nt reads).

Analyses were performed using previously described tools and methods[88]. Reads were aligned against hg19 (Feb., 2009) using bwa mem v. 0.7.12-r1039 [RRID:SCR_010910] with flags –t 16 –f, and mapping rates, fraction of multiply-mapping reads, number of unique 20-mers at the 5′ end of the reads, insert size distributions, and fraction of ribosomal RNAs were calculated using bedtools v. 2.25.0 [RRID:SCR_006646][89]. In addition, each resulting bam file was randomly down-sampled to a million

reads, which were aligned against hg19, and read density across genomic features were estimated for RNA-Seq-specific quality control metrics. For mapping and quantitation, reads were scored against GRCh38/ENSEMBL 101 annotation using Salmon v.1.3 with flags quant -p 8 -l ISR –validate-Mappings[90]. The resulting quant.sf files were imported into the R statistical environment using the tximport library (tximport function, option "salmon"), and gene-level counts and transcript per-milllion (TPM) estimates were calculated for protein-coding genes. Samples were clustered based on genes with average log2 TPM >0.1 across all samples (n=6320 genes) based on complete linkage clustering of the Cosine correlation among samples. Samples with similarity score <0.94, which were clear outliers from the rest, were excluded from further analysis (n=5).

Differential expression was also analyzed in the R statistical environment (R v.3.5.1) using Bioconductor's DESeq2 package on the protein-coding genes only [RRID:SCR_000154][91]. Dataset parameters were estimated using the estimateSizeFactors(), and estimateDispersions() functions; read counts across conditions were modeled based on a negative binomial distribution, and a Wald test was used to test for differential expression (nbinomWaldtest(), all packaged into the DESeq() function), using the treatment type as a contrast. Shrunken $\log_2$ fold-changes were calculated using the lfcShrink function, based on a normal shrinkage estimator[91]. Fold-changes and *p*-values were reported for each protein-coding gene. Upregulation was defined as a change in expression level >1.5-fold relative to the basal environment with a non-adjusted *p*-value < 10^{-5}. Gene ontology analyses were performed using the online DAVID server, according to tools and methods presented by Huang and co-workers[88].

Gene set enrichment analysis (GSEA). Differential expression results from DESeq2 were retrieved, and the "stat" column was used to pre-rank genes for GSEA analysis. These "stat" values reflect the Wald's test performed on read counts as modeled by DESeq2 using the negative binomial distribution. Genes that were not expressed were excluded from the analysis. GSEA (linux desktop version, v4.1)[92,93] was run in the pre-ranked mode against MSigDB 7.4 C5 (Gene Ontology) set, and ENSEMBL IDs were collapsed to gene symbols using the Human_ENSEMBL_Gene_ID_MSigDB.v7.4.chip (resulting in 12706 unique genes for par and 12141 for sg4, respectively). In addition, a weighted scoring scheme, meandiv normalization, and cutoffs on MSigDB signatures sizes (between 5 and 2000 genes, resulting in 8496 gene sets retained) were applied and 5000 permutations were run for *p*-value estimation.

Generating A549^cHSF1(p53-Lib) cells. A549^cHSF1 cells were infected with titered p53 library lentivirus at a multiplicity of infection of 0.25 (4×10^7 cells mixed with 1×10^7 lentiviral particles) in the presence of 8 µg/mL polybrene. Following transduction, cells were selected with 2 µg/mL puromycin (Gibco).

Deep mutational scanning. A549^{cHSF1}(p53-Lib) cells were seeded in 15 cm tissue culture plates at a density of 3×10^6 cells/plate. In order to maintain library diversity throughout selection, three plates were used per treatment for a total of 9×10^6 cells. Cells were treated with 0.01% DMSO, 1 µg/mL dox, 0.01% DMSO and 2.5 µM nutlin-3, or 1 µg/mL dox and 2.5 µM nutlin-3. Cells were trypsinized, counted, and re-seeded in three plates each at 3×10^6 cells/plate every 3 d. Following 12 d of treatment, cell pellets were harvested by centrifugation at 1,000 rpm for 5 min. Aliquots of 9×10^6 cells were snap-frozen in liquid N$_2$(g) in Eppendorf tubes and stored at –80 °C for subsequent DNA extraction. The deep mutational scanning experiment was repeated independently for a total of three biological replicates from the same p53-Lib cell line.

To prepare samples for Illumina sequencing, genomic DNA was purified from aliquots of frozen cells using the QIAamp Blood Midi Kit (Qiagen) and final DNA concentration was determined using a Qubit (Fischer). PCR amplicons of p53 were prepared using 2.0 µg of genomic DNA over 25 cycles and with Herculase II as the DNA polymerase (Agilent). The primers used were 5' ATTCTCCTTGGAATTT-GCCCTT 3' and 5' CATAGCGTAAAAGGAGCAACA 3'. Twelve PCR reactions were performed per sample, and the reactions were pooled and cleaned up using a PCR clean-up kit (Omega). The p53 amplicons were further gel-purified using a Pippin prep system (Sage Science) prior to library preparation via Nextera Flex. The resulting libraries were quantified using the Fragment Analyzer before they were pooled and sequenced on an Illumina NovaSeq with 2 × 150 bp paired-end reads.

Deep mutational scanning data analysis. The software ORFCall v1.0 [https://github.com/broadinstitute/ORFCall/releases/tag/v1.0] was used with flags -p -Q 30 to align the deep-sequencing reads against the *TP53* wild-type sequence and count the number of times each codon mutation was observed in each selection condition. The mutational fold-change for each variant was calculated by normalizing raw read counts to the total read count at each position. Next, the log$_2$ fold-change in selection versus mock conditions was calculated by taking the log of mutational fold-change in the selection condition normalized to the mutational fold-change in the mock condition. Mutational fitness in each condition was then determined by averaging the log$_2$ fold-change from three biological replicates. RSA was calculated using the software DSSP on chain A of the p53 DNA-binding domain crystal structure (PDBID 2OCJ)[94, 95]. DSSP calculates the solvent-accessible surface area of the monomer (ASA) and the RSA is calculated by dividing the ASA by the total theoretical solvent accessibility area[96]. Sites were classified as buried if the RSA was <0.2 and exposed if the RSA was >0.2.

Rosetta analysis. The calculations for ΔΔ*G* of protein stability upon substitution were performed using the cartesian_ddg application in Rosetta version 3.13[72]. The crystal structure of the DNA-binding domain of p53 (PDB ID: 2OCJ, chain A) was used as the initial structure for the ΔΔ*G* calculations[95]. The

initial p53 structure was relaxed using the Rosetta FastRelax application to generate a total of 20 relaxed decoys. The Rosetta FastRelax application performed five cycles of side-chain repacking and energy minimization with the Rosetta energy function ref2015_cart[72, 97-99]. The lowest energy structure of the 20 decoys was used as the wild-type structure for the cartesian_ddg calculation. In the cartesian_ddg calculation, the target residue was substituted with each of the 20 natural amino acids, and any neighboring residues within a 9-Å radius were repacked and energy-minimized using the ref2015_cart energy function. This calculation process was performed five times to generate five energy scores for the mutant and for the wild-type. The average wild-type scores were subtracted from the average mutant scores to calculate the $\Delta\Delta G$ values. The $\Delta\Delta G$ values were then scaled by a factor of 0.34; this scale factor was previously calculated by fitting Rosetta-predicted $\Delta\Delta G$ values to experimental $\Delta\Delta G$ values in units of kcal/mol, and is used here to better relate predicted $\Delta\Delta G$ values to experimental values[72].

Flow cytometry-based pairwise competition assay. A549[cHSF1] and U2OS[cHSF1] cells were transduced with concentrated lentivirus co-encoding a hygromycin-resistance gene, wild-type or a variant of p53, and a fluorescent protein. Transduced cells were then selected using 1 mg/mL of hygromycin-B for A549[cHSF1] cells or 100 µg/mL of hygromycin-B for U2OS[cHSF1] cells for 5 d. 25,000 wildtype + BFP cells with 25,000 variant + GFP cells were plated in one well of a 12-well plate, and the remaining cells were used for flow analysis to quantify the pre-competition mix. Cells were then treated with DMSO as vehicle or dox (1 µg/mL for A549[cHSF1] cells and 0.01 µg/mL for U2OS[cHSF1] cells), 2.5 µM nutlin-3, 500 nM STA-9090 (MedChem Express), 5 µM VER-155008 (Med-Chem Express) 24 h after plating. 48–96 h post-treatment, the GFP+ fraction of the cells was evaluated via flow cytometry. Fluorescence-based measurements for the validation of p53 variants were performed using the BD LSRFortessa Cell Analyzer in tube or plate reader format, using the BD FACSDiva v.3.0 software for data collection. Log_2 fold-change of the GFP+ cell fraction was calculated by first normalizing to the pre-competition mix and then to either the average of the vehicle-treated or nutlin-3-alone-treated cells.

Immunohistochemistry and confocal microscopy. A549[cHSF1] cells expressing p53 variants and a fluorescent protein were plated in a glass bottom 6-well plate (Cellvis). Cells were treated with drugs 24 h after plating. Cells were washed with PBS then fixed with 4% paraformaldehyde (Electron Microscopy Sciences) for 25 min at RT. Samples were treated with 0.1% Triton X-100 (Thermo Scientific) for 30 min to permeabilize the cells. Samples were then incubated in a solution of 5% BSA (Gibco) in PBS at RT for 30 min to block non-specific antibody binding. Cells were labeled with anti-p53 antibody (Santa-Cruz, sc-126, DO-1) in 1% BSA for 1 h at RT. Samples were washed 3× with PBS. Samples were incubated with Alexa Fluor 568-conjugated anti-mouse (Invitrogen) in 1% BSA for 1 h at RT. Samples were

stained with DAPI (Thermo Fisher). Images were acquired at the Swanson Biotechnology Center at the Koch Institute on an Evident FV4000 with a 100× oil-immersion objective (UPLSAPO100XS). The cellSens FV software was used for image acquisition. The excitation lasers to capture the images were 405, 488 and 561 nm. Image processing was performed using ImageJ.

Analysis of TCGA data. Human data from the TCGA lung adenocarcinoma study, particularly *TP53* mutations and *HSPA1A, HSPA6,* and *HSPA1B* mRNA expression data, were downloaded from cBioPortal. Study name: Lung adenocarcinoma TCGA, PanCancer Atlas, 507 total samples. We split the data into patients expressing wild-type and variants of p53. We then selected patients with variants of p53 that were found to have a mutational $\log_2$ fold-change nutlin-3 and HSF1 versus nutlin-3 alone >0.05 in our DMS experiment.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses. All experiments were performed in biological triplicate. Statistical analyses were done in Prisim GraphPad, Jupyter Notebook, and R. Site and mutational $\log_2$ fold-change (**Figure 2**) were calculated using a Wilcoxon signed-rank test. Statistical significance in mutational $\log_2$ fold-change in missense mutations and buried versus exposed sites (**Figures 4B**, **S2C**, **S2D**, **S6B**, and **S6D**) were calculated using a Welch's *t*-test for independent samples with Bonferroni correction, and significance from null was determined using a Wilcoxon signed-rank test. As a non-parametric test, the Wilcoxon's rank-sum test is particularly adept at assessing whether the sample comes from a population that is symmetrically distributed around a center (in our case, 0) without assuming normality of the distribution. All correlations were determined by calculating Pearson correlation coefficients using a two-tailed test. Statistical significance between exposed and buried sites (**Figure 4C**) and mRNA expression change between wild-type and HSF1 potentiated variants (**Figure 8C**) were calculated using a Wilcoxon-signed rank test. The statistical significance between solvent RSA classes or mutation types within a solvent accessibility class (**Figures S6C** and **S6D**) were evaluated using ANOVA, while comparisons between select conditions were calculated using Welch's *t*-test for independent samples with Bonferroni correction. We applied pairwise Welch-corrected *t*-tests in case of potential unequal variance among samples. Additionally, to control for type I errors, we applied a stringent Bonferroni correction to the resulting pairwise-comparison *p*-values. Statistical significance between stabilizing and destabilizing mutations based on experimental measurements (**Figure S7A**) as well as for significance between buried and exposed regions (**Figure 5B**) were calculated using a Welch's *t*-test for independent variables. ANOVA was also used to calculate statistical significance between RSA class and amino acid changes and $\Delta\Delta G$ (**Figure 5E**) and between missense mutations in the DMS experiment and

TP53 database (**Figures 8A** and **8B**). Statistical significance for flow cytometry-based competition assays was calculated using Fisher's exact test for counts data (biological replicates were summed prior to calculating significance) (**Figures 5E–H**, **7A**, and **7B**). Complete outcomes of Fisher's exact tests with odds ratio and 95% confidence interval are reported in **Table S9**.

References

1. Ostroverkhova, D., Przytycka, T.M. & Panchenko, A.R. Cancer driver mutations: Predictions and reality. *Trends Mol Med.* **29**, 554-566 (2023).
2. Hanahan, D. & Weinberg, R.A. The hallmarks of cancer. *Cell* **100**, 57-70 (2000).
3. Hanahan, D. & Weinberg, R.A. Hallmarks of cancer: The next generation. *Cell* **144**, 646-674 (2011).
4. Nagel, R., Semenova, E.A. & Berns, A. Drugging the addict: Non-oncogene addiction as a target for cancer therapy. *Embo Rep* **17**, 1516-1531 (2016).
5. Luo, J., Solimini, N.L. & Elledge, S.J. Principles of cancer therapy: Oncogene and non-oncogene addiction. *Cell* **136**, 823-837 (2009).
6. Dufey, E., Urra, H. & Hetz, C. ER proteostasis addiction in cancer biology: Novel concepts. *Semin Cancer Biol* **33**, 40-47 (2015).
7. Dai, C.K., Dai, S.Y. & Cao, J.Y. Proteotoxic stress of cancer: Implication of the heat-shock response in oncogenesis. *J Cell Physiol* **227**, 2982-2987 (2012).
8. Guang, M.H.Z. et al. Targeting proteotoxic stress in cancer: A review of the role that protein quality control pathways play in oncogenesis. *Cancers* **11** (2019).
9. Mayer, M.P. et al. Stress biology: Complexity and multifariousness in health and disease. *Cell Stress Chaperones* **29**, 143-157 (2024).
10. Bagratuni, T. et al. XBP1s levels are implicated in the biology and outcome of myeloma mediating different clinical outcomes to thalidomide-based treatments. *Blood* **116**, 250-253 (2010).
11. Chen, X. et al. XBP1 promotes triple-negative breast cancer by controlling the HIF1alpha pathway. *Nature* **508**, 103-107 (2014).
12. Akerfelt, M., Morimoto, R.I. & Sistonen, L. Heat shock factors: Integrators of cell stress, development and lifespan. *Nat Rev Mol Cell Biol.* **11**, 545-555 (2010).
13. Alasady, M.J. & Mendillo, M.L. The multifaceted role of HSF1 in tumorigenesis. *Adv Exp Med Biol.* **1243**, 69-85 (2020).
14. Mendillo, M.L. et al. HSF1 drives a transcriptional program distinct from heat shock to support highly malignant human cancers. *Cell* **150**, 549-562 (2012).
15. Scherz-Shouval, R. et al. The reprogramming of tumor stroma by HSF1 is a potent enabler of malignancy. *Cell* **158**, 564-578 (2014).
16. Santagata, S. et al. Tight coordination of protein translation and HSF1 activation supports the anabolic malignant state. *Science* **341**, 1238303 (2013).
17. Dai, C., Whitesell, L., Rogers, A.B. & Lindquist, S. Heat shock factor 1 is a powerful multifaceted modifier of carcinogenesis. *Cell* **130**, 1005-1018 (2007).
18. Whitesell, L. et al. HSP90 empowers evolution of resistance to hormonal therapy in human breast cancer models. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 18297-18302 (2014).
19. Desai, S. et al. Heat shock factor 1 (HSF1) controls chemoresistance and autophagy through transcriptional regulation of autophagy-related protein 7 (ATG7). *J Biol Chem.* **288**, 9165-9176 (2013).
20. Li, Z.N. & Luo, Y. HSP90 inhibitors and cancer: Prospects for use in targeted therapies (Review). *Oncol Rep.* **49**, 6 (2023).
21. Dong, B. et al. Targeting therapy-resistant prostate cancer via a direct inhibitor of the human heat shock transcription factor 1. *Sci Transl Med.* **12**, eabb5647 (2020).
22. Vilaboa, N. et al. New inhibitor targeting human transcription factor HSF1: Effects on the heat shock response and tumor cell survival. *Nucleic Acids Res.* **45**, 5797-5817 (2017).
23. Yoon, J., Patrick, J.E., Ogbunugafor, C.B. & Shoulders, M.D. Viral evolution shaped by host proteostasis networks. *Annu Rev Virol.* **10**, 77-98 (2023).
24. DePristo, M.A., Weinreich, D.M. & Hartl, D.L. Missense meanderings in sequence space: A biophysical view of protein evolution. *Nat Rev Genet.* **6**, 678-687 (2005).

25. Rutherford, S.L. & Lindquist, S. Hsp90 as a capacitor for morphological evolution. *Nature* **396**, 336-342 (1998).
26. Geller, R., Pechmann, S., Acevedo, A., Andino, R. & Frydman, J. Hsp90 shapes protein and RNA evolution to balance trade-offs between protein stability and aggregation. *Nat Commun.* **9**, 1781 (2018).
27. Karras, G.I. et al. HSP90 shapes the consequences of human genetic variation. *Cell* **168**, 856-866 e812 (2017).
28. Phillips, A.M. et al. Host proteostasis modulates influenza evolution. *eLife* **6**, e28652 (2017).
29. Phillips, A.M. et al. Enhanced ER proteostasis and temperature differentially impact the mutational tolerance of influenza hemagglutinin. *eLife* **7**, e38795 (2018).
30. Phillips, A.M. et al. Destabilized adaptive influenza variants critical for innate immune system escape are potentiated by host chaperones. *PLoS Biol.* **16**, e3000008 (2018).
31. Yoon, J. et al. The endoplasmic reticulum proteostasis network profoundly shapes the protein sequence space accessible to HIV envelope. *PLoS Biol.* **20**, e3001569 (2022).
32. Nekongo, E.E. et al. HSF1 activation can restrict HIV replication. *ACS Infect Dis.* **6**, 1659-1666 (2020).
33. Yoon, J. et al. The immune-evasive proline-283 substitution in influenza nucleoprotein increases aggregation propensity without altering the native structure. *Sci Adv.* **10**, eadl6144 (2024).
34. Mantovani, F., Collavin, L. & Del Sal, G. Mutant p53 as a guardian of the cancer cell. *Cell Death Differ.* **26**, 199-212 (2019).
35. Mendiratta, G. et al. Cancer gene mutation frequencies for the U.S. population. *Nat Commun.* **12**, 5961 (2021).
36. Baugh, E.H., Ke, H., Levine, A.J., Bonneau, R.A. & Chan, C.S. Why are there hotspot mutations in the TP53 gene in human cancers? *Cell Death Differ.* **25**, 154-160 (2018).
37. Boettcher, S. et al. A dominant-negative effect drives selection of TP53 missense mutations in myeloid malignancies. *Science* **365**, 599-604 (2019).
38. Gencel-Augusto, J. & Lozano, G. p53 tetramerization: At the center of the dominant-negative effect of mutant p53. *Genes Dev.* **34**, 1128-1146 (2020).
39. De Vries, A. et al. Targeted point mutations of p53 lead to dominant-negative inhibition of wild-type p53 function. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 2948-2953 (2002).
40. Dahiya, V. et al. Coordinated conformational processing of the tumor suppressor protein p53 by the HSP70 and HSP90 chaperone machineries. *Mol Cell* **74**, 816-830 e817 (2019).
41. Boysen, M., Kityk, R. & Mayer, M.P. HSP70- and HSP90-mediated regulation of the conformation of p53 DNA binding domain and p53 cancer variants. *Mol Cell* **74**, 831-843 e834 (2019).
42. Muller, L., Schaupp, A., Walerych, D., Wegele, H. & Buchner, J. Hsp90 regulates the activity of wild type p53 under physiological and elevated temperatures. *J Biol Chem.* **279**, 48846-48854 (2004).
43. Kaida, A. & Iwakuma, T. Regulation of p53 and cancer signaling by heat shock protein 40/J-domain protein family members. *Int J Mol Sci.* **22**, 13527 (2021).
44. Schulz-Heddergott, R. & Moll, U.M. Gain-of-Function (GOF) Mutant p53 as Actionable Therapeutic Target. *Cancers (Basel)* **10**, 188 (2018).
45. Wiech, M. et al. Molecular mechanism of mutant p53 stabilization: The role of HSP70 and MDM2. *PLoS One* **7**, e51426 (2012).
46. Li, D. et al. Functional inactivation of endogenous MDM2 and CHIP by HSP90 causes aberrant stabilization of mutant p53 in human cancer cells. *Mol Cancer Res.* **9**, 577-588 (2011).
47. Giard, D.J. et al. In vitro cultivation of human tumors: establishment of cell lines derived from a series of solid tumors. *J Natl Cancer Inst.* **51**, 1417-1423 (1973).
48. Lehman, T.A. et al. p53 mutations, ras mutations, and p53-heat shock 70 protein complexes in human lung carcinoma cell lines. *Cancer Res.* **51**, 4090-4096 (1991).
49. Sebastian, R.M. & Shoulders, M.D. Chemical Biology Framework to Illuminate Proteostasis. *Annu Rev Biochem.* **89**, 529-555 (2020).

50. Voellmy, R. Dominant-positive and dominant-negative heat shock factors. *Methods* **35**, 199-207 (2005).

51. Moore, C.L. et al. Transportable, chemical genetic methodology for the small molecule-mediated inhibition of Heat Shock Factor 1. *ACS Chem Biol.* **11**, 200-210 (2016).

52. Ryno, L.M. et al. Characterizing the altered cellular proteome induced by the stress-independent activation of heat shock factor 1. *ACS Chem Biol.* **9**, 1273-1283 (2014).

53. Liebelt, F. et al. SUMOylation and the HSF1-Regulated Chaperone Network Converge to Promote Proteostasis in Response to Heat Shock. *Cell Rep.* **26**, 236-249 e234 (2019).

54. Kovacs, D. et al. HSF1Base: A comprehensive database of HSF1 (heat shock factor 1) target genes. *Int J Mol Sci.* **20**, 5815 (2019).

55. Ying, W. et al. Ganetespib, a unique triazolone-containing Hsp90 inhibitor, exhibits potent antitumor activity and a superior safety profile for cancer therapy. *Mol Cancer Ther.* **11**, 475-484 (2012).

56. Riss, T.L. et al. in Assay Guidance Manual (Bethesda (MD); 2004).

57. Liberzon, A. et al. The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst.* **1**, 417-425 (2015).

58. Prieto, C. & De Las Rivas, J. APID: Agile Protein Interaction DataAnalyzer. *Nucleic Acids Res.* **34**, W298-302 (2006).

59. Giacomelli, A.O. et al. Mutational processes shape the landscape of TP53 mutations in human cancer. *Nat Genet.* **50**, 1381-1387 (2018).

60. Shen, H. & Maki, C.G. Pharmacologic activation of p53 by small-molecule MDM2 antagonists. *Curr Pharm Des.* **17**, 560-568 (2011).

61. Bouaoun, L. et al. TP53 variations in human cancers: New lessons from the IARC TP53 database and genomics data. *Hum Mutat.* **37**, 865-876 (2016).

62. de Andrade, K.C. et al. The TP53 Database: transition from the International Agency for Research on Cancer to the US National Cancer Institute. *Cell Death Differ* **29**, 1071-1073 (2022).

63. Doud, M.B. & Bloom, J.D. Accurate measurement of the effects of all amino-acid mutations on influenza hemagglutinin. *Viruses* **8**, 155 (2016).

64. Joerger, A.C. & Fersht, A.R. Structural biology of the tumor suppressor p53. *Annu Rev Biochem.* **77**, 557-582 (2008).

65. Natan, E. et al. Interaction of the p53 DNA-binding domain with its N-terminal extension modulates the stability of the p53 tetramer. *J Mol Biol.* **409**, 358-368 (2011).

66. Bullock, A.N., Henckel, J. & Fersht, A.R. Quantitative analysis of residual folding and DNA binding in mutant p53 core domain: Definition of mutant states for rescue in cancer therapy. *Oncogene* **19**, 1245-1256 (2000).

67. Butler, J.S. & Loh, S.N. Structure, function, and aggregation of the zinc-free form of the p53 DNA binding domain. *Biochemistry* **42**, 2396-2403 (2003).

68. Blanden, A.R. et al. Zinc shapes the folding landscape of p53 and establishes a pathway for reactivating structurally diverse cancer mutants. *eLife* **9**, e61487 (2020).

69. Joerger, A.C., Ang, H.C., Veprintsev, D.B., Blair, C.M. & Fersht, A.R. Structures of p53 cancer mutants and mechanism of rescue by second-site suppressor mutations. *J Biol Chem.* **280**, 16030-16037 (2005).

70. Nikolova, P.V., Henckel, J., Lane, D.P. & Fersht, A.R. Semirational design of active tumor suppressor p53 DNA binding domain with enhanced stability. *Proc. Natl. Acad. Sci. U. S. A.* **95**, 14675-14680 (1998).

71. Nikolova, P.V., Wong, K.B., DeDecker, B., Henckel, J. & Fersht, A.R. Mechanism of rescue of common p53 cancer mutations by second-site suppressor mutations. *EMBO J.* **19**, 370-378 (2000).

72. Park, H. et al. Simultaneous optimization of biomolecular energy functions on features from small molecules and macromolecules. *J Chem Theory Comput.* **12**, 6201-6212 (2016).

73. Ponten, J. & Saksela, E. Two established in vitro cell lines from human mesenchymal tumours. *Int J Cancer* **2**, 434-447 (1967).

74. Diller, L. et al. p53 functions as a cell cycle control protein in osteosarcomas. *Molecular and Cellular Biology* **10**, 5772-5781 (1990).

75. Willis, A., Jung, E.J., Wakefield, T. & Chen, X. Mutant p53 exerts a dominant negative effect by preventing wild-type p53 from binding to the promoter of its target genes. *Oncogene* **23**, 2330-2338 (2004).

76. Massey, A.J. et al. A novel, small molecule inhibitor of Hsc70/Hsp70 potentiates Hsp90 inhibitor induced apoptosis in HCT116 colon carcinoma cells. *Cancer Chemother Pharmacol* **66**, 535-545 (2010).

77. Cerami, E. et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* **2**, 401-404 (2012).

78. de Bruijn, I. et al. Analysis and Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE Biopharma Collaborative in cBioPortal. *Cancer Res* **83**, 3861-3867 (2023).

79. Gao, J. et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* **6**, pl1 (2013).

80. Dai, C. & Sampson, S.B. HSF1: Guardian of proteostasis in cancer. *Trends Cell Biol.* **26**, 17-28 (2016).

81. Solimini, N.L., Luo, J. & Elledge, S.J. Non-oncogene addiction and the stress phenotype of cancer cells. *Cell* **130**, 986-988 (2007).

82. Tchenio, T., Havard, M., Martinez, L.A. & Dautry, F. Heat shock-independent induction of multidrug resistance by heat shock factor 1. *Mol Cell Biol.* **26**, 580-591 (2006).

83. Vydra, N., Toma, A., Glowala-Kosinska, M., Gogler-Piglowska, A. & Widlak, W. Overexpression of Heat Shock Transcription Factor 1 enhances the resistance of melanoma cells to doxorubicin and paclitaxel. *BMC Cancer* **13**, 504 (2013).

84. Alexandrova, E.M. et al. Improving survival by exploiting tumour dependence on stabilized mutant p53 for treatment. *Nature* **523**, 352-356 (2015).

85. Jaeger, A.M. et al. Rebalancing Protein Homeostasis Enhances Tumor Antigen Presentation. *Clin Cancer Res* **25**, 6392-6405 (2019).

86. Soumillon, M., Cacchiarelli, D., Semrau, S., van Oudenaarden, A. & Mikkelsen, T.S. Characterization of directed differentiation by high-throughput single-cell RNA-Seq *bioRxiv* (2014).

87. Struntz, N.B. et al. Stabilization of the max homodimer with a small molecule attenuates myc-driven transcription. *Cell Chem Biol.* **26**, 711-723 e714 (2019).

88. Huang, D.W., Sherman, B.T. & Lempicki, R.A. Bioinformatics enrichment tools: Paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37**, 1-13 (2009).

89. Quinlan, A.R. & Hall, I.M. BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010).

90. Patro, R., Duggal, G., Love, M.I., Irizarry, R.A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).

91. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

92. Subramanian, A. et al. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545-15550 (2005).

93. Mootha, V.K. et al. PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet.* **34**, 267-273 (2003).

94. Kabsch, W. & Sander, C. Dictionary of protein secondary structure: Pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **22**, 2577-2637 (1983).

95. Wang, Y., Rosengarth, A. & Luecke, H. Structure of the human p53 core domain in the absence of DNA. *Acta Crystallogr D Biol Crystallogr.* **63**, 276-281 (2007).

96.   Tien, M.Z., Meyer, A.G., Sydykova, D.K., Spielman, S.J. & Wilke, C.O. Maximum allowed solvent accessibilites of residues in proteins. *PLoS One* **8**, e80635 (2013).

97.   Maguire, J.B. et al. Perturbing the energy landscape for improved packing during computational protein design. *Proteins* **89**, 436-449 (2021).

98.   Alford, R.F. et al. The Rosetta all-atom energy function for macromolecular modeling and design. *J Chem Theory Comput.* **13**, 3031-3048 (2017).

99.   Khatib, F. et al. Algorithm discovery by protein folding game players. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 18949-18953 (2011).

Figure 1: Characterization of chemical genetic HSF1 regulation and construction of the A549^cHSF1(p53-Lib) cell line.

(**A**) qPCR results showing transcript-level consequences of dox-mediated HSF1 activation for the HSF1 target genes *DNAJB1* and *HSPA1A* in A549^cHSF1 cells. Statistical significance was calculated using two-tailed students *t*-test where **** represents a *p*-value of <0.0001. Treatment with the HSP90 inhibitor STA-9090 was used as a positive control for HSR activation, and to ensure pathways were activated within the regime accessible to endogenous HSF1 transcriptional activity. (**B**) Volcano plot for RNA-Seq analysis of changes in gene transcription following HSF1 activation in A549^cHSF1 cells as compared to vehicle treatment. Proteins within the Agile Protein Interaction DataAnalyzer (APID) that have been identified as interactors with p53 are labeled and shown in red. (**C**) Heat map depicting log_2 fold-change of transcript expression of selected p53 interacting genes highlighted in (**B**). (**D**) Creation of A549^cHSF1(p53-Lib) cells: A549 cells expressing endogenous wild-type p53 were transduced at a low multiplicity of infection with a lentiviral population encoding all possible single amino acid substitutions within *TP53*.

Figure 2: HSF1 activation enhances mutational fitness of dominant-negative p53 variants.
(**A**) Selection of dominant-negative variants using nutlin-3. Treatment with nutlin-3 prevents MDM2-mediated degradation of both wild-type and mutant p53. Co-expression of dominant-negative p53 variants prevents induction of cell cycle arrest by endogenous wild-type p53, thereby allowing cell proliferation. (**B**) Heat map of p53 mutational frequency $\log_2$ fold-change averaged over three biological replicates for each amino acid substitution for nutlin-3-mediated dominant-negative p53 selection in an HSF1-activated environment as compared to nutlin-3-mediated dominant-negative p53 selection in a basal proteostasis environment (*top*). Sum of the mutational $\log_2$ fold-change at each site. Color indicates net positive (*red*) or negative (*blue*) site $\log_2$ fold-change (*bottom*). (**C**) Total site $\log_2$ fold-change (all substitutions for a given site) for amino acid substitutions in the selection conditions in (**B**) subdivided across each p53 domain. (**D**) $\log_2$ fold-change for each individual DNA-level mutation in the *TP53* gene subdivided for each domain, presented here for the selection conditions in (**B**) and separating missense (*colored*) from synonymous mutations (*grey*). Significance was calculated using a Wilcoxon signed-rank test, with *, ***,

and **** representing adjusted two-tailed p-values of <0.05, <0.001, and <0.0001, respectively, and ns indicating non-significant. AD1, activation domain 1; AD2, activation domain 2; PRD, proline-rich domain; DBD, DNA-binding domain; TD, tetramerization domain; RD, regulatory domain.

Figure 3: HSF1 activation in the context of nutlin-3-mediated selection alters mutational fitness within the p53 DNA-binding domain.
(**A**) Structure of the p53 DNA-binding domain (PDB 2OCJ). Residues are colored according to the net site log$_2$ fold-change. (**B**) Consequences of HSF1 activation at selected, individually labeled hot-spot sites associated with cancer. Color indicates net positive (*red*) or negative (*blue*) site log$_2$ fold-change. (**C**) Sequence logo plot of mutational log$_2$ fold-change for selected sites, full sequence logo plot for DNA binding domain is available in **Figure S5**. p53 hot-spot sites are highlighted in *yellow* and Zn^{2+} coordination sites are highlighted in *blue*. Alleles displaying opposing signs between biological replicates were removed, so all data shown are for variants that displayed a consistent response across biological replicates.

Figure 4: HSF1 most strongly increases fitness of destabilizing substitutions involving biophysically disruptive replacement of nonpolar residues with polar or charged residues in buried regions of the p53 DNA-binding domain.

(**A**) Mutational $\log_2$ fold-change in variant fitness for conservative amino acid substitutions, non-conservative nonpolar to polar substitutions, and non-conservative polar to nonpolar substitutions at buried (RSA < 0.2) versus exposed (RSA > 0.2) sites in the p53 DNA binding domain. (**B**) Theoretical $\Delta\Delta G$ calculated using Rosetta analysis for p53 DNA-binding domain substitutions in buried, partially exposed, or exposed sites. Amino acids in buried regions were generally thermodynamically less tolerant, and especially less tolerant to non-conservative nonpolar-to-polar/charged substitutions than sites in exposed regions. For (**A**) and (**B**), statistical significance between solvent accessibility classes or mutation types within a solvent accessibility class was evaluated using ANOVA, while comparisons between select conditions were calculated using Welch's t-test for independent samples with Bonferroni correction. *, ***, and **** represent adjusted two-tailed p-values of <0.05, <0.001, and <0.0001, respectively (**C**) Theoretical $\Delta\Delta G$ of buried and exposed variants binned according to mutational $\log_2$ fold-change in HSF1-activated versus basal proteostasis environments during nutlin-3 selection. Individual box plots represent bins of 0.2 mutational $\log_2$ fold-change greater than the lower limit and up to and including the upper limit. Statistical significance was calculated using a Wilcoxon signed-rank test, with *, ***, and **** representing adjusted two-tailed p-values of <0.05, <0.001, and <0.0001, respectively.

Figure 5: HSF1-potentiated dominant-negative p53 variants identified in the DMS experiment increase in fitness upon HSF1 activation in multiple cancer cell lines.
(**A**) Summary table of selected p53 variants. (**B**) qPCR results showing transcript expression of p53 target genes *BAX* and *CDKN1A* of cells expressing wild-type p53 and an HSF1-potentiated dominant-negative p53 variant during HSF1 activation and nutlin-3 treatment, confirming dominant-negative behavior of selected variants. (**C**) Workflow of flow cytometry-based pairwise competition assay. Log$_2$ fold-change of GFP frequency as a readout of mutant p53 enrichment under nutlin-3 selection versus vehicle (**D**) and nutlin-3 and HSF1 activation versus nutlin-3 only (**E**) in A549^{cHSF1} cells. Log$_2$ fold-change of GFP+ cells as a readout of mutant p53 enrichment under nutlin-3 selection versus vehicle (**F**) and nutlin-3 and HSF1 activation versus nutlin-3 only (**G**) in U2OS^{cHSF1} cells. Significance was assessed using Fisher's exact test for count data, with **** representing an adjusted two-tailed *p*-value <0.0001.
.

Figure 6: The destabilizing p53 V173Y variant forms aggregates in A549 cells that are cleared by HSF1 activation and exacerbated with HSP90 inhibition.

A549^cHSF1 cells stably expressing GFP (to visualize transduction efficiency), wild-type p53, and p53^V173Y were treated with either vehicle, dox to activate HSF1, or STA-9090 to inhibit HSP90. Confocal images of the cells stained for p53 and nuclear stain DAPI reveal that p53 aggregation in the nucleus under basal conditions was cleared by activation of HSF1 while aggregation of p53 found in the nucleus was exacerbated with HSP90 inhibition.

Figure 7: Inhibition of chaperones downstream of HSF1 compromises p53 mutant fitness.
Log$_2$ fold-change of mutant p53 with HSP70 (**A**) and HSP90 (**B**) inhibition in a pairwise competition assay. Significance was calculated using a Fisher's exact test for count data, with **** representing adjusted two-tailed *p*-values of <0.0001.

Figure 8: HSF1-potentiated p53 variants are enriched in patients.
(**A**) Box plot comparing the effects of HSF1 in a nutlin-3 environment (mutational $\log_2$ fold-change nutlin-3 and HSF1 activation versus nutlin-3 only) of missense mutations found in the DMS experiment (*orange*) compared to missense mutations with (*green*) and without (*blue*) the dominant-negative loss of function (DNE_LOF) annotation in the TP53 database (r21 release). Statistical significance was calculated using a Wilcoxon-signed rank test with **** representing a *p*-value <0.0001. (**B**) Violin plots depicting mRNA expression of sentinel genes downstream of HSF1 (*HSPA1A*, *HSPA1B* and *HSPA6*) in patients expressing wild-type p53 or an HSF1-potentiated p53 variant. Statistical significance was calculated using a Wilcoxon sign rank test with ** and *** representing adjusted two-tail *p*-values <0.01 and <0.001, respectively.