══ R E V I E W C O M M O N S M A N U S C R I P T ══

IMPORTANT:

- Manuscripts submitted to Review Commons are peer reviewed in a journal-agnostic way.
- Upon transfer of the peer reviewed preprint to a journal, the referee reports will be available in full to the handling editor.
- The identity of the referees will NOT be communicated to the authors unless the reviewers choose to sign their report.
- The identity of the referee will be confidentially disclosed to any affiliate journals to which the manuscript is transferred.

GUIDELINES:

- For reviewers: https://www.reviewcommons.org/reviewers
- For authors: https://www.reviewcommons.org/authors

CONTACT:

The Review Commons office can be contacted directly at: office@reviewcommons.org

Lymphopenia drives T cell exhaustion in immunodeficient STING gain-of-function mice

Damien Freytag [a], Stéphane Giorgiutti [a,b], Nadège Wadier [a], Sabine Depauw [a], Grégoire

Hopsomer [a], Philippe Mertz [a], Fabrice Augé [c], Raphaël Carapito [d], Isabelle Couillin [e], Anne-

Sophie Korganow [a,b], Francesca Pala [f], Marita Bosticardo [f], Luigi D. Notarangelo [f], Frédéric

Rieux-Laucat [g], Nicolas Riteau [e,h], Peggy Kirstetter [i], Pauline Soulas-Sprauel [a,b,*]

[*] Corresponding author: e-mail address: soulaspa@unistra.fr (P. Soulas-Sprauel); postal

address: Inserm U1109, CRBS, 1 rue Eugène Boeckel, F-67000 Strasbourg, France.

Affiliations:

[a] Laboratoire d'ImmunoRhumatologie Moléculaire, INSERM UMR_S1109, F-67000

Strasbourg, France; Strasbourg University, Faculty of Pharmacy and Faculty of Medicine,

Strasbourg, France.

[b] Department of Clinical Immunology and Internal Medicine, National Reference Center for

Systemic Autoimmune Diseases (CNR RESO), Tertiary Center for Primary

Immunodeficiency, Strasbourg University Hospital, F-67000 Strasbourg, France.

[c] IBMC, Institut de Biologie Moléculaire et Cellulaire, F-67000 Strasbourg, France.

[d] Laboratoire d'ImmunoRhumatologie Moléculaire, Plateforme GENOMAX, INSERM

UMR_S 1109, Faculté de Médecine, Fédération Hospitalo-Universitaire OMICARE, ITI

TRANSPLANTEX NG, Strasbourg Federation of Translational Medicine (FMTS),

Strasbourg University, Strasbourg 67091, France

[e] Experimental and Molecular Immunology and Neurogenetics Laboratory, University of

Orleans, Centre National de la Recherche Scientifique (CNRS), UMR7355, 45100 Orleans,

France.

^f Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and

Infectious Diseases (NIAID)/NIH, Bethesda, MD, USA.

^g Université Paris Cité, Laboratoire d'immunogénétique des maladies auto-immunes

pédiatriques, Institut Imagine, INSERM UMR_S1163, Paris, France.

^h Immune Health Laboratory, 'Regulation of host responses and immune health' IRL2029,

French National Centre for Scientific Research (CNRS) and Ribeirão Preto Medical School

(FMRP) of the São Paulo University (USP).

ⁱ Université de Strasbourg, IGBMC UMR 7104- UMR-S 1258, F-67400 Illkirch, France;

IGBMC, Institut de Génétique et de Biologie Moléculaire et Cellulaire, F-67400 Illkirch,

France.

Classification: Biological Sciences – Immunology and Inflammation

Keywords:

STING, T cell exhaustion, Lymphopenia, Homeostasis

Abstract

STING gain-of-function (GOF) mutations are associated with the severe autoinflammatory

disease designated STING Associated Vasculopathy with onset in Infancy (SAVI). Mice with

the STING GOF V154M mutation develop profound T cell lymphopenia, partly due to a

blockage of T cell development in the thymus. To better characterize the mechanisms of

peripheral T cell dysfunctions, we conducted a transcriptomic and phenotypic analysis on

sorted splenic CD4+ and CD8+ mature T cells from STING GOF V154M mice. We identify a

T cell exhaustion phenotype that manifests at a terminal stage, acquired early in life but only

after reaching the peripheral environment. This phenotype is independent of type I interferons

and does not rely on intrinsic STING activation in either T cells or stromal cells. Mechanistically, the limited number of mature T cells that reach the periphery appear to be quickly impacted by the lymphopenic environment, experiencing heightened stimulation of the IL-7 receptor and TCR pathways, including the NFAT pathway, a key factor in T cell exhaustion. By performing transplantation experiments with STING GOF long term-hematopoietic stem cells (LT-HSCs) along with supportive wild-type bone marrow (BM) cells, we were able to prevent the T cell exhaustion of STING GOF T cells in the resulting non-lymphopenic context, demonstrating that lymphopenia is a major driver of T cell exhaustion in STING GOF mice. T cell exhaustion, although less severe, was also observed in lymphopenic mice carrying *Rag1* hypomorphic mutations. In conclusion, our results, which highlight T cell exhaustion induced by lymphopenia, could have important implications for the management of patients with severe immune deficiencies.

Highlights

- We describe a phenotype of T cell exhaustion in STING GOF V154M mice, which is acquired early in life and in the periphery.

- STING GOF-associated T cell exhaustion is independent of type I IFNs, and STING GOF/activation in T cells or in stromal cells is not sufficient for T cell exhaustion.

- Lymphopenia is a major driver of T cell exhaustion in STING GOF mice, and increased antigenic/IL-7 stimulation of T cells in the lymphopenic context of STING GOF mice could be implicated in the induction of T cell exhaustion.

Introduction

Stimulator of Interferon Genes (STING) belongs to cytosolic double-stranded-DNA-

sensing pathways in innate immunity and has emerged as a central player in antiviral,

antibacterial, and anti-tumor immunity (1). When activated by cyclic dinucleotides produced

by cyclic GMP-AMP synthase (cGAS), STING translocates from endoplasmic reticulum

(ER) to ER-Golgi intermediate compartment (ERGIC)/Golgi where it activates TANK-

binding kinase 1 (TBK1), which then phosphorylates interferon regulatory factor 3 (IRF3)

and induces the expression of type I interferons (IFNs). In turn, type I IFNs activate the type I

IFNs receptor (IFNAR), inducing the expression of a set of interferon-stimulated genes (ISG)

and triggering inflammation. The NF-kB pathway is also engaged following STING

activation, participating in the inflammatory response (2).

Heterozygous gain-of-function (GOF) mutations in STING-coding gene *STING1* lead

to constitutive activation of the protein and have been described in patients with an

autoinflammatory disease designated STING Associated Vasculopathy with onset in Infancy

(SAVI) (3, 4), classified as a type I interferonopathy, with increased ISGs expression (5). This

disease is associated with damages to the skin small blood vessels and an interstitial lung

disease at the origin of morbidity and mortality (6–8). To study the role of STING in the

pathophysiology of the disease, our team (9) and others (10–16) developed STING GOF

mouse models including mice carrying the heterozygous V154M mutation (9, 10),

corresponding to the most common human V155M mutation (about 60%) in SAVI patients.

STING GOF V154M mouse models develop type I IFN-independent auto-inflammatory

manifestations, notably lung disease (10, 17), and also a high decrease of T cell counts in a

context of severe combined immunodeficiency (SCID) (9, 10), reminiscent of T cell

lymphopenia cases among SAVI patients. T cell lymphopenia was also described in STING

GOF N153S models (10, 11). In STING GOF N153S and V154M mice, the T cell

lymphopenia is partly explained by an impairment of T cell development at double negative

(DN)1 and DN2 stages (9, 11).

In addition, we showed that the rare mature T cells found in the periphery exhibited

both defective proliferation and increased apoptosis, which maintain the lymphopenic state

(9). These defects are at least partly STING-intrinsic as its anti-proliferative effect was

confirmed in human T cells overexpressing the STING GOF V155M mutant (18). In addition,

a pro-apoptotic effect of STING in lymphocytes was also reported in other models, and

involves TBK1 (19), endoplasmic reticulum (ER) stress (20) or p53-dependent pathways (21).

However, treatment of STING N153S mice with a pan-caspase inhibitor or ER stress blocker

only partially reverse T cell lymphopenia (20). Thus, other mechanisms must explain the

functional defects of peripheral T cells and maintenance of lymphopenia in STING GOF

mice.

To better characterize these mechanisms, we perform in the present study a

transcriptomic analysis on sorted T cells from STING GOF mice compared to their wild-type

(WT) littermate controls. An enrichment of transcriptomic signatures associated with T cell

exhaustion is revealed and confirmed by the overexpression of key transcription factors and

several surface inhibitory receptors (like PD-1, TIGIT, TIM-3 and LAG-3). We further

demonstrate that T cell exhaustion is associated with enhanced TCR (including NFAT

signaling) and IL-7R signaling in the lymphopenic environment. Importantly, we do not

observe any sign of T cell exhaustion in STING GOF T cells developing in non-lymphopenic

mice. Finally, a similar T cell exhaustion phenotype (albeit less severe than in STING GOF

mice) is observed in other lymphopenic conditions, as shown in mice carrying hypomorphic

mutations of *Rag1* gene (22). Together, these observations demonstrate that T cell exhaustion

phenotype in STING V154M mice is mostly triggered by lymphopenia and likely reinforce

this lymphopenia by a self-maintaining loop. While T cell exhaustion has primarily been

described as a consequence of chronic T cell stimulation in the context of persistent infections

and cancers (23, 24), our study reveals that lymphopenia should also be considered as a

context that can trigger T cell exhaustion, echoing recent findings of T cell exhaustion in

patients with severe combined immunodeficiencies (SCID) (25, 26).

Results

CD4$^+$ and CD8$^+$ T cells from STING GOF mice display an exhausted phenotype,

acquired in the periphery in a type I IFN independent manner

Mice carrying the heterozygous STING V154M mutation are characterized by a SCID

phenotype associated with marked T cell dysfunctions, including impaired proliferation, and

enhanced apoptosis (9). To decipher the mechanisms underlying this phenotype, we

performed a transcriptomic analysis by RNA-seq on sorted splenic CD4$^+$ and CD8$^+$ T cells

from STING GOF mice compared to their WT littermate controls. This analysis revealed a

transcriptomic signature of T cell exhaustion in CD4$^+$ and CD8$^+$ T cells in the STING GOF

group, as shown by a negative enrichment of the transcripts upregulated in naive versus

exhausted T cells, as well as a positive enrichment of the genes downregulated in naive versus

exhausted T cells (GSE9650 (27)) (**Fig. 1A**). T cell exhaustion is an altered differentiation

state of chronically activated T cells, notably caused by persistent antigenic stimulation and

sustained inflammation, and is commonly occurring in chronic viral infections or cancers (23,

24). Exhausted T cells co-express inhibitory receptors (like PD-1, TIGIT, TIM-3 and LAG-3)

in a sustained manner (24). Furthermore, exhausted T cells progressively lose their

proliferative and effector functions, which is reminiscent of the functional defects of mature

STING GOF T cells (9, 10).

Analysis of gene expression for key transcription factors associated with T cell

exhaustion in STING GOF T cells revealed overexpression of *Nfatc1 and Nfatc2* compared to

WT cells, without concurrent induction of the AP-1 cofactors *Fos* and *Jun* (**Fig. 1B**). This

pattern is indicative of a preferential activation of the transcriptional program of exhaustion

(28). We also observed increased expression of T cell exhaustion-associated transcription

factors, like EOMES (*Eomes*) and T-BET (*Tbx21*), as well as TOX (*Tox*), the latter being

more involved in the most advanced stages of T cell exhaustion program (**Fig. 1B**). Moreover,

transcripts of multiple genes encoding for inhibitory receptors (*Pdcd1*, *Tigit*, *Havcr2*, *Lag3*,

Cd244, *Entpd1*, *Ctla4* and *Cd160*) were also overexpressed (**Fig. 1C**). Accordingly, the

percentages of splenic $CD4^+$ and $CD8^+$ T cells positive for PD-1, TIGIT, TIM-3 or LAG-3

expression were significantly increased in STING GOF mice compared to their WT controls

(**Fig. 1D**). Then, we analyzed the expression of additional markers to define more precisely

the exhaustion stage of T cells in STING GOF mice. As *Tcf7* transcript was downregulated in

our RNA-seq data (**Fig. 1B**), we hypothesized that STING GOF T cells could be terminally

exhausted. We analyzed co-expression of the inhibitory receptors PD-1 and TIM-3, as well as

the transcription factor TCF-1 (encoded by *Tcf7* gene), to identify progenitor exhausted (PE)

T cells ($PD-1^+TCF-1^+TIM-3^-$) and terminally exhausted (TE) T cells ($PD-1^+TCF-1^-TIM-3^+$)

(29). An increased proportion of TE T cells was observed among $CD4^+$ and $CD8^+$ T cells

from STING GOF mice as compared to control mice. Interestingly, the STING GOF T cell

exhaustion phenotype appeared to be more profound in $CD4^+$ than in $CD8^+$ T cells (**Fig. 1E**).

Together, our findings revealed an unexpected phenotype of T cell exhaustion in STING GOF

mice, which reaches the stage of terminal exhaustion, and might contribute to the T cell

dysfunctions described in these mice.

Given that STING V154M mice develop a lung disease partly mediated by T cells

(17), we also investigated potential T cell exhaustion in the lungs. Interestingly, as in the

spleen, the proportion of $CD4^+$ and $CD8^+$ T cells infiltrating the lungs that expressed PD-1

(data available for $CD4^+$ T cells), TIGIT, TIM-3 or LAG-3 was significantly higher in STING

GOF mice compared to WT controls (**Fig. S1A**).

To determine if this exhaustion phenotype has a central origin, we examined whether

the single positive (SP) T cells, the most terminally differentiated T cell progenitors in the

thymus, were already exhibiting the phenotype. We observed only minimal increase of cells

expressing inhibitory receptors, for both thymic SP CD4$^+$ and CD8$^+$ progenitors (**Fig. S1B**).

In contrast, we demonstrated that splenic T cells from very young (2-week-old) mice, which

represent cells that have just left the thymus for the periphery, were already displaying the

exhaustion phenotype of adult mice (**Fig. S1C**). Altogether, these results demonstrate that the

T cell exhaustion phenotype in STING GOF mice is acquired very early after birth, but only

once T cells have reached the periphery.

Since STING drives type I IFNs, which have recently shown to regulate coinhibitory

receptors expression on T cells (30), we then explored the role of these cytokines in T cell

exhaustion in STING GOF mice. Overall, the T cell exhaustion phenotype was not reversed

by IFNAR deficiency in STING GOF mice (**Fig. S2**).

Ca^{2+}-NFAT pathway is activated by STING, but is not sufficient to induce T cell

exhaustion

We then searched for signals that could induce the exhaustion phenotype in STING

GOF T cells. Considering its crucial role in T cell exhaustion (31) and its upregulation in our

RNA-seq data (**Fig. 1B**), we first focused on NFAT signaling pathway.

Transcriptomic analysis confirmed a significant enrichment of NFAT signature from

the Pathway Interaction Database (PID) in STING GOF T cells, compared to control T cells

(**Fig. S3A**). Upregulation of the total (nuclear and cytosolic) NFATc1 transcription factor

expression was confirmed by flow cytometry (**Fig. S3B**). NFAT activation relies on its

dephosphorylation by the calcium (Ca^{2+})-calmodulin-calcineurin pathway, and Ca^{2+}

homeostasis is known to be disrupted in STING GOF T cells (20). Using Fura Red™-

mediated ratiometric measurement to study basal Ca^{2+} levels, we demonstrated that basal Ca^{2+}

levels were higher in CD4$^+$ and CD8$^+$ T cells from STING GOF mice compared to their WT

controls, suggesting that Ca^{2+}-NFAT signaling pathway in STING GOF T cells participates to

the induction of their exhaustion (**Fig. S3C**).

We next wondered whether the activation of the Ca^{2+}-NFAT signaling pathway could

be due to the STING activation *per se*. Therefore, we treated WT splenocytes with STING

agonist 5,6-dimethylxanthenone-4-acetic acid (DMXAA) *in vitro*. After 4 hours, DMXAA-

treated $CD4^+$ and $CD8^+$ T cells presented an increase of their basal Ca^{2+} levels, similar to

those observed in untreated STING GOF T cells (**Fig. 2A**). After 24 hours, DMXAA-treated

cells also showed upregulation of NFATc1 to the same extent as untreated STING GOF T

cells (**Fig. 2B**). However, STING activation *in vitro* was not sufficient to induce an increase

of key inhibitory receptors PD-1 and TIM-3 (**Fig. 2C**).

In conclusion, even if it can contribute to the Ca^{2+}-NFAT activation, STING activation

in T cells is not sufficient to induce their exhaustion, reinforcing the importance of an

environmental factor that was already suggested comparing thymic versus splenic exhaustion

profile from 2-weeks old STING GOF mice (**Fig. S1C**).

STING GOF T cells displayed features of TCR and IL-7R engagement mediated by the

lymphopenic environment

We then hypothesized that NFAT activation in STING GOF could be the consequence

of TCR signaling secondary to antigenic stimulations (31). Indeed, beyond NFAT

overexpression, transcriptomic analysis confirmed a significant enrichment of a broader TCR

signaling pathway signature from the Wikipathways (WP) in STING GOF T cells compared

to control T cells (**Fig. 3A**). In addition, we highlighted a significant downregulation of CD3

expression (consistent with TCR stimulation) on the surface of STING GOF T cells compared

to control T cells (**Fig. 3B**).

This antigenic stimulation was not secondary to an acute or chronic infection status, as

STING GOF mice were bred and held under specific opportunist pathogen free (SOPF)

environment, and was not associated with cancer as no tumors were detected in these mice.

Thus, we hypothesized that increased TCR stimulation by conventional dendritic cells (cDCs)

could be responsible for this phenotype, similar to what is observed during homeostatic

proliferation of T cells in lymphopenic environments, promoting the reconstitution of the

lymphoid compartment in the periphery (32). Interestingly, we showed a much higher number

of cDCs per CD4$^+$ T cell, and, to a lesser extent per CD8$^+$ T cell, in STING GOF mice

compared to their WT controls, which could enhance antigen-mediated TCR activation (**Fig.**

3C).

Besides antigenic stimulation, IL-7 could also affect T cell fate in STING GOF T-cell

lymphopenic environment, since IL-7R engagement promote T cell homeostatic proliferation

(32). A marked down-regulation of the alpha chain of the receptor (IL-7Rα), reminiscent of

IL-7R engagement, was observed for both CD4$^+$ and CD8$^+$ T cells (**Fig. 3D**). IL-7Rα

expression was already significantly downregulated in CD4$^+$ and CD8$^+$ naive T cells from

STING GOF mice (**Fig. 3E**), indicating that T-cell lymphopenia impacts T cell fate at early

stage in the periphery of STING GOF mice.

Taken together, our results suggest that the lymphopenic environment impacts T cells

in STING GOF mice through TCR and IL-7R engagement, which are known to govern

lymphopenia-induced homeostatic proliferation.

T cell exhaustion phenotype of STING GOF T cell is reversed in a non-lymphopenic

environment

We next wondered whether STING GOF T cells would display an exhausted profile in

a non-lymphopenic context. To this purpose, we took advantage of bone marrow

transplantation experiments. We sorted long term-hematopoietic stem cells (LT-HSCs) from bone marrow (BM) of donor STING GOF or WT littermate control mice (CD45.2$^+$, donor) and co-transplanted them along with supportive WT BM cells (CD45.1$^+$, support) into irradiated WT recipient mice (CD45.1$^+$CD45.2$^+$, host) (**Fig. 4A**). First, we confirmed the hematopoietic intrinsic effect of the STING GOF mutation on splenic T cell reconstitution (10), as very few T cells were derived from the STING GOF LT-HSCs (red dots), while reconstitution was effective from WT LT-HSCs (black dots). However, the lack of T cell reconstitution from the STING GOF LT-HSCs was compensated by transplantation of supportive WT BM-derived T cells (grey dots) (**Fig 4B**). Consequently, absolute numbers of total (including both CD45.1$^+$ and CD45.2$^+$) CD4$^+$ and CD8$^+$ T cells were similar in STING GOF (STING GOF→WT) and WT LT-HSCs (WT→WT) transplantation settings (**Fig 4C**). Interestingly, STING GOF LT-HSCs-derived CD4$^+$ and CD8$^+$ T cells did not overexpress the inhibitory receptors PD-1, TIGIT, TIM-3 or LAG-3 when compared to WT LT-HSCs-derived T cells, or to WT supportive BM-derived T cells (**Fig 4D**). Consistent with this, there were no terminally exhausted (TE) T cells among STING GOF LT-HSC-derived CD4$^+$ and CD8$^+$ T cells (**Fig 4E**).

Nevertheless, because of the WT background of recipient mice in LT-HSC transplantations, radio-resistant stroma was composed of WT cells (**Fig 4A**), which is not the case in STING GOF mice. In order to determine the potential impact of the STING GOF stroma on the T cell exhaustion phenotype, we performed transplantation experiments in which total BM cells from WT donor mice (CD45.1$^+$, donor) were transplanted into STING GOF or WT irradiated recipient mice (CD45.2$^+$, host) (**Fig. S4A**). As expected, a good splenic T-cell reconstitution were observed upon transplantation of WT donor BM cells into STING GOF irradiated host (**Fig. S4B**). In addition, the absolute numbers of CD4$^+$ and CD8$^+$ T cells were comparable in the two groups (WT→WT, and WT→STING GOF) of mice (**Fig.**

S4C). Moreover, the proportion of CD4$^+$ and CD8$^+$ T cells expressing the inhibitory receptors

PD-1, TIGIT, TIM-3 or LAG-3 was similar following transplantation of WT donor BM cells

into irradiated STING GOF mice or WT recipients (**Fig. S4D**), and the proportion of TE T

cells was extremely low in either group of mice (**Fig. S4E**).

Together, these data indicate that STING GOF stroma does not trigger T cell

exhaustion phenotype and rather identify T cell lymphopenia as the main mechanism driving

T cell exhaustion in STING GOF mice.

Mice carrying Rag1 hypomorphic mutations also display lymphopenia-associated T cell

exhaustion

In order to determine if lymphopenia-mediated T cell exhaustion in STING GOF mice

is specific to the STING GOF mutation, we investigated whether another lymphopenic mouse

model might also show T cell exhaustion. To this purpose, we studied two mouse models

carrying hypomorphic *Rag1* mutations (*Rag1*$^{R972Q/R972Q}$ and *Rag1*$^{R972W/R972W}$ mice, thereafter

referred to as RAG1 R972Q and RAG1 R972W mice), also presenting T cell lymphopenia of

central origin, based on early block in T cell development (22) as in STING GOF mice.

Indeed, RAG1 R972Q and R972W mutant mice similar levels of CD4$^+$ and CD8$^+$ T cell

lymphopenia in spleen, compared to STING GOF mice (**Fig. S5A**).

Interestingly, a significant proportion of CD4$^+$ and CD8$^+$ T cells from RAG1 R972Q

and R972W mice was expressing inhibitory receptors compared to their WT controls, albeit

not to the same extent as observed in STING GOF T mice for the majority of inhibitory

receptors studied (**Fig. 5**). TE T cells were also increased among CD4$^+$ and CD8$^+$ T cells for

both RAG1 mutants compared to their WT controls, without reaching the same proportion as

in STING GOF mice (**Fig. S5B**). In conclusion, lymphopenia seems to be sufficient to induce

T cell exhaustion. However, as T cell exhaustion appeared to be more severe in STING GOF

mice compared to RAG1 hypomorphic mutant mice, additional factors beyond central T cell

lymphopenia must contribute to the higher degree of T cell exhaustion observed in STING

GOF mice.

Discussion

Our data revealed an unexpected T cell exhaustion phenotype in STING GOF mice. Usually associated with chronic T cell stimulation (as in the case of chronic viral infections and cancers), this altered state of T cell differentiation turned out to be of particular interest since it could account for some of the functional defects of mature STING GOF T cells, especially their lack of proliferation in response to TCR activation (9). Moreover, we show here that T cell exhaustion in STING GOF mice is type I IFN-independent, as was the T cell proliferation defect (9). Overall, T cell exhaustion could contribute to T cell proliferation defects and apoptosis, in addition to intrinsic effects of the STING protein in mature T cells (18–21).

T cell exhaustion is also associated with a progressive loss of effector properties. In particular, T cell exhaustion may limit IFN-γ production by T cells, which is known to be increased by STING activation (17, 33) and to contribute to the lung disease mortality (17). Therefore, T cell exhaustion could thus be beneficial in STING GOF mice by limiting tissue immunopathology, similarly to the effect of T cell exhaustion in other auto-inflammatory and autoimmune diseases (34). In accordance with this, we also showed an increased proportion of exhausted T cells in lungs of STING GOF mice.

In addition to T cell exhaustion associated with prolonged antigenic stimulations as chronic viral infections or tumoral context (35), we describe here a T cell exhaustion state whose main driver is represented by lymphopenic environment. Like the two other contexts, T cell exhaustion in STING GOF mice appeared to be driven mainly by the environment rather than an intrinsic effect in T cells. Indeed, acute treatment of WT splenocytes with DMXAA *in vitro* had a pre-activating effect on T cells without inducing an exhaustion phenotype. In addition, bone marrow transplantation experiments showed that STING GOF LT-HSCs-derived T cells developing in a predominantly WT and non-lymphopenic

environment do not manifest the exhaustion phenotype. Among environmental factors

potentially implicated in the T cell exhaustion of STING GOF mice, type I IFNs were shown

to be dispensable, despite the potential role of these cytokines suggested in the literature (30,

36, 37). When studying other environmental factors, the STING GOF stroma does not appear

to be sufficient to induce T cell exhaustion.

STING GOF T cells display signs of TCR and IL-7R engagement, consistent with

lymphopenia-associated signals provided by (auto)antigen and IL-7-mediated stimulation,

respectively. While these signals are known to lead to generation of memory-like T cells

through homeostatic proliferation (32), T cell exhaustion appeared as a new phenomenon,

presumably linked to the persistency of these signals. Of note, the observation that the

lymphopenia leads to a higher number of cDCs per CD4$^+$ T cells than for CD8$^+$ T cells may

help explain the more severe T cell exhaustion phenotype observed for CD4$^+$ T cells. Finally,

the observation that mice carrying hypomorphic *Rag1* mutations are lymphopenic and display

obvious signs of T cell exhaustion further supports the notion that lymphopenia may represent

an important mechanism leading to T cell exhaustion.

Nevertheless, T cell exhaustion in STING GOF mice appears to be more severe than

the one observed in the two RAG1 hypomorphic mutant models studied. This difference does

not depend on the severity of the lymphopenia, since the latter was even more profound,

especially in the RAG1 R972W mutant. Thus, the STING GOF mutation may synergize with

lymphopenia in promoting and aggravating T cell exhaustion. An important contributory

mechanism to the phenomenon may be represented by the intrinsic effect of STING on T cell

activation, as suggested by the upregulation of Ca^{2+}-NFAT pathway upon DMXAA treatment.

This NFAT activation had been already highlighted by a previous transcriptomic analysis,

also indicating its independence from type I IFN (38). Thus, the STING GOF mutation could

place the T cells in a pre-activated state, making them not only more sensitive to endoplasmic

reticulum stress and apoptosis (20), but also to exhaustion. Even if we demonstrated that

STING GOF radioresistant cells (stroma) do not trigger T cell exhaustion phenotype by

themselves, synergistic effects of the STING GOF mutation on the antigen presentation

capacity of both non-hematopoietic and hematopoietic cells, especially cDC which are also

included in the radioresistant cell population, could reinforce TCR stimulation in the

lymphopenic context and thus contribute to exhaustion. The impact of this mutation on major

histocompatibility complex (MHC) molecules expression has already been demonstrated in

endothelial cells within the lung environment in STING GOF mice (17). Moreover, STING

activation by agonists in mice has been associated with the loss of cDC1 (39), a subpopulation

that plays a crucial role in maintaining and isolating progenitor exhausted (PE) T cells and

preventing their transition into terminally exhausted (TE) T cells (40).

Overall, our data show that the T cell exhaustion phenotype in STING V154M mice is

the result of poor central production of T cells and will potentially reinforce this lymphopenia

by a self-maintaining loop. These data should be considered in the context of severe T cell

lymphopenia as severe combined immunodeficiencies (SCID). Importantly, recent

descriptions of SCID patients developing T cell exhaustion have reinforced a lymphopenia-

mediated T cell exhaustion mechanism (25, 26). In particular, the analysis of 61 SCID

patients receiving hematopoietic cell transplantation (HCT) recently showed that patients with

poor T cell reconstitution after HCT display an increased frequency of exhausted T cells,

compared to patients with normal T cell counts after HCT, therefore directly linking

lymphopenia with T cell exhaustion (25). In addition, T cell depletion treatments were

identified as an important factor associated with PD1 expression on T cells in patients

receiving HCT (41).

In conclusion, the data and lessons drawn from the STING GOF mice could therefore

be of key importance for the follow up of patients receiving HCT on the one hand, but also

for a better comprehension of potential T cell exhaustion in lymphopenic murine models on

the other hand. Finally, STING GOF mice constitute a good model for the study of T cell

exhaustion in non-infectious and non-tumoral contexts.

Materials and methods

Detailed information on the materials and methods can be found in SI Appendix,

Supporting Materials and Methods.

Mice and cells

Detailed information on mice and cells can be found in SI Appendix, Supporting

Materials and Methods.

Flow cytometry and cell sorting

Splenic, thymic, BM or lung single-cell suspensions were analyzed by flow cytometry

according to standards protocol. Cells were stained with the antibodies listed in SI Appendix

(Supporting Materials and Methods) in PBS (Gibco) supplemented with 2% v/v FBS

(Dutscher). Intracellular of total (nuclear and cytosolic) TCF-1 and NFATc1 stainings were

performed with anti-TCF-1 AF647 (S33-966, BD Pharmingen) and anti-NFATc1 PE (7A6,

BioLegend) antibodies, after fixation and permeabilization of cells using Foxp3/Transcription

Factor Staining Buffer Set (Thermo Fisher Scientific), according to the manufacturer's

instructions. Viability was assessed by Fixable Viability Dye eFluor™ 450 (eBioscience) or

DAPI (1 μM/mL Sigma) stainings. Cells were analyzed using an Attune NxT™ flow

cytometer (Thermo Fisher Scientific). Data were analyzed using FlowJo software (TreeStar).

RNA-seq on sorted CD4$^+$ and CD8$^+$ T cells

Splenocytes from STING GOF mice (n=5) and WT littermate controls (n=5) were

stained as described above with anti-CD3e FITC (clone 145–2 C11, BD Biosciences), anti-

CD4 AF700 (RM4-5, BD Pharmingen), anti-CD8a PE (53-6 .7, BD Pharmingen) and DAPI

(1 μM/mL, Sigma) before positive sorting of live CD3$^+$CD4$^+$ or CD3$^+$CD8$^+$ T cells using a

BD FACSAria™ Fusion flow cytometer (IGBMC Flow Cytometry Facility, Strasbourg,

France). Isolated cells were at least 95% pure. Total RNA was extracted using RNeasy® Plus

Micro Kit (Qiagen) according to the manufacturer's instructions. RNAseq analyses were

performed by the Genomax Facility (INSERM U1109, ImmunoRhumatologie Moléculaire,

Université de Strasbourg, France). Libraries were prepared from 10 ng RNA using SMARTer

Stranded Total RNA-Seq Kit, Pico Input Mammalian (Takara) following the manufacturer's

instructions. Briefly, random primers were used for first-strand synthesis, and ribosomal

cDNA was cleaved by ZapR V.2 in the presence of mammalian R-probes V.2. Libraries were

pooled and sequenced (paired-end 2×75 bp) on a NextSeq 500 using the NextSeq 500/550

High Output Kit V.2 according to the manufacturer's instructions (Illumina). For each

sample, quality control was carried out and assessed with the next-generation sequencing

(NGS) Core Tools FastQC. Reads were aligned against *Mus musculus* mm10 reference

genome using TopHat 2 Aligner (42) and gene expression levels were estimated using

Cufflinks V.2.1.1 (43). Differential expression analysis was performed with Cuffdiff V.2.1.1

after exclusion of fragments per kilobase million below 9. Gene Set Enrichment Analysis

(GSEA) was carried out according to the developer's recommendations. The STING GOF

versus WT FPKMs of all genes detected in the RNA-seq were used for the ranked genes list

and compared with the indicated signatures from the Molecular Signatures Database

(MSigDB), using GSEA Software (UC San Diego, Broad Institute) (44, 45). Raw RNAseq

data have been deposited in the EMBL-EBI ArrayExpress archive (accession no. E-MTAB-

13658).

Cytosolic Ca^{2+} level measurements

For cytosolic Ca^{2+} level measurements, splenic single-cell suspensions were stained as

described above with Fixable Viability Dye eFluor™ 450 (eBioscience), anti-CD4 AF700

(RM4-5, BD Pharmingen) and anti-CD8a PE-CD584 (53-6 .7, BD Pharmingen) before being

loaded with 2 µM Fura Red™ acetoxymethyl ester (AM) (Thermo Fisher Scientific) in PBS

2% FBS containing 0.02% pluronic F-127 (Thermo Fisher Scientific) for 30 min at 37°C,

avoiding light. Loaded cells were washed, resuspended in DMEM (Gibco) containing 1.8 mM

Ca^{2+} and incubated for 10 min at 37°C avoiding light, before acquisition using an Attune

NxT™ flow cytometer (Thermo Fisher Scientific). After an initial 50-sec acquisition

sequence, ionomycin (1 µg/mL, InvivoGen) was added, as positive control, before the

acquisition of a second 60-sec sequence. Ca^{2+} levels were determined according to time by

ratiometric analyzes between VL4 (405 nm-excitation; 660 nm-emission) and BL3 (488 nm-

excitation; 695 nm-emission) channels using FlowJo software (TreeStar).

Splenocyte culture

In vitro splenocytes stimulations were performed in complete RPMI-1640 medium

containing L-glutamine (Lonza) supplemented with 10% v/v FBS (Dutscher), 50 mmol/L β-

mercaptoethanol (Gibco), 1% penicillin/streptomycin (Gibco), and 10 mmol/L HEPES

(Lonza). Splenocytes were stimulated for the indicated times at 37°C, under 5% CO_2, with the

STING agonist 5,6-dimethylxanthenone-4-acetic acid (DMXAA) (10 µg/mL, InvivoGen).

LT-HSCs transplantations with supportive BM cells

For LT-HSCs sorting, total BM cells were stained for lineage marker expression with

biotinylated anti-CD3 (17A2, BioLegend), anti-CD4 (GK1.5, BioLegend), anti-CD8 (53.6.7,

eBioscience), anti-CD11b (M1/70, BioLegend), anti-CD19 (1D3, BioLegend), anti-

CD45R/B220 (RA3-6B2, BioLegend), anti-CD49b (DX5, BioLegend), anti-Gr-1 (Ly-6G/C)

(RB6-8C5, BioLegend), anti-TER-119 (TER-119, BioLegend), before depletion with

Dynabeads sheep anti-rat IgG beads (Thermo Fisher Scientific). Lineage-negative cells were

stained as described above with Streptavidin AF700 (Invitrogen), anti-CD48 FITC (HM48-1,

BioLegend), anti-CD117 (c-Kit) APC (2B8, BioLegend), anti-CD150 (SLAM) PE (TC15-

12F12.2, BioLegend), anti-Sca-1 (Ly-6A/E) PE-Cy7 (D7, BioLegend) and DAPI (1 μM/mL,

Sigma) before positive sorting of live LT-HSCs ($CD48^-CD150^+Lin^-Sca1^+cKit^+$) using a BD

FACSAria™ Fusion flow cytometer (IGBMC Flow Cytometry Facility, Strasbourg, France).

Isolated LT-HSCs were at least 98% pure. 250 STING GOF or WT LT-HSCs from 6-weeks

old $CD45.2^+$ donor mice were i.v. injected with 5×10^5 supporting $CD45.1^+$ WT BM cells

into 9-Gy lethally irradiated $CD45.1^+CD45.2^+$ WT hosts. The spleens of the mice were

harvested 6 months after transplantation and the splenocytes analyzed by flow cytometry as

described above.

Wild-type (WT) total bone marrow cells transplantations

1×10^6 WT total BM cells from 10 to 15-weeks old $CD45.1^+$ donor mice were i.v.

injected into 9-Gy lethally irradiated $CD45.2^+$ STING GOF or WT hosts. Mouse spleens were

harvested 6 months after transplantation and the splenocytes analyzed by flow cytometry as

described above.

Statistical analysis

All data are presented as means +/- SEMs. Statistical analyses were performed with

Prism 8.2.1 software (GraphPad). Statistical significance was calculated with the non-

parametric test of Mann-Whitney test, except for NFATc1 MFI ratio, where statistical

significance was calculated with Wilcoxon signed-rank test with a hypothetical value of 1

(see details in each Figure legends).

Acknowledgements

We thank Fabrice Augé, Christian Galmiche, Delphine Lamon, Manon Lecointe, Fabien

Lhericel, Sophie Reibel-Foisset for excellent animal care. We thank Vincent Gies, Frédéric

Gros, Aurélien Guffroy, Sophie Jung and Thierry Martin (INSERM UMR - S1109,

Strasbourg University, France), Vincent Flacher and Christopher Mueller (CNRS UPR 3572,

Institute of Molecular and Cellular Biology (IBMC), Strasbourg, France) for scientific

discussions. We thank the Flow Cytometry facility of IGBMC (Illkirch, France) for cell

sorting. This work was supported by the Hôpitaux Universitaires de Strasbourg (HUS), by the

"Direction de la Recherche Clinique et de l'Innovation" (HUS) and by grants from Strasbourg

University (UdS), by the Agence Nationale de la Recherche (ANR-14-CE14-0026-04,

Lumugene; ANR-19-CE15-0028, LYMPHO-STING; and ANR-11-EQPX-022). The study

was also supported by the Institut National de la Santé et de la Recherche Médicale

(INSERM), by the Atip-Avenir, by government grants managed by the Agence National de la

Recherche as part of the "Investment for the Future" program (Institut Hospitalo-Universitaire

Imagine, grant ANR-10-IAHU-01, Recherche Hospitalo-Universitaire, grant ANR-18-RHUS-

0010). Luigi D. Notarangelo is supported by the Division of Intramural Research, National

Institute of Allergy and Infectious Diseases, National Institutes of Health.

References

1.   A. Ablasser, Z. J. Chen, cGAS in action: Expanding roles in immunity and
inflammation. *Science* **363**, eaat8657 (2019).

2.   K.-P. Hopfner, V. Hornung, Molecular mechanisms and cellular functions of cGAS–
STING signalling. *Nat Rev Mol Cell Biol* **21**, 501–521 (2020).

3.   Y. Liu, *et al.*, Activated STING in a Vascular and Pulmonary Syndrome. *N Engl J
Med* **371**, 507–518 (2014).

4.   N. Jeremiah, *et al.*, Inherited STING-activating mutation underlies a familial
inflammatory syndrome with lupus-like manifestations. *J. Clin. Invest.* **124**, 5516–
5520 (2014).

5.   Y. J. Crow, D. B. Stetson, The type I interferonopathies: 10 years on. *Nat Rev
Immunol* (2021).

6.   M.-L. Frémond, *et al.*, Overview of STING-Associated Vasculopathy with Onset in
Infancy (SAVI) Among 21 Patients. *The Journal of Allergy and Clinical Immunology:
In Practice* **9**, 803-818.e11 (2021).

7.   M.-L. Frémond, Y. J. Crow, STING-Mediated Lung Inflammation and Beyond. *J Clin
Immunol* **41**, 501–514 (2021).

8.   C. David, M.-L. Frémond, Lung Inflammation in STING-Associated Vasculopathy
with Onset in Infancy (SAVI). *Cells* **11**, 318 (2022).

9.   D. Bouis, *et al.*, Severe combined immunodeficiency in stimulator of interferon genes
(STING) V154M/wild-type mice. *Journal of Allergy and Clinical Immunology* **143**,
712-725.e5 (2019).

10.   M. Motwani, *et al.*, Hierarchy of clinical manifestations in SAVI N153S and V154M
mouse models. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 7941–7950 (2019).

11.   J. D. Warner, *et al.*, STING-associated vasculopathy develops independently of IRF3
in mice. *Journal of Experimental Medicine* **214**, 3279–3292 (2017).

12.   H. Luksch, *et al.*, STING-associated lung disease in mice relies on T cells but not type
I interferon. *Journal of Allergy and Clinical Immunology* **144**, 254-266.e8 (2019).

13.   B. G. Bennion, *et al.*, A Human Gain-of-Function STING Mutation Causes
Immunodeficiency and Gammaherpesvirus-Induced Pulmonary Fibrosis in Mice. *J
Virol* **93**, e01806-18 (2019).

14.   G. R. Martin, *et al.*, Expression of a constitutively active human *STING* mutant in
hematopoietic cells produces an *Ifnar1* -dependent vasculopathy in mice. *Life Sci.
Alliance* **2**, e201800215 (2019).

15.   H. Siedel, A. Roers, A. Rösen-Wolff, H. Luksch, Type I interferon-independent T cell
impairment in a Tmem173 N153S/WT mouse model of STING associated

vasculopathy with onset in infancy (SAVI). *Clinical Immunology* **216**, 108466 (2020).

16.   B. G. Bennion, *et al.*, STING Gain-of-Function Disrupts Lymph Node Organogenesis
534   and Innate Lymphoid Cell Development in Mice. *Cell Reports* **31**, 107771 (2020).

535   17.   K. M. Gao, M. Motwani, T. Tedder, A. Marshak-Rothstein, K. A. Fitzgerald,
536   Radioresistant cells initiate lymphocyte-dependent lung inflammation and IFNγ-
537   dependent mortality in STING gain-of-function mice. *Proc. Natl. Acad. Sci. U.S.A.*
538   **119**, e2202327119 (2022).

539   18.   S. Cerboni, *et al.*, Intrinsic antiproliferative activity of the innate sensor STING in T
540   lymphocytes. *Journal of Experimental Medicine* **214**, 1769–1785 (2017).

541   19.   M. F. Gulen, *et al.*, Signalling strength determines proapoptotic functions of STING.
542   *Nat Commun* **8**, 427 (2017).

543   20.   J. Wu, *et al.*, STING-mediated disruption of calcium homeostasis chronically activates
544   ER stress and primes T cell death. *Journal of Experimental Medicine* **216**, 867–883
545   (2019).

546   21.   A. R. Concepcion, *et al.*, The volume-regulated anion channel LRRC8C suppresses T
547   cell function by regulating cyclic dinucleotide transport and STING–p53 signaling.
548   *Nat Immunol* **23**, 287–302 (2022).

549   22.   L. M. Ott De Bruin, *et al.*, Hypomorphic Rag1 mutations alter the preimmune
550   repertoire at early stages of lymphoid development. *Blood* **132**, 281–292 (2018).

551   23.   E. J. Wherry, M. Kurachi, Molecular and cellular insights into T cell exhaustion. *Nat*
552   *Rev Immunol* **15**, 486–499 (2015).

553   24.   L. M. McLane, M. S. Abdel-Hakeem, E. J. Wherry, CD8 T Cell Exhaustion During
554   Chronic Viral Infection and Cancer. *Annu. Rev. Immunol.* **37**, 457–495 (2019).

555   25.   R. Labrosse, *et al.*, Aberrant T-cell exhaustion in severe combined immunodeficiency
556   survivors with poor T-cell reconstitution after transplantation. *Journal of Allergy and*
557   *Clinical Immunology* **151**, 260–271 (2023).

558   26.   W. Dong, *et al.*, Exhaustion-like dysfunction of T and NKT cells in an X-linked severe
559   combined immunodeficiency patient with maternal engraftment by single-cell
560   analysis. *Int J Mol Med* **51**, 25 (2023).

561   27.   E. J. Wherry, *et al.*, Molecular Signature of CD8+ T Cell Exhaustion during Chronic
562   Viral Infection. *Immunity* **27**, 670–684 (2007).

563   28.   G. J. Martinez, *et al.*, The Transcription Factor NFAT Promotes Exhaustion of
564   Activated CD8 + T Cells. *Immunity* **42**, 265–278 (2015).

565   29.   Y. Guo, *et al.*, Metabolic reprogramming of terminally exhausted CD8+ T cells by IL-
566   10 enhances anti-tumor immunity. *Nat Immunol* **22**, 746–756 (2021).

567   30.   T. S. Sumida, *et al.*, Type I interferon transcriptional network regulates expression of
568   coinhibitory receptors in human T cells. *Nat Immunol* **23**, 632–642 (2022).

31. W. Seo, C. Jerin, H. Nishikawa, Transcriptional regulatory network for the
establishment of CD8+ T cell exhaustion. *Exp Mol Med* **53**, 202–209 (2021).

32. C. D. Surh, J. Sprent, Homeostasis of Naive and Memory T Cells. *Immunity* **29**, 848–
862 (2008).

33. I. Benoit-Lizon, *et al.*, CD4 T cell-intrinsic STING signaling controls the
differentiation and effector functions of T $_H$ 1 and T $_H$ 9 cells. *J Immunother Cancer*
10, e003459 (2022).

34. E. F. McKinney, J. C. Lee, D. R. W. Jayne, P. A. Lyons, K. G. C. Smith, T-cell
exhaustion, co-stimulation and clinical outcome in autoimmunity and infection. *Nature*
523, 612–616 (2015).

35. C. U. Blank, *et al.*, Defining 'T cell exhaustion.' *Nat Rev Immunol* **19**, 665–674
(2019).

36. S. Terawaki, *et al.*, IFN-α Directly Promotes Programmed Cell Death-1 Transcription
and Limits the Duration of T Cell-Mediated Immunity. *The Journal of Immunology*
186, 2772–2779 (2011).

37. E. B. Wilson, *et al.*, Blockade of Chronic Type I Interferon Signaling to Control
Persistent LCMV Infection. *Science* **340**, 202–207 (2013).

38. J. Wu, N. Dobbs, K. Yang, N. Yan, Interferon-Independent Activities of Mammalian
STING Mediate Antiviral Response and Tumor Immune Evasion. *Immunity* **53**, 115-
126.e5 (2020).

39. B. Jneid, *et al.*, Selective STING stimulation in dendritic cells primes antitumor T cell
responses. *Sci. Immunol.* **8**, eabn6612 (2023).

40. S. Dähling, *et al.*, Type 1 conventional dendritic cells maintain and guide the
differentiation of precursors of exhausted T cells in distinct cellular niches. *Immunity*
55, 656-670.e8 (2022).

41. F. Simonetta, *et al.*, Dynamics of Expression of Programmed Cell Death Protein-1
(PD-1) on T Cells After Allogeneic Hematopoietic Stem Cell Transplantation. *Front.*
Immunol. **10**, 1034 (2019).

42. D. Kim, *et al.*, TopHat2: accurate alignment of transcriptomes in the presence of
insertions, deletions and gene fusions. *Genome Biol* **14**, R36 (2013).

43. C. Trapnell, *et al.*, Transcript assembly and quantification by RNA-Seq reveals
unannotated transcripts and isoform switching during cell differentiation. *Nat*
Biotechnol **28**, 511–515 (2010).

44. V. K. Mootha, *et al.*, PGC-1α-responsive genes involved in oxidative phosphorylation
are coordinately downregulated in human diabetes. *Nat Genet* **34**, 267–273 (2003).

45. A. Subramanian, *et al.*, Gene set enrichment analysis: A knowledge-based approach
for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U.S.A.* **102**,
15545–15550 (2005).

FIGURES

Figure 1. Splenic CD4+ and CD8+ T cells from STING GOF mice display an exhausted phenotype.

(**A-C**) Analysis of RNA-seq performed on sorted splenic CD4$^+$ or CD8$^+$ T cells from STING GOF mice (n=5) and their WT littermate controls (n=5). (**A**) Gene Set Enrichment Analysis (GSEA) of the signatures of genes upregulated or downregulated in naive versus exhausted CD8$^+$ T cells (GSE9650) among genes deregulated in STING GOF versus WT CD4$^+$ or CD8$^+$ T cells. Enrichment plots, normalized enrichment score (NES) and nominal p-value (p-value) are shown for each analysis. (**B**) Heatmap representing T cell exhaustion/dysregulated transcription factors mRNA expression (Z-score) by CD4$^+$ or CD8$^+$ T cells in STING GOF mice compared to WT mice. (**C**) Heatmap representing inhibitory receptors mRNA expression (Z-score) by CD4$^+$ or CD8$^+$ T cells in STING GOF and WT mice. (**D-E**) Immunophenotyping of splenic T cells from STING GOF mice and their WT littermate

controls by flow cytometry. **(D)** Proportion of PD-1-, TIGIT-, TIM-3- and LAG-3-expressing
cells among CD4$^+$ or CD8$^+$ T cells. Representative histograms are shown. **(E)** Proportion of
terminally exhausted (TE) T cells among CD4$^+$ or CD8$^+$ T cells. Representative contour plots
are shown. Each data point corresponds to one mouse; mean $\pm$ SEM are shown per population
for seven to twelve mice from multiple independent experiments. Statistical significances are
calculated with two-tailed Mann-Whitney test: ***, P < 0.001; ****, P < 0.0001.

Figure 2. Ca^{2+}-NFAT pathway is activated by STING, but is not sufficient to induce T cell exhaustion.

(A-C) Splenic WT splenocytes were treated *in vitro* with DMXAA (10 μg/mL) and compared to untreated (NT) WT and STING GOF splenocytes. After 4 hours (calcium levels measurement) or 24 hours (immunophenotyping), cells were analyzed by flow cytometry. **(A)** Relative cytosolic Ca^{2+} levels monitored by Fura Red™ ratio in CD4$^+$ or CD8$^+$ T cells Splenocytes were treated, recorded, and plotted as in Fig. S3C. Each data point represents the mean of three independent experiments, and their error bars the SEM. **(B)** Ratio of total NFATc1 MFI in CD4$^+$ or CD8$^+$ T cells. Ratio was normalized on the corresponding WT untreated control. **(C)** Proportion of PD-1- and TIM-3-expressing cells among CD4$^+$ or CD8$^+$ T cells. Each data point corresponds to one mouse; mean ± SEM are shown per population for four mice from four independent experiments. For NFATc1 MFI ratio, statistical significances are calculated with Wilcoxon signed-rank test with a hypothetical value of 1: ns, $P > 0.05$. For PD-1$^+$ and TIM-3$^+$ proportions, statistical significances are calculated with two-tailed Mann-Whitney test: ns, $P > 0.05$; *, $P < 0.05$.

Figure 3. STING GOF T cells displayed features of TCR and IL-7R engagement mediated by the lymphopenic environment.

(**A**) GSEA of the TCR pathway signature (WP) among genes deregulated in STING GOF versus WT CD4$^+$ or CD8$^+$ T cells. Enrichment plots, normalized enrichment score (NES) and nominal p-value (p-value) are shown for each analysis. (**B**) Ratio of CD3 mean fluorescence intensity (MFI) on splenic CD4$^+$ or CD8$^+$ T cells from STING GOF mice and their WT littermate controls. Ratio was normalized on the mean of WT controls of each analysis. Representative histograms are shown. (**C**) Ratio of cDCs absolute number per CD4$^+$ or CD8$^+$ T cells numbers from the spleen of 4-week-old mice. (**D-E**) Proportion of IL-7Rα-expressing cells among splenic total (**D**) and naive (CD44$^-$CD62L$^+$) (**E**) CD4$^+$ or CD8$^+$ T cells from STING GOF mice and their WT littermate controls. Gating strategy of naive T cells and representative histograms are shown. Each data point corresponds to one mouse; mean ± SEM are shown per population for six to nine mice from multiple independent experiments. Statistical significances are calculated with two-tailed Mann-Whitney test: **, P < 0.01; ****, P < 0.0001.

Figure 4. T cell exhaustion phenotype of STING GOF T cell is reversed in a non-lymphopenic environment.

(A-E) LT-HSCs from STING GOF mice and their WT littermate controls (CD45.2[+]) were transplanted with WT BM supportive cells (CD45.1[+]) in WT irradiated hosts to generate STING GOF→WT and WT→WT mice. 6 months later, spleens were assessed for T cell reconstitution and immunophenotyping. (A) Strategy of LT-HSCs transplantations with supportive BM cells. (B) Proportion of cells derived from donor (CD45.2[+]) LT-HSCs or WT

BM supportive cells (CD45.1$^+$) among splenic CD4$^+$ or CD8$^+$ T cells from STING
GOF→WT and WT→WT mice. **(C)** Absolute numbers of total (CD45.2$^+$ and CD45.1$^+$)
splenic CD4$^+$ or CD8$^+$ T cells from STING GOF→WT and WT→WT mice. **(D)** Proportion of
PD-1-, TIGIT-, TIM-3- and LAG-3-expressing cells among splenic CD4$^+$ or CD8$^+$ T cells **(E)**
Proportion of terminally exhausted (TE) T cells among splenic CD4$^+$ or CD8$^+$ T cells. Each
data point corresponds to one mouse; mean ± SEM are shown per population for seven to
eighteen mice from multiple independent experiments. Statistical significances are calculated
with two-tailed Mann-Whitney test: ns, P > 0.05; *, P < 0.05; ****, P < 0.0001.

Figure 5. Mice carrying Rag1 hypomorphic mutations also display lymphopenia-associated T cell exhaustion.

Proportion of PD-1-, TIGIT-, TIM-3- and LAG-3-expressing cells among splenic CD4$^+$ or CD8$^+$ T cells from hypomorphic RAG1 R972Q and R972W mice and their WT littermate controls by flow cytometry. Data were compared with previous results obtained from STING GOF mice and their WT littermate controls. Each point corresponds to one mouse; mean ± SEM are shown per population for four to twelve mice from multiple independent experiments. Statistical significances are calculated with two-tailed Mann-Whitney test: ns, P > 0.05; *, P < 0.05; **, P < 0.01; ***, P < 0.001.

34