Two methodologies and timescales show no promotion of antibiotic resistance from in-water coral disease treatments

Karen L. Neely[1*], Christina A. Kellogg[2], Julie J. Voelschow[2], Allison R. Cauvin[3], Sydney A.
M. Reed[4], Ewelina Rubin[4], Julie L. Meyer[4]

[1] National Coral Reef Institute, Nova Southeastern University, Dania Beach FL, USA

[2] U.S. Geological Survey, St. Petersburg Coastal and Marine Science Center, St. Petersburg FL,
USA

[3] School of Natural Resources and Environment, University of Florida, Gainesville, FL, USA

[4] Soil, Water, and Ecosystem Sciences Department, University of Florida, Gainesville, FL, USA

* Correspondence
Karen L. Neely
kneely0@nova.edu

Running Page Head: ARGs in corals
12
Abstract

The decimation of reefs caused by stony coral tissue loss disease prompted the use of a topical
amoxicillin treatment to prevent coral mortality. Application of this treatment led to concerns
about unintentional impacts such as potential alteration of the coral microbiome and possible
spread of antibiotic resistance. We used two different methodologies – microbial RNA
sequencing and microbial qPCR array – to assess these concerns. RNA sequencing was
conducted on coral mucus samples collected before and 24 hours after amoxicillin application on
wild *Montastraea cavernosa*. No differences in expression of antibiotic resistance genes (ARGs)
were detected. Additionally, there were no notable changes in the microbial communities
between the before and after samples. Microbial qPCR array was used to assess differences in
ARGs over longer timescales in wild *Colpophylia natans*, comparing never-treated corals with
ones treated a single time seven months prior and with those treated multiple times seven months
and more prior. No clinically relevant ARGs represented in the arrays were detected across any
samples. A small number of above-detection reads (4 in the never-treated corals, 2 in the once-
treated corals, and 0 in the multi-treated corals) may indicate weak amplification of similar
environmental (non-anthropogenic) ARGs in the corals. Results indicate that the localized
topical application of amoxicillin to prevent mortality of SCTLD-affected corals is neither
disrupting the microbiome of treated corals nor driving the proliferation of antibiotic resistance.
31

32 **Keywords:** antibiotic resistance genes, ARG, AMR, amoxicillin, stony coral tissue loss disease,
33 microarray, metatranscriptomics, RNAseq, coral reef, Florida

Introduction

The outbreak of stony coral tissue loss disease (SCTLD) has caused unprecedented mortality across the majority of stony coral species in the Caribbean, including more than 20 of the approximately 45 species of reef-builders and 5 of the U.S. Endangered Species-listed Caribbean coral species (Florida Coral Disease Response Research & Epidemiology Team, 2018). The disease originated in South Florida in 2014 (Precht et al., 2016), but since 2017 has spread throughout much of the greater Caribbean region (Papke et al., 2024).

Mortality rates of 67-100% of some species (Precht et al., 2016), along with significant losses to coral cover, diversity, and ecosystem function, triggered unprecedented efforts to save individual colonies using novel methods. Antibiotics added to an aquarium with SCTLD-affected corals halted disease progression (Miller et al., 2020), and a topical paste was subsequently developed specifically for delivering amoxicillin to diseased coral tissues. The paste, termed Base2b, was tested in the field on various species and found to be 67–100% effective at halting SCTLD lesions (Neely et al., 2020; Neely et al., 2021; Shilling et al., 2021; Walker et al., 2021). Though in some instances new lesions can appear on previously-treated corals, sites that are regularly visited can address these, leading to high survival of corals (Neely, 2023). The application of this antibiotic paste has since become the standard for treatment of SCTLD throughout the Caribbean. As of October 2024, over 25,000 corals have been treated with this method on Florida's Coral Reef (Florida Fish and Wildlife Research Institute, 2024).

Though these successes are encouraging, there are management concerns about the deployment of antibiotics onto coral reefs. These include the short- and/or long-term disruption of the coral microbiome, as well as the potential short- and/or long-term development of antibiotic resistance genes (ARGs). Assessments of these potential unintended consequences were identified by the four leading state and federal agencies as research priorities for Florida's Coral Disease Response in 2021, 2022, and 2023. The concerns also delayed or prohibited use of these treatments in some Caribbean jurisdictions (Lee Hing et al., 2022).

The widespread usage of antibiotics in medical and agricultural fields has prompted great concerns over the development of antibiotic resistance in our environment (Kümmerer, 2003; Zhang et al., 2022). The appearance of resistant microbes over the course of more than 80 years of human use has given the incorrect impression that development of antibiotic/antimicrobial resistance genes (ARGs, also known as AMR) is a recent occurrence. However, studies have revealed that antibiotic resistance is ancient, detecting a variety of ARGs in 30,000-year old permafrost and deep within caves that have been isolated for millions of years (D'Costa et al., 2011; Bhullar et al., 2012). The likely ubiquitous presence of ARGs in natural systems relates to their extensive role in common competitive microbe-microbe interactions (Granato et al., 2019) as well as additional physiological functions outside of their protective role (e.g., encoding efflux pumps or signal trafficking; Martínez (2008)). Nevertheless, anthropogenic antibiotic use strongly influences microbial genetic selection which can result in clinically relevant antibiotic resistance.

Antibiotics are commonly used in the treatment of human ailments as well as illnesses of pets and livestock. Additionally, they are frequently used prophylactically in agriculture and aquaculture or as growth promoters (Kümmerer, 2003; Cabello, 2006; Kümmerer, 2009a).

Estimates from the late 1990s and early 2000s placed worldwide consumption of antibiotics at
100,000 to 200,000 tons per year, with as little as 10% being for human clinical use (Kümmerer,
2009b). Due to concerns that overuse of antibiotics in animal feeds could compromise their use
in human medicine, their use as growth promoters has been banned in a number of European
countries (Kümmerer, 2009b). Given that a bacterium under optimal conditions can divide as
rapidly as every 20 to 30 min, it only takes about 20 generations to reach over 1 million cells; so
theoretically a cell containing an ARG could become dominant in a population within as little as
10 hours (Pray, 2008). Use of antibiotics for treatment of wildlife diseases, however, is
uncommon. In general, disease outbreaks in wildlife are considered natural and left to run their
course. Exceptions include when those outbreaks may also affect human, pet, or agricultural
health, or are impacting highly endangered or managed species (as reviewed in Neely et al.
(2021)).

It is currently unknown what sort of selective effects the use of topical amoxicillin may have on
coral microbiomes or ARGs, nor what kind of background levels and diversity of these genes
might already exist in corals. We tested whether the application of amoxicillin+Base2b on coral
disease lesions caused a change in the coral microbiome in tissues adjacent to the lesion. We also
used two different methods, metatranscriptomics and quantitative polymerase chain reaction
(qPCR) microarrays, to assess whether there was an increase in the abundance or diversity of
ARGs as a result of these small, periodic treatments.

96
Methods:
Sample Sites and Collections
Coral tissue and mucus samples were taken from two patch reefs within the Upper Keys section
of the Florida Keys National Marine Sanctuary: Hen and Chickens reef, and Cheeca Rocks. The
two reefs are 7.5 km from each other (Figure 1a-c). Both lie 2-3 km from the nearest point of
land, range in depth from 1.5 – 6.0 meters, and contain relatively high coral cover and coral

Figure 1. Map of sampling sites within the Florida Keys National Marine Sanctuary (a-c). Cores for microbial qPCR array were taken using a hole-punch (d) to remove a small area of tissue near previously treated stony coral tissue loss disease lesions (e).

103 species diversity compared to other Florida reefs. Here we compare the results of two
104 independently designed and executed experiments, one from Hen and Chickens reef and one
105 from Cheeca Rocks, to assess the in situ selective pressure for ARGs during antibiotic treatments
106 for SCTLD.
107
At Hen and Chickens reef, sampling was designed to assess whether there were immediate
changes in microbial community structure or in the expression of ARGs in corals treated with
antibiotics outside of the treatment area. Ten SCTLD-affected colonies of *Montastraea*
cavernosa, all of which had never previously been observed with SCTLD and which had never
been treated, were sampled on July 19, 2022. Mucus-tissue slurry samples were collected by
aspiration with a sterile needle-less 20-ml syringe using gentle agitation of the surface mucus
layer from apparently healthy tissue at least 30 cm from active SCTLD lesions. The surface
mucus layer was chosen as the sample type since it is the interface between the coral holobiont
and the environment and contains the highest density of prokaryotic cells (Garren and Azam,
2010), thus providing the most active microbial community for detection of short-term changes
in microbiome composition and gene expression. After sampling, any SCTLD lesions on the
coral were treated with the standard amoxicillin + Base2b mixture (1:8 ratio by weight). The
compound was topically applied directly to active disease lesions, overlapping the live-dead
tissue margin and adhering to the skeleton and tissue. Twenty-four hours later, the same corals
were revisited and sampled again, at a distance of three coral polyps from the treatment margin.
This sampling time frame was chosen because the treatment was designed to deliver a quick
burst of antibiotic on the first day, followed by slower release over the next two days. Lab
experiments show that more than half of the amoxicillin in the Base2b ointment is delivered
within 24 hours and all of the amoxicillin is delivered within 72 hours of application (Favero and
Curtis, 2020). In addition, 24 hours was hypothesized to be sufficient time for changes to be
detectable in both microbiome composition and gene expression. The distance of three coral
polyps (approximately 2 cm) was chosen as a representative distance for impacts to the coral
microbiome directly outside of the treatment area. On board the dive vessel, syringe samples
were stored in a highly insulated cooler with dry ice immediately. While this method keeps the
samples very cold, the saltwater typically does not freeze. On shore, syringes with mucus-tissue
slurries were set upright inside the cooler to allow settlement from concurrently sampled
seawater. Excess seawater was decanted from the syringes, and the remaining mucus-tissue
slurries were transferred to sterile conical tubes with Zymo DNA/RNA Shield. Samples were
transferred frozen to the University of Florida and stored at -80°C until processing.
137
At Cheeca Rocks, sampling was designed to assess whether antibiotic treatments were creating
long-term selective pressure that would result in detectable antibiotic resistance, and whether
single versus multiple treatments through time would affect this. Samples was taken from corals
at Cheeca Rocks on November 17, 2022. The experimental design included sampling four
colonies (biological replicates) within each of three categories: (i) never treated with amoxicillin
+ Base2b, (ii) treated a single time seven months prior to sampling, and (iii) treated at three
sample times, at least seven months prior (Figure 6). The rationale for sampling several months
after treatment(s) was to ensure that any clinically relevant ARG detected were not due to an
immediate response to the antibiotic, but rather development of lasting resistance, particularly in
the case of corals that had been treated multiple times. A total of 12 samples from *Colpophyllia*
natans were taken. Samples were also taken from 12 *Montastraea cavernosa* colonies following

the same procedures, but the *M. cavernosa* samples were not able to be analyzed due to DNA
extractions yielding largely coral host DNA rather than microbial DNA, and so are not reported
on further.
152
Samples were collected between a depth of 2.7 – 3.9 meters. Samples were collected using a 2
cm hole-punch hammered through the live tissue. For corals that had been previously treated,
samples were taken within 5 cm of the tissue margin which had previously been treated for
SCTLD (Figure 1d-e). All cores were taken on angled slopes of the colonies rather than the tops
or vertical sides. The hole-punch was cleaned between colonies using a disinfecting wipe. Each
extracted core sample was placed into a sterile plastic bag. Upon reaching the surface, seawater
was poured out of each bag and replaced with RNAlater. Bags were sealed and kept on ice back
to land, then placed in a refrigerator for 24 hours to allow fixative to infiltrate the tissue, then
placed in a -20°C freezer until shipment. Preserved samples were shipped frozen with cold packs
to the U.S. Geological Survey's Coral Microbial Ecology laboratory, arriving November 23,
2022 and stored in a -20°C freezer until analysis.
164
165 *Processing and Analyses: Microbial RNAseq and 16S rRNA amplicons (Hen and Chickens Reef)*
166
167 <u>Co-extraction of nucleic acids</u>
168 Mucus-tissue slurries were homogenized using 0.1- and 0.5-mm Bashing Beads lysis tubes
169 (Zymo Research) and vortexed horizontally for 5 min at 2500 rpm on a multi-tube vortex. DNA
170 and RNA were co-extracted from the homogenized slurry with a ZymoBIOMICS DNA/RNA
171 Miniprep Kit (Zymo Research) following the manufacturer instructions.
172
<u>Processing of RNAseq libraries</u>
After extraction, total RNA and DNA were quantified spectrophotometrically using the DS-11
spectrophotometer-fluorometer tool (DeNovix). RNA integrity numbers were determined before
cDNA library construction using Tape Station (Agilent). Barcoded metatranscriptomic libraries
were prepared using Universal Prokaryotic RNA-seq, Any-Deplete kit (NuGen) and sequenced
on two lanes of the Illumina NovaSeq6000 SP-PE-150-bp flow cell at the University of Florida's
Interdisciplinary Center for Biotechnology Research (RRID:SCR_019152). The quality of raw
and filtered sequencing reads was assessed with FastQC v. 0.11.5 (Andrews, 2017). Illumina
adapters and quality trimming were completed using Trimmomatic v. 0.36 (Bolger et al., 2014)
with the following settings: 1) a sliding window of 4 nucleotides averaging to a Phred score of
20, 2) leading and trailing sequence thresholds set to 3 nucleotides, and 3) a minimum of 50 bp
in length. Ribosomal RNA sequences were removed from trimmed sequencing reads using
SortMeRNA v. 4.0 (Kopylova et al., 2012). Non-ribosomal RNA sequence reads were then co-
assembled with Megahit v. 1.1.4 (Li et al., 2016). Protein-coding sequences for the assembled
transcripts were predicted with Prodigal v. 2.6.3 (Hyatt et al., 2010). The translated protein-
coding sequences were queried against NCBI's non-redundant protein database (downloaded in
November 2022) using the Diamond blastp tool v. 2.0.14 (Buchfink et al., 2015). The top
Diamond matches (highest bit score) were used for taxonomic classification with TaxonKit v.
0.9.0 (Shen and Xiong, 2019). Proteins from metatranscriptomes were grouped into six
taxonomic groups (Bacteria, Viruses, Archaea, Cnidaria, Symbiodinicaeace, other Eukaryotes).
Antimicrobial resistance genes in proteins assigned to Bacteria were identified with RGI v. 6.0.2
with the Comprehensive Antibiotic Resistance Database (Alcock et al., 2023). Mapping to all

contigs was done using Bowtie2 v. 2.4.5 (Langmead and Salzberg, 2012) mapper and the
transcript quantification (TPMs – transcript per million) was retrieved using Salmon v. 1.8.0
(Patro et al., 2017). The transcript quantification table was filtered to include only bacterial
transcripts, and differential gene expression analysis was determined between untreated and
treated samples using DESeq2 v. 1.42.1 (Love et al., 2014). The R script for analysis and plotting
of the differential gene expression is available at
https://github.com/meyermicrobiolab/AMR_deseq.
202
Processing of 16S rRNA amplicon libraries
The bacterial community composition of both RNA and DNA fractions of each sample was
characterized by amplicon sequencing of the V4 region of the 16S rRNA gene. Extracted RNA
was converted to cDNA with a ProtoScript II First strand cDNA synthesis kit (New England
Biolabs) before use as a template for PCR. Amplicon libraries were generated as previously
described (Meyer et al., 2019) using the 515F (Parada et al., 2016) and 806RB (Apprill et al.,
2015) Earth Microbiome primers. Amplification was performed in triplicate 25-µl reactions for
each sample with Phusion High-Fidelity Master Mix (New England Biolabs) with 3% dimethyl
sulfoxide, 0.25µM of each primer, and 2µl template. Triplicate reactions were cleaned with a
Qiagen MinElute PCR purification kit and quantified with a DeNovix DS-11 spectrophotometer.
One amplicon pool was generated with 240 ng of each cleaned, barcoded PCR product for
sequencing at the University of Florida's Interdisciplinary Center for Biotechnology Research
(RRID:SCR_019152) on an Illumina MiSeq with 2 x 150bp v. 2 cycle format. Illumina adapters
and primers were removed from raw sequencing reads with cutadapt v. 1.8.1 (Martin, 2011) and
remaining analyses were completed in RStudio v. 2023.12.1+402 with R version 4.3.2.
Amplicon sequence variants were determined with DADA2 v. 1.30.0 (Callahan et al., 2016).
Taxonomy was assigned with the Silva nr99 v. 138.1 database (Yilmaz et al., 2013). Low
abundance amplicon sequence variants with a mean read count across all samples of less than
five were removed from the analysis. Sequencing read count data were transformed by center-
log-ratio and the Aitchison distance of CLR-transformed data was determined with CoDaSeq
(Gloor et al., 2017) and microbial communities were analyzed with phyloseq v. 1.46.0
(McMurdie and Holmes, 2013) and vegan v. 2.6-4 (Dixon, 2003). The full reproducible R script
is available at https://github.com/meyermicrobiolab/AMR-16S-rRNA.
226
227 *Processing and Analyses: Microbial qPCR Array (Cheeca Rocks)*
228
Using sterile technique, preserved coral biopsies were thawed and crushed using a sterile
hammer, and approximately 0.2g of each sample was transferred to a bead-beating tube. DNA
extractions were conducted using Qiagen DNeasy PowerBiofilm kits following the
manufacturer's protocol. DNA concentrations were quantified using the dsDNA Broad Range
Qubit 3.0 fluorometric assay. Samples that yielded DNA concentrations below 10 ng/µL were re-
extracted and quantified.
235
236 DNA samples were screened for antibiotic resistance genes using Qiagen Microbial DNA qPCR
237 array for Antibiotic Resistance Genes (ARGs). The assays in each of the 96-wells of the
238 microarray contain predispensed PCR primers and 5'-hydrolysis probe sets that enable
239 quantitative real-time PCR. Each microarray plate contains assays for 87 antibiotic resistance
240 genes belonging to aminoglycoside, beta-lactam, erythromycin, fluorquinolone, macrolide-

lincosamide-streptogramin B, tetracycline, vancomycin, and multidrug resistance classifications. Beta-lactam resistance genes protect bacteria by hydrolyzing the beta-lactam ring of various compounds, including amoxicillin. The array contains reactions for 57 beta-lactam resistance genes of different classes based on functions and sequence similarity. Additionally, each plate includes positive control wells for the presence of bacterial DNA and positive PCR controls to test for the presence of inhibitors and confirm the efficiency of the reaction. For each array, a reaction mixture was made consisting of 1,275 µL of Qiagen Microbial qPCR Mastermix, 94 µL of sample DNA, and 1,181 µL of Microbial DNA-Free Water, for a total volume of 2,550 µL which allows 25 µL of reaction mix to be added to each well using a multichannel pipet and reagent reservoir. The manufacturer recommends using a minimum of 500 ng genomic DNA from metagenomic samples.

Given that it is not possible to determine how much of the sample DNA concentration was coral host DNA versus bacterial microbiome DNA, we applied the maximum concentration available from each sample (range: 1,100–10,246 ng/µL; median: 2,634 ng/µL). The microarrays were run on an Applied Biosystems StepOnePlus running software version 2.2.2, with baseline set as 8–20 cycles and a threshold setting of 0.2. The cycling conditions consisted of a single initial PCR activation step of 95°C for 10 min, followed by 40 cycles of 95°C for 15 sec denaturation and 60°C for 2 min annealing and extension. Raw C_t values were exported to the Microbial qPCR analysis Excel template to detect the presence of ARGs. The identification criteria were $C_T <32$ = positive, $C_t >37$ = negative, and $C_t >32–<37$ = inconclusive (Zhao et al., 2017). Quality controls included that the six PanBacteria wells had $C_t <37$ (confirming there was sufficient bacterial DNA in the sample to detect ARGs if present in sufficient gene copy numbers) and the positive PCR control (PPC) wells needed a $C_t = 22\pm2$ to confirm the instrument and mastermix performed correctly and that there were no PCR inhibitors in the sample. Per the manufacturer's documentation, most of the ARGs assays had a sensitivity of 20–50 gene copies per target, with a small number of targets requiring 100–200 gene copies for detection. Exact sample DNA concentration values, raw fluorescence values, and raw threshold cycle (C_t) values are available for each sample in the U.S. Geological Survey data release (Kellogg and Voelschow, 2024).

Results:

Microbial RNAseq and 16S rRNA amplicons (Hen and Chickens Reef)
Nucleic acids were extracted from a total of twenty mucus-slurries that were collected from ten *M. cavernosa* colonies before and 24 hours after the application of an antibiotic treatment. RNAseq libraries were successfully generated from 13 out of the 20 samples with RNA integrity numbers > 0.8. A total of 514,841,463 sequencing reads were generated for the 13 RNAseq libraries, with an average of 34,838,281 reads per sample after quality-filtering. Assembled RNAseq libraries with annotations are publicly available in the Joint Genome Institute's Integrated Microbial Genomes and Microbiomes database under the accession numbers listed in the github repository https://github.com/meyermicrobiolab/AMR_deseq. Amplicon libraries were successfully generated from 19 out of 20 DNA fractions and 18 out of 20 RNA fractions. A total of 1,428,607 sequencing reads were generated for the 37 amplicon libraries, with an average of 15,409 reads per sample after quality-filtering. Raw sequencing reads are available in NCBI under Bioproject PRNJA931253.

A total of 42,770
bacterial genes
encoding
antimicrobial
resistance
mechanisms were
expressed. The
most commonly
expressed resistance
mechanisms were
antibiotic efflux and
antibiotic
inactivation (Figure
2). A total of 12,824
transcripts for
antibiotic efflux
genes were detected,
which are used to

Figure 2. Proportion of expressed antimicrobial resistance genes per resistance mechanism for coral colonies before and after treatment.

actively pump out a broad range of antimicrobial compounds. A total of 17,901 transcripts for
antibiotic inactivation genes were detected, which act upon specific classes of antimicrobial
compounds. The third most commonly expressed resistance mechanism was antibiotic target
alteration. Of the 6,726 transcripts annotated as antibiotic target alteration, 57 were transcripts
for penicillin-binding protein mutations conferring resistance to beta-lactam antibiotics. A total
of 261 gene families for antibiotic inactivation were expressed, of which 213 gene families were
beta-lactamases that can inactivate beta-lactam antibiotics like amoxicillin. Expressed
antimicrobial inactivation gene families were very diverse and the most common included OXA
beta-lactamase and nitroimidazole reductase (Figure 3).
314
To determine if the
amoxicillin
treatments induced
short-term changes
in the expression of
ARGs, we tested
differential
expression between
samples collected
before the treatment
and those collected
one day later. No
differentially
expressed ARGs
were detected.
Furthermore, when
ARGs were grouped
by drug class,

Figure 3. Proportion of expressed antimicrobial inactivation gene families in coral colonies before and after treatment. The ten most abundant gene families by gene count are shown; the remaining 203 antimicrobial inactivation gene families are included as "other".

Figure 4. Principal component analysis of microbial communities in *Montastraea cavernosa* corals before and after amoxicillin treatment, faceted by nucleic acid fraction. Half of the variation among communities is shown by the first two principal components (PC1 and PC2).

333 resistance mechanism, or ARGs gene family, no differentially expressed classes, mechanisms, or
334 families were detected.
335
In addition to determining if ARGs expression was impacted by amoxicillin treatment, we also
examined the microbiome composition based on both DNA and RNA fractions before and after
amoxicillin treatment. Overall, 11% of the variation in the microbial community structure was
explained by the nucleic acid fraction (PERMANOVA $R^2 = 0.11$, p = 0.002) while only 7% of
the variation was explained by when the sample was taken (before or after treatment)
(PERMANOVA $R^2 = 0.07$, p = 0.018) (Figure 4). Thus, 82% of the community variation was not
explained by either factor.
343
In both the RNA and DNA fractions, *Montastraea cavernosa* microbial communities at Hen and
Chickens reef were composed primarily of Alphaproteobacteria and Gammaproteobacteria
(Figure S1). Microbial communities in the DNA fraction were composed predominantly of
members of the SAR11 clade, Flavobacteriales, and Blastocatellales, while the RNA fraction
was predominantly composed of Pseudomonadales, Sphingomonadales, and an unclassified
order of Proteobacteria, (Figure 5).
350
Microbial qPCR Array (Cheeca Rocks):
All array plates that were run met the quality requirement that the three positive PCR control
(PPC) replicate assays reached a $C_t = 22 \pm 2$ to confirm the instrument and mastermix performed
correctly and that there were no PCR inhibitors in the samples. However, the *M. cavernosa* DNA
samples did not meet the criteria of $C_t < 37$ for the triplicate assays for the PanBacteria 1 and
PanBacteria 3 controls and therefore the *M. cavernosa* data were not considered valid. Note that
this is likely due to *M. cavernosa* having large polyps which generate a lot of host DNA that can

Figure 5. Relative abundance of amplicon sequence variants, colored by Order or lowest taxonomic classification possible, in corals before and after amoxicillin treatment. Community composition of the DNA fraction is shown on top while composition of the RNA fraction is shown below for each colony.

swamp the microbial DNA signal. Conversely, the *C. natans* DNA samples did meet the criteria
for C_t <37 in the triplicate assays for the PanBacteria 1 and PanBacteria 3 controls, confirming
there was sufficient bacterial DNA in the samples for ARGs detection: PanBacteria1 wells had a
range of 24.18–31.6, with a median value of 28.41 and PanBacteria 3 wells had a range of
25.58–33.38, with a median value of 30.49. Based on the stated identification criteria of C_t <32 =
positive, C_t >37 = negative, and C_t >32–<37 = inconclusive, there were zero ARGs detected in
the 12 *C. natans* samples. However, there were a small number of above-threshold detections of
three genes, *ermB* (20 copy sensitivity), *mefA* (100 copy sensitivity), and *msrA* (100 copy
sensitivity) all of which belong to the classification macrolide-lincosamide-streptogramin_b
(MLS) (Figure 6). Since these assays had extremely high C_t values (38.32–39.51) they may have
resulted from non-specific amplification. It is, however, interesting to note that of the six times
they occurred, four were in samples from corals that had never been treated, and the remaining
two were in corals that had been treated once; none occurred in corals that had been treated
multiple times.
372
Discussion
These experiments were designed to assess the real-world impacts of topical antibiotic
applications on wild SCTLD-affected corals. We specifically targeted two distinct time points -
24 hours after application and seven months after application - to address both immediate and
long-term potential for the development of antibiotic resistance and impacts of intervention
efforts on the coral microbiome.

Treatment History	Sample	2021-05	2021-06	2021-07	2021-08	2021-09	2021-10	2021-11	2021-12	2022-01	2022-02	2022-03	2022-04	2022-05	2022-06	2022-07	2022-08	2022-09	2022-10	2022-11	ermB	mefA	msrA
Never treated	2																				39.51	38.41	-
Never treated	8																				-	-	39.31
Never treated	13																				-	-	-
Never treated	23																				-	39.44	-
Treated Once	7												x							Sampled	39.45	-	-
Treated Once	9												x								-	-	-
Treated Once	14												x								-	-	38.32
Treated Once	17												x								-	-	-
Treated Many	11	x		x									x								-	-	-
Treated Many	12	x		x									x								-	-	-
Treated Many	21			x	x								x								-	-	-
Treated Many	24								x		x		x								-	-	-

Figure 6. Treatment history and above-threshold ARGs detections for *Colpophyllia natans* coral colonies sampled for microbial qPCR array. Shaded boxes with an x indicate date (year-month) when a coral was treated prior to November 2022 sampling. Genes observed outside the acceptable identification criteria ($C_t > 37$) are listed with their C_t values.

379
There are several common analytical methods used to investigate ARGs. In clinical settings it is
common to first isolate candidate microbes and then screen them for resistance to specific
antibiotics. While this technique can be applied to coral-associated bacteria (e.g., Galkiewicz et
al. (2011)), it is limited to those bacterial taxa that are amenable to cultivation. For
environmental assessments, molecular techniques including targeted quantitative polymerase
chain reaction (qPCR) and metagenomics are the preferred methods (Scott et al., 2020). While
molecular techniques cannot assess bacterial viability, they can be scaled to enable high-
throughput of larger sample sizes. Targeted genomic analysis via the commercial qPCR
microarray used in this study allows for rapid screening of more than 80 clinically relevant
ARGs at detection limits ranging from 20–100 gene copies. Running a sample takes roughly 2
hours, followed by minimal analysis using a spreadsheet. However, this system relies on having
prior sequence knowledge of the target ARGs and can only detect the specifically chosen ARGs
on the array, thus limiting the scope of detection to these clinically relevant ARGs. In contrast,
metagenomic analysis is an open-ended tool that can reveal previously undiscovered diversity of
ARGs in an environmental microbial community. However, the post-sequencing bioinformatic
analysis is much more time- and computation-intensive. In addition, metagenomic studies
examining bacterial genes remain technically challenging to conduct with coral microbiota
because of the overwhelming genetic signature of the host (Messer et al., 2024). Therefore, we
used a metatranscriptomic approach which more effectively targets the bacterial RNA fraction
and provides a snapshot of actively transcribed genes.
400
The rationale for using a microarray with 87 target ARGs, of which 57 are beta-lactamase genes
(relevant to amoxicillin; inclusive of classes A, B, C, and D) was to determine if: 1) there was an
increase in abundance or diversity of ARGs directly linked to amoxicillin resistance and 2)
whether, due to multiple ARGs being carried on plasmids, this led to an increase in ARGs to
other classes of antibiotics. Instead, there were zero verified detections for clinical ARGs based
on microarrays. We did observe late amplification (past the cut off criteria of $C_t < 37$; Figure 6)

in five samples for genes (*ermB*, *mefA*, *msrA*) within the macrolide-lincosamide-streptogramin
(MLS) group of resistance genes. These only occurred in the never-treated or treated-once corals.
This might be non-specific amplification, but it is worth noting that this group of genes is
extremely common in environmental bacteria (defined as genera and species primarily found
outside of humans and animals) (Roberts, 2011). We surmise that these late amplifications could
be instances of array primers picking up environmental genes that are similar but not exact
matches to their optimized targets (Aminov, 2009). Persistence of certain bacteria during
laboratory experiments involving antibiotic treatment of corals implied the presence of some
naturally antibiotic-resistant bacteria in coral microbiomes (Sweet et al., 2011; Connelly et al.,
2022; Connelly et al., 2023).
417
To date, only a handful of studies have examined ARGs associated with corals or coral reef
ecosystems, but these all suggest that the presence of ARGs in marine bacteria are likely
common and widespread. For example, isolates of the coral pathogen *Vibrio coralliilyticus* have
demonstrated antimicrobial resistance to 26 clinically relevant antibiotics, including amoxicillin
(Vizcaino et al., 2010). In addition, ARGs for eight clinically relevant antibiotics were detected
by quantitative PCR in microorganisms from both seawater (Liu et al., 2020) and microplastics
(Zhou et al., 2024) from coral reefs in China. Here, even with fewer than 20 metatranscriptomic
libraries we detected the expression of over 42,000 unique bacterial ARGs. Given that the
majority of currently known antibiotic-inactivation ARGs encode resistance to beta-lactams, it is
not surprising that we found genes for resistance to amoxicillin, even in untreated corals. These
findings suggest that expressed ARGs are extremely diverse and common in the environment,
regardless of small-scale, localized treatments.
430
Overuse and misuse of antibiotics in human and veterinary medicine, agriculture, and
aquaculture are considered the main cause of the rise in antimicrobial-resistant pathogens and
development of environmental reservoirs of clinically-relevant ARGs (Thanner et al., 2016;
Dadgostar, 2019). The contributions of topical antimicrobials to this rise are largely unknown
due to insufficient data (Blackburn et al., 2023). Beta-lactam antibiotics like amoxicillin make up
the largest share of human use antibiotics in most countries, accounting for more than half of
total antibiotic use (Kümmerer, 2009a). Beta-lactam ARGs are the most commonly detected
resistance genes in the marine environment, based on a meta-analysis of metagenomic data from
different biomes (Zhang et al., 2022). However, high amounts of clinically relevant ARGs are
primarily detected in areas of high human interaction, such as hospitals, wastewater, and areas
directly impacted by large scale farming (Zhang et al., 2022). The amount of selective pressure
being exerted by a spatially limited dose of antibiotic paste, compared to these sources, or to an
aquaculture situation where higher doses of antibiotics are being consistently employed either in
medicated feed or via bath immersion (Chen et al., 2018), is a comparatively negligible addition
to the environment.

The reflexive expectation that treating corals with antibiotic paste would result in selective
pressure favoring antibiotic-resistant bacteria and promotion of horizontal gene transfer of ARGs
via plasmids needs to be examined. We show here that the small, focused treatments used to treat
SCTLD-affected corals are not increasing the diversity or abundance of ARGs at either
immediate (24 hour) nor longer-term (7+ month) time frames. As elevated ARGs are not found
even in the directly-treated coral tissues, it is extremely unlikely that they could be dispersing
farther afield into surrounding organisms, sediment, or the water column. Beta-lactams are

typically only detected in low concentrations in the environment and are easily hydrolyzed at
ambient temperature (Längin et al., 2009; Bergheim et al., 2010). This suggests that any
amoxicillin that leaches out of the Base2b paste is not only rapidly diluted by seawater but also
chemically broken down.

There is evidence from a variety of studies that the coral-associated microbial community can be
resilient to environmental stressors and can return to the original community composition after
disturbance (Garren et al., 2009; Ziegler et al., 2019; Strudwick et al., 2022). After deliberate
removal of >99% of the coral-associated bacteria via a 6-day immersion treatment of
ciprofloxacin, Sweet et al. (2011) identified substantial recovery of the bacterial community by
day 4 (the end of the experimental monitoring). In another experiment that treated corals in
aquaria with an antibiotic mixture for 8 days and then returned them to the reef, most of the
treated corals succumbed to bleaching or mortality, but the few survivors regained a mucus
microbiome similar to pretreatment by 28 days (Glasl et al., 2016). A similar experiment that
exposed corals to an antibiotic cocktail for a week found that the mucus microbiome recovered
to its initial state within 2 weeks (Bent et al., 2021). Our results suggest a negligible effect of
treatments on the microbiome, probably due to the topical nature of application. The use of a
slow-release antibiotic paste applied only to lesions, rather than full colony immersion in liquid
antibiotics for 3 to 7 days, did not result in the dramatic reduction in bacterial diversity seen in
prior laboratory experiments (Sweet et al., 2011; Gilbert et al., 2012; Sweet et al., 2014; Glasl et
al., 2016; Connelly et al., 2022; Connelly et al., 2023). The effects of antibiotics on coral health
via disruption of the microbiome observed in prior tank experiments appear to be a direct
consequence of both the longer antibiotic exposure times and the choice to uniformly expose the
entire coral fragment.

SCTLD is known to be extremely lethal to affected corals (Precht et al., 2016; Aeby et al., 2019;
Thome et al., 2021), and in-water topical antibiotic intervention offers the only known tool for
preventing that mortality. As managers are considering the use of this tool and weighing the pros
with the potential cons of its use, the spread of antibiotic resistance may be a topic of discussion.
This study identifies that within the parameters (location, species, and timeframes) we tested,
development of antibiotic resistance through these treatments is unlikely to be a factor of
concern.

Acknowledgments
We are grateful for additional field assistance by Sydney Gallagher and Michelle Dobler. This
work was supported by the U.S. Geological Survey's Ecosystems Biological Threats and
Invasive Species Research Program (CAK). Funding support was also provided by Florida
Department of Environmental Protection Office of Resilience and Coastal Protection – Southeast
Region (JM). Collections were conducted opportunistically under a Florida DEP grant (KLN).
Additional time was donated by the author (KLN). All work was conducted under Florida Keys
National Marine Sanctuary permits FKNMS-2022-066-A1 and FKNMS-2019-177-A3. Any use
of trade, firm, or product names is for descriptive purposes only and does not imply endorsement
by the U.S. Government.

Data Availability Statement
494 The raw data supporting the conclusions of this article will be made available by the authors,
495 without undue reservation. Raw sequencing reads from the RNAseq and 16S rRNA amplicon

studies are publicly available at NCBI under BioProject PRJNA931253. Raw fluorescence data and cycle threshold data from microarray study is publicly available via USGS data release https://doi.org/10.5066/P9AST28V.

References

Aeby, G., Ushijima, B., Campbell, J.E., Jones, S., Williams, G., Meyer, J.L., Hase, C., and Paul, V. (2019). Pathogenesis of a tissue loss disease affecting multiple species of corals along the Florida Reef Tract. *Frontiers in Marine Science* 6.

Alcock, B.P., Huynh, W., Chalil, R., Smith, K.W., Raphenya, A.R., Wlodarski, M.A., Edalatmand, A., Petkau, A., Syed, S.A., and Tsang, K.K. (2023). CARD 2023: expanded curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Research* 51**,** D690-D699.

Aminov, R.I. (2009). The role of antibiotics and antibiotic resistance in nature. *Environmental Microbiology* 11**,** 2970-2988.

Andrews, S. (2017). "FastQC: a quality control tool for high throughput sequence data. 2010", in: *http://www.bioinformatics.babraham.ac.uk/projects/fastqc/*.).

Apprill, A., Mcnally, S., Parsons, R., and Weber, L. (2015). Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology* 75**,** 129-137.

Bent, S.M., Miller, C.A., Sharp, K.H., Hansel, C.M., and Apprill, A. (2021). Differential patterns of microbiota recovery in symbiotic and aposymbiotic corals following antibiotic disturbance. *mSystems* 6**,** 10.1128/msystems. 01086-01020.

Bergheim, M., Helland, T., Kallenborn, R., and Kümmerer, K. (2010). Benzyl-penicillin (penicillin G) transformation in aqueous solution at low temperature under controlled laboratory conditions. *Chemosphere* 81**,** 1477-1485.

Bhullar, K., Waglechner, N., Pawlowski, A., Koteva, K., Banks, E.D., Johnston, M.D., Barton, H.A., and Wright, G.D. (2012). Antibiotic resistance is prevalent in an isolated cave microbiome. *PloS One* 7**,** e34953.

Blackburn, J., Ousey, K., Patton, D., Moore, Z., and Avsar, P. (2023). What is the evidence that there is antimicrobial resistance associated with the use of topical antimicrobial preparations? *Wound Practice and Research* 31**,** 40-48.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30**,** 2114-2120.

Buchfink, B., Xie, C., and Huson, D.H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nature Methods* 12**,** 59-60.

Cabello, F.C. (2006). Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environmental microbiology* 8**,** 1137-1144.

Callahan, B., Mcmurdie, P., Rosen, M., Han, A., Johnson, A., and Dada, S.H. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13**,** 581-583.

Chen, B., Lin, L., Fang, L., Yang, Y., Chen, E., Yuan, K., Zou, S., Wang, X., and Luan, T. (2018). Complex pollution of antibiotic resistance genes due to beta-lactam and aminoglycoside use in aquaculture farming. *Water Research* 134**,** 200-208.

Connelly, M.T., Mcrae, C.J., Liu, P.-J., Martin, C.E., and Traylor-Knowles, N. (2022).
Antibiotics Alter *Pocillopora* Coral-Symbiodiniaceae-Bacteria Interactions and Cause
Microbial Dysbiosis During Heat Stress. *Frontiers in Marine Science* 8.
Connelly, M.T., Snyder, G., Palacio-Castro, A.M., Gillette, P.R., Baker, A.C., and Traylor-
Knowles, N. (2023). Antibiotics reduce *Pocillopora* coral-associated bacteria diversity,
decrease holobiont oxygen consumption and activate immune gene expression.
Molecular ecology 32**,** 4677-4694.
D'costa, V.M., King, C.E., Kalan, L., Morar, M., Sung, W.W., Schwarz, C., Froese, D., Zazula,
G., Calmels, F., and Debruyne, R. (2011). Antibiotic resistance is ancient. *Nature* 477**,**
457-461.
Dadgostar, P. (2019). Antimicrobial resistance: implications and costs. *Infection and Drug*
Resistance 12**,** 3903-3910.
Dixon, P. (2003). VEGAN, a package of R functions for community ecology. *Journal of*
Vegetation Science 14**,** 927-930.
Favero, M., and Curtis, K. (2020). "Creation and Release Profiles of Novel Biodegradable
Ointments for Lesion Based and Whole Colony Treatment Methods to be Utilized in the
Response to Stony Coral Tissue Loss Disease (SCTLD)". (Miami, FL: Florida DEP).
Florida Coral Disease Response Research & Epidemiology Team (2018). "Case Definition:
Stony Coral Tissue Loss Disease".
(https://floridadep.gov/sites/default/files/Copy%20of%20StonyCoralTissueLossDisease_
CaseDefinition%20final%2010022018.pdf).
Florida Fish and Wildlife Research Institute (2024). "Coral Disease Intervention Dashboard", in:
https://arcg.is/0L1LWX.).
Galkiewicz, J.P., Pratte, Z.A., Gray, M.A., and Kellogg, C.A. (2011). Characterization of
culturable bacteria isolated from the cold-water coral *Lophelia pertusa*. *FEMS*
Microbiology Ecology 77**,** 333-346.
Garren, M., and Azam, F. (2010). New Method for Counting Bacteria Associated with Coral
Mucus. *Applied and Environmental Microbiology* 76**,** 6128-6133.
Garren, M., Raymundo, L., Guest, J., Harvell, C.D., and Azam, F. (2009). Resilience of coral-
associated bacterial communities exposed to fish farm effluent. *PLoS One* 4**,** e7319.
Gilbert, J.A., Hill, R., Doblin, M.A., and Ralph, P.J. (2012). Microbial consortia increase thermal
tolerance of corals. *Marine Biology* 159**,** 1763-1771.
Glasl, B., Herndl, G.J., and Frade, P.R. (2016). The microbiome of coral surface mucus has a key
role in mediating holobiont health and survival upon disturbance. *The ISME Journal* 10**,**
2280-2292.
Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., and Egozcue, J.J. (2017). Microbiome
datasets are compositional: and this is not optional. *Frontiers in Microbiology* 8**,** 294209.
Granato, E.T., Meiller-Legrand, T.A., and Foster, K.R. (2019). The Evolution and Ecology of
Bacterial Warfare. *Current Biology* 29**,** R521-R537.
Hyatt, D., Chen, G.-L., Locascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010).
Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC*
Bioinformatics 11**,** 119.
Kellogg, C.A., and Voelschow, J.J. (2024). "Screening coral tissue samples (*Colpophyllia*
natans) collected from the Florida Keys using a quantitative polymerase chain reaction
(qPCR) antibiotic resistance gene microarray - raw data".

Kopylova, E., Noé, L., and Touzet, H. (2012). SortMeRNA: fast and accurate filtering of
586   ribosomal RNAs in metatranscriptomic data. *Bioinformatics* 28**,** 3211-3217.
587   Kümmerer, K. (2003). Significance of antibiotics in the environment. *Journal of Antimicrobial*
588   *Chemotherapy* 52**,** 5-7.
589   Kümmerer, K. (2009a). Antibiotics in the aquatic environment–a review–part I. *Chemosphere*
590   75**,** 417-434.
591   Kümmerer, K. (2009b). Antibiotics in the aquatic environment–a review–part II. *Chemosphere*
592   75**,** 435-441.
593   Längin, A., Alexy, R., König, A., and Kümmerer, K. (2009). Deactivation and transformation
594   products in biodegradability testing of ß-lactams amoxicillin and piperacillin.
595   *Chemosphere* 75**,** 347-354.
596   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*
597   *Methods* 9**,** 357-359.
598   Lee Hing, C., Guifarro, Z., Dueñas, D., Ochoa, G., Nunez, A., Forman, K., Craig, N., and
599   Mcfield, M. (2022). Management responses in Belize and Honduras, as stony coral tissue
600   loss disease expands its prevalence in the Mesoamerican Reef. *Frontiers in Marine*
601   *Science* 9:883062.
602   Li, D., Luo, R., Liu, C.-M., Leung, C.-M., Ting, H.-F., Sadakane, K., Yamashita, H., and Lam,
603   T.-W. (2016). MEGAHIT v1. 0: a fast and scalable metagenome assembler driven by
604   advanced methodologies and community practices. *Methods* 102**,** 3-11.
605   Liu, S., Su, H., Pan, Y.-F., and Xu, X.-R. (2020). Spatial and seasonal variations of antibiotics
606   and antibiotic resistance genes and ecological risks in the coral reef regions adjacent to
607   two typical islands in South China Sea. *Marine Pollution Bulletin* 158**,** 111424.
608   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
609   dispersion for RNA-seq data with DESeq2. *Genome Biology* 15**,** 1-21.
610   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
611   *EMBnet.journal* 17**,** 3.
612   Martínez, J.L. (2008). Antibiotics and antibiotic resistance genes in natural environments.
613   *Science* 321**,** 365-367.
614   Mcmurdie, P., and Holmes, S. (2013). phyloseq: an R package for reproducible interactive
615   analysis and graphics of microbiome census data. *PLoS One* 8**,** e61217.
616   Messer, L.F., Bourne, D.G., Robbins, S.J., Clay, M., Bell, S.C., Mcilroy, S.J., and Tyson, G.W.
617   (2024). A genome-centric view of the role of the *Acropora kenti* microbiome in coral
618   health and resilience. *Nature Communications* 15**,** 2902.
619   Meyer, J.L., Castellanos-Gell, J., Aeby, G.S., Häse, C., Ushijima, B., and Paul, V.J. (2019).
620   Microbial community shifts associated with the ongoing stony coral tissue loss disease
621   outbreak on the Florida Reef Tract. *bioRxiv***,** 626408.
622   Miller, C., May, L., Moffitt, Z., and Woodley, C. (2020). "Exploratory treatments for stony coral
623   tissue loss disease: pillar coral (*Dendrogyra cylindrus*)". (Charleston, SC: NOAA
624   Technical Memorandum NOS NCCOS 245 and CRCP 37).
625   Neely, K. (2023). "Florida Keys Coral Disease Intervention Strike Teams: Final Report".
626   (Miami, FL: Florida DEP).
627   Neely, K.L., Macaulay, K.A., Hower, E.K., and Dobler, M.A. (2020). Effectiveness of topical
628   antibiotics in treating corals affected by Stony Coral Tissue Loss Disease. *PeerJ* 8**,**
629   e9289.

Neely, K.L., Shea, C.P., Macaulay, K.A., Hower, E.K., and Dobler, M.A. (2021). Short- and
Long-Term Effectiveness of Coral Disease Treatments. *Frontiers in Marine Science* 8.
Papke, E., Carreiro, A., Dennison, C., Deutsch, J.M., Isma, L.M., Meiling, S.S., Rossin, A.M.,
Baker, A.C., Brandt, M.E., and Garg, N. (2024). Stony coral tissue loss disease: a review
of emergence, impacts, etiology, diagnostics, and intervention. *Frontiers in Marine*
Science 10**,** 1321271.
Parada, A.E., Needham, D.M., and Fuhrman, J.A. (2016). Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. *Environmental Microbiology* 18**,** 1403-1414.
Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods* 14**,** 417-419.
Pray, L. (2008). Antibiotic resistance, mutation rates and MRSA. *Nature Education* 1**,** 30.
Precht, W.F., Gintert, B.E., Robbart, M.L., Fura, R., and Van Woesik, R. (2016). Unprecedented
Disease-Related Coral Mortality in Southeastern Florida. *Scientific Reports* 6:31374.
Roberts, M.C. (2011). Environmental macrolide–lincosamide–streptogramin and tetracycline
resistant bacteria. *Frontiers in Microbiology* 2**,** 9102.
Scott, L.C., Lee, N., and Aw, T.G. (2020). Antibiotic resistance in minimally human-impacted
environments. *International Journal of Environmental Research and Public Health* 17**,**
3939.
Shen, W., and Xiong, J. (2019). TaxonKit: a cross-platform and efficient NCBI taxonomy
toolkit. *bioRxiv* 513523.
Shilling, E.N., Combs, I.R., and Voss, J.D. (2021). Assessing the effectiveness of two
intervention methods for stony coral tissue loss disease on *Montastraea cavernosa*.
Scientific Reports 11**,** 1-11.
Strudwick, P., Seymour, J., Camp, E.F., Edmondson, J., Haydon, T., Howlett, L., Le Reun, N.,
Siboni, N., and Suggett, D.J. (2022). Impacts of nursery-based propagation and out-
planting on coral-associated bacterial communities. *Coral Reefs***,** 1-18.
Sweet, M., Croquer, A., and Bythell, J. (2011). Dynamics of bacterial community development
in the reef coral *Acropora muricata* following experimental antibiotic treatment. *Coral*
Reefs 30**,** 1121-1133.
Sweet, M.J., Croquer, A., and Bythell, J.C. (2014). Experimental antibiotic treatment identifies
potential pathogens of white band disease in the endangered Caribbean coral *Acropora*
cervicornis. *Proceedings of the Royal Society B - Biological Sciences* 281.
Thanner, S., Drissner, D., and Walsh, F. (2016). Antimicrobial Resistance in Agriculture. *mBio*
7**,** 10.1128/mbio.02227-02215.
Thome, P.E., Rivera-Ortega, J., Rodríguez-Villalobos, J.C., Cerqueda-García, D., Guzmán-
Urieta, E.O., García-Maldonado, J.Q., Carabantes, N., and Jordán-Dahlgren, E. (2021).
Local dynamics of a white syndrome outbreak and changes in the microbial community
associated with colonies of the scleractinian brain coral *Pseudodiploria strigosa*. *PeerJ* 9**,**
e10695.
Vizcaino, M.I., Johnson, W.R., Kimes, N.E., Williams, K., Torralba, M., Nelson, K.E., Smith,
G.W., Weil, E., Moeller, P.D., and Morris, P.J. (2010). Antimicrobial resistance of the
coral pathogen *Vibrio coralliilyticus* and Caribbean sister phylotypes isolated from a
diseased octocoral. *Microbial Ecology* 59**,** 646-657.

Walker, B.K., Noren, H., Buckley, S., and Pitts, K. (2021). Optimizing stony coral tissue loss
disease (SCTLD) intervention treatments on *Montastraea cavernosa* in an endemic zone.
Frontiers in Marine Science 8**,** 746.
Yilmaz, P., Parfrey, L.W., Yarza, P., Gerken, J., Pruesse, E., Quast, C., Schweer, T., Peplies, J.,
Ludwig, W., and Glöckner, F.O. (2013). The SILVA and "All-species Living Tree
Project (LTP)" taxonomic frameworks. *Nucleic Acids Research* 42**,** D643-D648.
Zhang, Z., Zhang, Q., Wang, T., Xu, N., Lu, T., Hong, W., Penuelas, J., Gillings, M., Wang, M.,
and Gao, W. (2022). Assessment of global health risk of antibiotic resistance genes.
Nature Communications 13**,** 1553.
Zhao, Z., Wang, J., Han, Y., Chen, J., Liu, G., Lu, H., Yan, B., and Chen, S. (2017). Nutrients,
heavy metals and microbial communities co-driven distribution of antibiotic resistance
genes in adjacent environment of mariculture. *Environmental Pollution* 220**,** 909-918.
Zhou, Z., Tang, J., Tang, K., An, M., Liu, Z., Wu, Z., Cao, X., and He, C. (2024). Selective
enrichment of bacteria and antibiotic resistance genes in microplastic biofilms and their
potential hazards in coral reef ecosystems. *Chemosphere* 352**,** 141309.
Ziegler, M., Grupstra, C.G., Barreto, M.M., Eaton, M., Baomar, J., Zubier, K., Al-Sofyani, A.,
Turki, A.J., Ormond, R., and Voolstra, C.R. (2019). Coral bacterial community structure
responds to environmental change in a host-specific manner. *Nature Communications* 10**,**
3092.
693
694

695 **Supplementary Information**
696
697 Figure S1. Relative abundance of amplicon sequence variants, colored by Class, in corals before
698 and after amoxicillin treatment. Community composition of the DNA fraction is shown on top
699 while composition of the RNA fraction is shown below for each colony.
700