Molecular and Functional Asymmetry in Cckar-expressing Vagal Sensory Neurons

Hailey F. Welch[1], Ishwarya Sankaranarayanan[1,2], Veronica M. Hong[1,2], Haniya Qavi[1],
Khadijah Mazhar[1,2], Benedict J. Kolber[1,2], Theodore J. Price[1,2], *Catherine A. Thorn[1]
[1]Department of Neuroscience, The University of Texas at Dallas, 800 W. Campbell Rd. Richardson, TX, 75080, USA
[2]Center for Advanced Pain Studies, The University of Texas at Dallas, 800 W. Campbell Rd.,
Richardson, TX, 75080, USA
*Lead contact / Correspondence: catherine.thorn@utdallas.edu
7

SUMMARY

The vagus nerves are important carriers of sensory information from the viscera to the brain.
Emerging evidence suggests that sensory signaling through the right, but not the left, vagus
nerve evokes striatal dopamine release and reinforces appetitive behaviors. However, the
extent to which differential gene expression within vagal sensory neurons may underlie this
asymmetric reward-related signaling is unknown. Here, we use single-nucleus RNA sequencing,
in situ hybridization, and calcium imaging to identify a single cluster of neurons, defined by co-
expression of *Chrna3* (nicotinic acetylcholine receptor subunit 3) and *Cckar* (cholecystokinin 1
receptor), that is preferentially expressed in the right nodose ganglia of rats and mice. This
neuronal population also expresses several genes implicated in digestive signaling, including
Glp1r and *Sctr*, consistent with a gut-innervating subtype. Our results suggest that right-biased
expression of this unique chemosensing vagal cell type may contribute to asymmetric encoding
of interoceptive rewards by the vagus nerves.

KEYWORDS

23 Vagus nerve, transcriptomics, asymmetry, lateralization, interoception, cholecystokinin

INTRODUCTION

The vagus nerves (cranial nerve X) transmit sensory information from the viscera, including cardiovascular, respiratory, and gastrointestinal (GI) organs, to the central nervous system to maintain homeostatic control over autonomic processes such as heart rate, respiration, and digestion [1–3]. The sensory neurons of the vagus nerve are situated in the jugular and nodose ganglia (NG), located near the base of the skull. These neurons detect a wide range of visceral signals, including mechanical stretch and chemical stimuli, and relay this information to the brainstem to mediate appropriate autonomic and behavioral responses [4–8].

Several studies have demonstrated striking left-right asymmetries in sensorimotor vagal pathways, with the left and right vagus nerves exhibiting distinct patterns of innervation across visceral organs, including the heart, lungs, and gastrointestinal (GI) tract [9–14]. These anatomical differences are thought to contribute to divergent functional outcomes following left vs. right vagal stimulation [15–19]. Of particular interest is the emerging evidence for lateralized processing of visceral reward-related signals. Stimulation of the right vagus nerve, but not the left, is seen to reinforce behavior and activate midbrain dopaminergic reward nuclei [18], and gut-innervating vagal afferents likely play a key role in this lateralized reward signaling [16]. These recent studies raise additional questions about the cellular origins of asymmetric reward encoding by the right versus left vagus nerve.

Genetic sequencing studies of the NG in mice have revealed incredible diversity among NG cell types [6,20–23]. Transcription patterns of vagal neurons have been used to identify distinct subclasses of NG cells and their hypothesized sensory functions [20,21,24]. This work has typically been performed on pooled samples from left and right NG, however, and cannot address the question of which NG genes or neuronal subtypes exhibit asymmetric left-right expression that might contribute to lateralized vagal signaling. Recently, a more targeted investigation of left vs. right NG expression of several genes associated with digestive signaling showed that, in mice, neurons expressing *Cckar* (cholecystokinin 1 receptor), *Glp1r* (glucagon-like peptide 1 receptor), and *Npy2r* (neuropeptide Y receptor Y2; peptide YY receptor 2) are more numerous in right NG compared to left [12]. Differential left-right signaling through these receptors was not observed, however, leaving open the possibility of additional functionally relevant lateralized signaling pathways. In the current study, we use single-nucleus RNA sequencing (snRNA-seq) to identify genes of interest that are differentially expressed between the left and right NG that could contribute to right-biased vagus-mediated reward encoding. Because of their larger size compared to mice, rats are a more commonly used laboratory model to investigate translational vagal therapies such as vagus nerve stimulation (VNS). We thus performed comprehensive genetic profiling of left vs. right NG neurons in rats and compare our results to a publicly available mouse dataset [25]. Computational results were functionally validated using RNA fluorescence *in situ* hybridization (RNA-FISH) and calcium imaging.

Our results identify a single cluster of neurons, marked by co-expression of *Cckar* and *Chrna3*, that is preferentially found in the right NG of both rats and mice. Our data further suggest that this right-biased neuronal subtype exhibits a genetic profile consistent with a gut-innervating population, including co-expression of *Glp1r* (glucagon like peptide 1 receptor) and *Sctr* (secretin receptor). We additionally find that population responses to cholecystokinin 1 receptor (CCK1R)

activation likewise exhibit significant left-right asymmetry, with more neurons responding to
CCK1R agonism in the right NG compared to left. These findings suggest that differential right >
left expression of chemosensory vagal neurons expressing CCK1Rs could give rise to asymmetric
transduction of visceral sensory signals by the vagus nerve. Combined, our study raises the
possibility that right-biased digestive peptide signaling within this cell population, including at
CCK1Rs, may contribute to the lateralized encoding of interoceptive rewards.

RESULTS

Diverse mechanosensitive and chemosensitive cell types identified in rat nodose ganglia
We performed snRNA-seq on 12 replicate groups of NG cell suspensions. Each replicate group
was formed by pooling the right or left NG from six male or female rats (**Figure 1A**). In all,
36,741 nuclei were sequenced. Data were filtered according to standard quality control (QC)
metrics, and data from all replicate groups were integrated prior to clustering (**Figure S1**) [26,27].
Non-neuronal clusters (expressing *Ttn*, *Myh4*, *Mpz*, *Mbp*, *Apoe*, *Sparc*, *Emcn*, and/or *Tyrp1*)
and jugular neuronal clusters (expressing *Pdrm12*) were identified and removed during an initial
4 rounds of cluster filtering (**Figure S2**). After QC and cluster filtering, the remaining 23,428
nuclei were reclustered, resulting in 19 clusters corresponding to putative NG neuron subtypes,
which were included in subsequent analyses (**Figure 1B-C**).

Prior sequencing of mouse NG has identified *Piezo2* expressing (*Piezo2*+) putative
mechanosensory cells and *Scn10a* expressing "nociceptor-like" cells in the NG [21]. Our data, in
rats, replicate these fundamental results. We observed minimal co-expression of two sodium
channel genes, *Scn1a* (encoding $Na_v1.1$) and *Scn10a* (encoding $Na_v1.8$) (**Figure 1D**).
Expression of *Scn1a* typically coincided with expression of *Piezo2* and *Cntn1* (**Figure 1D**, left),
suggesting a myelinated and mechanosensory phenotype for the majority of these clusters. A
smaller group of two *Piezo2*+ putative mechanosensory clusters was also identified in our
dataset, which preferentially expressed *Scn10a* over *Scn1a*, and co-expressed *Tmem233*
(transmembrane protein 233). Most *Scn10a*-expressing clusters were not found to exhibit genes
associated with mechanosensation, however. Rather, they co-expressed *Oprm1* (mu-1 opioid
receptor) and many also co-expressed *Trpc4* (transient receptor potential cation channel
subfamily C member 4), pointing to a chemosensory phenotype for the majority of *Scn10a*-
expressing clusters (**Figure 1D**, right). Approximately 20% of the cells in our dataset (n =
5,006/23,428; 21.4%) were sorted into one of 8 *Piezo2*+, putative mechanosensory clusters,
whereas the majority of NG cells (n = 18,422/23,428; 78.6%) were instead sorted into one of the
11 *Piezo2*-negative putative chemosensory clusters. Our computational results suggest that in
rats, as has been shown previously in mice, there is little overlap of expression of *Scn1a* and
Scn10a in NG neurons, and that expression of the *Scn1a* vs. *Scn10a* is highly (though not
exclusively) correlated with putative mechanosensory vs. chemosensory functions, respectively.

Nodose ganglia clusters express both unique and overlapping sensory receptor
transcripts
We next examined the cluster-defining transcripts for each of our 19 identified NG clusters
(**Figure 2A**). Among these genes were a large number that encode membrane proteins
previously associated with sensory functions. These include several ion channels including

Grin2a (NMDA receptor subunit A; defining Cluster 12), *Chrna3* (nicotinic acetylcholine receptor
subunit 3; defining Cluster 4), *Trpa1* (transient receptor potential cation channel subfamily A
member 1; defining Cluster 3), and *Htr3b* (5-hydroxytryptomine receptor 3B; defining Cluster 5)
[28–32]. Other functional transmembrane proteins thought to be involved in visceral sensory
signaling were also prominent among cluster defining genes, including *Tmem233* (defining
Cluster 9)*, Cckar* (defining Cluster 4), and *Gpr65 (*G protein coupled receptor 65; Cluster 5) [33–
35]. Across all clusters, expression of the top 5 cluster-defining genes was highly correlated
between right and left NG (R^2 = 0.956, p < 0.000, Pearson's correlation), consistent with
excellent integration of data from right and left NG samples prior to clustering (**Figure 2B**).

We further examined the expression of several genes previously identified in the literature to be
strongly expressed in NG neurons (**Figure 2C**). Based on literature suggesting that lateralized
vagal reward signaling arises from gut-innervating NG neurons [16], we focused on genes known
to be associated with digestive signaling in the GI tract [36–38]. However, it is worth noting that
many of these receptors, including *Oprm1*, *P2rx3* (P2X purinergic receptor 3), and *Trpv1*
(transient receptor potential cation channel subfamily V member 1), for example, are not
exclusive to gut-innervating NG neurons, and have been shown to impact vagal signaling in
multiple organ systems [32,39–41]. Several of these known NG-expressed genes, including *Gpr65*,
Trpa1, and *Cckar* are also included among the cluster-defining genes shown in **Figure 2A**,
supporting the validity of our computational clustering approach. In this targeted subset of
genes, we find that the majority are expressed to some degree in both putative
mechanosensing and chemosensing clusters, though overall expression patterns support the
hierarchical clustering of our subtypes along these functional lines (**Figure 2C**). Consistent with
published literature, we find that expression of *Oprm1*, *P2rx3*, and *Cnr1* (endocannabinoid
receptor 1), in particular, are near-ubiquitously expressed across almost all NG clusters [42–44]

Few genes exhibit significant left-right differential expression within NG clusters
To begin to examine the possible molecular basis of lateralized vagal reward signaling, we
performed a pseudobulk analysis of differential gene expression within each of our 19 clusters [45].
This analysis identified five clusters in which any genes were found to be significantly differentially
expressed between right and left NG (Clusters 7, 3, 1, 2, and 5; **Figure 3A**) In the UMAP
visualization and hierarchical clustering tree, these *Scn10a+* chemosensing clusters were
situated near each other (**Figure 1B & 1D**), suggesting a degree of similarity in their genetic
profiles. Across all five clusters, 15 unique genes were identified as significantly differentially
expressed with at least a 1.5-fold difference in expression between left and right sides and a
Bonferroni-adjusted p < 0.01 (**Figure 3A**). Expression of two of these genes, *Gria1* (AMPA
receptor subunit 1) and *Shisa9* (Shisa family member 9), was found to be significantly left-NG
biased in more than one cluster. Overall, however, the effect sizes associated with identified
differentially expressed genes were small, rarely exceeding a 2-fold difference in expression
between sides. Our pseudobulk analysis suggests that, in naïve rats, cell-type specific expression
of differential sensory receptors may not be a major driver of lateralized vagal reward signaling.

A *Chrna3* and *Cckar* co-expressing cluster is preferentially found in right nodose ganglia
To test whether the identified cell types themselves might be differentially expressed between
right and left NG, we analyzed cell counts from each cluster to determine if that cluster
contained a greater-than-expected number of cells from the left or right side. Across the entire

dataset, we observed that the percentage of cells from right NG was slightly higher than the
percentage of cells from left NG (56% vs. 44%, respectively), which we attribute to a slight bias
in sample collection, as right NG were collected first during dissection in 20 of 36 rats. For each
individual cluster, we compared the percentage of cells from the right or left NG to this expected
distribution using multiple Bonferroni-corrected binomial tests. This statistical analysis revealed
four clusters with significantly lateralized expression (**Figure 3B**), only one of which was a right-
biased cluster. Cluster 4 comprised 7.2% of the overall NG population (n=1,694/23,428 nuclei)
and made up a significantly larger percentage of right NG cells compared to left (**Figure 3C**;
Left: 8.8 ± 0.30%, Right 5.2 ± 0.27%, p = 0.0006, paired t-test). This cluster was defined by high
levels of expression of *Cckar* and *Sctr*—genes known to play key roles in nutrient sensing and
digestive signaling [46–48]—as well as *Trpc6* (transient receptor potential cation channel subfamily
C member 6), *Chrna3, and Xylt1* (xylosyltransferase 1) (**Figure 3D**). Left-biased expression of
Clusters 8, 3, and 5 was also observed. These clusters were defined by the expression of
several genes involved in inflammation (e.g., *Cysltr2* and *S1pr3*, Clusters 10 and 8,
respectively) and noxious stimuli (e.g., *Trpa1* and *Htr3b*, Clusters 8 & 9, respectively) known to
be expressed in vagal fibers innervating the respiratory and gastrointestinal systems [21,37,49–51].

CCK1Rs are strongly expressed in gut-innervating NG neurons and are known to regulate
satiety and macronutrient sensing [52–55]. CCK has also been shown to play a role in the vagal
signaling pathways that shape nutrient preferences [25,56,57]. Right-biased expression of a NG
population co-expressing *Cckar*, *Glp1*, and *Npy2r* was recently shown in mice (Lansbury et al.,
2025), and lesions to *Cckar*+ neurons in the right, but not left NG, block the satiety-inducing
effects of systemically administered CCK [16]. Right-biased expression of a *Cckar*-expressing
cluster of NG neurons thus provides a strong candidate for lateralized transduction of reward-
related interoceptive signaling in rodents. To validate right-biased expression of our *Cckar*-
expressing Cluster 4, we used RNA-FISH to quantify *Cckar*, *Chrna3*, and *Xylt1* expression in left
versus right NG from six rats (**Figure 4A-C**). Our results confirmed overlapping expression of
these 3 genes in both left and right NG cells (**Figure 4D-E**). *Cckar*+ neurons were found to be
significantly more numerous in right NG compared to left (**Figure 4F**; *Cckar*+, cells/mm^2: Left:
96.04 ± 7.64, Right: 138.94 ± 12.12; p = 0.0245, paired t-test). We also found that while most
Chrna3+ neurons co-expressed *Cckar* and/or *Xylt1* (n = 2,513/4,309; 58.3%), 42.5% expressed
Chrna3 alone, without these other Cluster 4 marker genes (**Figure 4E**). Differential expression
of *Chrna3*+ neurons exhibited a non-significant trend toward right-side biased expression
(**Figure 4G**; Left: 121.62 ± 7.05, Right: 166.98 ± 12.34; p = 0.0605, paired t-test). Thus, while
the majority of *Cckar*-expressing cells co-express *Chrna3* (n = 2,468/3,483; 70.9%), and this
population is seen to be more numerous in right NG compared to left, we also find a population
of *Chrna3*+ cells that do not express *Cckar* and do not exhibit strong differential left-right
expression. *Xylt1*+ neurons were less numerous overall than predicted by the computational
data, though this may in part be attributable to quantification difficulties resulting from high
background staining in this probe channel. Nonetheless, the majority of *Xylt1*+ neurons (n =
974/1,490; 65.4%) were found to co-express *Cckar* and *Chrna3* (**Figure 4J**), and *Xylt1*+
neurons were significantly more numerous in the right NG (**Figure 4H**; *Xylt1*+, cells/mm^2: Left:
38.72 ± 4.08, Right: 56.67 ± 6.87; p = 0.0265, paired t-test), as were triple-labeled cells (**Figure
4I**; *Cckar*+|*Chrna3*+|*Xylt1*+ co-labeled, cells/mm^2: Left: 22.66 ± 1.02, Right: 40.99 ± 5.15; p =

0.0086, paired t-test). Overall, our RNA-FISH results confirm the presence of a
Cckar+|Chrna3+|Xylt1+ neuronal subtype that is preferentially expressed in the right NG.

To estimate the percentage of NG neurons comprised by our Cluster 4 subtype, we performed
additional RNA-FISH experiments to probe expression of the neuronal marker *Snap25*, along with
Cckar and *Chrna3* in left vs. right NG slices (**Figure 4J-R**). As in the previous experiment, we
found significant right-biased expression of *Cckar+* neurons (**Figure 4O**; Percent *Cckar+*: Left:
11.86 ± 0.91%, Right: 19.60 ± 1.83%; p = 0.0162, paired t-test) but differential expression of
Chrna3+ did not quite reach statistical significance (**Figure 4P**; Left: 19.97 ± 2.28% , Right: 24.77
± 2.51%; p = 0.0661, paired t-test). The average density of cells expressing *Snap25* did not differ
between right and left NG (**Figure 4Q**; *Snap25+*, cells per mm^2: Left: 759.20 ± 78.21, Right:
804.84 ± 70.15; p = 0.4538, paired t-test). Consistent with our computational results, *Cckar* and
Chrna3 co-labeled neurons, putatively corresponding to our Cluster 4, were found to be ca. 1.7
times more numerous in the right NG compared to the left (**Figure 4R**; Percent *Cckar+|Chrna3+*
co-labeled: Left: 9.00 ± 0.90%, Right: 14.78 ± 1.76%, p= 0.0204, paired t-test).

Cckar expression partially overlaps with expression of Glp1r and Cartpt
Right-biased co-expression of several gut peptide receptor genes, including *Cckar*, *Glp1r*, and
Npy2r, was recently demonstrated in NG of mice (Lansbury et al., 2025), delineating
lateralization of a neuronal subpopulation previously identified in transcriptomics data in that
species [21]. Our transcriptomics data from rats, however, suggests that while several genes
associated with gastrointestinal peptidergic signaling, including *Glp1r*, *Cartpt*, and *Lepr*, for
example, are likely present in our *Cckar+|Chrna3+* Cluster 4, expression of these genes may be
more prominent in *Cckar-* clusters. We thus performed additional RNA-FISH experiments to
quantify whether a subset of these genes, namely *Glp1r* and *Cartpt*, also exhibit differential
expression in left vs. right NG in rats (**Figure 5A-D**), and the extent to which these genes are
co-expressed in *Cckar+* NG cells (**Figure 5E**). Our results again confirm strong right-biased
Cckar expression (**Figure 5F**; *Cckar+*, cells/mm^2: Left: 75.35 ± 13.11, Right: 131.68 ± 16.40; p =
0.0202, paired t-test). While the majority of *Cckar+* cells were found to co-express *Cartpt* (n =
2,213/3,250, 68.1%), the reverse was not true—most *Cartpt+* cells did not co-express *Cckar* or
Glp1r (**Figure 5E**; n=6,866/9,387, 73.1%), and differential expression of *Cartpt+* cells did not
reach statistical significance (**Figure 5G**; Left: 255.04 ± 24.39, Right: 316.09 ± 15.23; p =
0.0598, paired t-test). Consistent with our computational results, and similar to what has
recently been reported for mice [12], just over half of *Cckar+* cells were found to co-express *Glp1r*
(n = 1,720/3,250, 52.9%). *Glp1r+* cells were found to be significantly more numerous in right NG
compared to left (**Figure 5H**; Left: 92.69 ± 5.64, Right: 134.12 ± 10.38; p = 0.0321, paired t-
test), as were triple-labeled cells (**Figure 5I**; Left: 31.05 ± 5.64, Right: 51.44 ± 8.93; p = 0.0296,
paired t-test). These results indicate that *Cartpt* expression labels a much broader population of
NG neurons than *Cckar* and *Chrna3* co-expression. By contrast, there is significant overlap and
similar right-biased expression in *Glp1r+* and *Cckar+|Chrna3+* NG neuron populations in rats.

Cckar and Chrna3 co-expressing NG neurons exhibit right-biased expression in mice
By integrating our rat data with a publicly available dataset from mice [58], we examined whether
gene expression patterns and right-side bias of a *Cckar+* cluster could be similarly observed in
both species. Mouse and rat datasets were first filtered using quality control metrics (**Figure
S3A-B)**. After QC filtering, the mouse and rat datasets were integrated and clustered, applying

the same computational pipeline as before. Clusters representing jugular neurons (*Pdrm12*+)
and non-neuronal cell types (expressing *Ttn*, *Myh4*, *Mpz*, *Mbp*, *Hand2*, *Daam2*, and/or *Mecom*)
were removed through 3 rounds of initial clustering (**Figure S4**). After removal of non-neuronal
and jugular neuron clusters and subsequent reclustering, 16 putative NG clusters were
identified in the integrated dataset (**Figure 6A**), with good overlap between the mouse and rat
data (**Figures 6B-C**). In the integrated dataset, Cluster 7 was observed to be defined by
expression of *Trpc6* and *Chrna3*, and to express high levels of *Cckar*, similar to Cluster 4 in our
rat-only data (**Figure 6D**). In both mice and rats, Cluster 7 made up ca. 5-6% of the total sample
(mouse: n=181/3,664 nuclei, 4.9%; rat: n=1,766/29,117 nuclei, 6.1%). Statistical analysis
confirmed that neurons in this cluster were significantly more numerous in right NG samples
than in left in both mice (n = 169/181, 93%, compared to 62.5% right NG neurons total in mouse
samples; p < 0.0000, binomial test), and rats (n = 1,228/1,766; 69.5%, compared to 59.7% right
NG neurons total in rat samples; p < 0.0000, binomial test). While *Chrna3* and *Cckar* co-
expression defined integrated Cluster 7 and exhibited strong expression in both the mouse and
rat datasets (**Figure 6E; Figure S3C**), *Xylt1* appeared weakly expressed within this cluster in
mouse samples. Several other well-known NG genes associated with gut-brain signaling,
including *Cartpt*, *Npy2r*, and *Vip*, were observed to be much more strongly expressed in Cluster
7 in mice compared to rats (**Figure 6E**), suggesting significant species specificity in gene
expression patterns within this cluster.

To validate right-biased expression of our integrated Cluster 7 in mice, we performed RNA-FISH
to probe for *Cckar*, *Chrna3,* and *Snap25* expression in mouse NG slices (**Figure 6F-N**). As in
rats, most *Cckar*+ neurons in mice also co-expressed *Chrna3* (**Figure 6J**; n = 763/1,058,
72.1%), and *Cckar*+ neurons were found to be significantly more numerous in right NG
compared to left (**Figure 6K**; Left: 10.18 ± 1.30%, Right: 19.20 ± 1.19%; p = 0.0120, paired t-
test). Only ca. 40% of *Chrna3* neurons were found to co-express *Cckar* (n = 763/1,821, 41.9%).
The overall *Chrna3*+ population was found to be significantly larger in right NG (**Figure 6L**; Left:
17.35 ± 3.77%, Right: 32.14 ± 1.74%; p = 0.0214, paired t-test) as was the *Cckar*+|*Chrna3*+ co-
labeled population (**Figure 6N**; Left: 5.62 ± 1.30%, Right: 14.91 ± 0.81%; p = 0.0024, paired t-
test). We found no difference in *Snap25*+ neuron density between right and left NG (**Figure 6M**;
Snap25+, cells per mm^2: Left: 1,386.22 ± 117.63, Right: 1,548.02 ± 191.49; p = 0.4030, paired t-
test).

Combined, our results demonstrate that a *Cckar* and *Chrna3* co-expressing subpopulation of
neurons is preferentially expressed in the right NG over the left in both mice and rats.

CCK1R agonism preferentially activates right NG neurons
To validate functional lateralization of CCK1R signaling in the NG, we performed calcium
imaging on left and right NG cultures from 10 rats (6 male, 4 female). Responses to bath
application of the selective CCK1R agonist, A71623 (100 nM), were recorded for 1,233 total
neurons (**Figure 7A-B**). Consistent with our transcriptomic and RNA-FISH results, we observed
significant asymmetry in population responsiveness to CCK1R agonism among left vs. right NG
neurons. Approximately twice as many A71623-responsive neurons were found in right NG
cultures compared to left (**Figure 7C**; Right: n=69/495, 13.94%; Left: n = 50/738, 6.78%; p <
0.0001, Chi-square test). Among responders, there was no difference between left and right NG
neurons in the magnitude of the peak response to CCK1R agonism (**Figure 7D**; Left: 49.41 ±

3.68%, Right: 45.42 ± 2.03%, p = 0.3117, unpaired t-test). These findings are consistent with
our computational data and suggest that protein expression levels are similar for CCK1R-
expressing left and right NG neurons. However, the CCK1R-expressing population itself is more
numerous in the right NG compared to the left.

Taken together, our transcriptomic, RNA-FISH, and calcium imaging results demonstrate
significant right > left asymmetry in the expression of a *Cckar*+ vagal neuron subtype, and in
functional responsiveness to CCK1R activation in the NG.

DISCUSSION

Our data represent the first snRNA-seq analysis of NG neurons in rats, providing an important
resource to the neuroscientific community. In the current study, we aimed to utilize this dataset
to determine whether genetic differences in right vs. left NG neurons could contribute to right-
biased encoding of visceral reward-related signaling by the vagus nerve. Our analyses identified
a single neuronal cluster, defined by *Cckar* and *Chrna3* co-expression, that exhibits preferential
expression in the right NG of both rats and mice. Our transcriptomic and RNA-FISH results
further indicate that, in rats, neurons in this population are further defined by *Sctr* co-expression,
and often co-express *Cartpt* and *Glp1r*, pointing to a role for this neuronal subtype in digestive
signaling. Finally, our calcium imaging experiments provide novel evidence of functional right-
biased asymmetry in CCK1R-mediated NG signaling. Combined, the current study identifies a
unique population of right-biased *Cckar* and *Chrna3* co-expressing NG neurons as a strong
candidate for contributing to the lateralized vagal encoding of interoceptive rewards.

Cholecystokinin (CCK) is a peptide released by enteroendocrine cells in the gastrointestinal
tract in response to food intake, and CCK1Rs are strongly expressed in gut-innervating NG
neurons [52–54]. CCK1Rs play an important role in regulating satiety and sensing macronutrients
through gut-brain vagal signaling pathways. CCK has also been found to play a critical role in
shaping the preference for macronutrients including fat, sugar, and amino acids [25,56,57].
Glucagon-like peptide-1 (GLP1) and secretin are likewise known to contribute to vagal-
dependent nutrient detection, satiety signaling, and the modulation of reward-related signaling
pathways [59–62]. In light of this prior work, our data support a model of sensory transduction of
reward-related interoceptive cues by a right-side biased population of neurons engaged in
detection of CCK and other digestive peptides.

Previous studies have shown that stimulation of the right, but not left, vagus nerve increases
dopamine release and reinforces behavior, consistent with lateralization of interoceptive reward
processing [16,18]. Gut-innervating vagal neurons, in particular, likely give rise to this left-right
asymmetry in vagal-mediated reward signaling [16]. Using an unbiased transcriptomic approach,
we identified a single cluster of NG neurons, distinguished by co-expression of *Cckar* and
Chrna3 in both rats and mice, that is significantly more abundant in the right NG compared to
the left. Our transcriptomic data indicate that this neuronal subtype is also distinguished by high
levels of expression *Trpc6* and *Sctr*. This gene expression profile is highly similar to that of a
putative gut-innervating IGLE population previously identified by Kupari et al. [21] in mice. Our

study provides novel evidence that this cluster is conserved across mice and rats and that it exhibits strongly right-biased expression that is unique among NG subtypes.

Expression of this *Cckar* and *Chrna3* co-expressing subpopulation is ca. 1.5 to 2 times greater in the right NG compared to the left, comprising up to ca. 20% of right NG neurons in our data, and ca. 5-10% of left NG neurons. It is notable, then, that VNS has been observed to exhibit rewarding properties only when applied to the right side. Several factors may contribute to these findings. At parameters typically used for neurological indications, VNS is thought to preferentially activate myelinated cervical vagal fibers [63], and thus may only weakly activate *Cckar*+ IGLEs [37]. Clinically and preclinically, VNS is predominantly applied to the left vagus nerve, which our results indicate is likely to harbor fewer of these *Cckar*+ gut-innervating, putative reward-transducing neurons. As a result, left-sided stimulation may fail to sufficiently engage this population, and/or the effects of activation may be masked by co-activation of mechanosensitive or nociceptive-like subtypes. On the right side, the impact of stimulation of the larger population of vagal reward-sensitive neurons may be further amplified by asymmetrical connectivity within the central nervous system, as has been described by Han et al. [16], producing dopaminergic signaling that more readily drives behavioral reinforcement. The current study presents a novel characterization of prominent neural asymmetry in left vs. right vagal signaling and highlights the importance of considering this molecular and functional lateralization in future mechanistic studies of interoceptive reward processing. By demonstrating significant differential left-right expression of neuronal subtypes in the NG, our work additionally provides fundamental insights to inform the development and optimization of VNS or pharmacological strategies designed to modulate vagal signaling. These insights may be particularly relevant for therapeutic strategies aimed at treating conditions involving dopamine dysregulation, including obesity, depression, and addiction.

Limitations of the study

In this study, we investigated whether genetic differences in right vs. left NG neurons could contribute to right-biased encoding of visceral reward-related signaling by the vagus nerve. We identified a single cluster, marked by the expression of *Cckar* and *Chrna3*, that exhibits a strong right-biased expression in the NG. Combined with prior literature, our results suggest that lateralized reward-related interoceptive signaling may be supported by this unique neuronal subtype. Our snRNA-seq data indicate that this *Cckar*+ and *Chrna3*+ population co-expresses genes encoding several digestive peptide receptors, consistent with a gut-innervating subtype. However, the extent to which the gut receives preferential innervation from *Cckar*+ neurons in the right NG rather than the left, remains to be anatomically confirmed. Additional research is also needed to causally test whether activation of this right-biased NG subpopulation is indeed sufficient to drive dopaminergic midbrain activity and reinforce behavior, as well as to determine the specific cellular signaling pathways that participate in the lateralized transduction of visceral rewards by the vagus nerve.

ACKNOWLEDGMENTS

This work was funded by NIH R21 DA055166 (C.A.T.), R01 NS065926 (T.J.P.), R01 DK115478 (B.J.K.), and F31 NS129269 (V.M.H.). We thank Dr. Adriana Shembel, Dr. Robert Morrison, and Dr. I-Fan Mau for their technical assistance in establishing NG dissection protocols. We thank the UTD Imaging and Histology Core for their expertise in imaging methods. We thank Dr. Ashley Plumb and Dr. Katelyn Sadler for their assistance with calcium imaging experiments.

AUTHOR CONTRIBUTIONS

H.F.W. participated in conceptualization, investigation, formal analysis, visualization, data curation, and writing – original draft; V.M.H., I.S., H.Q., and K.M. participated in investigation and writing – review & editing; B.J.K. and T.J.P. participated in supervision, writing – review & editing, and funding acquisition; C.A.T. participated in conceptualization, formal analysis, data curation, writing – original draft, project administration, and funding acquisition.

DECLARATION OF INTERESTS

The authors declare no competing interests.

METHODS

EXPERIMENTAL MODEL DETAILS

A total of 52 rats (27 male, 25 female) and 6 mice (3 male, 3 female) were used in the study. All procedures were conducted in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and were approved by the University of Texas Institutional Animal Care and Use Committee (protocol number 2024-0126).

METHOD DETAILS

Experimental design: The right and left NG of experimentally naïve, 8-week-old male (n=18) and female (n=18) rats were dissected, and single-nucleus RNA sequencing (snRNA-Seq) was performed. snRNA-seq was performed on NG samples from 12 replicate groups (3 Male|Left, 3 Male|Right, 3 Female|Left, 3 Female|Right); each replicate group was created by pooling left or right NG samples from six rats (**Figure 1A**). Samples were prepared using a Parse Biosciences Evercode Nuclei Fixation Kit v2 (ECF2003) and Parse Single Cell Whole Transcriptome Kit v2 (ECW02130) and sequenced on an Illumina Nextseq 2000 at the Genome Center at the University of Texas at Dallas. Clustering and statistical analyses were performed in R version 4.4.1 [65] using Seurat version 4.3.0 (Hao et al., 2021). Computational results were validated using RNA fluorescence *in situ* hybridization (RNA-FISH) to quantify overlap in expression of key transcripts as well as side-specific expression in NG samples collected from an additional 6 rats (3 male, 3 female) and 6 mice (3 male, 3 female). Calcium imaging was performed on cultured NG cells from an additional 10 rats (6 male, 4 female) to validate functional asymmetry in NG CCK1R signaling.

snRNA-Seq Sample Preparation and Sequencing: Rats were deeply anesthetized with 5% isoflurane and sacrificed via rapid decapitation. The right and left NG were dissected and stored in separate PCR tubes before flash-freezing on dry ice. NG samples were stored in a -80 ºC freezer until sample preparation.

Individual NG were pooled to prepare three replicate groups for each of the four treatment groups (Male|Left; Male|Right; Female|Left; Female|Right). Each replicate group was created by pooling the right or left NG samples from 6 rats (**Figure 1A**). Nuclei were fixed and barcoded according to Parse Biosciences protocols (Evercode Version 2 Nuclei Fixation Kit, Version 2 WT Kit). The samples were split into 8 sublibraries and were sequenced via paired-end sequencing using the Illumina Nextseq 2000 at the Genome Center at the University of Texas at Dallas. In all, an estimated 90,000-100,000 nuclei were prepared for sequencing.

snRNA-Seq Analysis: The 12 replicate groups generated a total of 36,741 sequenced cells with an average of ca. 32,000 reads/cell and ca. 1,500 genes detected per cell. Cells were filtered using standard quality control metrics (less than 6,500 genes per cell, less than 20,000 molecules per cell, and less than 5% mitochondrial genes), removing 281 poor-quality cells. Data from each treatment group was integrated, scaled, and clustered using default parameters in Seurat. Additional filtering was performed to remove clusters expressing high levels of marker genes for muscle (*Ttn, Myh4*), myelin (*Mbp, Mpz*), endothelial cells (*Emcn*), melanocytes (*Tryp1*), satellite glia (*Apoe, Sparc*), and jugular neurons (*Prdm12*). This iterative process of removing clusters, reclustering, and re-examining non-neuronal cell and jugular neuron marker gene expression in the rat dataset was performed over 4 initial rounds of cluster filtering (**Figure S2**). The resulting dataset of putative NG-specific neurons was comprised of 23,428 nuclei and was reclustered once more prior to analysis.

Cluster-specific marker genes were identified using Wilcoxon rank-sum tests with Bonferroni corrections. To test for differential gene expression between right and left samples, pseudobulk analysis was used [45]. For each cluster, data from all replicate groups was aggregated and

differential right vs. left gene expression was calculated using Bonferroni-corrected Wilcoxon
rank-sum tests.

To compare the lateralized expression of NG clusters in rats and mice, we integrated our rat
snRNA-seq data with a publicly available mouse NG sequencing dataset of 5,507 cells in which
left and right samples were separately identified (GEO #GSE185173) [25]. The rat and mouse
datasets were individually filtered using standard quality control metric thresholds (less than 6,500
genes per cell, less than 20,000 molecules per cell, and less than 5% mitochondrial genes).
Mouse and rat data were then integrated, scaled, and clustered using default parameters in
Seurat. Clusters in the integrated dataset were further filtered to restrict analysis to putative NG
neurons by removing clusters expressing high levels of marker genes for muscle (*Ttn*, *Myh4*),
myelin (*Mbp*, *Mpz*), sympathetic neurons (*Hand2*), oligodendrocytes (*Daam2*), endothelial cells
(*Mecom*), and jugular neurons (*Prdm12*). There were high levels of *Apoe* across multiple clusters
in the mouse data, which could be a general sign of cellular damage or stress [69,70], rather than a
reliable marker of satellite glial cells. Thus, we did not remove *Apoe+* clusters in the integrated
dataset as we did for the rat-only data. This iterative process of removing clusters, reclustering,
and re-examining non-neuronal cell and jugular neuron marker gene expression in the integrated
dataset was performed over 3 initial rounds of cluster filtering (**Figure S4**). Cluster filtering
resulted in an integrated dataset comprised of 32,781 nuclei (mouse: n = 3,664; rat: n = 29,117).

In situ hybridization: Rats and mice were deeply anesthetized with sodium pentobarbital and
transcardially perfused with cold phosphate buffered saline (PBS) followed by 4%
paraformaldehyde (PFA) in PBS. NG were dissected, embedded in OCT, quickly frozen over dry
ice and stored at -80 ºC until sectioning. Slices were made at 20 μm thickness using a Leica
CN1860 Cryostat and directly mounted on slides. Slides were stored at -80 ºC until RNA-FISH
was performed.

Hybridization chain reaction RNA fluorescence *in situ* hybridization (HCR RNA-FISH v3.0;
Molecular Instruments) was used to validate differential right vs. left NG expression of the
computationally identified right-biased *Chrna3+* and *Cckar+* cluster. Hybridization probes sets
targeting rat *Chrna3*, *Cckar*, and *Xylt1* were custom designed based on the transcript sequences
found in NCBI (NCBI Accession numbers: *Cckar* (rat)*:* NM_012688, *Chrna3* (rat)*:* NM_052805,
Xylt1 (rat): NM_022295). To validate neuronal expression of the *Cckar* and *Chrna3* co-expressing
populations in rats and mice, probe sets targeting *Cckar* and *Chrna3*, as well as the neuronal
marker *Snap25*, were used (NCBI Accession numbers: *Snap25* (rat)*:* NM_030991; *Cckar*
(mouse): NM_009827, *Chrna3* (mouse): NM_145129, *Snap25* (mouse): NM_011428). To
quantify overlapping and right-biased expression of select genes associated with digestive
peptide signaling in rats, probes targeting *Cartpt*, *Glp1r*, and *Cckar* were used (NCBI Accession
numbers: *Cartpt* (rat): NM_017110, *Glp1r* (rat): NM_012728). Multiplexed staining was performed
using an HCR RNA-FISH kit including three-probe sets, probe channel-specific HCR amplifiers
(each consisting of a pair of fluorophore-tethered hairpins), as well as hybridization, wash, and
amplification buffers.

HCR[TM] RNA-FISH was performed according to adapted vendor protocols [64]. After removing
samples from -80°C, mouse slides were baked for 30 minutes at 60°C using HybEZ II oven
(#240200, ACD) to improve tissue adhesion (rat slides were not baked). The procedure for mouse
samples continued as written. Slides were immersed at room temperature in 100% ethanol for 15
minutes, followed by immersion in 100% ethanol for 5 minutes, then immersed once in 1X PBS
briefly to wash off remaining OCT. Slides were then air dried and a hydrophobic barrier was drawn
around the tissue using the ImmEdge® Hydrophobic Barrier PAP pen (H-4000, Vector
Laboratories). Probe hybridization buffer was carefully added on top of the samples and allowed
to pre-hybridize for 10 minutes in a humidified chamber using HybEZ II oven at 37 °C. A probe

solution was prepared by adding 2.5 μL of each hybridization probe set (1 μM stock) in 100 μL of
pre-heated probe hybridization buffer solution at 37 °C per slide. Pre-hybridization solution was
removed from slides, and probe solution was added on top of the samples and incubated
overnight (> 20 hrs) at 37 °C. As a negative control, one slide was treated with hybridization
solution with no gene probes so there would be no initiator to trigger growth of the fluorescent
amplification polymer.

The following day, slides were washed in a series of mixed and pre-heated probe wash buffer
and 5X Saline Sodium Citrate buffer (SSCT) (BP1325, Fisher) solutions at 37 °C for 15 min each
in the following order: (1) 75% probe wash buffer with 25% 5X SSCT, (2) 50% probe wash buffer
with 50% 5X SSCT, (3) 25% probe wash buffer with 75% 5X SSCT, and (4) 100% 5X SSCT. This
was followed by a 5 min wash in 100% 5X SSCT at room temperature. Amplification buffer was
added on top of the samples, and slides were pre-amplified in the humidified chamber for 30
minutes at room temperature. Hairpin dyes were snap-cooled by incubating at 95°C for 90
seconds, then immediately cooled at room temperature in the dark for at least 30 minutes. The
snap-cooled hairpins were added to the amplification buffer to create a hairpin solution. The pre-
amplification buffer was removed from the slides, and the hairpin solution was added on top of
the samples and incubated overnight in a humidified chamber at room temperature.

The next day, slides were washed in 100% 5X SSCT twice for 30 min and then once for 5 min.
An autofluorescence quenching solution was created by making a 1:1:1 solution of reagents in
the Vector® TrueVIEW® Autofluorescence Quenching Kit (SP-8400-15, Vector Laboratories). This
solution was added on top of the samples and incubated for 3.5 minutes, then slides were washed
in 1X PBS for 5 minutes at room temperature. Slides were cover slipped with VectaShield®
Vibrance® Antifade Mounting Medium with DAPI (H-1800-10, Vector Laboratories) and sealed
with nail polish. Slides were imaged within 24 hours.

FISH imaging and image analysis: An Olympus FV3000RS or Evident FV4000 confocal
microscope was used to capture images. The same acquisition parameters were used for all
experimental samples and negative controls. Two to six slices per ganglia were imaged at 20X
(rat) or 60X (mouse) for semi-automated cell counting. Multiple 16-bit images with overlapping
fields of view were taken to capture the entirety of each slice, and images were stitched together
to reconstruct whole-slice images in each channel [71]. For some images, the built-in multi-area
time lapse (MATL) feature was used to stitch whole images. Image analysis was performed
using FIJI [67]. Image quantification was performed by a blinded experimenter. Stitched images
were smoothed using a 3x3 mean filter, then thresholded using minimum pixel intensity cutoffs
based on probe-specific signal-to-noise ratios (*Cckar* (rat): Mean+2.5*STD, *Chrna3* (rat):
Mean+1.5*STD, *Xylt1* (rat): Mean+2.5*STD, *Snap25* (rat): Mean+1.5*STD, *Cartpt* (rat):
Mean+1*STD, *Glp1r* (rat): Mean+2*STD; *Cckar* (mouse): Mean+1*STD, *Chrna3* (mouse):
Mean+1.5*STD, *Snap25* (mouse): Mean+1*STD). Cells were then automatically detected using
particle analysis for particles larger than 50 pixels^2. This process generated image masks
comprised of detected cells, from which cell counts for each individual channel were obtained.
For each slice, probe co-expression was manually quantified by an experimenter blinded to
sample side and sex, using overlaid and dual- or tri-color channel masks. Single and co-labeled
cell counts for each slice were normalized by dividing by the area of the slice (mm^2), and these
normalized values were averaged across slices to obtain an average cell density for each
animal. For slices with *Snap25* label, the numbers of *Cckar*+, *Chrna3*+, and *Cckar*+|*Chrna3*+
co-labeled cells were divided by the number of *Snap25*+ neurons for each slice, and these
percentages were averaged across slices for each animal. To illustrate the overlap in probe
labeling for each probe set, semi-quantitative Venn diagrams were created using DeepVenn [68].
Circle area in the diagrams is proportional to the average number of labeled cells per slice in the

relevant probe channel. This average was calculated by first computing the mean number of
cells per slice for each subject, then averaging across all subjects.

Primary NG neuron isolation and culture: Rats were deeply anesthetized with 5% isoflurane and
sacrificed via rapid decapitation. The left and right NG were dissected and placed in separate 1.5
mL Eppendorf tubes containing 1 mL Hanks Balanced Salt Solution (HBSS). Samples were
immediately used for cell culture.

Dissected samples in HBSS were transported to a biosafety cabinet. HBSS was removed from
samples. NG were physically dissociated with scissors and incubated for 40 minutes in
Dulbecco's Modified Eagle Medium (DMEM) and 1 mg/mL collagenase type IV. Solution was
removed, and NG then incubated for 45 minutes in DMEM and 1 mg/mL trypsin. NG were
plated onto laminin-coated glass coverslips within a 12-well plate and incubated for 1 hour.
Neurons were fed with media containing DMEM with 10% heat-inactivated horse serum, 2 mM
L-glutamine, 1% glucose, and 100 units/mL pen-strep. Samples were incubated at 37 °C
overnight and imaged the next day.

Calcium imaging: Calcium imaging was performed on neurons using the ratiometric calcium
indicator dye Fura-2-AM. Neurons were washed with 2.5 µg/mL Fura-2 in 2% BSA for 45 minutes
at room temperature and then washed with extracellular buffer for 15 minutes. Extracellular buffer
(7.4 pH, 320 Osm) contained 150 mM NaCl, 10 mM HEPES, 8 mM glucose, 5.6 mM KCl, 2 mM
$CaCl_2$, and 1 mM $MgCl_2$. Coverslips were mounted onto an inverted fluorescence microscope and
perfused with room temperature extracellular buffer at a rate of 6 mL/min. Cells were perfused
with extracellular buffer for 1.5 minutes, A71623 for 2 minutes (100 nM in extracellular buffer),
extracellular buffer for 2 minutes, KCl for 1 minute (50 mM in extracellular buffer), and extracellular
buffer for 30 seconds. Fluorescence images were captured at 340 and 380 nm every 0.5 seconds.

Calcium imaging analysis: Cells were excluded from further analysis if there was a greater than
25% increase in the 340/380 nm ratio during the last 60 seconds of the initial extracellular buffer
perfusion. The 340/380 nm ratio traces were then smoothed using a 15-point moving average
filter prior to testing for drug responses. Cells that exhibited a greater than 25% increase in the
340/380 nm ratio during both A71623 and KCl treatment periods, compared to the 3.5 second
pre-treatment windows, were counted as responders. Cells were considered non-responders to
A71623 if they exhibited a greater than 25% increase in the 340/380 nm ratio during the KCl
treatment period only. Cells that failed to respond to KCl treatment were excluded from analysis.
For A71623-responsive neurons, a 3-point running average was computed for the ±1 sec period
around the maximum raw 340/380 nm ratio during the A71623 treatment period. The maximum
value of the 3-point smoothed data within this ±1 sec window was taken as the peak response.
A chi-squared test was used to test whether there was a difference in the frequency of A71623-
responsive neurons from left vs. right NG cultures.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data were preprocessed using Trailmaker™ (Parse Biosciences) to obtain cell-gene count
matrices for each replicate group. Sequencing data were then analyzed with R (version 4.4.1)
using Seurat (version 4.3.0). For the pseudobulk analysis, statistically significant left-right
differential gene expression is reported for genes with at least a 1.5-fold difference in gene
expression between the two sides, and a Bonferroni corrected $p < 0.01$. To test for lateralized cell
counts for each identified cluster in the rat dataset, Bonferroni-corrected binomial tests were
performed. For each cluster identified as having significant left or right-side biased counts
(corrected $p < 0.05$), additional binomial tests were used to validate that this bias was observed
in samples from both males and females. In the rat + mouse integrated dataset, we tested for

lateralized expression of our *Chrna3* and *Cckar* co-expressing cluster of interest using a binomial
test. Statistical significance is reported for $p < 0.05$.

For analysis of RNA-FISH data, paired t-tests were used to test for differential left-right expression
of labeled neurons. Statistical significance is reported for $p < 0.05$. MATLAB was used to plot data
and perform statistical tests.

In the calcium imaging experiments, a chi-squared test was used to test whether there was a
difference in the frequency of A71623-responsive neurons from left vs. right NG cultures
(GraphPad Prism). Peak amplitudes for left vs. right A71623-responsive neurons were compared
using an unpaired t-test (MATLAB). Statistical significance is reported for $p < 0.05$.

Unless otherwise noted, summary data are presented as mean $\pm$ SEM. Additional statistical
details, including precise p values and numerical n values, can be found in the text and figure
legends.

REFERENCES

1. Prescott, S.L., and Liberles, S.D. (2022). Internal senses of the vagus nerve. Preprint, https://doi.org/10.1016/j.neuron.2021.12.020 https://doi.org/10.1016/j.neuron.2021.12.020.

2. Berthoud, H.R., and Neuhuber, W.L. (2000). Functional and chemical anatomy of the afferent vagal system. In Autonomic Neuroscience: Basic and Clinical https://doi.org/10.1016/S1566-0702(00)00215-0.

3. Zhuo, H., Ichikawa, H., and Helke, C.J. (1997). Neurochemistry of the nodose ganglion. Prog Neurobiol *52*. https://doi.org/10.1016/S0301-0082(97)00003-8.

4. Capilupi, M.J., Kerath, S.M., and Becker, L.B. (2020). Vagus nerve stimulation and the cardiovascular system. Cold Spring Harb Perspect Med *10*. https://doi.org/10.1101/cshperspect.a034173.

5. Maniscalco, J.W., and Rinaman, L. (2018). Vagal interoceptive modulation of motivated behavior. Preprint, https://doi.org/10.1152/physiol.00036.2017 https://doi.org/10.1152/physiol.00036.2017.

6. Mazzone, S.B., and Undem, B.J. (2016). Vagal afferent innervation of the airways in health and disease. Physiol Rev *96*. https://doi.org/10.1152/physrev.00039.2015.

7. Schwartz, G.J. (2000). The role of gastrointestinal vagal afferents in the control of food intake: Current prospects. In Nutrition https://doi.org/10.1016/S0899-9007(00)00464-0.

8. Teckentrup, V., and Kroemer, N.B. (2024). Mechanisms for survival: vagal control of goal-directed behavior. Preprint, https://doi.org/10.1016/j.tics.2023.11.001 https://doi.org/10.1016/j.tics.2023.11.001.

9. Prechtl, J.C., and Powley, T.L. (1985). Organization and distribution of the rat subdiaphragmatic vagus and associated paraganglia. Journal of Comparative Neurology *235*. https://doi.org/10.1002/cne.902350204.

10. Zandstra, T.E., Notenboom, R.G.E., Wink, J., Kiès, P., Vliegen, H.W., Egorova, A.D., Schalij, M.J., De Ruiter, M.C., and Jongbloed, M.R.M. (2021). Asymmetry and Heterogeneity: Part and Parcel in Cardiac Autonomic Innervation and Function. Preprint, https://doi.org/10.3389/fphys.2021.665298 https://doi.org/10.3389/fphys.2021.665298.

11. Verzele, N.A., Chua, B.Y., Short, K.R., Moe, A.A.K., Edwards, I.N., Bielefeldt-Ohmann, H., Hulme, K.D., Noye, E.C., Tong, M.Z., Reading, P.C., et al. (2024). Evidence for vagal sensory neural involvement in influenza pathogenesis and disease. PLoS Pathog *20*, e1011635.

12. Landsbury, E.L., Vana, V., Lund, M.L., Ludwig, M.Q., Mamedova, E., Gautron, L., Arnold, M., Egerod, K.L., Kuhre, R.E., Holst, J.J., et al. (2025). Neurons Co-Expressing GLP-1, CCK, and PYY Receptors Particularly in Right Nodose Ganglion and Innervating Entire GI Tract in Mice. Int J Mol Sci *26*.

13. Wang, F. Bin, and Powley, T.L. (2000). Topographic inventories of vagal afferents in gasointestinal muscle. Journal of Comparative Neurology *421*. https://doi.org/10.1002/(sici)1096-9861(20000605)421:3<302::aid-cne2>3.0.co;2-n.

14. Jotz, G.P., De Campos, D., Rodrigues, M.F., and Xavier, L.L. (2011). Histological asymmetry of the human recurrent laryngeal nerve. Journal of Voice *25*. https://doi.org/10.1016/j.jvoice.2009.06.007.

15.   Ardell, J.L., and Randall, W.C. (1986). Selective vagal innervation of sinoatrial and atrioventricular
nodes in canine heart. Am J Physiol Heart Circ Physiol *251*.
https://doi.org/10.1152/ajpheart.1986.251.4.h764.

16.   Han, W., Tellez, L.A., Perkins, M.H., Perez, I.O., Qu, T., Ferreira, J., Ferreira, T.L., Quinn, D., Liu,
Z.W., Gao, X.B., et al. (2018). A Neural Circuit for Gut-Induced Reward. Cell *175*.
https://doi.org/10.1016/j.cell.2018.08.049.

17.   Roubeau, B., Lefaucheur, J.P., Moine, A., and St Guily, J.L. (1998). Asymmetry of the laryngeal
reflex responses to superior laryngeal nerve stimulation unrelated to the length of the recurrent
nerves in the porcine model. Acta Otolaryngol *118*. https://doi.org/10.1080/00016489850182620.

18.   Brougher, J., Aziz, U., Adari, N., Chaturvedi, M., Jules, A., Shah, I., Syed, S., and Thorn, C.A.
(2021). Self-Administration of Right Vagus Nerve Stimulation Activates Midbrain Dopaminergic
Nuclei. Front Neurosci *15*. https://doi.org/10.3389/fnins.2021.782786.

19.   Chhetri, D.K., Neubauer, J., and Sofer, E. (2014). Influence of asymmetric recurrent laryngeal
nerve stimulation on vibration, acoustics, and aerodynamics. Laryngoscope *124*.
https://doi.org/10.1002/lary.24774.

20.   Bai, L., Mesgarzadeh, S., Ramesh, K.S., Huey, E.L., Liu, Y., Gray, L.A., Aitken, T.J., Chen, Y.,
Beutler, L.R., Ahn, J.S., et al. (2019). Genetic Identification of Vagal Sensory Neurons That
Control Feeding. Cell *179*. https://doi.org/10.1016/j.cell.2019.10.031.

21.   Kupari, J., Häring, M., Agirre, E., Castelo-Branco, G., and Ernfors, P. (2019). An Atlas of Vagal
Sensory Neurons and Their Molecular Specialization. Cell Rep *27*.
https://doi.org/10.1016/j.celrep.2019.04.096.

22.   Prescott, S.L., Umans, B.D., Williams, E.K., Brust, R.D., and Liberles, S.D. (2020). An Airway
Protection Program Revealed by Sweeping Genetic Control of Vagal Afferents. Cell *181*.
https://doi.org/10.1016/j.cell.2020.03.004.

23.   Zhao, Q., Yu, C.D., Wang, R., Xu, Q.J., Dai Pra, R., Zhang, L., and Chang, R.B. (2022). A
multidimensional coding architecture of the vagal interoceptive system. Nature *603*.
https://doi.org/10.1038/s41586-022-04515-5.

24.   Berthoud, H.R., Albaugh, V.L., and Neuhuber, W.L. (2021). Gut-brain communication and obesity:
Understanding functions of the vagus nerve. Preprint, https://doi.org/10.1172/JCI143770
https://doi.org/10.1172/JCI143770.

25.   Buchanan, K.L., Rupprecht, L.E., Kaelberer, M.M., Sahasrabudhe, A., Klein, M.E., Villalobos, J.A.,
Liu, W.W., Yang, A., Gelman, J., Park, S., et al. (2022). The preference for sugar over sweetener
depends on a gut sensor cell. Nat Neurosci *25*. https://doi.org/10.1038/s41593-021-00982-7.

26.   Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y., Stoeckius,
M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*.
https://doi.org/10.1016/j.cell.2019.05.031.

27.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*.
https://doi.org/10.1016/j.cell.2021.04.048.

28.   Zhang, Y.-Y., Zhu, D.-X., Wang, M.-Y., Yi, Y.-T., Feng, Y.-H., Zhou, C., Li, C.-J., Liu, F., and Shen,
J.-F. (2025). Activation of NR2A-Wnt-TLR2 Signaling Axis in Satellite Glial Cells of the Dorsal Root
Ganglion Contributes to Neuropathic Pain Induced by Nerve Injury in Diabetic Mice. Mol Neurobiol
62.

29. Prato, V., Taberner, F.J., Hockley, J.R.F., Callejo, G., Arcourt, A., Tazir, B., Hammer, L., Schad,
P., Heppenstall, P.A., Smith, E.S., et al. (2017). Functional and Molecular Characterization of
Mechanoinsensitive "Silent" Nociceptors. Cell Rep *21*.
https://doi.org/10.1016/j.celrep.2017.11.066.

30. Kashio, M., and Tominaga, M. (2022). TRP channels in thermosensation. Preprint,
https://doi.org/10.1016/j.conb.2022.102591 https://doi.org/10.1016/j.conb.2022.102591.

31. Morales, M., and Wang, S. De (2002). Differential composition of 5-hydroxytryptamine3 receptors
synthesized in the rat CNS and peripheral nervous system. Journal of Neuroscience *22*.
https://doi.org/10.1523/jneurosci.22-15-06732.2002.

32. Kim, S.H., Patil, M.J., Hadley, S.H., Bahia, P.K., Butler, S.G., Madaram, M., and Taylor-Clark, T.E.
(2022). Mapping of the Sensory Innervation of the Mouse Lung by Specific Vagal and Dorsal Root
Ganglion Neuronal Subsets. eNeuro *9*. https://doi.org/10.1523/ENEURO.0026-22.2022.

33. Jami, S., Deuis, J.R., Klasfauseweh, T., Cheng, X., Kurdyukov, S., Chung, F., Okorokov, A.L., Li,
S., Zhang, J., Cristofori-Armstrong, B., et al. (2023). Pain-causing stinging nettle toxins target
TMEM233 to modulate NaV1.7 function. Nat Commun *14*. https://doi.org/10.1038/s41467-023-
37963-2.

34. Cai, H., Schnapp, W.I., Mann, S., Miscevic, M., Shcmit, M.B., Conteras, M., and Fang, C. (2024).
Neural circuits regulation of satiation. Appetite *200*.

35. Williams, E.K.K., Chang, R.B.B., Strochlic, D.E.E., Umans, B.D.D., Lowell, B.B.B., and Liberles,
S.D.D. (2016). Sensory Neurons that Detect Stretch and Nutrients in the Digestive System. Cell
166. https://doi.org/10.1016/j.cell.2016.05.011.

36. Dockray, G.J., and Burdyga, G. (2011). Plasticity in vagal afferent neurones during feeding and
fasting: Mechanisms and significance. Preprint, https://doi.org/10.1111/j.1748-1716.2010.02219.x
https://doi.org/10.1111/j.1748-1716.2010.02219.x.

37. Egerod, K.L., Petersen, N., Timshel, P.N., Rekling, J.C., Wang, Y., Liu, Q., Schwartz, T.W., and
Gautron, L. (2018). Profiling of G protein-coupled receptors in vagal afferents reveals novel gut-to-
brain sensing mechanisms. Mol Metab *12*. https://doi.org/10.1016/j.molmet.2018.03.016.

38. Yu, X., Yu, M., Liu, Y., and Yu, S. (2016). TRP channel functions in the gastrointestinal tract.
Preprint, https://doi.org/10.1007/s00281-015-0528-y https://doi.org/10.1007/s00281-015-0528-y.

39. Wang, Y., Li, G., Yu, K., Liang, S., Wan, F., Xu, C., Gao, Y., Liu, S., and Lin, J. (2009).
Expressions of P2X2 and P2X3 receptors in rat nodose neurons after myocardial ischemia injury.
Auton Neurosci *145*. https://doi.org/10.1016/j.autneu.2008.11.006.

40. Brouns, I., Adriaensen, D., Burnstock, G., and Timmermans, J.P. (2000). Intraepithelial vagal
sensory nerve terminals in rat pulmonary neuroepithelial bodies express P2X3 receptors. Am J
Respir Cell Mol Biol *23*. https://doi.org/10.1165/ajrcmb.23.1.3936.

41. Lee, L.Y., Hsu, C.C., Lin, Y.J., Lin, R.L., and Khosravi, M. (2015). Interaction between TRPA1 and
TRPV1: Synergy on pulmonary sensory nerves. Pulm Pharmacol Ther *35*.
https://doi.org/10.1016/j.pupt.2015.08.003.

42. Burdyga, G., Varro, A., Dimaline, R., Thompson, D.G., and Dockray, G.J. (2010). Expression of
cannabinoid CB1 receptors by vagal afferent neurons: Kinetics and role in influencing
neurochemical phenotype. Am J Physiol Gastrointest Liver Physiol *299*.
https://doi.org/10.1152/ajpgi.00059.2010.

43.   Búzás, B., and Cox, B.M. (1997). Quantitative analysis of Mu and delta opioid receptor gene
700   expression in rt brain and peripheral ganglia using competitive polymerase chain reaction.
701   Neuroscience *76*. https://doi.org/10.1016/S0306-4522(96)00242-4.

702   44.   Xiang, Z., Bo, X., and Burnstock, G. (1998). Localization of ATP-gated P2X receptor
703   immunoreactivity in rat sensory and sympathetic ganglia. Neurosci Lett *256*.
704   https://doi.org/10.1016/S0304-3940(98)00774-5.

705   45.   Zimmerman, K.D., Espeland, M.A., and Langefeld, C.D. (2021). A practical solution to
706   pseudoreplication bias in single-cell studies. Nat Commun *12*. https://doi.org/10.1038/s41467-021-
707   21038-1.

708   46.   Varga, G., Bálint, A., Burghardt, B., and D'Amato, M. (2004). Involvement of endogenous CCK
709   and CCK 1 receptors in colonic motor function. Preprint, https://doi.org/10.1038/sj.bjp.0705769
710   https://doi.org/10.1038/sj.bjp.0705769.

711   47.   Chaudhri, O., Small, C., and Bloom, S. (2006). Gastrointestinal hormones regulating appetite.
712   Preprint, https://doi.org/10.1098/rstb.2006.1856 https://doi.org/10.1098/rstb.2006.1856.

713   48.   Chey, W.Y., and Chang, T.M. (2003). Secretin, 100 years later. Preprint,
714   https://doi.org/10.1007/s00535-003-1235-3 https://doi.org/10.1007/s00535-003-1235-3.

715   49.   Patil, M.J., Meeker, S., Bautista, D., Dong, X., and Undem, B.J. (2019). Sphingosine-1-phosphate
716   activates mouse vagal airway afferent C-fibres via S1PR3 receptors. Journal of Physiology *597*.
717   https://doi.org/10.1113/JP277521.

718   50.   Nassenstein, C., Kwong, K., Taylor-Clark, T., Kollarik, M., Macglashan, D.M., Braun, A., and
719   Undem, B.J. (2008). Expression and function of the ion channel TRPA1 in vagal afferent nerves
720   innervating mouse lungs. Journal of Physiology *586*. https://doi.org/10.1113/jphysiol.2007.148379.

721   51.   Gu, Q., and Lee, L.Y. (2021). TRP channels in airway sensory nerves. Neurosci Lett *748*.
722   https://doi.org/10.1016/j.neulet.2021.135719.

723   52.   Cawthon, C.R., and de La Serre, C.B. (2021). The critical role of CCK in the regulation of food
724   intake and diet-induced obesity. Peptides (N.Y.) *138*.
725   https://doi.org/10.1016/j.peptides.2020.170492.

726   53.   Dockray, G.J. (2009). Cholecystokinin and gut-brain signalling. Preprint,
727   https://doi.org/10.1016/j.regpep.2009.03.015 https://doi.org/10.1016/j.regpep.2009.03.015.

728   54.   Rogers, R.C., and Hermann, G.E. (2008). Mechanisms of action of CCK to activate central vagal
729   afferent terminals. Peptides (N.Y.) *29*. https://doi.org/10.1016/j.peptides.2008.06.023.

730   55.   Smith, G.P., Jerome, C., Cushin, B.J., Eterno, R., and Simansky, K.J. (1981). Abdominal
731   vagotomy blocks the satiety effect of cholecystokinin in the rat. Science (1979) *213*.
732   https://doi.org/10.1126/science.7268408.

733   56.   Li, M., Tan, H.E., Lu, Z., Tsang, K.S., Chung, A.J., and Zuker, C.S. (2022). Gut–brain circuits for
734   fat preference. Nature *610*. https://doi.org/10.1038/s41586-022-05266-z.

735   57.   McDougle, M., de Araujo, A., Singh, A., Yang, M., Braga, I., Paille, V., Mendez-Hernandez, R.,
736   Vergara, M., Woodie, L.N., Gour, A., et al. (2024). Separate gut-brain circuits for fat and sugar
737   reinforcement combine to promote overeating. Cell Metab *36*.
738   https://doi.org/10.1016/j.cmet.2023.12.014.

739   58.   Buchanan, K.L., Rupprecht, L.E., Kaelberer, M.M., Sahasrabudhe, A., Klein, M.E., Villalobos, J.A.,
740   Liu, W.W., Yang, A., Gelman, J., Park, S., et al. (2022). The preference for sugar over sweetener
741   depends on a gut sensor cell. Nat Neurosci *25*. https://doi.org/10.1038/s41593-021-00982-7.

742   59.   Hayes, M.R., and Schmidt, H.D. (2016). GLP-influences food and drug reward. Preprint,
743   https://doi.org/10.1016/j.cobeha.2016.02.005 https://doi.org/10.1016/j.cobeha.2016.02.005.

744   60.   Rocca, A.S., and Brubaker, P.L. (1999). Role of the vagus nerve in mediating proximal nutrient-
745   induced glucagon- like peptide-1 secretion. Endocrinology *140*.
746   https://doi.org/10.1210/endo.140.4.6643.

747   61.   Chu, J.Y.S., Cheng, C.Y.Y., Sekar, R., and Chow, B.K.C. (2013). Vagal Afferent Mediates the
748   Anorectic Effect of Peripheral Secretin. PLoS One *8*.
749   https://doi.org/10.1371/journal.pone.0064859.

750   62.   Sekar, R., Wang, L., and Chow, B.K.C. (2017). Central control of feeding behavior by the secretin,
751   pacap, and glucagon family of peptides. Preprint, https://doi.org/10.3389/fendo.2017.00018
752   https://doi.org/10.3389/fendo.2017.00018.

753   63.   Yoo, P.B., Lubock, N.B., Hincapie, J.G., Ruble, S.B., Hamann, J.J., and Grill, W.M. (2013). High-
754   resolution measurement of electrically-evoked vagus nerve activity in the anesthetized dog. J
755   Neural Eng *10*. https://doi.org/10.1088/1741-2560/10/2/026003.

756   64.   Choi, H.M.T., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha,
757   A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction: Multiplexed,
758   quantitative, sensitive, versatile, robust. Development (Cambridge) *145*.
759   https://doi.org/10.1242/dev.165753.

760   65.   R Core Team (2023). R: A language and environment for statistical computing. Preprint at R
761   Foundation for Statistical Computing.

762   66.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
763   Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*.
764   https://doi.org/10.1016/j.cell.2021.04.048.

765   67.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
766   Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: An open-source platform for biological-
767   image analysis. Preprint, https://doi.org/10.1038/nmeth.2019 https://doi.org/10.1038/nmeth.2019.

768   68.   Hulsen, T. (2022). DeepVenn -- a web application for the creation of area-proportional Venn
769   diagrams using the deep learning framework Tensorflow. ArXiv.

770   69.   Gillen, C., Gleichmann, M., Spreyer, P., and Müller, H.W. (1995). Differentially expressed genes
771   after peripheral nerve injury. J Neurosci Res *42*. https://doi.org/10.1002/jnr.490420203.

772   70.   Xu, Q., Bernardo, A., Walker, D., Kanegawa, T., Mahley, R.W., and Huang, Y. (2006). Profile and
773   regulation of apolipoprotein E (ApoE) expression in the CNS in mice with targeting of green
774   fluorescent protein gene to the ApoE locus. Journal of Neuroscience *26*.
775   https://doi.org/10.1523/JNEUROSCI.5476-05.2006.

776   71.   Preibisch, S., Saalfeld, S., and Tomancak, P. (2009). Globally optimal stitching of tiled 3D
777   microscopic image acquisitions. Bioinformatics *25*. https://doi.org/10.1093/bioinformatics/btp184.

Figure 1. Putative mechanosensing and chemosensing clusters identified in left and right NG neurons in rats. A) Cartoon illustrating pooling of right and left NG from 18 male and 18 female rats into replicate groups. Created in BioRender. Thorn, C. (2024) BioRender.com/w17y601. B) After quality control and removal of non-neuronal and jugular ganglia clusters, 19 NG clusters were identified from integrated right and left samples (n = 23,428 nuclei). C) All NG clusters met quality control criteria and expressed high levels of the neuronal marker *Snap25*. D) Clusters can be divided into putative mechanosensing (gray boxes) versus chemosensing subtypes based on expression of *Piezo2.* Left: *Piezo2* expression overlaps with expression of *Scn1a and Cntn1* in most mechanosensory clusters; Right: In most chemosensory clusters, *Snc10a* expression overlaps with high levels of *Oprm1* and low levels of *Piezo2*.

Figure 2. Unique genetic profiles define NG neuronal clusters. A) Expression of the top 2 cluster-defining genes for each cluster. B) Across all clusters, expression of the top 5 cluster-defining genes was highly correlated between left and right NG (R^2 = 0.956, p < 0.0000, Pearson's correlation), indicating strong integration of data from the two sides prior to clustering; colors represent cluster identities as in Fig. 1B. C) Expression of select genes known to be expressed in gut-innervating NG neurons. For dot plots, average expression is plotted if the gene was expressed in > 10% of cells in the cluster.

A

←— Left   Right —→

-log10(p_val_adj)

log2FC

Cntn5
Bmper
Gria1
Shisa9
Sorcs1
Crtac1
Shisa9
Opcml Unc13c
Adora1 Rasgef1b
Gria1 Kcna2
Tns Adgrl2
Dgki Cdh10

B

100%

Male
Female

Left
Right

R   L   L   L

0%

14 9 18 12 10 11 16 15 6 17 4 0 8 13 7 3 1 2 5

Identity

C

Cluster 4 (% total cells)

***

Left   Right

D

Expression Level

3
Trpc6
0

4
Chrna3
0

3
Cckar
0

3
Sctr
0

5
Xylt1
0

14 9 18 12 10 11 16 15 6 17 4 0 8 13 7 3 1 2 5

Identity

Figure 3. Identification of right-side biased expression of a Chrna3 and Cckar co-expressing cluster. A) Pseudobulk analysis identified 15 genes with significant left-right differential expression distributed across Clusters 7, 3, 1, 2, and 5. B) Percent of cells in each cluster originating from left (light colors) versus right (dark colors) NG samples from male (blue) vs. female (magenta) rats. Dashed line denotes percentage of total cells across all clusters obtained from right NG samples. Cluster 4 (cyan arrow) was the only cluster that exhibited significantly greater than expected representation in right NG samples. R/L above bars denotes clusters found to have significantly greater than expected representation in right (R) or left (L) NG. C) The percentage of Cluster 4 cells in left-NG vs right-NG replicate groups. Paired data represent percentage of Cluster 4 cells for each replicate from male (blue) and female (magenta) rats; error bars denote SEM; ***p < 0.001, paired t-test. D) Cluster 4 is distinguished by expression of *Trpc6*, *Chrna3*, *Cckar*, *Sctr*, and *Xylt1*.

Figure 4. RNA-FISH validation of right-biased expression of NG neurons co-expressing Chrna3 and Cckar. A-D) RNA-FISH confirms the presence of a *Cckar* (A, red), *Chrna3* (B, blue), and *Xylt1* (C, green) co-labeled (D) population of cells in the left and right NG of rats. Merged images (D) also show DAPI-stained nuclei (cyan) and scale bar (100 μm) for A-D. E) Semi-quantitative Venn diagrams illustrating overlapping *Cckar*, *Chrna3*, and *Xylt1* labeling in left and right NG; circle area for each probe is proportional to the number of labeled cells per slice, averaged across rats. F-I) Density of cells in left vs. right NG that expressed *Cckar* (F), *Chrna3* (G), *Xylt1* (H), or were co-labeled for all 3 mRNAs (I). J-M) RNA-FISH confirms that all *Cckar*+ (J) and *Chrna3*+ (K) cells co-express the neuronal marker *Snap25* (L). Merged image (M) shows DAPI-labeled nuclei and scale bar (100 μm) for J-M. N) Semiquantitative Venn diagrams illustrating overlap of *Cckar*, *Chrna3*, and *Snap25* in left and right NG slices. O-Q) The percentage of *Cckar*+ (O) and *Chrna3*+ (P) cells in left vs. right NG, expressed as a percentage of the *Snap25*+ neuronal population (Q). R) Percentage of *Cckar*+ and *Chrna3*+ co-labeled neurons in left vs. right NG slices. Paired data in F-I and O-R represent averages across left vs. right NG slices for each rat (magenta: females; blue: males); error bars denote SEM; *p < 0.05, **p < 0.01, paired t-test.

Figure 5. Cckar+ NG neurons overlap with populations expressing Cartpt and Glp1r. A-D) RNA-FISH was used to probe for overlapping expression and right-side biased expression of *Cckar*+ (A, red), *Cartpt*+ (B, green), *Glp1r*+ (C, blue), and triple-labeled neurons (D) in left and right NG slices from rats. Merged images (D) also show DAPI-stained nuclei (cyan) and scale bar (100 µm) for A-D. E) Semiquantitative Venn diagrams, as in Figure 4E, illustrating overlap in *Cckar*, *Glp1r*, and *Cartpt* labeling in left and right NG. F-I) Density of cells in left vs. right NG that expressed *Cckar* (F), *Cartpt* (G), *Glp1r* (H), or were co-labeled for all 3 mRNAs (I). Paired data represent averages across left vs. right NG slices for each rat (magenta: females; blue: males); error bars denote SEM; $*p < 0.05$, paired t-test.

Figure 6. Mice also exhibit preferential expression of a Chrna3+ and Cckar+ subtype in right NG. A) Clustering identified 16 putative neuronal clusters in the integrated rat + mouse snRNA-seq dataset (top). Putative mechanosensitive versus chemosensitive clusters can be distinguished based on expression of *Piezo2*. As in rats, *Piezo2* expression overlaps with expression of *Scn1a* in most mechanosensory clusters, whereas most *Scn10a+* clusters in the integrated dataset are *Piezo2-* (bottom). B-C) Overlapping UMAP representations (B) and percentages of cells from each dataset distributed across identified clusters (C) indicate rat and mouse data were well integrated. D) Dot plot illustrating expression of top 2 marker genes for each cluster, plus *Cckar* expression across all clusters. Cluster 7, defined by high expression of *Trpc6* and *Chrna3*, and expressing high levels of *Cckar*, is highlighted (red arrow and shading). E) In both mouse and rat, co-expression of *Chrna3* and *Cckar* is predicted within Cluster 7, though expression of other marker genes (*Xylt1*) and digestive peptide signaling related genes (*Cartpt*, *Npy2r*, and *Vip*) diverges between the two species. F-I) RNA-FISH was used to quantify *Cckar* (F), *Chrna3* (G), and *Snap25* (H) expression, in left (top) and right (bottom) NG slices from mice; DAPI staining and scale bar (100 um) for F-I is shown in merged images (I). J) Semiquantitative Venn diagrams, as in Figure 4E, illustrating overlap in *Snap25*, *Cckar*, and *Chrna3* labeling in mouse NG slices. K-M) Percentage of *Cckar+* (K) and *Chrna3+* (L) cells in left vs. right NG, expressed as a percentage of the *Snap25+* neuronal population (M). N) Percentage of *Cckar+* and *Chrna3+* co-labeled neurons in left vs. right NG slices. Paired data in K-N represent averages across left vs. right NG slices for each mouse (magenta: females; blue: males); error bars denote SEM; *p < 0.05, **p < 0.01, paired t-test.

Figure 7. Right-biased responses to CCK1R activation in rat NG. In vitro calcium imaging was performed on primary cultures from left and right NG of rats during bath application of the CCK1R agonist, A71623. A) Example traces from neurons classified as A71623-responsive (top) or non-responsive (bottom) based on the presence or absence of an increase in calcium signal following drug application. B) Mean calcium signals of A71623-responsive (top) and non-nonresponsive (bottom) neurons from the left (blue) and right (red) NG. Shading represents 95% confidence intervals. C) Significantly more A71623-responsive neurons were found in right-NG cultures than in left-NG cultures; ***$p < 0.001$, Chi-square. D) Among neurons exhibiting A71623 responses, the peak amplitude of the calcium signal did not differ between left and right NG samples.

Figure S1. Quality control filtering and integration of replicate groups for rat dataset,
Related to Figure 1. A) Nuclei with greater than 6,500 features (left), greater than
20,000 molecules (center), or greater than 5% mitochondrial genes (right) were
removed prior to clustering. Before filtering, mean nFeature_RNA = 1857.668, mean
nCount_RNA = 3804.594, and mean mitochondrial gene percentage = 0.179%. B)
Samples from all 12 replicates (3 each from Male|Left (ML), Male|Right (MR),
Female|Left (FL), and Female|Right (FR) treatment groups) were integrated prior to
clustering. C) Samples from male and female rats exhibited overlapping UMAP profiles
after sample integration.

Figure S2. Filtering of clusters representing non-neuronal subtypes and jugular nodose

neurons in rat data, Related to Figure 1. Four rounds of clustering and filtering were

performed. A-C) In Round 1, two of 27 clusters were removed based on strong expression of marker genes associated with striated muscle (*Ttn*, *Myh4*; cluster 27) and myelin (*Mpz*, *Mbp*; cluster 12). A: Round 1 UMAP; B: dotplot showing expression of *Ttn*, *Myh4*, *Mpz*, and *Mbp* within each cluster; C: feature plots showing expression of targeted marker genes within filtered clusters. D-E) As in A-C for Round 2, in which 2 of 28 clusters were removed based on weak expression of the neuronal marker *Snap25* in conjunction with strong expression of the endothelial marker gene *Emcn* (cluster 2) and/or strong expression of the satellite glial marker genes *Apoe* and *Sparc* (clusters 2 & 3). G-I) As in A-C for Round 3 of cluster filtering, in which 4 of 24 clusters were removed based on strong expression of the melanocyte marker gene *Tyrp1* in conjunction with *Apoe* and *Sparc* (clusters 3, 4, 8, & 9). J-L) As in A-C for Round 4 of cluster filtering, in which 1 of 22 clusters was removed based on strong co-expression of *Tyrp*, *Apoe*, and *Sparc* (cluster 9), and 3 were removed based on strong expression of the jugular ganglia neuronal marker gene *Pdrm12*.

Figure S3. Quality control filtering of the rat + mouse integrated dataset, Related to Figure 6. A) For both datasets, nuclei with greater than 6,500 features (left), greater than 20,000 molecules (center), or greater than 5% mitochondrial genes (right) were removed prior to clustering. Dashed lines denote QC thresholds for each metric. Before filtering, mouse (red) mean nFeature_RNA = 2858.082, mean nCount_RNA = 7851.702, and mean percent mitochondrial genes = 2.922%. Rat data (blue) is as in Figure S1. B) After QC filtering and removal of non-neuronal cells and jugular ganglia neurons, remaining clusters all met quality control criteria and exhibited strong expression of the neuronal marker *Snap25*. C) Cluster 7 was the only cluster that exhibited expression of both *Cckar* and *Chrna3* in rats, and co-expression of these genes was likewise highest in mice.

Figure S4. Filtering of clusters representing non-neuronal subtypes and jugular nodose

neurons in the integrated rat + mouse dataset, Related to Figure 6. Three rounds of

clustering and filtering were performed. A-C) In Round 1, two of 24 clusters were

removed based on strong expression of marker genes associated with striated muscle

(*Ttn*, *Myh4*; cluster 24) and myelin (*Mpz*, *Mbp*; cluster 4). A: Round 1 UMAP; B: dotplot

showing expression of *Ttn*, *Myh4*, *Mpz*, and *Mbp* within each cluster; C: feature plots showing expression of targeted marker genes within filtered clusters. D-F) As in A-C for Round 2, in which 1 of 25 clusters was removed based on strong expression of the sympathetic neuron marker gene *Emcn* (cluster 9). G-I) As in A-C for Round 3 of cluster filtering, in which 1 of 22 clusters was removed based on strong expression of the endothelial marker gene *Mecom*, 1 cluster was removed based on strong expression of the oligodendrocyte marker gene *Daam2*, and 4 clusters were removed based on expression of the jugular ganglia neuronal marker gene *Prdm12* (clusters 13, 20, 21, and 22).