*Research Article*

**Tumoral Hypoxic Extracellular Vesicles Foster a Protective Microenvironment in**

**Triple-Negative Breast Cancer**

Bianca C. Pachane[1,2*], Pedro H. T. Bottaro[1], Aline M. Machado[1], Cynthia A. de Castro[3],

Gabriela Guerra[1], Larissa T. Gozzer[1,4], Marina M. Grigoli[5,6], Artur D. Zutião[5], Angelina M.

Fuzer[5], Marcia R. Cominetti[5], Wanessa F. Altei[2,7], Heloisa S. Selistre-de-Araujo[1]

[1] Biochemistry and Molecular Biology Laboratory, Department of Physiological Sciences, Universidade Federal
de São Carlos – UFSCar, São Carlos, SP, Brazil.

[2] Molecular Oncology Research Center, Barretos Cancer Hospital, Barretos, SP, Brazil.

[3] Pathology and Biocompatibility Laboratory, Department of Morphology and Pathology, Universidade Federal
de São Carlos - UFSCar, São Carlos, SP, Brazil.

[4] Universidade de Aveiro, Aveiro, Portugal

[5] Department of Gerontology, Universidade Federal de São Carlos - UFSCar, São Carlos, SP, Brazil.

[6] Departament of Psychiatry and Psychotherapy, University Medical Center Mainz, University of Johannes
Gutenberg of Mainz – JGU, Mainz, Germany

[7] Radiation Oncology Department, Barretos Cancer Hospital, Barretos, SP, Brazil

[*] Corresponding author: Bianca Cruz Pachane. Department of Physiological Sciences, Universidade Federal de
São Carlos – UFSCar, Rod. Washington Luis, km 235 – SP-310; 13565-905, São Carlos, SP, Brazil.
biancapachane@gmail.com.

Running Head: EVh protection of TNBC microenvironment

ORCID:

BCP: 0000-0002-7302-4498
PHTB: 0000-0002-6456-6244
AMM: 0000-0001-8047-8482
CAC: 0000-0002-2675-8069
GG: 0009-0008-9975-1125
MMG: 0000-0002-8212-9574
ADZ: 0009-0005-8844-8866
AMF: 0000-0003-0673-8287
MRC: 0000-0001-6385-7392
WFA: 0000-0002-6636-7951
HSSA: 0000-0002-2372-7814

**Graphical Abstract**

# Abstract

The highly metastatic triple-negative breast cancer (TNBC) relies on the tumor

microenvironment (TME) to maintain phenotypic heterogeneity and progression.

Extracellular vesicles from hypoxic TNBC (EVh) have been previously shown to facilitate

tumoral invasion; however, their function in the tumor microenvironment remains unclear.

We used a novel method to investigate the TME *in vitro* called multicellular circulating co-

culture, to characterize how EVh interferes with tumoral and endothelial cells, fibroblasts,

monocytes and macrophages. EVh promoted monocyte differentiation to M2-like

macrophages and inhibited macrophage-derived phagocytosis in endothelial and tumoral

cells. The protection of endothelial, tumoral and stromal cellular integrity by EVh increased

pro-tumoral and pro-angiogenic signaling, collagen matrix synthesis and showed a potential

differentiation to cancer-associated fibroblasts. Our findings highlight the critical role of EVh

in protecting tumor cells, indicating its cooperation towards a protective TME, which was

demonstrated by the multicellular circulating co-culture and conventional co-culture

protocols. These findings lead to an adequate system with potential for investigating other

tumor-related processes, including circulating tumor cells and metastasis.

*Keywords: phagocytosis; extracellular vesicles; hypoxia; tumor microenvironment.*

# Background

Triple-negative breast cancer (TNBC) is mainly characterized by the lack of expression of

receptors for estrogen, progesterone, and the human epidermal growth factor (HER2) (1).

Comprised of a heterogeneous mass with a high risk of metastasis, TNBC affects over 15%

of all breast tumors and presents poor overall survival (2). Its development is plagued with

hypoxia, a condition in which oxygen distribution is reduced due to the increased cellular

activity within the tumoral tissue, the inefficient tumoral vasculature, and the constant

competition for nutrients within the mass (3,4). Hypoxia and tumorigenesis favor the

secretion of extracellular vesicles (EV), nanoparticles (30-300 nm) released by all known

cells, with a major function in intercellular communication (5,6). We previously showed that

tumor-derived hypoxic EVs (EVh) favor pro-invasive pathways when compared to normoxic

tumoral EVs (7), but how they affect the TNBC tumor microenvironment (TME) remains

unknown.

The TME is a myriad of cellular and acellular components that evolve as the tumor

phenotypic heterogeneity grows according to the stages of cancer progression (8–10). In the

pre-invasive TME, autocrine signaling by epithelial (EGF), hepatocyte (HGF), and

transforming growth factors (TGF-β) favors cell proliferation, migration, and plasticity (11).

The extracellular matrix (ECM) is primarily modified by fibroblasts, specifically cancer-

associated fibroblasts (CAF), through protein synthesis and the expression of remodeling

enzymes, which alters the protein content, activity, and crosslinking to favor collagen

synthesis and deposition (12,13). Pro-inflammatory mediators induce monocyte

differentiation into macrophages and their subsequent polarization into tumor-associated

macrophages (TAM), whose phenotype may vary from the classically polarized, pro-

inflammatory M1 subtype to the alternatively polarized, anti-inflammatory M2 phenotype

(13). The increased cell activity of the tumor stimulates the release of reactive oxygen

species, which favors DNA damage and the secretion of pro-angiogenic stimuli, including

VEGF and fibroblast growth factor (FGF-2) (11). Once the invasive profile is achieved, cell

motility is favored by the polarization of the cytoskeleton, the interplay between adhesion

molecules, and ECM remodeling by matrix metalloproteases (MMPs) (14). These changes in

cell behavior result from the secretion of TGF-β, interleukins (IL-6 and IL-8), and VEGF by

CAFs, which favors the epithelial-mesenchymal transition, survival, and maintenance of

immune control (11).

The collective complexity of the TME renders its study challenging, particularly when using

*in vitro* techniques such as 2D cell culture. Due to the intrinsic method limitations, including

the restriction in the number of simultaneous cell cultures and the low correspondence with

the physiology of clinical TME, there has been a shift towards 3D cell culture models, which

add a novel layer of complexity to *in vitro* studies. However, one pitfall of these models is the

difficulty in observing individual cellular responses from the mass (15). In this work, we

describe a novel method for studying the TME *in vitro* called the multicellular circulating co-

culture (MC-CC), to determine how TNBC-derived hypoxic EVs influence individual

immortalized cellular models in a TME setting. Using this approach, we demonstrated for the

first time that EVh protects cells from macrophage-derived phagocytosis. This pro-

tumorigenic function of EVh confirms previous suppositions on the tumoral function of EVs

and supports the development of novel EV-based strategies to overcome resistance to

anticancer therapies.

## Materials and Methods

*Cell culture*

Human cells from triple-negative mammary adenocarcinoma MDA-MB-231 (ATCC® CRM-

HTB-26™) were grown between passages 36 and 45 in Leibovitz L-15 media supplemented

with 10% FBS (pH 7.0, Vitrocell) at 37 ℃. Dermal fibroblasts HDFa (ATCC® PCS-201-

012™) and umbilical cord endothelial cells HUVEC (ATCC® CRL-1730™) were maintained

between passages 6 and 15 in DMEM media supplemented with 10% FBS and 1% pen/strep

(pH 7.4, Vitrocell) at 37 ℃, 5% $CO_2$. Monocytes THP-1 (ATCC® TIB-202™) were kept

under suspension in RPMI 1640 media with added 10% FBS and 1% pen/strep (pH 7.0,

Vitrocell) at 37 ℃, 5% $CO_2$. Cells were grown under sterile conditions for no longer than 4

weeks after thawing and were frequently tested for *Mycoplasma* using the MycoAlert® Assay

kit (Lonza). For co-culture, cells were maintained in OptiMEM I Reduced Serum Media

supplemented with 1% pen/strep (Gibco).

*EVh separation and characterization*

MDA-MB-231 cells (3000 cells/cm$^2$) were seeded in EV-depleted media (antibiotic-free

DMEM supplemented with 10% UC-FBS, 4.6 g/L glucose and 1 mM sodium pyruvate) and

maintained for 24 h at a hypoxic atmosphere (1% $O_2$, 5% $CO_2$, 37 ℃; H35 Hypoxystation,

Don Whitley Scientific). The media was replaced by antibiotic-free OptiMEM for 48 hours,

and cells were further maintained for 48 hours in hypoxia. Hypoxic EVs (EVh) were

separated by differential ultracentrifugation and filtration, as described previously (7).

Following the MISEV2023 guidelines (16), EV preparations were characterized by

nanoparticle tracking analysis, transmission electron microscopy, protein quantification and

Western blotting.

*Nanoparticle Tracking Analysis (NTA)*

EVh samples were diluted in PBS (1:1000, v/v) and injected into the module of a Nanosight

NS300 (Malvern). Capture settings were set at camera level 16, slide shutter 1300, slider gain

512, FPS 25.0, and 749 frames, acquired in three-60s events. Analysis was performed on

NTA software (v. 3.2 Dev Build 3.2.16) using the detection threshold of 4.

*Protein quantification*

EVh samples were lysed with 2% SDS, diluted in deionized water (4:150) and added to a 96-

well plate with an albumin standard curve with known values (0-16 µg.ml). Using the

microBCA kit (Thermo-Fisher Scientific), the working solution was prepared (25A:24B:1C)

and added to samples, which were incubated for 2 hours at 37 ℃. Plates were evaluated using

a SpectraMax i3 plate reader (Molecular Devices) at $O.D._{562nm}$ and quantification was

considered successful if $R^2 < 0.99$.

*Western blotting*

MDA-MB-231 EVh (10 µl) and cell (10 µg) lysates were mixed with Laemmli buffer under

reducing conditions (1:4, v/v) and boiled at 100 ℃ for 5 minutes. Precision Plus Protein

Dual-Color (Bio-Rad) was used as the loading control. Samples were added to 10% SDS-

PAGE gels for electrophoresis at 100 V and proteins were transferred to nitrocellulose

membranes (0.45 µm, Bio-Rad) at 100 V, 4 ℃, for 2.5 hours in buffer (2.5 mM Tris, 2 M

glycine, 20% methanol; pH 7.6). Transfer efficiency was confirmed by Ponceau S membrane

staining. Membranes were blocked with 5% milk-TBST for 1 hour under agitation and

probed overnight for ALIX (1:1000, 186429, Abcam), Calnexin (1:1000, mAb 2679, Cell

Signaling), CD63 (1:1000, 59479, Abcam) or Flotillin-1 (1:1000, 61020, BD Biosciences).

After three 5-minute washes with TBST, membranes were incubated with respective

secondary antibodies (Goat anti-Rb, 1:15.000, #205718, Abcam; or Goat anti-Ms, 1:10.000,

#97040, Abcam) for 1 hour at RT under agitation. Membranes were washed four times with

PBST for 5 minutes and exposed to ECL substrates (Clarity™ Western ECL Substrate, Bio-

Rad and SuperSignal™ West Femto, Thermo-Fisher Scientific) for 2 minutes. The

chemiluminescence reaction was documented using ChemiDoc™ XRS$^+$ (Bio-Rad).

*Transmission Electron Microscopy (TEM)*

An EVh dilution (1:2, v/v in PBS) was deposited in copper grids covered with formvar and

carbon (Lot 051115, 01800 F/C, 200 mesh Cu, Ted Pella Inc.) for 20 min, RT. Samples were

fixed in 2% paraformaldehyde (PFA-PBS) for 20 min at RT and washed extensively in

deionized water. Grids were exposed to 4% Uranyl-acetate (pH 4) and 2% methylcellulose

solution for 10 min on ice and in the dark. Samples were air-dried and visualized in an FEI

TECNAI G2 F20 HRTEM microscope at 40,000x magnification.

*Monocyte differentiation in macrophages*

THP-1 cells were seeded in 6-well plates ($6x10^5$ cells/ml) in OptiMEM medium

supplemented with 1% pen/strep and treated with the vehicle control (i.e., PBS), EVh ($10^9$

particles/ml) or phorbol-12-myristate-13-acetate (PMA, 100 nM), following a previously

established protocol (17). Cells were incubated for 72 h at 37 ℃, 5% $CO_2$, and differentiation

to macrophages was determined by the adhesion of cells to the well, followed by phenotype

modification, assessed by light microscopy at 20x (Lux 2, CytoSMART).

*Cell viability assay*

At the end of cellular assays, a 10% resazurin solution (0.1 ml/ml, #199303, Sigma-Aldrich)

was prepared in OptiMEM and replaced the conditioned media of each well for 4 h at 37 ℃,

5% $CO_2$. A blank no-cell control and a 100%-reduced control, obtained from the reduction of

resazurin by heat, were maintained in parallel. The supernatant was transferred to a new plate

and analyzed by fluorescence ($\lambda_{Exc}$ 555 nm, $\lambda_{Em}$ 585 nm) using a plate reader (SpectraMax i3,

Molecular Devices). Cell viability was determined after normalization by the blank no-cell

control and displayed as the % of reduced resazurin.

*Time-lapse cell motility assay*

In a 24-well dish, MDA-MB-231 ($5x10^4$ cells/ml), HUVEC ($5x10^4$ cells/ml) or HDFa ($10^4$

cells/ml) were seeded in complete growth media (Leibovitz L-15 or DMEM supplemented

with 10% FBS and 1% pen/strep) for adhesion over 24 h at 37 ℃, 5% $CO_2$, followed by

media replacement with OptiMEM 1% pen/strep. THP-1 cells were diluted in OptiMEM 1%

pen/strep to achieve a $10^5$ cells/ml concentration. Cells were treated with EVh ($10^9$

particles/ml) for 24h at 37 ℃, 5% $CO_2$ and kept in parallel with an untreated control (i.e.,

PBS). Motility was evaluated under time-lapse light microscopy (Lux2, CytoSMART), with

snaps taken every 30 minutes, and single-cell tracking was processed using the *TrackMate*

plugin in FIJI (18,19).

*Enzyme-Linked Immunosorbent Assay - ELISA*

To evaluate the levels of cytokines IL-6, IL-10, TNF-α and IL-1β from the monocyte

differentiation assay and the multicellular co-culture assay, we used high-affinity microplates

sensitized with monoclonal anti-cytokine antibodies (IL-6: 555220; IL-10: 555157; TNF-α:

555212; IL-1β: 557953, OptEIA, BD Biosciences) for 16-18 h. Plates were washed thrice

(0.05% Tween 20-PBS), incubated in 4% FBS-PBS for 1 hour at RT, and washed again. The

conditioned media and a standard curve of recombinant cytokines were added to the wells for

2 h, RT, and the excess was washed off. Next, the anti-cytokine antibodies conjugated with

peroxidase were incubated for 1.5 h (RT) followed by five washes with 0.05% Tween 20. A

developing solution (3,3',5,5'-tetramethylbenzidine, TMB) was added to wells, and the

reaction was stopped with sulfuric acid. Plates were screened at $O.D._{450nm}$ (SpectraMax i3,

Molecular Devices) and sample concentrations were calculated from the titration curve of the

cytokine standards, expressed in pg/ml.

*Fluorescent gelatin invasion assay*

Black 96-well plates were coated with fluorescein-conjugated gelatin from pig skin (0.2

mg/ml, G13187, Molecular Probes) before cell seeding. MDA-MB-231, HUVEC, HDFa, or

THP-1 cells in their respective culture media were added to the gelatin-coated wells and

incubated overnight at 37 ℃, 5% $CO_2$. The media was replaced with OptiMEM 1% pen/strep

containing EVh ($10^9$ particles/ml) or PBS for 24 h at 37 ℃, 5% $CO_2$. After incubation, cell

viability was assessed using a resazurin-based fluorimetric assay, while the area of gelatin

degradation and cell morphology metrics were evaluated on FIJI (ImageJ) using

epifluorescence microscopy images, as previously described (7). The full protocol is

available on the protocols.io platform (20).

*Cell invasion in transwell co-culture*

To evaluate cell invasion in co-culture without contact between cell lines, fluorescein-

labelled gelatin-coated coverslips were prepared as described previously (7) and pre-

conditioned in 24-well plates with Leibovitz L-15 medium for 30 min at 37 ℃ before the

seeding of MDA-MB-231 cells (5 x $10^4$ cells/ml). In inserts with 0.3 µm membranes,

HUVEC ($5x10^4$ cells/ml) or HDFa ($10^4$ cells/ml) cells were seeded on the upper chamber in

DMEM 10% FBS. Inserts and coverslips were incubated for 24 hours at 37 ℃, 5% $CO_2$ for

cell adhesion. Once the monolayers were stable, a Boyden's chamber was assembled to

contain the gelatin-seeded MDA-MB-231 in the bottom chamber and an insert in the upper

chamber. The system was maintained in OptiMEM for 24h at 37 ℃, 5% $CO_2$. Cells and

conditioned media were processed following the full protocol described elsewhere (23).

Downstream analysis included the evaluation of cell morphology metrics, tumoral invasion

into fluorescent gelatin and matrix metalloprotease secretion by zymography.

*Gelatin zymography*

To assess MMP-2 and MMP-9 activity, conditioned media samples (10 µg) were mixed with

non-reducing Laemmli buffer (2:1, v/v) and loaded in technical duplicates into 10% gelatin

(100 µg/ml) SDS-PAGE gels with the Precision Plus™ Dual-Color Protein Ladder (Bio-Rad)

as loading control. A normalizing FBS sample (10 µg) was added to all gels for band

quantification. Electrophoresis was run at 4 ℃ under 80 V. Gels were washed with 2.5%

Triton X-100 for 40 min, RT, and submersed in refolding buffer (20 mM Tris, 5 mM $CaCl_2$, 1

µM $ZnCl_2$, pH 8.0) for 20 h at 37 ℃. Gels were stained with Coomassie Brilliant Blue

overnight, de-stained for band revelation, and photographed in ChemiDoc™ XRS[+] (Bio-Rad)

(22). Band densitometry was performed on FIJI (ImageJ).

*Gelatin invasion in direct co-culture*

MDA-MB-231 ($5x10^4$ cells/ml), HUVEC ($5x10^4$ cells/ml), HDFa ($10^4$ cells/ml), or THP-1

($10^5$ cells/ml) cells were stained either with CellTracker™ Red CMTPX (5 µM, C34552,

Invitrogen) or CellTrace™ CSFE (5 µM, C34554, Invitrogen) in OptiMEM 1% pen/strep

following the product datasheets. A choice of two cell lineages was simultaneously seeded

into wells of a fluorescein-conjugated gelatin-coated 96-well black plate. A control group

(PBS) was maintained alongside an EVh-treated group ($10^9$ particles/ml) at 37 ℃, 5% $CO_2$

for 24 h. After incubation, cell viability was assessed using a resazurin-based fluorometric

assay and the area of gelatin degradation and cell morphology metrics were evaluated on FIJI

(ImageJ) using epifluorescence microscopy images, as previously described (7). The full

protocol is available on the protocols.io platform (21).

*Multicellular circulating co-culture (MC-CC)*

MDA-MB-231 ($5x10^4$ cells/ml), HUVEC ($5x10^4$ cells/ml) and HDFa ($10^4$ cells/ml) cells

were seeded on gelatin, Matrigel, and FN-coated coverslips, respectively, overnight at 37 ℃,

5% $CO_2$. After adhesion, they were individually assembled in sterile QV500 (*Kirkstall*)

culture chambers with 1 ml of OptiMEM 1% pen/strep. THP-1 cells ($10^5$ cells/ml) were pre-

stained with CellTracker™ Red CMTPX (5 µM, C34552, Invitrogen) and transferred to the

reservoir containing EVh ($10^9$ particles/ml) or vehicle (PBS). The tubing system was closed

and connected to the MDA-MB-231, HUVEC, and HDFa chambers, respectively, and

propelled by a peristaltic pump set to a 50 µl/s flow rate. The circuit was incubated at 37 ℃,

5% $CO_2$ for 24h. Coverslips were individually processed for cell staining and

immunofluorescence, following the full protocol published elsewhere (24). The conditioned

media was collected, spun a 300 *x g* for 10 min, and the supernatant was probed for cytokines

via ELISA and MMPs by gelatin zymography.

*Individual static and circulating culture controls*

Cells seeded in ECM-coated coverslips (24) were assembled individually in a sterile QV500

(*Kirkstall*) culture chamber with 1 ml of OptiMEM 1% pen/strep and connected with the

tubing system to a reservoir containing 13 ml of media. Following the previous method, the

system was maintained with a peristaltic pump, parallel to a static individual control.

Coverslips were removed from the compartments, counterstained, and imaged at 40x

magnification (ImageXpress Micro XRS, Molecular Devices). Cell circularity index and

gelatin degradation were quantified using FIJI (7,18).

*Statistical analysis*

Cellular assays were repeated on three distinct occasions in technical triplicates. MC-CC was

repeated four times with both groups running simultaneously. Each sample was applied twice

on zymograms, and average band densitometry was used for statistics. Datasets were checked

for outliers using ROUT's test and distribution by Shapiro-Wilk (n < 9) or D'Agostino-

Pearson omnibus K2 (n ≥ 9). Parametric data was evaluated using unpaired t-tests (2 groups)

or ANOVA one-way with Tukey's multiple comparison test (3+ groups). Non-parametric

data was analyzed using the Mann-Whitney test (2 groups) or Kruskal-Wallis analysis of

variance with Dunn's multiple comparison test (3+ groups). Values of $p < 0.05$ were

considered statistically relevant (GraphPad Prism (v. 9.3)).

## 269 Results

### 270 *EVh characterization*

In this follow-up study, we maintained the isolation of EVh samples protocol published

before, where samples were thoroughly characterized by label-free proteomics in comparison

with their normoxic counterparts (7). Incorporating the recommendations outlined in the

MISEV2023 guidelines (16), EVh samples were separated from $2.2 \times 10^7$ MDA-MB-231

cells grown in hypoxia in serum-free media, leading to a population with modal size of 130.2

nm (± 78.7) and concentration of $3.69 \times 10^{11}$ particles/ml (± $8.74 \times 10^{10}$) (Fig. 1A). Samples

contained 236.6 µg/ml of protein and yielded 167,346.94 particles per cell or $1.07 \times 10^{-5}$ µg

protein per cell. Known EV biomarkers CD63, ALIX and flotillin-1 were abundant in EV

samples, whereas calnexin was absent (Fig. 1B). Visualization by TEM confirmed EVh

abundance and typical morphology (Fig. 1C).

**Figure 1. EVh characterization**. (A) Size dispersion and EV abundance by nanoparticle tracking analysis of MDA-MB-231-derived EVh. (B) Detection of ALIX, calnexin (CANX), CD63 and flotillin-1. (C) Wide-field and close-up TEM images of EVh. Scale bars: 200 nm and 100 nm.

### EVh-mediated THP-1 differentiation into M2-like macrophages

To evaluate whether EVh affected the homeostatic conditions of cells found in the TME, we designed individual experiments to assess behaviors like macrophage differentiation, cell viability, invasion and motility. Using the THP-1 monocytic cell line, which grows in suspension and acquires an adherent and broader phenotype upon differentiation into macrophages, we observed that both EVh- and PMA-treated groups became adherent after 16-24 hours of treatment, while untreated cells remained in suspension for up to 72 h (Fig. 2A and Suppl. 1A,B). In contrast, while the full well of PMA-treated cells differentiated into unpolarized macrophages, a reduced population of monocytes exposed to EVh fulfilled the

differentiation, which indicates a distinct differentiation mechanism between groups (Fig.

2B).

**Figure 2. EVh-mediated macrophage differentiation**. **A -** Timelapse tracking of THP-1 cells treated with

EVh. Snaps from the same site were compiled in 2-hour increments over 3 days (scale bar: 50 µm). **B –**

Representative images of THP-1 treated with vehicle (PBS), EVh or PMA after 72h differentiation (nuclei in

cyan and F-actin in gray; scale bar: 20 µm). **C –** Representative images of THP-1 cells (nuclei in cyan, F-actin

in magenta) treated with PBS or EVh on a gelatin coating (gray), with degradation spots identified by white

arrows (scale bar: 30 µm). **D –** Quantification of degraded gelatin area ($\mu m^2$). **E -** Cell circularity index for

quantification of THP-1 morphology. **F -** Cell viability of cells under invasion assay. **G –** Tracking of THP-1

trajectory over 24h. **H –** Average track speed of cells in 24 h (µm/s). **I -** Total distance traveled by cells in 24 h

(µm). **J-M -** Quantification of cytokines TNF-α, IL-1β, IL-6, and IL-10, respectively. Assays were repeated over

three different occasions with technical duplicates or triplicates. *p* values indicated above comparative bars with

statistical significance.

The EVh-induced group showed a reduction in gelatin invasion compared to the

undifferentiated control (Fig. 2C,D), but no changes were observed regarding cell

morphology metrics and viability (Fig. 2E,F). Cell tracking revealed undirected movement in

both untreated and PMA-differentiated cells, while EVh-differentiated cells exhibited

directional migration with higher mean speed and greater average distance traveled (Fig. 2G-

I). Inflammatory cytokine profiling of THP-1 conditioned media detected TNF-α and IL-1β

only in the PMA group (Fig. 2J,K), a non-significant trend of IL-6 reduction in the EVh

group (Fig. 2L), and continuity of IL-10 expression in all three groups (Fig. 2M). Overall,

these results suggest the macrophage-like cells derived from EVh signaling display a pro-

tumoral behavior, which is indicative of an M2-like phenotype.

***Tumoral reprogramming of endothelial cells in vitro to promote vascular permeability***

Regarding the endothelial cell model HUVEC, EVh promoted gelatin invasion and elongated

cell morphology metric, which is consistent with a pro-invasive phenotype (Fig. 3A-C).

Despite the clear indication of cell polarization, cell viability remained unaffected (Fig.

Suppl. 2A). Unlike THP-1, there was non-directional motility in EVh-exposed HUVEC (Fig.

3D). Although the average speed of movement was similar between the groups, a higher

distance traveled by cells was seen in EVh groups (Fig. 3E,F).

**Figure 3. Endothelial cells respond to tumoral signaling**. **A -** Representative images of HUVEC (nuclei in

cyan, F-actin in magenta) treated with PBS or EVh on a gelatin coating (gray), with degradation spots in black

(scale bar: 50 µm). **B –** Quantification of degraded gelatin area ($\mu m^2$) in individual HUVEC cultures. **C –**

HUVEC circularity index in individual culture. **D -** Tracking of HUVEC trajectory over 24h. **E –** Average track

speed of cells in 24 h (µm/s). **F -** Total distance traveled by cells in 24 h (µm). **G –** Experimental design of the

indirect co-culture assay. **H –** Representative images of HUVEC (nuclei in cyan, F-actin in yellow; scale bar: 50

µm) in indirect co-culture with MDA-MB-231 (nuclei in cyan, F-actin in magenta; scale bar: 100 µm). **I –** Cell

circularity index for quantification of HUVEC and MDA-MB-231 morphology in direct co-culture. **J –**

Representative images of gelatin matrix (gray) with degradation spots identified by white arrows. **K -** MMP-2

detection by gelatin zymography and densitometry analysis of indirect co-culture. **L –** Quantification of

degraded gelatin area ($µm^2$) in indirect co-culture. **M -** Experimental design of the direct co-culture assay. **N –**

Representative images of HUVEC (cytoplasm in yellow) in direct co-culture with MDA-MB-231 (nuclei in

cyan, F-actin in magenta) seeded atop a gelatin coating (gray) (scale bar: 100 µm). **O -** Cell circularity index for

quantification of combined cell morphology in direct co-culture. **P -** Quantification of degraded gelatin area

($µm^2$) in direct co-culture. **Q –** Cell viability of HUVEC and MDA-MB-231 direct co-culture. Assays were

repeated on three different occasions with technical triplicates. *p* values indicated above comparative bars with

statistical significance.

To explore a more complex level of tumoral communication, the pro-tumoral effect of EVh

in endothelial cells was further investigated using indirect and direct co-culture assays with

MDA-MB-231 cells. In indirect co-culture, where tumor cells were separated from

endothelial cells by a porous membrane (Fig. 3G), HUVEC exhibited typical morphology and

significant intercellular interaction, while the tumoral cells showed multiple cytoskeleton

extensions, which gave them a downy appearance (Fig. 3H). In EVh-treated conditions, the

cytoskeleton retracted as endothelial cells visibly lost intercellular interactions, and tumor

cells showed a more elongated phenotype, without F-actin extensions (Fig. 3I). Tumoral

gelatin invasion was detected in both co-culture groups, likely due to MMP-2 activation,

which was detected only in its active form (Fig. 3J-K; Suppl. Fig. 2B). Both MMP-2 activity

and gelatin degraded area were slightly higher in EVh-treated groups (Fig. 3L). Adding to

previous results (7), we found EVh directs movement in MDA-MB-231, which facilitates

tumoral invasion in gelatin matrices (Fig. Suppl. 2C-E).

In the direct co-culture approach, where both cell types were seeded in the same well

simultaneously (Fig. 3M), close proximity triggered a stress response characterized by a star-

shaped rearrangement of the endothelial cytoskeleton, while tumor cells appeared more

cohesive (Fig. 3N). Upon treatment with EVh, cells regained their original morphological

features (Fig. 3N; Suppl. Fig. 2F,G). These alterations were evident in the circularity index,

highlighting the role of EVh in modulating intercellular interactions within the TME (Fig.

3O). Similar to the indirect co-culture model, EVh-treated cells also showed enhanced gelatin

invasion (Fig. 3P–Q). Under these conditions, cell viability was unaffected (Fig. 3R).

***EVh modulates fibroblast behavior and suppresses tumor-associated invasion***

Fibroblasts play a pivotal role in the tumor microenvironment by supporting tumor

progression and remodeling the extracellular matrix. In this context, HDFa cells exhibited

pronounced gelatinase activity, which was further enhanced following EVh treatment (Fig.

4A). This was accompanied by a migratory phenotype characterized by increased gelatin

matrix degradation and a significant elevation in cell viability upon EVh administration (Fig.

4B–D). EVh also induced a polarized migration pattern in HDFa, reflected by decreased track

speed and increased total distance traveled (Fig. 4E–G).

To further assess fibroblast responses in distinct tumor-associated contexts, an indirect co-

culture assay was established between HDFa and gelatin-precoated MDA-MB-231 cells. In

this setup, fibroblasts displayed only modest alterations in response to tumor-derived cues,

regardless of EVh treatment (Fig. 4H–I). A reduction in cell area was observed, leading to a

broader morphology compared to monoculture controls, while gelatin degradation by tumor

cells remained largely unaltered (Fig. 4J–K). These morphological changes were supported

by a decreased circularity index. Zymographic analysis indicated that EVh treatment reduced

MMP-2 activity in this context (Fig. 4L).

In a direct co-culture model, EVh appeared to suppress pro-invasive characteristics. Both

fibroblasts and tumor cells displayed distinct morphological alterations (Fig. 4M, Suppl. Fig.

2H), with increased cellular circularity (Fig. 4N) and reduced gelatinase activity following

EVh exposure (Fig. 4O). Notably, cell viability was unaffected under these conditions (Suppl.

Fig. 2I).

**Figure 4. Fibroblast response to tumoral signaling**. **A -** Representative images of HDFa (nuclei in cyan, F-actin in magenta) treated with PBS or EVh on a gelatin coating (gray), with degradation spots identified by white arrows (scale bar: 100 µm). **B –** Cell circularity index for quantification of HDFa morphology in individual culture. **C -** Quantification of degraded gelatin area ($\mu m^2$) in individual HUVEC cultures. **D –** HDFa cell viability in individual culture. **E -** Tracking of HDFa trajectory over 24h. **F –** Average track speed of cells in 24 h (µm/s). **G -** Total distance traveled by cells in 24 h (µm). **H –** Representative images of HDFa (nuclei in cyan, F-actin in yellow; scale bar: 100 µm) in indirect co-culture with MDA-MB-231 (nuclei in cyan, F-actin in magenta; scale bar: 30 µm). **I –** Cell circularity index for quantification of HDFa and MDA-MB-231 morphology in indirect co-culture. **J -** Representative images of gelatin matrix (gray) with degradation spots identified by white arrows. **K -** Quantification of degraded gelatin area ($\mu m^2$) in indirect co-culture. **L –** MMP-2 detected by gelatin zymography and densitometry analysis to identify gelatinase activity in indirect co-culture. **M –** Representative images of MDA-MB-231 (cytoplasm in yellow) in direct co-culture with HDFa (nuclei in cyan, F-actin in magenta) seeded atop a gelatin coating (gray) (scale bar: 100 µm). **N -** Cell circularity index for quantification of combined cell morphology in direct co-culture. **O –** Quantification of degraded gelatin area ($\mu m^2$) in direct co-culture. Assays were repeated on three different occasions with technical triplicates. *p* values indicated above comparative bars with statistical significance.

## *EVh attenuates phagoptosis and promotes pro-tumoral features in a multicellular circulating co-culture model*

A multicellular circulating co-culture (MC-CC) was developed to recreate the cellular complexity of a TNBC tumor *in vitro* for evaluating EVh response in a TME-like setting (Fig. 5A). In the system, each TME cell type was seeded in a compartment for individual assessment of targets, while maintaining communication by a constant flow of 50 µl/s, set by previous optimization steps containing isolated cells (Fig. Suppl. 3A). A closed system comprised of MDA-MB-231, HUVEC, HDFa and THP-1 cells circulated for 24 hours showed high levels of phagoptosis (i.e., death by macrophage phagocytosis) (25), especially in the TNBC and endothelial cell chambers (Fig. 5B). In these chambers, integer nuclei and cytoplasm were seldomly found. EVh exposure led to more frequently sighted cells with integer nuclei and cytoplasm throughout chambers, suggesting phagoptosis inhibition. Cell

morphological plasticity was confirmed by the circularity indexes of MDA-MB-231 and

HUVEC (Fig. Suppl. 3B-C). The media flow reduced gelatin invasion in TNBC cells

compared to individual static controls. EVh treatment partially recovered the cell invasive

ability (Fig. 5C). The conditioned media was tested for gelatinases, and MMP-2 activity

remained unchanged (Fig. 5D, Suppl. 3D). Cytokine IL-1β was present in similar levels in

both groups, but IL-6 was more abundant in the EVh-treated system (Fig. 5E,F).

**Figure 5. EVh inhibits phagoptosis in the TME**. **A –** Experimental design of the multicellular circulating co-culture assay (MC-CC) using the QV500 system (Kirstall). **B -** Representative images of MC-CC chambers containing MDA-MB-231, HUVEC or HDFa treated with PBS or EVh. (scale bars: 30 µm and 50 µm). **C -** Quantification of degraded gelatin area (µm²) in the MC-CC MDA-MB-231 chamber. **D –** Band densitometry of MMP-2 detected by gelatin zymography from the MC-CC conditioned media. **E,F –** Cytokines IL-1β and IL-6 probed from the MC-CC conditioned media by ELISA. **G,H –** Relative fluorescence intensity of β-catenin and VEGFR2 in HUVEC, respectively. **I,J –** Relative fluorescence intensity of collagen II and III in HDFa,

respectively. Assays were repeated on four different occasions with a single replicate. *p* values indicated above

comparative bars with statistical significance.

Pro-tumoral features were assessed through quantitative immunofluorescence analysis. In

endothelial cells, the expression of β-catenin and VEGFR2—key mediators of intercellular

adhesion and the angiogenic cascade, respectively—was upregulated in the EVh-treated

group (Fig. 5G,H). Among the cell types evaluated, fibroblasts exhibited the highest degree

of structural preservation, with only minor morphological alterations detected (Fig. Suppl.

4E). Regarding extracellular matrix (ECM) remodeling, EVh exposure led to increased

expression of collagen types II and III, predominantly localized around the pericellular region

(Fig. 5I,J). Due to technical limitations, collagen type I expression was not analyzed.

Macrophage distribution within the MC-CC model revealed a predominant localization of

THP-1 cells in the MDA-MB-231 chamber (Fig. 6A). Notably, the frequency of macrophage

presence was reduced in EVh-treated conditions compared to controls, supporting the

hypothesis that EVh mitigates phagoptosis within the tumor microenvironment. To further

validate this observation, direct co-culture assays between THP-1 cells and each individual

cell type were conducted. These experiments recapitulated the MC-CC findings, with

elevated levels of phagoptosis observed in MDA-MB-231 and HUVEC cultures, and

evidence of cellular debris present in fibroblast cultures (Fig. 6B). EVh treatment attenuated

macrophage phagocytic activity and preserved the original morphology of all cell types.

Interestingly, THP-1 cells formed a peripheral layer around fibroblasts, indicative of potential

bidirectional communication. In this condition, fibroblasts acquired a more branched,

polarized morphology, consistent with an activated state (Fig. 6C). Cell viability was

unaffected by EVh across all co-culture settings (Fig. Suppl. 3F). However, EVh treatment

enhanced gelatin invasion in tumor and endothelial cells following co-culture with THP-1,

while fibroblast invasiveness was reduced (Fig. 6D).

**Figure 6. Macrophage remodeling by EVh in TME models *in vitro* A –** Representative images of THP-1 (magenta, stained with CellTracker CMPTX) attachment to MDA-MB-231 MC-CC chambers (nuclei in cyan, F-actin in gray; scale bar: 30 µm). **B -** Representative images of THP-1 cells (magenta, stained with CellTrace CSFE) in direct co-culture with MDA-MB-231, HUVEC or HDFa (nuclei in cyan, F-actin in gray; scale bars: 50 µm). **C -** Cell circularity index for quantification of combined cell morphology in THP-1 direct co-culture with MDA-MB-231, HUVEC and HDFa. **D –** Quantification of degraded gelatin area (µm²) in THP-1 direct co-culture with MDA-MB-231, HUVEC and HDFa. Assays were repeated on three different occasions with technical triplicates. *p* values indicated above comparative bars with statistical significance.

# Discussion

The role of EVs in tumor progression and development is well described, and as new facets of tumorigenesis are revealed, their importance to cancer continuity becomes apparent.

Tumor-associated EVs have been implicated in promoting angiogenesis (26), immune

modulation (27), tumoral invasion (7,28), migration (29) and organotropic metastasis (30).

The establishment of hypoxia in tumor progression promotes EV secretion in breast cancer,

with distinct composition and characteristics (5–7). After describing the pro-invasive

potential of hypoxic EV in comparison to normoxic EVs from TNBC cell line MDA-MB-231

(7), we show in this work that EVh differentiated THP-1 cells in M2-like macrophages,

downregulating phagoptosis in tumor microenvironment models *in vitro*. Moreover, under the

MC-CC model, EVh protected endothelial and tumoral cellular integrity, promoted pro-

tumoral responses in endothelial cells and induced collagen synthesis in fibroblasts.

Monocyte recruitment and activation are crucial steps of the tumor-induced immune cascade,

often triggered by the macrophage colony-stimulating factor (M-CSF) and other

environmental cues (31). The differentiation of monocytes to macrophages *in vitro* relies on

PMA to generate non-polarized cells, which may be induced to a pro-inflammatory M1

phenotype or an anti-inflammatory M2 profile, each with a distinct cytokine secretion pattern

(17,32). We found that PMA-induced cells were enriched with pro-inflammatory components

IL-1β, IL-6 and TNF-$\square$, suggesting an M1-like profile, and the subtle reduction in IL-6 and

sustained expression of anti-inflammatory IL-10 suggested that EVh promoted an M2-like

phenotype (32–35).

The macrophage profiles were consistently maintained during co-culture with tumoral and

endothelial cells, where phagoptosis was abundant in the control groups but avoided with

EVh treatment. Cell death by phagocytosis results from the imbalance between the eat me

and don't eat me signals from the target cell and their recognition by receptors in the

phagocytic cell (25). In our models, EVh likely modulated don't eat me signals, such as

CD47 and MHC-I, to improve cell survival, corroborating previous findings in which

hypoxia inhibited macrophage phagocytosis in a HIF-1□-dependent manner (36,37). Further

studies are required to confirm this hypothesis.

Studying the TME *in vitro* is challenging due to methodological, practical, and technological

limitations. The recent development of techniques such as multicellular co-cultures,

microfluidics and 3D culture has improved the complexity with which we can investigate

basic mechanisms (38–40). A significant limitation of multicellular 3D co-cultures is the

difficulty of evaluating responses in individual cells from the cultured mass, which often

relies upon more refined and less accessible techniques (15). To circumvent this issue, we

described a novel method to study the TME *in vitro* where cells are cultured *in tandem* in a

commercially available system, henceforth named a multicellular circulating co-culture (MC-

CC). Results obtained from the MC-CC model were complementary to other co-culture

techniques, including direct and indirect co-culture, but we have not investigated whether it is

a better alternative to 3D culture. This method was previously described for studying the

blood-brain barrier but has several limitations, including the number of available co-culture

chambers, flow shear force, and lack of direct contact between chambers which reduces our

ability to evaluate certain cellular responses (41). We acknowledge that different

experimental designs, including other cell types, order of chambers, incubation conditions,

flow rate, and media composition, may interfere with the results and have not been

investigated in the present study.

Our experimental design permitted the evaluation of cell morphology alongside cytokine and

enzyme secreted in the circulating medium across all MC-CC chambers. EVh cause stromal

and tumor cells to favor a more pro-tumoral phenotype, likely due to the increased levels of

IL-6, a cytokine whose overexpression is closely related to the development of various

tumors, regulation of the acute phase of inflammation, and modulation of T and B

lymphocytes (42). Furthermore, the high levels of IL-6 and IL-1β favor the inflammatory

phenotype of cancer-associated fibroblasts, which is consistent with our evaluation of

fibroblast response (43). There was a non-statistical tendency for an increase in IL-1β levels

in EVh-treated cells, suggesting fibroblast activation, but additional testing to detect

differences in □-smooth muscle actin (□-SMA) is required to confirm this supposition.

EVh in endothelial cells under MC-CC increased the expression of markers for cell-cell

adhesion and proangiogenicity, including VEGFR2 and β-catenin (44). This may facilitate

tumor development and metastatic spread by increasing the connections between endothelial

and tumoral cells and encouraging angiogenesis (44,45). Intercellular junctions are crucial for

endothelial integrity and depend on the function of multiple proteins, especially VE-cadherin

(46). With nearly exclusive expression in vascular endothelium, VE-cadherin is

intracellularly linked to β-catenin, which directly connects to the cytoskeleton (47,48). In the

indirect co-culture assay, we could not identify which adhesive molecules were influenced by

EVh; however, multicellular co-culture under flow suggested an increase in β-catenin

expression in HUVEC under EVh action. One possible interpretation for the increased β-

catenin is that, since HUVECs are not fully confluent and VE-cadherin localization at the cell

membrane is not evident, β-catenin may be predominantly cytoplasmic or even translocated

to the nucleus, where it can promote transcriptional programs associated with cell migration

rather than strengthening cell-cell junctions. This scenario aligns with the observed increase

in VEGFR2 expression and suggests that β-catenin upregulation may contribute to enhanced

endothelial migration and angiogenic activity rather than contradicting the pro-adhesive

environment. These findings underscore the complexity of endothelial responses within the

tumor microenvironment and warrant further investigation. Furthermore, we suggest that

EVh disrupts endothelial cell adhesion to aid EVh-mediated tumor cell invasion.

Fibroblasts are structural support cells that can associate with tumors, transforming into

cancer-associated fibroblasts (CAFs) through prolonged tumor-associated inflammation

(TAI) signaling (43). TAI is sustained by continuous tumor mass growth and hypoxia, which

have been described as determinants of the inflammatory phenotype of fibroblasts in

pancreatic cancer (49). EVh increased fibroblast activity, modified their morphology and, in

MC-CC, promoted the synthesis of type II and type III collagen, Both collagens are fibrillar

and associated with increased ECM stiffness, a crucial factor for metastasis (50). Hypoxic

extracellular vesicles also play a significant role in modulating fibroblast activity, leading to

increased production of collagen types II and III. This process of extracellular matrix

remodeling through MMP production and excessive collagen formation has been shown to

promote tumor growth, progression, and resistance to treatments such as radiotherapy

(44,45,51,52). In this setting, we show that hypoxic extracellular vesicles from triple-negative

breast cancer impact fibroblast activity.

## Conclusion

In summary, we found that tumor-derived hypoxic EVs show a distinct immune-modulating

function, allowing the differentiation of M2-like macrophages in a TNBC in the TME *in vitro*

and inhibiting endothelial and tumoral phagoptosis. EVh administration protected the

endothelial and tumoral cellular integrity from phagoptosis. It promoted pro-tumoral and pro-

angiogenic responses in endothelial cells, inducing collagen synthesis in fibroblasts and their

potential differentiation to CAFs. In addition, our work describes a novel method for studying

the TME *in vitro* using a commercial system with great potential for investigating other

tumor-related processes, including circulating tumor cells and metastasis.

## Author's contribution

BCP: Conceptualization; Data curation; Formal Analysis; Funding acquisition; Investigation;

Methodology; Project administration; Software; Writing – original draft. PHTB: Formal

Analysis; Investigation; Visualization; AMM: Formal Analysis; Investigation; Writing –

review & editing; CAC: Formal Analysis; Investigation; Methodology; Writing – review &

editing; GG: Investigation; Visualization; LTG: Methodology; Visualization; MMG:

Investigation; Writing – review & editing; ADZ: Investigation; Visualization; AMF:

Conceptualization; Resources; Writing – review & editing; MRC: Funding acquisition;

Resources; Software; Writing – review & editing; WFA: Conceptualization; Resources;

Supervision; Writing – review & editing; HSSA: Conceptualization; Funding acquisition;

Resources; Supervision; Writing – review & editing.

## Declarations

570 The authors (BCP, PHTB, AMM, CAC, GG, LTG, MMG, ADZ, AMF, MRC, WFA, HSSA)

571 declare this study complies with the highest ethical guidelines and biosafety regulations. This

572 study does not involve clinical or pre-clinical samples and did not require consent to

573 participate. All authors declare no competing interests in this study and consent to

574 publication. This study does not contain artificial intelligence-generated data.

## Acknowledgments

The authors thank members from LMBBM and LABEN for critical advice. We thank the

Laboratory of Structural Characterization (LCE/DEMa/UFSCar) for the electron microscopy

facilities; Otavio Thiemann and Glaucius Oliva (IFSC/USP) for Beckman ultracentrifuge and

rotors; Fausto Bruno dos Reis Almeida for the use of Nanosight NS300; Monica Rosa da

Costa Iemma for the HDFa cell lineage; Gerson Jhonatan Rodrigues and Luis Henrique

Oliveira de Moraes for the HUVEC cell lineage.

## Data availability

583 The data generated in this study are available within the article and its supplementary data

584 files. Raw files are available upon reasonable request to the corresponding author. Detailed

protocols have been submitted to the "protocols.io" platform (20,21,23,24). We have

submitted all relevant data of our EV experiments to the EV-TRACK knowledgebase (EV-

TRACK ID: EV240149) (53).

## Funding

This work was funded by the São Paulo Research Foundation through grants 2019/05149-9,

2021/01983-4, and 2022/04146-9 (to BCP); 2022/12307-2 (to PHTB); 2020/11328-0 (to

LTG); 2017/01287-2 (to AMF); 2021/01863-9 and 2021/14673-3 (to MRC); 2019/11437-7

(to HSSA); Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), code

001; and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) through

grants 143120/2023-9 (ID 1618, to GG); 174872/2023-2 (to AMF). The funding agencies had

no direct involvement in the study's design, conduct, analysis, or reporting.

## References

1. Foulkes William D., Smith Ian E., Reis-Filho Jorge S. Triple-Negative Breast Cancer. New England Journal of Medicine. 2010;363(20):1938–48.

2. Altei WF, Pachane BC, Souza C, Marques MMC, Selistre-de-Araújo H. New insights into the discovery of drugs for triple-negative breast cancer metastasis. Expert Opinion on Drug Discovery. 2022 Apr 3;17(4):365–76.

3. Majmundar AJ, Wong WJ, Simon MC. Hypoxia-Inducible Factors and the Response to Hypoxic Stress. Molecular Cell. 2010 Oct 22;40(2):294–309.

4. Zhang Y, Zhang H, Wang M, Schmid T, Xin Z, Kozhuharova L, et al. Hypoxia in Breast Cancer—Scientific Translation to Therapeutic and Diagnostic Clinical Applications. Front Oncol. 2021 Mar 11;11:652266.

5. King HW, Michael MZ, Gleadle JM. Hypoxic enhancement of exosome release by breast cancer cells. BMC Cancer. 2012 Sep 24;12(1):421.

6. Xu R, Rai A, Chen M, Suwakulsiri W, Greening DW, Simpson RJ. Extracellular vesicles in cancer — implications for future improvements in cancer care. Nat Rev Clin Oncol. 2018 Oct;15(10):617–38.

7. Pachane BC, Nunes ACC, Cataldi TR, Micocci KC, Moreira BC, Labate CA, et al. Small Extracellular Vesicles from Hypoxic Triple-Negative Breast Cancer Cells Induce

Oxygen-Dependent Cell Invasion. International Journal of Molecular Sciences. 2022
Jan;23(20):12646.

8. Jin MZ, Jin WL. The updated landscape of tumor microenvironment and drug
repurposing. Sig Transduct Target Ther. 2020 Aug 25;5(1):1–16.

9. Kucharzewska P, Belting M. Emerging roles of extracellular vesicles in the adaptive
response of tumour cells to microenvironmental stress. Journal of Extracellular Vesicles.
2013;2(1):20304.

10. Dagogo-Jack I, Shaw AT. Tumour heterogeneity and resistance to cancer therapies. Nat
Rev Clin Oncol. 2018 Feb;15(2):81–94.

11. Visser KE de, Joyce JA. The evolving tumor microenvironment: From cancer initiation to
metastatic outgrowth. Cancer Cell. 2023 Mar 13;41(3):374–403.

12. Amos SE, Choi YS. The Cancer Microenvironment: Mechanical Challenges of the
Metastatic Cascade. Front Bioeng Biotechnol [Internet]. 2021 Feb 12 [cited 2024 Apr
6];9. Available from: https://www.frontiersin.org/articles/10.3389/fbioe.2021.625859

13. Yuan Z, Li Y, Zhang S, Wang X, Dou H, Yu X, et al. Extracellular matrix remodeling in
tumor progression and immune escape: from mechanisms to treatments. Molecular
Cancer. 2023 Mar 11;22(1):48.

14. Hamidi H, Ivaska J. Every step of the way: integrins in cancer progression and
metastasis. Nat Rev Cancer. 2018 Sep;18(9):533–48.

15. Dey M, Ayan B, Yurieva M, Unutmaz D, Ozbolat IT. Studying tumor angiogenesis and
cancer invasion in a 3D vascularized breast cancer microenvironment. Adv Biol (Weinh).
2021 Jul;5(7):e2100090.

16. Welsh JA, Goberdhan DCI, O'Driscoll L, Buzas EI, Blenkiron C, Bussolati B, et al.
Minimal information for studies of extracellular vesicles (MISEV2023): From basic to
advanced approaches. Journal of Extracellular Vesicles. 2024;13(2):e12404.

17. Starr T, Bauler TJ, Malik-Kale P, Steele-Mortimer O. The phorbol 12-myristate-13-
acetate differentiation protocol is critical to the interaction of THP-1 macrophages with
Salmonella Typhimurium. PLoS One. 2018 Mar 14;13(3):e0193601.

18. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012 Jul;9(7):676–82.

19. Ershov D, Phan MS, Pylvänäinen JW, Rigaud SU, Le Blanc L, Charles-Orszag A, et al.
TrackMate 7: integrating state-of-the-art segmentation algorithms into tracking pipelines.
Nat Methods. 2022 Jul;19(7):829–32.

20. Pachane BC, Araujo HSS de. Fluorescent Gelatin Degradation Assay to Evaluate EVh
Action in TME Cells. 2024 Sep 9 [cited 2024 Sep 16]; Available from:
https://www.protocols.io/view/fluorescent-gelatin-degradation-assay-to-evaluate-
dkt34wqn

21. Pachane BC, Araujo HSS de. Cell Invasion in Direct Co-Culture. 2024 Sep 9 [cited 2024
Sep 16]; Available from: https://www.protocols.io/view/cell-invasion-in-direct-co-
culture-dkws4xee

22. Leber TM, Balkwill FR. Zymography: A Single-Step Staining Method for Quantitation of
Proteolytic Activity on Substrate Gels. Analytical Biochemistry. 1997 Jun 15;249(1):24–
8.

23. Pachane BC, Araujo HSS de. Indirect Co-Culture Assay using Boyden Chambers. 2024
Sep 10 [cited 2024 Sep 16]; Available from: https://www.protocols.io/view/indirect-co-
culture-assay-using-boyden-chambers-dkxc4xiw

24. Pachane BC, Bottaro PHT, Altei WF, Araujo HSS de. Multicellular Circulating Co-
Culture. 2024 Sep 10 [cited 2024 Sep 16]; Available from:
https://www.protocols.io/view/multicellular-circulating-co-culture-dkxv4xn6

25. Brown GC. Cell death by phagocytosis. Nat Rev Immunol. 2024 Feb;24(2):91–102.

26. Gao J, Zhang X, Jiang L, Li Y, Zheng Q. Tumor endothelial cell-derived extracellular
vesicles contribute to tumor microenvironment remodeling. Cell Communication and
Signaling. 2022 Jun 25;20(1):97.

27. Marar C, Starich B, Wirtz D. Extracellular vesicles in immunomodulation and tumor
progression. Nat Immunol. 2021 May;22(5):560–70.

28. Hoshino D, Kirkbride KC, Costello K, Clark ES, Sinha S, Grega-Larson N, et al.
Exosome Secretion Is Enhanced by Invadopodia and Drives Invasive Behavior. Cell
Reports. 2013 Dec 12;5(5):1159–68.

29. Sung BH, von Lersner A, Guerrero J, Krystofiak ES, Inman D, Pelletier R, et al. A live
cell reporter of exosome secretion and uptake reveals pathfinding behavior of migrating
cells. Nat Commun. 2020 Apr 29;11(1):2092.

30. Hoshino A, Costa-Silva B, Shen TL, Rodrigues G, Hashimoto A, Tesic Mark M, et al.
Tumour exosome integrins determine organotropic metastasis. Nature. 2015
Nov;527(7578):329–35.

31. Italiani P, Boraschi D. From Monocytes to M1/M2 Macrophages: Phenotypical vs.
Functional Differentiation. Frontiers in Immunology. 2014 Oct 17;5:514.

32. Jayasingam SD, Citartan M, Thang TH, Mat Zin AA, Ang KC, Ch'ng ES. Evaluating the
Polarization of Tumor-Associated Macrophages Into M1 and M2 Phenotypes in Human
Cancer Tissue: Technicalities and Challenges in Routine Clinical Practice. Front Oncol.
2020 Jan 24;9:1512.

33. Kartikasari AER, Huertas CS, Mitchell A, Plebanski M. Tumor-Induced Inflammatory
Cytokines and the Emerging Diagnostic Devices for Cancer Detection and Prognosis.
Front Oncol [Internet]. 2021 Jul 7 [cited 2024 Apr 30];11. Available from:
https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2021.692142/full

34. Yeh CR, Slavin S, Da J, Hsu I, Luo J, Xiao GQ, et al. Estrogen receptor α in cancer
associated fibroblasts suppresses prostate cancer invasion via reducing CCL5, IL6 and
macrophage infiltration in the tumor microenvironment. Mol Cancer. 2016 Jan 20;15:7.

35. Yunna C, Mengru H, Lei W, Weidong C. Macrophage M1/M2 polarization. European
Journal of Pharmacology. 2020 Jun 15;877:173090.

36. Anand RJ, Gribar SC, Li J, Kohler JW, Branca MF, Dubowski T, et al. Hypoxia causes
an increase in phagocytosis by macrophages in a HIF-1alpha-dependent manner. J
Leukoc Biol. 2007 Nov;82(5):1257–65.

37. Martins F, Oliveira R, Cavadas B, Pinto F, Cardoso AP, Castro F, et al. Hypoxia and
Macrophages Act in Concert Towards a Beneficial Outcome in Colon Cancer. Cancers.
2020 Mar 28;12(4):818.

38. Casali BC, Baptista MP, Pachane BC, Cortez AA, Altei WF, Selistre-de-Araújo HS.
Blockage of αvβ3 integrin in 3D culture of triple-negative breast cancer and endothelial
cells inhibits migration and discourages endothelial-to-mesenchymal plasticity.
Biochemistry and Biophysics Reports. 2024 Jul 1;38:101686.

39. Park JV, Chandra R, Cai L, Ganguly D, Li H, Toombs JE, et al. Tumor Cells Modulate
Macrophage Phenotype in a Novel In Vitro Co-Culture Model of the NSCLC Tumor
Microenvironment. J Thorac Oncol. 2022 Oct;17(10):1178–91.

40. Rebelo SP, Pinto C, Martins TR, Harrer N, Estrada MF, Loza-Alvarez P, et al. 3D-3-
culture: A tool to unveil macrophage plasticity in the tumour microenvironment.
Biomaterials. 2018 May 1;163:185–97.

41. Miranda-Azpiazu P, Panagiotou S, Jose G, Saha S. A novel dynamic multicellular co-
culture system for studying individual blood-brain barrier cell types in brain diseases and
cytotoxicity testing. Sci Rep. 2018 Jun 8;8(1):8784.

42. Weng YS, Tseng HY, Chen YA, Shen PC, Al Haq AT, Chen LM, et al. MCT-1/miR-
34a/IL-6/IL-6R signaling axis promotes EMT progression, cancer stemness and M2
macrophage polarization in triple-negative breast cancer. Molecular Cancer. 2019 Mar
18;18(1):42.

43. Kennel KB, Bozlar M, De Valk AF, Greten FR. Cancer-Associated Fibroblasts in
Inflammation and Antitumor Immunity. Clinical Cancer Research. 2023 Mar
14;29(6):1009–16.

44. Liu Y, Shi K, Chen Y, Wu X, Chen Z, Cao K, et al. Exosomes and Their Role in Cancer
Progression. Front Oncol [Internet]. 2021 Mar 22 [cited 2024 Oct 21];11. Available from:
https://www.frontiersin.org/journals/oncology/articles/10.3389/fonc.2021.639159/full

45. Becker A, Thakur BK, Weiss JM, Kim HS, Peinado H, Lyden D. Extracellular Vesicles
in Cancer: Cell-to-Cell Mediators of Metastasis. Cancer Cell. 2016 Dec 12;30(6):836–48.

46. Duong CN, Vestweber D. Mechanisms Ensuring Endothelial Junction Integrity Beyond
VE-Cadherin. Front Physiol [Internet]. 2020 May 21 [cited 2024 Jun 21];11. Available
from:
https://www.frontiersin.org/journals/physiology/articles/10.3389/fphys.2020.00519/full

47. Dejana E. Endothelial cell–cell junctions: happy together. Nat Rev Mol Cell Biol. 2004
Apr;5(4):261–70.

48. Guo M, Breslin JW, Wu MH, Gottardi CJ, Yuan SY. VE-cadherin and β-catenin binding
dynamics during histamine-induced endothelial hyperpermeability. American Journal of
Physiology-Cell Physiology. 2008 Apr;294(4):C977–84.

49. Mello AM, Ngodup T, Lee Y, Donahue KL, Li J, Rao A, et al. Hypoxia promotes an
inflammatory phenotype of fibroblasts in pancreatic cancer. Oncogenesis. 2022 Sep
15;11(1):1–9.

50. Winkler J, Abisoye-Ogunniyan A, Metcalf KJ, Werb Z. Concepts of extracellular matrix
remodelling in tumour progression and metastasis. Nat Commun. 2020 Oct 9;11(1):5120.

51. Chulpanova DS, Kitaeva KV, James V, Rizvanov AA, Solovyeva VV. Therapeutic
Prospects of Extracellular Vesicles in Cancer Treatment. Front Immunol [Internet]. 2018
Jul 3 [cited 2024 Oct 21];9. Available from:
https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2018.01534/ful
l

52. Taghizadeh-Hesary F. "Reinforcement" by Tumor Microenvironment: The Seventh "R"
of Radiobiology. International Journal of Radiation Oncology, Biology, Physics. 2024 Jul
1;119(3):727–33.

53. Van Deun J, Mestdagh P, Agostinis P, Akay Ö, Anand S, Anckaert J, et al. EV-TRACK:
transparent reporting and centralizing knowledge in extracellular vesicle research. Nat
Methods. 2017 Mar;14(3):228–32.

**A** Flow Direction (5) μl/s

Peristaltic Pump

THP-1

MDA MB 231  HUVEC  HDFa
Gelatin coating  Matrigel coating  Fibronectin coating

**B**                                    MC-CC chambers

MDA-MB-231          HUVEC          HDFa

PBS

*Nuclei  Gelatin  F-actin*   *Nuclei  VEGFR2  β-catenin*   *Nuclei  Col II  Col III*

EVh

**C** Gelatin Invasion
MDA-MB-231 on MC-CC

Degraded area (μm²)
<0.0001
<0.0001
0.0014

Static  50 μl  PBS  EVh
Static  |  MC-CC

**D** MMP-2 Activity

Relative Band Density (AU)
ns

PBS  EVh

**E** IL-1β
MC-CC

IL-1β (pg/mL)
ns

PBS  EVh

**F** IL-6
MC-CC

IL-6 (pg/mL)
0.0431

PBS  EVh

**G** β-catenin
HUVEC on MC-CC

Relative Fluorescence
Intensity (AU)
<0.0001

PBS  EVh

**H** VEGFR2
HUVEC on MC-CC

Relative Fluorescence
Intensity (AU)
0.0059

PBS  EVh

**I** Col II
HDFa on MC-CC

Relative Fluorescence
Intensity (AU)
<0.0001

PBS  EVh

**J** Col III
HDFa on MC-CC

Relative Fluorescence
Intensity (AU)
<0.0001

PBS  EVh

**A** THP-1 in MC-CC

**B** THP-1 in Direct Co-culture

MDA-MB-231     HUVEC     HDFa

PBS

EVh

*Nuclei*   *F-actin*   *CellTracker CMPTX*

*Nuclei*   *F-actin*   *CellTrace CSFE*

**C** **Cell Morphology**
<u>THP-1 Direct Co-Culture</u>

Circularity Index

*ns*    *ns*    0.0295

PBS EVh   PBS EVh   PBS EVh
MDA-MB-231   HUVEC   HDFa

**D** **Gelatin Invasion**
<u>THP-1 Direct Co-Culture</u>

Gelatin Degradation (μm²)

*<0.0001*   *<0.0001*   *<0.0001*

PBS EVh   PBS EVh   PBS EVh
MDA-MB-231   HUVEC   HDFa