Title: Integrative analysis of CAM photosynthesis reveals its impact on primary metabolism in *Yucca*

Short title: Impacts of CAM on primary metabolism in Yucca

David Wickell[1*], Richard Field[2,3*], Karl Weitz[4], Rosalie Chu[4], Jesse Trejo[5], Nikola Tolic[4], Nathalie Munoz Munoz[4], Hope Hundley[6], Emily Savage[6], Anna Lipzen[6], Jim Leebens-Mack[2], Karolina Heyduk[1]

[1] Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, 06269
[2] Department of Plant Biology, University of Georgia, Athens, GA, 30602
[3] USDA-ARS, Athens, GA, 30605
[4] Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, WA, 99354
[5] Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, 99354
[6] US Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720
*D.W. and R.C. contributed equally to this work

Corresponding author: David Wickell (david.wickell@uconn.edu)

Abstract:

Crassulacean Acid Metabolism (CAM) is an adaptation that temporally separates carbon uptake at night from photosynthesis during the day. CAM has evolved repeatedly across vascular plants, suggesting its emergence may depend on relatively simple changes to deeply conserved regulatory mechanisms. Modern CAM research relies heavily on the interpretation of transcriptomic data. This approach may bias mechanistic explanations of CAM toward transcriptional regulation, though regulation is known to occur at multiple levels following transcription. Additionally, while the majority of research to date has focused on a handful of genes and metabolites in the core CAM pathway, the co-option of conserved regulatory and functional genes is bound to have wide ranging effects on other aspects of primary metabolism. Thus, taking a broad and integrative approach to understanding the downstream effects associated with the emergence of CAM is essential to explaining its convergent evolution as well as to efforts to engineer carbon concentrating mechanisms in crops. In this study, we integrate genomic, transcriptomic, proteomic, and metabolomic data to compare primary metabolism between the CAM species *Yucca aloifolia* and closely related C_3 species, *Y. filamentosa*. We observe minimal correlation between protein abundance and mRNA expression, suggesting significant post-transcriptional regulation in CAM species. We also find evidence of shifts in gene expression and metabolite accumulation outside of the central CAM pathway suggesting that the shift to CAM has cascading effects across primary metabolism, especially nitrogen metabolism. Our findings provide insights into the metabolic shifts associated with CAM evolution and highlight the complexity of its regulation at multiple biological levels.

Introduction

Crassulacean Acid Metabolism (CAM) is a variation on the more common C_3 metabolism that allows plants to temporally separate carbon uptake and storage from photosynthesis. Briefly, in C_3 plants, uptake of atmospheric CO_2 and carbon fixation by Rubisco co-occur during the day. In CAM plants, however, CO_2 is taken up at night and stored in the vacuole as malate prior to being liberated and fed into the Calvin cycle during the day (Fig. 1). CAM photosynthesis allows terrestrial plants to keep their stomata closed during the hottest part of the day, greatly reducing transpirational water loss. This, in combination with reduced rates of photorespiration due to the enrichment of CO_2 in photosynthetic tissues during the day, results in

dramatic improvement to water use efficiency in CAM lineages relative to C_3 plants (Winter &

Smith, 2022).

CAM is distinct from other carbon concentrating mechanisms in that it temporally

separates CO_2 uptake from Rubisco-mediated fixation, whereas C_4 photosynthesis and

pyrenoids achieve this separation spatially, relying on specialized anatomy (Sage et al., 2012) or

subcellular compartments (Robison et al., 2024), respectively. Accordingly, CAM physiology

appears to arise from regulatory modifications to deeply conserved pathways common to all

green photosynthetic organisms. Still, there is some debate as to whether the emergence of CAM

requires complex regulatory rewiring across the genome (Wai et al., 2019), if it is the product of

minimal changes resulting in increased metabolic flux through existing C_3 pathways (Bräutigam

et al., 2017), or some combination of the two. Despite the potential complexity of such

fundamental alterations to primary metabolism, CAM has evolved over 60 times across the plant

phylogeny (Gilman et al., 2023). It is found in every major lineage of vascular plants, in diverse

taxa occupying an equally diverse range of habitats from deserts to lakes to rainforest canopies.

While the central components of the CAM pathway are present in highly divergent taxa, studies

have found a great deal of lineage specific variation as well. CAM plants can vary in their

mechanism of day-time decarboxylation (Ramachandra Reddy & Rama Das, 1978), gene copy

recruitment (Heyduk et al., 2022; Wickell et al., 2021), the role of circadian regulation (L.-Y.

Chen et al., 2020; Wai et al., 2019; Wickell et al., 2021), and citrate accumulation (Franco et al.,

1992; Freschi et al., 2010; Lüttge, 1988). This high degree of interspecific variation in the details

of CAM makes comparison of closely related CAM and C_3 species essential to our

understanding of a complex syndrome that has evolved repeatedly in a diverse array of genomic

backgrounds.

Figure 1: Flow chart of the core CAM pathway (black arrows) and its interactions with other primary metabolic pathways (gray arrows). Alternatively, malate can be decarboxylated by MDH and PEPCK during the day (not shown) and conversion of citrate to 2-OG may occur in the cytosol (as shown) or mitochondrion (not shown). Abbreviations: 2-OG, 2-oxoglutarate; MDH, malate dehydrogenase; ME, malic enzyme; OAA, oxaloacetate; PEP, phosphoenolpyruvate; PEPC, phosphoenolpyruvate carboxylase; PPDK, pyruvate-phosphate dikinase

To date, the majority of genome-enabled research on CAM species has focused on the central carboxylation mechanisms that fix CO_2 into malate for storage at night and decarboxylation processes that release CO_2 during the day (e.g. (Abraham et al., 2016; Ming et al., 2015; Wickell et al., 2021). These core CAM pathways involve genes, proteins, and metabolites that are also central components of primary metabolism in C_3 plants where they replenish intermediates in the citric acid cycle. As such, alterations to their expression and

abundance are expected to have complex downstream effects on pathways outside of CAM. A

notable example of the effects of CAM outside of the canonical pathway is the night time

accumulation of citrate in many CAM plants, often exhibiting a pattern similar to malate (Franco

et al., 1992; Freschi et al., 2010; Kornas et al., 2009; Miszalski et al., 2013). Authors have

suggested multiple potential explanations that integrate citrate into the existing CAM framework

including recapture of respired CO_2 (Franco et al., 1992), control of redox potential (Kornas et

al., 2009), and mitigation of photoinhibition and photorespiration via efficient concentration of

CO_2 around Rubisco (Kornas et al., 2009; Miszalski et al., 2013). However, unlike malate, citrate

synthesis is not associated with a net carbon gain (Lüttge, 1988). Thus, despite its frequent

association with CAM, the precise role of nocturnal citrate accumulation and how frequently it

occurs across CAM lineages remain understudied and ultimately unresolved.

Organic acid accumulation, including malate and citrate, has been previously shown to be

strongly impacted by both the relative abundance and source of nitrogen and several studies have

sought to characterize a mechanistic relationship between CAM and nitrogen metabolism

(Gonçalves & Mercier, 2021; Pereira et al., 2018; Pereira & Cushman, 2019; Santos & Salema,

1991). Specifically, researchers have found that some species including *Guzmania monostachia*

and several *Kalanchoë* species exhibit increased CAM expression and nocturnal acidification

under nitrogen deficiency, further underscoring connections between carbon and nitrogen

metabolism (Nobel, 1983; Ota, 1988; Rodrigues et al., 2014; Winter et al., 1982). However, few

studies have incorporated gene expression data and detailed comparisons of N-metabolism

between closely related CAM and C_3 lineages are rare. Similarly, few papers have investigated

the photorespiratory pathway in CAM lineages (Duarte & Lüttge, 2007; Lüttge, 2011) despite it

being widely accepted that the reduction of photorespiration is one of the primary factors driving

the convergent evolution of CAM in vascular plants.

The Agavoideae has emerged as an important model group for studying the evolutionary

origins of CAM due to its high degree of photosynthetic diversity, containing C_3, CAM, and

physiologically intermediate "C_3 + CAM" species (Heyduk et al., 2023). In particular, the genus

Yucca within the Agavoideae is well suited to comparative analyses of CAM evolution as it

comprises both C_3 and CAM species with a single origin of CAM inferred roughly 6 mya in

section Yucca (Heyduk, McKain, et al., 2016; McKain et al., 2016). The presence of closely

related species that differ in photosynthetic pathway allows researchers to investigate CAM

through comparative analyses while controlling for the confounding effects of shared ancestry.

As a result, considerable research has been done in *Yucca* investigating the genomic (Heyduk et

al., 2019, 2022), physiological (Heyduk, Burrell, et al., 2016; Heyduk et al., 2020), and

anatomical (Heyduk et al., 2020; Heyduk, McKain, et al., 2016) changes associated with the

transition to CAM. A recent study found evidence of shared expression patterns of core CAM

genes between closely related C_3 and CAM species (Heyduk et al., 2019). Similarly, ancestral

state reconstruction in Agavoideae indicated that key features of CAM anatomy such as leaf

thickness, cell succulence, and 3D venation may have evolved prior to strong CAM (Heyduk,

McKain, et al., 2016). These studies seem to corroborate work suggesting that the evolution of

CAM may merely require an increase in flux through existing carboxylation pathways

(Bräutigam et al., 2017). Even so, Heyduk et al. (2019) also found widespread differences in

gene expression within and outside of the central CAM pathway alongside differential

abundance of several primary metabolites between CAM and C_3 species, indicative of potentially

complex downstream effects of CAM across primary metabolism.

Characterizing the wide ranging effects of CAM on primary metabolism requires

simultaneous comparative analysis of multiple pathways and molecules. While this would ideally

integrate functional genetic analysis, these types of studies are still in their infancy in CAM

plants; *Kalanchoë* is the only CAM lineage where knockdown studies have confirmed genetic

function (Agarie et al., 2020; Boxall et al., 2020; Dever et al., 2015; Ferrari et al., 2020). As

such, comparative multi-omics approaches potentially offer a more accessible way to investigate

modification to primary metabolism in a greater diversity of CAM species. In this study we

integrated chromosome level genome assemblies with transcriptomic, proteomic, and

metabolomic time series data for *Yucca aloifolia* (CAM) and *Y. filamentosa* (C_3) to investigate

shifts in primary metabolism associated with the transition from C_3 to CAM. Comparison of

transcriptomic and proteomic datasets showed little correlation between fluctuations in transcript

and protein abundance over time, suggesting that post-translational modification and

activation/inhibition by other proteins and metabolites play a significant role in regulation of

CAM following translation of mRNA into proteins. We also found alterations to protein

abundance, gene expression, and metabolite abundance related to citrate metabolism and

nitrogen metabolism between *Y. aloifolia* (CAM) and *Y. filamentosa* (C_3). However, despite

interspecies differences in the abundance of metabolites associated with photorespiration, there

were few clear differences in protein abundance or gene expression between the two species.

This may suggest a diminished role for photorespiration in the evolution of CAM and support the

theory that the evolution of CAM may be driven by increased flux through existing C_3 pathways

as opposed to extensive regulatory rewiring (Bräutigam et al., 2017). Our findings underscore the

utility of comprehensive multi-omics analyses in unraveling the complex metabolic shifts

associated with CAM, providing valuable insights into gene regulation, and generating testable

hypotheses that pave the way for further research of CAM evolution and its underlying

mechanisms across vascular plants.

Results

Broad comparison of -omics datasets

172   Gas exchange data (Supplementary Fig. S1) and malate accumulation differences

between species support prior evidence that showed *Y. aloifolia* and *Y. filamentosa* are CAM and

C_3, respectively. Data also corroborated prior work that demonstrated *Y. aloifolia*'s reliance on

CAM slightly decreases under drought stress (Heyduk et al., 2019). Transcriptomic analysis of *Y.*

aloifolia and *Y. filamentosa* found that 83.4% (34,663/41,577) of *Y. aloifolia* genes and 50.4%

(22,195/43,996) of *Y. filamentosa* genes exhibited time-structured mRNA expression in well-

watered and/or drought-stressed treatments (Fig. 2, Supplementary Tables 2 and 3). When

analysis was restricted to genes that cycle in both treatments, 54% (22,523/41,577) and 28%

(12,357/43,996) of genes cycled in *Y. aloifolia* and *Y. filamentosa*, respectively (Fig. 2,

Supplementary Fig. S2, Supplementary Tables 2 and 3).

182   Proteomic analysis of leaf tissue recovered abundance data for far fewer genes than

transcriptomic analysis. Abundance was recorded for 6,354 proteins in *Y. aloifolia* and 6,859

proteins in *Y. filamentosa*, representing 15.2% and 15.6% of annotated genes, respectively. Also

in contrast to analyses of transcriptomic data, relatively few proteins exhibited time-structured

abundance patterns (Fig. 2, Supplementary Fig. S2, Supplementary Tables 4 and 5). In proteins,

linear regression identified 50.5% (3,214/6,354) of proteins in *Y. aloifolia* and only 35%

(2,400/6,859) of proteins in *Y. filamentosa* as cycling over time in at least one treatment. When

the analysis was restricted to genes that cycle in both treatments, far fewer proteins were

identified with 9.8% (622/6,354) and 4.5% (310/6,859) proteins cycling in *Y. aloifolia* and *Y.*

filamentosa, respectively (Fig. 2, Supplementary Fig. S2, Supplementary Tables 4 and 5).

Transcription factors (TFs) were notably scarce in our proteomic dataset with only 256 out of

5,636 annotated TFs observed in *Y. aloifolia* and 349 out of 5,521 annotated TFs observed in *Y.*

filamentosa.

Untargeted metabolomic analysis identified 72 metabolites in *Y. aloifolia* and 81

metabolites in *Y. filamentosa* of which 48 were found in both species (Supplementary Table 6).

In *Y. aloifolia* 21/72 metabolites showed time-structured abundance over the day-night period in

at least one treatment compared to 10/81 in *Y. filamentosa* (Supplementary Table 6). Of the

metabolites identified in both species, just 5/48 metabolites cycled in at least one treatment in

both species (Supplementary Table 6).

Figure 2. Overlap in temporal patterns of protein abundance and mRNA expression in

Yucca. Barplots show the total number of genes that exhibit cycling expression and or protein

abundance for *Y. aloifolia* (YA) and *Y. filamentosa* (YF). Proteins with abundance peaks

shifted ≥4 h relative to its associated mRNA transcript are designated as exhibiting

shifted peaks.

The CAM pathway

Of CAM associated proteins (Fig. 3a), relatively few exhibited cycling abundance. While

no copies of β-CA cycled in *Y. aloifolia,* two copies cycled in *Y. filamentosa*, one of which had

no clear ortholog in *Y. aloifolia.* (Fig. 3b, Supplementary Fig S3). Notably, *Y. aloifolia* exhibits

cycling mRNA expression of two distinct forms of PEPC (Heyduk et al., 2022) commonly

referred to as plant-type PEPC (PTPC), and bacterial-type PEPC (BTPC; Supplementary Fig.

S3). Two copies of PTPC cycled in each *Y. aloifolia* and *Y. filamentosa* with one cycling in both.

Curiously, the copy in *Y. aloifolia* corresponding to the most highly expressed mRNA transcript

– typically associated with CAM – did not cycle in protein abundance, while its ortholog in *Y.*

filamentosa did. In the CAM pathway, only BTPC abundance both cycled and appeared to track

mRNA expression under drought-stressed and well-watered conditions in *Y. aloifolia* where two

copies showed similar cycling abundance (Fig 3b, Supplementary Fig. S3). BTPC abundance did

not cycle in *Y. filamentosa*. PPCK, required for activation of PEPC, was notably absent from

the proteomic data in both species, potentially due to its low abundance (Hartwell et al., 1999).

We found 2 copies of cytoplasmic NAD-MDH that cycled in abundance in *Y. filamentosa* but

none in *Y. aloifolia* (Fig 3b, Supplementary Fig. S3). In the light reactions, CAM associated

copies of NADP-ME and PPDK cycled in protein abundance in *Y. aloifolia*, though peak

expression was shifted 4 to 8 hours later in both relative to mRNA expression (Fig. 3b). The

same copy of NADP-ME also cycled in *Y. filamentosa* with peak abundance just before dusk,

despite the fact that its transcript did not cycle. A second paralog of NADP-ME cycled in *Y.*

aloifolia with peak protein abundance during the dark period despite its hypothesized

involvement in the daytime decarboxylation reactions in CAM (Supplementary Fig. S3).

In contrast to protein abundance, the majority of CAM associated mRNA transcripts

exhibited cycling expression in *Y. aloifolia* (33/38) relative to *Y. filamentosa* (18/36;

Supplementary Table 6). While at least one copy, and occasionally multiple copies, of canonical

CAM genes cycled in *Y. aloifolia*, their temporal expression patterns varied considerably (Fig.

3b, Supplementary Fig. S3) . In some cases this was expected, as in the partitioning of genes

involved in nighttime carboxylation from those involved in daytime decarboxylation. Similarly,

this was true of PEPC and its activator PPCK in the CAM carboxylation pathway which

frequently exhibit distinct expression patterns in other CAM species (Heyduk et al., 2019; Ming

et al., 2015; Wai et al., 2019; Wickell et al., 2021; Yang et al., 2017). As expected, both BTPC

and PTPC exhibited similar expression patterns in *Y. aloifolia* with peak expression just before

dusk (Fig. 3b). In contrast to PEPC and PPCK, cytosolic copies of NAD-MDH in *Y. aloifolia*

showed distinct expression patterns between well-watered and drought-stressed conditions

peaking at dawn and during the middle of the dark period, respectively (Fig. 3b, Supplementary

Fig. S3). While some CAM species rely on NAD-MDH to convert malate to OAA and PEPCK

for subsequent daytime decarboxylation, transcriptomic data suggests that *Y. aloifolia* appears to

rely on NADP-ME for decarboxylation followed by PPDK for regeneration of PEP, based on

their higher, cycling expression relative to NAD-MDH and PEPCK. However, similar to many

nighttime carboxylation genes, NADP-ME and PPDK generally exhibited distinct expression

patterns from one another in our MaSigPro analysis with peak expression occurring near dawn in

PPDK and shifted 4 to 8 hours later in NADP-ME (Fig. 3b, Supplementary Table 6).

In *Y. filamentosa*, multiple copies of *PEPC* cycled in mRNA expression, including one

paralog which resembled the most highly expressed copy in *Y. aloifolia*, peaking near midday

(Supplementary Fig. S2) but we found no correlation in expression patterns between *BTPC* and

PTPC and overall expression was much lower in *Y. filamentosa* (Fig. 3b). A single copy of

PPCK in *Y. filamentosa* shows similar, cycling expression to the most highly expressed copy of

PPCK in *Y. aloifolia,* though expression is roughly 10x lower in *Y. filamentosa* (Fig. 3b). *NAD-*

MDH expression was also generally ~5-10x lower in *Y. filamentosa* than in *Y. aloifolia* and only

a single copy appeared to cycle (Fig. 3b, Supplementary Table 6). Similarly, most copies of

NADP-ME and *PPDK* in *Y. filamentosa* were lowly expressed and did not cycle (Supplementary

Fig. S2, Supplementary Table 6).

Malate was the only core CAM metabolite present in our metabolomic dataset. As

expected, malate cycled strongly in *Y. aloifolia* relative to *Y. filamentosa*, accumulating

throughout the night and decreasing in abundance during the day (Fig. 3c). Despite some

similarity between species in expression of CAM genes such as *PPCK* and at least one copy of

PTPC, no nighttime accumulation of malate was observed in *Y. filamentosa*. This result was

consistent with measurements of stomatal conductance in *Y. aloifolia* relative to *Y. filamentosa*

(Supplementary Fig. S1). In *Y. aloifolia*, but not *Y. filamentosa*, phosphate abundance also cycled

strongly and tracked malate abundance, increasing at night and decreasing during the day (Fig.

3c). Cycling of phosphate in *Y. aloifolia* likely correlates with its release during the

carboxylation of PEP to OAA at night, its essential role in activation of PEPC, and high demand

during the day for regenerating compounds such as ATP and NADP.

Figure 3: Comparison of CAM photosynthesis in *Y. aloifolia* and *Y. filamentosa*. (a.) Flow-

chart of core reactions of the CAM pathway in *Y.aloifolia*. **(b.)** Plots of z-scaled mean protein

abundance (solid line, SD represented by shaded area) with mean expression in TPM overlaid

(dashed line) for comparison. Filled and unfilled circles indicate significantly ($p < 0.05$) time-

structured protein abundance and gene expression, respectively. **(c.)** Plots of abundance over

time for malate and phosphate, (indicated in flow-chart by an '*') shaded areas represent the

standard deviation. Filled circles indicate significantly ($p < 0.05$) time-structured metabolite

abundance and daggers (†) indicate metabolites that are significantly ($p < 0.05$) different between

species. Abbreviations: b-CA, beta-carbonic anhydrase; PEP, phosphoenolpyruvate; PPCK,

phosphoenolpyruvate carboxylase kinase; PEPC, phosphoenolpyruvate carboxylase; PTPC,

plant-type phosphoenolpyruvate carboxylase; BTPC, bacterial-type phosphoenolpyruvate

carboxylase; OAA, oxaloacetate; NAD-MDH, NAD dependent malate dehydrogenase; NADP-

ME, NADP dependent malic enzyme; PPDK, pyruvate phosphate dikinase.

Citrate accumulation

288   Citrate exhibited similar expression to malate in *Y. aloifolia*, as previously reported in

several other CAM taxa (Borland & Griffiths, 1989; Franco et al., 1992; Freschi et al., 2010)

leading us to investigate genes associated with citrate metabolism in both species (Fig. 4). In

plants, citrate is typically produced by citrate synthase (CS) in the peroxisome as part of the

glyoxylate cycle and in the mitochondria, as part of the tricarboxylic acid cycle (TCA) cycle. In

the mitochondria, citric acid is subsequently converted to isocitrate and isocitrate to 2-

oxoglutarate (2-OG) by aconitase (ACO) and NAD-dependent isocitrate dehydrogenase (NAD-

IDH), respectively (Fig. 4a). Abundance of citrate in *Y. aloifolia* increased throughout the night

and decreased during the day, compared to *Y. filamentosa* where citrate levels were relatively

stable between day and night (Fig. 4c). We also observed differences between species in genes

related to citrate metabolism.

299   As in core CAM pathway genes, relatively few proteins associated with citrate

production cycled in abundance in either *Y. aloifolia* or *Y. filamentosa* and even fewer were

associated with transcripts that also cycled in expression. A single copy of mitochondrial CS

protein cycled in *Y. aloifolia*, but did not correspond to the most highly expressed cycling

transcript in that species (Fig 4b, Supplementary Fig. S4). One mitochondrial and one

peroxisomal copy of CS protein appeared to cycle in *Y. filamentosa*. However both exhibited

shifts in their abundance profiles relative to transcript expression, particularly under drought-

stress. ACO and IDH followed a similar trend, in that the most highly expressed cycling

transcripts were not associated with cycling protein abundance, though several copies exhibited

cycling protein abundance under at least one treatment (Fig 4b, Supplementary Fig. S4,

Supplementary Table 6).

In contrast to our proteomic data, 35 out of 40 TCA cycle genes cycled in transcript

expression in *Y. aloifolia* and exhibited a diverse array of expression patterns, while the

relatively few transcripts that cycled in *Y. filamentosa* showed highly similar expression patterns

in our MaSigPro analysis (cluster 1, Supplementary Fig. S4, Supplementary Table 6). Notably, in

Y. aloifolia, one copy of *CS*, two copies of *ACO*, one copy of *NAD-IDH*, and one copy of *NADP-*

IDH all had similar cycling expression that tracked temporal patterns of citrate abundance with

peak expression during the dark period (Fig. 4b, Supplementary Fig. S4). The most highly

expressed copy of mitochondrial *CS* cycled in *Y. aloifolia* but not in *Y. filamentosa*, exhibiting

peak expression at dawn. Two copies of *ACO* exhibited strong cycling expression in *Y. aloifolia*,

with expression decreasing during the morning before increasing and remaining high throughout

the night (Fig. 4b, Supplementary Fig. S4). Expression of *ACO* in *Y. filamentosa* did not have a

clear temporal pattern and was 3-4 times lower than in *Y. aloifolia*. Differences in expression

were further observed in both NAD-dependent (mitochondrial) and NADP-dependent (cytosolic)

copies of *IDH*. *NAD-IDH* expression was relatively stable over time in both species and

treatments but roughly twice as high in *Y. aloifolia* (Supplementary Fig. S4). Cytosolic *NADP-*

IDH exhibited peak expression in both species near dusk but remained high throughout the night

in *Y. aloifolia*, strongly resembling the expression of *ACO* (Fig. 4b).

Figure 4: Comparison of nitrogen metabolism in *Y. aloifolia* and *Y. filamentosa*. (a.) Flow-chart of core reactions of the nitrogen metabolism pathway and its interactions with CAM, photorespiration, and the citric acid cycle. Conversion of citrate to 2-OG may occur in the cytosol (not shown) or mitochondrion (as shown). **(b.)** Plots of z-scaled mean protein abundance (solid line, SD represented by shaded area) with mean expression in TPM overlaid (dashed line) for comparison. Filled and unfilled circles indicate significantly ($p < 0.05$) time-structured protein abundance and gene expression, respectively. **(c.)** Plots of abundance over time for glutamate, aspartate, and citrate (indicated in flow-chart by an '*') shaded areas represent the standard deviation. Filled circles indicate significantly ($p < 0.05$) time-structured metabolite abundance and daggers (†) indicate metabolites that are significantly ($p < 0.05$) different between species. Abbreviations: 2-OG, 2-oxoglutarate; AAT, aspartate aminotransferase; ACO, aconitase; CS, citrate synthase; GDH, glutamate dehydrogenase; GOGAT, glutamine oxoglutarate aminotransferase; GS, glutamine synthetase; IDH, isocitrate dehydrogenase;

NADP-IDH, NADP-dependent isocitrate dehydrogenase; NiR, nitrite reductase; NR, nitrate

reductase; OAA, oxaloacetate.

Nitrogen metabolism

Outside of the CAM pathway, our multi-omics analysis allowed us to explore differences

in protein abundance and gene expression associated with nitrogen metabolism (Fig. 4). Multiple

proteins were found to cycle in abundance in *Y. aloifolia* and to a lesser extent in *Y. filamentosa*.

Neither nitrate reductase (NR) nor nitrite reductase (NiR) cycled in abundance in *Y. aloifolia*,

though the single-copy NiR did cycle in *Y. filamentosa*, peaking in abundance at midday (Fig.

4b). In the chloroplast, multiple copies of glutamine synthetase (GS) exhibited cycling protein

abundance in both species, while ferredoxin-dependent glutamate synthase (Fd-GOGAT) and

aspartate aminotransferase (AAT) each had multiple paralogs that cycled in *Y. aloifolia* alone

(Fig. 4b, Supplementary Fig. S5, Supplementary Table 6). A single copy of glutamate

dehydrogenase (GDH) cycled in protein abundance in both species but showed inverse patterns

of protein abundance, peaking at dawn in *Y. aloifolia* and just before dusk in *Y. filamentosa* (Fig.

4b). Notably, one copy of GS cycled in both species with similar abundance peaking during the

dark period in *Y. aloifolia* in well-watered and drought-stressed conditions as well as *Y.*

filamentosa under drought (Fig. 4b). In addition, another copy of GS cycled in abundance in *Y.*

aloifolia under well-watered conditions and two copies cycled in *Y. filamentosa* under well-

watered and drought conditions (Supplementary Fig. S5, Supplementary Table 6). Both copies of

Fd-GOGAT exhibited cycling abundance in *Y. aloifolia* (Fig. 4b, Supplementary Fig. S5,

Supplementary Table 6) with higher abundance during the dark period, but not in *Y. filamentosa*.

Similarly, 4 out of 5 AAT proteins cycled in *Y. aloifolia* with peak expression near dusk (Fig. 4b,

Supplementary Fig. S4, Supplementary Table 6). None of the copies of AAT present in *Y.*

filamentosa displayed time structured protein abundance.

Several nitrogen metabolism genes showed clear differences in transcript expression

between species and treatments, even in cases where their corresponding proteins did not (Fig.

4b, Supplementary Fig. S4, Supplementary Table 6). For example, *NR* and *NiR*, which catalyze

the initial reduction of nitrate to ammonia, exhibited cycling expression in *Y. aloifolia*, peaking

near dawn under both drought stress and well-watered conditions (Fig. 4b). In *Y. filamentosa*, a

similar pattern was observed under drought stress, but expression remained relatively stable in

well-watered plants (Fig. 4b). *GDH*, which catalyzes the reversible conversion of glutamate to 2-

OG and ammonia in the mitochondria, also showed distinct temporal expression, peaking near

dawn in *Y. aloifolia* and around midday in *Y. filamentosa* in both treatments (Fig. 4b). Despite

the difference in gene expression between species, this was the only clear case where the most

highly expressed gene cycled in both transcript and protein abundance in both species. However,

the peak in gene expression was shifted 8 hours earlier than corresponding protein abundance in

Y. aloifolia and similarly under well-watered conditions in *Y. filamentosa* (Fig. 4b). In contrast to

NR and *NiR*, other chloroplast-localized genes in the nitrogen metabolism pathway, including

GS, *Fd-GOGAT*, and *AAT*, showed little variation in timing of peak expression across species.

In addition, two copies of *AAT* and *Fd-GOGAT* cycled in *Y. filamentosa* transcripts despite none

of the proteins cycling in abundance (Fig. 4b, Supplementary Fig. S5, Supplementary Table S6).

Expression levels of *GS* and *Fd-GOGAT* were generally lower in *Y. aloifolia*, especially under

well-watered conditions, while *AAT* expression was nearly four times higher in *Y. aloifolia*

across both treatments (Fig. 4b).

Two primary products of nitrogen metabolism, aspartate and glutamate, were identified

by metabolomic analysis (Fig 4c). The production of aspartate and glutamate by AAT and Fd-

GOGAT, respectively, plays a central role in primary nitrogen metabolism and uses OAA, a

primary product of CAM dark reactions and TCA cycle intermediate, as a substrate. Despite the

similarity in gene expression of *AAT* and *Fd-GOGAT* between CAM and C_3 species, their

immediate products appeared to cycle in *Y. aloifolia* but not in *Y. filamentosa*. *GDH*, which

catalyzes the reversible formation of glutamate from 2-OG and ammonia in mitochondria, is

differentially expressed between the two species. In *Y. aloifolia*, aspartate and glutamate

exhibited inverse abundance with one another, with aspartate peaking during the dark period and

glutamate peaking during the light period (Fig. 4c).

Figure 5: Comparison of photorespiration in *Y. aloifolia* and *Y. filamentosa*. (a.) Flow-chart of core reactions of the photorespiratory pathway. **(b.)** Plots of z-scaled mean protein abundance (solid line, SD represented by shaded area) with mean expression in TPM overlaid (dashed line) for comparison. Filled and unfilled circles indicate significantly ($p < 0.05$) time-structured protein abundance and gene expression, respectively. **(c.)** Plots of abundance over time for serine and glycerate (indicated in flow-chart by an '*') shaded areas represent the standard deviation. Filled circles indicate significantly ($p < 0.05$) time-structured metabolite abundance and daggers (†) indicate metabolites that are significantly ($p < 0.05$) different between species. Abbreviations: GDH/GDC-H, glycine decarboxylase complex - subunit H; GGAT, glutamate:glyoxylate aminotransferase; GLYK, glycerate kinase; GOX, glycolate oxidase; HPR, hydroxypyruvate

reductase; PGP, phosphoglycolate phosphatase; Rubisco, ribulose bisphosphate carboxylase;

SGAT, serine:glyoxylate aminotransferase; SHMT, serine hydroxymethyl transferase.

Photorespiration

Very few genes in the photorespiratory pathway (Fig. 5a) displayed cycling protein

abundance in either species of *Yucca*. Out of 8 core genes involved in photorespiration only

mitochondrial serine hydroxymethyl transferase (SHMT) and glycine decarboxylase complex -

subunit H (GDC-H) cycled in protein abundance in both species (Fig. 5b). SHMT and the GDC

enzyme complex facilitate the conversion of glycine to serine in the mitochondria as part of the

photorespiratory pathway (Fig. 5a). Specifically, as a component of the GDC enzyme complex,

GDC-H plays an essential role in the decarboxylation of glycine and subsequent release of

photorespiratory CO_2 and ammonia (Timm et al., 2012).

Overall expression of genes in the photorespiratory pathway was relatively similar

between *Y. aloifolia* and *Y. filamentosa* with many genes having peak expression near dawn or

during the dark period (Fig. 5b). A notable exception is phosphoglycolate phosphatase (*PGP*),

the enzyme that catalyzes the first step of the photorespiratory pathway, converting 2-

phosphoglycolate to glycolate in the chloroplast. Both species possess two copies of *PGP*, and

both copies exhibit similar expression under drought-stressed and well-watered treatments in *Y.*

filamentosa, with elevated expression throughout the daylight hours and reduced expression at

night (Fig 5b., Supplementary Fig. S6). This pattern is consistent with a presumed response to

elevated photorespiratory stress during the day. Conversely, in *Y. aloifolia*, a single copy of *PGP*

is expressed. Under well watered conditions, *PGP* exhibits peak expression at midday but

increases again to similar levels during the dark period. Under drought, expression peaks at dawn

in *Y. aloifolia*, steadily decreasing throughout the day before increasing again during the dark

period (Fig. 5b). In addition to the relatively similar expression between species, cycling genes in

the photorespiratory pathway were consistently assigned to the same MaSigPro clusters within

each species. 15 out of 24 cycling transcripts including 6 out of 8 of the most highly expressed

copies in *Y.aloifolia* were assigned to a single cluster with peak expression at the end of the night

period (cluster 1, Supplementary Fig. S6, Supplementary Table 6). In *Y. filamentosa*, a similar

pattern was observed with 10 out of 20 cycling transcripts assigned to a single cluster with peak

expression during the dark period (cluster 6, Supplementary Fig. S6, Supplementary Table 6) and

5 of 20 were assigned to another cluster with peak expression at midday (cluster 3,

Supplementary Fig. S6, Supplementary Table 6). There were no genes assigned to a

corresponding midday cluster in *Y. aloifolia*. *SHMT* expression was uniquely diverse among

photorespiratory genes. Nearly every transcript (6 out of 8 in *Y. aloifolia*, 4 out of 5 in *Y.*

filamentosa) that deviated from the dominant expression patterns in either species were paralogs

of *SHMT* (Supplementary Fig S6, Supplementary Table 6).

Y. aloifolia and *Y. filamentosa* exhibited distinct metabolite abundance profiles for serine

and glycerate. Serine, glycerate, and glycolate are all produced in plants as intermediates in the

photorespiratory pathway (Fig. 5a). Serine exhibited elevated abundance during the dark period

in drought-stressed and well-watered conditions in *Y. aloifolia,* though the change in abundance

was much stronger under drought stress (Fig. 5c). Glycerate and glycolate showed relatively

steady abundance throughout the day and night under well-watered conditions in *Y. aloifolia* but

under drought, both decreased in abundance throughout the day and increased during the dark

period. In contrast, *Y. filamentosa* peaked in glycerate abundance during daylight hours, as

expected given that photorespiration is a light dependent reaction. Glycolate was not detected in

Y. filamentosa.

Discussion

Through the use of integrated -omics data, this study takes advantage of two closely-

related but photosynthetically distinct species of *Yucca* to understand the regulation of CAM

photosynthesis at the cellular level. As expected based on prior work, *Y. aloifolia* has nocturnal

CO_2 assimilation (Supplementary Fig. S1) (though, notably, daytime CO_2 assimilation is not

zero) and a significant accumulation of malate over the course of the night (Fig. 3c). In contrast,

Y. filamentosa has no ability to assimilate CO_2 at night, nor does it have a significant

accumulation of malate under either treatment condition. The different photosynthetic pathway

used by these two *Yucca* species as close evolutionary comparisons not only in transcription—

which has been studied in these taxa among other CAM species—but also metabolomics and

proteomics. Our results highlight core differences in nitrogen metabolism and photorespiration

between the CAM and C_3 species, and also suggest relatively low dynamic changes to protein

abundance over the course of the diel cycle in contrast to transcript abundance.

Protein abundance shows reduced temporal expression relative to mRNA

Relatively few proteins cycled in abundance and protein abundance rarely cycled along

with their associated transcripts, or in cases where they did, they frequently exhibited shifts in

peak abundance relative to mRNA expression (Fig. 3). The lack of correlation between proteins

and their transcripts was particularly apparent in the core CAM pathway, to the extent that it

might be difficult to distinguish between CAM and C_3 on the basis of protein abundance alone

(Fig 3; Supplementary Fig. S3). In any case, our results align with previous studies that found

relatively low correlation between mRNA expression and protein abundance in other plants

(Baerenfaller et al., 2012; de Sousa Abreu et al., 2009; Guan et al., 2024; Kundrátová et al.,

2021). The lack of correlation between transcript expression and protein abundance is potentially

driven by multiple factors. First, far fewer gene products were represented in our protein

abundance dataset relative to our transcript data. Many genes with low expression (below 100

TPM average) did not have associated protein data. As a result, it is possible that some

correlations were not detected. For example, proteins that are commonly present in low

abundances including transcription factors (e.g. PRR1, CCA1/LHY,) and protein kinases (e.g.

PPCK), were absent from our proteomic dataset regardless of their expression at the transcript

level. Many of these proteins are key regulators of metabolism that are associated with

transcripts that cycle in expression throughout the day and are similarly expected to exhibit

strong cycling abundance themselves (Hartwell et al., 1999). Second, there is ample evidence

that shifts in protein abundance are driven by factors other than mRNA expression including

post-transcriptional regulation, translation, and degradation of protein products (Vogel &

Marcotte, 2012; Wu et al., 2024). Finally, it is important to keep in mind that enzymatic rates are

not simply regulated by protein abundance. Processes such as protein-protein interaction (Rugen

et al., 2021; Zhang et al., 2018), allosteric regulation by metabolites (Hartman et al., 2023), and

post-translational modification (PTM) have significant effects on primary metabolism (Balparda

et al., 2023). For instance, several proteins in the CAM pathway, including PEPC (Giglioli-

Guivarc'h et al., 1996), MDH (Carr et al., 1999), and PPDK (Y.-B. Chen et al., 2014), are

subject to post-translational interactions and modifications that affect their activity. Likewise,

multiple enzymes involved in nitrogen metabolism (Costa-Broseta et al., 2020; Kaiser & Huber,

2001; Seabra et al., 2013) and photorespiration (Jossier et al., 2019; Liu et al., 2020; Palmieri et

al., 2010) are known to be regulated by PTMs.

Modification of primary metabolism in Y. aloifolia

The evolution of CAM involves regulatory rewiring of gene expression, accompanied by

shifts in primary metabolite accumulation. These broad modifications to core processes of

carbon fixation and circadian regulation likely necessitate simultaneous adjustments across other

pathways that share metabolic substrates and overlap in regulation. In particular, PEPC activity

is strongly associated with N-metabolism in C_3 plants as it is responsible for replenishing carbon

skeletons diverted from the TCA cycle for amino acid synthesis (O'Leary & Plaxton, 2015). Our

investigation into downstream effects of CAM in *Yucca* reveals significant differences in N-

metabolism between *Y. aloifolia* and *Y. filamentosa*, in agreement with previous studies

highlighting links between CAM photosynthesis and nitrogen assimilation in various CAM and

C_3+CAM lineages (Freschi et al., 2010; Gonçalves & Mercier, 2021; Pereira et al., 2018; Santos

& Salema, 1991). Under well-watered conditions, we observed distinct changes in gene

expression related to nitrate reduction and amino acid synthesis. Specifically, there were clear

differences in accumulation of glutamate and aspartate, immediate downstream products of

primary nitrogen metabolism. Accumulation of glutamate during the light period in *Y. aloifolia*

was associated with a shift in expression of GDH from midday to early morning. Conversely, the

sharp increase in aspartate abundance coupled with the drop in glutamate abundance overnight

was not accompanied by any changes in expression of AAT, which is capable of converting

OAA and glutamate to 2-OG and aspartate. Thus, increased nocturnal accumulation of OAA

may be sufficient to explain nocturnal synthesis of aspartate via AAT without any alteration to

transcript expression or protein abundance. Prior research has shown that, along with malate,

aspartate is a major transporter of assimilated carbon into the bundle sheath cells of C_4 plants

(Meister et al., 1996; Schlüter et al., 2019). Additionally, aspartate transport in and out of the

vacuole has been demonstrated in C_3 plants (Snowden et al., 2015). Accordingly, it is plausible

that at least some CAM plants including *Y. aloifolia* may use aspartate as an alternative form of

carbon storage during the night, supplementing or buffering traditional malate-based carbon

pools during CAM cycling.

Alongside well established differences in expression in genes associated with the central

CAM carboxylation and decarboxylation pathways, we observed differences in accumulation of

organic acids other than malate, such as citrate, malonic acid, and maleic acid in the CAM

species *Y. aloifolia* relative to the C_3 species *Y. filamentosa* (Supplementary Table 5). Nocturnal

accumulation of citrate has been documented in multiple CAM species (Borland & Griffiths,

1989; Franco et al., 1992; Freschi et al., 2010; Kornas et al., 2009; Lüttge, 1988, 1990).

However, its role in CAM is less clear than that of malate, as fixation of CO_2 into citrate does not

result in a net carbon gain (Lüttge, 1988). A number of theories have been put forth regarding

putative benefits of citrate accumulation in CAM plants. In the CAM pathway, the production of

malate by MDH is dependent on NADH, as opposed to citrate synthesis which results in a net

gain of NADH. As such, nocturnal citrate production may provide additional reducing power for

reduction of malate while also serving as a secondary storage mechanism for excess CO_2 as

internal stores of NADH are exhausted (Trípodi & Podestá, 2003). Additionally, in *Yucca*, we

observed a difference in expression of NR and NiR from midday in *Y. filamentosa* to the end of

the night in *Y. aloifolia*. This may suggest that the additional reducing power generated by citrate

synthesis is driving reduction of nitrate during the dark period in *Y. aloifolia,* as opposed to C_3

plants which primarily generate reducing power for N-assimilation via photosynthesis during the

day (Foyer et al., 2001). It is also possible that accumulation of citrate is merely an atavistic trait

retained in CAM plants from their C_3 ancestors (Bräutigam et al., 2017). However, this scenario

seems unlikely as we do not observe any nocturnal accumulation of citrate in *Y. filamentosa*.

In C_3 species, the decarboxylation of citrate during the day is thought to supply the

carbon skeletons for amino acid synthesis (Scheible et al., 2000). Citrate's role in linking carbon

and nitrogen metabolism in C_3 plants may offer an explanation for the inverse cycling of

aspartate and glutamate observed in *Y. aloifolia* throughout the day and night. In C_3 plants, 2-OG

is supplied for nitrogen assimilation from the decarboxylation of citrate in the mitochondria or

the cytosol (Szal & Podgórska, 2012). Additionally, there is evidence that 2-OG used for day

time nitrogen assimilation in C_3 plants originates, at least in part, from citrate accumulated

during the night (Tcherkez et al., 2009). As 2-OG is the direct substrate for glutamate synthesis

by GDH and GOGAT, the high accumulation of glutamate during the day in *Y. aloifolia* relative

to *Y. filamentosa* could be driven by the increase in citrate production during the dark period and

its subsequent degradation during the day (Fig. 4).

In both CAM and C_3 plants, there is a clear association between citrate production and

various types of abiotic stress(Lüttge, 2006; Tahjib-Ul-Arif et al., 2021). Citrate:malate ratios are

known to increase in CAM plants in response to high light stress in *Tillandsia (Borland &*

Griffiths, 1989) and high light stress and drought stress in *Clusia* (Franco et al., 1992). Citrate

production is more energetically favorable than that of malate, as the conversion of hexose to

citrate results in a net gain of ATP and NAD(P)H relative to fixation of CO_2 into malate (Lüttge,

1988). In addition, the conversion of citric acid to OAA releases two molecules of CO_2 whereas

the conversion of malate to pyruvate during the light reactions of CAM releases only a single

molecule of CO_2. As such, production of citrate may represent a way to recycle respiratory CO_2

during periods of drought where stomatal conductance is severely reduced during the day and

night, limiting access to atmospheric CO_2 (Lüttge, 1988). Likewise, under high light,

accumulation of citrate at night and the subsequent release of CO_2 upon its breakdown during the

day is thought to minimize photorespiration and photoinhibition by concentrating CO_2 around

rubisco as supplies of malate are exhausted. Finally, it has been suggested in *Clusia* that

accumulating citrate at night could increase the buffering capacity of the vacuole, leading to an

increase in storage capacity for CO_2 (Franco et al., 1992). Our observations in *Y. aloifolia* were

consistent with those in other species that accumulate citrate with the ratio of citrate to malate

increasing under drought stress. This may be due to citrate's role in minimizing oxidative stress

or alternatively may reflect the lower energetic requirements of citrate production relative to

malate (Lüttge, 1988). Regardless, our findings highlight a potentially important link between

citrate metabolism and CAM in *Yucca*.

Comparison of transcript and protein abundance over time between *Y. aloifolia* and *Y.*

filamentosa demonstrates that the metabolic changes underpinning the evolution of CAM occur

at many levels with shifts in protein kinetics and post-translational modification potentially

having a significant impact on CAM and other downstream pathways. Furthermore, analysis of

gene expression alongside fluctuations in metabolite abundance demonstrate a clear effect on

overlapping pathways of nitrogen assimilation and to a lesser extent, photorespiration. The CAM

metabolite OAA seems to play a key role in driving shifts in primary metabolism and may act as

a link between regulation of carbon and nitrogen metabolism (L.-H. Chen et al., 2022). Multiple

metabolites that exhibit day-night fluctuations in abundance in *Y. aloifolia*, including those in the

nitrogen cycle, rely on OAA as a substrate. The cycling abundance of these primary metabolites

is notable as OAA is produced in great abundance by carboxylation of PEP by the enzyme PPC

in CAM plants. The subsequent reduction of OAA to malate by MDH is dependent on the

presence of NADH. Under conditions of high light or drought, it's possible that OAA synthesis

by PPC outstrips the ability of MDH to convert OAA to malate. This could potentially result in

OAA being funneled into other processes that either generate reducing power, such as citrate

synthesis, or at the very least do not require NADH, such as aspartate synthesis. As such, it's

possible that cycling of multiple metabolites outside of the central CAM pathway may initially

be driven by greater flux through OAA rather than extensive regulatory rewiring outside of the

core CAM pathway.

Photorespiratory pathway largely conserved between CAM and C$_3$ plants

The mitigation of photorespiration is thought to play a major role in CAM evolution; by

using PEPC as the initial carboxylating enzyme to concentrate carbon around RuBisCO, CAM

plants are thought to avoid or reduce photorespiration by preventing RuBisCO's interaction with

O_2. It is therefore assumed that photorespiration decreases alongside the evolution of carbon

concentrating mechanisms like CAM and C$_4$. However, the photorespiratory pathway itself has

only rarely been the direct focus of CAM research, and studies have shown conflicting results. In

a strong CAM species, internal concentrations of CO_2:O_2 were high enough during the day to

largely suppress photorespiration, while a facultatively CAM species demonstrated high levels of

photorespiration even when CAM was induced (Duarte & Lüttge, 2007; Spalding et al., 1979).

Our results show that while there are subtle differences in gene expression and protein

abundance related to photorespiratory pathways between CAM and C$_3$ species of *Yucca*, both

remain largely conserved between species in drought-stressed and well-watered conditions. A

notable exception to this similarity in expression is found in PGP, which has daytime expression

in *Y. filamentosa* but is highly expressed at night in *Y. aloifolia* under both treatments. The

difference is particularly pronounced under drought stress where expression of PGP is nearly

inverse between species. Conserved nocturnal expression of multiple photorespiratory genes

between *Y. aloifolia* and *Y. filamentosa* combined with altered expression pattern of PGP in *Y.*

aloifolia may suggest that different portions of the photorespiratory pathway are under distinct

controls. Furthermore, it is possible that differences in expression of PGP, particularly under

drought stress could be linked to lower rates of photorespiration in *Y. aloifolia*.

Despite the broad similarities in gene expression and protein abundance between species

and treatments, we observed nocturnal accumulation of glycerate and serine, two products of the

photorespiratory pathway, exclusively under drought stress in *Y. aloifolia*. In addition, we did not

observe strong accumulation of serine or glycerate during the day in either species under either

treatment in contrast to what has been observed in other C_3 plants (Sato et al., 2008; Timm et al.,

2013). Serine is an important substrate for 1C metabolism in plants where it serves as a major

donor of one-carbon units for the biosynthesis of nucleotides, amino acids, and other essential

metabolites. Serine is primarily produced via photorespiration in C_3 plants. However, the

nocturnal accumulation of serine in *Y. aloifolia* is unlikely to be driven by photorespiratory flux,

as photorespiration is a light-dependent process (Foyer et al., 2009). Alternative pathways of

serine biosynthesis exist in plants commonly referred to as the "phosphorylated serine pathway"

and the "non-phosphorylated (glycerate) serine pathway" (Igamberdiev & Kleczkowski, 2018).

Based on the time structured nature of both glycerate and serine under drought, along with the

absence of phosphorylated intermediates in our metabolomic data, nocturnal accumulation of

serine and glycerate in *Y. aloifolia* could be driven by the poorly characterized, non-

phosphorylated serine pathway of serine biosynthesis. This pathway is thought to be an

important source of serine in C_4 plants where photorespiration is greatly reduced (Kleczkowski

& Givan, 1988; Randall et al., 1971). Many obligate CAM plants such as *Y. aloifolia* open their

stomata during the early morning and late evening (Males & Griffiths, 2017), providing an

opportunity for normal, C_3 levels of photorespiration to occur at the beginning and end of day.

However, drought stress can induce a further reduction in stomatal conductance (Herrera et al.,

2000), restricting gas exchange entirely to the dark period in *Y. aloifolia* (Supplementary Fig. S1)

and essentially eliminating fixation of O_2 by Rubisco. Thus it is likely that nocturnal

accumulation of serine and glycerate under drought stress in *Y. aloifolia* does not reflect flux

through the photorespiratory pathway, but production of serine by an alternative pathway in

response to reduced photorespiration. It is also notable that Serine has been shown to be a

potentially powerful transcriptional regulator of multiple genes in the photorespiratory pathway

(Timm et al., 2013). Specifically, external application of serine has been shown to deregulate the

expression of GDC, SHMT, and GLYK, resulting in significantly higher expression during the

night than the day in Arabidopsis. Expression of PGP was also altered but remained higher

during the day. Additionally, the deregulation of expression of photorespiratory genes resulted in

substantial discordance between expression of transcripts and abundance of their corresponding

protein products (Timm et al., 2013). This pattern is similar to what we observe in *Y. aloifolia*

and *Y. filamentosa*, potentially indicating a link between non-photorespiratory serine production

and alterations to gene expression and protein abundance in the photorespiratory pathway in

Yucca. However, the extent to which this might be tied to the evolution of CAM in *Yucca*

remains unclear.

It is also of interest that when *Y. filamentosa* experiences drought, the levels of aspartate

and glutamate have similar abundance patterns as those in *Y. aloifolia* (Fig. 3c). However, the

similarity between C_3 and CAM species in metabolite abundance under drought stress does not

appear to extend to photorespiratory metabolites found in our data. This may suggest that the

evolution of CAM was not primarily driven by shifts in the photorespiratory pathway, as drought

stress induces "CAM-like" behavior in nitrogen metabolism but not photorespiration. The similar

expression of photorespiratory genes indicates that significant changes in gene expression

outside the CAM pathway may be less critical to its establishment in *Yucca*. Instead, CAM

evolution may only initially require changes to expression of a handful of genes, which primarily

enhance metabolic flux through existing C_3 pathways rather than requiring complex, global

regulatory shifts across multiple pathways. Additional, detailed comparisons of molecular time

series and functional analysis of photorespiratory genes in related CAM and C_3 species are

needed to fully understand the dynamics we observe in *Y. aloifolia* and *Y. filamentosa*.

Considerations for future studies in CAM evolution

Our study shows the importance of conducting genomics research in closely related C_3

and CAM taxa and future studies can continue to expand the scope of our knowledge of CAM

evolution by conducting similar studies in other lineages. In addition to continuing to develop

molecular resources in the Agavoideae and other model CAM taxa, researchers should endeavor

to incorporate an ever greater diversity of taxa including aquatic CAM species and

physiologically diverse C_3 + CAM species. While C_3 + CAM should not necessarily be treated as

a transitional state, its study is likely to provide valuable insights into initial molecular changes

that selection might act upon to drive the emergence of CAM in a C_3 background. Furthermore,

the overall lack of concordance between transcript expression, protein abundance, and metabolite

abundance seen in *Yucca* suggests that while analyses based on transcriptomic data are highly

accessible and capable of providing valuable insight into CAM evolution, they are but one piece

of a much larger puzzle. Future studies would do well to keep this in mind and supplement

traditional, bulk-transcriptomic analyses with other molecular data types. In particular, the

discordance between transcriptomic and proteomic datasets in our study highlights the potential

importance of post-translational mechanisms to regulation within and outside of the CAM

pathway. As such, incorporating methods that directly examine post-translational modification

(Schiller et al., 2025) will be essential to characterizing the transition from C3 to CAM across

vascular plants.

In addition, while comparative genomics and transcriptomics are powerful tools for

identifying candidate genes associated with CAM evolution, relying on sequence homology to

distantly related C☐ plants to infer gene function can be unclear or misleading.While core

functions of genes involved in primary metabolism are likely conserved across deep evolutionary

time, many of the genes involved in nitrogen metabolism, photorespiration, and other primary

metabolic pathways are present as multiple copies that may differ in regulation, interaction, or

function between C☐ and CAM lineages. This is particularly true for regulatory genes such as

kinases and transcription factors that are often members of large, diverse gene families. Thus,

while genomic approaches are a relatively accessible and effective method for identifying

candidate genes, targeted functional research is necessary to determine which genes are central to

shifts in CAM physiology and how they are regulated. This will necessitate the development of

experimental plant transformation resources in CAM and closely related C_3 lineages to directly

test gene roles and regulatory mechanisms underlying the evolution of CAM.

Our research advances the study of CAM evolution by identifying genes that are

promising targets for focused investigations into metabolism beyond the core CAM pathway.

However the complex dynamics evident in comparisons of our transcript, protein and metabolite

datasets underscore the limitations of inferences drawn from comparative genomic analyses

alone. While continued expansion of genome and transcriptome data for related plants with

contrasting modes of carbon acquisition is of great value, advancing our understanding of CAM

physiology will ultimately require a shift from broad comparative analyses to targeted multi-

omic and experimental investigations of the molecular mechanisms and regulatory controls that

promote the convergent evolution of CAM.

Methods

Physiology

Six mature *Y. aloifolia* plants, each a different genotype, were selected from a collection

of wild-collected plants at the University of Georgia (UGA) Plant Biology greenhouse. Six

mature *Y. filamentosa* plants, also each a different genotype, were obtained either from the UGA

collection or were purchased from landscape nurseries. Plants obtained from nurseries were re-

potted in 5-gallon containers with a 50/50 mixture of standard potting soil and sand (the same

mixture in which the UGA collection plants were grown and maintained) and allowed to

acclimate to the greenhouse for at least 4 weeks. Plants were introduced into a growth chamber

at the UGA Plant Biology greenhouse in randomized positions one week before the onset of

sampling to allow entrainment to the light regime and were watered daily. Growth chamber

conditions were programmed with the following conditions: the light period commenced at 7:00

am with relative humidity of 40%, ambient temperature at 30° C, and photosynthetically active

radiation (PAR) approximately 270 µmol m^{-2} s^{-1} measured at leaf level. The dark period

commenced at 7:00 pm with relative humidity of 40%, ambient temperature at 17° C, and

photosynthetically active radiation (PAR) 0 µmol m^{-2} s^{-1}.

Gas exchange and leaf tissue sampling for the well-watered treatment commenced on

December 03, 2021. Beginning one hour after the onset of the light period ("Zeitgeber time 1",

ZT1), instantaneous gas exchange measurements were obtained with a LiCOR-6400XT portable

photosynthesis system on mid-leaf sections of mature leaves. LiCOR chamber conditions were

set to match ambient conditions in the growth chamber, including temperature, light, and

humidity; the fan speed was set to maximum and flow was set at 500 umol s^{-1}. Gas exchange

measurements were repeated on the same leaf, and a random leaf was sampled for each plant, in

the same order every four hours for the next five time points (ZT5, ZT9, ZT13, ZT17, ZT21).

Immediately following gas exchange measurements, a randomly selected, mature leaf was

excised and snap frozen in liquid N_2 and stored at -80° C. Soil moisture was measured at the end

of the 24-hour sampling period with a digital soil moisture probe inserted into the top ~3 inches

of soil for each plant. Watering was then withheld for the remainder of the experiment to induce

the drought stress. Soil moisture levels were periodically checked over the next 18 days. At day

13, it was observed that the *Y. aloifolia* species' soil was drying down faster than that of *Y.*

filamentosa. To maintain comparable soil conditions, each fast-drying plant was top-watered

with 500 mL water, bringing those plants back to similar soil water content as the others. By day

18 (December 20[th], 2021), the average soil moisture for each species was below 5% and gas

exchange and leaf tissue sampling were repeated in the same manner as the well-watered

samples.

RNA-sequencing and analysis

Whole leaf tissue for each sample was ground to a powder in a mortar and pestle under

liquid N_2, transferred to a 15 mL Falcon tube and stored in a -80° C freezer. RNA extraction for

each sample from an aliquot of approximately 100 mg of powdered leaf tissue was carried out

with the Qiagen Plant RNeasy kit with DNase I on-column digestion according to the Qiagen

standard protocol. RNA quantity was measured with a Qubit fluorometer, and RNA integrity was

measured on an Agilent 2100 Bioanalyzer with an RNA Nano 6000 kit. RNA samples were

shipped on dry ice to the DOE's Joint Genome Institute (JGI) labs in Berkeley, CA for cDNA

library construction and paired-end 150 bp Illumina sequencing on an Illumina NovaSeq

sequencer. All but five libraries produced deep sequence data (total PE150 libraries = 139,

median fragment count = 28,066,771, mean Q30 = 0.95).

RNA-seq read quantification

RNA sequence data was downloaded from the JGI server onto the Georgia Advanced

Computing Research Center computing cluster. For each species, gene annotations were used

from the draft genomes on Phytozome (Yucca aloifolia Ya24Inoko v2.1 DOE-JGI,

https://phytozome.jgi.doe.gov/info/YaloifoliaYa24Inoko_v2_1 and Yucca filamentosaC3 pri

v2.1 DOE-JGI, https://phytozome.jgi.doe.gov/info/YfilamentosaC3pri_v2_1) (pre-publication

permission from JGI was obtained for *Yucca* annotation use). Reads for each species were

pseudo-aligned to their respective primary transcript nucleotide fasta sequences and

quantification was calculated with *kallisto* (Bray et al., 2016) with default parameters. Raw read

counts were imported into expression matrices in R with *tximport* (Soneson et al., 2016).

Inference of homology with Orthofinder

Homology was assessed with Orthofinder2 (Emms & Kelly, 2019) using CDS sequences

from the *Y. aloifolia* and *Y. filamentosa* genome assemblies from Phytozome along with genomic

CDS sequences from the following taxa: *Acorus americanus* (Acorus americanus v1.1, DOE-

JGI, http://phytozome-next.jgi.doe.gov/), *Ananas comosus* (Ming et al., 2015), *Asparagus*

officinalis (Harkess et al., 2017), *Agave tequiliana* (Agave tequilana var. Weber's Blue v2.1

DOE-JGI, https://phytozome-next.jgi.doe.gov/info/Atequilanavar_WebersBlueHAP1_v2_1),

Arabidopsis thaliana (Cheng et al., 2017), *Amborella trichopoda (Amborella Genome Project,*

2013), Brachypodium distachyon (International Brachypodium Initiative, 2010), Kalanchoe

fedtschenkoi (Yang et al., 2017), *Oryza sativa* (Ouyang et al., 2007), *Sorghum bicolor*

(McCormick et al., 2018), *Solanum lycopersicum* (Tomato Genome Consortium, 2012), and *Zea*

mays (Hufford et al., 2021). CDS sequences were translated to amino acids prior to running

Orthofinder using the default settings.

Time-structured expression profiling of transcriptomic data

To assess differential gene expression patterns due to watering regimes over the 24-hour

sampling period, differential expression analysis taking time and treatments as variables was

carried out in the R software package *maSigPro* (Nueda et al., 2014). For each species, trimmed

mean of *M* values (TMM) were calculated and used to normalize read counts within species, and

counts per-million (CPM) were computed from the normalized libraries in the R software

package *edgeR* (Robinson et al. 2009). Genes with less than 0.5 CPM across all time points were

removed. Outlier libraries were checked with multidimensional scaling plots. Two and six

libraries were removed from *Y. aloifolia* and *Y. filamentosa*, respectively. Genes were tested for

significant time structure with functions *p.vector (counts=TRUE)*, and the *T.fit(step method =*

"backward", alpha = 0.05). Genes with high variability between replicates were classified as

overly influential by *maSigPro*, removed, and the analyses were rerun on the filtered data set.

Genes with significant treatment effects across time were called with *sig.siggenes(rsq = 0.6)*. To

sort genes into clusters based on time-structured expression similarity, *maSigPro* requires the

user to input the expected numbers of clusters (*k*). To estimate *k,* within-group sums of squares

were calculated and plotted so the *k* value at which marginal reductions in the sums of squares

can be determined. *k*=6 was chosen for both species and genes were sorted into clusters using

see.genes(cluster.method="hclust").

Metabolite extraction and protein isolation

2.0 ml tubes (Safe-Lock, Eppendorf) were filled with leaf material to approximately ¼

capacity, covered with tube membranes, and samples were lyophilized overnight. Lyophilized

leaves were then homogenized in a Tissue Lyser II (Qiagen) using tungsten beads, specifically

two-3mm beads and one-5mm in each tube. Samples were lysed for 2 min at a frequency of 23

Hz until ground to a powder. Metabolites from 20 mg of the ground leaves were extracted with

800 µl of a 4:1 methanol: water mix, and each sample spiked with 50 µl of a 330 µg/ml solution

of glycine d5 (Cambridge Isotopes). Samples were then placed in a thermomixer and shaken at

1200 rpm and 21°C for 1 hr. Following centrifugation at 5000 x g, supernatant containing the

metabolite fraction was removed and 10% of the total metabolite extract was transferred to a

glass vial to be dried under vacuum for subsequent metabolomic analysis (described below).

After removing the metabolite fraction, protein pellets were washed twice by adding 500 µl

100% MeOH, vortexing for 15 seconds, and centrifuging at 10000 x g and 21°C for 10 min. The

resulting pellet was dried in a fume hood until remaining MeOH had completely evaporated and

stored at -80°C until protein digestion.

Protein pellets were resuspended in 120 µl of an aqueous solution of 8 M urea and 50

mM HEPES by vortexing and sonication for 30 sec. Following resuspension, 6.3 μL 1 M

aqueous solution of DTT was added to each sample and samples were placed in a thermomixer

and shaken at 500 rpm and 37°C for 30 min for reduction. Next, 40 μL 0.5 M aqueous solution

of iodoacetamide was added to each sample and samples were placed in a thermomixer and

shaken for another hour at 500 rpm and 37°C for alkylation. Samples were then diluted with 800

μL 50 mM aqueous solution of HEPES prior to digestion with trypsin (1:50 w/w trypsin:protein)

for 3 hrs at 500 rpm and 37°C. Sample clean-up was performed using C18 SPE media and

samples were concentrated to ~100 μL and peptide concentration was determined via BCA

assay. Finally, 30 μg of each sample were transferred to a new 1.5 mL tube, dried down in a

speedvac and stored at -80°C for TMT labeling.

TMT isobaric tag labeling

Samples were designed into 12 TMT sets (6 per plant) including two pools, one

consisting of a mix of the well-watered plants and one of all the drought-stressed plants to ensure

correlation across sets. Each set included 1 replicate of each condition and blanks to establish a

base background noise signal and then each set completely randomized. Each sample was diluted

in 40 µl 500 mM HEPES, pH 8.5 and labeled using amine-reactive Tandem Mass Tag Isobaric

Mass Tagging Kits (Thermo Scientific, Rockford, IL) according to the manufacturer's

instructions. Briefly, 250 µL of anhydrous acetonitrile was added to each 5 mg reagent, vortexed,

and allowed to dissolve for 5 mins with occasional vortexing. Reagents (5 µl) were then added to

each sample and incubated for 1 hour at room temperature with shaking at 400 rpm. Each sample

was then diluted with 30 µl 20% acetonitrile. A portion from each sample was collected as a

premix to run on a mass spectrometer to ensure complete labeling. The samples were frozen at -

80°C until the results showed efficient labeling. At that point the frozen samples were thawed

and the reaction was quenched by adding 8 µL of 5 % hydroxylamine to the sample with

incubation for 15 mins at room temperature with shaking at 400 rpm. The samples within each

set were then combined and completely dried in the speedvac. Each of the samples were then

resuspended in 0.1% trifluoroacetic acid solution and cleaned using Discovery C18 50 mg/1 mL

solid phase extraction tubes as described above and once again assayed with bicinchoninic acid

to determine the final peptide concentration and vialed for microfractionation.

Microfractionation

Peptide mixtures were separated by high resolution reversed phase UPLC using a

nanoACQUITY UPLC® system (Waters Corporation, Milford, MA) equipped with an

autosampler. Capillary columns, 200 µm i.d. × 65 cm long, were packed with 300Å 3.0 µm p.s.

Jupiter C18 bonded particles (Phenomenex, Torrence, CA). Separations were performed at a

flow rate of 2.2 µL/min on binary pump systems, using 10 mM ammonium formate (pH 7.5) as

mobile phase A and 100% acetonitrile as mobile phase B. 20 µL of TMT labeled peptide

mixtures (0.5 µg/µL) were loaded onto the column and separated using a binary gradient of 1 %

B for 35 min, 1-10 % B in 2 min, 10-15 % B in 5 min, and 15-25 % B in 35 min, 25-32 % B in

25min, 32-40 % B in 13 min, 40-90 % B in 43 min, held at 90 % B for 2 min (column washed

and equilibrated from 90-50 % B in 2 min, 50-95% B in 2 min, held at 95 % B for 2 min and 95-

1 % in 4 min). The capillary column eluent was automatically deposited every minute into 12x32

mm polypropylene vials (Waters Corporation, Milford, MA) starting at minute 60 and ending at

minute 170, over the course of the 180 min LC run. Fractions were concatenated by collecting

fractions from vial 1 to vial 24 and then returning to vial 1 back to vial 24 (etc.). Prior to peptide

fraction collection, 100 µL of water was added to each vial to aid the addition of small droplets

into the vial. Each vial was completely dried in a vacuum concentrator (Labconco, Kansas City,

MO), reconstituted in 15 µL 2 % acetonitrile, 0.1 % formic acid and stored at -20°C until LC-

MS/MS analysis.

TMT quantitation of isolated peptides

A Waters nano-Acquity dual pumping UPLC system (Milford, MA) was configured for

on-line trapping of a 5 µL injection at 5 µL/min for 5 min with reverse-flow elution onto the

analytical column at 200 nL/min using mobile phase A. The analytical column was slurry packed

in-house using Waters BEH C18 1.7 um particles (Waters Chromatography, Drinagh, Ireland)

into a 25 cm PicoFrit 360um od x 75um id column (New Objective, Littleton, MA) and held at

45C during use by a 15 cm AgileSleeve flexible capillary heater (Analytical Sales and Services,

Inc., Flanders, NJ). The trapping column was slurry packed in-house using Jupiter 5um C18

particles (Phenomenex, Torrence, CA) into a 4cm long piece of 360um od x 150um id fused

silica (Polymicro Technologies, Phoenix, AZ) and solgel fritted at both ends. Mobile phases

consisted of (A) 0.1% formic acid in water and (B) 0.1% formic acid in acetonitrile with the

following gradient profile (min, %B): 0, 1; 10, 8; 105, 25; 115, 35; 120, 75; 123, 95; 129, 95;

130, 50; 132, 95; 138, 95; 140, 1.

MS analysis of 24 fractions per plex set was performed using a Orbitrap Fusion Lumos

Tribrid MS (ThermoFisher Scientific) outfitted with a homemade nano electrospray ionization

interface. Electrospray emitters were homemade using 150 µm o.d. × 20 µm i.d. chemically

etched fused silica. The ion transfer tube temperature and spray voltage were 250°C and 2.2 kV,

respectively. Peptides were separated on an in-house prepared 40 cm x 75 um 1.7-um Waters

BEH C18 column, mobile phases consisted of 0.1% formic acid in water (MP-A) and 0.1%

formic acid in acetonitrile (MP-B). Data were collected for 120 min following a 10 min delay

after completion of sample trapping and start of LC gradient.

FT MS spectra were acquired from 350 to 1800 m/z at a resolution of 60 k (AGC

target 4e5) and the FT HCD MS/MS spectra were acquired in data dependent mode for ions

with detected charge states 2-6 and minimum intensity of 2.5e3 using a normalized collision

energy of 30. Other MS/MS parameters were as follows - isolation window was set to 0.7 m/z at

a resolution of 50 k, AGC target 1e5, and dynamic exclusion time was 45 s.

Acquired raw spectra were processed using automated MSGF+ software (S. Kim et al.,

2008) in target/decoy mode with 20 ppm parent ion mass tolerance, partial tryptic rule with

oxidation of methionine (+15.9949) as dynamic modification and static modifications

carbamidomethylation of cysteine (+57.0215) and TMTpro modification of lysine on peptide N

termini (+304.2071). Each set of samples was matched with corresponding protein collections

for *Y. aloifolia* (43,466 protein sequences) and *Y. filamentosa* (45,266 protein sequences).

Common contaminant proteins from trypsin and keratins (16 protein sequences) were considered

in each search.

Top peptide-spectral matches (PSM) passing 1% FDR cut-off established for the whole

datasets of *Y. aloifolia* and *Y. filamentosa* were used for subsequent TMT quantification. MASIC

(Monroe et al., 2008) software was used to pull abundances of the sample reporter ions for all

PSMs. Measured reporter ion abundances were log transformed to remove skewness of

abundance distribution and normalized using mean central tendency normalization. Peptide

abundances from all fractions were summed prior to normalization, normalization was performed

using InfernoRDN (Polpitiya et al., 2008) software. Protein level roll-up of normalized peptide

abundances was normalized again using the same strategy and software.

Proteomic abundance analysis

Normalized protein abundance data was filtered to remove proteins supported by less

than two peptide matches as well as proteins that were missing abundance data for more than ⅓

of samples. After filtering, abundance values were converted to z-scores in R to facilitate

removal of outlier data points, identification of proteins that fluctuate in abundance throughout

the day (a.k.a. "cycling" proteins), and high level comparisons between species and timepoints.

For each gene and treatment, individual abundance values more than two standard deviations

above or below the next closest abundance value at a given time point were considered outliers

and removed prior to identifying cycling proteins.

Cycling proteins were identified and peak mean expression was calculated using a

custom R script. To identify cycling proteins, filtered, z-scaled abundance data was divided into

drought-stressed and well-watered conditions prior to fitting polynomial functions for linear,

quadratic, cubic, and quartic models. The best fitting model was determined using an ANOVA

test. Proteins were considered cycling if any model provided a significantly better fit than the

linear model ($p < 0.05$). R scripts used for z-scaling, outlier removal, and identification of

cycling proteins are available on GitHub at https://github.com/dawickell/Yucca-omics.

Untargeted metabolomic quantitation and analysis

Dried metabolite extracts were chemically derivatized as previously reported (Y.-M. Kim

et al., 2015). Briefly, the extracts were derivatized by methoxyamination and trimethylsilyation

(TMS), then the samples were analyzed by GC-MS. Samples were acquired in an Agilent GC

7890A using a HP-5MS column (30 m × 0.25 mm × 0.25 μm; Agilent Technologies, Santa

Clara, CA) coupled with a single quadrupole MSD 5975C (Agilent Technologies). One

microliter of sample was injected into a splitless port at constant temperature of 250°C. The GC

temperature gradient started at 60 °C, with a hold of temperature for 1 minute after injection,

followed by increase to 325 °C at a rate of 10 °C/minute and a 10-minute hold at this

temperature. A fatty acid methyl ester standard mix (C8-28) (Sigma-Aldrich) was analyzed in

parallel as standard for retention time calibration.

GC-MS raw data files were processed using the Metabolite Detector software (Hiller et

al., 2009). Retention indices (RI) of detected metabolites were calculated based on the analysis

of a FAMEs mixture, followed by their chromatographic alignment across all analyses after

deconvolution. Metabolites were initially identified by matching experimental spectra to a PNNL

augmented version of Agilent GC-MS metabolomics Library, containing spectra and validated

retention indices for over 1200 metabolites. Then, the unknown peaks were additionally matched

with the NIST20/Wiley11 GC-MS library. All metabolite identification and quantification ions

were validated and confirmed to reduce deconvolution errors during automated data-processing

and to eliminate false identifications. Peak areas for identified metabolites and unknowns were

normalized to the peak area of the internal standard glycine d5, reported and used for further

statistical analysis.

To assess shifts in metabolite abundance over time in *Y. aloifolia* and *Y. filamentosa* we

used the MaSigPro package in R. Spectral counts of identified metabolite abundance were tested

for significant time structure with the *p.vector (counts=TRUE)* function.

Accession Numbers

RNA seq data for our gene expression profiling analyses can be found in NCBI's

Sequencing Read Archive; SRA BioProject numbers: 'PRJNA1162856', 'PRJNA1162850',

'PRJNA1162828', 'PRJNA1162846', 'PRJNA1162829', 'PRJNA1162847', 'PRJNA1162862',

'PRJNA1162861', 'PRJNA1162852', 'PRJNA1162845', 'PRJNA1162853', 'PRJNA1162857'.

Supplementary Data

Supplementary Figure 1: Stomatal conductance varies by treatment and condition in *Yucca*.

Supplementary Figure 2: Detailed comparison of cycling transcripts and proteins in *Y. aloifolia*

and *Y. filamentosa*.

Supplementary Figure 3: Detailed mRNA expression and protein abundance data for all

paralogs of genes in the CAM pathway.

Supplementary Figure 4: Detailed mRNA expression and protein abundance data for all

paralogs of genes associated with citrate metabolism.

Supplementary Figure 5: Detailed mRNA expression and protein abundance data for all

paralogs of genes associated with nitrogen metabolism.

Supplementary Figure 6: Detailed mRNA expression and protein abundance data for all

paralogs of genes associated with photorespiration.

Supplementary Table 1: Summary of maSigPro results and peak mean expression for *Y.*

aloifolia.

Supplementary Table 2: Summary of maSigPro results and peak mean expression for *Y.*

filamentosa.

Supplementary Table 3: Results of linear modeling on protein data for *Y. aloifolia*.

Supplementary Table 4: Results of linear modeling on protein data for *Y. filamentosa*.

Supplementary Table 5: Summary of metabolomics data fo *Y. aloifolia* and *Y. filamentosa*

Supplementary Table 6: Summary statistics for *Y. aloifolia* and *Y. filamentosa* genes associated

with CAM, N-metabolism, photorespiration, and citrate metabolism.

Acknowledgements

The authors would like to thank Jen Liddle and Jeremy Balsbaugh for consultation on proteomic

analysis, Kerrie Berry for project management at JGI, and Amir Ahkami and Sarah Leichty for

project management at EMSL. We would also like to thank the Department of Energy Joint

1002

Author Contributions

1004   R.F., J.L-M, and K.H designed the research; R.F., K.W., R.C., J.T., N.T., N.M.M., H.H., E.S.,

1005   and A.L. contributed to data collection, including physiology, transcriptomic, metabolomic, and

1006   proteomic; D.W. led analyses with assistance from R.F. and K.H.; D.W. wrote the paper with

1007   assistance from K.H.

1008

References:

Abraham, P. E., Yin, H., Borland, A. M., Weighill, D., Lim, S. D., De Paoli, H. C., Engle, N.,

Jones, P. C., Agh, R., Weston, D. J., Wullschleger, S. D., Tschaplinski, T., Jacobson, D.,

Cushman, J. C., Hettich, R. L., Tuskan, G. A., & Yang, X. (2016). Transcript, protein and

metabolite temporal dynamics in the CAM plant Agave. *Nature Plants*, *2*, 16178.

Agarie, S., Umemoto, M., Sunagawa, H., Anai, T., & Cushman, J. C. (2020). An Agrobacterium-

mediated transformation via organogenesis regeneration of a facultative CAM plant, the

common ice plant *Mesembryanthemum crystallinum* L. *Plant Production Science*, *23*(3),

343–349.

Amborella Genome Project. (2013). The Amborella genome and the evolution of flowering

plants. *Science*, *342*(6165), 1241089.

Baerenfaller, K., Massonnet, C., Walsh, S., Baginsky, S., Bühlmann, P., Hennig, L.,

Hirsch☐Hoffmann, M., Howell, K. A., Kahlau, S., Radziejwoski, A., Russenberger, D.,

Rutishauser, D., Small, I., Stekhoven, D., Sulpice, R., Svozil, J., Wuyts, N., Stitt, M.,

Hilson, P., … Gruissem, W. (2012). Systems☐based analysis of Arabidopsis leaf growth

reveals adaptation to water deficit. *Molecular Systems Biology*, *8*(1), 606.

Balparda, M., Bouzid, M., Martinez, M. D. P., Zheng, K., Schwarzländer, M., & Maurino, V. G.

(2023). Regulation of plant carbon assimilation metabolism by post-translational

modifications. *The Plant Journal: For Cell and Molecular Biology*, *114*(5), 1059–1079.

Borland, A. M., & Griffiths, H. (1989). The Regulation of Citric Acid Accumulation and Carbon

Recycling During CAM in *Ananas comosus*. *Journal of Experimental Botany*, *40*(1), 53–60.

Boxall, S. F., Kadu, N., Dever, L. V., Kneřová, J., Waller, J. L., Gould, P. J., & Hartwell, J.

(2020). *Kalanchoë* PPC1 is essential for crassulacean acid metabolism and the regulation of

core circadian clock and guard cell signaling genes. *The Plant Cell*, *32*(4), 1136–1160.

Bräutigam, A., Schlüter, U., Eisenhut, M., & Gowik, U. (2017). On the Evolutionary Origin of

CAM Photosynthesis. *Plant Physiology*, *174*(2), 473–477.

Carr, P. D., Verger, D., Ashton, A. R., & Ollis, D. L. (1999). Chloroplast NADP-malate

dehydrogenase: structural basis of light-dependent regulation of activity by thiol oxidation

and reduction. *Structure* , *7*(4), 461–475.

Cheng, C.-Y., Krishnakumar, V., Chan, A. P., Thibaud-Nissen, F., Schobel, S., & Town, C. D.

(2017). Araport11: a complete reannotation of the Arabidopsis thaliana reference genome.

The Plant Journal: For Cell and Molecular Biology, *89*(4), 789–804.

Chen, L.-H., Cheng, Z., Xu, M., Yang, Z., & Yang, L. (2022). Effects of nitrogen deficiency on

the metabolism of organic acids and amino acids in Oryza sativa. *Plants*, *11*.

https://doi.org/10.3390/plants11192576

Chen, L.-Y., Xin, Y., Wai, C. M., Liu, J., & Ming, R. (2020). The role of cis-elements in the

evolution of crassulacean acid metabolism photosynthesis. *Horticulture Research*, *7*, 5.

Chen, Y.-B., Lu, T.-C., Wang, H.-X., Shen, J., Bu, T.-T., Chao, Q., Gao, Z.-F., Zhu, X.-G.,

Wang, Y.-F., & Wang, B.-C. (2014). Posttranslational Modification of Maize Chloroplast

Pyruvate Orthophosphate Dikinase Reveals the Precise Regulatory Mechanism of Its

Enzymatic Activity. *Plant Physiology*, *165*(2), 534–549.

Costa-Broseta, Á., Castillo, M., & León, J. (2020). Nitrite Reductase 1 Is a Target of Nitric

Oxide-Mediated Post-Translational Modifications and Controls Nitrogen Flux and Growth

in Arabidopsis. *International Journal of Molecular Sciences*, *21*(19), 7270.

de Sousa Abreu, R., Penalva, L. O., Marcotte, E. M., & Vogel, C. (2009). Global signatures of

protein and mRNA expression levels. *Molecular bioSystems*, *5*(12), 1512–1526.

Dever, L. V., Boxall, S. F., Kneřová, J., & Hartwell, J. (2015). Transgenic perturbation of the

decarboxylation phase of Crassulacean acid metabolism alters physiology and metabolism

but has only a small effect on growth. *Plant Physiology*, *167*(1), 44–59.

Duarte, H. M., & Lüttge, U. (2007). Correlation between photorespiration, CO2-assimilation and

spatiotemporal dynamics of photosynthesis in leaves of the C3-photosynthesis/crassulacean

acid metabolism-intermediate species *Clusia minor* L. (Clusiaceae). *Trees* , *21*(5), 531–540.

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for

comparative genomics. *Genome Biology*, *20*(1), 238.

Ferrari, R. C., Bittencourt, P. P., Nagumo, P. Y., Oliveira, W. S., Rodrigues, M. A., Hartwell, J.,

& Freschi, L. (2020). Developing Portulaca oleracea as a model system for functional

genomics analysis of C4/CAM photosynthesis. *Functional Plant Biology: FPB*, *48*(7), 666–

682.

Foyer, C. H., Bloom, A. J., Queval, G., & Noctor, G. (2009). Photorespiratory metabolism:

genes, mutants, energetics, and redox signaling. *Annual Review of Plant Biology*, *60*(1),

455–484.

Foyer, C. H., Ferrario-Méry, S., & Noctor, G. (2001). Interactions between carbon and nitrogen

metabolism. In *Plant Nitrogen* (pp. 237–254). Springer Berlin Heidelberg.

Franco, A. C., Ball, E., & Lüttge, U. (1992). Differential effects of drought and light levels on

accumulation of citric and malic acids during CAM in *Clusia*. *Plant, Cell & Environment*,

15(7), 821–829.

Freschi, L., Rodrigues, M. A., Tiné, M. A. S., & Mercier, H. (2010). Correlation between citric

acid and nitrate metabolisms during CAM cycle in the atmospheric bromeliad *Tillandsia*

pohliana. *Journal of Plant Physiology*, *167*(18), 1577–1583.

Giglioli-Guivarc'h, N., Pierre, J. N., Brown, S., Chollet, R., Vidal, J., & Gadal, P. (1996). The

Light-Dependent Transduction Pathway Controlling the Regulatory Phosphorylation of C4

Phosphoenolpyruvate Carboxylase in Protoplasts from *Digitaria sanguinalis*. *The Plant*

Cell, *8*(4), 573–586.

Gilman, I. S., Smith, J. A. C., Holtum, J. A. M., Sage, R. F., Silvera, K., Winter, K., & Edwards,

E. J. (2023). The CAM lineages of planet Earth. *Annals of Botany*, *132*(4), 627–654.

Gonçalves, A. Z., & Mercier, H. (2021). Transcriptomic and Biochemical Analysis Reveal

Integrative Pathways Between Carbon and Nitrogen Metabolism in *Guzmania monostachia*

(Bromeliaceae) Under Drought. *Frontiers in Plant Science*, *12*, 715289.

Guan, Q., Kong, W., Tan, B., Zhu, W., Akter, T., Li, J., Tian, J., & Chen, S. (2024). Multiomics

unravels potential molecular switches in the C3 to CAM transition of Mesembryanthemum

crystallinum. *Journal of Proteomics*, *299*(105145), 105145.

Harkess, A., Zhou, J., Xu, C., Bowers, J. E., Van der Hulst, R., Ayyampalayam, S., Mercati, F.,

Riccardi, P., McKain, M. R., Kakrana, A., Tang, H., Ray, J., Groenendijk, J., Arikit, S.,

Mathioni, S. M., Nakano, M., Shan, H., Telgmann-Rauber, A., Kanno, A., … Chen, G.

(2017). The asparagus genome sheds light on the origin and evolution of a young Y

chromosome. *Nature Communications*, *8*(1), 1279.

Hartman, M. D., Rojas, B. E., Iglesias, A. A., & Figueroa, C. M. (2023). The involvement of

allosteric effectors and post-translational modifications in the control of plant central carbon

metabolism. *The Plant Journal: For Cell and Molecular Biology*, *114*(5), 1037–1058.

Hartwell, J., Gill, A., Nimmo, G. A., Wilkins, M. B., Jenkins, G. I., & Nimmo, H. G. (1999).

Phosphoenolpyruvate carboxylase kinase is a novel protein kinase regulated at the level of

expression. *The Plant Journal: For Cell and Molecular Biology*, *20*(3), 333–342.

Herrera, A., Fernández, M., & Taisma, M. A. (2000). Effects of drought on CAM and water

relations in plants of Peperomia carnevalii. *Annals of Botany*, *86*, 511–517.

Heyduk, K., Burrell, N., Lalani, F., & Leebens-Mack, J. (2016). Gas exchange and leaf anatomy

of a C3-CAM hybrid, *Yucca gloriosa* (Asparagaceae). *Journal of Experimental Botany*,

67(5), 1369–1379.

Heyduk, K., McAssey, E. V., Field, R., & Leebens-Mack, J. (2023). The Agavoideae: an

emergent model clade for CAM evolutionary biology. *Annals of Botany*, *132*(4), 727–737.

Heyduk, K., McAssey, E. V., & Leebens-Mack, J. (2022). Differential timing of gene expression

and recruitment in independent origins of CAM in the Agavoideae (Asparagaceae). *The*

New Phytologist, *235*(5), 2111–2126.

Heyduk, K., McKain, M. R., Lalani, F., & Leebens-Mack, J. (2016). Evolution of a CAM

anatomy predates the origins of Crassulacean acid metabolism in the Agavoideae

(Asparagaceae). *Molecular Phylogenetics and Evolution*, *105*, 102–113.

Heyduk, K., Ray, J. N., Ayyampalayam, S., Moledina, N., Borland, A., Harding, S. A., Tsai, C.-

J., & Leebens-Mack, J. (2019). Shared expression of crassulacean acid metabolism (CAM)

genes pre-dates the origin of CAM in the genus *Yucca*. *Journal of Experimental Botany*,

70(22), 6597–6609.

Heyduk, K., Ray, J. N., & Leebens-Mack, J. (2020). Leaf anatomy is not correlated to CAM

function in a C3+CAM hybrid species, *Yucca gloriosa*. *Annals of Botany*, *127*(4), 437–449.

Hiller, K., Hangebrauk, J., Jäger, C., Spura, J., Schreiber, K., & Schomburg, D. (2009).

MetaboliteDetector: comprehensive analysis tool for targeted and nontargeted GC/MS

based metabolome analysis. *Analytical Chemistry*, *81*(9), 3429–3439.

Hufford, M. B., Seetharam, A. S., Woodhouse, M. R., Chougule, K. M., Ou, S., Liu, J., Ricci, W.

A., Guo, T., Olson, A., Qiu, Y., Della Coletta, R., Tittes, S., Hudson, A. I., Marand, A. P.,

Wei, S., Lu, Z., Wang, B., Tello-Ruiz, M. K., Piri, R. D., … Dawe, R. K. (2021). De novo

assembly, annotation, and comparative analysis of 26 diverse maize genomes. *Science (New*

York, N.Y.), *373*(6555), 655–662.

Igamberdiev, A. U., & Kleczkowski, L. A. (2018). The glycerate and phosphorylated pathways

of Serine synthesis in plants: The branches of plant glycolysis linking carbon and nitrogen

metabolism. *Frontiers in Plant Science*, *9*, 318.

International Brachypodium Initiative. (2010). Genome sequencing and analysis of the model

grass Brachypodium distachyon. *Nature*, *463*(7282), 763–768.

Jossier, M., Liu, Y., Massot, S., & Hodges, M. (2019). Enzymatic Properties of Recombinant

Phospho-Mimetic Photorespiratory Glycolate Oxidases from *Arabidopsis thaliana* and *Zea*

mays. *Plants*, *9*(1), 27.

Kaiser, W. M., & Huber, S. C. (2001). Post‐translational regulation of nitrate reductase:

mechanism, physiological relevance and environmental triggers. *Journal of Experimental*

Botany, *52*(363), 1981–1989.

Kim, S., Gupta, N., & Pevzner, P. A. (2008). Spectral probabilities and generating functions of

tandem mass spectra: a strike against decoy databases. *Journal of Proteome Research*, *7*(8),

3354–3363.

Kim, Y.-M., Nowack, S., Olsen, M. T., Becraft, E. D., Wood, J. M., Thiel, V., Klapper, I., Kühl,

M., Fredrickson, J. K., Bryant, D. A., Ward, D. M., & Metz, T. O. (2015). Diel

metabolomics analysis of a hot spring chlorophototrophic microbial mat leads to new

hypotheses of community member metabolisms. *Frontiers in Microbiology*, *6*, 209.

Kleczkowski, L. A., & Givan, C. V. (1988). Serine formation in leaves by mechanisms other

than the glycolate pathway. *Journal of Plant Physiology*, *132*(6), 641–652.

Kornas, A., Fischer-Schliebs, E., Lüttge, U., & Miszalski, Z. (2009). Adaptation of the obligate

CAM plant *Clusia alata* to light stress: Metabolic responses. *Journal of Plant Physiology*,

166(17), 1914–1922.

Kundrátová, K., Bartas, M., Pečinka, P., Hejna, O., Rychlá, A., Čurn, V., & Červeň, J. (2021).

Transcriptomic and Proteomic Analysis of Drought Stress Response in Opium Poppy Plants

during the First Week of Germination. *Plants*, *10*(9), 1878.

Liu, Y., Guérard, F., Hodges, M., & Jossier, M. (2020). Phosphomimetic T335D Mutation of

Hydroxypyruvate Reductase 1 Modifies Cofactor Specificity and Impacts Arabidopsis

Growth in Air. *Plant Physiology*, *183*(1), 194–205.

Lüttge, U. (1988). Day□night changes of citric□acid levels in crassulacean acid metabolism:

phenomenon and ecophysiological significance. *Plant, Cell & Environment*, *11*(6), 445–

451.

Lüttge, U. (1990). Nocturnal citrate accumulation and its response to environmental stress in the

CAM plant *Kalanchoe pinnata* (Lam.) Pers. *Plant, Cell & Environment*, *13*(9), 977–982.

Lüttge, U. (2006). Photosynthetic flexibility and ecophysiological plasticity: questions and

lessons from *Clusia*, the only CAM tree, in the neotropics. *The New Phytologist*, *171*(1), 7–

25.

Lüttge, U. (2011). Photorespiration in Phase III of Crassulacean Acid Metabolism: Evolutionary

and Ecophysiological Implications. In U. E. Lüttge, W. Beyschlag, B. Büdel, & D. Francis

(Eds.), *Progress in Botany 72* (pp. 371–384). Springer Berlin Heidelberg.

Males, J., & Griffiths, H. (2017). Stomatal biology of CAM Plants1[CC-BY]. *Plant Physiology*,

174, 550–560.

McCormick, R. F., Truong, S. K., Sreedasyam, A., Jenkins, J., Shu, S., Sims, D., Kennedy, M.,

Amirebrahimi, M., Weers, B. D., McKinley, B., Mattison, A., Morishige, D. T., Grimwood,

J., Schmutz, J., & Mullet, J. E. (2018). The Sorghum bicolor reference genome: improved

assembly, gene annotations, a transcriptome atlas, and signatures of genome organization.

The Plant Journal: For Cell and Molecular Biology, *93*(2), 338–354.

McKain, M. R., McNeal, J. R., Kellar, P. R., Eguiarte, L. E., Pires, J. C., & Leebens-Mack, J.

(2016). Timing of rapid diversification and convergent origins of active pollination within

Agavoideae (Asparagaceae). *American Journal of Botany*, *103*(10), 1717–1729.

Meister, M., Agostino, A., & Hatch, M. (1996). The roles of malate and aspartate in C4

photosynthetic metabolism of Flaveria bidentis (L.). *Planta*, *199*(2).

https://doi.org/10.1007/bf00196567

Ming, R., VanBuren, R., Wai, C. M., Tang, H., Schatz, M. C., Bowers, J. E., Lyons, E., Wang,

M.-L., Chen, J., Biggers, E., Zhang, J., Huang, L., Zhang, L., Miao, W., Zhang, J., Ye, Z.,

Miao, C., Lin, Z., Wang, H., … Yu, Q. (2015). The pineapple genome and the evolution of

CAM photosynthesis. *Nature Genetics*, *47*(12), 1435–1442.

Miszalski, Z., Kornas, A., Rozpądek, P., Fischer-Schliebs, E., & Lüttge, U. (2013). Independent

fluctuations of malate and citrate in the CAM species *Clusia hilariana* Schltdl. under low

light and high light in relation to photoprotection. *Journal of Plant Physiology*, *170*(5), 453–

458.

Monroe, M. E., Shaw, J. L., Daly, D. S., Adkins, J. N., & Smith, R. D. (2008). MASIC: a

software program for fast quantitation and flexible visualization of chromatographic profiles

from detected LC-MS(/MS) features. *Computational Biology and Chemistry*, *32*(3), 215–

217.

Nobel, P. S. (1983). Nutrient levels in cacti-relation to nocturnal acid accumulation and growth.

American Journal of Botany, *70*(8), 1244–1253.

O'Leary, B., & Plaxton, W. C. (2015). The central role of glutamate and aspartate in the post-

translational control of respiration and nitrogen assimilation in plant cells. In J. P. F.

D'Mello (Ed.), *Amino acids in higher plants* (1st ed., pp. 277–297). CAB International.

Ota, K. (1988). CAM Photosynthesis under Drought Conditions in *Kalanchoe blossfeldiana*

Grown with Nitrate or Ammonium as the Sole Nitrogen Source. *Plant & Cell Physiology*,

29(5), 801–806.

Ouyang, S., Zhu, W., Hamilton, J., Lin, H., Campbell, M., Childs, K., Thibaud-Nissen, F.,

Malek, R. L., Lee, Y., Zheng, L., Orvis, J., Haas, B., Wortman, J., & Buell, C. R. (2007).

The TIGR Rice Genome Annotation Resource: improvements and new features. *Nucleic*

Acids Research, *35*(Database issue), D883–D887.

Palmieri, M. C., Lindermayr, C., Bauwe, H., Steinhauser, C., & Durner, J. (2010). Regulation of

plant glycine decarboxylase by s-nitrosylation and glutathionylation. *Plant Physiology*,

152(3), 1514–1528.

Pereira, P. N., & Cushman, J. C. (2019). Exploring the Relationship between Crassulacean Acid

Metabolism (CAM) and Mineral Nutrition with a Special Focus on Nitrogen. *International*

Journal of Molecular Sciences, *20*(18), 4363.

Pereira, P. N., Gaspar, M., Smith, J. A. C., & Mercier, H. (2018). Ammonium intensifies CAM

photosynthesis and counteracts drought effects by increasing malate transport and

antioxidant capacity in *Guzmania monostachia*. *Journal of Experimental Botany*, *69*(8),

1993–2003.

Polpitiya, A. D., Qian, W.-J., Jaitly, N., Petyuk, V. A., Adkins, J. N., Camp, D. G., 2nd,

Anderson, G. A., & Smith, R. D. (2008). DAnTE: a statistical tool for quantitative analysis

of -omics data. *Bioinformatics* , *24*(13), 1556–1558.

Ramachandra Reddy, A., & Rama Das, V. S. (1978). The Decarboxylating Systems in Fourteen

Taxa Exhibiting CAM Pathway. *Zeitschrift Für Pflanzenphysiologie*, *86*(2), 141–146.

Randall, D. D., Tolbert, N. E., & Gremel, D. (1971). 3-phosphoglycerate phosphatase in plants:

II. Distribution, physiological considerations, and comparison with P-glycolate

phosphatase: II. Distribution, physiological considerations, and comparison with P-

glycolate phosphatase. *Plant Physiology*, *48*(4), 480–487.

Robison, T. A., Oh, Z., Lafferty, D. J., Xu, X., Villarreal, J. C. A., Gunn, L. H., & Li, F.-W.

(2024). Hornworts reveal a spatial model for pyrenoid-based CO_2-concentrating

mechanisms in land plants. *bioRxiv*. https://doi.org/10.1101/2024.06.26.600872

Rodrigues, M. A., Freschi, L., Pereira, P. N., & Mercier, H. (2014). Interactions Between

Nutrients and Crassulacean Acid Metabolism. In U. Lüttge, W. Beyschlag, & J. Cushman

(Eds.), *Progress in Botany: Vol. 75* (pp. 167–186). Springer Berlin Heidelberg.

Rugen, N., Schaarschmidt, F., Eirich, J., Finkemeier, I., Braun, H.-P., & Eubel, H. (2021).

Protein interaction patterns in Arabidopsis thaliana leaf mitochondria change in dependence

to light. *Biochimica et Biophysica Acta, Bioenergetics*, *1862*(8), 148443.

Sage, R. F., Sage, T. L., & Kocacinar, F. (2012). Photorespiration and the evolution of C4

photosynthesis. *Annual Review of Plant Biology*, *63*(1), 19–47.

Santos, I., & Salema, R. (1991). Nitrogen nutrition and the level of Crassulacean acid

metabolism in *Kalanchoë lateritia* Engl. *Plant, Cell & Environment*, *14*(3), 311–317.

Sato, S., Arita, M., Soga, T., Nishioka, T., & Tomita, M. (2008). Time-resolved metabolomics

reveals metabolic modulation in rice foliage. *BMC Systems Biology*, *2*(1), 51.

Scheible, W.-R., Krapp, A., & Stitt, M. (2000). Reciprocal diurnal changes of

phosphoenolpyruvate carboxylase expression and cytosolic pyruvate kinase, citrate synthase

and NADP☐isocitrate dehydrogenase expression regulate organic acid metabolism during

nitrate assimilation in tobacco leaves. *Plant, Cell & Environment*, *23*(11), 1155–1167.

Schiller, K., Janshoff, S., Zenker, S., Viehöver, P., Hartwell, J., Eirich, J., Finkemeier, I., &

Bräutigam, A. (2025). Regulation of Crassulacean acid metabolism at the protein level in

Kalanchoë laxiflora. *Plant Physiology*, *197*(4), kiaf095.

Schlüter, U., Bräutigam, A., Droz, J.-M., Schwender, J., & Weber, A. P. M. (2019). The role of

alanine and aspartate aminotransferases in C4 photosynthesis. *Plant Biology (Stuttgart,*

Germany), *21 Suppl 1*(S1), 64–76.

Seabra, A. R., Silva, L. S., & Carvalho, H. G. (2013). Novel aspects of glutamine synthetase

(GS) regulation revealed by a detailed expression analysis of the entire GS gene family of

Medicago truncatula under different physiological conditions. *BMC Plant Biology*, *13*(1),

137.

Snowden, C. J., Thomas, B., Baxter, C. J., Smith, J. A. C., & Sweetlove, L. J. (2015). A

tonoplast Glu/Asp/GABA exchanger that affects tomato fruit amino acid composition. *The*

Plant Journal: For Cell and Molecular Biology, *81*(5), 651–660.

Spalding, M. H., Stumpf, D. K., Ku, M. S. B., Burris, R. H., & Edwards, G. E. (1979).

Crassulacean acid metabolism and diurnal variations of internal CO2 and O2 concentrations

in Sedum praealtum DC. *Functional Plant Biology: FPB*, *6*(4), 557.

Szal, B., & Podgórska, A. (2012). The role of mitochondria in leaf nitrogen metabolism:

Mitochondria and N metabolism. *Plant, Cell & Environment*, *35*(10), 1756–1768.

Tahjib-Ul-Arif, M., Zahan, M. I., Karim, M. M., Imran, S., Hunter, C. T., Islam, M. S., Mia, M.

A., Hannan, M. A., Rhaman, M. S., Hossain, M. A., Brestic, M., Skalicky, M., & Murata,

Y. (2021). Citric Acid-Mediated Abiotic Stress Tolerance in Plants. *International Journal of*

Molecular Sciences, *22*(13), 7235.

Tcherkez, G., Mahé, A., Gauthier, P., Mauve, C., Gout, E., Bligny, R., Cornic, G., & Hodges, M.

(2009). *In folio* respiratory fluxomics revealed by 13C isotopic labeling and H/D isotope

effects highlight the noncyclic nature of the tricarboxylic acid "cycle" in illuminated leaves.

Plant Physiology, *151*(2), 620–630.

Timm, S., Florian, A., Arrivault, S., Stitt, M., Fernie, A. R., & Bauwe, H. (2012). Glycine

decarboxylase controls photosynthesis and plant growth. *FEBS Letters*, *586*(20), 3692–

3697.

Timm, S., Florian, A., Wittmiß, M., Jahnke, K., Hagemann, M., Fernie, A. R., & Bauwe, H.

(2013). Serine acts as a metabolic signal for the transcriptional control of photorespiration-

related genes in Arabidopsis. *Plant Physiology*, *162*(1), 379–389.

Tomato Genome Consortium. (2012). The tomato genome sequence provides insights into fleshy

fruit evolution. *Nature*, *485*(7400), 635–641.

Trípodi, K. E. J., & Podestá, F. E. (2003). Purification and characterization of an NAD-

dependent malate dehydrogenase from leaves of the crassulacean acid metabolism plant

Aptenia cordifolia. *Plant Physiology and Biochemistry*, *41*(2), 97–105.

Vogel, C., & Marcotte, E. M. (2012). Insights into the regulation of protein abundance from

proteomic and transcriptomic analyses. *Nature Reviews. Genetics*, *13*(4), 227–232.

Wai, C. M., Weise, S. E., Ozersky, P., Mockler, T. C., Michael, T. P., & VanBuren, R. (2019).

Time of day and network reprogramming during drought induced CAM photosynthesis in

Sedum album. *PLOS Genetics*, *15*(6), e1008209.

Wickell, D., Kuo, L.-Y., Yang, H.-P., Dhabalia Ashok, A., Irisarri, I., Dadras, A., de Vries, S., de

Vries, J., Huang, Y.-M., Li, Z., Barker, M. S., Hartwick, N. T., Michael, T. P., & Li, F.-W.

(2021). Underwater CAM photosynthesis elucidated by *Isoetes* genome. *Nature*

Communications, *12*(1), 6348.

Winter, K., Foster, J. G., Schmitt, M. R., & Edwards, G. E. (1982). Activity and quantity of

ribulose bisphosphate carboxylase-and phosphoenolpyruvate carboxylase-protein in two

Crassulacean acid metabolism plants in relation to leaf age, nitrogen nutrition, and point in

time during a day/night cycle. *Planta*, *154*(4), 309–317.

Winter, K., & Smith, J. A. C. (2022). CAM photosynthesis: the acid test. *The New Phytologist*,

233(2), 599–609.

Wu, H.-Y. L., Jen, J., & Hsu, P. Y. (2024). What, where, and how: Regulation of translation and

the translational landscape in plants. *The Plant Cell*, *36*(5), 1540–1564.

Yang, X., Hu, R., Yin, H., Jenkins, J., Shu, S., Tang, H., Liu, D., Weighill, D. A., Cheol Yim,

W., Ha, J., Heyduk, K., Goodstein, D. M., Guo, H.-B., Moseley, R. C., Fitzek, E., Jawdy,

S., Zhang, Z., Xie, M., Hartwell, J., … Tuskan, G. A. (2017). The *Kalanchoë* genome

provides insights into convergent evolution and building blocks of crassulacean acid

metabolism. *Nature Communications*, *8*(1), 1899.

Zhang, Y., Swart, C., Alseekh, S., Scossa, F., Jiang, L., Obata, T., Graf, A., & Fernie, A. R.

(2018). The Extra-Pathway Interactome of the TCA Cycle: Expected and Unexpected

Metabolic Interactions. *Plant Physiology*, *177*(3), 966–979.

1305