Genomic Architecture of the *Resistance to Phytophthora Cactorum 2* (*RPc2*) Locus in

Strawberry (*Fragaria × ananassa*)

Hyeondae Han[1#], Youngjae Oh[1,2#], Yoon Jeong Jang[1], Saket Chandra[1,3], Upinder Gill[1,4], Sujeet

Verma[1], Young-Hee Noh[1], Sachiko Isobe[5], Dominique D. A. Pincot[6], Glenn S. Cole[6], Randi A.

Famula[6], Mitchell J. Feldmann[6], Steven J. Knapp[6], Vance M. Whitaker[1] and Seonghee Lee[1*]

[1]Gulf Coast Research and Education Center, Institute of Food and Agricultural Sciences,

University of Florida, Wimauma, FL 33598, USA

[2] Department of Horticulture, Chungbuk National University, Cheongju 28644, Republic of

Korea

[3] School of Biotechnology, NIT Calicut, Kozhikode, Kerala 673601, India,

[4] Department of Plant Pathology, North Dakota State University, Fargo, ND 58102, USA

[5] Kazusa DNA Research Institute, Kisarazu, Chiba 292-0818, Japan

[6] Department of Plant Sciences, University of California, Davis, One Shields Avenue, Davis,

California 95616, USA

[*] Correspondence: Seonghee Lee, e-mail: seonghee105@ufl.edu

#Authors equally contributed to the manuscript.

Word counts:

Introduction: 1,217

Materials and Methods: 1,934

Results: 2,707

Discussion: 1,423

Number of Figures: 6, all figures need to be published in color.

Abstract

Phytophthora crown rot (PhCR), caused by *Phytophthora cactorum*, is a soilborne disease with broad impacts to cultivated strawberries in the United States and worldwide. While a resistance locus, *RPc2*, has been identified in octoploid strawberries, the underlying genomic architecture and mechanism remain unclear. Here, we fine-mapped the *RPc2* region to 546 kb, containing 92 genes, and constructed a chromosome-scale haplotype-phased genome of resistant breeding selection FL 16.33-8. Comparative genome analyses with high-quality octoploid reference genomes, 'Florida Brilliance' (FaFB1) and 'Royal Royce' (FaRR1), identified two candidate genes, *Wall Associated Kinase 1* (*WAK1*) and *Cyclic Nucleotide Gated Channel 1* (*CNGC1*). Gene functions were validated using an Agrobacterium-mediated transient expression assay. Furthermore, by leveraging the population structure of the *RPc2* locus among genetically diverse breeding populations, we unveiled the complexity of genetic architecture and recent selective sweeps associated with *RPc2*. Notably, the predominant resistant haplotype (*RPc2*-H3) is prevalent in most commercial strawberry varieties in the US. The findings from this study are facilitating the advancement of genome-assisted breeding strategies for resistance to PhCR in strawberries.

Summary

- Phytophthora crown rot (PhCR), caused by the soilborne pathogen *Phytophthora cactorum*, is a disease with broad impacts to cultivated strawberries worldwide. The *Resistance to Phytophthora Cactorum 2* (*RPc2*) locus is widespread in important cultivars and is a target of selection in breeding programs. Despite its importance, the genomic architecture and mechanisms of *RPc2* remain poorly characterized.

- We performed fine-mapping and delimited the genomic region to a 546 kb segment containing 92 genes. We developed a chromosome-scale haplotype-phased genome from an elite resistant line, FL16.33-8, and made comparisons to reference genomes of 'Florida Brilliance' (FaFB1) and 'Royal Royce' (FaRR1) to uncover gene sequence and structural variants in resistant accessions.

- Through comparative genomics and transcriptome profiling, we identified three candidate genes, *Wall Associated Kinase 1* (*WAK1*) and *Cyclic Nucleotide Gated Channel 1* (*CNGC1*) and *CNGC2*. The functions of these genes were validated using a simplified *Agrobacterium*-mediated transformation method for transient gene expression in strawberry crowns and roots.

- An examination of the population structure of the *RPc2* locus among genetically diverse accessions (N=1029) revealed that the predominant *H3* resistant haplotype is widespread in breeding accessions in the US, supporting the findings of a previous study on recent selective sweeps associated with *RPc2*. Results from this study are advancing genome-assisted breeding strategies for enhancing resistance to PhCR in strawberry.

Key words: Cyclic-nucleotide-gated calcium channels, fine-mapping, phased genome assemblies, QTL, wall associated kinase

Introduction

Phytophthora crown rot disease (PhCR) of cultivated strawberry (*Fragaria ×ananassa*) is caused by *Phytophthora cactorum*, a hemi-biotrophic, oomycete plant pathogen. PhCR is considered one of the most significant soilborne diseases affecting strawberries in the temperate regions of the world and causes production losses of up to 40% (Stensvand *et al.*, 1999). Soil fumigations using methyl bromide, dazomet, and chloropicrin 1,3-dichloropropene are effective ways to control *P. cactorum* (Cal *et al.*, 2004). However, increasing limitations on the use of fumigants have led to deterioration in the control of *P. cactorum* in recent years (Cal *et al.*, 2004; Holmes *et al.*, 2020). The development of resistant cultivars has become important for strawberry breeding programs and a critical strategy for combating PhCR in an era where fumigation alternatives are decreasing while global consumption of strawberries continues to increase.

The resistance locus *Phytophthora cactorum 1* (*RPc-1*) was initially reported in *F. vesca* (Davik *et al.*, 2015) but has not been found in cultivated strawberry. In a study conducted at UF, *FaRPc2* (hereafter referred to as *RPc2*) on chromosome 7B was identified using a pedigree-based analysis in complex and multiparent population sets (Mangandi *et al.*, 2017). It was described that two primary haplotypes, *RPc2-H2* and *RPc2-H3* (hereafter referred to as *H2* and *H3*) were associated with resistance against *P. cactorum*. Each resistant haplotype explained about 40% of the phenotypic variance in the tested populations (Mangandi *et al.*, 2017). These two resistant haplotypes seem to be originated from different sources and may represent distinct functional alleles. While all varieties with the H3 haplotype are resistant to PhCR, some susceptible varieties contain the H2 haplotype. This suggests that within the H2 haplotype, there may be very small allelic changes or sequence variants responsible for PhCR resistance. Although the exact cause is unknown, future studies should be conducted to elucidate the genomic region of *RPc2-H2*. he *H2* haplotype exhibits strong resistance to PhCR but is more poorly characterized.

Recently, Jiménez *et al.* (Jiménez *et al.*, 2023) reported minimal genetic gains in breeding for Phytophthora crown rot (PhCR) resistance, with only 3% of cultivars exhibiting high resistance in a genetically diverse training population. However, through multivariate genome-wide association studies (GWAS), multiple significant loci contributing to PhCR resistance were uncovered, notably including the *RPc2* locus. Also, the observed differences in phenotypic and

breeding values between UCD and non-UCD individuals in the training population align with

insights from previous genome-wide studies, indicating a decrease in genetic variation in the

UCD population due to strong directional selection, breeding bottlenecks, and selective sweeps

(Jiménez *et al.*, 2023). The result from this study also suggests a decline in resistance to PhCR in

the UCD population over the last half-century, with non-UCD individuals harboring both

favorable and unfavorable alleles not present in the UCD population or commonly found in gene

banks (Jiménez *et al.*, 2023). Following the searching of the genomic region near the SNP

marker AX-184109190 associated with *RPc2*, (Jiménez *et al.*, 2023) described several potential

defense related candidate genes for *RPc2* In another study from a European breeding group,

three resistance loci, *RPc6C, RPc6D,* and *RPc7D*, were identified using a bi-parental population

('Emily' × 'Fenella') (Nellist *et al.*, 2019). The *RPc7D* locus was shown to locate in the same

genomic region as *RPc2*, suggesting that these are same QTL.

The cultivated strawberry has a complex allo-octoploid genome (2n = 8x = 56),

consisting of four subgenomes with total genome size of 850 Mb (Hirakawa *et al.*, 2014;

Hummer *et al.*, 2011; Tennessen *et al.*, 2014). The first chromosome-scale reference genome was

assembled from 'Camarosa'(Edger *et al.*, 2019), and subsequent genomes were published from

the Japanese cultivar 'Reikou' (Shirasawa *et al.*, 2021) and a highly homozygous inbred line

'Wongyo3115' (Lee *et al.*, 2021). Recently, two fully haplotype-phased genomes were published

from a day-neutral cultivar, 'Royal Royce' (FaRR1) and short-day cultivar 'Florida Brilliance'

(FaFB1). 'Florida Brilliance' is susceptible to PhCR, and the FaFB1 genome does not possess

the resistant haplotype of *RPc2*. It has been reported that the cultivar 'Royal Royce' is

moderately resistant to Verticillium wilt (*Verticillium dahlia*), moderate susceptibility to

Phytophthora crown rot (*Phytophthora cactorum*) and Fusarium wilt (*Fusarium oxysporum*), and

susceptible to Charcoal rot (*Macrophomina phaseolina*) (https://itc.ucdavis.edu/wp-

content/uploads/UCD-Royal-Royce-Cultivar.pdf). To determine the genomic architecture of

RPc2, it will be necessary to construct a chromosome-scale haplotype-phased genome for a

resistant cultivar.

The molecular mechanisms underlying *RPc2*-mediated resistance against *P. cactorum* are

largely unexplored. Although a transcriptome analysis has been conducted in diploid strawberry

(*F. vesca*) challenged with *P. cactorum*, it is important to note that the regulation of defense

responses during the pathogen infection in octoploid strawberry appear to differ from that in *F.*

vesca due to diverse gene expression patterns across homoeologous copies (Toljamo *et al.*,

2016). Studies have highlighted the importance of genes encoding cell wall strengthening

proteins, particularly those involved in lignin biosynthesis enzymes associated with Damage-

Associated Molecular Patterns (DAMPs)-mediated defense pathways, in responding to

Phytophthora infection in plants. Additionally, there have been reports of the differential

expression of plant surface receptors, including 25 receptor-like proteins and 224 receptor-like

kinases, in tomato plants infected with *Phytophthora cactorum* (Zhou *et al.*, 2023). Jiménez *et al.*

(2023) described that the *RPc2* locus spanning 1.26 Mb contained 233 annotated genes with 18

of them associated with plant defense responses. Notably, these genes include intracellular

leucine-rich repeat (LRR)-type immune receptors, membrane-localized immune receptors, Ca2+-

dependent protein kinases, and cyclic-nucleotide-gated calcium channels (*CNGCs*), known to

play important roles in plant disease resistance. Moreover, a gene homologous to *WAK1*,

encoding a wall-associated receptor kinase galacturonan-binding protein, was highlighted as a

promising candidate (Jiménez *et al.*, 2023).

Despite the abundance of genetic information available for *RPc2*, there is currently no

study specifically characterizing causal variants, sequence structural variants, or testing of gene

function for this locus. Exploring these aspects could offer valuable insights into mechanisms of

resistance and facilitate the improvement of resistance to PhCR through genome-assisted

breeding approaches. In this study, we finely mapped the genomic region for *RPc2* using a

diverse mapping panel. A haplotype-phased genome of an elite resistant accession (FL 16.33-8)

was developed in order to characterize the resistant haplotypes and comparatively analyzed with

other publicly available high-quality haplotype-phased genomes of octoploid strawberry.

Furthermore, a genome-wide transcriptome profiling analysis highlighted two candidate genes,

Wall Associated Kinase 1 (*WAK1*) and *Cyclic Nucleotide Gated Channel 1* (*CNGC1*), that appear

to play pivotal roles in regulating resistance to PhCR in strawberry. Using an efficient

Agrobacterium-mediated transient gene expression method, we successfully validated the

functions of these candidates in the crowns and roots of strawberry plants. To understand

population genetic structure of the *RPc2* locus, we conducted a survey of globally diverse

breeding accessions (N=1029), revealing recent positive selection of the *H3* haplotype.

Materials and Methods

Genome sequencing and assembly

An elite resistant breeding selection FL16.33-8, which contains the major resistant haplotype

RPc2-H3 was identified for genome assembly. FL16.33-8 originated from a 2016 cross between

unreleased selection FL13.27-142 and FL12.93-4. FL13.27-142 was chosen as a resistant source

of *H3*. Etiolated young white leaf tissues were used for extracting high molecular weight DNA.

The single-molecule real-time sequencing (SMRT) bell library construction and Hi-Fi long read

PacBio DNA sequencing were conducted at DNA Link (Seoul, South Korea). The genome of FL

16.33-8 was assembled using the Hifiasm (Cheng *et al.*, 2021), which is a de novo assembler that

utilizes long high-fidelity sequence reads to represent the haplotype information in a phased

assembly graph. Two parents' Illumina paired-end (PE) data was utilized to phase haplotype

scaffolds. The genome continuity of the Illumina draft genome assemblies was assessed with

QUAST (Gurevich *et al.*, 2013). A BUSCO analysis was performed using default parameters to

assess the core gene set (Simão *et al.*, 2015). The de novo genome assembly, raw genomic reads

were used as input data for Ragtag (Alonge *et al.*, 2022), a tool designed for contig-ordering and

scaffolding genome assemblies. Genome quality and completeness was assessed using Merqury

ver 1.3 (Rhie *et al.*, 2020), which evaluates assembly based on efficient *k*-mer set operations. The

collinearity with the genetic linkage map for FL 16.33-8 and 'Florida Brilliance' was visually

inspected using Circa (https://omgenomics.com/circa/). The LTR Assembly Index for each

haplotype-phased genome was calculated based on the custom repeat library (Ou *et al.*, 2018).

To identify and classify repetitive elements in the genome, LTR retrotransposon

candidates were searched using EDTA (Su *et al.*, 2021). For the gene annotation, two haplotype-

phased assembly was annotated using Genome Sequence Annotation Server (Humann *et al.*,

2019). Two methodologies in parallel were used to mask and annotate the repeat library. A

library of predicted repeats detected with RepeatModeler (version 2.0.1; www.repeatmasker.org),

was combined with a library of repeats identified using RepeatMasker (version 4.1.1;

www.repeatmasker.org) and an *Arabidopsis thaliana* repeat library within GenSAS to create a

consensus library of repeats and mask polished assembly. Nucleotide sequences were aligned to

NCBI refseq plant using BlastN ver. 2.7.1 (Camacho *et al.*, 2009) as well as Diamond (Buchfink

et al., 2015). These alignments were combined with results of several gene prediction modeling

including AGUSTUS ver 3.3.1 (Hoff and Stanke, 2019) to generate an official gene set and

identify predicted transcripts within each masked assembly using EVidenceModeler (Haas *et al.*,

2008).

Scaffold validation using a genetic linkage map

A high-density genetic map was developed using a total of 169 F_1 individuals from a cross

between 'Florida Brilliance' and breeding selection FL 16.33-8. Axiom™ IStraw35 SNP arrays

were used to genotype all 169 F_1 individuals. Markers were filtered to have approach was

applied to identify singletons to fix the marker order and regions with low marker density or gaps

caused by segregation distortion. A final linkage map consisting of 10,269 SNP markers was

used to validate the scaffolds from the whole genome assembly. SNP probes sequences used for

the construction of linkage maps were mapped to the FL 16.33-8 assembly sequence using blastn

procedure. Alignments were filtered to retain markers if they matched to unique sequence

position in the phased genome assembly and with a maximum of 2 mismatches in the second-

best hit. The alignments were queried to detect potentially problematic scaffolds mapped with

SNP probes from different LGs. The number of scaffolds with SNP probes mapped from

different LGs was used as a metric in the quality assessment.

Fine-mapping

A total of 339 accessions with three seasons field phenotypic data (2013-2015) were used for

fine-mapping analysis (Supplementary Table S1). DNA of additional strawberry accessions was

extracted by the simplified CTAB (cetyltrimethylammonium bromide) method with minor

modifications (Keb-Llanes *et al.*, 2002; Noh *et al.*, 2017). To further fine map the *RPc2* region,

we developed sub-genome specific high-resolution melting (HRM) markers by selecting

FanaSNP array probes every 10 kb in the *RPc2* region, and selected HRM markers

discriminating H2 and H3 alleles through HRM analysis. The HRM markers representing 15 of

Axiom® IStraw90 SNP probes and 10 FanaSNP array probes encompassing *RPc2* region

(Hardigan *et al.*, 2020; Mangandi *et al.*, 2017; Noh *et al.*, 2018) were used to genotype 339

breeding accessions (Supplementary Table S2). Amplification products were resolved by high-

resolution melting analysis in LightCycler® 480 system II (Roche Life Science, Germany).

Melting curve data was analyzed by Melt Curve Genotyping and Gene Scanning software for

Roche LightCycler 480 II system.

RNA sequencing analysis

The plants were grown for four weeks in growth chamber under ideal conditions (73 ± 2°F,

Relative humidity 57 - 63%, 16 h light and 8 h darkness). Total of four breeding accessions were

used for RNA sequencing study: FL11.28-34 (H3:H1; heterozygous resistant haplotype),

FL12.75-77 (H3:H3; homozygous resistant haplotype), FL12.82-44 (H1:H1; homozygous

susceptible haplotype) and FL13.22-336 (H1:H1; homozygous susceptible haplotype). All

accessions were highly connected and comprises two haplotypes (H1 and H3), as each of them

has arose from elite breeding population and had undergone repeated selection for >15

generations (Mangandi *et al.*, 2017). The pathogen inoculation and spore concentration were

prepared according to the method as exactly mentioned in previous study (Mangandi *et al.*,

2017). Crown tissues were collected at 0, 24-, 48-, 72-, and 96-hour post inoculation (hpi) with at

least three biological replications.

For RNA extraction, crown tissues from each five time points were collected, and

samples were ground into fine powder using liquid nitrogen in mortar and pestle. Total RNA was

extracted according to the protocol of the Spectrum^{TM} Plant Total RNA Kit (Sigma-Aldrich, St.

Louis, MO, United States). To remove any traces of DNA, the isolated RNA was treated with

DNAse I (Invitrogen) and resuspended in a total volume of 50 μl of RNase-free water. In total of

1 μg total RNA was used for first-strand cDNA synthesis using Transcriptor First Strand cDNA

synthesis kit (Roche, Switzerland) following the manufacturer's instructions. The RNA samples

of 0 and 96 hpi were selected for RNA-seq study. The paired-end library construction and

sequencing was conducted on the Nextseq 500 Illumina sequencing platform (Novogene, USA).

Illumina adapters were removed from the raw reads using Trimmomatic ver. 0.33 (Bolger *et al.*,

2014). After quality assessment with FastQC (https://www.bioinformatics.babraham.

ac.uk/projects/fastqc/), the filtered reads were then mapped to each haplotype-phased genome

assembly using HISAT2 (Kim *et al.*, 2019). The sequence alignment map (SAM) format was

converted to Binary Alignment Map (BAM) format. To import the BAM files into R,

'Rsamtools' package was used. Differentially expressed genes in resistant vs susceptible were

identified using the DESeq2 pipeline. DESeq2 software was employed to calculate the

expression between genotypes (Love *et al.*, 2014). Pairwise comparison was performed between

H1-water control (WC) vs. *H1*-pathogen inoculation (PI), *H1*-PI vs. *H3*-PI, *H3*-WC vs. *H3*-PI,

and total resistance *vs.* total susceptibility. The genes with a | log2 fold change| $\geq$ 1 and FDR-

adjusted *p*-value (q-value) < 0.05 were selected as DEGs. Heatmap was generated using R

package 'gplots' using expression data for genes locating at interval of *RPc2*. The differentially

expressed sequences were annotated with the help of sequence homology search using blastx

program in BLAST+ package (Camacho *et al.*, 2009). The sequences were analyzed at an e-

value of $1e^{-6}$ to NCBI non-redundant protein database. The homologues were selected on the

basis of the e-value of less than $1e^{-6}$ and highest high-scoring segment pairs (HSP).

Comparative genomic data analysis

For the comparative genomic analysis of the two different haplotypes of *H1* and *H3*, we selected

the 546 kb *H3* region identified from fine-mapping. This region was extracted from the genomes

of 'Royal Royce' (homozygous *H3*), 'Florida Brilliance' (homozygous *H1*), and the phase 1 and 2

assemblies of FL16.33-8 (*H3*), respectively. Genomic sequences were compared with Mauve

software in the Geneious Prime software 2020.2.4. The nucleotide and amino acid sequences of

candidate genes, *WAK1*, *CNGC1*, and *CNGC2*, were compared using MAFFT alignment in

Geneious Prime software version 2020.2.4.

Agrobacterium-based transient assay

To validate candidate gene function in strawberries, crown and root transient overexpression and

RNAi transient assays were performed using *A. tumefaciens*, with modifications to the previous

method (Zhong *et al.*, 2016). We used three-week-old strawberry daughter plants of 'Florida

Brilliance' (susceptible) for overexpression transient assays and 'Florida Beauty' (moderately

resistant) for RNAi transient assays. The pMDC32 and pK7GWIWG(II) gateway vector systems

were used for overexpression and RNAi transient assays (Curtis and Grossniklaus, 2003; Karimi

et al., 2002). The final cloning products of pMDC32::*WAK1*, pMDC32::*CNGC1*, pK7::*WAK1*,

and pk7::*CNGC1/CNGC2* were introduced *A. tumefaciens* EHA105.

Transformed *A. tumefaciens* were grown at 28°C overnight in LB medium containing

antibiotics appropriate for each vector. When the culture density reached to OD_{600} of 1.0, *A.*

tumefaciens were harvested by centrifuge at 4500rpm for 25min and resuspended in adjusted to

an OD_{600} of 0.6 in MMA activation buffer (10mM 2-N-morpholino) ethanesulphonic acid

(MES), 10mM $MgCl_2$, pH5.6 and 200 μM acetosyringone) and shaken 50-70rpm for 3hr at room

temperature before infiltration of root and crown. To enhance the transformation efficiency, we

applied the vacuum infiltration method during the overexpression and RNAi transient assays.

Briefly, roots and crowns were gently scraped with a syringe needle before vacuum infiltration,

then transferred to a suspension *A. tumefaciens* with 0.05% silwet L-77. Vacuum infiltration was

conducted for five minutes. Subsequently, the inoculated plants were soaked in the same solution

for three hours at room temperature. Afterward, the plants were transferred to 4-inch pots in a

humid box in a growth room for four days (12/12hr light/night, 23-25°C). Four days later, the

pathogen inoculum (zoospore suspension with a final concentration of 1×10^4 zoospores/ml)

was prepared according to the method in previous study (Mangandi *et al.*, 2017). Inoculations

were performed by immersing the plant roots for five seconds in 500 ml of the zoospore

suspension up to the base of the crown immediately before transferring to pots. The RNAi and

overexpression experiments were conducted twice over time. Each experiment was conducted

with three biological replicates with empty vector controls. After 17 days post-infection, plant

crowns were halved, and photos were taken from each plant. The percentage of symptomatic

areas was assessed using ImageJ software to quantify the areas showing typical *P. cactorum*

symptoms on the crown and roots, including the areas of halved crowns and the internal

symptomatic areas. Subsequent statistical analysis was performed using R software, conducting

One-way ANOVA, and Tukey's honestly significant difference (HSD) test was used to compare

the means of all treatments to each other and statistically validate the data ($p \leq 0.05$). The target

genes were *FaWAK* and *FaCNGC*, while *FaGAPDH2* was selected as housekeeping gene

control. Primers were designed using Primer 3 (https://primer3.ut.ee/) tool, and all primer

sequences were represented in Supplementary Table S3. The qRT-PCR experiment was

performed using LightCycler® 480 system (Roche Diagnostics, Mannheim, Germany) using

Forget-Me-Not™ EvaGreen® qPCR Master Mix (Biotium, CA, USA).

Population structure analysis and phylogenetic analysis

The 1,029 octoploid accessions (Supplementary Dataset2 and 3). A total of 17 SNP markers in

the 546 kb of *RPc2* region genotype matrices were used to evaluate population structure by

principal component analysis and clustering with an admixture model using STRUCTURE

(v.2.3.4) (Pritchard *et al.*, 2000). The STRUCTURE analysis tested for K=2-10 subpopulations

(10,000 burn-in steps and 10,000 Markov-Chain Monte Carlo steps with 10 replicates per K

value). The optimal subpopulation (K) value was determined using STRUCTURE HARVESTER

(v0.6.94) (Earl and VonHoldt, 2012). The results were visualized using pophelpher (Francis,

2017). Principal component analysis was conducted using TASSEL5 (Bradbury *et al.*, 2007).

Cladogram was generated using fastTree (Price *et al.*, 2009) on Geneious Prime. After estimating

genetic distance matrices were generated, a maximum-likelihood phylogenetic tree was

constructed.

Results

Fine-Mapping of *RPc2* Conferring Resistance to Phytophthora Crown Rot in

Octoploid Strawberry

In our previous investigation, we mapped *RPc2* to a 1.12 Mb segment using 15 SNP markers

derived from Axiom® IStraw90 SNP probes (Noh *et al.*, 2018) (Fig. 1A, Supplementary Tables

S1 and S2). To refine this interval for effective marker-assisted selection in resistance breeding,

we assembled an association mapping panel consisting of 339 accessions. This panel comprised

206 accessions used in the previous study for QTL analysis (Noh *et al.*, 2018) and an additional

133 accessions selected based on consistent field data for mortality due to PhCR collected over

three seasons (Supplementary Table S1). Employing these association mapping panels, family-

based linkage studies were performed to identify within-family recombination events. To further

enhance the resolution of the identified interval, 15 sub-genome-specific SNP or InDel-based

high-resolution melt (HRM) markers were employed, including four newly developed markers

using the 50K FanaSNP Array (Hardigan *et al.*, 2021) (Fig. 1A, Supplementary Table S1). The

genotyping of all 339 accessions with these 15 HRM markers revealed recombination events at

markers AX-89845076 and AX-184173618, resulting in a refined *RPc2* region reduced from 1.12

Mb to 546 kb (Fig. 1A, Supplementary Table S1). The markers between AX-18423389 and AX-

184173618 did not accurately predict the phenotype of some individuals containing the H2

haplotype.

Fig. 1. Fine-mapping of the *RPc2* genomic region conferring resistance to Phytophthora

crown rot resistance. (A) Preliminary mapping of *RPc2* region spans approximately 1.12 Mb

and is based on 11 markers (black) derived from Axiom® IStraw90 SNP probes (Mangandi et

al., 2017). Using the additional subgenome-specific HRM markers (blue) derived from FanaSNP

50k array, *RPc2* was delimited to 546 Kb. (B) HRM marker genotype results (N: negative, P:

positive) associated with the resistance haplotype, *RPc2-H3*; (a) showcases the diplotypes of

accessions in the *RPc2* region from Mangandi et al. (2017). *H1* and *H2*: susceptible haplotype,

H3: resistance haplotype (b) is based on the area under the disease progress curve (AUDPC)

values reported in Mangandi et al. (2017), with R: resistance, MR: moderate resistance, HS: high

susceptibility, S: susceptibility.

Haplotype-Phased Genome Assembly and Annotation of Resistant Breeding Selection 'FL 16.33-8'

The trio-binning method was applied to develop a high-quality haplotype-phased genome of the resistant accession FL 16.33-8 using PacBio HiFi long reads and parental short reads. The contig N50 values for FL 16.33-8-Phase1 and FL 16.33-8-Phase2 were 16.4 Mb and 14.3 Mb (Table 1). The assembly spans a total length of 782.4 Mb for FL 16.33-8-Phase1 and 778.1 Mb in FL 16.33-8-Phase2, distributed across 56 sub-chromosomal pseudomolecules (Supplementary Table S4). These pseudomolecules were numbered following the convention of reference genomes from 'Camarosa', 'Royal Royce'- FaRR1, and 'Florida Brilliance' - FaFB1 (Fig. 2A, Supplementary Table S4). In FL 16.33-8-Phase1, the shortest and longest chromosomes were chromosome 7D (19.35 Mb) and chromosome 6B (43.16 Mb), while in FL 16.33-8-Phase2, they were chromosome 7C (21.75Mb) and chromosome 6C (34.25 Mb) (Supplementary Table S4). Genome assembly and annotation completeness showed 98.2% of complete BUSCO in both FL 16.33-8-Phase1 and FL 16.33-8-Phase2, indicating a high-quality chromosome-scale assembly (Supplementary Table S5).

To validate the accuracy of haplotype phasing in the genome assembly, we employed a high-density genetic linkage map developed from a cross of 'Florida Brilliance' by 'FL 16.33-8'. The genetic map consisted of 13,910 SNP markers at 10,269 distinct positions across 31 linkage groups (LGs) with a total genetic length of 2,382.03 cM (Supplementary Table S6; Supplementary Dataset 1). The 31 LGs from LG1A to LG7D showed synteny with one of the 28 sub-chromosomal pseudomolecules from the assembly with less than 0.5% of markers exhibiting conflicts with their positions in the assembled chromosomes (Fig. 2B, Supplementary Table S6). Sub-genome specific FanaSNP markers used in constructing the linkage map were anchored to sub-chromosomal pseudomolecules of the reference genome and assigned to each subgenome from chromosome 1-1 or 1A to chromosome 7-1 or 7D based on nomenclature of 'Camarosa' or 'FaRR1' reference genomes (Fig. 2B). Trio binning with hifiasm (v0.12) successfully classified both haplotypes of FL 16.33-8, resulting in k-mer distributions for the read bins and an assembly that fully resolved both parental haplotypes (Fig. 2C). This observation was further confirmed by the spectra plot for the combined assembly, where homozygous regions predominantly contained 2-copy k-mer, and heterozygous regions mostly contain 1-copy k-mer. This distribution aligns

with expectations based on the presence of both complete parental haplotypes and minimal

artefactual duplication (Supplementary Fig. 1). Additionally, conserved synteny blocks between

F. vesca and FL 16.33-8 genomes were confirmed (Fig. 2D). When comparing three putative

ancestral diploid strawberry genomes, *F. vesca*, *F. iinumae*, and *F. viridis*, the haplotype-phased

genome of FL 16.33-8 showed a high level of collinearity with all diploid strawberry genomes

(Supplementary Figs. S2 and S3).

Table 1. Statistics on assembled haplotype-phased genome of a resistant accession, FL 16.33-8.

Metrics	Value	
	Phase 1	Phase 2
Number of contigs (>= 10,000 bp)	1,468	592
Number of contigs (>= 25,000 bp)	1,447	587
Number of contigs (>= 50,000 bp)	637	456
Total length (>= 10,000 bp)	869,351,047	826,921,887
Total length (>= 25,000 bp)	868,872,493	826,804,304
Total length (>= 50,000 bp)	837,337,314	821,510,393
Number of contigs	1,468	592
Largest contig	27,317,239	27,253,556
Total length	869,351,047	826,921,887
GC (%)	40.1	39.99
N50	16,418,230	14,284,693
N75	10,170,993	9,465,336
L50	23	22
L75	39	40
Number of N's per 100 kbp	0	0

The total TE content of haplotype-phased genomes of FL 16.33-8 were 345 Mb in 16.33-8-

Phase1 and 343 Mb in 16.33-8-Phase2, including long terminal repeat (LTR) retrotransposons

(Class I: 55.95% and 58.53%) and terminal inverted repeat (TIR) elements (Class II: 31.59% and

27.02%), respectively. TEs were evenly distributed across the 28 chromosomes of FL 16.33-8

with a small increase in content near the centromeres (Fig. 2A and Supplementary Table S7). The

most abundant Class I TEs were LTR retrotransposons, particularly the *Gypsy* superfamily

followed by *Copia*. Mutator was the most abundant among Class II TEs. Comparing TEs in both

16.33-8-Phase1 and 16.33-8-Phase2 revealed differences between each super-families, indicating

a high level of divergence in TE families between the haplotype genomes. This divergence

resulted in structural and sequential variations in protein coding genes and their regulatory

sequences. The genome quality was further assessed by examining assembly continuity in repeat

space using the LTR Assembly Index (LAI) (Ou *et al.*, 2018). The adjusted LAI score of each

16.33-8-Phase1 and 16.33-8-Phase2 are 15 and 14, which fall within the "reference" standard

range according to the LAI classification. Estimation of regional LAI in 1 Mb sliding windows

demonstrated uniform and high-quality assembly continuity across the entire genome (Fig. 2A).

Fig. 2. The haplotype-phased genome assembly of an elite resistant line, FL 16.33-8

containing the resistance haplotype *H3*. (A) Genome visualization of haplotype-phased

genome A of FL 16.33-8. The tracks from outside to inside are: a = haplotype-phased subgenome

name, b = length (Mb) of sub-genome, c = number of genes per 1 Mb window, d = density of

long term repeated (LTR) per 1 Mb window, e = LTR assembly index (LAI) in sub-genomes

from 1A to 7D plotted in 1 Mb window, f = The indel position differing by more than 30,000 bp

among 'Royal Royce', 'Wonkyo 3115', and FL 16.33-8, g = synteny comparison within the haplotype-phased genome 1. (B) Comparison of the collinearity between a genetic linkage map and haplotype-phased genome 1 of FL 16.33-8. Genetic positions from a linkage map for 'Florida Brilliance' × FL 16.33-8 were plotted against the corresponding physical positions in the 'Florida Brilliance' genome assembly, FaFB1 (blue points). The X-axis indicates the genetic linkage map in centimorgans (cM), while the Y-axis represents genome coordinates in the haplotype-phased genome A of FL 16.33-8. (C) The k-mer blob Plot illustrates the distribution of maternal (red) and paternal (blue) k-mer within each scaffold of the respective assemblies. The size of the blob corresponds to the total number of k-mers in that scaffold. (D) Dotplot of FL 16.33-8-phase1 assembly against diploid *Fragaria vesca* ver. 4.0. Dot plots are produced using the DGENIE software and alignments with minimap2.

Comparative Transcriptome and Genome Analysis Revealed Candidate Genes for the *RPc2*-Associated Resistance Against *P. cactorum*

Genome-wide transcriptome profiling analysis was conducted on two highly resistant (FL 12.75-77 and FL 11.28-34) and two highly susceptible accessions (FL 12.82-44 and FL 13.22-336) 96 hours after *P. cactorum* infection (Fig. 3A and Supplementary Table S8). The whole genome heatmap of differentially expressed gene (DEGs) revealed significant patterns of gene expression between the resistant (*H3*) and susceptible (*h3*) accessions post-pathogen infection (PI) (Fig. 3A, Supplementary Table S9). Infected roots were observed in both resistant and susceptible accessions after 96 hours of inoculation, but interestingly, newly developed roots were more frequently observed in resistant accessions compared to susceptible ones. Additionally, more severe disease symptoms were observed in the roots of susceptible accessions (Fig. 3A). A total of 409 DEGs (gray) were upregulated in resistant accessions after the pathogen infection (Fig. 3B and Supplementary Table S10). Notably, among the 34 DEGs identified from the comparison of resistant and susceptible accessions (*H3*-P.c Inoculation vs. *h3*-P.c Inoculation; gray ∩ orange), three DEGs (Fxa7Bg964820, Fxa7Bg964830, and Fxa2Bg234610) and four DEGs (Fxa3Ag419260, Fxa2Cg141260, Fxa2Dg203300, Fxa2Bg262750) were found to be closely associated with plant defense signaling and lignin biosynthesis pathways (Supplementary Fig. S4 and S5). In the volcano plot analysis of whole-genome DEGs, we identified distinct gene expression patterns. The two *cyclic nucleotide-gated ion channel 1-like* (*CNGCs*) genes: Fxa7Bg964820 and Fxa7Bg964830 were the most significantly upregulated in response to the

pathogen inoculation. (Fig. 3C). There were number of upregulated DEGs with up to 10-fold

changes including *WAK1*. In this study, we focused on *CNGC1*, *CNGC2*, and *WAK1*, as these

DEGs are located in the *RPc2* region. Global analyses of DEGs in each pairwise comparison

demonstrated significant alterations in gene expressions associated with 'Binding' and 'Catalytic

Activity' across all comparisons following the inoculation of *P. cactorum* (Supplementary Fig.

S6).

Fig. 3. **Differentially expressed genes associated with the resistance against *P. cactorum*.**
(A) Heatmap showing differentially expressed genes at the whole-genome level in response to
inoculation with an oospore suspension. Disease symptoms were observed in the roots of
resistant and susceptible genotypes 96 hours post-inoculation. (B) Venn diagram of upregulated
DEGs in the water and oospore treatments between susceptible and resistance accessions. (C)
Heatmap visualization of the genes present in the fine-mapped region (546 kb) of *RPc2*. Red
box: susceptible genotypes, Blue box: resistant genotypes. The genes highly upregulated in the
resistance genotype compared to the susceptible genotype, were highlighted in yellow. (D)
Volcano plot comparing gene expression levels between susceptible and resistance accessions
after infection. *CNGC1*, *CNGC2*, and *WAK1* are marked with purple triangle.

The transcriptome expression profiling was conducted on 92 genes located within the

genomic region of the resistant haplotype, *H3*, in response to *P. cactorum* infection. Two *cyclic*

nucleotide-gated ion channel 1-like (*CNGC1* and *CNGC2*; Fxa7Bg964820, Fxa7Bg964830)

genes and *One Wall-Associated Kinase 1-like* (*WAK1*; Fxa7Bg964800) were constitutively

upregulated in the resistant accessions compared to susceptible accessions (Fig. 3C,

Supplementary Table S11). To identify genomic structure and causal variants associated with

RPc2, the resistant haplotype genome, 16.33-8-Phase1 containing *H3*, was comparatively

analyzed with 'Florida Brilliance' (FaFB1) and 'Royal Royce' (FaRR1). The genomic region of

the resistant haplotype *H3* from FL 16.33-8 closely resembled that of 'Royal Royce', suggesting

that 'Royal Royce' possesses the resistant haplotype *H3*. On the other hand, there were

significant structural and sequence variations between the genomes of resistant (16.33-8-Phase1,

FaRR1-Phase1 and FaRR1-Phase2) and susceptible (FaFB1 both haplotypes) accessions (Fig.

4A). The sequence variations of candidate genes *WAK1*, *CNGC1*, and *CNGC2* were specifically

examined. The amino acid sequence of *WAK1* from 16.33-8-Phase1 was identical with both

FaRR1 haplotypes and was 91.25% consistent with the *WAK1* in FaFB1 (Supplementary Fig.

S7). Two copies of the *CNGC* gene (*CNGC1* and *CNGC2*) were exclusively identified in the

genomes of 16.33-8-Phase1 and FaRR1, both carrying the resistance haplotype *H3*. The amino

acid sequence similarity between CNGC1 and CNGC2 was 98.67% (Supplementary Fig. S8).

Two homoeologous copies of WAK1 were found in chromosomes 7C and 7D, sharing 86%

similarity. However, the amino acid sequences of WAK1 present in chromosome 7B were 30%

consistent with a copy on chromosome 7C and 45% consistent with another copy on

chromosome 7D (Supplementary Fig. S9). Two homoeologous copies of *CNGC1* and *CNGC2*

were present in chromosomes 7A and 7D. The amino acid sequences of CNGC1 were 84.58%

consistent with homoeologous copy on chromosome 7A, and 62.93% consistent with another

homoeologous copy on chromosome 7D (Supplementary Fig. S10).

To examine the functions of *WAK1*,*CNGC1*, and *CNGC2*, we conducted *Agrobacterium*-

mediated transient gene expression assays for RNAi-mediated knockdown and overexpression

(Figs. 4D, and E). Due to the high sequence identity between *CNGC1* and *CNGC2*, it was not

possible to target only one gene individually. Therefore, we designed the RNAi construct to

knockdown the expression of both genes. For the overexpression study, however, we specifically

cloned *CNGC1* into the overexpression vector to assess its effects for PhCR phenotype in a

susceptible variety. The qRT-PCR results showed a significant downregulation of *WAK1* and

CNGC1/CNGC2 gene expression levels in RNAi plants and an upregulation in overexpression

plants compared to the empty vector controls (Supplementary Fig. S11). The differences in

disease symptoms between the RNAi and overexpression treatment groups were assessed using

one-way ANOVA. The RNAi-mediated knockdown of *WAK1* and *CNGC1/CNGC2* in the

moderately resistant 'Florida Beauty' (*H3*) displayed significantly increased symptomatic area

compared to the control plants (empty vector) against *P. cactorum* (Figs. 4D; Supplementary Fig.

11B). The overexpression of *WAK1* and *CNGC1* in susceptible 'Florida Brilliance' (*h3*) resulted

in 27.3% and 20.7% reductions in symptomatic area of crown respectively compared to control

plants with empty vector treatments following *P. cactorum* infection (Figs. 4E; Supplementary

Fig. 11A; Supplementary Table S12). However, the overexpression of *CNGC1* appears to be

statistically less significant compared to the control plants (empty vector), likely due to

variations between replicated experiments. The *WAK1*-RNAi plants shown in Fig. 4D are from

the moderately resistant 'Florida Beauty' PhCR, while Fig. 4E features the susceptible 'Florida

Brilliance' (*h3*) group for *WAK1* and *CNGC1*. These plants represent different genotypes, leading

to distinct phenotypes. In Fig. 4D, *WAK1*-RNAi knockdown plants display intermediate

browning in the crown and a lack of new bare roots, indicating a partial compromise in

resistance. Conversely, Fig. 4E shows 'Florida Brilliance' with browning starting at the crown

and also a failure to produce new bare roots, highlighting its susceptibility. Taken together, the

results from comparative genomic and transcriptomic analyses indicated *WAK1*, *CNGC1*, and

CNGC2 as candidate genes involved in *RPc2*-mediated resistance to PhCR in strawberry.

Fig. 4. Identification of candidate genes for *RPc2* and functional validations in strawberry crowns and roots. (A) FL 16.33-8 phase1, 'Royal Royce' FaRR1 phase1 and phase2 represent the structure of resistance haplotype *RPc2*-H3, and 'Florida Brilliance' FaFB1 represents the structure of susceptible haplotype *RPc2*-H1 (referred to as *rpc-h3*). Mauve alignment of genomic region of *RPc2* between 'Florida Brilliance' (FaFB1), FL 16.33-8 phase1 and 2, 'Royal Royce'

(FaRR1) phase1 and phase2. Each color indicates the locally collinear blocks. Same color of
blocks represents local collinearity. Molecular marker linked to the candidate genes is indicated
with a blue arrow. (B) Physical locations of WAK1, CNGC1, and CNGC2 in genomes of
'Florida Brilliance' and FL 16.33-8. (C) Sequence comparisons were conducted for WAK1
between resistant and susceptible varieties. Additionally, CNGC1 and CNGC2 sequences present
in FL 16.33-8 Phase 1 genome were compared each other. Different functional domains and
motifs were marked and described separately using different colors. 3D protein structures of
WAK1, CNGC1, CNGC2 from FL 16.33-8, and WAK1 from 'Florida Brilliance'. "(D)
Agrobacterium-mediated transient gene expression of *WAK1* and *CNGC1/CNGC2* for RNAi-
mediated knockdown in the moderately resistant 'Florida Beauty' (RPc2-H3). (E) overexpression
of *WAK1* and *CNGC1* in susceptible 'Florida Brilliance' (*rpc2-h3*).
547

Population structure and selection signatures of the *RPc2* locus among genetically

diverse strawberry accessions

We assessed the population structure of *RPc2* across diverse breeding populations (UCD: n=501,

UF: n=61, other USA: n=267, Canada: n=42, Europe: n=91, Japan: n=46, Taiwan: n=3, Chile:

n=6, Ecuador: n=1). Comparative genetic structure and principal component analysis (PCA)

were performed using 76 SNP markers spanning the genomic region of *RPc2*, approximately 1.2

Mb. The results of the PCA showed that the accessions were segregated into four distinct groups,

based on the presence of either the resistant or susceptible haplotype of *RPc2* (Fig. 5A,

Supplementary Dataset2, Supplementary Table S13). Additionally, the dosage of resistant

haplotype (*H3*) or susceptible haplotype groups (*h3*) was clearly discriminated by the PCA

analysis. The group of susceptible haplotypes (*h3*) clustered in the intermediate region between

homozygous- and heterozygous-*RPc2-H3*. As shown in Figure 5b, the phylogenetic analysis of

1,029 accessions largely categorized them into the groups with and without the resistant

haplotype *H3* (Supplementary Dataset 3; Fig. 5B).

Bayesian inference of population structure was also performed. The optimal number of

clusters (K) was determined to be k=4, revealing 340 accessions as homozygous *RPc2-H3*

(cluster 2) and 376 as heterozygous *RPc2-H3* (cluster 1 and cluster 4) (Fig. 5C, Supplementary

Fig. S12, Supplementary Table S14). The cluster 1 (*RPc2-H3/rpc2-h2*) and cluster 4 (*RPc2-

H3/rpc2-h1*) mainly comprise accessions with the heterozygous resistant *RPc2-H3* haplotype

with the susceptible haplotypes. Investigation of the geographical origin revealed that relatively

high frequency of resistant haplotype *RPc2-H3* was present in the US strawberry breeding

accessions, respectively (Supplementary Table S15). This finding indicates that resistant *RPc2*

alleles have undergone positive selection during strawberry breeding in the US. Additionally, the

resistant haplotype *RPc2-H3* is well preserved in 829 (80.5%) accessions, with 60 out of 91

cultivars bred in Europe possessing *RPc2*-H3. This distribution supports the notion of a selective

sweep driven by continuous selection of *P. cactorum*-resistant individuals in breeding programs

worldwide.

Fig. 5. Genetic structure analysis of octoploid strawberry populations using 76 SNP

markers in the 1.2 Mb of *RPc2* region. (A) Principal component analysis that shows different

genetic clusters with and without the resistance haplotype, *RPc2-H3*. (B) Phylogenetic analysis

revealed three groups characterized by *RPc2-h1*, *RPc2-h2*, and *RPc2-H3*. (C) Admixture analysis

using Structure ver. 2.3.4 created four clusters with and without the resistance haplotype *RPc2-*

H3.

Discussion

In previous studies, a large-effect locus, *RPc2*, for resistance to crown rot caused by *P. cactorum* was initially identified in the UF breeding program through pedigree-based QTL detection (Mangandi et al., 2017). Subsequent multivariate genome-wide association studies (GWAS) confirmed the significance of the locus on chromosome 7B, describing a substantial portion of genetic variance and suggesting its necessity, though not sufficiency for resistance to PhCR (Jiménez *et al.*, 2023). Despite its importance, this genomic region has not been fully characterized; thus this study highlights the need for further research to uncover potential resistance mechanisms and facilitate more effective breeding strategies to control PhCR.

This study developed a high-quality phased genome for the resistant accession FL16.33-8 and compared the *RPc2* locus across multiple genomes, revealing shared origins of the resistant haplotype between California and Florida breeding programs. Building upon previous studies (Lin *et al.*, 2023; Lozada *et al.*, 2021; Wang *et al.*, 2021), the high-quality genome information would provide a foundation for identifying candidate genes associated with resistance traits and for elucidating the molecular mechanisms underlying the resistance to the soilborne pathogen. In this study, we identified two *Cyclic Nucleotide-Gated Channels* (*CNGC*) and one *Wall-Associated Kinase 1* (*WAK1*) gene for the resistance of PhCR.. Previous studies have highlighted the significance of *CNGC* genes in plant defense against pathogens (Yoshioka *et al.*, 2006). It was also mentioned that calcium channel and other candidate genes with known plant defense functions could be associated with the *RPc2* resistance to PhCR (Jiménez *et al.*, 2023). Similar findings have been reported in other plant species, where overexpression of *WAK* and *CNGC* genes conferred enhanced resistance to various pathogens (He *et al.*, 1998; Navarro *et al.*, 2004). These results underscore the importance of *WAK1* and *CNGC1* as key genetic factors in *RPc2*-mediated resistance in strawberries and provide valuable insights into the molecular mechanisms underlying resistance to *P. cactorum*.

The *WAK* play vital role in expansion of cell, protection against heavy metal and imparts resistance against plant-pathogens (He *et al.*, 1998; Hou *et al.*, 2005; Lally *et al.*, 2001; Tang *et al.*, 2017; Verica and He, 2002b). Protein kinase has also been reported to have function related to signaling during osmotic stress, keeping plant-specific cell wall and cytoskeleton continuum and also helps in closure of stomata by combining the environmental stimulus with cellular

action with the help of H_2O_2 (Baluška *et al.*, 2003; Desikan *et al.*, 2004; Urao *et al.*, 1999). These

reports indicate that *WAK* has an important role against fungal pathogen inside the root cortex by

initiating the defense response by modulating cytoskeleton-cell wall border. Also, *Phytophthora*

species exudes hydrolase that damages the plant cell wall and leads to the degradation of

homogalacturonan that ultimately releases Oligogalacturonides and these further binds with *WAK*

to initiate defense response (Ferrari *et al.*, 2013). The upregulation of *WAK* in response to

Phytophthora has also been found in previous studies (Burra *et al.*, 2014; Khatib *et al.*, 2004;

Toljamo *et al.*, 2016; Xiao *et al.*, 2019). *WAK* genes are receptor-like kinases localized in the cell

wall, involved in recognizing pathogen-associated molecular patterns (PAMPs) and activating

downstream defense responses (Supplementary Fig. S13) (Anderson *et al.*, 2001; Verica and He,

2002a). Upon recognition of fungal PAMPs, *WAK* genes initiate signaling cascades that lead to

the activation of various defense mechanisms, including the production of reactive oxygen

species (ROS), the induction of pathogenesis-related (PR) genes, and the reinforcement of cell

walls through callose deposition (Brutus *et al.*, 2010; Kohorn and Kohorn, 2012).

During the sample collection of RNA-seq experiment, we often observed new root

initiation near the crown in resistant plants after pathogen inoculation, while susceptible plants

showed more severe root damage and lacked new root development near the crown as shown in

Figure 3A. This suggests a potential role for *CNGC* and *WAK* in promoting root growth and

development, particularly in response to soilborne pathogen infection, in resistant plant

genotypes. MapMan analysis for the DEGs displayed upregulated genes involved in cell wall,

phytohormone jasmonic acid, and wax in the resistant individuals in response to the pathogen

(Fig. 6). Interestingly, the key genes in lignin biosynthesis are upregulated after the infection of

P. cactorum, suggesting the important role of lignin in resistance of PhCR in strawberry. This

observation has been also reported in Arabidopsis exposed to *Xanthomonas campestris* and flax

(*Linum usitatissimum*) cell cultures treated with *Botrytis cinerea* where the expression of genes

involved in lignin biosynthesis were observed (Hano *et al.*, 2006; Lauvergeat *et al.*, 2001;

Zimmermann *et al.*, 1999). The phytohormone jasmonic acid (JA), which plays an important role

as a signaling molecule in plant defense mechanisms, was also upregulated in resistant plants

(Campos *et al.*, 2014).

Fig. 6. The MapMan-based classification of differentially expressed genes (DEGs) in response to *P. cactorum*. (A) Expression profiles of DEGs involved in biotic and abiotic stresses. (B) Expression profiles of DEGs involved in secondary metabolism. The scale bar represents positive (red) and negative (blue) regulation of gene expression based on log 2 fc scores.

CNGCs are known to have role in transport of ions, like Ca^{2+}, through the plasma membrane and helps in regulating the defense signal (Deising, 2009). *CNGCs*, on the other hand, are plasma membrane-localized ion channels that regulate calcium influx in response to pathogen perception (Ma *et al.*, 2009). Calcium signaling is a central component of plant defense signaling, triggering downstream immune responses such as the activation of mitogen-activated protein kinase (MAPK) cascades and the expression of defense-related genes (Ali *et al.*, 2017; Ma *et al.*, 2009). Additionally, *CNGCs* have been recognized for their involvement in mediating ion fluxes across the plasma membrane, which contribute to the generation of ROS and the establishment of an alkaline cytoplasmic pH, creating an unfavorable environment for fungal growth (Ali *et al.*, 2017; Moeder *et al.*, 2011). This gene has also been found to be an important defense-related gene in plants against *Phytophthora* infection (Ali *et al.*, 2017; Toljamo *et al.*, 2016; Wang *et al.*, 2019). As shown in Figure 5, the plants with transient overexpression of *CNGC1* exhibited decreased levels of crown infection symptoms with newly developed roots but not in the empty vector control plants. CNGCs are not directly involved in lignin production in the plant cell wall, but they can play a role in signaling pathways that respond to biotic stresses, which may influence lignin biosynthesis (Hedrich and Neher et al., 2018). Genes directly involved in lignin biosynthesis and cell wall signaling pathways were expressed in resistant

varieties against the PhCR pathogen (Figure 6). This finding demonstrates the involvement of the

WAK-CNGC defense pathway would act as a frontline defense mechanism against soilborne

pathogens, initiating rapid and effective immune responses to restrict the pathogen invasion and

colonization in plant roots.

By examining the genetic structure of the *RPc2* locus in a large set of genetically diverse

breeding populations, we were able to understand the population structures controlling the

genetic variance for resistance to PhCR. Jiménez *et al.* (2023) evaluated genetically diverse

strawberry populations for resistance to California pathogen isolates, identifying five highly

resistant individuals (1.3%) and 27 moderately resistant individuals (5.5%). Our analysis of 30

individuals revealed that 27 belonged to the resistant *H3* haplotype. Importantly, many

historically resistant cultivars, including 'Cyclone', 'Massey', 'Addie', 'MD683', and 'Camarosa'

were developed over 30 years ago, suggesting a longstanding presence of the PhCR resistance.

Positive selection for disease resistance genes during the breeding process has been well-

documented in various crop species (Gross and Olsen, 2010; Mondragón-Palomino *et al.*, 2002;

Turner-Hissong *et al.*, 2020). The selective sweep observed in our study, where the *H3* haplotype

was found to be well-preserved in breeding populations globally, further supports the notion of

positive selection for disease resistance alleles during breeding. This is consistent with the

previous finding of the extremely high frequency of the *RPc2*-associated favorable allele in the

UCD training population. Our genetic structure analysis of the *RPc2* locus across diverse

breeding populations also revealed a high frequency of the resistant haplotype *H3*, particularly in

strawberries bred in the US and Europe. This skewed distribution of *H3* suggests ongoing

selection for resistance to PhCR in breeding programs worldwide. The identification and

preservation of resistance alleles, whether known or unknown, through selective breeding

strategies underscore the importance of continued efforts to enhance disease resistance in

strawberry. These efforts could be important for sustainable agriculture, particularly in the face

of evolving pathogen populations and changing environmental conditions.

In conclusion, our study has effectively elucidated the genetic architecture of *RPc2*

underlying resistance against *P. cactorum*. The development of a high-quality haplotype-phased

genome for the elite resistant accession containing the resistant haplotype *H3* facilitated the

discovery of candidate genes, *WAK1*, *CNGC1*, and *CNGC2*, for resistance to PhCR. These

findings facilitate understanding of the molecular mechanisms underlying the resistance against

the soilborne pathogen, *P. cactorum*, in strawberries and provide valuable insights for developing

strategies to enhance resistance in breeding programs. Furthermore, the integration of pan-

genomic analyses confirmed the historical sharing of the *RPc2* genomic region during strawberry

breeding in the US. Our study supports the evidence of positive selection for a favorable

resistance allele, demonstrated by the widespread distribution of the *H3* haplotype across

breeding populations in the US and potentially worldwide. These efforts highlight the importance

of continued research and genomic-driven breeding efforts to effectively manage the disease and

ensure the long-term sustainability of strawberry.

Acknowledgements

This research is supported by grants from the United States Department of Agriculture National

Institute of Food and Agriculture (NIFA) Specialty Crops Research Initiative (SCRI) "Delivering

Breeding and Management Solutions to Prevent Losses to Emerging and Expanding Disease

Threats in Strawberry" under award number (#2022-51181-38328). We also acknowledge

Florida Strawberry Growers Association and the strawberry breeding lab members at UF/IFAS

Gulf Coast Research and Education Center for their support of this study.

Competing interests

718 The authors declare no competing interests.

Author contributions

H.H., S.C., Y.J., S.L conceived the hypothesis and original idea and designed the experiments.

H.H., S.C., and U.G. performed the genome assembly and transcriptome analysis. Y.J., Y-H.N,

and S.I. designed molecular markers to conduct the fine mapping analysis. S.V. constructed

genetic linkage map to validate the collinearity. D.P., G.C., R.F., M.F., and S.K. provided field

phenotype data of Phytophthora crow rot and SNP genotyping data. S.L., S.K., V.M. reviewed

and edited the manuscript. All authors reviewed the manuscript.

Data availability

RNA sequencing data for strawberry is available in NCBI SRA database with accession numbers of BioProject number PRJNA890188. DNA sequencing data for strawberry is available in NCBI SRA. The octoploid genome assembly and annotation files are available at Genome Database for Rosaceae (GDR; https://www.rosaceae.org/)

Supplementary Data

Supplementary Fig. S1. Merqury's k-mer-based assembly validation (A) The *k*-mer blob Plot illustrates the distribution of maternal (red) and paternal (blue) k-mer within each scaffold of the respective assemblies. The size of the blob corresponds to the total number of *k*-mers in that scaffold. (B) The combined spectra plot of inherited *k*-mers displays three peaks. The first peak (grey) signifies *k*-mers present in the raw reads but absent from the assembly due to sequencing errors. The second peak corresponds to *k*-mers from heterozygous regions, while the third peak corresponds to *k*-mers from homozygous regions. These plots indicate a complete and haplotype-resolved assembly.

Supplementary Figure Legend

Supplementary Fig. S1. Merqury's k-mer-based assembly validation. The combined spectra plot of inherited *k*-mers displays three peaks. The first peak (grey) signifies *k*-mers present in the raw reads but absent from the assembly due to sequencing errors. The second peak corresponds to *k*-mers from heterozygous regions, while the third peak corresponds to *k*-mers from homozygous regions. These plots indicate a complete and haplotype-resolved assembly.

Supplementary Fig. S2. Dotplot visualization of whole-genome alignment of the FL 16.33-8-phase1 assembly to the diploid *F. vesca* genome assembly ver. 4.0 using MCScanX.

Supplementary Fig. S3. Dotplot of FL 16.33-8-phase1 assembly to diploid *Fragaria vesca* ver. 4.0. (A), *F. iinumae* (B), and *F. viridis* (C). Dot plots are produced using the DGENIE software and alignments with minimap2.

Supplementary Fig. S4. KEGG pathway maps for plant-pathogen interaction in three pairwise comparison.

Supplementary Fig. S5. MapMan visualization of DEGs related to lignin biosynthesis pathways. The log2 fold changes of significant DEGs were imported and visualized in MapMan software (3.5.1 R2).

Supplementary Fig. S6. Gene ontology (GO) of enriched genes across all comparisons following the inoculation of *P. cactorum.*

Supplementary Fig. S7. Genomic, CDS, and peptide sequence alignment of *WAK1* gene derived from 'Florida Brilliance', FL 16.33-8-phase1, and 'Royal Royce' phase1 and 2 assemblies.

Supplementary Fig. S8. Genomic, CDS, and peptide sequence alignment of *CNGC* genes derived from FL 16.33-8-phase1, and 'Royal Royce' phase1 and 2 assemblies.

Supplementary Fig. S9. Genomic, CDS, and peptide sequence alignment of *WAK1* and homoeologous genes from FL 16.33-8-phase1 assembly.

Supplementary Fig. S10. Genomic, CDS, and peptide sequence alignment of *CNGC* and homoeologous genes from FL 16.33-8-phase1 assembly.

Supplementary Fig. S11. Gene expression analysis of *WAK1* and *CNGC1* by qRT-PCR in strawberry resistance to *P. cactorum* pathogens. (A) The transient expression assay was conducted employing the pMDC32 overexpression vector to drive the expression of *WAK1* and *CNGC1* genes. (B) The transient expression assay was conducted employing the pK7GWIWG(II) RNAi vector to drive the knockdown expression of *WAK1* and *CNGC1* genes. The x-axis represents the vector composition used. EV denotes 'Empty Vector, 'Error bars represent standard deviation. Asterisks indicate significant outcomes with P values of Student's t-test (*P < 0.05, **P < 0.01, ***P < 0.001, Student's t-test), respectively.

Supplementary Fig. S12. Delta K values for STRUCTURE analysis of octoploid strawberries.

Supplementary Fig. S13. A hypothetical model for the enhanced resistance against *Phytophthora cactorum* in the *WAK1*, *CNGC1* knockdown strawberry root. Differentially expressed genes identified in *WAK1* and *CNGC1* knockdown crown in response to *P. cactorum* were represented. DAMPs such as Oligogalacturonides, and pectin fractions are sensed by *WAK1*, leading to the activation of *FaRLCK*. Subsequently, the activated *FaRLCK*

800   phosphorylates *CNGC1* induces extracellular Ca^{2+} influx and then triggers PTI-related gene
801   expressions, which result in disease resistance in strawberry.
802
803   **Supplementary Dataset1.** Detailed information on Genetic linkage map for 'Florida Brilliance'
804   and FL 16.33-8
805
806   **Supplementary Dataset2.** Dataset used for phylogenetic analysis and principal component
807   analysis.
808
809   **Supplementary Dataset3.** Dataset used for structure analysis.
810

REFERENCES

Ali SS, Shao J, Lary DJ, Kronmiller BA, Shen D, Strem MD, Amoako-Attah I, Akrofi AY, Begoude BD, Ten Hoopen GM. 2017. Phytophthora megakarya and Phytophthora palmivora, closely related causal agents of cacao black pod rot, underwent increases in genome sizes and gene numbers by different mechanisms. Genome biology and evolution **9**, 536-557.

Alonge M, Lebeigle L, Kirsche M, Jenike K, Ou S, Aganezov S, Wang X, Lippman ZB, Schatz MC, Soyk S. 2022. Automated assembly scaffolding using RagTag elevates a new tomato system for high-throughput genome editing. Genome biology **23**, 1-19.

Anderson DE, Scuteri A, Agalakova N, Parsons DJ, Bagrov AY. 2001. Racial differences in resting end-tidal CO2 and circulating sodium pump inhibitor. American journal of hypertension **14**, 761-767.

Baluška F, Šamaj J, Wojtaszek P, Volkmann D, Menzel D. 2003. Cytoskeleton-plasma membrane-cell wall continuum in plants. Emerging links revisited. Plant physiology **133**, 482-491.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics **30**, 2114-2120.

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. TASSEL: software for association mapping of complex traits in diverse samples. Bioinformatics **23**, 2633-2635.

Brutus A, Sicilia F, Macone A, Cervone F, De Lorenzo G. 2010. A domain swap approach reveals a role of the plant wall-associated kinase 1 (WAK1) as a receptor of oligogalacturonides. Proceedings of the National Academy of Sciences **107**, 9452-9457.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. Nature methods **12**, 59-60.

Burra DD, Berkowitz O, Hedley PE, Morris J, Resjö S, Levander F, Liljeroth E, Andreasson E, Alexandersson E. 2014. Phosphite-induced changes of the transcriptome and secretome in Solanum tuberosum leading to resistance against Phytophthora infestans. BMC plant biology **14**, 254.

Cal Ad, Martinez-Treceno A, Lopez-Aranda J, Melgarejo P. 2004. Chemical alternatives to methyl bromide in Spanish strawberry nurseries. Plant disease **88**, 210-214.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. BMC bioinformatics **10**, 421.

Campos ML, Kang J-H, Howe GA. 2014. Jasmonate-triggered plant immunity. Journal of chemical ecology **40**, 657-675.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. Nature methods **18**, 170-175.

Curtis MD, Grossniklaus U. 2003. A gateway cloning vector set for high-throughput functional analysis of genes in planta. Plant physiology **133**, 462-469.

Davik J, Eikemo H, Brurberg MB, Sargent DJ. 2015. Mapping of the RPc-1 locus for Phytophthora cactorum resistance in Fragaria vesca. Molecular Breeding **35**, 1-11.

Deising HB. 2009. *Plant relationships*: Springer Science & Business Media.

Desikan R, Cheung M-K, Clarke A, Golding S, Sagi M, Fluhr R, Rock C, Hancock J, Neill S. 2004. Hydrogen peroxide is a common signal for darkness-and ABA-induced stomatal closure in Pisum sativum. Functional Plant Biology **31**, 913-920.

Earl DA, VonHoldt BM. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. Conservation genetics resources **4**, 359-361.

Edger PP, Poorten TJ, VanBuren R, Hardigan MA, Colle M, McKain MR, Smith RD, Teresi SJ, Nelson AD, Wai CM. 2019. Origin and evolution of the octoploid strawberry genome. Nature genetics **51**, 541-547.

Ferrari S, Savatin DV, Sicilia F, Gramegna G, Cervone F, De Lorenzo G. 2013. Oligogalacturonides: plant damage-associated molecular patterns and regulators of growth and development. Frontiers in plant science **4**, 49.

Francis RM. 2017. pophelper: an R package and web app to analyse and visualize population structure. Molecular ecology resources **17**, 27-32.

Gross BL, Olsen KM. 2010. Genetic perspectives on crop domestication. Trends in plant science **15**, 529-537.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. Bioinformatics **29**, 1072-1075.

Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR, Wortman JR. 2008. Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. Genome biology **9**, 1-22.

Hano C, Addi M, Bensaddek L, Crônier D, Baltora-Rosset S, Doussot J, Maury S, Mesnard F, Chabbert B, Hawkins S. 2006. Differential accumulation of monolignol-derived compounds in elicited flax (Linum usitatissimum) cell suspension cultures. Planta **223**, 975-989.

Hardigan MA, Feldmann MJ, Lorant A, Bird KA, Famula R, Acharya C, Cole G, Edger PP, Knapp SJ. 2020. Genome synteny has been conserved among the octoploid progenitors of cultivated strawberry over millions of years of evolution. Frontiers in Plant Science **10**, 1789.

Hardigan MA, Lorant A, Pincot DD, Feldmann MJ, Famula RA, Acharya CB, Lee S, Verma S, Whitaker VM, Bassil N. 2021. Unraveling the complex hybrid ancestry and domestication history of cultivated strawberry. Molecular biology and evolution **38**, 2285-2305.

He ZH, He D, Kohorn BD. 1998. Requirement for the induced expression of a cell wall associated receptor kinase for survival during the pathogen response. The Plant Journal **14**, 55-63.

Hirakawa H, Shirasawa K, Kosugi S, Tashiro K, Nakayama S, Yamada M, Kohara M, Watanabe A, Kishida Y, Fujishiro T. 2014. Dissection of the octoploid strawberry genome by deep sequencing of the genomes of Fragaria species. DNA research **21**, 169-181.

Hoff KJ, Stanke M. 2019. Predicting genes in single genomes with AUGUSTUS. Current protocols in bioinformatics **65**, e57.

Holmes GJ, Mansouripour SM, Hewavitharana SS. 2020. Strawberries at the crossroads: Management of soilborne diseases in California without methyl bromide. Phytopathology **110**, 956-968.

Hou X, Tong H, Selby J, DeWitt J, Peng X, He Z-H. 2005. Involvement of a cell wall-associated kinase, WAKL4, in Arabidopsis mineral responses. Plant physiology **139**, 1704-1716.

Humann JL, Lee T, Ficklin S, Main D. 2019. Structural and functional annotation of eukaryotic genomes with GenSAS. *Gene prediction*: Springer, 29-51.

Hummer K, Bassil N, Njuguna W. 2011. Fragaria//Wild Crop Relatives: Genomic and Breeding Resources, Temperate Fruits/Ed. C. Kole. Berlin: Springer.

Jiménez NP, Feldmann MJ, Famula RA, Pincot DD, Bjornson M, Cole GS, Knapp SJ. 2023. Harnessing underutilized gene bank diversity and genomic prediction of cross usefulness to enhance resistance to Phytophthora cactorum in strawberry. The plant genome **16**, e20275.

Karimi M, Inzé D, Depicker A. 2002. GATEWAY™ vectors for Agrobacterium-mediated plant transformation. Trends in plant science **7**, 193-195.

Keb-Llanes M, González G, Chi-Manzanero B, Infante D. 2002. A rapid and simple method for small-scale DNA extraction in Agavaceae and other tropical plants. Plant Molecular Biology Reporter **20**, 299-300.

Khatib M, Lafitte C, Esquerré-Tugayé MT, Bottin A, Rickauer M. 2004. The CBEL elicitor of Phytophthora parasitica var. nicotianae activates defence in Arabidopsis thaliana via three different signalling pathways. New Phytologist **162**, 501-510.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology **37**, 907-915.

Kohorn BD, Kohorn SL. 2012. The cell wall-associated kinases, WAKs, as pectin receptors. Frontiers in Plant Science **3**, 25457.

Lally D, Ingmire P, Tong H-Y, He Z-H. 2001. Antisense expression of a cell wall–associated protein kinase, WAK4, inhibits cell elongation and alters morphology. The Plant Cell **13**, 1317-1332.

Lauvergeat V, Lacomme C, Lacombe E, Lasserre E, Roby D, Grima-Pettenati J. 2001. Two cinnamoyl-CoA reductase (CCR) genes from Arabidopsis thaliana are differentially expressed during development and in response to infection with pathogenic bacteria. Phytochemistry **57**, 1187-1195.

Lee H-E, Manivannan A, Lee SY, Han K, Yeum J-G, Jo J, Kim J, Rho IR, Lee Y-R, Lee ES. 2021. Chromosome level assembly of homozygous inbred line 'Wongyo 3115'facilitates the construction of a high-density linkage map and identification of QTLs associated with fruit firmness in octoploid strawberry (Fragaria× ananassa). Frontiers in Plant Science **12**, 696229.

Lin X, Jia Y, Heal R, Prokchorchik M, Sindalovskaya M, Olave-Achury A, Makechemu M, Fairhead S, Noureen A, Heo J. 2023. Solanum americanum genome-assisted discovery of immune receptors that detect potato late blight pathogen effectors. Nature genetics **55**, 1579-1588.

Love M, Anders S, Huber W. 2014. Differential analysis of count data–the DESeq2 package. Genome Biol **15**, 10-1186.

Lozada DN, Nunez G, Lujan P, Dura S, Coon D, Barchenger DW, Sanogo S, Bosland PW. 2021. Genomic regions and candidate genes linked with Phytophthora capsici root rot resistance in chile pepper (Capsicum annuum L.). BMC plant biology **21**, 601.

Ma W, Qi Z, Smigel A, Walker RK, Verma R, Berkowitz GA. 2009. Ca2+, cAMP, and transduction of non-self perception during plant immune responses. Proceedings of the National Academy of Sciences **106**, 20995-21000.

Mangandi J, Verma S, Osorio L, Peres NA, van de Weg E, Whitaker VM. 2017. Pedigree-based analysis in a multiparental population of octoploid strawberry reveals QTL alleles conferring resistance to Phytophthora cactorum. G3: Genes, Genomes, Genetics **7**, 1707-1719.

Moeder W, Urquhart W, Ung H, Yoshioka K. 2011. The role of cyclic nucleotide-gated ion channels in plant immunity. Molecular plant **4**, 442-452.

Mondragón-Palomino M, Meyers BC, Michelmore RW, Gaut BS. 2002. Patterns of positive selection in the complete NBS-LRR gene family of Arabidopsis thaliana. Genome research **12**, 1305-1315.

Navarro JF, Buron E, Martín-López M. 2004. Behavioral profile of L-655,708, a selective ligand for the benzodiazepine site of GABA-A receptors which contain the α5 subunit, in social encounters between male mice. Aggressive Behavior: Official Journal of the International Society for Research on Aggression **30**, 319-325.

Nellist CF, Vickerstaff RJ, Sobczyk MK, Marina-Montes C, Wilson FM, Simpson DW, Whitehouse AB, Harrison RJ. 2019. Quantitative trait loci controlling Phytophthora cactorum resistance in the cultivated octoploid strawberry (Fragaria× ananassa). Horticulture Research **6**.

Noh Y-H, Lee S, Whitaker VM, Cearley KR, Cha J-S. 2017. A high-throughput marker-assisted selection system combining rapid DNA extraction high-resolution melting and simple sequence repeat analysis: Strawberry as a model for fruit crops. Journal of Berry Research **7**, 23-31.

Noh Y-H, Oh Y, Mangandi J, Verma S, Zurn JD, Lu Y-T, Fan Z, Bassil N, Peres N, Cole G. 2018. High-throughput marker assays for FaRPc2-mediated resistance to Phytophthora crown rot in octoploid strawberry. Molecular Breeding **38**, 104.

Ou S, Chen J, Jiang N. 2018. Assessing genome assembly quality using the LTR Assembly Index (LAI). Nucleic acids research **46**, e126-e126.

Price MN, Dehal PS, Arkin AP. 2009. FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. Molecular biology and evolution **26**, 1641-1650.

Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus genotype data. Genetics **155**, 945-959.

Rhie A, Walenz BP, Koren S, Phillippy AM. 2020. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. Genome biology **21**, 1-27.

Shirasawa K, Hirakawa H, Nakayama S, Sasamoto S, Tsuruoka H, Minami C, Watanabe A, Kishida Y, Kohara M, Yamada M. 2021. A chromosome-scale strawberry genome assembly of a Japanese variety, Reikou. BioRxiv.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics **31**, 3210-3212.

Stensvand A, Herrero M, Talgø V. 1999. Crown rot caused by Phytophthora cactorum in Norwegian strawberry production. EPPO Bulletin **29**, 155-158.

Su W, Ou S, Hufford MB, Peterson T. 2021. A tutorial of EDTA: extensive de novo TE annotator. Plant Transposable Elements, 55-67.

Tang D, Wang G, Zhou J-M. 2017. Receptor kinases in plant-pathogen interactions: more than pattern recognition. The Plant Cell **29**, 618-637.

Tennessen JA, Govindarajulu R, Ashman T-L, Liston A. 2014. Evolutionary origins and dynamics of octoploid strawberry subgenomes revealed by dense targeted capture linkage maps. Genome biology and evolution **6**, 3295-3313.

Toljamo A, Blande D, Kärenlampi S, Kokko H. 2016. Reprogramming of strawberry (Fragaria vesca) root transcriptome in response to Phytophthora cactorum. PLoS One **11**, e0161078.

Turner-Hissong SD, Bird KA, Lipka AE, King EG, Beissinger TM, Angelovici R. 2020. Genomic prediction informed by biological processes expands our understanding of the genetic architecture underlying free amino acid traits in dry Arabidopsis seeds. G3: Genes, Genomes, Genetics **10**, 4227-4239.

Urao T, Yakubov B, Satoh R, Yamaguchi-Shinozaki K, Seki M, Hirayama T, Shinozaki K. 1999. A transmembrane hybrid-type histidine kinase in Arabidopsis functions as an osmosensor. The Plant Cell **11**, 1743-1754.

Verica JA, He Z-H. 2002a. The cell wall-associated kinase (WAK) and WAK-like kinase gene family. Plant physiology **129**, 455-459.

Verica JA, He Z-H. 2002b. The Cell Wall-Associated Kinase (WAK) andWAK-Like Kinase Gene Family. Plant physiology **129**, 455-459.

Wang J, Liu X, Zhang A, Ren Y, Wu F, Wang G, Xu Y, Lei C, Zhu S, Pan T. 2019. A cyclic nucleotide-gated channel mediates cytoplasmic calcium elevation and disease resistance in rice. Cell research **29**, 820-831.

Wang W, Chen L, Fengler K, Bolar J, Llaca V, Wang X, Clark CB, Fleury TJ, Myrvold J, Oneal D. 2021. A giant NLR gene confers broad-spectrum resistance to Phytophthora sojae in soybean. Nature Communications **12**, 6263.

Xiao C, Gao J, Zhang Y, Wang Z, Zhang D, Chen Q, Ye X, Xu Y, Yang G, Yan L. 2019. Quantitative proteomics of potato leaves infected with phytophthora infestans provides insights into coordinated and altered protein expression during early and late disease stages. International journal of molecular sciences **20**, 136.

Yoshioka K, Moeder W, Kang H-G, Kachroo P, Masmoudi K, Berkowitz G, Klessig DF. 2006. The chimeric Arabidopsis CYCLIC NUCLEOTIDE-GATED ION CHANNEL11/12 activates multiple pathogen resistance responses. The Plant Cell **18**, 747-763.

Zhong L, Zhang Y, Liu H, Sun G, Chen R, Song S. 2016. Agrobacterium-mediated transient expression via root absorption in flowering Chinese cabbage. SpringerPlus **5**, 1-7.

Zhou X, Wen K, Huang S-X, Lu Y, Liu Y, Jin J-H, Kale SD, Chen X-R. 2023. Time-Course Transcriptome Profiling Reveals Differential Resistance Responses of Tomato to a Phytotoxic Effector of the Pathogenic Oomycete Phytophthora cactorum. Plants **12**, 883.

Zimmermann S, Ehrhardt T, Plesch G, Müller-Röber B. 1999. Ion channels in plant signaling. Cellular and Molecular Life Sciences CMLS **55**, 183-203.

A

B

RPc2-H3 Control vs.
RPc2-H3 P.o Inoculation

34 ← Genes involved in PNCR Resistance

rpc2-h3 Control vs.
rpc2-h3 P.o Inoculation

rpc2-h3 P.o Inoculation vs.
RPc2-H3 P.o Inoculation

D

C

A

AX-89845076

AX-184173618

FL 16.33-8-1
Royal Royce-1
Royal Royce-2
Florida Brilliance-1
FL 16.33-8-2

B

22.99 Mb

Florida Brilliance
(*farpc2-h1*)

WAK
3,079 bp

22.07 Mb

FL 16.33-8
(*FaRPc2-H3*)

CNGC 2 (Fxa7Bg202971)   CNGC 1 (Fxa7Bg202973)   WAK

3,387 bp   20,524 bp   2,967 bp   13,987 bp   2,823 bp

C

WAK1

WAK (FL 16.33-8)   wak (FL Brilliance)

CNGC1 & 2

CNGC1 (FL 16.33-8)   CNGC2 (FL 16.33-8)

D

Empty

pK7::*FaWAK*

pK7::*FaCNGC*

E

Empty

pMDC32::*FaWAK*

pMDC32::*FaCNGC*

A

Biotic stress

Abiotic stress

Recognition

R genes

Respiratory burst

Hormone signaling

Auxins
Brassinost.
ABA
Ethylene
SA
JA

Putative involvement in biotic stress

Misc. functions

HSPs

Cell wall

Proteolysis

Signaling

MAPK

Transcription factors

Defense genes

PR-proteins

Secondary metabolites

Redox state
Peroxidases
Glutathione-S-transferase

Signaling

MAPK

ERF
bZIP
WRKY
MYB
DOF

Transcription factors

Putative involvement in biotic stress

Heat shock proteins

Secondary metabolites

B

Secondary metabolism

Non MVA pathway

Shikimate pathway

MVA pathway

Terpenoids

Tocopherol

Phenlypropanoids

Simple phenols

Alkaloids-like

Carotenoids

Lignin and lignans

Betains

Wax

Flavonoids

Chalcones

Isoflavonoids

Flavonols

Dihydroflavonols

Flavonones

Anthocyanins

Glucosinolates

Cyanogenic glycosides

Sulfur containing