TITLE: MEF2C controls segment-specific gene regulatory networks that direct heart tube morphogenesis

AUTHOR LIST: Jonathon M. Muncie-Vasic[1], Tanvi Sinha[2], Alexander P. Clark[3], Emily F. Brower[1], Jeffrey J. Saucerman[3,4], Brian L. Black[2,5], Benoit G. Bruneau[1,6,7,8,9],*

AFFILIATIONS: [1]Gladstone Institutes, San Francisco, CA; [2]Cardiovascular Research Institute, University of California, San Francisco, CA; [3]Department of Biomedical Engineering, University of Virginia, Charlottesville, VA; [4]Robert M. Berne Cardiovascular Research Center, University of Virginia, Charlottesville, VA; [5]Department of Biochemistry and Biophysics, University of California, San Francisco, CA; [6]Roddenberry Center for Stem Cell Biology and Medicine, Gladstone Institutes, San Francisco, CA; [7]Department of Pediatrics, University of California, San Francisco, CA; [8]Institute for Human Genetics, University of California, San Francisco, CA; [9]Eli and Edythe Broad Center for Regeneration Medicine and Stem Cell Research, University of California, San Francisco, CA; *Author for correspondence (benoit.bruneau@gladstone.ucsf.edu)

ABSTRACT

The gene regulatory networks (GRNs) that control early heart formation are beginning to be understood, but lineage-specific GRNs remain largely undefined. We investigated networks controlled by the vital transcription factor MEF2C, with a time course of single-nucleus RNA- and ATAC-sequencing in wild-type and *Mef2c*-null embryos. We identified a "posteriorized" cardiac gene signature and chromatin landscape in the absence of MEF2C. Integrating our multiomics data in a deep learning-based model, we constructed developmental trajectories for each of the outflow tract, ventricular, and inflow tract segments, and alterations of these in *Mef2c*-null embryos. We computationally identified segment-specific MEF2C-dependent enhancers, with activity in the developing zebrafish heart. Finally, using inferred GRNs we discovered that the *Mef2c*-null heart malformations are partly driven by increased activity of the nuclear hormone receptor NR2F2. Our results delineate lineage-specific GRNs in the early heart tube and provide a generalizable framework for dissecting transcriptional networks governing developmental processes.

A recurrent observation in developmental biology is that the same transcription factor (TF)
can play different roles depending on the cells or developmental stage in which it is
expressed. The cell type- or stage-specific activity of a TF is dependent on a number of
factors, including its own expression level, the availability of potential co-factors, the
presence or absence of other TFs that may act co-operatively or antagonistically, and the
dynamic state of the chromatin landscape in which the TF is acting[1–4]. Together, these
phenomena can be conceptualized as transcriptional or gene regulatory networks (GRNs)
that can be used to understand the role of a particular TF within a given developmental
context[5]. However, for most TFs, the precise GRNs underlying their pleiotropic activity in
distinct cell types and at different developmental timepoints remain undefined.

The developing heart presents an excellent model for exploring this question.
Several TFs critical for cardiac development have been identified and their expression
patterns are well-defined[6–8]. Moreover, lineage tracing studies have revealed distinct
progenitor cell types that contribute to the various segments of the linear heart tube[7–11]
(Fig. 1a). To explore in detail how a specific TF can have distinct roles within different
parts of a developing organ, we focused on a key cardiac transcription factor, MEF2C[12],
and its role in each of the developing heart tube segments. In mice, loss of MEF2C leads
to cardiac defects and embryonic lethality by mid-gestation at embryonic day (E) 10.5[13].
Interestingly, although MEF2C is expressed in both first heart field (FHF) and second
heart field (SHF) progenitors in the cardiac crescent at E7.75, and throughout the
developing heart tube from E8.5 to E9, the loss of MEF2C causes distinct defects in
different segments of the heart tube (Fig. 1b-c, Extended Data Fig. 1a). At E9, the outflow
tract, which gives rise to the aorta and pulmonary trunk, is severely hypoplastic and only
a single hypoplastic ventricle has formed. By contrast, the inflow tract, precursor to the
atria, appears to be expanded and improperly patterned (Fig. 1c). These segment-specific
defects suggest that MEF2C plays distinct regulatory roles in the outflow tract, ventricles,
and inflow tract.

To dissect the segment-specific regulation of heart tube morphogenesis by
MEF2C, we performed combined single-nucleus RNA-seq and ATAC-seq (snRNA-seq
and snATAC-seq) on wild-type (WT) and *Mef2c* knockout (KO) mouse embryos at key
stages throughout heart tube development. Taking advantage of the ability to match the

transcriptomic and chromatin accessibility data for every cell in the dataset, we applied a

suite of complementary bioinformatics tools to construct GRNs for cardiac progenitor

lineages, precisely defined the role of MEF2C in regulating gene expression and

chromatin accessibility in the developing heart tube, identified novel MEF2C-dependent

enhancers, and discovered a genetic interaction that underlies the inflow tract

malformation in *Mef2c* KO embryos (Fig. 1d).

RESULTS

Loss of MEF2C reduces expression of key cardiomyocyte genes across all heart

tube segments and alters expression of anterior/posterior markers

To assess the role of MEF2C in controlling gene expression in the developing heart tube,

we performed snRNA-seq on stage-matched WT and *Mef2c* KO embryos at E7.75, E8.5,

and E9 (n = 2 embryos for each genotype at each stage, Extended Data Fig. 1b). All

embryos carried the *Smarcd3*-F6-eGFP transgene reporter to mark cardiac progenitors[14].

At E7.75, we harvested the entire embryos, whereas at E8.5 and E9 we removed the

headfolds and posterior trunk via microdissection to increase the proportion of cardiac

progenitors captured for sequencing. We created separate Seurat objects[15] for the

embryos profiled at each timepoint (E7.75, E8.5, and E9). Following pre-processing,

quality control, normalization with scTransform[16], and dimensionality reduction, we

performed unbiased Louvain clustering and plotted the data in uniform manifold

approximation and projection (UMAP) space. We then used the expression of known

marker genes to identify the cell types represented by each cluster (Extended Data Fig.

2a-f, Supplementary Table 1).

To resolve specific cardiomyocyte subtypes, we subset and re-clustered cell types

of interest, including cardiac progenitors, cardiomyocytes, and related mesoderm cell

types (Fig. 2a-c, Extended Data Fig. 2g-i, Supplementary Table 2). By doing so, we were

able to identify clusters representing FHF and early differentiating cardiomyocytes

(CMs/FHF), SHF, and juxtacardiac field (JCF) progenitors[17] at E7.75 (Fig. 2a). At E8.5,

we could distinguish inflow tract cardiomyocytes (IFT-CMs), ventricular cardiomyocytes

(V-CMs), and outflow tract cardiomyocytes (OFT-CMs) (Fig. 2b). At E9 we identified the

IFT-CMs more specifically as either atrial or atrioventricular canal (A- or AVC-CMs) (Fig.

2c). We next performed differential gene expression analysis between *Mef2c* KO and WT

cells within each of these cell types (Fig. 2d-f, Supplementary Table 3). Overall, more

genes were down-regulated than up-regulated in the *Mef2c* KO relative to WT embryos,

consistent with the notion that MEF2C is an activator of the myogenic transcriptional

program[18]. Among the top differentially expressed genes (DEGs), we found several genes

that have been associated with high confidence to congenital heart defects[19], including

Myh6, *Myh7*, *Actc1*, and *Foxp1* (Extended Data Fig. 2j). Furthermore, nearly all these top

DEGs were identified as the nearest genes to MEF2C-occupied peaks in a previously

published MEF2C ChIP-seq dataset of embryonic hearts at E14[20], suggesting that these

genes are direct targets of MEF2C in the developing heart.

Closer examination of our DEG data (Supplementary Table 3) revealed two

categories of DEGs. The first category consisted of key cardiomyocyte genes that were

substantially down-regulated across all three heart tube segments in the *Mef2c* KO

embryos, including genes encoding sarcomeric proteins such as *Tnnt2* and *Ttn,* the

cardiac transcription factor *Nkx2-5*, and the natriuretic peptide *Nppa* (Fig. 2g-i). These

data confirm the importance of MEF2C in early cardiomyocyte fate specification, as a

direct activator of contractile protein gene expression and by regulating additional TFs

and functional proteins necessary for proper cardiomyocyte development. The second

category of DEGs consisted of genes involved in anterior/posterior patterning of the heart

tube. Specifically, genes such as *Tbx5*, *Gata4*, and *Wnt2,* which are normally expressed

only in the posterior IFT segment of the heart tube, were not only up-regulated in the IFT-

CMs, but their expression was also expanded into the V-CMs in *Mef2c* KO (Fig. 2g-i).

Additionally, expression of the anterior OFT-specific gene *Tdgf1* was completely lost in

Mef2c KO (Fig. 2h-i), as previously demonstrated[21]. Consistent with this apparent

posteriorization of gene expression in the heart tube, we also observed a larger proportion

of A-CMs and AVC-CMs in *Mef2c* KO embryos compared to WT at E9, with a concomitant

reduction in the proportion of V-CMs and OFT-CMs (Extended Data Fig. 2k). We validated

the observed changes in gene expression with whole-mount embryo fluorescence *in situ*

hybridization, confirming that *Mef2c* KO leads to broad reduction in the cardiomyocyte

transcriptional program and a notable posteriorization of the developing heart tube (Fig.

2j).

MEF2C regulates chromatin accessibility broadly throughout the heart tube and in a segment-specific manner

To investigate how MEF2C regulates the observed segment-specific gene expression
changes, we integrated our chromatin accessibility and gene expression datasets using
ArchR[22]. We performed dimensionality reduction using ArchR's latent semantic index
(LSI) implementation on the snRNA-seq and snATAC-seq modalities and then plotted the
cells in UMAP space using the combined dimensions (Extended Data Fig. 3a-c). Matching
cell barcodes in the snRNA-seq and snATAC-seq libraries allowed us to transfer the cell
type labels from the Seurat objects to the integrated ArchR datasets. As we did with the
gene expression data, we subset the cardiac cell types and associated mesoderm in
order to better resolve cardiomyocyte subtypes. Finally, we created pseudobulk groups
consisting of WT and *Mef2c* KO cells for each given cell type label in the subset datasets
and called peaks representing accessible regions using MACS2[23].

Our initial analyses revealed few changes to chromatin accessibility at E7.75
(Extended Data Fig. 3d). This was not entirely unexpected, given that there is no
discernable phenotype in the *Mef2c* KO embryos at this timepoint, and agrees with the
relatively few changes in gene expression we observed in CMs/FHF, SHF, and JCF cells
at E7.75 (Fig. 2d). Thus, we focused on the E8.5 dataset to identify the earliest substantial
changes in chromatin accessibility induced by the loss of MEF2C (Fig. 3a). Using our
pseudobulked data, we identified differentially accessible regions (DARs) of chromatin
between *Mef2c* KO and WT cells in each of the subset cell types and found that, as
expected, cardiomyocytes displayed the largest numbers of DARs (Fig. 3b). We also
found that in each of the CM subtypes (IFT-CMs, V-CMs, and OFT-CMs), *Mef2c* KO CMs
lost more DARs than they gained (Fig. 3b), and these lost DARs were highly enriched for
MEF2 binding motifs (Extended Data Fig. 3e). These data further reinforce MEF2C's role
as an activator of the contractile gene transcriptional program. Interestingly, when we
examined the lost DARs in *Mef2c* KO embryos, we found 675 regions that demonstrated
lost accessibility in two or more heart tube segments, and 2,549 regions that lost
accessibility in a segment-specific manner (Fig. 3c). The former are likely regions involved
in the regulation of general cardiomyocyte genes, while the latter may control segment-

specific gene regulation. By contrast, the gained DARs in *Mef2c* KO embryos were almost

exclusively unique to each heart tube segment (Fig. 3c). These segment-specific DARs

are interesting because they provide evidence that distinct transcriptional networks exist

for each of the three heart tube segments and that each segment-specific network is

uniquely altered by the loss of MEF2C.

To identify the DARs most likely to represent *bona fide* regulatory elements, we

performed a peak-to-gene linkage (P2G) analysis, which identifies statistically significant

correlations between the accessibility of individual peaks and the expression of potential

target genes within 250 kilobases of each peak. We found 817 P2G links in the OFT-CMs,

1,201 in the V-CMs, and 2,257 in the IFT-CMs. We then intersected the P2G links with

the MEF2C-dependent DEGs at each timepoint (Supplementary Table 4) and plotted

each P2G link according to the fold change of the peak's accessibility, the fold change of

linked gene's differential expression, and the correlation strength of the P2G link (Fig. 3d).

These P2G DAR-DEG correlation plots reveal a number of interesting

observations. Genes such as *Nkx2-5*, *Myh6*, and *Myh7*, whose expression decreased

throughout the heart tube in *Mef2c* KO embryos, are linked to DARs in all three segments

(Fig. 3d). In particular, the *Myh6*/*Myh7* locus contains multiple regions that have P2G links

and clearly altered accessibility in OFT-CMs, V-CMs, and IFT-CMs (Fig. 3e). This

suggests that these DARs are likely to be regulatory elements that control expression of

Myh6 and *Myh7* and are sensitive to the expression of *Mef2c* in each segment of the

heart tube. By contrast, there are also P2G links between peaks with lost accessibility

and decreased gene expression that are unique to the individual segments, such as links

to *Wnt11* in the OFT-CMs, *Nppa* in the V-CMs, and *Cacna1c* in the IFT-CMs (Fig. 3d).

Furthermore, examining the links between peaks with gained accessibility and genes with

increased expression in *Mef2c* KO reveals additional evidence for the posteriorized

transcriptional program we previously described. Namely, P2G links to *Tbx5* in the V-CMs,

as well as *Gata4* and *Wnt2* in the IFT-CMs (Fig. 3d and 3f). These P2G links represent

potential regulatory elements that are activated or over-activated in response to the loss

of MEF2C and, at least in part, promote the increased posteriorization of the heart tube.

Each heart tube segment exhibits a distinct MEF2C-depedent developmental trajectory

Our multiomic datasets containing precisely staged embryos at different developmental timepoints provided us the opportunity to dissect how lineage trajectories unfold in WT embryos and are altered by loss of MEF2C. To this end, we applied multimodal Models for Integrated Regulatory Analysis (MIRA)[24], which uses deep learning and probabilistic graphical models to construct sets of "topics" that represent either gene expression or peak accessibility. These topics can then be used to plot cells in UMAP space, assign cell identities, perform pseudotime analyses, and construct cell state trees (Fig. 4a).

We trained topic models using cardiac progenitors and cardiomyocyte subtypes from both genotypes at each of our three timepoints (E7.75: CMs/FHF, SHF, JCF; E8.5: aSHF, pSHF, OFT-CMs, V-CMs, IFT-CMs; E9: OFT-CMs, V-CMs, A-CMs, AVC-CMs) and then used those models to construct joint representation UMAPs of the outflow tract, ventricular, and inflow tract lineages (Fig. 4b, 4f, 4j). We next calculated pseudotime values for cells within the UMAPs and used the results to construct cell state trajectories for each of the three lineages (Fig. 4c-d, 4g-h, 4k-l).

We observed clear differences in the fate specification of *Mef2c* KO cells compared to WT cells, as represented by branch points in the trajectories (Fig. 4d, 4h, 4l). Most interestingly, we noticed that the structure of these trajectories, along with the manner and timing by which *Mef2c* KO cells diverged from WT cells within the trajectories, was different for all three lineages. For instance, within the outflow tract lineage, *Mef2c* KO and WT cells proceeded similarly up until E8.5, and then diverged as development proceeded to E9 (Fig. 4d). Within the ventricular lineage, there was no branch point. Instead, *Mef2c* KO V-CMs from both E8.5 and E9 were located earlier in the trajectory than even E8.5 WT V-CMs (Fig. 4h). This indicates that loss of MEF2C in V-CMs resulted in a marked delay or termination of normal development rather than a divergence to an alternative cell state. The situation was most complex for the inflow tract lineage, in which there were two points of divergence between *Mef2c* KO and WT cells (Fig. 4l). First, a subset of *Mef2c* KO cells branched away from the WT trajectory as they differentiated from posterior SHF (pSHF) progenitors to IFT-CMs at E8.5. A second set of *Mef2c* KO cells proceeded along the WT trajectory to E9, before ultimately diverging from WT cells

at the "terminal" AVC-CM and A-CM states. This complex branching behavior may reflect
the notion that both FHF and pSHF progenitors contribute to the IFT[11] and suggests that
MEF2C likely controls distinct GRNs within these two IFT progenitor subpopulations.
Together, these data reinforce our working hypothesis that MEF2C regulates
development of the heart tube in a segment-specific manner via distinct regulatory
networks.

We gained further insights regarding the segment-specific regulation of heart tube
development by examining the top 200 genes and the top ranked TF binding motifs
enriched in the gene expression and chromatin accessibility topics (Supplementary Table
5). As WT cells diverged from *Mef2c* KO cells in the outflow tract lineage, their gene
expression topics became enriched for canonical outflow tract markers *Tdgf1* and
Wnt11[21,25], and their accessibility topics for MEF2 and FOS binding motifs (Fig. 4e). By
contrast, the *Mef2c* KO cells were marked by gene expression topics enriched for Notch
and Wnt signaling components such as *Dlk1* and *Fzd1*, and transcription factors *Meis1*
and *Hand2,* and exhibited enrichment of an accessibility topic containing ISL, NKX, and
TEAD binding motifs (Fig. 4e), suggesting these cells may retain a more progenitor-like
state in the absence of MEF2C. The gene expression and accessibility topics enriched
along the ventricular lineage trajectory supports our interpretation that the *Mef2c* KO cells
experience a substantial delay or termination of their normal differentiation. Notably,
topics that were enriched for CM genes (e.g. *Myl2*, *Tnnt1*) and cardiac TF binding motifs
(e.g. TBX, MEF2, NKX) marked the terminal end of the differentiation trajectory consisting
of only WT cells (Fig. 4i). Interestingly, the mid-point of the ventricular trajectory containing
E8.5 and E9 *Mef2c* KO cells is marked by an accessibility topic enriched for the nuclear
receptor NR2F2 binding motif, a transcription factor that is restricted to the IFT in WT
embryos[26] (Fig. 4i). In the complex branching inflow tract trajectory, terminal WT cells
were enriched for topics containing expected atrial genes, such as *Angpt1* and *Nr2f1*, and
binding motifs for TBX5 and SNAI, among others (Fig. 4m, Extended Data Fig. 4a-b). By
contrast, the *Mef2c* KO cells streamed into one of two terminal branches enriched for
genes such as *Wnt6, Jarid2, Tbx2,* and *Zeb1* and binding motifs for ERF, GATA, and
TEAD factors (Fig. 4m, Extended Data Fig. 4a-b). Together, these data illustrate the
lineage-specific changes in gene expression and chromatin accessibility that accompany

heart tube development and the distinct impact of the loss of MEF2C on these lineage-
specific programs.

Candidate regulatory elements with MEF2C-dependent chromatin accessibility display enhancer activity in zebrafish

Our integrated chromatin accessibility data (Fig. 3) and developmental trajectory analyses
with MIRA (Fig. 4) highlighted changes in chromatin accessibility during heart tube
development that were MEF2C-dependent and segment-specific. We hypothesized that
these regions contain regulatory elements that are necessary for driving segment-specific
gene expression. To identify enhancers, we applied a series of filtering criteria to our
integrated multiomic data and selected 12 candidates of the highest priority to screen
based on the Log2FC value of peak accessibility and linkage to genes of interest (Fig. 5a,
Supplementary Table 6). We enumerated these candidate regions with the prefix "MVEB,"
after the initials of authors Muncie-Vasic and Brower. Overlaying our chromatin
accessibility data at these regions with published ChIP-seq datasets[20,27] confirmed that
each of our candidates had clear and strong occupancy of both MEF2C and H3K27ac, a
histone post-translational modification that marks active enhancers (Fig. 5b). Of note, one
of our ventricular-specific candidates, MVEB8, was located within the *Nppa/Nppb* super
enhancer cluster[28]. Furthermore, two of the candidates we identified from differential
accessibility in IFT-CMs overlapped with regions found in the VISTA enhancer database[29],
with demonstrated enhancer activity in the hearts of E11.5 mouse embryos (Fig. 5c-d).

To determine whether our candidate regions are *bona fide* enhancers with activity
in early heart tube development, we employed Tol2 transgenesis in zebrafish[30,31], a
relatively high-throughput *in vivo* screening system (Fig. 5e). We reasoned that because
Mef2c has orthologs in zebrafish (*mef2ca* and *mef2cb*), and because there is a high
degree of overlap between the key cardiac transcriptional regulators in both species[32],
there should be concordance between the activity of our candidate regions in mouse and
zebrafish. Indeed, specific regulatory elements have been shown to exhibit activity in the
heart of both species[33–35].

Upon screening, seven of eleven candidates displayed enhancer activity in
developing zebrafish hearts (Fig. 5f, Extended Data Fig. 5a-b), with one of our candidates

being excluded due to a highly repetitive "GT" sequence. The two candidates that
overlapped with regions found in the VISTA enhancer database (MVEB1, MVEB2)
exhibited activity in the heart by 72 hours post-fertilization (hpf) (Fig. 5f). Additionally, two
candidates from our IFT-CM data (MVEB3 and MVEB6), two from our V-CM data (MVEB7
and MVEB8), and one from our OFT-CM data (MVEB12) demonstrated robust and
specific activity in the zebrafish heart at 72 hpf (Fig. 5f, Extended Data Fig. 5a-b). To our
knowledge, four of these latter candidates (MVEB3, MVEB6, MVEB7, and MVEB12) have
not been previously reported and represent novel, MEF2C-dependent enhancers with
activity in the developing heart. Interestingly, we also observed a range in the timing of
enhancer activity in the embryonic zebrafish heart. Candidates MVEB2 and MVEB6 were
clearly detectable as early as 24 hpf, while the others were not active until 72 hpf (Fig.
5g-h). This variation in the onset of enhancer activity could result from differences in the
timing of accessibility of these regulatory elements, different sensitivity to MEF2C levels,
or cooperation with other TFs or transcriptional regulators.
276
Loss of MEF2C induces an overactive posteriorized gene regulatory network driven
by NR2F2 and GATA4
Our data indicate that loss of MEF2C causes a posteriorized gene signature (Fig. 2h-j)
and chromatin accessibility landscape (Fig. 3d, 3f) in the developing heart tube. Moreover,
as previously described[36], the expanded but aberrantly formed inflow tract that forms in
Mef2c KO embryos (Fig. 1c) indicates that loss of MEF2C leads to a more complex heart
tube phenotype than a simple failure of cardiac differentiation and growth. To determine
what transcriptional regulators may drive the inflow tract malformation, we performed TF
binding motif enrichment analysis on the regions of chromatin that gain accessibility in
the *Mef2c* KO IFT-CMs at E8.5. We found that Nuclear Receptor (NR) and GATA motifs
were among the most enriched (Fig. 6a). From this observation, we hypothesized that
MEF2C co-binds certain regulatory elements with NR or GATA factors in the WT
condition, and that these NR and GATA factors become free to bind different regulatory
elements with loss of MEF2C. Consistent with this hypothesis, we found that nearly half
of the sites of accessible chromatin that mark WT IFT-CMs and contain an NR or GATA

motif also contain a MEF2 motif, whereas the vast majority of sites that gain accessibility
in *Mef2c* KO and contain an NR or GATA motif do not (Fig. 6b, Extended Data Fig. 6a).

These data are particularly interesting given the importance of GATA factors, and
in particular GATA4, in cardiac development[37] and the knowledge that nuclear receptor
subfamily 2 group F member 2 (NR2F2, also known as COUP-TFII) is a necessary driver
of atrial development[26,38]. Using publicly available ChIP-seq datasets[38,39], we found
NR2F2 and GATA4 occupancy at a number of the regions of chromatin that gained
accessibility in the *Mef2c* KO IFT-CMs at E8.5 (Extended Data Fig. 6b-c), suggesting that
these DARs could indeed be sites of potential NR2F2 and GATA4 regulation. To
determine whether altered NR2F2 and GATA4 activity may drive the posteriorization of
the heart tube, we next asked if NR2F2 and GATA4 target genes were upregulated in the
inflow tract of *Mef2c* KO embryos. Using publicly available datasets to identify the target
genes of each TF[40,41], we calculated the odds ratio (OR) of finding NR2F2 or GATA4
target genes compared to non-target genes amongst our up-regulated DEGs at E8.5. We
found strong evidence that NR2F2 (OR = 1.52, p = 1.79e-4) and GATA4 (OR = 4.23, p <
2.2e-16) targets were enriched amongst the upregulated genes (Fig. 6c, Extended Data
Fig. 6d). Next, we took the list of NR2F2 and GATA4 target genes that were upregulated
(Supplementary Table 7) and calculated a module score[42] for the expression of these
genes in our dataset. In agreement with our working model that these NR2F2 and GATA4
targets are driving the posteriorized heart tube phenotype in *Mef2c* KO, we found the
NR2F2 and GATA4 module scores were increased in all segments of the heart tube at
E8.5 (Fig. 6d, Extended Data Fig. 6e).

To understand how altered NR2F2 and GATA4 activity upon loss of MEF2C affects
the entire transcriptional network, we used CellOracle[43] to construct GRNs for the WT
and *Mef2c* KO IFT-CMs at E8.5 and E9 (Fig. 6e). In brief, CellOracle uses integrated gene
expression and chromatin accessibility data to infer interactions between TFs and target
genes based on the accessibility of a given TF's binding motif near a potential target gene
and the correlation of expression between the TF and the potential target. To validate that
these inferred CellOracle networks effectively captured the GRNs that govern inflow tract
development, we simulated the knockout of *Mef2c in silico* in the E8.5 WT IFT-CMs
network and compared the predicted changes in gene expression to the measured

changes captured in our E9 dataset (Fig. 6f-g). We found that the networks correctly
predicted the gene expression changes with 74% accuracy, even with a relatively
permissive DEGs cutoff of Log2FC >= 0.75 (Fig. 6g). As expected, the prediction
accuracy increased with more stringent Log2FC cutoffs, and vice versa (Supplementary
Table 8).

We next visualized the subnetworks of 12 core cardiac TFs and the top 100 DEGs
in the WT and *Mef2c* KO inflow tracts (Fig. 6h). These subnetworks revealed that loss of
MEF2C causes a network-wide reorganization of gene regulation. First, we observed that
MEF2C may be critical for coordinating cooperative regulation by multiple cardiac TFs.
For instance, in the WT network, MEF2C, MEF2A, and TBX20 all have an inferred
activating effect on *Ttn* expression. However, in the *Mef2c* KO network, both MEF2A and
TBX20 lose their activating interaction with *Ttn.* Similarly, in the WT network, MEF2C,
MEF2A, and NKX2-5 are all inferred activators of *Myh6*, but in the *Mef2c* KO network,
MEF2A and NKX2-5 are no longer predicted to activate *Myh6* expression. These data
suggest that MEF2C is required for cooperative activation with other cardiac TFs in the
developing heart tube, similar to what was shown to occur during cardiomyocyte
reprogramming *in vitro*[44]. Additionally, we found that loss of MEF2C may alter the scope
and strength of interactions between other cardiac TFs and their gene targets. For
example, in the WT network, GATA4 and MEIS2 exhibit competitive regulation of *Fbxo32*,
which encodes a muscle-specific ubiquitin-E3 ligase that is critical for sarcomeric function
and associated with dilated cardiomyopathy[45,46]. This competitive relationship is
maintained in the *Mef2c* KO network, but the interactions are strengthened in both
directions. Finally, we observed that the activity of NR2F2 is substantially altered in the
Mef2c KO network, with a larger number of gene targets and apparent strengthening of
both activating and inhibitory interactions. Most notably, the activating interaction between
NR2F2 and *Angpt*1 expression is much stronger in the *Mef2c* KO network than in the WT
network. Interestingly, the activation of *Nr2f2* itself by TBX5 is lost upon *Mef2c* KO.

Next, we further narrowed in on the direct interactions between NR2F2 and GATA4
and their target genes in the WT and *Mef2c* KO inflow tract networks (Fig. 6i, Extended
Data Fig. 6f). From these analyses we observe similar behavior for both TFs –
approximately half of the WT interactions are maintained in the *Mef2c* KO, while nearly

twice as many new direct interactions are established. This is consistent with our working
model that in response to loss of MEF2C, aberrant NR2F2 and GATA4 activity regulate
additional genes that drive the inflow tract phenotype in the posterior of the heart tube.
357
Reduction of NR2F2 dosage partially rescues the heart tube phenotype in *Mef2c*
KO embryos
Given our observation that loss of MEF2C causes aberrant NR2F2 activity in the IFT-CM
GRNs, we theorized that reducing the dosage of NR2F2 may rescue the inflow tract
phenotype in *Mef2c* KO embryos. To test this possibility, we generated *Mef2c*$^{+/-}$;*Nr2f2*$^{+/-}$
male mice and crossed them to *Mef2c*$^{+/-}$ females. Although *Nr2f2*$^{+/-}$ embryos have no
apparent heart tube phenotype at E9.5 (Fig. 6j, Extended Data Fig. 7a), pups of this
genotype were recovered at lower numbers than expected Mendelian ratios[26], suggesting
that *Nr2f2* is a dosage-sensitive gene. We observed a notable partial rescue in ventricle
and inflow tract development at E9.5 (18-24 somites) in the *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$ embryos
compared to *Mef2c*$^{-/-}$ (Fig. 6j-k), although the extent of the rescue varied (Extended Data
Fig. 7b-c). The *Mef2c* KO single-ventricle phenotype persisted in *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$
embryos, but these ventricles were larger and more expanded than in *Mef2c*$^{-/-}$ *embryos.*
Moreover, the posterior structures of the heart tube in *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$ embryos more
closely resembled WT inflow tracts with expanded and looping prospective atria,
compared to the completely disrupted inflow tract in *Mef2c*$^{-/-}$ embryos (Fig. 6j-k). This
partial rescue of the ventricle and inflow tract in *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$ embryos is evidence that
the *Mef2c* KO phenotype in the posterior heart tube is driven, at least in part, by an altered
gene regulatory network with increased and aberrant NR2F2 activity.
377
DISCUSSION
In this study we demonstrated that MEF2C controls complex, segment-specific GRNs that
regulate gene expression and chromatin accessibility in the developing linear heart tube.
We found that loss of MEF2C causes a posteriorization of the heart tube and revealed
novel segment-specific regulatory elements with enhancer activity. These results address
the long-standing question of how a given TF can play distinct regulatory roles in different
cell types and at particular developmental stages. For instance, decades of work have

established that FHF progenitors give rise to the left ventricle and atria, while the SHF is
subdivided into the aSHF, which produces the outflow tract and right ventricle, and the
pSHF, which contributes to the atria[7–11]. Although these progenitor cell types have
different molecular signatures and give rise to individual structures with specific
morphology and function, they largely rely on a shared set of cardiac TFs to regulate their
development. Our data illustrate that these core cardiac TFs operate in unique GRNs for
each of the inflow tract, ventricular, and outflow tract lineages in the developing heart
tube. We found that distinct chromatin landscapes and regulatory elements are important
determinants for setting up these GRNs, but other embryological phenomena are likely
to be involved, such as morphogen signaling from differing neighbor cell types,
heterochronic timing of differentiation[47], and mechanical phenomena introduced by tissue
morphogenesis and the initiation of cardiomyocyte beating[48–50].

Using the GRNs inferred from our multiomic data, we identified a genetic
interaction between *Mef2c* and *Nr2f2* in the inflow tract of the developing heart tube.
Previous work from our lab showed that TF occupancy at specific loci is highly dependent
upon both TF-TF and TF-DNA affinities, and that TF-TF interactions not only serve to
regulate gene expression, but prevent the redistribution of one or both TFs to exogenous
loci[51]. The results we present here lend further support to those concepts. By removing
MEF2C, the set of possible TF-TF interactions is limited and causes altered activity of
partner TFs, including NR2F2 and GATA4. Our comparison of the WT and *Mef2c* KO
GRNs revealed that the transcriptional networks regulating early heart development are
highly sensitive to TF perturbations, with loss of MEF2C causing a reorganization of heart
tube GRNs that is not limited to direct MEF2C connections, and reduction of NR2F2
dosage capable of partially rescuing the heart tube malformations. Thus, these GRNs
may be implemented as a powerful tool for predicting the network-wide consequences of
genetic mutations that underlie CHDs, potentially leading to unexpected molecular targets
for intervention and correction.

Our approach provides a generalizable framework for integrating snRNA-seq and
snATAC-seq data to build GRNs that provide specific and comprehensive models of gene
regulation. By constructing these GRNs using both WT and mutant data, we were able to
immediately test predictions made from the WT networks and validated the inferred

regulatory interactions. These tools also allow specific networks to be built for particular
cell or tissue types, provided there are enough cells in the group of interest to generate
robust inferences. Moreover, the experimental mutant data revealed how the overall
networks were altered – both in ways that would and would not be predicted from the WT
networks. These unexpected differences help to illuminate indirect or transactivating gene
regulation. Discoveries from these networks lead to many new hypotheses about
regulatory elements, TF interactions, DNA binding, and chromatin organization. As these
hypotheses are experimentally tested, the GRNs can be refined, leading to higher-fidelity
models with greater predictive power. Such higher-fidelity networks will yield important
new insights into the transcriptional regulation of developmental processes and will have
the ability to reveal precisely how these processes are altered in the context of disease-
causing mutations.

MATERIALS AND METHODS

Mouse Models

All mouse studies were performed in strict compliance with the UCSF Institutional Animal Care and Use Committee. Mice were housed in a standard 12-hour light/dark animal husbandry barrier facility at the Gladstone Institutes. The *Mef2c*[+/-] allele used was the same as the original generated by Lin et al.[13]. The *Smarcd3*-F6-eGFP reporter allele was generated in our lab and described previously[14]. The *COUP-TFII lacZ* knock-in allele (abbreviated here as *Nr2f2*[+/-]) was obtained from the Mutant Mouse Resource & Research Centers and was originally generated by Takamoto et al.[52]. The *Mef2c*[+/-] and *Nr2f2*[+/-] lines were maintained in the C57BL6/J background (Jackson Laboratory #664). The *Smarcd3*-F6-eGFP allele was crossed into our *Mef2c*[+/-] line from a mixed background of C57BL6/J (Jackson Laboratory #664) and CD-1 (Charles River #022).

Timed Embryo Dissections and Genotyping

To achieve timed matings, male and female mice were housed together in the evening and pregnancy was assessed by vaginal plug the following morning. Gestational stage was determined relative to noon on the day of plug detection, defined as day E0.5. Embryos were dissected and, at later stages when yolk was present, also de-yolked, in ice-cold PBS (Life Technologies, 14190250) with 1% fetal bovine serum (FBS; Thermo Fisher Scientific, 10439016) on ice. The posterior trunk was removed by microdissection for all embryos at E8.5 or later that were collected for imaging (IF or FISH via HCR) to ensure unobstructed views of the heart tube. DNA for genotyping was extracted using QuickExtract DNA Extraction Solution (Lucigen, QE09050) from harvested yolk sac tissue, if available, or else from a micro-dissected piece of the extra-embryonic anterior proximal region. For embryos collected for 10x Multiome library construction, rapid genotyping was performed while embryos remained on ice using Phire Green Hot Start II PCR Master Mix (Thermo Fisher Scientific, F126S), according to manufacturer's protocols. For all other experiments, genotyping was performed after embryo fixation using GoTaq Green Master Mix (VWR, PAM7122), according to manufacturer's protocols. PCR products were run on a 1% agarose gel (Thermo Fisher Scientific, BP164-500) with Biotium GelRed Nucleic Acid Gel Stain (Thermo Fisher Scientific, NC9594719) used for

DNA detection. Primers used for genotyping were ordered from Integrated DNA

Technologies (IDT). For genotyping the *Mef2c* allele, a three-primer mix was used (P1:

ACTGGCTGAGAGTTGTACCCAC, P2: GCGTGATTTGCTAGATAGTGGTAGAC, P3:

ATGTGGAATGTGTGCGAGGC), resulting in a single 314 bp band for WT, a single 242

bp band for *Mef2c*[-/-], and both bands for *Mef2c*[+/-]. For genotyping the *Nr2f2* allele, a three-

primer mix was used (P4: CATCCGGGATATGTTACTGTCCGG, P5:

TGGGGAAGCTAAGTGTTGATGTGATTCC, P6: GCCGTGGGTTTCAATATTGGCTTC),

resulting in a single 786 bp band for WT, a single ~1600 bp band for *Nr2f2*[-/-], and both

bands for *Nr2f2*[+/-].

Whole-Mount Embryo Immunostaining and Imaging

Embryos collected for immunostaining were fixed for 1 h at room temperature in 4%

paraformaldehyde (PFA) freshly diluted from 16% weight/volume PFA aqueous solution

(Thermo Fisher Scientific, 043368-9M) in PBS (Life Technologies, 14190250). Embryos

were stored at 4°C for up to 1 month prior to immunostaining. To permeabilize and block

non-specific antigens, embryos were incubated at 37°C for 2 h with gentle rotation in a

blocking buffer consisting of PBS (Life Technologies, 14190250), 5% normal donkey

serum (Sigma Aldrich, S30-M), and 0.8% Triton X-100 (Sigma Aldrich, X-100-5ML). This

was then removed and replaced with primary antibodies diluted in the same blocking

buffer and incubated overnight at 37°C with gentle rotation. The next day, primary

antibody solution was removed and embryos were washed 3x 30 min in the same blocking

buffer at 37°C with gentle rotation. Secondary antibodies and DAPI (Abcam, ab228549,

used at 1 µg/mL) were added in a blocking buffer consisting of PBS (Life Technologies,

14190250), 5% normal donkey serum (Sigma Aldrich, S30-M), and 0.4% Triton X-100

(Sigma Aldrich, X-100-5ML), and incubated for 3-4 h at 37°C with gentle rotation.

Secondary antibody solution was removed and embryos were washed 3x 30 min in PBS

(Life Technologies, 14190250) with 0.1% Tween-20 (Thermo Fisher Scientific,

J20605.AP) at 37°C with gentle rotation. Embryos were imaged in PBS using an upright

epifluorescence microscope (Leica MZFLIII, Leica DFC 3000G, Lumen Dynamics XCite

120LED) and acquisition software LASX (Leica). Primary antibodies used were sheep

polyclonal MEF2C (R&D Systems, AF6786, used at 1:250) and rabbit polyclonal cardiac

Troponin T (Thermo Fisher Scientific, 15513-1-AP, used at 1:250). Secondary antibodies

used were Donkey anti-Sheep IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa

Fluor™ 647 (Thermo Fisher Scientific, A-21448, used at 1:1000) and Donkey anti-Rabbit

IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 555 (Thermo

Fisher Scientific, A-31572, used at 1:1000).

Whole-Mount Embryo Fluorescence *In Situ* Hybridization via Hybridization Chain

Reaction and Imaging

The protocol for *in situ* hybridization via hybridization chain reaction (HCR™) was adapted

from the optimized manufacturer's protocol for whole-mount mouse embryos (Molecular

Instruments). Embryos were fixed overnight at 4°C in 4% PFA freshly diluted from 16%

weight/volume PFA aqueous solution (Thermo Fisher Scientific, 043368-9M) in PBS (Life

Technologies, 14190250). Embryos were then washed 3x 5 min in PBS (Life

Technologies, 14190250) with 0.1% Tween-20 (Thermo Fisher Scientific, J20605.AP),

abbreviated PBST, and then processed through 10 min serial incubations in a dehydration

series of 25%, 50%, 75%, and 100% methanol solutions in PBST on ice. Embryos were

stored in 100% methanol at −20°C for up to 6 months prior to *in situ* hybridization.

Embryos were then rehydrated through 10 min serial incubations in a series of 75%, 50%,

25%, and 0% methanol solutions in PBST on ice. Embryos were then brought to room

temperature and permeabilized by 10 min incubation with 10 µg/mL Proteinase K (Sigma

Aldrich, 70663-5) in PBST, washed 2x 5 min in PBST at room temperature, post-fixed in

4% PFA freshly diluted from 16% weight/volume PFA aqueous solution (Thermo Fisher

Scientific, 043368-9M) in PBS (Life Technologies, 14190250), and then washed 3x 5 min

in PBST at room temperature. Embryos were pre-hybridized in Probe Hybridization Buffer

(Molecular Instruments) for 5 min at room temperature followed by an additional 30 min

at 37°C with fresh buffer. Single or multiplexed hybridization probes were then diluted in

pre-warmed Probe Hybridization Buffer (Molecular Instruments) and embryos were

hybridized for 48 h at 37°C. Embryos were then washed 4x 15 min with pre-warmed Probe

Wash Buffer (Molecular Instruments) at 37°C, followed by 2x 5 min with 5xSSCT (diluted

in nuclease-free water from 20xSSCT, Thermo Fisher Scientific, 15557044) at room

temperature. During washes, amplification hairpins were snap-cooled by heating to 95°C

for 90 s and then allowed to cool to room temperature in the dark. Embryos were pre-
amplified in Amplification Buffer (Molecular Instruments) for 5 min at room temperature.
Snap-cooled hairpins and DAPI (Abcam, ab228549, used at 1 μg/mL) were diluted in
Amplification Buffer (Molecular Instruments) and embryos were incubated in this solution
overnight in the dark at room temperature. The following day, embryos were washed
through a series of 5xSSCT washes (diluted in nuclease-free water from 20xSSCT,
Thermo Fisher Scientific, 15557044) in the dark at room temperature (2x 5 min, 2x 30
min, 1x 5 min). Embryos were imaged in PBS using an upright epifluorescence
microscope (Leica MZFLIII, Leica DFC 3000G, Lumen Dynamics XCite 120LED) and
acquisition software LASX (Leica). To maximize the utility of collected embryos, probes
were then stripped by overnight incubation in 80% formamide in water (Thermo Fisher
Scientific, BP228100) at room temperature, followed by 3x 5 min washes in PBST at room
temperature. Embryos could then be probed for additional targets, beginning the protocol
again at the pre-hybridization step. Probes used in this study were ordered from the
Molecular Instruments catalog, when possible, or custom-designed and manufactured by
Molecular Instruments. Hybridization probes used were: *Tnnt2*-B3 (used at 4 nM), *Ttn*-B2
(used at 4 nM), *Nppa*-B2 (used at 20 nM), *Nkx2.5*-B1 (used at 20 nM), *Tbx5*-B1 (used at
20 nM), *Gata4*-B1 (used at 20 nM), *Wnt2*-B2 (used at 20 nM), *Tdgf1*-B2 (used at 20 nM).
Amplification probes used were all used at 60 nM: B1-Alexa Fluor™ 647, B2-Alexa
Fluor™ 546, B2-Alexa Fluor™ 647, B3-Alexa Fluor™ 488.
541

Embryo Preparation for 10x Multiome snRNA-seq and snATAC-seq
For E7.75 embryos, whole embryos were dissected and harvested for single-nucleus
library generation. For E8.5 and E9 embryos, the head folds and posterior trunk were
removed by microdissection prior to harvesting for library generation to enrich the relative
capture of cardiac cell types. Two embryos were collected per genotype per embryonic
stage. Following dissection and rapid PCR genotyping, embryo samples selected for the
multiome experiment were incubated in 200 μl TrypLE Select (Thermo Fisher Scientific,
12563-011) for 5 min at 37°C, triturated gently by pipetting up and down, and then
incubated an additional 3 min at 37°C. The dissociated cell suspension was quenched
with 600 μl of PBS (Life Technologies, 14190250) with 1% FBS (Thermo Fisher Scientific,

10439016), singularized by passage through the 35 μm mesh of a 5 mL Falcon™ Round-
Bottom Polystyrene Test Tube with Cell Strainer Snap Cap (Thermo Fisher Scientific, 08-
771-23), pelleted by centrifugation at 300 g for 5 min at 4°C, and resuspended in 50 μl of
PBS (Life Technologies, 14190250) with 0.04% bovine serum albumin (BSA, Sigma
Aldrich, A1595). At this stage, the manufacturer's protocol for Nuclei Isolation for Single
Cell Multiome ATAC + Gene Expression Sequencing, Appendix, Low Cell Input Nuclei
Isolation (10x Genomics, CG000365 Rev B) was followed exactly to prepare nuclei for
library generation.

Generation of snRNA-seq and snATAC-seq Libraries and Sequencing
Libraries for snRNA-seq and snATAC-seq were prepared according to the manufacturer's
protocol for Chromium Next GEM Single Cell Multiome ATAC + Gene Expression (10x
Genomics, CG000338 Rev B), using the 10x Genomics Chromium controller, Chromium
Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Bundle (10x
Genomics, PN-1000283), Chromium Next GEM Chip J Single Cell Kit (10x Genomics,
PN-1000234), Single Index Kit N Set A (10x Genomics, PN-1000212), and Dual Index Kit
TT Set A (10x Genomics, PN-1000215). Nuclei isolated from multiome samples were
processed directly into the snATAC-seq transposition reaction and then captured in 10x
Genomics GEMs via the Chromium Controller. GEMs were stored at -80°C for up to 4
weeks, allowing for collection of replicate embryos of each genotype at each development
stage. All subsequent library preparation steps were performed together for all embryos
of a given developmental stage to reduce the likelihood of batch artifacts. A targeted
maximum recovery of 10,000 cells per sample were loaded onto the 10x Genomics
Chromium controller instrument and each sample was indexed with a unique sample
identifier (Single Index Kit N Set A for snATAC-seq libraries, Dual Index Kit TT Set A for
snRNA-seq libraries). Final libraries were quality-controlled using an Agilent Bioanlyzer
instrument with Agilent High Sensitivity DNA Kit (Agilent, 5067-4626). The DNA
concentration of each library was measured using KAPA Library Quantification qPCR Kit
(Thermo Fisher Scientific, 50-196-5234), and then libraries were pooled and sequenced
on NovaSeq6000 S4 lanes (Illumina). All snRNA-seq libraries were sequenced to a depth
of at least 20,000 mean raw aligned reads per cell and snATAC-seq libraries were

sequenced to a depth of at least 35,000 mean raw aligned read pairs per cell (most
>60,000).

Processing of Raw Sequencing Data
Raw sequencing reads were processed using the 10x Genomics Cell Ranger Arc v2.0.0
pipeline. Reads were demultiplexed using cellranger-arc mkfastq and aligned with
cellranger-arc count to the mm10 reference genome containing an additional sequence
for eGFP to map reads of the *Smarcd3*-F6-eGFP reporter transcript.

Seurat Analysis of snRNA-seq Data
Gene expression data were analyzed using Seurat v4.3.0[15] in R. Datasets for each
developmental stage (E7.75, E8.5, E9) were analyzed as separate Seurat objects.
Counts matrices from each sample that were output by the Cell Ranger Arc pipeline were
used as inputs to the Read10X() and CreateSeuratObject() functions. Data were filtered
for quality control based on number of detected genes, unique molecular identifier (UMI)
counts, percent ribosomal transcripts, and percent mitochondrial transcripts.
SCTransform v2[16] was used to normalize and scale data, with regressions performed with
respect to percentage of ribosomal transcripts and percentage of mitochondrial
transcripts. Principal component analysis (PCA) was performed, followed by batch
correction with Harmony[53], and cells were clustered and visualized as UMAP embeddings
based on the top 30 principal components using FindNeighbors(), FindClusters(),
RunUMAP(), and DimPlot(). Cell types were annotated using the FindAllMarkers()
function with Wilcoxon rank-sum test (min.pct = 0.25, logfc threshold = 0.25, p_val_adj <
0.05) to identify cluster-specific marker genes (Supplementary Table 1, Extended Data
Fig. 2d-f). Cardiac progenitors, cardiomyocytes, and related mesoderm cell types were
subset into new Seurat objects. PCA analysis, batch correction, visualization, and
clustering were then performed on these new objects as described above. Cell types in
the subset objects were annotated using the FindAllMarkers() function with Wilcoxon
rank-sum test (min.pct = 0.25, logfc threshold = 0.5, p_val_adj < 0.05) to identify cluster-
specific marker genes that enabled more specific cell type labeling than in the full datasets
(Supplementary Table 2, Extended Data Fig. 2g-i). Differential gene expression was

performed between *Mef2c* KO and WT cell types of interest using the FindMarkers()
function with Wilcoxon rank-sum test (min.pct = 0.1, logfc threshold = 0.25, p_val_adj <
0.05). The DotPlot() and RidgePlot() functions was used to visualize gene expression
levels of differentially expressed genes across cell types of interest. The
AddModuleScore() function(42) was used to add module scores for up-regulated (*Mef2c*
KO-vs-WT) NR2F2 and GATA4 target genes and the RidgePlot() function was used to
visualize these module scores in cell types of interest.

ArchR Analysis of Integrated snRNA-seq and snATAC-seq Data
Initial Dataset Processing
Chromatin accessibility and integrated multiome data was analyzed using the ArchR
software package v1.0.2[22] in R. The fragments files output by the Cell Ranger Arc pipeline
were used as inupts to the createArrowFiles() function to create Arrow Files for each
sample. These Arrow Files were used to create individual ArchR projects for
developmental stage (E7.75, E8.5, E9). The gene expression data was added using the
import10xFeatureMatrix() and addGeneExpressionMatrix() functions, using the matching
cell barcodes to combine the snRNA-seq and snATAC-seq data for each cell. Data were
filtered for quality control based on the number of unique fragments, TSS Enrichment
score, expression UMI counts, and number of detected genes. Doublets were filtered
using the addDoubletScores() and filterDoublets() functions. Dimensionality reduction
was performed using ArchR's iterative latent semantic index (LSI) implementation
individually for the snATAC-seq data (varFeatures = 25,000, dimsToUse = 1:30) and
snRNA-seq data (varFeatures = 5,000, dimsToUse = 1:30). Then, the reduced
dimensions were combined using addCombinedDims() and batch correction was
performed on the combined dimensions using Harmony[53]. Cells were then embedded in
UMAP space with addUMAP(), clustered using addClusters(), and visualized using
plotEmbedding(). Cell types were annotated by transferring metadata labels from the
previously analyzed Seurat objects. The getMarkerFeatures() function with Wilcoxon
rank-sum test (FDR <= 0.01, Log2FC >= 0.5) was used to identify cluster-specific marker
genes to confirm cell type labels. Cardiac progenitors, cardiomyocytes, and related
mesoderm cell types were subset into new ArchR projects. Dimensionality reduction,

batch correction, clustering, and visualization were performed as described above. Cell

types in the subset projects were annotated by transferring metadata labels from the

previously analyzed subset Seurat objects. The getMarkerFeatures() function with

Wilcoxon rank-sum test (FDR <= 0.01, Log2FC >= 0.5) was used to identify cluster-

specific marker genes to confirm subset cell type labels. Due to the small number of *Mef2c*

KO OFT-CMs in the E8.5 dataset, an ArchR project with combined E8.5 and E9 samples

was created and processed as described above for downstream analyses/comparisons

of OFT-CMs.

Peak Calling, Differential Peak Testing, and Motif Enrichment

Cells were pseudobulked into replicates based on their cell type and genotype labels (e.g.

WT V-CMs), with two pseudobulked replicates created corresponding to the two embryos

collected for each genotype at each developmental stage. Replicates were created using

the addGroupCoverages() function (minCells = 20, maxCells = 500, sampleRatio = 0.8).

Peaks were then called using addReproduciblePeakSet() with MACS2(23)

(reproducibility = 2, peaksPerCell = 500, minCells = 20). Differential peak accessibility

testing was performed using getMarkerFeatures() with Wilcoxon rank-sum test (FDR <=

0.15, Log2FC >= 0.5 or Log2FC <= -0.5) for each cell type of interest with *Mef2c* KO cells

used as the group of interest and WT cells used as the background. Transcription factor

binding motifs were annotated using addMotifAnnotations() with the Vierstra motif

archetypes[54] and motif enrichments for regions of gained or lost accessibility were

calculated using peakAnnoEnrichment() with cutoffs of FDR <= 0.15 and Log2FC <= -0.5

or Log2FC >= 0.5.

Peak2Gene Linkages and Browser Track Visualization

Peak2Gene Linkages were identified using the addPeak2GeneLinks() and

getPeak2GeneLinks() functions. Code written in this study was used to filter Peak2Gene

links based on presence of a MEF2C motif and linkage to specific genes (i.e. DEGs).

Peak2Gene links were visualized using the plotBrowserTrack() function. The

getGroupBW() function was used to export the accessibility coverage data in bigwig

format for visualization along with published ChIP-seq data using the IGV software
package[55].

MIRA Analysis of Integrated snRNA-seq and snATAC-seq Data

Preparing Merged Datasets

We used the Multimodal models for Integrated Regulatory Analysis (MIRA)[24] python
package to examine dynamic changes in gene expression and chromatin accessibility in
WT and *Mef2c* KO embryos across the time course of developmental stages that we
collected (E7.75, E8.5, E9). We first created a merged Seurat object and ArchR project
containing only our cardiac cell types of interest. We then used sceasy[56] to convert our
Seurat object and the peak matrix from the ArchR project to AnnData formats, which were
used as inputs for the MIRA analyses.

Topic Modeling

The first step of the MIRA pipeline is to train and tune topic models for both the gene
expression and chromatin accessibility data. Model training and tuning was performed
with GPU support on nodes of the Wynton HPC Co-Op cluster at UCSF. For our gene
expression topic model, we used the 4,088 most variable genes (min_disp = 0.1) as the
"exogenous" gene set and top 1,959 variable genes (dispersion threshold > 0.7) as the
"endogenous" gene set for model training. We then set the minimum and maximum
learning rate bounds, tuned the model (iters = 64, max_topics = 55), and MIRA's tuner
selected the best model that was used for all subsequent analyses, which contained 33
gene expression topics. To aid in the interpretation of results, we exported the top 200
genes for each of the 33 gene expression topics. We then repeated these steps to train
and tune the accessibility topic model, using all features (i.e. there are no "exogenous" or
"endogenous" peak sets), which resulted in a model containing 17 accessibility topics. To
aid in the interpretation of these topics, we performed TF binding motif enrichment
analysis on each topic and exported the TF motif enrichment scores (top_quantile = 0.2)
for each of the 17 accessibility topics.

Pseudotime Analysis and Lineage Trajectories

For each of the heart tube segment lineages that we analyzed (OFT, V, IFT), we created
separate subsetted AnnData objects containing only the cell types that contribute to each
lineage. We generated UMAP embeddings for each lineage based on joint
representations of gene expression and accessibility topics. We then performed
pseudotime analysis by calculating the diffusion map, selecting the appropriate number
of diffusion map components, and generated a K nearest neighbor graph based on these
diffusion map components. To create pseudotime trajectories, we randomly selected a
WT cell in the earliest progenitor cell type of each lineage to serve as the initial cell, and
then manually selected terminal cells from the appropriate cell types based on the
structure of each trajectory (for OFT lineage: E9 WT OFT-CM and E9 KO OFT-CM; for V
lineage: E9 WT V-CM; for IFT lineage: E8.5 KO IFT-CM, E9 WT A-CM, E9 WT AVC-CM,
E9 KO A-CM, E9 KO AVC-CM). Given these initial and terminal cells, along with the KNN
pseudotime graph, the probability of reaching each terminal state was calculated for every
cell and these probabilities we parsed to create the bifurcating tree lineage trajectory plots
(stream plots). Gene expression and accessibility topics of interest were then plotted
along these stream plots to interpret the dynamic changes in gene expression and
chromatin accessibility that occur in WT and *Mef2c* KO cells as they differentiate through
each lineage.

Cloning of Reporter Constructions for Zebrafish Transgenesis
To identify candidate enhancer regions to clone, we utilized our E8.5 ArchR dataset
containing integrated chromatin accessibility and gene expression data. We identified all
peaks that lost accessibility in the *Mef2c* KO (3,224), then filtered to the peaks that
demonstrated altered accessibility in only one of the three heart tube segments (2,549),
and that contained a MEF2 binding motif (1,059). Next, we utilized ArchR's Peak2Gene
functionality and filtered down to peaks correlated with DEGs in the *Mef2c* KOs (130) and
were not located in promoter regions (119). Finally, from this list of 119 candidates, we
selected 12 of the highest priority to screen based on the Log2FC value of the change in
peak accessibility and linkage to genes of interest.
To generate the Tg(*MVEB:egfp*)[sfc26] zebrafish lines, 650-900 nucleotide fragments
containing the candidate enhancers were cloned by PCR from mouse genomic DNA.

Candidate nucleotide sequences and the primers used to amplify them are listed in
Supplementary Table 6. The resulting fragments were cloned into the Tol2 transgenic
vector E1b-eGFP-Tol2[30] (RRID: Addgene_37845) using the In-Fusion Cloning system
(Takara Bio). All plasmid sequences were confirmed by long-read sequencing before
zebrafish injection (Plasmidsaurus). We were unable to successfully clone one of our
twelve candidates (MVEB10) due to a highly repetitive "GT" sequence, so our final screen
consisted of eleven candidate enhancers.
744
Zebrafish Transgenesis Injections, Screening, and Imaging
All zebrafish experiments were reviewed and approved by the UCSF Institutional Animal
Care and Use Committee and were performed in accordance with the Public Health
Service Policy on the Humane Care and Use of Laboratory Animals. Zebrafish were
raised under standard laboratory conditions at 28°C. The outbred wildtype zebrafish EKW
(*Danio rerio-Ekwill*) and Tg(*cmlc2:mCherry*)[s890] lines have been described previously[57].
One-cell zebrafish embryos obtained from Tg(*cmlc2:mCherry*)[s890] and wildtype EKW
(Ekwill strain) parents were injected with 1nl of injection mix containing 15pg of candidate-
enhancer-eGFP and 15pg of ß-actin-BFP reporter plasmids along with 30pg of *Tol2*
mRNA. The ß -actin promoter drives expression of BFP strongly in somite muscles and
was used as a positive injection control. As a negative control, we also injected and
screened embryos with an empty reporter construct containing only the minimal E1b
promoter and eGFP. Injected embryos were screened at 24 and 72 hours post-fertilization
(hpf) for eGFP and BFP expression. Representative embryos were imaged for eGFP
expression as a readout of candidate enhancer activity and for mCherry expression as a
myocardial marker. Enhancer activity was scored as a percentage of embryos expressing
eGFP in the heart compared to the total number of BFP-positive embryos. A minimum of
at least 50 BFP+ embryos were screened per construct, except for MVEB1, for which only
17 BFP+ embryos were obtained. The exact numbers screened for each construct are
displayed in Extended Data Fig. 5b.
765
Comparative Analyses with Published Datasets

NR2F2 and GATA4 target genes were downloaded from the Harmonizome[40] and TFLink[41]

databases, respectively. Fisher's exact test was used to determine the Odd's Ratios and

p-values for identifying NR2F2 and GATA4 targets among genes upregulated in *Mef2c*

KO IFT-CMs at E8.5. Publicly available ChIP-seq datasets were downloaded from the

Gene Expression Omnibus (GEO) in wig or bigwig format for MEF2C (GSE124008)[20],

H3K27ac (GSE52386)[27], GATA4 (GSE52123)[39], and NR2F2 (GSE46498)[38]. Datasets

provided in wig format were converted to bigwig format using the Galaxy[58] web tool

wigToBigWig. CrossMap[59] was used together with chain files from the UCSC Genome

Browser[60] to lift genome annotations from mm9 to mm10 reference genome assemblies,

and vice versa. For the browser tracks in Fig. 5b-c, the MEF2C and H3K27ac ChIP-seq

bigwigs were lifted from mm9 to mm10, the Galaxy[58] web tool bigwigAverage was used

to generate a single average MEF2C and H3K27ac occupancy track from the individual

replicate tracks, and the results were displayed along with our pseudobulked snATAC-

seq data using the IGV software package[55]. For the heat maps displayed in Extended

Data Fig. 6b-c, we first created Browser Extensible Data (BED) format files for the gained

DARs in *Mef2c* KO IFT-CMs containing NR or GATA motifs. We then lifted these BED

regions from mm10 to mm9 and used the Galaxy[58] web tools computeMatrix and

plotHeatmap with the downloaded NR2F2 and GATA4 ChIP-seq data to visualize NR2F2

and GATA4 occupancy at the DARs. At the computeMatrix step, missing data was

converted to zeroes. At the plotHeatmap step, kmeans clustering was used to generate

two clusters of DARs (effectively "occupied" and "not occupied") and these were sorted

by decreasing mean value prior to being plotted.

Gene Regulatory Network Inference and Analysis

We used CellOracle[43] to infer GRNs for WT and *Mef2c* KO IFT-CMs at E8.5 and E9. The

GRN inference process included the following steps:

Base GRN Construction

We used OFT-CM, V-CM, IFT-CM, and A-CM snATAC-seq data to construct base GRNs

that define the accessible TF binding motifs near genes for each of the following

conditions: E8.5-WT, E8.5-KO, E9-WT, and E9-KO. Pooling the OFT-CM, V-CM, and A-

CM/IFT-CMs at each timepoint mitigated artifacts caused by data sparsity. The binarized

PeakMatrix output from ArchR was used as an input to Cicero's[61] run_cicero() function

(with default parameters) to compute co-accessibility scores among peak pairs within 500

kb. Output scores were from -1 to 1, where higher scores indicate stronger co-

accessibility. Each dataset generated over 20 million peak pairs for co-accessibility

comparisons. We annotated transcription start sites (TSSs) with their corresponding

genes for accessible peaks using the motif_analysis.get_tss_info() function in CellOracle.

Unlabeled peaks were considered potential regulatory elements. Subsequently, we

applied motif_analysis.integrate_tss_peak_with_cicero() to link unlabeled peaks with

genes, filtering for peak-to-gene pairs with co-accessibility scores > 0.8. TF binding motifs

within each 500 bp peak were identified using the motif_analysis.TFinfo() function in

CellOracle, which converts peak IDs into DNA sequences using the mm10 reference

genome. TF motif scanning used the gimme.vertebrate.v5.0[62] library to produce a set of

TF motifs present in accessible regions within 500kb of each gene. Finally, differentially

accessible chromatin regions in KO-vs-WT were determined for each timepoint (FDR <

0.01) and inaccessible regions were pruned from the base GRNs. The TFs that bind to

motifs present in regions that are co-accessible with and within 500 kb of the TSS of each

gene are inferred to have the potential to regulate that gene.

GRN Inference

Using CellOracle, we constructed GRNs in which edge strengths are quantified based on

the TF-to-gene correlation, constrained by the accessible regions defined in the base

GRN. We inferred GRNs for E8.5 WT, E8.5 KO, E9 WT, and E9 KO IFT-CMs, using the

computed base GRNs to determine permissible TF-to-gene links. GRN inference used

the top 5,000 differentially expressed genes in the snRNA-seq dataset. We used the

CellOracle oracle.get_links() function, with alpha set to 1,000 and otherwise default

parameters, fitting linear models with L2 weight regularization to derive coefficients

relating TF expression to target gene expression. The Bagging Ridge model was used to

fit coefficients, providing posterior distributions for each TF-to-gene relationship. Positive

coefficients indicated activating, while negative coefficients indicated inhibiting

relationships. We then filtered the inferred network to include 12,000 edges with the
largest magnitude coefficients.

Simulation of Mef2c Knockout Effects

To predict the effect of *Mef2c* KO on gene expression, we identified genes directly
regulated by MEF2C in the E8.5 WT IFT-CM network. We predicted that reduced
expression of *Mef2c* in the network would downregulate genes directly activated by
MEF2C and upregulate genes directly inhibited by MEF2C in the E8.5 WT IFT-CM
network. The simulation results were compared against differentially expressed genes in
the E9 snRNA-seq IFT-CM experimental data (KO-vs-WT). Simulations were performed
and prediction accuracy tested with varying Log2FC cutoffs for differentially expressed
genes, as summarized in Supplementary Table 8.

Network Visualization

Whole-network visualizations were generated using Gephi[63], applying the Yifan Hu
layout, with node sizes scaled by betweenness centrality. PyVis[64] was used to visualize
subnetworks that included manually selected TFs (*Mef2a, Mef2c, Nr2f2, Tbx5, Gata4,*
Meis2, Plagl1, Isl1, Pitx2, Tbx20, Hand2, Nkx2-5) and the top 100 differentially expressed
genes (ranked in ascending order by p value) identified by comparing the E8.5 WT and
Mef2c KO IFT-CM snRNA-seq experimental data. Visualizations for direct interactions
between TFs (*Nr2f2* and *Gata4*) and targets in the E8.5 WT and *Mef2c* KO IFT-CMs were
developed using Python's NetworkX package[65], with a circular layout and edge widths
scaled by coefficient values.

Phenotypic Analysis of *Mef2c$^{-/-}$* and *Mef2c$^{-/-}$; Nr2f2$^{+/-}$* Embryos

We conducted comparative phenotypic analysis of *Mef2c$^{-/-}$* and *Mef2c$^{-/-}$; Nr2f2$^{+/-}$* embryos
by performing immunofluorescent staining of cardiac Troponin T and analyzing the
apparent heart tube phenotypes, initially unblinded to embryo genotype. We determined
that the heart tube phenotypes ranged in apparent severity and fell into roughly three
categories, with Category 1 being the most severely malformed and Category 3 the least

severe. We then blinded ourselves to the embryo genotypes and re-examined the

embryos, placing each into the appropriate category of severity.

ACKNOWLEDGEMENTS

J.M.M-V. is grateful for assistance from Dr. Kavitha Rao in learning Seurat, as well as from Drs Ryan Corces and Christina Theodoris for assistance with ArchR and MIRA, respectively. J.M.M-V. is also grateful to all his colleagues in the Bruneau Lab for many insightful discussions, important feedback on directions of the project, and help with experiments. We thank the Gladstone Genomics Core for use and maintenance of the Agilent Bioanalyzer and 10x Genomics Chromium Controller. Portions of this work were performed on the Wynton HPC Co-Op cluster, which is supported by UCSF research faculty and UCSF institutional funds. The authors wish to thank the UCSF Wynton team for their ongoing technical support of the Wynton environment.

AUTHOR CONTRIBUTIONS

J.M.M-V. and B.G.B. conceived and designed the experiments presented in this study. J.M.M-V. performed embryo dissections, whole-mount embryo immunostaining, whole-mount embryo FISH via HCR, embryo imaging, preparation of sequencing libraries, and data analysis via Seurat, ArchR, and MIRA. E.F.B. performed mouse colony maintenance and genotyping. J.M.M-V. and E.F.B. performed cloning of reporter constructs for zebrafish transgenesis. T.S. performed zebrafish transgenesis injections and imaging. A.P.C. performed GRN analyses using CellOracle and newly written code. B.G.B., B.L.B., and J.J.S. supervised and advised. J.M.M-V. prepared the figures and wrote the manuscript with input from the co-authors.

DECLARATION OF INTERESTS

B.G.B. is a founder, shareholder, and advisor of Tenaya Therapeutics and an advisor for Silver Creek Pharmaceuticals. The work presented here is not related to the interests of these commercial entities.

FUNDING

This work was funded by grants from the National Heart, Lung, and Blood Institute (NHLBI) of the National Institutes of Health (F32HL162450 to J.M.M-V.; T32HL007284 to A.P.C.; R01HL160665 and R01HL162925 to J.J.S.; R01HL177462 to B.L.B.;

R01HL114948 and R01HL155906 to B.G.B.). The content of this manuscript is solely the

responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. This research was also supported by the Roddenberry

Foundation (B.G.B.), Additional Ventures (B.G.B.), and the Younger Family Fund

(B.G.B.). J.M.M-V. was also supported by a postdoctoral fellowship from the American

Heart Association together with The Children's Heart Foundation (Grant ID:

24POST1191660, DOI: https://doi.org/10.58275/AHA.24POST1191660.pc.gr.190926).

Sequencing performed at the UCSF CAT was supported by UCSF PBBR, RRP IMIA, and

NIH 1S10OD028511-01 grants. This work was also supported by an NIH/NCRR grant

(C06 RR018928) to the J. David Gladstone Institutes.

DATA AND CODE AVAILABILITY

Raw and processed data for snRNA-seq and snATAC-seq datasets has been deposited

in the NCBI Gene Expression Omnibus (GEO) under accession number GSE280587.

Code used for multiomics analyses, GRNs, and plot generation is available in the GitHub

repository associated with this manuscript: https://github.com/jmmuncie/Muncie-

Vasic_2025.

FIGURE CAPTIONS

Figure 1: MEF2C is expressed throughout the developing heart tube and its loss causes segment-specific defects.

a) Schematic of cardiac progenitors and their contributions to linear heart tube development from cardiac crescent (E7.75) to looped heart tube (E9) stage. b) Immunofluorescent staining of MEF2C (cyan) and cardiac Troponin T (cTnT, magenta) in E7.75, E8.5, and E9 WT embryos. c) Representative images of WT and *Mef2c* KO embryos at E7.75, E8.5 and E9. Cardiac progenitors are marked by the *Smarcd3*-F6-eGFP reporter transgene (green). BF, brightfield. d) Schematic of the methodology and biological insights presented in the current study. Elements of this panel were created in BioRender. B, B. (2024) https://BioRender.com/i72e213. Scale bars = 200 µm. FHF, first heart field; aSHF, anterior second heart field; pSHF, posterior second heart field; LV, left ventricle; RV, right ventricle; V, ventricle; IFT, inflow tract; OFT, outflow tract.

Figure 2: Loss of MEF2C reduces expression of key cardiomyocyte genes across all heart tube segments and alters expression of anterior/posterior markers.

a-c) UMAPs of snRNA-seq data for cardiac progenitors, cardiomyocytes, and related mesoderm subtypes from E7.75 (a), E8.5 (b), and E9 (c) embryos labeled by cell type (left) and genotype/sample ID (right). d-f) Bar plots displaying the number of up-regulated and down-regulated genes in *Mef2c* KO relative to WT in cell types of interest at E7.75 (d), E8.5 (e), and E9 (f). g-i) Dot plots displaying expression of key CM genes and anterior/posterior (A/P) markers at E7.75 (g), E8.5 (h), and E9 (i). j) Fluorescence *in situ* hybridization of key CM genes and A/P markers in E8.5-E9 WT and *Mef2c* KO embryos. Note the reduced expression of CM genes *Tnnt2*, *Ttn*, *Nppa*, and *Nkx2-5*, the expanded expression of posterior markers *Tbx5*, *Gata4*, and *Wnt2*, and the loss of anterior OFT marker *Tdgf1* in *Mef2c* KO embryos compared to WT. Scale bars = 200 µm. CMs, cardiomyocytes; FHF, first heart field; SHF, second heart field; JCF, juxtacardiac field; CrM, cranial mesoderm; PrxM, paraxial mesoderm; LPM, lateral plate mesoderm; SoM, somitic mesoderm; NMPs, neuromesodermal progenitors; KPs, kidney progenitors; ExM, extraembryonic mesoderm; HSCs, hematopoietic stem cells; V-CMs, ventricular cardiomyocytes; IFT-CMs, inflow tract cardiomyocytes; OFT-CMs, outflow tract

cardiomyocytes; aSHF, anterior second heart field; pSHF, posterior second heart field;

PostM, posterior mesoderm; PhM, pharyngeal mesoderm; MixM, mixed mesoderm; A-

CMs, atrial cardiomyocytes; AVC-CMs, atrioventricular canal cardiomyocytes; Pe,

proepicardium; VP, venous pole; *, Genes known to be associated with CHDs[19]; #, Direct

targets of MEF2C based on MEF2C ChIP-seq data[20].

Figure 3: MEF2C regulates chromatin accessibility broadly throughout the heart

tube and in a segment-specific manner.

a) UMAP of integrated snRNA-seq and snATAC-seq data for cardiac progenitors,

cardiomyocytes, and related mesoderm subtypes at E8.5 labeled by cell types

determined from snRNA-seq clustering (left) and genotype/sample ID (right). b) Bar plots

displaying the number of gained and lost DARs in *Mef2c* KO relative to WT cell types of

interest at E8.5. c) Venn diagrams displaying unique and overlapping DARs in the three

heart tube segments. d) Scatter plots displaying the relationship between the Log2 fold

change (Log2FC) values of the DEG and DAR analyses for all identified Peak2Gene

(P2G) links in the three heart tube segments. Dots are colored by the P2G correlation

score. e-f) Genome browser tracks displaying snATAC-seq accessibility profiles at the

Myh6/*Myh7* (e) and *Wnt2* (f) loci for the indicated pseudobulked cell types. DARs are

highlighted and the peaks are indicated by red bars (*Mef2c* KO-vs-WT IFT-CMs). Loops

indicate P2G links, colored by the P2G correlation score. V-CMs, ventricular

cardiomyocytes; IFT-CMs, inflow tract cardiomyocytes; OFT-CMs, outflow tract

cardiomyocytes; aSHF, anterior second heart field; pSHF, posterior second heart field;

LPM, lateral plate mesoderm; PostM, posterior mesoderm; PhM, pharyngeal mesoderm;

NA, cells not available in the snRNA-seq dataset; DAR, differentially accessible region;

DEG, differentially expressed gene.

Figure 4: Each heart tube segment exhibits a distinct MEF2C-depedent

developmental trajectory.

a) Schematic of multimodal Models for Integrated Regulatory Analysis (MIRA)[24] pipeline.

b) UMAP of outflow tract lineage cells plotted by MIRA topic models labeled by cell type

(top) and genotype (bottom). c) Pseudotime plot of outflow tract lineage cells. d) Lineage

trajectory stream plots for outflow tract lineage cells labeled by cell type (top) and
genotype (bottom). e) Stream plots displaying the flow of gene expression (top) and
chromatin accessibility (bottom) topics in the outflow tract lineage cells. Examples of the
top genes and TF binding motifs for dynamic topics are labeled. f-i) Same as (b-e), but
for ventricular lineage cells. j-m) Same as (b-e) and (f-i), but for inflow tract lineage cells.
975

Figure 5: Candidate regulatory elements with MEF2C-dependent chromatin accessibility display enhancer activity in zebrafish.

a) Schematic of selection process to identify candidate MEFC-dependent enhancers from
integrated snRNA-seq and snATAC-seq data. b) Genome browser tracks displaying
snATAC-seq accessibility profiles, MEF2C ChIP-seq occupancy profiles[20], and H3K27ac
ChIP-seq occupancy profiles[27] at example loci containing IFT-specific (left), V-specific
(middle), and OFT-specific (right) candidate enhancers (yellow highlights). c) Genome
browser tracks displaying snATAC-seq accessibility profiles, MEF2C ChIP-seq
occupancy profiles[20], and H3K27ac ChIP-seq occupancy profiles[27] at the *Myh6*/*Myh7*
locus, which contains two candidate enhancers with IFT-specific altered accessibility
(yellow highlights, MVEB1 and MVEB2) that overlap with regions found in the VISTA
Enhancer Browser database[29]. d) Images of E11.5 mouse embryos from the Vista
Enhancer Browser database[29] demonstrating positive enhancer activity of regions that
overlap with candidates MVEB1 and MVEB2. e) Schematic of the Tol2 transgenesis
assay used to screen candidate enhancers in zebrafish. Elements of this panel were
created in BioRender. B, B. (2024) https://BioRender.com/h83r503. f) Representative
ventral view images of Tg(*cmlc2:mCherry*) zebrafish embryos at 72 hours post-
fertilization (hpf) injected with candidate enhancers that demonstrated positive activity in
the heart. Boxed area in the representative brightfield image (left) indicates the
anatomical region of interest captured in the fluorescent images. g) Schematic
representation of the observed onset of enhancer activity for candidate enhancers that
demonstrated positive activity in the heart. h) Representative ventral view images of
Tg(*cmlc2:mCherry*) zebrafish embryos at 24 and 72 hpf injected with *MVEB2:eGFP* or
MVEB6:eGFP reporter constructs. Scale bars = 100 µm.
1000

Figure 6: Loss of MEF2C induces an overactive posteriorized gene regulatory network that is partially rescued by reduced NR2F2 dosage.

a) TF binding motif enrichment analysis for gained DARs in *Mef2c* KO relative to WT IFT-CMs at E8.5. b) Pie charts showing the proportion of lost or gained DARs (*Mef2c* KO-vs-WT) containing NR and MEF2C motifs or only NR motifs. c) Odds ratio analysis for NR2F2 target genes[40] amongst DEGs up-regulated in *Mef2c* KO IFT-CMs. p-value calculated using Fisher's exact test. d) Ridge plot displaying module scores for up-regulated NR2F2 targets in *Mef2c* KO and WT heart tube segments. e) Inferred GRNs constructed for *Mef2c* KO and WT IFT-CMs at E8.5 and E9. Boxed region of E8.5 WT GRN is shown at higher magnification in (f). f) Schematic of *in silico* simulated *Mef2c* KO in the E8.5 WT GRN. g) Results of GRN validation displaying the high accuracy (74%) of predicted relative to measured gene expression changes at E9. h) Visualizations of subnetworks consisting of 12 cardiac TFs and the top 100 DEGs within the WT and *Mef2c* KO E8.5 IFT-CM GRNs. i) Visualization of direct NR2F2 interactions in the WT and *Mef2c* KO E8.5 IFT-CM GRNs. Direct interactions that occur upon *Mef2c* KO are highlighted. #, mis-regulated DEG in *Mef2c* KO IFT-CMs at E8.5; *, Direct target of NR2F2[40]. j-k) Immunofluorescent staining of cardiac Troponin T (cTnT, green) in representative E9.5 embryos (18-24 somites) collected from *Mef2c*$^{+/-}$;*Nr2f2*$^{+/-}$ to *Mef2c*$^{+/-}$ crosses. Boxed regions in (j) are shown at higher magnification in (k). Arrows point to the ventricle, which is expanded in *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$ embryos compared to *Mef2c*$^{-/-}$ embryos. Asterisks mark the atria, which are better developed and have undergone more looping in *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$ embryos compared to *Mef2c*$^{-/-}$ embryos. n=5 *Mef2c*$^{-/-}$ and n=7 *Mef2c*$^{-/-}$;*Nr2f2*$^{+/-}$ embryos from 7 independent litters. Scale bars = 200 µm

REFERENCES

1. Hafner A, Boettiger A. The spatial organization of transcriptional control. Nat Rev Genet. 2023;24(1):53–68.

2. Wagh K, Stavreva DA, Upadhyaya A, Hager GL. Transcription Factor Dynamics: One Molecule at a Time. Annu Rev Cell Dev Biol. 2023;39(1):277–305.

3. Spitz F, Furlong EEM. Transcription factors: from enhancer binding to developmental control. Nat Rev Genet. 2012;13(9):613–26.

4. Lee TI, Young RA. Transcriptional Regulation and Its Misregulation in Disease. Cell. 2013;152(6):1237–51.

5. Mercatelli D, Scalambra L, Triboli L, Ray F, Giorgi FM. Gene regulatory network inference resources: A practical overview. Biochim Biophys Acta (BBA) - Gene Regul Mech. 2020;1863(6):194430.

6. Bruneau BG. Signaling and Transcriptional Networks in Heart Development and Regeneration. Cold Spring Harb Perspect Biol. 2013;5(3):a008292.

7. Kelly RG, Buckingham ME, Moorman AF. Heart Fields and Cardiac Morphogenesis. Cold Spring Harb Perspect Med. 2014;4(10):a015750.

8. Evans SM, Yelon D, Conlon FL, Kirby ML. Myocardial Lineage Development. Circ Res. 2010;107(12):1428–44.

9. Lescroart F, Chabab S, Lin X, Rulands S, Paulissen C, Rodolosse A, et al. Early lineage restriction in temporally distinct populations of Mesp1 progenitors during mammalian heart development. Nat Cell Biol. 2014;16(9):829–40.

10. Devine WP, Wythe JD, George M, Koshiba-Takeuchi K, Bruneau BG. Early patterning and specification of cardiac progenitors in gastrulating mesoderm. eLife. 2014;3:e03848.

11. Meilhac SM, Esner M, Kelly RG, Nicolas JF, Buckingham ME. The Clonal Origin of Myocardial Cells in Different Regions of the Embryonic Mouse Heart. Dev Cell. 2004;6(5):685–98.

12. Martin JF, Schwarz JJ, Olson EN. Myocyte enhancer factor (MEF) 2C: a tissue-restricted member of the MEF-2 family of transcription factors. Proc Natl Acad Sci. 1993;90(11):5282–6.

13. Lin Q, Schwarz J, Bucana C, Olson EN. Control of Mouse Cardiac Morphogenesis and Myogenesis by Transcription Factor MEF2C. Science. 1997;276(5317):1404–7.

14. Krup AL, Winchester SAB, Ranade SS, Agrawal A, Devine WP, Sinha T, et al. A Mesp1-
dependent developmental breakpoint in transcriptional and epigenomic specification of early
cardiac precursors. Development. 2023;150(9):dev201229.

15. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis
of multimodal single-cell data. Cell. 2021;184(13):3573-3587.e29.

16. Choudhary S, Satija R. Comparison and evaluation of statistical error models for scRNA-seq.
Genome Biol. 2022;23(1):27.

17. Tyser RCV, Ibarra-Soria X, McDole K, Jayaram SA, Godwin J, Brand TAH van den, et al.
Characterization of a common progenitor pool of the epicardium and myocardium. Science.
2021;371(6533).

18. Black BL, Olson EN. Transcriptional control of muscle development by myocyte enhancer
factor-2 (MEF2) proteins. Annu Rev Cell Dev Biol. 1998;14(1):167–96.

19. Yang A, Alankarage D, Cuny H, Ip EKK, Almog M, Lu J, et al. CHDgene: A Curated
Database for Congenital Heart Disease Genes. Circ: Genom Precis Med. 2022;15(3):e003539.

20. Akerberg BN, Gu F, VanDusen NJ, Zhang X, Dong R, Li K, et al. A reference map of
murine cardiac transcription factor chromatin occupancy identifies dynamic and conserved
enhancers. Nat Commun. 2019;10(1):4907.

21. Barnes RM, Harris IS, Jaehnig EJ, Sauls K, Sinha T, Rojas A, et al. MEF2C regulates
outflow tract alignment and transcriptional control of Tdgf1. Development. 2016;143(5):774–9.

22. Granja JM, Corces MR, Pierce SE, Bagdatli ST, Choudhry H, Chang HY, et al. ArchR is a
scalable software package for integrative single-cell chromatin accessibility analysis. Nat Genet.
2021;53(3):403–11.

23. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based
Analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137.

24. Lynch AW, Theodoris CV, Long HW, Brown M, Liu XS, Meyer CA. MIRA: joint
regulatory modeling of multimodal expression and chromatin accessibility in single cells. Nat
Methods. 2022;19(9):1097–108.

25. Zhou W, Lin L, Majumdar A, Li X, Zhang X, Liu W, et al. Modulation of morphogenesis by
noncanonical Wnt signaling requires ATF/CREB family–mediated transcriptional activation of
TGFβ2. Nat Genet. 2007;39(10):1225–34.

26. Pereira FA, Qiu Y, Zhou G, Tsai MJ, Tsai SY. The orphan nuclear receptor COUP-TFII is
required for angiogenesis and heart development. Genes Dev. 1999;13(8):1037–49.

27. Nord AS, Blow MJ, Attanasio C, Akiyama JA, Holt A, Hosseini R, et al. Rapid and
Pervasive Changes in Genome-wide Enhancer Usage during Mammalian Development. Cell.
2013;155(7):1521–31.

28. Man JCK, Duijvenboden K van, Krijger PHL, Hooijkaas IB, Made I van der, Vries C de G
de, et al. Genetic Dissection of a Super Enhancer Controlling the Nppa-Nppb Cluster in the
Heart. Circ Res. 2021;128(1):115–29.

29. Visel A, Minovitsky S, Dubchak I, Pennacchio LA. VISTA Enhancer Browser—a database
of tissue-specific human enhancers. Nucleic Acids Res. 2007;35(suppl_1):D88–92.

30. Birnbaum RY, Clowney EJ, Agamy O, Kim MJ, Zhao J, Yamanaka T, et al. Coding exons
function as tissue-specific enhancers of nearby genes. Genome Res. 2012;22(6):1059–68.

31. Fisher S, Grice EA, Vinton RM, Bessling SL, Urasaki A, Kawakami K, et al. Evaluating the
biological relevance of putative enhancers using Tol2 transposon-mediated transgenesis in
zebrafish. Nat Protoc. 2006;1(3):1297–305.

32. Hinits Y, Pan L, Walker C, Dowd J, Moens CB, Hughes SM. Zebrafish Mef2ca and Mef2cb
are essential for both first and second heart field cardiomyocyte differentiation. Dev Biol.
2012;369(2):199–210.

33. Zlatanova I, Sun F, Wu RS, Chen X, Lau BH, Colombier P, et al. An injury-responsive
mmp14b enhancer is required for heart regeneration. Sci Adv. 2023;9(48):eadh5313.

34. Yan R, Cigliola V, Oonk KA, Petrover Z, DeLuca S, Wolfson DW, et al. An enhancer-based
gene-therapy strategy for spatiotemporal control of cargoes during tissue repair. Cell Stem Cell.
2023;30(1):96-111.e6.

35. Kang J, Hu J, Karra R, Dickson AL, Tornini VA, Nachtrab G, et al. Modulation of tissue
repair by regeneration enhancer elements. Nature. 2016;532(7598):201–6.

36. Verzi MP, McCulley DJ, Val SD, Dodou E, Black BL. The right ventricle, outflow tract, and
ventricular septum comprise a restricted expression domain within the secondary/anterior heart
field. Dev Biol. 2005;287(1):134–45.

37. Tremblay M, Sanchez-Ferras O, Bouchard M. GATA transcription factors in development
and disease. Development. 2018;145(20):dev164384.

38. Wu S pin, Cheng CM, Lanz RB, Wang T, Respress JL, Ather S, et al. Atrial Identity Is
Determined by a COUP-TFII Regulatory Network. Dev Cell. 2013;25(4):417–26.

39. He A, Gu F, Hu Y, Ma Q, Ye LY, Akiyama JA, et al. Dynamic GATA4 enhancers shape the
chromatin landscape central to heart development and disease. Nat Commun. 2014;5(1):4907.

40. Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG, et al.
The harmonizome: a collection of processed datasets gathered to serve and mine knowledge
about genes and proteins. Database. 2016;2016(2016):baw100.

41. Liska O, Bohár B, Hidas A, Korcsmáros T, Papp B, Fazekas D, et al. TFLink: an integrated
gateway to access transcription factor–target gene interactions for multiple species. Database.
2022;2022:baac083.

42. Tirosh I, Izar B, Prakadan SM, II MHW, Treacy D, Trombetta JJ, et al. Dissecting the
multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. Science.
2016;352(6282):189–96.

43. Kamimoto K, Stringa B, Hoffmann CM, Jindal K, Solnica-Krezel L, Morris SA. Dissecting
cell identity via network inference and in silico gene perturbation. Nature. 2023;614(7949):742–
51.

44. Stone NR, Gifford CA, Thomas R, Pratt KJB, Samse-Knapp K, Mohamed TMA, et al.
Context-Specific Transcription Factor Functions Regulate Epigenomic and Transcriptional
Dynamics during Cardiac Reprogramming. Cell Stem Cell. 2019;25(1):87-102.e9.

45. Al-Yacoub N, Shaheen R, Awad SM, Kunhi M, Dzimiri N, Nguyen HC, et al. FBXO32,
encoding a member of the SCF complex, is mutated in dilated cardiomyopathy. Genome Biol.
2016;17(1):2.

46. Ghasemi S, Mahdavi M, Maleki M, Salahshourifar I, Kalayinia S. A novel likely pathogenic
variant in the FBXO32 gene associated with dilated cardiomyopathy according to whole-exome
sequencing. BMC Méd Genom. 2022;15(1):234.

47. Rowton M, Perez-Cervantes C, Hur S, Jacobs-Li J, Lu E, Deng N, et al. Hedgehog signaling
activates a mammalian heterochronic gene regulatory network controlling differentiation timing
across lineages. Dev Cell. 2022;57(18):2181-2203.e9.

48. Tyser RC, Miranda AM, Chen C mun, Davidson SM, Srinivas S, Riley PR. Calcium
handling precedes cardiac differentiation to initiate the first heartbeat. eLife. 2016;5:e17113.

49. Barriga EH, Franze K, Charras G, Mayor R. Tissue stiffening coordinates morphogenesis by
triggering collective cell migration in vivo. Nature. 2018;554(7693):523–7.

50. Caldarelli P, Chamolly A, Villedieu A, Alegria-Prévot O, Phan C, Gros J, et al. Self-
organized tissue mechanics underlie embryonic regulation. Nature. 2024;1–8.

51. Luna-Zurita L, Stirnimann CU, Glatt S, Kaynak BL, Thomas S, Baudin F, et al. Complex
Interdependence Regulates Heterotypic Transcription Factor Distribution and Coordinates
Cardiogenesis. Cell. 2016;164(5):999–1014.

52. Takamoto N, You LR, Moses K, Chiang C, Zimmer WE, Schwartz RJ, et al. COUP-TFII is essential for radial and anteroposterior patterning of the stomach. Development. 2005;132(9):2179–89.

53. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and accurate integration of single-cell data with Harmony. Nat Methods. 2019;16(12):1289–96.

54. Vierstra J, Lazar J, Sandstrom R, Halow J, Lee K, Bates D, et al. Global reference mapping of human transcription factor footprints. Nature. 2020;583(7818):729–36.

55. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24–6.

56. Cakir B, Prete M, Huang N, Dongen S van, Pir P, Kiselev VY. Comparison of visualization tools for single-cell RNAseq data. NAR Genom Bioinform. 2020;2(3):lqaa052.

57. Zhang R, Han P, Yang H, Ouyang K, Lee D, Lin YF, et al. In vivo cardiac reprogramming contributes to zebrafish heart regeneration. Nature. 2013;498(7455):497–501.

58. Community TG, Abueg LAL, Afgan E, Allart O, Awan AH, Bacon WA, et al. The Galaxy platform for accessible, reproducible, and collaborative data analyses: 2024 update. Nucleic Acids Res. 2024;52(W1):W83–94.

59. Zhao H, Sun Z, Wang J, Huang H, Kocher JP, Wang L. CrossMap: a versatile tool for coordinate conversion between genome assemblies. Bioinformatics. 2014;30(7):1006–7.

60. Raney BJ, Barber GP, Benet-Pagès A, Casper J, Clawson H, Cline MS, et al. The UCSC Genome Browser database: 2024 update. Nucleic Acids Res. 2023;52(D1):D1082–8.

61. Pliner HA, Packer JS, McFaline-Figueroa JL, Cusanovich DA, Daza RM, Aghamirzaie D, et al. Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin Accessibility Data. Mol Cell. 2018;71(5):858-871.e8.

62. Bruse N, Heeringen SJ van. GimmeMotifs: an analysis framework for transcription factor motif analysis. bioRxiv. 2018;474403.

63. Bastian M, Heymann S, Jacomy M. Gephi: An Open Source Software for Exploring and Manipulating Networks. ICWSM. 2009;3:361–2.

64. Perrone G, Unpingco J, Lu H minn. Network visualizations with Pyvis and VisJS. arXiv. 2020;

65. Hagberg AA, Schult DA, Swart PJ. Exploring Network Structure, Dynamics, and Function using NetworkX. In: Varoquaux G, Vaught T, Millman J, editors. Proceedings of the 7th Python in Science Conference (SciPy 2008). 2008. p. 11–6.

Figure 1

a

Cardiac Crescent
E7.75

Linear Heart Tube
E8.5

Looped Heart Tube
E9

FHF → LV/Atria
aSHF → RV/OFT
pSHF → IFT/Atria

b

MEF2C
DAPI

cTnT
DAPI

E7.75

E8.5

E9

c

WT

Mef2c KO

Smarcd3-F6
BF

E7.75

E8.5

E9

OFT
RV
LV
IFT

OFT
V
IFT

d

10x Multiome on Dissociated Embryos

10x Bead

snRNA-seq

snATAC-seq

Suite of Complementary Bioinformatics Analyses

mira
(Deep-Learning Models for Pseudotime Trajectory Inference)

SEURAT
(Gene Expression)

ArchR
(Integrated Gene Expression and Chromatin Accessbility)

CellOracle
(Gene Regulatory Networks)

Biological Insights

Define Heart Tube GRNs

Dissect Normal vs. Aberrant TF Activity

Determine Molecular Basis of Morphological Defects

Identify Novel Segment-Specific Enhancers

Figure 2

Figure 3

Figure 4

a

Multimodal Models for Integrated Regulatory Analysis (MIRA)
(Lynch, Theodoris, et al. 2022)

Input Features Linked by Cell Barcode

Gene Expression   Chromatin Accessibility

GCTAT...

Generate Topic Models via Deep Learning and Probabilisitc Graphical Models

Topics   Genes   Regions

Cells   GEX Topics   ATAC Topics

Construct Joint Representation (KNN) for Data Visualization

Cell States   Lineage Trajectories   Topic Compositions

Outflow Tract Lineage

b

UMAP2 / UMAP1

■ E8.5 aSHF ■ E8.5 OFT-CMs ■ E9 OFT-CMs

■ WT ■ *Mef2c* KO

c

Pseudotime (Arbitrary Units) 0–20

d

■ E8.5 aSHF ■ E8.5 OFT-CMs ■ E9 OFT-CMs

■ WT ■ *Mef2c* KO

e

Enriched Genes
Tdgf1, Wnt11, Mybpc1, etc.
Fzd1, Dlk1, etc.
Meis1, Hand2, etc.

Gene Expression Topics
■ Topic 13 ■ Topic 17 ■ Topic 20
■ Topic 24 ■ Topic 26 ■ Topic 8

Enriched TF Motifs
MEF2, FOS, etc.
ERF, STP, etc.
TEAD, ISL, NKX, etc.

Accessibility Topics
■ Topic 8 ■ Topic 1 ■ Topic 7
■ Topic 3 ■ Topic 2 ■ Topic 5

Pseudotime

Ventricular Lineage

f

UMAP2 / UMAP1

■ E7.75 CMs/FHF ■ E8.5 V-CMs ■ E9 V-CMs

■ WT ■ *Mef2c* KO

g

Pseudotime (Arbitrary Units) 0–10

h

■ E7.75 CMs/FHF ■ E8.5 V-CMs ■ E9 V-CMs

■ WT ■ *Mef2c* KO

i

Enriched Genes
Myl2, Tnnt1, etc.
Tbx2, Zeb1, etc.
Wnt6, Jarid2, etc.

Gene Expression Topics
■ Topic 31 ■ Topic 28 ■ Topic 15
■ Topic 17 ■ Topic 23 ■ Topic 21

Enriched TF Motifs
MEF2, TBX5 NKX, etc.
CTCF, NR2F2, etc.

Accessibility Topics
■ Topic 0 ■ Topic 7 ■ Topic 13
■ Topic 9 ■ Topic 1 ■ Topic 16

Pseudotime

Inflow Tract Lineage

j

UMAP2 / UMAP1

■ E8.5 pSHF ■ E8.5 IFT-CMs ■ E9 AVC-CMs ■ E9 A-CMs

■ WT ■ *Mef2c* KO

k

Pseudotime (Arbitrary Units) 0–25

l

■ E8.5 pSHF ■ E8.5 IFT-CMs ■ E9 AVC-CMs ■ E9 A-CMs

■ WT ■ *Mef2c* KO

m

Enriched Genes
Bmp2k, Fzd6, etc.
Wnt6, Jarid2, etc.
Gjb3, Wnt4, etc.
Angpt1, Nr2f1, etc.
Hand1, Notch2, etc.
Tbx2, Zeb1, etc.

Gene Expression Topics
■ Topic 4 ■ Topic 18 ■ Topic 3 ■ Topic 17
■ Topic 19 ■ Topic 9 ■ Topic 16 ■ Topic 28

Enriched TF Motifs
ERF, RAP, etc.
ERF, STP, etc.
GATA, ELT, etc.
SNAI, TBX5, etc.
GATA, TEAD, etc.

Accessibility Topics
■ Topic 14 ■ Topic 9 ■ Topic 13 ■ Topic 16
■ Topic 12 ■ Topic 6 ■ Topic 5

Pseudotime

Figure 5

Figure 6