Heart tube morphogenesis is regulated by segment-specific gene regulatory networks controlled by MEF2C

Jonathon M. Muncie-Vasic[1], Tanvi Sinha[2], Alexander P. Clark[3], Emily F. Brower[1], Jeffrey J. Saucerman[3,4], Brian L. Black[2,5], Benoit G. Bruneau[1,6,7,8,9,*]

[1]Gladstone Institutes, San Francisco, CA; [2]Cardiovascular Research Institute, University of California, San Francisco, CA; [3]Department of Biomedical Engineering, University of Virginia, Charlottesville, VA; [4]Robert M. Berne Cardiovascular Research Center, University of Virginia, Charlottesville, VA; [5]Department of Biochemistry and Biophysics, University of California, San Francisco, CA; [6]Roddenberry Center for Stem Cell Biology and Medicine, Gladstone Institutes, San Francisco, CA; [7]Department of Pediatrics, University of California, San Francisco, CA; [8]Institute for Human Genetics, University of California, San Francisco, CA; [9]Eli and Edythe Broad Center for Regeneration Medicine and Stem Cell Research, University of California, San Francisco, CA; *Author for correspondence (benoit.bruneau@gladstone.ucsf.edu)

ABSTRACT

The transcription factor MEF2C plays a critical role in the development of the linear heart tube, but the specific transcriptional networks controlled by MEF2C remain largely undefined. To address this, we performed combined single-nucleus RNA- and ATAC-sequencing on wild type and MEF2C-null embryos at distinct stages of development. We identified a broadly "posteriorized" cardiac gene signature and chromatin landscape throughout the heart tube in the absence of MEF2C. By integrating our gene expression and chromatin accessibility data in a deep-learning based model, we were able to construct developmental trajectories for each of the outflow tract, ventricular, and inflow tract lineages and determined how each of these segment-specific trajectories were distinctly altered in the MEF2C-null embryos. We computationally identified potential segment-specific MEF2C-dependent enhancers, and from these candidates, identified novel enhancers with activity in the developing heart tube using transgenesis in zebrafish. Finally, using inferred gene regulatory networks we discovered a genetic interaction between *Mef2c* and the atrial nuclear hormone receptor *Nr2f2*, revealing that the MEF2C-null heart malformations are partly driven by a transcriptional network with increased NR2F2 activity. These studies not only provide a rich description of the genomic regulation of early heart tube development, but provide a generalizable framework for using genetic mutants to dissect the transcriptional networks that govern developmental processes.

INTRODUCTION

Transcription factors (TFs) regulate gene expression to direct cell fate decisions and control developmental processes. Over the course of embryonic development, TFs are typically expressed in specific spatiotemporal patterns that dictate the cell and tissue types that they regulate[1–3]. The activity of a given TF is dependent on a number of factors, including its own expression level, the availability of potential co-factors, the presence or absence of other TFs that may act co-operatively or antagonistically, and the dynamic state of the chromatin landscape in which the TF is acting[4–7]. Together, these phenomena can be conceptualized as transcriptional or gene regulatory networks (GRNs) that can be used to understand the role of a particular TF within a given developmental context[8]. These GRNs can be particularly useful for untangling the complexities of gene regulation and answering open questions in the field of developmental biology, such as a how a single TF may have different activity within its expression domain.

The developing linear heart tube presents an excellent model for exploring how a single TF may play distinct roles within different regions of a developing tissue. The TFs that are critical for cardiac development have been clearly identified and their expression patterns are well-defined[1,9–11]. Moreover, lineage tracing studies and subsequent molecular analyses have revealed the distinct progenitor cell types that contribute to the various segments of the linear heart tube (Fig. 1A). Cardiac progenitors originate in the mesoderm that arises during gastrulation[12–14]. By E7.75 in the mouse, this cardiogenic mesoderm forms a morphologically distinct ridge of cells underneath the early head folds termed the cardiac crescent[15]. The cells that make up the protruding rostral aspect of the crescent arrive to this position first and are termed the first heart field (FHF), which later gives rise to the left ventricle and atria[1,10,11,16,17]. Cells located more posterior and medial to the FHF in the cardiac crescent are termed the second heart field (SHF) and primarily produce the outflow tract and right ventricle, along with contributing to the atria[10,11,16–18].

To explore in detail how a specific TF can have distinct roles within different parts of a developing organ, we focused on a key cardiac transcription factor, MEF2C. In mice, loss of MEF2C leads to cardiac defects that are embryonic lethal by mid-gestation at embryonic day E10.5[19]. Interestingly, although MEF2C is expressed throughout the

entire cardiac crescent (FHF and SHF) and developing heart tube from E7.75 to E9 (Fig.

1B, Fig. S1A), the loss of MEF2C causes distinct defects in different segments of the

heart tube. At E9, the outflow tract is severely hypoplastic and only a single hypoplastic

ventricle has formed, but by contrast, the inflow tract appears to be expanded and

improperly patterned (Fig. 1C). These segment-specific defects suggest that MEF2C

plays distinct regulatory roles in the outflow tract, ventricles, and inflow tract.

In order to dissect the segment-specific regulation of heart tube morphogenesis by

MEF2C, we used the 10x Multiome platform to perform combined single-nucleus RNA-

seq and ATAC-seq (snRNA-seq and snATAT-seq) on wild-type (WT) and MEF2C

knockout (KO) mouse embryos at key stages throughout heart tube development.

Specifically, we collected two WT and two MEF2C KO stage-matched embryos at E7.75,

E8.5, and E9 (Fig. S1B). All embryos carried the *Smarcd3*-F6-eGFP transgene reporter

to mark cardiac progenitors[20]. At E7.75, we harvested the entire embryos, whereas, at

E8.5 and E9 we removed the headfolds and posterior trunk via microdissection in order

to increase the proportion of cardiac progenitors captured in the 10x Multiome assay. By

taking advantage of the ability to match the transcriptomic and chromatin accessibility

data for every cell in the dataset, we applied a suite of complementary bioinformatics tools

to construct GRNs for cardiac progenitor lineages, precisely defined the role of MEF2C

in regulating gene expression and chromatin accessibility in the developing heart tube,

identified novel MEF2C-dependent enhancers, and discovered a genetic interaction that

underlies the molecular basis of the expanded inflow tract phenotype in MEF2C KO

embryos (Fig. 1D).

RESULTS

Loss of MEF2C leads to broad changes in gene expression throughout the heart

tube, including mis-regulation of key cardiomyocyte genes and anterior/posterior

markers

To begin examining how MEF2C regulates the morphogenesis of specific heart tube

segments, we used Seurat[21] to determine the changes in gene expression that result

from MEF2C KO. We began by creating separate Seurat objects for the embryos profiled

at each timepoint (E7.75, E8.5, and E9). Following pre-processing, quality control,

normalization with scTransform[22], and dimensionality reduction, we performed unbiased
Louvain clustering and plotted the data in uniform manifold approximation and projection
(UMAP) space. We then used the expression of marker genes to identify the cell types
represented by each cluster (Fig. S2A-F, Table S1A-C). In order to resolve specific
cardiomyocyte subtypes, we subset and re-clustered cell types of interest, including
cardiac progenitors, cardiomyocytes, and related mesoderm cell types (Fig. 2A-C, Fig.
S2G-I, Table S2A-C). By doing so, we were able to identify clusters representing FHF and
early differentiating cardiomyocytes (CMs/FHF), SHF, and juxtacardiac field[23] (JCF)
progenitors at E7.75 (Fig. 2A). Similarly, at E8.5 we could distinguish inflow tract
cardiomyocytes (IFT-CMs), ventricular cardiomyocytes (V-CMs), and outflow tract
cardiomyocytes (OFT-CMs) (Fig. 2B). At E9 we could more specifically identify the IFT-
CMs as either atrial or atrioventricular canal (A- or AVC-CMs) (Fig. 2C). Having identified
our cell types of interest, we next performed differential gene expression analysis
between MEF2C KO and WT cells within each of these cell types (Fig. 2D-F, Table S3A-
J). Generally speaking, we found larger numbers of genes down-regulated than up-
regulated with loss of MEF2C, agreeing with the notion of MEF2C as an activator of the
myogenic transcriptional program[24]. Among the top differentially expressed genes
(DEGs) within each cell type of interest, we found a number of genes that have been
associated with high confidence to congenital heart defects[25], including *Myh6*, *Myh7*,
Actc1, and *Foxp1* (Fig. S2J). Furthermore, nearly all of these top DEGs were identified
as the nearest gene to peaks identified in a previously published MEF2C ChIP-seq
dataset of embryonic hearts at E14[26], demonstrating that these genes are indeed likely
to be direct targets of MEF2C regulation in the developing heart.

By more closely examining the results of our differential gene expression analysis
(Table S3A-J), two categories of DEGs emerged. First, we observed substantial down-
regulation of key cardiomyocyte genes across all three heart tube segments. This
included cardiac transcription factors such as *Nkx2.5*, *Tbx5*, and *Gata4*, as well as
sarcomeric proteins including *Tnnt2* and *Ttn* (Fig. 2G-I). This result further emphasizes
the importance of MEF2C in early cardiomyocyte fate specification, both by directly
driving expression of contractile proteins and by promoting additional transcription factors
involved in regulating the cardiomyocyte transcriptional program. The second category of

DEGs we identified were genes involved in anterior/posterior patterning of the heart tube.

Specifically, genes such as *Tbx5*, *Gata4*, and *Wnt2* that are normally expressed only in

the posterior inflow tract segment of the heart tube were not only up-regulated in the IFT-

CMs, but their expression also expanded into the V-CMs in MEF2C KO (Fig. 2G-I).

Additionally, expression of the anterior outflow tract-specific gene *Tdgf1* was completely

lost in MEF2C KO (Fig. 2H-I), as previously demonstrated[27]. Finally, we validated these

changes in gene expression with whole-mount embryo fluorescence *in situ* hybridization,

confirming that MEF2C KO leads to broad reduction in the cardiomyocyte transcriptional

program and leads to a notable "posteriorization" of the developing heart tube (Fig. 2J).

MEF2C regulates chromatin accessibility broadly throughout the heart tube and in

a segment-specific manner

In order to better understand how MEF2C regulates the observed segment-specific gene

expression changes, we next used ArchR[28] to integrate both the gene expression and

chromatin accessibility data. We performed dimensionality reduction using ArchR's latent

semantic index (LSI) implementation on the snRNA-seq and snATAC-seq modalities, and

then plotted the cells in UMAP space using the combined dimensions (Fig. S3A-C).

Because individual cells have matching barcodes in the 10x Multiome snRNA-seq and

snATAC-seq libraries, we simply transferred the cell type labels from our Seurat objects

to our ArchR datasets. As we did with the gene expression data, we subset the cardiac

cell types and associated mesoderm in order to better resolve cardiomyocyte subtypes.

Finally, we created pseudobulk groups consisting of WT and MEF2C KO cells for each

given cell type label in these subset datasets and called peaks in the accessibility data

using MACS2[29].

Our initial analyses revealed few changes to chromatin accessibility at E7.75 (Fig.

S3D). This was not entirely unexpected, given that there is no discernable phenotype in

the MEF2C KO embryos at this timepoint, and agrees with the relatively few changes in

gene expression we observed in CMs/FHF, SHF, and JCF cells at E7.75 (Fig. 2D). Thus,

we chose to focus on the E8.5 dataset in order to identify the earliest substantial changes

in chromatin accessibility that lead to the MEF2C KO phenotype (Fig. 3A). Using our

pseudobulked data, we identified differentially accessible regions (DARs) of chromatin

between MEF2C KO and WT cells in each of the subset cell types and found that, as
expected, the most changes occurred in the cardiomyocytes (Fig. 3B). We also found that
in each of the IFT-CMs, V-CMs, and OFT-CMs, there were more lost DARs than gained
DARs in MEF2C KO (Fig. 3B), and that these lost DARs were highly enriched for MEF2C
binding motifs (Fig. S3E). These data further reinforce MEF2C's role as an activator of
the cardiac transcriptional program. Interestingly, when we examined the lost DARs in
MEF2C KO embryos, we found that there were 675 regions that demonstrated lost
accessibility in two or more heart tube segments, and 2,549 regions that lost accessibility
in a segment-specific manner (Fig. 3C). This likely reflects two sets of DARs with different
biological functions: i) regions that are involved in the regulation of general cardiomyocyte
genes, and ii) regions that are involved in segment-specific gene regulation. By contrast,
the gained DARs in MEF2C KO were almost exclusively unique to each heart tube
segment (Fig. 3C). This is interesting because it provides evidence that distinct
transcriptional networks exist for each of the three heart tube segments, and thus, each
network is uniquely affected by loss of MEF2C.

We performed peak-to-gene linkage (P2G) analysis in order to determine which of
the identified differentially accessible regions were most likely to be *bona fide* regulatory
elements. Peak-to-gene linkage identifies statistical correlation between the accessibility
of a given peak and the expression of potential target genes within 250 kilobases of the
peak. This revealed 817 P2G links in the OFT-CMs, 1,201 in the V-CMs, and 2,257 in the
IFT-CMs. To further refine these results, we intersected the P2G links with our differential
gene expression analysis in Seurat to identify which P2G links are connected to genes
that exhibit altered expression in MEF2C KO (Table S4A-C). We then visualized this data
by plotting each P2G link according to the Log2 fold change values from the differential
peak accessibility and differential gene expression tests, and colored each point by the
correlation strength of the P2G link (Fig. 3D).

These DAR-P2G correlation plots reveal a number of interesting observations.
Genes such as *Nkx2.5*, *Myh6*, and *Myh7*, which have decreased expression throughout
the heart tube in MEF2C KO, are also linked to the accessibility of DARs in all three
segments (Fig. 3D). In particular, the *Myh6*/*Myh7* locus contains a number of regions that
have P2G links and clearly altered accessibility in OFT-CMs, V-CMs, and IFT-CMs (Fig.

3E). This suggests that these DARs are likely to be regulatory elements that drive
expression of these genes and are sensitive to the expression of MEF2C in the entire
heart tube. By contrast, there are also P2G links between peaks with lost accessibility
and decreased gene expression that are unique to the individual segments, such as links
to *Wnt11* in the OFT-CMs, *Nppa* in the V-CMs, and *Cacna1c* in the IFT-CMs (Fig. 3D).
Furthermore, examining the links between peaks with gained accessibility and genes with
increased expression in MEF2C KO reveals additional evidence for the increased
"posteriorized" transcriptional program we previously described. Namely, P2G links to
Tbx5 in the V-CMs, as well as *Gata4* and *Wnt2* in the IFT-CMs (Fig. 3D and 3F). These
P2G links represent potential regulatory elements that are activated or over-activated in
response to the loss of MEF2C and, at least in part, drive the increased posteriorization
of the heart tube.
187
Pseudotime analysis with MIRA reveals distinct MEF2C-depedent developmental
trajectories for each heart tube segment
In order to fully take advantage of our multiomic dataset containing precisely staged
embryos at consecutive embryonic timepoints, we used multimodal Models for Integrated
Regulatory Analysis[30] (MIRA) to examine the altered temporal dynamics of gene
expression and chromatin accessibility in the developing heart tubes of WT and MEF2C
KO embryos. MIRA's probabilistic topic modeling borrows concepts that were developed
to identify abstract topics within natural language that give meaning to a collection of text,
and applies them to snRNA-seq and snATAC-seq. In this case, MIRA uses deep learning
and probabilistic graphical models to construct sets of "topics" that represent either genes
or accessibility peaks, and then uses these topics to define cell identities. Finally, MIRA
uses these topics to construct a joint representation that can be used to plot cells in UMAP
space, perform pseudotime analysis, and construct cell state trees (Fig. 4A).
We were particularly interested in using MIRA's ability to construct cell state trees
to understand how the lineage trajectories of OFT-CMs, V-CMs, and IFT-CMs were
altered due to the loss of MEF2C. We trained topic models using cardiac progenitors and
cardiomyocyte subtypes from both genotypes at each of our three timepoints (E7.75:
CMs/FHF, SHF, JCF; E8.5: aSHF, pSHF, OFT-CMs, V-CMs, IFT-CMs; E9: OFT-CMs, V-

CMs, A-CMs, AVC-CMs) and then used those models to construct joint representation
UMAPs of the outflow tract, ventricular, and inflow tract lineages (Fig. 4B, 4F, 4J). We
next calculated pseudotime values for cells within the UMAPs and used the results to
construct cell state trajectories for each of the three lineages (Fig. 4C-D, 4G-H, 4K-L).

By labeling the cells within these lineage trajectories by both cell type and
genotype, we can observe clear differences in the fate specification of MEF2C KO cells,
compared to WT cells, as represented by branch points in the trajectories (Fig. 4D, 4H,
4L). Most interestingly, we notice that the structure of these trajectories, along with the
manner and timing by which MEF2C KO cells diverge from WT cells within the
trajectories, is different for all three lineages. Specifically, within the outflow tract lineage,
MEF2C KO and WT cells proceed similarly up until E8.5, and then completely diverge as
development proceeds to E9 (Fig. 4D). Within the ventricular lineage, there is no branch
point. Instead, MEF2C KO V-CMs from both E8.5 and E9 are located earlier in the
trajectory than even E8.5 WT V-CMs (Fig. 4H). This indicates that the result of knocking
out MEF2C in V-CMs resembles a marked delay or termination of normal development
more so than a divergence to an alternative cell state. The situation is most complex for
the inflow tract lineage, in which there are two points of divergence between MEF2C KO
and WT cells (Fig. 4L). First, there is a subset of MEF2C KO cells that branch away from
the WT trajectory as they differentiate from pSHF progenitors to IFT-CMs at E8.5. A
second set of MEF2C KO cells proceed along the WT trajectory to E9, before ultimately
diverging from WT cells at the "terminal" AVC-CM and A-CM states. Together, these data
reinforce our working hypothesis that MEF2C regulates development of the heart tube in
a segment-specific manner via distinct regulatory networks.

We gained further insights regarding the segment-specific regulation of heart tube
development by examining the top 200 genes and the top ranked TF binding motifs
enriched in the gene expression and chromatin accessibility topics (Table S5). As WT
cells diverge from MEF2C KO cells in the outflow tract lineage, they are marked by gene
expression topics enriched for canonical outflow tract markers *Tdgf1* and *Wnt11*[27,31], and
their accessibility topics are marked by peaks enriched for MEF2 and FOS binding motifs
(Fig. 4E). Interestingly, the MEF2C KO cells are marked by expression of Notch and Wnt
signaling components such as *Dlk1* and *Fzd1*, and transcription factors *Meis1* and *Hand2*

(Fig. 4E). The accessibility topic enriched for ISL, NKX, and TEAD is maintained to a
larger extent in the outflow tract of MEF2C KO cells (Fig. 4E), suggesting these cells may
retain a more progenitor-like state in the absence of MEF2C. The gene expression and
accessibility topics enriched along the ventricular lineage trajectory supports our
interpretation that the MEF2C KO cells experience a substantial delay or termination of
their normal differentiation. Notably, topics that are enriched for CM genes (e.g. *Myl2,*
Tnnt1) and cardiac TF binding motifs (e.g. TBX, MEF2, NKX) only mark the end of the
differentiation trajectory containing WT cells (Fig. 4I). In the complex branching inflow
tract trajectory, we noted that the terminal WT cells are enriched for topics containing
expected atrial genes, such as *Angpt1* and *Nr2f1*, and binding motifs for TBX5 and SNAI,
among others (Fig. 4M, Fig. S4A-B). Whereas, the MEF2C KO inflow tract cells stream
into one of two terminal branches enriched for genes such as *Wnt6, Jarid2, Tbx2,* and
Zeb1 and binding motifs for ERF, GATA, and TEAD factors (Fig. 4M, Fig. S4A-B).
Together, these data illustrate that lineage-specific changes in gene expression and
chromatin accessibility occur as heart tube development progresses and that these
lineage-specific programs are uniquely altered upon the loss of MEF2C.
253
Leveraging multiomic data to identify novel enhancers via zebrafish transgenesis
We next sought to determine whether the observed lineage-specific patterns of gene
regulation by MEF2C were driven by segment-specific enhancers that are active during
heart tube development. To identify candidate enhancers, we utilized our E8.5 ArchR
dataset containing integrated chromatin accessibility and gene expression data. We
began by identifying all peaks that lost accessibility in the MEF2C KO (3,224), then filtered
to the peaks that demonstrated altered accessibility in only one of the three heart tube
segments (2,549) and contained a MEF2C binding motif (1,059). Next, we again utilized
ArchR's Peak2Gene functionality and filtered down to peaks correlated with the
expression of genes that were mis-regulated in the MEF2C KOs (130) and were located
in promoter regions (119).   Finally, from this list of 119 candidates, we selected 12 of the
highest priority to screen based on the Log2FC value of the change in peak accessibility
and linkage to genes of interest (Fig. 5A, Table S6). We enumerated these candidate
regions with the prefix "MVEB," after the initials of authors Muncie-Vasic and Brower. To

further validate our selection of candidates, we overlayed our chromatin accessibility data

at these regions with published ChIP-seq datasets[26,32] to visualize the occupancy of

MEF2C and H3K27ac, a histone post-translational modification that marks active

enhancers. Importantly, each of our candidates had clear and strong occupancy of both

MEF2C and H3K27ac (Fig. 5B). Of note, one of our ventricular-specific candidates,

MVEB8, is located within the *Nppa/Nppb* super enhancer cluster[33]. Furthermore, two of

the candidates we identified form differential accessibility in IFT-CMs overlapped with

regions found in the VISTA enhancer database[34], with demonstrated enhancer activity

in the hearts of E11.5 mouse embryos (Fig. 5C-D).

In order to determine whether our candidate regions are *bona fide* enhancers with

activity in early heart tube development, we employed the zebrafish Tol2 Transgenesis

system[35,36], allowing us to screen candidates with relatively high-throughput. We

reasoned that because *Mef2c* has the orthologs *mef2ca* and *mef2cb* in zebrafish, and

because there is a high degree of overlap between the key cardiac transcriptional

regulators in zebrafish and mouse[37], there should be a fairly high concordance between

the activity of our candidate regions identified in the mouse and their activity in the fish.

Moreover, it has been previously demonstrated that regulatory elements can exhibit

activity in the heart of both species[38–40]. To perform our screen, we cloned our candidate

enhancer regions into vector backbones just upstream of a minimal E1b promoter and

the coding sequence for eGFP, flanked by Tol2 recognition sites. As a negative control,

we also tested an empty reporter construct consisting of only the minimal E1b promoter

and eGFP. To control for injection efficacy, we utilized a vector consisting of the ß-actin

promoter upstream of the minimal E1b promoter and BFP, flanked by Tol2 sites. These

vectors were injected along with transposase mRNA into zebrafish embryos from

Tg(*cmlc2:mCherry*)[s890] and WT parents, allowing us to screen F0 embryos for enhancer

activity (eGFP) within the heart (mCherry) (Fig. 5E). To determine whether a candidate

enhancer demonstrated activity, we quantified the percentage of BFP-positive embryos

that exhibited eGFP expression in the heart. We were unable to successfully clone one

of our twelve candidates (MVEB10) due to a highly repetitive "GT" sequence, so our final

screen consisted of eleven candidate enhancers.

Upon screening, we detected enhancer activity in the developing heart for seven
of the eleven candidates (Fig. 5F, Fig. S5A-B). Both candidates that overlapped with
regions found in the VISTA enhancer database (MVEB1, MVEB2) exhibited activity in the
heart by 72 hours post-fertilization (hpf) (Fig. 5F). Two additional candidates we identified
from differential accessibility in the IFT-CMs, MVEB3 and MVEB6, demonstrated robust
and specific activity in the zebrafish heart at 72 hpf (Fig. 5F, Fig. S5A-B). Finally, we also
observed activity from two candidates that we identified from differential accessibility in
the V-CMs, MVEB7 and MVEB8, and one candidate identified from differential
accessibility in the OFT-CMs, MVEB12 (Fig. 5F, Fig. S5A-B). To our knowledge, these
five candidates (MVEB3, MVEB6, MVEB7, MVEB8, MVEB12) have not been previously
reported and represent novel, MEF2C-depenedent enhancers with activity in the
developing heart. Interestingly, we also observed a range in the timing of enhancer activity
in the embryonic zebrafish heart. Candidates MVEB2 and MVEB6 were clearly detectable
as early as 24 hpf, while the others were not active until 72 hpf (Fig. 5G-H). This variation
in the onset of enhancer activity could result from differences in the timing of accessibility
of these regulatory elements, different levels of sensitivity to *Mef2c* expression, or
cooperation with other TFs or transcriptional regulators.

An overactive "posteriorized" gene regulatory network driven by NR2F2 and
GATA4 underlies the expanded inflow tract phenotype in MEF2C KO
The posteriorized gene signature (Fig. 2H-J) and chromatin accessibility landscape (Fig.
3F, 3H) that occurs in the heart tube of MEF2C KO embryos, together with the multi-
staged branching structure of the inflow tract lineage trajectory (Fig. 4L), reveals a
previously unappreciated complexity of the transcriptional network governing inflow tract
development that we sought to further dissect. Moreover, the expanded but aberrantly
formed inflow tract that forms in MEF2C KO embryos (Fig. 1C) suggests that loss of
MEF2C leads to a more complex heart tube phenotype than a simple failure of cardiac
differentiation and growth. To determine what transcriptional regulators may drive the
aberrant phenotype in the posterior of the heart tube, we performed TF binding motif
enrichment analysis on the regions of chromatin that gain accessibility in the MEF2C KO
IFT-CMs at E8.5 and found that Nuclear Receptor (NR) and GATA motifs are among the

most enriched (Fig. 6A). From this observation, we hypothesized that the loss of MEF2C in the posterior heart tube causes a redistribution of NR and GATA factors to regions of chromatin where they would normally not have activity. More specifically, we hypothesize that MEF2C co-binds certain regulatory elements along with NR or GATA factors in the WT condition, and that these NR and GATA factors become "free" to bind different regulatory elements in the MEF2C KO. If this were true, one would expect that sites of accessible chromatin that mark WT IFT-CMs and contain an NR or GATA motif would be likely to also contain a MEF2C motif, whereas the sites that gain accessibility in MEF2C KO and contain an NR or GATA motif would be more likely to not also contain a MEF2C motif. Indeed, this is exactly what we observed (Fig. 6B).

These data are particularly interesting given the importance of GATA factors, and in particular GATA4, in cardiac development[41] and the knowledge that nuclear receptor subfamily 2 group F member 2 (NR2F2, also known as COUP-TFII) is a necessary driver of atrial development[42,43]. Using the web tool Galaxy[44] and publicly available ChIP-seq datasets[42,45], we found NR2F2 and GATA4 occupancy at a number of the regions of chromatin that gained accessibility in the MEF2C KO IFT-CMs at E8.5 (Fig. S6A-B), suggesting that these DARs could indeed be sites of potential NR2F2 and GATA4 regulation. To determine whether NR2F2 and GATA4 may be redistributed and driving the "posteriorization" of the heart tube (Fig. 2H-I), we next asked whether NR2F2 and GATA4 target genes were upregulated in the inflow tract of MEF2C KO embryos. Using publicly available datasets to identify the target genes of each TF[46,47], we calculated the odds ratio (OR) of finding NR2F2 or GATA4 target genes compared to non-target genes amongst our upregulated differentially expressed genes at E8.5. We found strong evidence that NR2F2 (OR = 1.52, p = 1.79e-4) and GATA4 (OR = 4.23, p < 2.2e-16) targets were enriched amongst the upregulated genes (Fig. 6C). Next, we took the list of NR2F2 and GATA4 target genes that were upregulated (Table S7A-B) and calculated a "module score" for the expression of these genes in our dataset[48]. In agreement with our working model that these NR2F2 and GATA4 targets are driving the "posteriorized" heart tube phenotype in MEF2C KO, we found the NR2F2 and GATA4 module scores were increased in all segments of the heart tube at E8.5 (Fig. 6D).

To further examine the altered roles of NR2F2 and GATA4 in the transcriptional

networks governing heart tube development, we used CellOracle[49] to construct gene

regulatory networks for each segment of the heart tube at each timepoint in our dataset

(Fig. 6E). In brief, CellOracle uses integrated gene expression and chromatin accessibility

data to infer interactions between TFs and target genes based on the accessibility of a

given TF's binding motif near a potential target gene and the correlation of expression

between the TF and the potential target. To validate that these inferred CellOracle

networks effectively captured the true gene regulatory networks that govern inflow tract

development, we deleted *Mef2c in silico* in the E8.5 inflow tract network and compared

the predicted changes in gene expression to the measured changes in gene expression

in our E9 dataset (Fig. 6F). We found that the networks correctly predicted the gene

expression changes with 92% accuracy (Fig. 6G). Having validated the accuracy of the

CellOracle networks, we next visualized the core cardiac subnetworks of altered genes

in the WT and MEF2C KO inflow tracts (Fig. 6H). These subnetworks consist of key

cardiac transcription factors and their inferred interactions with the top 100 most

differentially regulated genes in the E8.5 MEF2C KO IFT-CMs. Next, we further narrowed

in on the direct interactions between NR2F2 and GATA4 and their target genes in the WT

and MEF2C KO inflow tract networks (Fig. 6I-J). From these plots we observe similar

behavior for both TFs – approximately half of the WT interactions are maintained in the

MEF2C KO, while nearly twice as many new direct interactions are established. This is

consistent with our working model that in response to MEF2C KO, aberrant NR2F2 and

GATA4 activity regulate additional genes that drive the inflow tract phenotype in the

posterior of the heart tube.

Given this hypothesis, we theorized that reducing the activity of NR2F2 may rescue

the inflow tract phenotype in the MEF2C KO heart tube. To test this, we generated

Mef2c^del/+;*Nr2f2*^del/+ male mice and crossed them to *Mef2c*^del/+ females. Although there is

no apparent heart tube phenotype at E9.5 for *Nr2f2*^del/+ (Fig. 6K), it was previously shown

that these animals are recovered at lower numbers than expected Mendelian ratios[43],

suggesting that *Nr2f2* is a dosage-sensitive gene and that *Nr2f2* heterozygosity can be

expected to alter transcriptional activity. Strikingly, we observed a notable partial rescue

in ventricle and inflow tract development at E9.5 (18-24 somites) in the

Mef2c^{del/del};*Nr2f2*^{del/+} embryos compared to *Mef2c*^{del/del} (Fig. 6K-L), although the extent of

the rescue did vary (Fig. S6C-D). The MEF2C KO single-ventricle phenotype was

maintained in *Mef2c*^{del/del};*Nr2f2*^{del/+} embryos, but these ventricles were larger and more

expanded than Mef2c^{del/del}. Moreover, the posterior structures of the heart tube in

Mef2c^{del/del};*Nr2f2*^{del/+} more closely resembled WT inflow tracts with expanded prospective

atria, compared to the completely disrupted inflow tract in *Mef2c*^{del/del} embryos (Fig. 6K-

L). This partial rescue of the ventricle and inflow tract in *Mef2c*^{del/del};*Nr2f2*^{del/+} embryos

provides clear evidence that the MEF2C KO phenotype in the posterior heart tube is

driven, at least in part, by an altered gene regulatory network with increased and aberrant

NR2F2 activity.

DISCUSSION

To date, MEF2C has been thought of as a broad activator of the myocyte transcriptional

program, important for driving the expression of sarcomeric proteins and related

contractile machinery[24,50]. Here, we demonstrate that in addition to this role, MEF2C is

involved in more complex, segment-specific networks that regulate gene expression and

chromatin accessibility in the developing linear heart tube. Overall, we found that loss of

MEF2C causes a "posteriorization" of the heart tube. Typical outflow tract markers were

downregulated in MEF2C KO mice, while inflow tract markers were upregulated and their

domain of expression expanded into the developing ventricle (Fig. 2H-J, Table S3). These

changes in gene expression were coupled to altered chromatin accessibility throughout

the developing heart tube (Fig. 3). In line with MEF2C functioning as both a myocyte

activator and a segment-specific regulator, we found sets of DARs that were linked to

altered gene expression of sarcomeric proteins throughout the entire heart tube, as well

as DARs that were linked to DEGs unique to specific heart tube segments (Fig. 3C-F).

Using a deep-learning based modeling approach[30] to construct pseudotime lineage

trajectories for WT and MEF2C KO CMs, we also found a previously unappreciated

complexity to the role of MEF2C in regulating development of the inflow tract (Fig. 4J-M).

Our analyses demonstrated that MEF2C is required for proper determination of inflow

tract fate at E8.5, as well as for proper differentiation to either atrial or atrioventricular

canal CM subtypes at E9. This was markedly different from what we observed in the

outflow tract and ventricular lineages, where loss of MEF2C resulted in either a single
branch point diverging from normal development (outflow tract) or a delay in differentiation
(ventricular).

Using our integrated snRNA-seq and snATAC-seq datasets, we identified seven
MEF2C-dependent enhancers with activity in the early zebrafish embryonic heart with a
hit-rate greater than 60% (Fig. 5). The accuracy of our screening approach resulted from
our ability to link accessibility directly to gene expression in a segment-specific manner
and prioritize our candidates using published MEF2C and H3K27ac ChIP-seq
datasets[26,32]. Interestingly, we observed a range in the timing of onset of enhancer
activity among the positive hits we identified. This variation is likely due to differences in
the underlying mechanisms of activity specific to each enhancer, which warrant further
study. Possible mechanisms include differences in the temporal dynamics of accessibility
of each enhancer region, differential sensitivity to *Mef2c* expression, and unique
cooperativity with other TFs and regulatory elements. Due to the mosaicism of zebrafish
transgenesis in F0 embryos, we were unable to determine whether our novel enhancers
demonstrated segment-specific activity in the zebrafish. We are currently in the process
of generating stable lines that will be better suited for examining the precise spatial
domains of enhancer activity. In the future, we also plan to perform transgenesis of our
novel enhancers in the mouse to more accurately determine their activity in the
mammalian heart.

By constructing lineage-specific gene regulatory networks, we identified a genetic
interaction between MEF2C and NR2F2 in the inflow tract of the developing linear heart
tube. Although *Nr2f2* is not differentially expressed in any CM subtype at E8.5 and is
modestly downregulated in A-CMs at E9 (Table S3), we identified an altered
transcriptional network occurs in MEF2C KO (Fig. 6H) that leads to increased gene
regulation by NR2F2 (Fig. 6I) at newly accessible regions of chromatin (Fig. 6A).
Interestingly, MEF2C and NR2F2 have been shown to physically interact to regulate gene
expression in Leydig cells in the male testes[51]. Our data suggest that MEF2C and NR2F2
also cooperate as co-factors in the linear heart tube. We envision a model in which the
loss of MEF2C leads to a redistribution of NR2F2 to other sites in the genome, altering
its transcriptional activity. In support of this model, we observed that newly accessible

regions of chromatin in MEF2C KO IFT-CMs containing an NR binding motif rarely also

contained a MEF2C motif (Fig. 6B). This suggests that in the WT condition, MEF2C may

be responsible for recruiting NR2F2 to the sites that are regulated by cooperativity

between MEF2C and NR2F2. Both MEF2C and NR2F2 contain well-characterized DNA

binding regions and regulate gene expression by binding DNA directly[52,53], so it remains

unclear why and how NR2F2 redistributes to new sites upon the loss of MEF2C. One

likely possibility is that interactions with new co-factors are involved in this process. It will

also be interesting to determine whether the normal interaction of MEF2C and NR2F2 in

the heart tube occurs directly or involves other key cardiac TFs or co-factors.

Our approach provides a generalizable framework for integrating snRNA-seq and

snATAC-seq data to build GRNs that provide specific and comprehensive models of gene

regulation. By constructing these GRNs using both WT and mutant data, we are able to

immediately test predictions made from the WT networks (Fig. 6F-G) and validate the

inferred regulatory interactions. These tools also allow specific networks to be built for

particular cell or tissue types, provided there are enough cells in the group of interest to

generate robust inferences. Moreover, the experimental mutant data reveals how the

overall networks are altered – both in ways that would and would not be predicted from

the WT networks. These unexpected differences help to illuminate indirect or

transactivating gene regulation. Insights from these networks lead to new hypotheses

about regulatory elements, TF interactions, DNA binding, and chromatin organization. As

these hypotheses are tested experimentally, the GRNs can be refined, leading to higher-

fidelity models that will have greater predictive power. These higher-fidelity networks will

yield important new insights into the transcriptional regulation of developmental

processes and will have the ability to reveal precisely how these processes are altered in

the context of disease-causing mutations.

MATERIALS AND METHODS

Mouse Models

Animal studies were performed in strict compliance with the UCSF Institutional Animal Care and Use Committee. Mice were housed in a standard 12-hour light/dark animal husbandry barrier facility at the Gladstone Institutes. The *Mef2c*[del/+] allele used was the same as the original generated by Lin et al.[19]. The *Smarcd3*-F6-eGFP reporter allele was generated in our lab and described previously[20]. The *COUP-TFII lacZ* knock-in allele (abbreviated here as Nr2f2[del/+]) was obtained from the Mutant Mouse Resource & Research Centers and was originally generated by Takamoto et al.[54]. The *Mef2c*[del/+] and *Nr2f2*[del/+] lines were maintained in the C57BL6/J background (Jackson Laboratory #664). The *Smarcd3*-F6-eGFP allele was crossed into our *Mef2c*[del/+] line from a mixed background of C57BL6/J (Jackson Laboratory #664) and CD-1 (Charles River #022).

Timed Embryo Dissections and Genotyping

To achieve timed matings, male and female mice were housed together in the evening and pregnancy was assessed by vaginal plug the following morning. Gestational stage was determined relative to noon on the day of plug detection, defined as day E0.5. Embryos were dissected and, at later stages when yolk was present, also de-yolked, in ice-cold PBS (Life Technologies, 14190250) with 1% fetal bovine serum (FBS; Thermo Fisher Scientific, 10439016) on ice. The posterior trunk was removed by microdissection for all embryos at E8.5 or later that were collected for imaging (IF or FISH via HCR) to ensure unobstructed views of the heart tube. DNA for genotyping was extracted using QuickExtract DNA Extraction Solution (Lucigen, QE09050) from harvested yolk sac tissue, if available, or else from a micro-dissected piece of the extra-embryonic anterior proximal region. For embryos collected for 10x Multiome library construction, rapid genotyping was performed while embryos remained on ice using Phire Green Hot Start II PCR Master Mix (Thermo Fisher Scientific, F126S), according to manufacturer's protocols. For all other experiments, genotyping was performed after embryo fixation using GoTaq Green Master Mix (VWR, PAM7122), according to manufacturer's protocols. PCR products were run on a 1% agarose gel (Thermo Fisher Scientific, BP164-500) with Biotium GelRed Nucleic Acid Gel Stain (Thermo Fisher Scientific, NC9594719) used for

DNA detection. Primers used for genotyping were ordered from Integrated DNA

Technologies (IDT). For genotyping the *Mef2c* allele, a three-primer mix was used (P1:

ACTGGCTGAGAGTTGTACCCAC, P2: GCGTGATTTGCTAGATAGTGGTAGAC, P3:

ATGTGGAATGTGTGCGAGGC), resulting in a single 314 bp band for WT, a single 242

bp band for $Mef2c^{del/del}$, and both bands for $Mef2c^{del/+}$. For genotyping the *Nr2f2* allele, a

three-primer mix was used (P4: CATCCGGGATATGTTACTGTCCGG, P5:

TGGGGAAGCTAAGTGTTGATGTGATTCC, P6: GCCGTGGGTTTCAATATTGGCTTC),

resulting in a single 786 bp band for WT, a single ~1600 bp band for $Nr2f2^{del/del}$, and both

bands for $Nr2f2^{del/+}$.

Whole-Mount Embryo Immunostaining and Imaging

Embryos collected for immunostaining were fixed for 1 hour at room temperature in 4%

paraformaldehyde (PFA) freshly diluted from 16% weight/volume PFA aqueous solution

(Thermo Fisher Scientific, 043368-9M) in PBS (Life Technologies, 14190250). Embryos

were stored at 4 °C for up to 1 month prior to immunostaining. To permeabilize and block

non-specific antigens, embryos were incubated at 37 °C for 2 hours with gentle rotation

in a blocking buffer consisting of PBS (Life Technologies, 14190250), 5% normal donkey

serum (Sigma Aldrich, S30-M), and 0.8% Triton X-100 (Sigma Aldrich, X-100-5ML). This

was then removed and replaced with primary antibodies diluted in the same blocking

buffer and incubated overnight at 37 °C with gentle rotation. The next day, primary

antibody solution was removed and embryos were washed 3x 30 minutes in the same

blocking buffer at 37 °C with gentle rotation. Secondary antibodies and DAPI (Abcam,

ab228549, used at 1 µg/mL) were added in a blocking buffer consisting of PBS (Life

Technologies, 14190250), 5% normal donkey serum (Sigma Aldrich, S30-M), and 0.4%

Triton X-100 (Sigma Aldrich, X-100-5ML), and incubated for 3-4 hours at 37 °C with gentle

rotation. Secondary antibody solution was removed and embryos were washed 3x 30

minutes in PBS (Life Technologies, 14190250) with 0.1% Tween-20 (Thermo Fisher

Scientific, J20605.AP) at 37 °C with gentle rotation. Embryos were imaged in PBS using

an upright epifluorescence microscope (Leica MZFLIII, Leica DFC 3000G, Lumen

Dynamics XCite 120LED) and acquisition software LASX (Leica). Primary antibodies

used were sheep polyclonal MEF2C (R&D Systems, AF6786, used at 1:250) and rabbit

polyclonal cardiac Troponin T (Thermo Fisher Scientific, 15513-1-AP, used at 1:250).

Secondary antibodies used were Donkey anti-Sheep IgG (H+L) Cross-Adsorbed

Secondary Antibody, Alexa Fluor™ 647 (Thermo Fisher Scientific, A-21448, used at

1:1000) and Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody,

Alexa Fluor™ 555 (Thermo Fisher Scientific, A-31572, used at 1:1000).

Whole-Mount Embryo Fluorescence *In Situ* Hybridization via Hybridization Chain Reaction and Imaging

The protocol for *in situ* hybridization via hybridization chain reaction (HCR™) was adapted

from the optimized manufacturer's protocol for whole-mount mouse embryos (Molecular

Instruments). Embryos were fixed overnight at 4 °C in 4% PFA freshly diluted from 16%

weight/volume PFA aqueous solution (Thermo Fisher Scientific, 043368-9M) in PBS (Life

Technologies, 14190250). Embryos were then washed 3x 5 minutes in PBS (Life

Technologies, 14190250) with 0.1% Tween-20 (Thermo Fisher Scientific, J20605.AP),

abbreviated PBST, and then processed through 10-minute serial incubations in a

dehydration series of 25%, 50%, 75%, and 100% methanol solutions in PBST on ice.

Embryos were stored in 100% methanol at −20 °C for up to 6 months prior to *in situ*

hybridization. Embryos were then rehydrated through 10-minute serial incubations in a

series of 75%, 50%, 25%, and 0% methanol solutions in PBST on ice. Embryos were

then brought to room temperature and permeabilized by 10-minute incubation with 10

μg/mL Proteinase K (Sigma Aldrich, 70663-5) in PBST, washed 2x 5 minutes in PBST at

room temperature, post-fixed in 4% PFA freshly diluted from 16% weight/volume PFA

aqueous solution (Thermo Fisher Scientific, 043368-9M) in PBS (Life Technologies,

14190250), and then washed 3x 5 minutes in PBST at room temperature. Embryos were

pre-hybridized in Probe Hybridization Buffer (Molecular Instruments) for 5 minutes at

room-temperature followed by an additional 30 minutes at 37 °C with fresh buffer. Single

or multiplexed hybridization probes were then diluted in pre-warmed Probe Hybridization

Buffer (Molecular Instruments) and embryos were hybridized for 48 hours at 37 °C.

Embryos were then washed 4x 15 minutes with pre-warmed Probe Wash Buffer

(Molecular Instruments) at 37 °C, followed by 2x 5 minutes with 5xSSCT (diluted in

nuclease-free water from 20xSSCT, Thermo Fisher Scientific, 15557044) at room

temperature. During washes, amplification hairpins were snap-cooled by heating to 95 °C
for 90 seconds and then allowed to cool to room temperature in the dark. Embryos were
pre-amplified in Amplification Buffer (Molecular Instruments) for 5 minutes at room
temperature. Snap-cooled hairpins and DAPI (Abcam, ab228549, used at 1 µg/mL) were
diluted in Amplification Buffer (Molecular Instruments) and embryos were incubated in this
solution overnight in the dark at room temperature. The following day, embryos were
washed through a series of 5xSSCT washes (diluted in nuclease-free water from
20xSSCT, Thermo Fisher Scientific, 15557044) in the dark at room temperature (2x 5
minutes, 2x 30 minutes, 1x 5 minutes). Embryos were imaged in PBS using an upright
epifluorescence microscope (Leica MZFLIII, Leica DFC 3000G, Lumen Dynamics XCite
120LED) and acquisition software LASX (Leica). To maximize the utility of collected
embryos, probes were then stripped by overnight incubation in 80% formamide in water
(Thermo Fisher Scientific, BP228100) at room temperature, followed by 3x 5 minutes
washes in PBST at room temperature. Embryos could then be probed for additional
targets, beginning the protocol again at the pre-hybridization step. Probes used in this
study were ordered from the Molecular Instruments catalogue, when possible, or custom-
designed and manufactured by Molecular Instruments. Hybridization probes used were:
Tnnt2-B3 (used at 4 nM), *Ttn*-B2 (used at 4 nM), *Nppa*-B2 (used at 20 nM), *Nkx2.5*-B1
(used at 20 nM), *Tbx5*-B1 (used at 20 nM), *Gata4*-B1 (used at 20 nM), *Wnt2*-B2 (used at
20 nM), *Tdgf1*-B2 (used at 20 nM). Amplification probes used were all used at 60 nM: B1-
Alexa Fluor™ 647, B2-Alexa Fluor™ 546, B2-Alexa Fluor™ 647, B3-Alexa Fluor™ 488.
591

Embryo Preparation for 10x Multiome
For E7.75 embryos, whole embryos were dissected and harvested for single-nucleus
library generation. For E8.5 and E9 embryos, the head folds and posterior trunk were
removed by microdissection prior to harvesting for library generation to enrich the relative
capture of cardiac cell types. Two embryos were collected per genotype per embryonic
stage. Following dissection and rapid PCR genotyping, embryo samples selected for the
multiome experiment were incubated in 200 µl TrypLE Select (Thermo Fisher Scientific,
12563-011) for 5 minutes at 37 °C, triturated gently by pipetting up and down, and then
incubated an additional 3 minutes at 37 °C. The dissociated cell suspension was

quenched with 600 µl of PBS (Life Technologies, 14190250) with 1% FBS (Thermo Fisher

Scientific, 10439016), singularized by passage through the 35 µm mesh of a 5 mL

Falcon™ Round-Bottom Polystyrene Test Tube with Cell Strainer Snap Cap (Thermo

Fisher Scientific, 08-771-23), pelleted by centrifugation at 300 g for 5 minutes at 4 °C,

and resuspended in 50 µl of PBS (Life Technologies, 14190250) with 0.04% bovine serum

albumin (BSA, Sigma Aldrich, A1595). At this stage, the manufacturer's protocol for Nuclei

Isolation for Single Cell Multiome ATAC + Gene Expression Sequencing, Appendix, Low

Cell Input Nuclei Isolation (10x Genomics, CG000365 Rev B) was followed exactly to

prepare nuclei for library generation.

Generation of snRNA-seq and snATAC-seq Libraries and Sequencing

Libraries for snRNA-seq and snATAC-seq were prepared according to the manufacturer's

protocol for Chromium Next GEM Single Cell Multiome ATAC + Gene Expression (10x

Genomics, CG000338 Rev B), using the 10x Genomics Chromium controller, Chromium

Next GEM Single Cell Multiome ATAC + Gene Expression Reagent Bundle (10x

Genomics, PN-1000283), Chromium Next GEM Chip J Single Cell Kit (10x Genomics,

PN-1000234), Single Index Kit N Set A (10x Genomics, PN-1000212), and Dual Index Kit

TT Set A (10x Genomics, PN-1000215). Nuclei isolated from multiome samples were

processed directly into the snATAC-seq transposition reaction and then captured in 10x

Genomics GEMs via the Chromium Controller. GEMs were stored at -80 °C for up to 4

weeks, allowing for collection of replicate embryos of each genotype at each development

stage. All subsequent library preparation steps were performed together for all embryos

of a given developmental stage to reduce the likelihood of technical batch artifacts. A

targeted maximum recovery of 10,000 cells per sample were loaded onto the 10x

Genomics Chromium controller instrument and each sample was indexed with a unique

sample identifier (Single Index Kit N Set A for snATAC-seq libraries, Dual Index Kit TT Set

A for snRNA-seq libraries). Final libraries were quality-controlled using an Agilent

Bioanlyzer instrument with Agilent High Sensitivity DNA Kit (Agilent, 5067-4626). The

DNA concentration of each library was measured using KAPA Library Quantification

qPCR Kit (Thermo Fisher Scientific, 50-196-5234), and then libraries were pooled and

sequenced on NovaSeq6000 S4 lanes (Illumina). All snRNA-seq libraries were

sequenced to a depth of at least 20,000 mean raw aligned reads per cell and snATAC-
seq libraries were sequenced to a depth of at least 35,000 mean raw aligned read pairs
per cell (most libraries >60,000).

Processing of Raw Sequencing Data
Raw sequencing reads were processed using the 10x Genomics Cell Ranger Arc v2.0.0
pipeline. Reads were demultiplexed using cellranger-arc mkfastq and aligned with
cellranger-arc count to the mm10 reference genome containing an additional sequence
for eGFP to map reads of the *Smarcd3*-F6-eGFP reporter transcript.

Seurat Analysis of snRNA-seq Data
Gene expression data were analyzed using Seurat v4.3.0 in R[21]. Datasets for each
developmental stage (E7.75, E8.5, E9) were analyzed as separate Seurat objects.
Counts matrices from each sample that were output by the Cell Ranger Arc pipeline were
used as inputs to the Read10X() and CreateSeuratObject() functions. Data were filtered
for quality control based on number of detected genes, unique molecular identifier (UMI)
counts, percent ribosomal transcripts, and percent mitochondrial transcripts.
SCTransform v2[22] was used to normalize and scale data, with regressions performed
with respect to percentage of ribosomal transcripts and percentage of mitochondrial
transcripts. Principal component analysis (PCA) was performed, followed by batch
correction with Harmony[55], and cells were clustered and visualized as UMAP
embeddings based on the top 30 principal components using FindNeighbors(),
FindClusters(), RunUMAP(), and DimPlot(). Cell types were annotated using the
FindAllMarkers() function with Wilcoxon rank-sum test (min.pct = 0.25, logfc threshold =
0.25, p_val_adj < 0.05) to identify cluster-specific marker genes (Table S1A-C, Fig. S2D-
F). Cardiac progenitors, cardiomyocytes, and related mesoderm cell types were subset
into new Seurat objects. PCA analysis, batch correction, visualization, and clustering were
then performed on these new objects as described above. Cell types in the subset objects
were annotated using the FindAllMarkers() function with Wilcoxon rank-sum test (min.pct
= 0.25, logfc threshold = 0.5, p_val_adj < 0.05) to identify cluster-specific marker genes
that enabled more specific cell type labeling than in the full datasets (Table S2A-C, Fig.

S2G-I). Differential gene expression was performed between MEF2C KO and WT cell

types of interest using the FindMarkers() function with Wilcoxon rank-sum test (min.pct =

0.1, logfc threshold = 0.25, p_val_adj < 0.05). The DotPlot() and RidgePlot() functions

was used to visualize gene expression levels of differentially expressed genes across cell

types of interest. The AddModuleScore() function[48] was used to add module scores for

up-regulated (MEF2C KO-vs-WT) NR2F2 and GATA4 target genes and the RidgePlot()

function was used to visualize these module scores in cell types of interest.

ArchR Analysis of Integrated snRNA-seq and snATAC-seq Data

Initial Dataset Processing

Chromatin accessibility and integrated multiome data was analyzed using the ArchR[28]

software package v1.0.2 in R. The fragments files output by the Cell Ranger Arc pipeline

were used as inupts to the createArrowFiles() function to create Arrow Files for each

sample. These Arrow Files were used to create individual ArchR projects for

developmental stage (E7.75, E8.5, E9). The gene expression data was added using the

import10xFeatureMatrix() and addGeneExpressionMatrix() functions, using the matching

cell barcodes to combine the snRNA-seq and snATAC-seq data for each cell. Data were

filtered for quality control based on the number of unique fragments, TSS Enrichment

score, expression UMI counts, and number of detected genes. Doublets were filtered

using the addDoubletScores() and filterDoublets() functions. Dimensionality reduction

was performed using ArchR's iterative latent semantic index (LSI) implementation

individually for the snATAC-seq data (varFeatures = 25,000, dimsToUse = 1:30) and

snRNA-seq data (varFeatures = 5,000, dimsToUse = 1:30). Then, the reduced

dimensions were combined using addCombinedDims() and batch correction was

performed on the combined dimensions using Harmony[55]. Cells were then embedded in

UMAP space with addUMAP(), clustered using addClusters(), and visualized using

plotEmbedding(). Cell types were annotated by transferring metadata labels from the

previously analyzed Seurat objects. The getMarkerFeatures() function with Wilcoxon

rank-sum test (FDR <= 0.01, Log2FC >= 0.5) was used to identify cluster-specific marker

genes to confirm cell type labels. Cardiac progenitors, cardiomyocytes, and related

mesoderm cell types were subset into new ArchR projects. Dimensionality reduction,

batch correction, clustering, and visualization were performed as described above. Cell

types in the subset projects were annotated by transferring metadata labels from the

previously analyzed subset Seurat objects. The getMarkerFeatures() function with

Wilcoxon rank-sum test (FDR <= 0.01, Log2FC >= 0.5) was used to identify cluster-

specific marker genes to confirm subset cell type labels. Due to the small number of

MEF2C KO OFT-CMs in the E8.5 dataset, an ArchR project with combined E8.5 and E9

samples was created and processed as described above for downstream

analyses/comparisons of OFT-CMs.

Peak Calling, Differential Peak Testing, and Motif Enrichment

Cells were pseudobulked into replicates based on their cell type and genotype labels (e.g.

WT V-CMs), with two pseudobulked replicates created corresponding to the two embryos

collected for each genotype at each developmental stage. Replicates were created using

the addGroupCoverages() function (minCells = 20, maxCells = 500, sampleRatio = 0.8).

Peaks were then called using addReproduciblePeakSet() with MACS2[29] (reproducibility

= 2, peaksPerCell = 500, minCells = 20). Differential peak accessibility testing was

performed using getMarkerFeatures() with Wilcoxon rank-sum test (FDR <= 0.15,

Log2FC >= 0.5 or Log2FC <= -0.5) for each cell type of interest with MEF2C KO cells

used as the group of interest and WT cells used as the background. Transcription factor

binding motifs were annotated using addMotifAnnotations() with the Vierstra motif

archetypes[56] and motif enrichments for regions of gained or lost accessibility were

calculated using peakAnnoEnrichment() with cutoffs of FDR <= 0.15 and Log2FC <= -0.5

or Log2FC >= 0.5.

Peak2Gene Linkages and Browser Track Visualization

Peak2Gene Linkages were identified using the addPeak2GeneLinks() and

getPeak2GeneLinks() functions. Code written in this study was used to filter Peak2Gene

links based on presence of a MEF2C motif and linkage to specific genes (i.e. differentially

expressed genes). Peak2Gene links were visualized using the plotBrowserTrack()

function. The getGroupBW() function was used to export the accessibility coverage data

in bigwig format for visualization along with published ChIP-seq data using the IGV

software package[57].

MIRA Analysis of Integrated snRNA-seq and snATAC-seq Data

Preparing Merged Datasets

We used the Multimodal models for Integrated Regulatory Analysis[30] (MIRA) python

package to examine dynamic changes in gene expression and chromatin accessibility in

WT and MEF2C KO embryos across the time course of developmental stages that we

collected (E7.75, E8.5, E9). We first created a merged Seurat object and ArchR project

containing only our cardiac cell types of interest. We then used sceasy[58] to convert our

Seurat object and the peak matrix from the ArchR project to AnnData formats, which were

used as inputs for the MIRA analyses.

Topic Modeling

The first step of the MIRA pipeline is to train and tune topic models for both the gene

expression and chromatin accessibility data. Model training and tuning was performed

with GPU support on nodes of the Wynton HPC Co-Op cluster at UCSF. For our gene

expression topic model, we used the 4,088 most variable genes (min_disp = 0.1) as the

"exogenous" gene set and top 1,959 variable genes (dispersion threshold > 0.7) as the

"endogenous" gene set for model training. We then set the minimum and maximum

learning rate bounds, tuned the model (iters = 64, max_topics = 55), and MIRA's tuner

selected the best model that was used for all subsequent analyses, which contained 33

gene expression topics. To aid in the interpretation of results, we exported the top 200

genes for each of the 33 gene expression topics. We then repeated these steps to train

and tune the accessibility topic model, using all features (i.e. there are no "exogenous" or

"endogenous" peak sets), which resulted in a model containing 17 accessibility topics. To

aid in the interpretation of these topics, we performed TF binding motif enrichment

analysis on each topic and exported the TF motif enrichment scores (top_quantile = 0.2)

for each of the 17 accessibility topics.

Pseudotime Analysis and Lineage Trajectories

For the heart tube segment lineages we analyzed (OFT, V, IFT), we created separate

subsetted AnnData objects containing only the cell types that contribute to each lineage.

We generated UMAP embeddings for each lineage based on joint representations of gene

expression and accessibility topics. We then performed pseudotime analysis by

calculating the diffusion map, selecting the appropriate number of diffusion map

components, and generated a K nearest neighbor graph based on these diffusion map

components. In order to create pseudotime trajectories, we randomly selected a WT cell

in the earliest progenitor cell type of each lineage to serve as the initial cell, and then

manually selected terminal cells from the appropriate cell types based on the structure of

each trajectory (for OFT lineage: E9 WT OFT-CM and E9 KO OFT-CM; for V lineage: E9

WT V-CM; for IFT lineage: E8.5 KO IFT-CM, E9 WT A-CM, E9 WT AVC-CM, E9 KO A-

CM, E9 KO AVC-CM). Given these initial and terminal cells, along with the KNN

pseudotime graph, the probability of reaching each terminal state was calculated for every

cell and these probabilities we parsed to create the bifurcating tree lineage trajectory plots

(stream plots). Gene expression and accessibility topics of interest were then plotted

along these stream plots to interpret the dynamic changes in gene expression and

chromatin accessibility that occur in WT and MEF2C KO cells as they differentiate through

each lineage.

Cloning of Reporter Constructions for Zebrafish Transgenesis

To generate the Tg(*MVEB:egfp*)[sfc26] zebrafish lines, 650-900 nucleotide fragments

containing the candidate enhancers were cloned by PCR from mouse genomic DNA.

Candidate nucleotide sequences and the primers used to amplify them are listed in Table

S6. The resulting fragments were cloned into the Tol2 transgenic vector E1b-eGFP-

Tol2[35] (RRID: Addgene_37845) using the In-Fusion Cloning system (Takara Bio). All

plasmid sequences were confirmed by long-read sequencing before zebrafish injection

(Plasmidsaurus).

Zebrafish Transgenesis Injections, Screening, and Imaging

Zebrafish were raised under standard laboratory conditions at 28 °C. The outbred

wildtype zebrafish EKW (*Danio rerio-Ekwill*) and Tg(*cmlc2:mCherry*)[s890] lines have been

described   previously[55].   One-cell   zebrafish   embryos   obtained   from

Tg(*cmlc2:mCherry*)[s890] and wildtype EKW (Ekwill strain) parents were injected with 1nl of

injection mix containing 15pg of candidate-enhancer-eGFP and 15pg of ß-actin-BFP

reporter plasmids along with 30pg of *Tol2* mRNA. The ß -actin promoter drives expression

of BFP strongly in somite muscles and was used as a positive injection control. As a

negative control, we also injected and screened embryos with an empty reporter construct

containing only the minimal E1b promoter and eGFP. Injected embryos were screened at

24 and 72 hpf for eGFP and BFP expression. Representative embryos were imaged for

eGFP expression as a readout of candidate enhancer activity and for mCherry expression

as a cardiac marker. Enhancer activity was scored as a percentage of embryos

expressing eGFP in the heart compared to the total number of BFP-positive (BFP+)

embryos. A minimum of at least 50 BFP+ embryos were screened per construct, except

for MVEB1, for which only 17 BFP+ embryos were obtained. The exact numbers screened

for each construct are displayed in Fig. S5B.

Comparative Analyses with Published Datasets

NR2F2 and GATA4 target genes were downloaded from the Harmonizome[46] and

TFLink[47] databases, respectively. Fisher's exact test was used to determine the Odd's

Ratios and p-values for identifying NR2F2 and GATA4 targets among genes upregulated

in MEF2C KO IFT-CMs at E8.5. Publicly available ChIP-seq datasets were downloaded

from the Gene Expression Omnibus (GEO) in wig or bigwig format for MEF2C

(GSE124008)[26], H3K27ac (GSE52386)[32], GATA4 (GSE52123)[45], and NR2F2

(GSE46498)[42]. Datasets provided in wig format were converted to bigwig format using

the Galaxy[44] web tool wigToBigWig. CrossMap[60] was used together with chain files from

the UCSC Genome Browser[61] to lift genome annotations from mm9 to mm10 reference

genome assemblies, and vice versa. For the browser tracks in Fig. 5B-C, the MEF2C and

H3K27ac ChIP-seq bigwigs were lifted from mm9 to mm10, the Galaxy[44] web tool

bigwigAverage was used to generate a single average MEF2C and H3K27ac occupancy

track from the individual replicate tracks, and the results were displayed along with our

pseudobulked snATAC-seq data using the IGV software package[57]. For the heat maps

displayed in Fig. S6A-B, we first created bed format files for the gained DARs in MEF2C

KO IFT-CMs containing NR or GATA motifs. We then lifted these bed regions from mm10

to mm9 and used the Galaxy[44] web tools computeMatrix and plotHeatmap with the

downloaded NR2F2 and GATA4 ChIP-seq data to visualize NR2F2 and GATA4

occupancy at the DARs. At the computeMatrix step, missing data was converted to

zeroes. At the plotHeatmap step, kmeans clustering was used to generate two clusters of

DARs (effectively "occupied" and "not occupied") and these were sorted by decreasing

mean value prior to being plotted.

Gene Regulatory Network Inference and Analysis

We used CellOracle[49] to infer GRNs for WT and MEF2C KO IFT-CMs at E8.5 and E9.

The GRN inference process included the following steps:

Base GRN Construction

We used OFT-CM, V-CM, and IFT-CM snATAC-seq data to construct a base GRN that

defines the accessible TF binding motifs near genes for each of the following conditions:

E8.5-WT, E8.5-KO, E9-WT, and E9-KO. Pooling the OFT-CM, V-CM, and IFT-CMs

mitigated artifacts caused by data sparsity. The binarized PeakMatrix output from ArchR

was used as an input to Cicero's[62] run_cicero() function (with default parameters) to

compute co-accessibility scores among peak pairs within 500kb. Output scores were from

-1 to 1, where higher scores indicate stronger co-accessibility. Each dataset generated

over 20 million peak pairs for co-accessibility comparisons. We annotated transcription

start sites (TSSs) with their corresponding genes for accessible peaks using the

motif_analysis.get_tss_info() function in CellOracle. Unlabeled peaks were considered

potential   regulatory   elements.   Subsequently,   we   applied

motif_analysis.integrate_tss_peak_with_cicero() to link unlabeled peaks with genes,

filtering for peak-to-gene pairs with co-accessibility scores >0.8. TF binding motifs within

each 500bp peak were identified using the motif_analysis.TFinfo() function in CellOracle,

which converts peak IDs into DNA sequences using the mm10 reference genome. TF

motif scanning used the gimme.vertebrate.v5.0[63] library to produce a set of TF motifs

present in accessible regions within 500kb of each gene. The TFs that bind to these motifs

have the potential to regulate the given gene.

GRN Inference

Using CellOracle, we constructed GRNs in which edge strengths are quantified based on the TF-to-gene correlation, constrained by the accessible regions defined in the base GRN. We inferred GRNs for E8.5 WT, E8.5 KO, E9 WT, and E9 KO IFT-CMs, using the computed base GRNs to determine permissible TF-to-gene links. GRN inference used the top 5,000 differentially expressed genes in the snRNA-seq dataset. We used the CellOracle oracle.get_links() function (with default parameters), fitting linear models with L2 weight regularization to derive coefficients relating TF expression to target gene expression. The Bagging Ridge model was used to fit coefficients, providing posterior distributions for each TF-to-gene relationship. Positive coefficients indicated activating, while negative coefficients indicated inhibiting relationships. We then filtered the inferred network to include 4,000 edges with the largest magnitude coefficients.

Simulation of MEF2C Knockout Effects

To predict the effect of MEF2C KO on gene expression, we identified genes directly regulated by MEF2C in the E8.5 WT IFT-CM network. We predicted that reduced expression of *Mef2c* in the network would downregulate genes directly activated by MEF2C and upregulate genes directly inhibited by MEF2C in the E8.5 WT IFT-CM network. The simulation results were compared against differentially expressed genes in the E9 snRNA-seq IFT-CM experimental data (WT-vs-KO). Differentially expressed genes were identified by selecting for genes with logfold change magnitude >1.5.

Network Visualization

Whole-network visualizations were generated using Gephi[64], applying the Yifan Hu layout, with node sizes scaled by betweenness centrality. PyVis[65] was used to visualize subnetworks that included manually selected TFs (*Mef2a, Mef2c, Nr2f2, Tbx5, Gata4, Meis2, Plagl1, Isl1, Pitx2, Tbx20, Hand2, Nkx2-5*) and the the top 100 differentially expressed genes (ranked in ascending order by p value) identified by comparing the E8.5 WT and MEF2C KO IFT-CM snRNA-seq experimental data. Visualizations for direct interactions between TFs (*Nr2f2* and *Gata4*) and targets in the E8.5 WT and KO IFT-CMs

879   were developed using Python's NetworkX package[66], with a circular layout and edge
880   widths scaled by coefficient values.

ACKNOWLEDGEMENTS

Sequencing was performed at the UCSF CAT, supported by UCSF PBBR, RRP IMIA, and NIH 1S10OD028511-01 grants. We thank the Gladstone Genomics Core for use and maintenance of the Agilent Bioanalyzer and 10x Genomics Chromium Controller. Portions of this work were performed on the Wynton HPC Co-Op cluster, which is supported by UCSF research faculty and UCSF institutional funds. The authors wish to thank the UCSF Wynton team for their ongoing technical support of the Wynton environment. J.M.M-V. is grateful for assistance from Dr. Kavitha Rao in learning Seurat, as well as from Dr.'s Ryan Corces and Christina Theodoris for assistance with ArchR and MIRA, respectively. J.M.M-V. is also grateful to all his colleagues in the Bruneau Lab for many insightful discussions, important feedback on directions of project, and help with experiments.

AUTHOR CONTRIBUTIONS

J.M.M-V. and B.G.B. conceived and designed the experiments presented in this study. J.M.M-V. performed embryo dissections, whole-mount embryo immunostaining, whole-mount embryo FISH via HCR, embryo imaging, preparation of sequencing libraries, and data analysis via Seurat, ArchR, and MIRA. E.B. performed mouse colony maintenance and genotyping. J.M.M-V. and E.B. performed cloning of reporter constructs for zebrafish transgenesis. T.S. performed zebrafish transgenesis injections and imaging. A.P.C. performed GRN analyses using CellOracle and newly written code. B.G.B., B.L.B., and J.J.S. supervised and advised. J.M.M-V. prepared the figures and wrote the manuscript with input from the co-authors.

DECLARATION OF INTERESTS

B.G.B. is a founder, shareholder, and advisor of Tenaya Therapeutics and an advisor for Silver Creek Pharmaceuticals. The work presented here is not related to the interests of these commercial entities.

FUNDING

This work was funded by grants from the National Heart, Lung, and Blood Institute (NHLBI; R01-HL160665 and R01-HL162925 to J.J.S.; R01-HL114948 and R01-

912 HL155906 to B.G.B.), the Roddenberry Foundation (B.G.B.), Additional Ventures
913 (B.G.B.), and the Younger Family Fund (B.G.B.). J.M.M-V. was supported by fellowships
914 from the NHLBI (F32-HL162450) and the American Heart Association together with The
915 Children's Heart Foundation (24POST1191660). A.P.C. was supported by the NHLBI
916 (T32-HL007284).

DATA AND CODE AVAILABILITY

918
919 Raw and processed data for snRNA-seq and snATAC-seq datasets has been deposited
920 in the NCBI Gene Expression Omnibus (GEO) under accession number GSE280587.
921 Code used for multiomics analyses and plot generation is available in the GitHub
922 repository associated with this manuscript: https://github.com/jmmuncie/Muncie-
923 Vasic_bioRxiv_2024.

FIGURES AND CAPTIONS

Figure 1

Figure 1: Although MEF2C is expressed throughout the developing heart tube, loss of MEF2C causes segment-specific defects.

A) Schematic of cardiac progenitors and their contributions to linear heart tube development from cardiac crescent (E7.75) to looped heart tube (E9) stage. B) Immunofluorescent staining of MEF2C (cyan) and cardiac Troponin T (cTnT, magenta) in E7.75, E8.5, and E9 WT embryos. C) Representative images of WT and MEF2C KO embryos at E7.75, E8.5 and E9. Cardiac progenitors are marked by the *Smarcd3*-F6-eGFP reporter transgene (green). BF, brightfield. D) Schematic of the methodology and biological insights presented in the current study. Elements of this panel were created in

937   BioRender. B, B. (2024) https://BioRender.com/i72e213. Scale bars = 200 µm. FHF, first

938   heart field; aSHF, anterior second heart field; pSHF, posterior second heart field; LV, left

939   ventricle; RV, right ventricle; V, ventricle; IFT, inflow tract; OFT, outflow tract.

Figure 2

Figure 2: Loss of MEF2C leads to broad changes in gene expression throughout the heart tube, including mis-regulation of key cardiomyocyte genes and anterior/posterior markers.

A-C) UMAPs of snRNA-seq data for cardiac progenitors, cardiomyocytes, and related mesoderm subtypes from E7.75 (A), E8.5 (B), and E9 (C) embryos labeled by cell type (left) and genotype/sample ID (right). D-F) Bar plots displaying the number of up-regulated and down-regulated genes (MEF2C KO-vs-WT) for cell types of interest at E7.75 (D), E8.5 (E), and E9 (F). G-I) Dot plots displaying gene expression of key CM genes and anterior/posterior (A/P) markers at E7.75 (G), E8.5 (H), and E9 (I). J

Fluorescence *in situ* hybridization of key CM genes and A/P markers in E8-E8.5 WT and MEF2C KO embryos. Scale bars = 200 μm. CMs, cardiomyocytes; FHF, first heart field; SHF, second heart field; JCF, juxtacardiac field; CrM, cranial mesoderm; PrxM, paraxial mesoderm; LPM, lateral plate mesoderm; SoM, somitic mesoderm; NMPs, neuromesodermal progenitors; KPs, kidney progenitors; ExM, extraembryonic mesoderm; HSCs, hematopoietic stem cells; V-CMs, ventricular cardiomyocytes; IFT-CMs, inflow tract cardiomyocytes; OFT-CMs, outflow tract cardiomyocytes; aSHF, anterior second heart field; pSHF, posterior second heart field; PostM, posterior mesoderm; PhM, pharyngeal mesoderm; MixM, mixed mesoderm; A-CMs, atrial cardiomyocytes; AVC-CMs, atrioventricular canal cardiomyocytes; Pe, proepicardium; VP, venous pole; *, Genes known to be associated with CHDs[25]; #, Direct targets of MEF2C based on MEF2C ChIP-seq data[26].

Figure 3

Figure 3: MEF2C regulates chromatin accessibility broadly throughout the heart tube and in a segment-specific manner.

A) UMAP of integrated snRNA-seq and snATAC-seq data for cardiac progenitors, cardiomyocytes, and related mesoderm subtypes at E8.5 labeled by cell types determined from snRNA-seq clustering (left) and genotype/sample ID (right). B) Bar plots displaying the number of gained and lost DARs (MEF2C KO-vs-WT) for cell types of interest at E8.5. C) Venn diagrams displaying unique and overlapping DARs in the three heart tube segments. D) Scatter plots displaying the relationship between the Log2 fold change (Log2FC) values of the DEG and DAR analyses for all identified Peak2Gene (P2G) links in the three heart tube segments. Dots are colored by the P2G correlation score. E-F) Genome browser tracks displaying snATAC-seq accessibility profiles at the *Myh6/Myh7* (E) and *Wnt2* (F) loci for the indicated pseudobulked cell types. DARs are highlighted and the peaks are indicated by red bars (MEF2C KO-vs-WT IFT-CMs). Loops indicate P2G links, colored by the P2G correlation score. V-CMs, ventricular

981   cardiomyocytes; IFT-CMs, inflow tract cardiomyocytes; OFT-CMs, outflow tract

982   cardiomyocytes; aSHF, anterior second heart field; pSHF, posterior second heart field;

983   LPM, lateral plate mesoderm; PostM, posterior mesoderm; PhM, pharyngeal mesoderm;

984   NA, cells not available in the snRNA-seq dataset.

Figure 4

Figure 4: Pseudotime analysis with MIRA reveals distinct MEF2C-depedent developmental trajectories for each heart tube segment.

A) Schematic of multimodal models for integrated regulatory analysis (MIRA) pipeline. B) UMAP of outflow tract lineage cells plotted by MIRA topic models labeled by cell type (top)

and genotype (bottom). C) Pseudotime plot of outflow tract lineage cells. D) Lineage

trajectory stream plots for outflow tract lineage cells labeled by cell type (top) and

genotype (bottom). E) Stream plots displaying the flow of gene expression (top) and

chromatin accessibility (bottom) topics in the outflow tract lineage cells. Examples of the

top genes and TF binding motifs for dynamic topics are labeled. F) UMAP of vericular

lineage cells plotted by MIRA topic models labeled by cell type (top) and genotype

(bottom). G) Pseudotime plot of ventricular lineage cells. H) Lineage trajectory stream

plots for ventricular lineage cells labeled by cell type (top) and genotype (bottom). E)

Stream plots displaying the flow of gene expression (top) and chromatin accessibility

(bottom) topics in the ventricular lineage cells. Examples of the top genes and TF binding

motifs for dynamic topics are labeled. J) UMAP of inflow tract lineage cells plotted by

MIRA topic models labeled by cell type (top) and genotype (bottom). K) Pseudotime plot

of inflow tract lineage cells. L) Lineage trajectory stream plots for inflow tract lineage cells

labeled by cell type (top) and genotype (bottom). M) Stream plots displaying the flow of

gene expression (top) and chromatin accessibility (bottom) topics in the inflow tract

lineage cells. Examples of the top genes and TF binding motifs for dynamic topics are

labeled.

Figure 5

1010

1011

Figure 5: Leveraging multiomic data to identify novel enhancers via zebrafish

transgenesis.

A) Schematic of selection process to identify candidate MEFC-dependent enhancers from

integrated snRNA-seq and snATAC-seq data. B) Genome browser tracks displaying

snATAC-seq accessibility profiles, MEF2C ChIP-seq occupancy profiles[26], and H3K27ac

ChIP-seq occupancy profiles[32] at example loci containing IFT-specific (left), V-specific

(middle), and OFT-specific (right) candidate enhancers (yellow highlights). C) Genome

browser tracks displaying snATAC-seq accessibility profiles, MEF2C ChIP-seq

occupancy profiles[26], and H3K27ac ChIP-seq occupancy profiles[32] at the *Myh6*/*Myh7*

locus, which contains two candidate enhancers (yellow highlights, MVEB1 and MVEB2)

with IFT-specific altered accessibility that overlap with regions found in the VISTA

Enhancer Browser database[34]. D) Images of E11.5 mouse embryos from the Vista

Enhancer Browser database[34] demonstrating positive enhancer activity of regions that

overlap with candidates MVEB1 and MVEB2. E) Schematic of the Tol2 Transgenesis

assay used to screen candidate enhancers in zebrafish. Elements of this panel were

created in BioRender. B, B. (2024) https://BioRender.com/h83r503. F) Representative

ventral view images of zebrafish embryos at 72 hpf injected with candidate enhancers

that demonstrated positive activity in the heart. Boxed area in the representative

brightfield image (left) indicates the anatomical region of interest captured in the

fluorescent images. G) Schematic representation of the observed onset of enhancer

activity for candidate enhancers that demonstrated positive activity in the heart. H)

Representative ventral view images of zebrafish embryos at 24 and 72 hpf injected with

MVEB2:eGFP or MVEB6:eGFP reporter constructs. Scale bars = 100 μm.

Figure 6

1036

1037

Figure 6: An overactive "posteriorized" GRN driven by NR2F2 and GATA4 underlies

the expanded inflow tract phenotype in MEF2C KO.

A) TF binding motif enrichment analysis for gained DARs (MEF2C KO-vs-WT) in IFT-CMs

at E8.5. B) Pie charts showing the proportion of gained or lost DARs (MEF2C KO-vs-WT)

containing NR and MEF2C motifs or only NR motifs (top), and those containing GATA and

MEF2C motifs or only GATA motifs (bottom). C) Odds ratio analysis for NR2F2 target

genes[46] (left) and GATA4 target genes[47] (right) amongst DEGs up-regulated in MEF2C

KO IFT-CMs. p-values were calculated using Fisher's exact test. D) Ridge plots displaying

module scores for up-regulated NR2F2 targets (left) and GATA4 targets (right) in MEF2C

KO and WT heart tube segments. E) Inferred GRNs constructed for MEF2C KO and WT

IFT-CMs at E8.5 and E9. F) Schematic of GRN validation by predicting gene expression

with *Mef2c* knockdown in the E8.5 WT network and comparing to measured gene

expression changes at E9. G) Results of GRN validation displaying predicted and

measured gene expression changes. H) Visualizations of cardiac subnetworks within the

WT and MEF2C KO E8.5 IFT-CM GRNs. I-J) Visualizations of direct NR2F2 (I) and

GATA4 (J) interactions in the WT and MEF2C KO E8.5 IFT-CM GRNs. Direct interactions

that occur upon MEF2C KO are highlighted. #, up-regulated DEG in MEF2C KO IFT-CMs

at E8.5; *, Direct target of NR2F2 or GATA4[46,47]. K-L) Immunofluorescent staining of

cardiac Troponin T (cTnT, green) in representative E9.5 embryos (18-24 somites)

collected from *Mef2c*$^{del/+}$;*Nr2f2*$^{del/+}$ to *Mef2c*$^{del/+}$ crosses. Boxed regions in (K) are shown

at higher magnification in (L). Arrows indicate more expanded ventricle phenotype in

Mef2c$^{del/del}$;*Nr2f2*$^{del/+}$ embryos. Asterisks indicate better developed and looping atria in

Mef2c$^{del/del}$;*Nr2f2*$^{del/+}$ embryos. n=5 *Mef2c*$^{del/del}$ and n=4 *Mef2c*$^{del/del}$;*Nr2f2*$^{del/+}$ embryos

from 5 independent litters. Scale bars = 200 μm.

REFERENCES

1. Bruneau BG. Signaling and Transcriptional Networks in Heart Development and Regeneration. Cold Spring Harb Perspect Biol. 2013;5(3):a008292.

2. Cohen M, Briscoe J, Blassberg R. Morphogen interpretation: the transcriptional logic of neural tube patterning. Curr Opin Genet Dev. 2013;23(4):423–8.

3. McQueen C, Towers M. Establishing the pattern of the vertebrate limb. Development. 2020;147(17):dev177956.

4. Hafner A, Boettiger A. The spatial organization of transcriptional control. Nat Rev Genet. 2023;24(1):53–68.

5. Wagh K, Stavreva DA, Upadhyaya A, Hager GL. Transcription Factor Dynamics: One Molecule at a Time. Annu Rev Cell Dev Biol. 2023;39(1):277–305.

6. Spitz F, Furlong EEM. Transcription factors: from enhancer binding to developmental control. Nat Rev Genet. 2012;13(9):613–26.

7. Lee TI, Young RA. Transcriptional Regulation and Its Misregulation in Disease. Cell. 2013;152(6):1237–51.

8. Mercatelli D, Scalambra L, Triboli L, Ray F, Giorgi FM. Gene regulatory network inference resources: A practical overview. Biochim Biophys Acta (BBA) - Gene Regul Mech. 2020;1863(6):194430.

9. Olson EN. Gene Regulatory Networks in the Evolution and Development of the Heart. Science. 2006;313(5795):1922–7.

10. Kelly RG, Buckingham ME, Moorman AF. Heart Fields and Cardiac Morphogenesis. Cold Spring Harb Perspect Med. 2014;4(10):a015750.

11. Evans SM, Yelon D, Conlon FL, Kirby ML. Myocardial Lineage Development. Circ Res. 2010;107(12):1428–44.

12. Parameswaran M, Tam PPL. Regionalisation of cell fate and morphogenetic
movement of the mesoderm during mouse gastrulation. Dev Genet. 1995;17(1):16–28.

13. Lopez-Sanchez C, Garcia-Martinez V, Schoenwolf GC. Localization of Cells of the
Prospective Neural Plate, Heart and Somites within the Primitive Streak and Epiblast of
Avian Embryos at Intermediate Primitive-Streak Stages. Cells Tissues Organs.
2001;169(4):334–46.

14. Stalsberg H, DeHaan RL. The precardiac areas and formation of the tubular heart in
the chick embryo. Dev Biol. 1969;19(2):128–59.

15. DeRuiter MC, Poelmann RE, Vries IV de, Mentink MMT, Groot ACG de. The
development of the myocardium and endocardium in mouse embryos. Anat Embryol.
1992;185(5):461–73.

16. Lescroart F, Chabab S, Lin X, Rulands S, Paulissen C, Rodolosse A, et al. Early
lineage restriction in temporally distinct populations of Mesp1 progenitors during
mammalian heart development. Nat Cell Biol. 2014;16(9):829–40.

17. Devine WP, Wythe JD, George M, Koshiba-Takeuchi K, Bruneau BG. Early patterning
and specification of cardiac progenitors in gastrulating mesoderm. eLife. 2014;3:e03848.

18. Meilhac SM, Kelly RG, Rocancourt D, Eloy-Trinquet S, Nicolas JF, Buckingham ME.
A retrospective clonal analysis of the myocardium reveals two phases of clonal growth in
the developing mouse heart. Development. 2003;130(16):3877–89.

19. Lin Q, Schwarz J, Bucana C, Olson EN. Control of Mouse Cardiac Morphogenesis
and Myogenesis by Transcription Factor MEF2C. Science. 1997;276(5317):1404–7.

20. Krup AL, Winchester SAB, Ranade SS, Agrawal A, Devine WP, Sinha T, et al. A
Mesp1-dependent developmental breakpoint in transcriptional and epigenomic
specification of early cardiac precursors. Development. 2023;150(9):dev201229.

21. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated
analysis of multimodal single-cell data. Cell. 2021;184(13):3573-3587.e29.

22. Choudhary S, Satija R. Comparison and evaluation of statistical error models for
scRNA-seq. Genome Biol. 2022;23(1):27.

23. Tyser RCV, Ibarra-Soria X, McDole K, Jayaram SA, Godwin J, Brand TAH van den,
et al. Characterization of a common progenitor pool of the epicardium and myocardium.
Science. 2021;371(6533).

24. Black BL, Olson EN. TRANSCRIPTIONAL CONTROL OF MUSCLE DEVELOPMENT
BY MYOCYTE ENHANCER FACTOR-2 (MEF2) PROTEINS. Annu Rev Cell Dev Biol.
1998;14(1):167–96.

25. Yang A, Alankarage D, Cuny H, Ip EKK, Almog M, Lu J, et al. CHDgene: A Curated
Database for Congenital Heart Disease Genes. Circ: Genom Precis Med.
2022;15(3):e003539.

26. Akerberg BN, Gu F, VanDusen NJ, Zhang X, Dong R, Li K, et al. A reference map of
murine cardiac transcription factor chromatin occupancy identifies dynamic and
conserved enhancers. Nat Commun. 2019;10(1):4907.

27. Barnes RM, Harris IS, Jaehnig EJ, Sauls K, Sinha T, Rojas A, et al. MEF2C regulates
outflow tract alignment and transcriptional control of Tdgf1. Development.
2016;143(5):774–9.

28. Granja JM, Corces MR, Pierce SE, Bagdatli ST, Choudhry H, Chang HY, et al. ArchR
is a scalable software package for integrative single-cell chromatin accessibility analysis.
Nat Genet. 2021;53(3):403–11.

29. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-
based Analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137.

30. Lynch AW, Theodoris CV, Long HW, Brown M, Liu XS, Meyer CA. MIRA: joint
regulatory modeling of multimodal expression and chromatin accessibility in single cells.
Nat Methods. 2022;19(9):1097–108.

31. Zhou W, Lin L, Majumdar A, Li X, Zhang X, Liu W, et al. Modulation of morphogenesis
by noncanonical Wnt signaling requires ATF/CREB family–mediated transcriptional
activation of TGFβ2. Nat Genet. 2007;39(10):1225–34.

32. Nord AS, Blow MJ, Attanasio C, Akiyama JA, Holt A, Hosseini R, et al. Rapid and
Pervasive Changes in Genome-wide Enhancer Usage during Mammalian Development.
Cell. 2013;155(7):1521–31.

33. Man JCK, Duijvenboden K van, Krijger PHL, Hooijkaas IB, Made I van der, Vries C
de G de, et al. Genetic Dissection of a Super Enhancer Controlling the Nppa-Nppb Cluster
in the Heart. Circ Res. 2021;128(1):115–29.

34. Visel A, Minovitsky S, Dubchak I, Pennacchio LA. VISTA Enhancer Browser—a
database of tissue-specific human enhancers. Nucleic Acids Res.
2007;35(suppl_1):D88–92.

35. Birnbaum RY, Clowney EJ, Agamy O, Kim MJ, Zhao J, Yamanaka T, et al. Coding
exons function as tissue-specific enhancers of nearby genes. Genome Res.
2012;22(6):1059–68.

36. Fisher S, Grice EA, Vinton RM, Bessling SL, Urasaki A, Kawakami K, et al. Evaluating
the biological relevance of putative enhancers using Tol2 transposon-mediated
transgenesis in zebrafish. Nat Protoc. 2006;1(3):1297–305.

37. Hinits Y, Pan L, Walker C, Dowd J, Moens CB, Hughes SM. Zebrafish Mef2ca and
Mef2cb are essential for both first and second heart field cardiomyocyte differentiation.
Dev Biol. 2012;369(2):199–210.

38. Zlatanova I, Sun F, Wu RS, Chen X, Lau BH, Colombier P, et al. An injury-responsive
mmp14b enhancer is required for heart regeneration. Sci Adv. 2023;9(48):eadh5313.

39. Yan R, Cigliola V, Oonk KA, Petrover Z, DeLuca S, Wolfson DW, et al. An enhancer-
based gene-therapy strategy for spatiotemporal control of cargoes during tissue repair.
Cell Stem Cell. 2023;30(1):96-111.e6.

40. Kang J, Hu J, Karra R, Dickson AL, Tornini VA, Nachtrab G, et al. Modulation of tissue
repair by regeneration enhancer elements. Nature. 2016;532(7598):201–6.

41. Tremblay M, Sanchez-Ferras O, Bouchard M. GATA transcription factors in
development and disease. Development. 2018;145(20):dev164384.

42. Wu S pin, Cheng CM, Lanz RB, Wang T, Respress JL, Ather S, et al. Atrial Identity Is
Determined by a COUP-TFII Regulatory Network. Dev Cell. 2013;25(4):417–26.

43. Pereira FA, Qiu Y, Zhou G, Tsai MJ, Tsai SY. The orphan nuclear receptor COUP-
TFII is required for angiogenesis and heart development. Genes Dev. 1999;13(8):1037–
49.

44. Community TG, Abueg LAL, Afgan E, Allart O, Awan AH, Bacon WA, et al. The Galaxy
platform for accessible, reproducible, and collaborative data analyses: 2024 update.
Nucleic Acids Res. 2024;52(W1):W83–94.

45. He A, Gu F, Hu Y, Ma Q, Ye LY, Akiyama JA, et al. Dynamic GATA4 enhancers shape
the chromatin landscape central to heart development and disease. Nat Commun.
2014;5(1):4907.

46. Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG,
et al. The harmonizome: a collection of processed datasets gathered to serve and mine
knowledge about genes and proteins. Database. 2016;2016(2016):baw100.

47. Liska O, Bohár B, Hidas A, Korcsmáros T, Papp B, Fazekas D, et al. TFLink: an
integrated gateway to access transcription factor–target gene interactions for multiple
species. Database. 2022;2022:baac083.

48. Tirosh I, Izar B, Prakadan SM, II MHW, Treacy D, Trombetta JJ, et al. Dissecting the
multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. Science.
2016;352(6282):189–96.

49. Kamimoto K, Stringa B, Hoffmann CM, Jindal K, Solnica-Krezel L, Morris SA.
Dissecting cell identity via network inference and in silico gene perturbation. Nature.
2023;614(7949):742–51.

50. Molkentin JD, Black BL, Martin JF, Olson EN. Cooperative activation of muscle gene
expression by MEF2 and myogenic bHLH proteins. Cell. 1995;83(7):1125–36.

51. Di-Luoffo M, Brousseau C, Tremblay JJ. MEF2 and NR2F2 cooperate to regulate
Akr1c14 gene expression in mouse MA-10 Leydig cells. Andrology. 2016;4(2):335–44.

52. Molkentin JD, Black BL, Martin JF, Olson EN. Mutational Analysis of the DNA Binding,
Dimerization, and Transcriptional Activation Domains of MEF2C. Mol Cell Biol.
1996;16(6):2627–36.

53. Pereira FA, Tsai MJ, Tsai* SY. COUP-TF orphan nuclear receptors in development
and differentiation. Cell Mol Life Sci CMLS. 2000;57(10):1388–98.

54. Takamoto N, You LR, Moses K, Chiang C, Zimmer WE, Schwartz RJ, et al. COUP-
TFII is essential for radial and anteroposterior patterning of the stomach. Development.
2005;132(9):2179–89.

55. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive
and accurate integration of single-cell data with Harmony. Nat Methods.
2019;16(12):1289–96.

56. Vierstra J, Lazar J, Sandstrom R, Halow J, Lee K, Bates D, et al. Global reference
mapping of human transcription factor footprints. Nature. 2020;583(7818):729–36.

57. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24–6.

58. Cakir B, Prete M, Huang N, Dongen S van, Pir P, Kiselev VY. Comparison of
visualization tools for single-cell RNAseq data. NAR Genom Bioinform.
2020;2(3):lqaa052.

59. Zhang R, Han P, Yang H, Ouyang K, Lee D, Lin YF, et al. In vivo cardiac
reprogramming contributes to zebrafish heart regeneration. Nature.
2013;498(7455):497–501.

60. Zhao H, Sun Z, Wang J, Huang H, Kocher JP, Wang L. CrossMap: a versatile tool for
coordinate conversion between genome assemblies. Bioinformatics. 2014;30(7):1006–7.

61. Raney BJ, Barber GP, Benet-Pagès A, Casper J, Clawson H, Cline MS, et al. The
UCSC Genome Browser database: 2024 update. Nucleic Acids Res.
2023;52(D1):D1082–8.

62. Pliner HA, Packer JS, McFaline-Figueroa JL, Cusanovich DA, Daza RM, Aghamirzaie
D, et al. Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin
Accessibility Data. Mol Cell. 2018;71(5):858-871.e8.

63. Bruse N, Heeringen SJ van. GimmeMotifs: an analysis framework for transcription
factor motif analysis. bioRxiv. 2018;474403.

64. Bastian M, Heymann S, Jacomy M. Gephi: An Open Source Software for Exploring
and Manipulating Networks. ICWSM. 2009;3:361–2.

65. Perrone G, Unpingco J, Lu H minn. Network visualizations with Pyvis and VisJS.
arXiv. 2020;

66. Hagberg AA, Schult DA, Swart PJ. Exploring Network Structure, Dynamics, and
Function using NetworkX. In: Varoquaux G, Vaught T, Millman J, editors. Proceedings of
the 7th Python in Science Conference (SciPy 2008). 2008. p. 11–6.