1 **The coordination between CTCF, cohesin and TFs impacts nucleosome repositioning**
2 **and chromatin insulation to define state specific 3D chromatin folding**
3

4 Catherine Do[1], Guimei Jiang[1], Giulia Cova[1], Paul Zappile[3], Adriana Heguy[3] & Jane A Skok[1,2,*]
5
6 1.   Department of Pathology, NYU Grossman School of Medicine, New York, NY, USA
7 2.   Perlmutter Cancer Center, NYU Langone Health, New York, NY, USA
8 3.   Genome Technology Center, NYU Grossman School of Medicine, New York, NY, USA
9
10 * Corresponding author: Jane.Skok@nyulangone.org

11
12
13
14

Abstract (250 words)

CTCF-mediated chromatin folding plays a key role in gene regulation, however the mechanisms controlling chromatin organization across cell states are not fully elucidated. Comprehensive analyses reveal that CTCF binding stability and cohesin overlap in mice and humans, are regulated by species specific differences in CTCF binding site (CBS) accessibility and enrichment of motifs corresponding to expressed TFs. By analyzing TFs, we confirm the co-operativity and competitiveness of TF/CTCF binding, which we further validate by allele specific analysis of SNPs. TF motif enrichment at CTCF bound sites is determined by cell state-specific transcriptional programs, which either stabilize or destabilize CTCF binding, as reflected by changing TF concentration. To examine CTCF binding in the context of nucleosome positioning, we performed single molecule nano-NOMe-seq. Subsetting reveals a continuum of binding states for CTCF, which are differentially represented at accessible versus inaccessible CBSs. As expected, CTCF degradation leads to a progressive loss of binding and nucleosome repositioning giving profiles similar to CTCF free CBSs. We also observe a similar time dependent effect when the cohesin subcomponent, SSC1 is degraded although CTCF remains bound, indicating that cohesin mediates CTCF-associated nucleosome repositioning. Stratified analysis of CTCF signal strength and accessibility reveals that in the presence of cohesin, CTCF strength contributes to nucleosome repositioning and chromatin insulation independent of accessibility. However, cobound TFs can uncouple the relationship between signal strength and nucleosome repositioning, without affecting the connection between repositioning and insulation. These studies identify mechanisms underlying cell state-specific CTCF profiles, linked to local and long-range chromatin organization.

Introduction

CTCF is an eleven zinc finger protein that makes contact with a 12-15bp DNA motif through its central ZFs, 3-7. Together CTCF and the cohesin complex promote the formation of chromatin loops within highly self-interacting topologically associated domain (TAD) structures, via a loop extrusion mechanism [1,2]. The two factors are enriched at TAD boundaries where they function as insulators promoting interactions between enhancers and promoters within a domain, and restricting the action of regulatory elements to target genes outside a domain. CTCF mediated chromatin folding plays a key role in gene regulation as exemplified by numerous studies showing that disruption of insulated TAD boundaries by deletion of CTCF binding sites (CBSs) can alter gene expression leading to defects in developmental processes and cancer initiation as a result of illegitimate enhancer promoter contacts [3-7].

CTCF dependent chromatin interactions can be conserved, or cell-state specific [8], however the mechanisms underlying CTCF's ability to act as a chromatin organizer in a developmental stage, or disease specific manner have not been fully elucidated. We and others have found that CTCF can bind both accessible and inaccessible sites, with differential functional outcomes [9,10]. By analyzing the interplay between CTCF, cohesin and ATAC-seq data in CTCF WT and mutant expressing cells, we demonstrated that accessibility is a key determinant of CTCF binding stability, and CTCF signal strength in turn is the strongest predictor of cohesin overlap and chromatin insulation score [9]. Specifically, when CTCF binds accessible sites, it has a stronger signal and is more efficient at blocking cohesin and participating in chromatin loop formation than CTCF at inaccessible sites [9]. However, the relationship between accessibility and CTCF binding stability is not clear cut, because a subset of inaccessible CTCF bound sites have strong signals with cohesin overlap, and conversely, CTCF at accessible sites can exhibit weak signals devoid of cohesin. Furthermore, there are numerous accessible CBSs that remain unbound. Thus, other factors must positively and negatively contribute to binding.

While previous studies have identified a handful of transcription factor (TF) cofactors that stabilize CTCF binding in specific cell types, a comprehensive analysis exploring the role of TFs in cell type-specific binding is lacking [11-16]. In this study, we demonstrate that DNA sequences found within a 35 bp window surrounding CBSs motifs corresponding to expressed cell type-specific TFs can differentially affect CTCF signal strength and its capacity to block cohesin, in a manner that depends on whether they are located in accessible or inaccessible chromatin. In both mice and humans, we identified enriched TF motifs showing strong similarity within specific windows,

suggesting potential redundancy in TF-mediated regulation of CTCF binding and a mechanism
for cell type-specific control. We also highlight differences between the two species: notably,
mouse and human differ respectively in the proportion of inaccessible versus accessible CBSs in
their genomes, which is reflected in the regulatory mechanisms that govern their binding. Although
TF motifs are present at most binding sites in both species, in mice they are enriched at
inaccessible sites, while in humans they are more abundant at accessible sites.

To examine the impact of transcriptional programming on CTCF binding, we looked at how
variations in individual TF levels across different cell states and cell types are associated with
changes in their motif enrichment at CTCF sites. In contrast to previous findings [11-16], we show
that TFs can function to both stabilize and destabilize CTCF binding, as reflected by the direction
of enrichment at increasing TF concentration. By analyzing a handful of TFs, we could confirm
both their co-operativity and competitiveness with CTCF binding at CBSs. Consistent with our
ChIP-seq analysis, AlphaFold3 predicts that at individual locations, TFs can respectively compete
or cobind with CTCF depending on whether TF binding overlaps the CBS or occurs up /
downstream of it. Allele specific validation further highlights that SNPs affecting either CTCF or
TF binding have reciprocal effects on the binding of each factor. Moreover, changes in binding of
either factor can impact cohesin binding and to a lesser extent, ATAC-seq signals. Consequently,
changes in TF expression levels can convert a weak CTCF-bound site into a strong site that halts
the progression of cohesin, or vice versa, depending on whether the adjacent TF binds
cooperatively or competitively with CTCF and whether its expression increases or decreases.
This duality helps explain why many accessible CBSs are not occupied by CTCF itself, but rather
by TFs that prevent its localization

Using long read, nano-NOME-seq single molecule footprinting, we analyzed the impact of CTCF
binding on nucleosome repositioning in different genomic contexts. The single molecule approach
allowed us to subset our data revealing a continuum of binding states for CTCF which are present
at individual peaks, and differentially represented at accessible and inaccessible sites. This
information has been missing from previous bulk analyses that average all the data. Using the
clustered analysis, we demonstrate that CTCF has a reduced ability to reposition nucleosomes
at inaccessible sites. Furthermore, we found that (i) the profiles of degraded CTCF and CTCF
free CBSs were very similar, reflecting nucleosome occupied CBSs or binding of TFs that can still
phase nucleosomes but cannot reposition them, and (ii) degradation of the SCC1 subcomponent
of cohesin gave rise to a similar profile while CTCF remains bound, pointing to the interesting

finding that cohesin and loop extrusion mediate nucleosome positioning and phasing by creating

or stabilizing the barrier effects at CTCF anchors.

A stratified analysis of CTCF signal strength and accessibility revealed that in the presence of

cohesin, CTCF strength contributes to nucleosome repositioning and insulation independent of

accessibility. However, cobound TFs can uncouple the relationship between signal strength and

nucleosome repositioning without affecting the connection between nucleosome repositioning

and insulation. These findings indicate that nucleosome repositioning is a more reliable predictor

of chromatin insulation than CTCF signal strength and that TF cobinding can modulate loop

extrusion at CTCF anchors. Together, these studies demonstrate that cell state-specific

transcriptional programs, shape CTCF profiles linked to local and long-range chromatin

organization and gene regulation.

Results

TF motifs enriched within 35 bps of CTCF bound sites stabilize CTCF binding

A number of studies have reported that CTCF binding can be stabilized by the cobinding of cell-

type specific TFs adjacent to CTCF [11-16], but there have been no comprehensive, unbiased

analyses to determine how cell type-specific cofactors regulate CTCF binding and its loop forming

capacity. In addition, previous studies did not take into account the chromatin accessibility of

CTCF binding sites, which can play an important role in CTCF cell-specific binding and function

as previously shown by us and others[9,10].

To investigate, we used mESC data previously generated by our lab [9] to examine the enrichment

of TFs at defined windows in the 35 bp region surrounding the CTCF binding site (**Figure 1A**).

We focused on this region as we observed little enrichment of TF motifs beyond this point (**Figure**

S1A), although a slight increase could be detected further upstream between 50-75bps, that could

impact CTCF binding via a different mechanism. In **Figure 1A** the x-axis shows the gap/distance

in bps between the end of the TF motif and the start of the CTCF motif for upstream regions, or

the start of the TF motif and the end of the CTCF motif for downstream regions. CTCF has eleven

zinc fingers of which ZFs 3-7 make up the core binding domain that contacts the 12-15bp

consensus motif. It has previously been reported that at some sites an Upstream (U) CTCF motif,

can engage ZFs 9 to 11 and a less prevalent Downstream (D) motif can impair the binding of ZFs

1 and 2 to alter binding stability [11,17-19]. We therefore defined CTCF upstream and downstream

regions as the U motif window (U win) with binding ending at the 6 to 8 bp upstream gap, the U

spacer as binding ending at the 0 to 7 bp upstream gap, the D spacer as binding beginning at the

0 to 9 bp downstream gap, and the D motif window (D win) as binding beginning at the 10 to 11

downstream gap.

TF motifs have distinct enrichment profiles in mice and humans

Overall, in the mESCs, the percentage of TF motif enrichment (excluding U motifs and U like

motifs) at both upstream and downstream regions highlights that TF motifs are predominantly

associated with inaccessible (71%) rather than accessible (59%) sites (**Figure 1A**). A similar

analysis in human WTC11 cells revealed that in contrast to mESCs, TF motifs are more enriched

overall at accessible (61%) rather than inaccessible (41%) sites (**Figure 1A**). The differences

between human and mouse could reflect the fact that there are proportionately more accessible

CTCF binding sites in human compared to mouse (**Figure S1B** and **Table S1 A-B**).

There is redundancy in the TFs that can bind a particular window

In **Figure 1A**, TFs with distinct motifs (green stacked bars) can be distinguished from U and D

motifs (red and blue stacked bars, respectively). For each window, the TF motifs are shown in

Figures S2-S3 for mESCs and **S4-S6** for human WTC11 cells. In the U window, TFs that

resemble U motifs are also detected, but the latter are less enriched in human compared to mouse

cells (2.79% and 1.73% in WTC11 versus 25.5% and 11.85% in mESCs at accessible and

inaccessible sites, respectively) as shown in **Figures S2A,** and **S4A-B**. Although CTCF is highly

conserved between human and mouse, this result suggests that there are differences in their

binding sites, such that in human there is less reliance on binding of ZFs 9-11 to the U motif.

Motifs of other TFs enriched in the U and D windows and spacers and their preference for

enrichment at accessible versus inaccessible sites are shown in **Figure 1B, S2-S3** for mESCs

and **Figure 1B, S4-S6** for human WTC11 cells. The frequency with which each TF motif is found

is shown as a percentage next to the motif in the figure. The majority of CTCF sites are enriched

in one or more TF motifs, with each factor having a small contribution. It is pertinent that motifs in

each window and group of TFs show similarities within and between TF families such as the ETS

family in the U spacer, or the HOX family in the U window, indicating there is redundancy in the

factors that could bind, thereby providing a potential mechanism for cell type-specific regulation.

TF enrichment for each window in WT mESCs is shown in **Table S2A**, while in **Table S2B,** TF

enrichment in WTC11 cells and other human tissues is shown to highlight that TF motif enrichment

at CTCF sites is detected across different cell types, but as expected, motif enrichment of TFs

varies, which could reflect each cell's transcription program.

Figure 1: Bound CBSs are enriched in TF motifs within 35bps of the CTCF motif in mouse and human. A: Bar graphs showing the percentage of enriched TF motifs found using spacing motif enrichment analysis (SpaMo) at accessible and inaccessible sites in WT mESCs (top) and human iPSCs (bottom). The x-axis shows the gaps/distances in bp between the end of the TF motifs and the start of the CTCF motif for upstream gaps, or the start of the TF motif and the end of the CTCF motif for downstream gaps. For the U and D motif windows, the percentage of U and D motifs and TFs similar to the U and D motif are distinguished from other TF motifs (red or blue stacked bars, respectively). **B**: Motif logos of the enriched TF motifs found in the U motif window at accessible sites in mESCs (left) and human iPSCs (right). TF motifs ressembling the U motif and enrichment for the other windows are shown in **Figure S2-S6**,. **C-D**: CTCF binding profiles from ChIP-seq at accessible and inaccessible sites in mESCs (**C**) and human iPSCs (**D**) at CBSs containing at least one enriched TF motif within 35 bp upstream, downstream or both upstream and downstream compared to CBSs without any TF motifs.

The presence of TF motifs abutting CTCF binding sites is associated with stronger CTCF
binding

To determine whether enrichment of TF motifs is associated with changes in CTCF signal
strength, we compared the binding profiles of CTCF sites where we identified TF motifs at
upstream or downstream regions to sites without TF motifs. The presence of TF motifs at
accessible CTCF bound sites in the upstream and downstream regions, or both together were
associated with increases in CTCF signals (**Figure 1C**). The overall CTCF background profiles
were lower at inaccessible CTCF bound sites (**Figure 1C**) and the impact of TFs upstream or
downstream had a distinct effect compared to accessible sites. The differential effect sizes could
be explained by the predilection of TF classes found within specific windows as shown in **Figure**
1B, S2-S3 and **S5-S6**. Importantly, enrichment of TF motifs is associated with a similar increase
in CTCF binding strength in human WTC11 cells as observed in mESCs, suggesting that TFs
play an equivalent role in stabilizing CTCF in the two species (**Figure 1D**). It is of note that motifs
in up and downstream windows contribute in a distinct manner, in the two species, likely due to
their distinct enrichment profiles.

In sum, the data in **Figure 1** shows that in human and mouse the majority of CTCF bound sites
are enriched in one or more TF motifs. Their enrichment appears to be functionally important for
increasing the stability of CTCF binding. Motifs within each window and group of TFs show
similarities within and between TF families indicating redundancy in the cofactors that can bind.
However, they are degenerate, thereby providing a mechanism for cell type-specific differences
in regulation. These findings suggest that cell type-specific binding of CTCF might be influenced
by a large pool of TFs that reflect the transcriptional program of each cell type, and our analyses
provide a resource for identifying which TFs have the potential to influence CTCF binding and
chromatin organization in different cell types (**Tables S2A, B**).

Cobound TFs strengthen CTCF signals
We have shown that enrichment of TF motifs in the 35bps surrounding CTCF bound sites is
associated with stronger CTCF signal (**Figure 1C** and **D**). To confirm the relationship between TF
binding and CTCF signal strength, we analyzed ChIP-seq binding of individual TFs (KLF4, OCT4,
MYC, SOX2 and YY1) in mESCs using data downloaded from O'Dwyer *et al.*, and Laugwitz *et*
al., [20,21]. Our analysis, focused on CBSs with the consensus CTCF motif, highlighting sites at
which CTCF binds alone, CTCF and TFs cobind and TFs bind alone. Heatmaps and profiles for
accessible and inaccessible sites are shown in **Figure 2A** and **Figure S7**, respectively. The ChIP-

seq profiles validate our motif enrichment analysis, demonstrating that individual TFs stabilize

CTCF binding, with an increased effect at inaccessible sites for most TFs.

The heatmaps and profiles also demonstrate that CTCF free CBSs occupied by TFs are

associated with a stronger TF signal than at cobound sites for most TFs analyzed (**Figure 2A and**

Figure 2: Individual TFs strengthen CTCF signal and disrupt CTCF-dependent nucleosome phasing. A: Heatmaps and profiles of CTCF (FLAG) and 5 TFs (including 4 pioneer TFs, KLF4, OCT4, MYC and SOX2) showing the effect of (i) TF binding on CTCF signal strength and (ii) CTCF binding on TF signal strength at sites where they are co-bound (green) compared to CTCF only (blue) and TF only (red) sites at accessible regions. This analysis is restricted to CTCF binding sites with the consensus CTCF motif and profiles are oriented on C to N terminal CTCF binding sites. The profiles at inaccessible sites are shown in **Figure S7**. **B**: AlphaFold3 predictions of the CTCF, YY1 and DNA complex at three cobound sites. The structures of CTCF with (colored) or without YY1 (grey) were aligned and compared by calculated the RMSD (root mean standard deviation) for the CTCF-DNA complex and separately for CTCF and DNA. **C**: AlphaFold3 predictions of the YY1 and DNA complex at three YY1 only bound CBSs. YY1 is shown in blue. CTCF binding predictions are overlayed to visualize the position of YY1 relative to CTCF. Predictions for the other TFs are shown in **Figure S8**.

219 **S7**). The peak summits of the TFs are almost centered on the C to N terminal oriented CTCF

220 motif, suggesting a competition between CTCF and the TF. At cobound sites however, we

221 observe binding of the TFs at sites overlapping as well as on either side of the CBS.

AlphaFold3 predicts CTCF/TF binding conformation at cobound and CTCF free CBSs

To get a clearer picture of how TFs and CTCF could affect each other's binding, we turned to

AlphaFold3 and analyzed TF binding at cobound and TF bound CTCF free CBSs for each of the

five TFs, KLF4, OCT4, MYC, SOX2 and YY1. In **Figure 2B**, we show three examples of YY1

binding with CTCF at cobound sites. YY1 can bind in the Upstream spacer between ZFs 7 and 9

as well as the upstream or downstream regions. At these locations, there is little change in the

CTCF conformation as reflected by the small change in the route mean square deviation (RMSD),

however cobinding can affect DNA conformation which could result in changes in the CTCF-DNA

interaction.

When we examined AlphaFold3 predictions of YY1 binding (blue) at CTCF free CBSs, we

observed a direct competition between YY1 and CTCF at their predicted binding sites (**Figure**

2C), which could explain the absence of bound CTCF. Examples showing AlphaFold3 predictions

of how CTCF and the remaining TFs could affect each other's binding are shown in **Figure S8A-**

H. As with YY1, each TF binds upstream, downstream or within the U spacer when cobound with

CTCF, while at CTCF free CBSs, TFs tend to bind to sites overlapping the CBS, preventing CTCF

binding. We also show examples of OCT4/SOX2 and MYC/MAX cobinding with CTCF.

Interestingly, at the cobound locus shown in **Figure S8G,** the binding of SOX2 causes

considerable DNA bending, which has been previously described [22]. At this locus, the predicted

binding is in the Upstream spacer between ZFs 7 and 9 which is flexible [23], allowing ZFs 9-11 to

be properly positioned on the DNA despite the DNA bending, highlighting the importance of ZF8

as a spacer.

Allele-specific analysis reveals both the cooperativity and competitiveness of CTCF-TF

cobinding

To further validate the relationship between TFs and CTCF, we first downloaded the 145

ENCODE TF ChIP-seq sets available for the GM12878 cell line (**Table S3**) and found that 69%

of the CTCF peaks overlap at least one TF, confirming that co-binding of CTCF with another TF

is a frequent event. We then performed allele-specific analysis using ENCODE data from

GM12878 to identify SNPs that impact CTCF or TF motifs (and therefore their binding affinity)

(**Table S4**). In **Figure 3A, B** we show examples of heterozygous SNPs at two separate CTCF

binding sites. In both cases CTCF binding is reduced on the allele with the base change, coupled

with a loss of the cobound TF, overlapping RAD21 (a component of the cohesin complex) and to

a lesser extent altered ATAC-seq signals. Thus, CTCF binding can affect binding of the TF,

suggesting that in some cases CTCF binding might be required to open up the site for TF binding.

Of note, the SNP in **Figure 3A** is linked in GWAS to colon cancer and cardiometabolic traits.

Figure 3. **SNPs reveal both the cooperativity and competitiveness of CTCF-TF cobinding**. **A**: Map showing allele-specific binding (ASB) of CTCF due to a SNP disrupting its motif on the maternal allele in the GM12878 cell line. RAD21 and ARNT binding are also disrupted on the CTCF unbound allele. The CTCF consensus motif and the sequence at this locus are shown on the bottom track along with the orientation. The rectangles highlight the heterozygous SNP. For visualization purpose, the sequence tracks are not to scale. M, maternal; P, paternal. **B**: Map showing ASB of CTCF due to a SNP disrupting its motif on the maternal allele, which is associated with the disruption of BHLHE40, cohesin binding and chromatin accessibility on the same allele. **C**: Map showing ASB of IKZF2 due to a SNP disrupting its motif on the maternal allele. CTCF, cohesin binding and to a lesser extent the chromatin accessibility are also disrupted on the IKZF2 unbound allele. **D**: Map showing ASB of BHLHE40 due to a SNP disrupting its motif on the paternal allele. At this locus, the unbound BHLHE40 allele is associated with CTCF and cohesin binding as well as more open chromatin. None of these regions overlap with known imprinting regions and the observed ASBs are likely be cis-dependent ASBs rather than parent-of-origin dependent.

Conversely, SNPs that affect binding of TFs on one allele can impact binding of CTCF at

overlapping sites. In **Figure 3C**, the base change in the maternal haplotype impairs IKAROS

binding, with a concomitant reduction in CTCF, RAD21 and ATAC-seq signals. Thus, in this

setting binding of the TF is required for opening up the site to enable CTCF binding. In contrast,

the SNP disrupting the BHLHE40 motif and binding on the paternal allele at the *KIAA1671* locus

is permissive for CTCF binding and associated with concomitant binding of RAD21 and more

open chromatin (**Figure 3D**). This highlights an example of a TF that binds competitively with

CTCF and together with the data in **Figure 3B**, demonstrates that a single TF can have both a

stabilizing and destabilizing effect as shown in **Figure 2A**.

Overall, we identified 131 regions with allele-specific binding of CTCF that are associated with

allele-specific binding of a cofactor and where the presence of a SNP disrupting either the CTCF

or the cofactor motif provides a causal explanation. In half of these cases, the perturbation

introduced by the SNP indicates that TF co-binding impacts CTCF binding, while in the other half,

the effect is reversed, confirming the reciprocal nature of TF-CTCF co-binding (**Table S4**).

Additionally, we observed a destabilizing effect in 10% of the cases, suggesting that binding

cooperativity is more prevalent than competitiveness.

Transcriptional reprogramming contributes to cell state-specific CTCF binding

Our analyses demonstrate that cofactor binding can influence CTCF's binding stability, however

the impact of changes in transcriptional programs on CTCF binding have not been investigated.

To address this question, we compared TF expression levels at accessible and inaccessible sites

in human IPSCs. Overall, we found that TFs that bind motifs abutting CTCF were expressed at a

higher level at accessible sites (**Figure 4A**) and that those sites are more sensitive to changes in

TF expression levels compared to inaccessible sites, as shown by the significant difference in the

correlation slopes in **Figure 4B**.

To examine the effect of increasing transcriptional output on CTCF, we analyzed the effect of

variations in TF levels across 84 different ENCODE cell lines on their enrichment at CTCF binding

sites (**Table S2B**). We identified a subset of TFs whose expression levels showed a significant

positive or negative correlation with motif enrichment around bound CTCF sites (**Figure 4C and**

Table S5). This suggests that, in addition to context-specific effects, TFs might exert intrinsic

stabilizing or destabilizing influences on CTCF binding and that the enrichment or depletion of a

particular TF at CTCF sites in each cell type is dependent on its expression level, highlighting a

mechanism by which cell type-specific transcriptional programs can impact CTCF binding profiles.

Figure 4: The percentage of TF motifs within 35 bp of CTCF consensus motif correlates with their expression level. A: Bar graph showing an increase in expression of TFs with a higher contribution at accessible versus inaccessible sites ($\geq$2 fold differential contribution) in human iPSCs, after controlling for the differential effects of the gap window and direction relative to CTCF motif. The error bars correspond to the standard error computed using a multivariate linear model. **B**: Fitted correlation between overall TF expression and the percentage of CTCF sites found with motifs after controlling for the differential effect of windows and motif direction at accessible and inaccessible sites in human iPSCs. The shaded areas correspond to the 95%-confidence intervals. **C**: Examples of significant correlations between TF expression level and the percentage of CTCF peaks associated with the TF motif at accessible and inaccessible sites across human ENCODE cells and tissues. The shaded areas correspond to the 95%-confidence intervals.

CTCF mediates nucleosome repositioning around the CBS

298 To ask how CTCF binding affects nucleosome positioning in a context-specific manner, we
299 performed long-read Nano-NOMe-seq in mESCs, which allowed us to assess nucleosome
300 positioning using exogenous 5mC at HGC (GC excluding CGC) at the single molecule (SM) level.
301
302 To assess the effect of CTCF motif orientation and to discriminate molecules with occupied and
303 unoccupied CTCF binding sites, we restricted our analyses to CTCF peaks with GpC-containing
304 CTCF motifs after confirming that nucleosome occupancy at these sites were similar (**Figure
305 S9A**).
306
Unlike other similar analyses, we separated our profiles into occupied and unoccupied molecules.
At CTCF ChIP-seq peaks, a substantial fraction of molecules was temporarily unoccupied, which
is expected since DNA-protein interactions are dynamic. By comparing temporarily unoccupied
to occupied molecules at CTCF bound sites - thereby controlling for context-specific confounding
factors between bound and unbound CBSs - we confirmed previous findings demonstrating the
displacement and phasing of flanking nucleosomes after CTCF binding (**Figure 5A**).
Nucleosomes appeared less phased at inaccessible compared to accessible CTCF sites and
exhibited slightly shorter periodicity as well as a reduced CTCF flanking nucleosome-free region
(NFR) (**Figure 5A**).
316

Subdivision of CTCF binding sites into 14 clusters reveals a continuum of binding states
at aggregated and individual peaks

To further understand the binding dynamics of CTCF binding on chromatin, we performed
unsupervised clustering of the single molecules, incorporating data from all experimental
conditions in mESC across both accessible and inaccessible regions. This generated 14 clusters,
based on 50 bp-binned nucleosome occupancy within 1 kb of CBSs (**Figure 5B**). Differential
enrichment of each cluster in specific conditions and genomic contexts underscores the biological
relevance of each (**Figure S9B**). The UMAP projections of clustered SMs revealed that UMAP 1
correlates with global nucleosome occupancy whereas UMAP 2 partially captures differences in
nucleosome phasing patterns, despite only subtle variations in phasing scores (**Figure 5B,** right
and **S9C**).
328
Trajectory inference using Slingshot [24] helped define the relationships and dynamics among
clusters. For this analysis, we set cluster 1 as the starting point, as the molecules in this cluster -
along with clusters 2 and 3 (black clusters) - exhibit the characteristics of highly accessible, newly

replicated chromatin, previously shown to lack well-defined nucleosome organization [25].
Interestingly, cluster 1 displays a weak CTCF footprint with no surrounding nucleosome phasing,
while clusters 2 and 3 exhibit weak phasing either on the left or right of the CTCF footprint, a
distinction that could reflect the direction of replication. This observation suggests that CTCF can
act as a barrier during redeposition of nucleosomes on nascent chromatin and might help to
organize nucleosomes on newly replicated DNA.
338

Slingshot identified a bifurcation point corresponding to stable CTCF binding (red clusters 9-10)
with two divergent branches (green and blue lines) along the UMAP 2 axis, leading to violet
(clusters 4-8) and light blue (clusters 11-13) groups, respectively (**Figure 5B**). This branching
pattern may reflect alternative temporary nucleosome configurations associated with CTCF
binding. Indeed, the red clusters, enriched at strong CTCF peaks, show a pronounced CTCF
footprint and well-defined flanking NFR, consistent with CTCF-associated nucleosome
repositioning [10]. The **violet** clusters (4-8) show a CTCF footprint on the left, middle or right side
of a nucleosome, supporting findings that CTCF can bind its motifs regardless of nucleosome
position [26]. In these clusters, the close proximity of nucleosomes to the CTCF motif and the
presence of partial NFRs suggest a less stable CTCF binding – supported by the enrichment in
weak CTCF peaks - or intermediate states prior to, or during the action of nucleosome remodelers
by CTCF [10]. The **light blue** clusters (11-13) maintain nucleosomes in an organized, phased
manner but with a narrower NFR around unbound CTCF sites (lacking a footprint), possibly
reflecting temporally unbound CTCF. A third branch (orange) along UMAP1 leads to the **dark**
blue cluster (14), which is enriched in the CTCF-depleted condition following IAA treatment. This
might represent sites at which CTCF rebinds weakly or does not return at all, leading to disruption
of nucleosome phasing. Finally, the purple path reflects the cyclical process between nascent
and mature chromatin configurations.
357

Importantly when we focus on individual CTCF peaks we can detect all fourteen clusters, as
shown for three representative examples (**Figure 5C**). These analyses are important for
demonstrating that the clusters are not simply an aggregation of data from different binding sites
enriched for one cluster or another across the genome. Instead, they represent a continuum of
different states of CTCF binding at each peak. The proportion of each cluster is shown for twelve
individual binding sites (**Figure 5D**).

Figure 5

Figure 5: Subdivision of CTCF binding sites into 14 clusters reveals distinct phases of binding. A: Profiles and single molecule heatmaps showing the nucleosome occupancy (NO) and CTCF binding footprinting from nano-NOMe-seq for accessible and inaccessible CTCF sites. Both occupied (black) and temporarily occupied (blue) molecules at CTCF bound sites (ChIP-seq peaks) are shown. The data are centered on the 15-bp oriented consensus motif. The y-axis of the profiles shows the percentage of exogenous 5mC detected in an HGC context. High exogenous methylation reflects TF or nucleosome free chromatin accessible to the GpC methyltransferase, while lower methylation reflects occupied regions. The heatmaps show the nucleosome occupancy and CTCF footprinting at the single molecule (SM) level for a random sample of 5000 molecules. Black (or blue) and white dots represent occupied and free GpCs, respectively. The red and orange lines correspond to the stable position of the first upstream and downstream nucleosome flanking the CTCF motif. **B**: SM heatmaps of 5000 molecules per condition and genomic context showing the nucleosome occupancy and CTCF binding footprinting at accessible and inaccessible CTCF sites by cluster (left), and UMAP of the SMs colored by cluster and phasing scores (right). The arrowed lines show the SM trajectories identified by Slingshot. **C**: Example of heatmaps showing the nucleosome occupancy and CTCF binding footprinting at three individual CTCF peaks. **D**: Bar graph showing examples of cluster distribution at individual CTCF peaks. **E**: SM heatmaps and bar graphs showing the representative proportion of each cluster at accessible and inaccessible sites.

Analysis of clustering at accessible and inaccessible sites highlights that the narrowing of the NFR

surrounding CTCF seen in **Figure 5A,** is reflected by a decrease in the red clusters (strongly

bound CTCF) and an increase in Cluster 14 (unbound or weakly bound CTCF with no nucleosome

repositioning (**Figure 5E**). Additionally, a reduced ability to reposition nucleosomes is reflected

by a more robust nucleosome presence on either side or overlapping the CBS (violet clusters). In

sum, single molecule nano-NOMe-seq data reveals the dynamics of CTCF binding states at both

aggregated and individual peak level, highlighting CTCF's ability to reposition nucleosomes to

create a NFR surrounding its binding sites and to act as a barrier that facilitates nucleosome

phasing, both of which are impaired at inaccessible CBSs.

Unoccupied CBSs are bound by nucleosomes or other factors

We next analyzed CBSs that are not bound by CTCF in mESCs. Most of the molecules at those

sites are occupied (59% and 72% respectively at accessible and inaccessible sites), either by a

nucleosome or a TF as suggested by the persistence of a narrow footprint (**Figure 6A**).

Nucleosomes are still phased around the CBS but there is a noticeable increase in the proportion

of violet and light blue clusters and a decrease in the red cluster compared to CTCF bound sites

(**Figure 5E**). These data indicate that binding of TFs to CTCF free sites can still phase

nucleosomes, but these are not associated with a NFR flanking the CBS. As before, we see a

more robust nucleosome presence at inaccessible sites, consistent with less TF binding at these

regions.

CTCF is required to reposition and phase nucleosomes

To confirm the effect of CTCF on nucleosome phasing and repositioning, we used an mESC

CTCF auxin inducible degron [9]. Loss of CTCF protein at 3, 18 and 48 hours resulted in an

Figure 6

Figure 6: *SCC1 mediates CTCF associated nucleosome repositioning and phasing.* **A:** Profiles and SM heatmaps stratified by cluster showing the nucleosome occupancy and CTCF binding footprinting for CTCF-free and motif-containing accessible and inaccessible CBSs. CTCF-free binding sites are defined as FIMO-detected CTCF motifs without CTCF or SMC3 ChIP-seq peaks. **B:** Profiles (for untreated, 3, 18 and 24 hours) and heatmaps of untreated and 18 hour IAA degraded mESCs at accessible and inaccessible sites. The heatmaps for 3 and 48 hours are shown in **Figure S10**. **C**: bar graphs showing the distribution of each cluster at CTCF peaks, CBS and after CTCF degradation (IAA). **D**: CTCF ChIP-seq profiles in untreated human Hela cells and after SCC1 degradation at 3 and 12 hours. **E:** Profiles of nucleosome occupancy in untreated human Hela cells and after SCC1 degradation (3 and 12 hours) at accessible and inaccessible sites. **F**: SM heatmaps and bar graphs showing the distribution of each cluster in untreated Hela cells and after IAA degradation.

increasing reduction in nucleosome repositioning at CBSs (profiles in **Figure 6B**). The more

granular analyses of the heatmaps and barplot show an increased proportion of cluster14 (dark

blue), with a less marked footprint at the CBS over time (**Figure 6B, C** and **Figure S10A**). As

expected, there is a notable decrease in the representation of the strong, red clustered CTCF

binding sites and at the violet clusters, we detect a weaker CTCF footprint and a more robust

nucleosome presence (**Figure S10A**). Interestingly, after CTCF degradation, CBSs show similar

proportions of violet, red and light blue clusters to CTCF free sites (**Figure 4A**). Given these

similarities, we speculate that the weak footprint at CTCF degraded sites could be due to the

presence of incoming, or previously bound TFs, although we cannot rule out that some of this

footprint is due to leftover, undegraded or weak CTCF [27].

SCC1 mediates CTCF associated nucleosome repositioning and phasing

Since CTCF degradation leads to loss of both CTCF and cohesin at CBSs (**Figure S10B**), the

CTCF degron experiment could not be used to determine which factor is responsible for mediating

the changes we observed after CTCF degradation. To address this question, we used an auxin

inducible degron of SCC1 (a component of the cohesin complex) in the human Hela cell line [28].

Since cohesin is required for mitosis, we assessed early degradation time points, 3 hours and 12

hours, where we could still collect live cells. Alongside nano-NOMe-seq we performed CTCF

ChIP-seq. CTCF binding profiles remained the same as untreated cells after IAA treatment at 3

hours and 12 hours (**Figure 4D**). In contrast, the nucleosome profiles for occupied and

unoccupied sites show that although there is little change in nucleosome clustering at the 3-hour

time point, despite unchanged CTCF profiles, we could detect a reduction in nucleosome

repositioning and phasing at CBSs by 12 hours (**Figure 4E**), similar to that observed after CTCF

degradation (**Figure 4B**). Looking at the heatmaps and barplots (**Figure 4F)** at both accessible

and inaccessible sites there is a noteable reduction in the strongly bound CTCF fraction (red

clusters) and an increase at weakly bound CTCF sites characterized by disorganized

nucleosomes surrounding the CBS (dark blue clusters). Violet clusters are also reduced but more

Figure 7: The CTCF flanking NFR is associated with insulation. A. Clustered heatmaps of SMs stratified by chromatin accessibility and CTCF signal strength. **B:** Profiles comparing the nucleosome occupancy and insulation score between weak versus strong CTCF sites stratified by chromatin accessibility. The averaged insulation score generated at 10 kb resolution from HIC data is shown for each genomic context. **C**: Heatmaps of SMs stratified by clusters at accessible and inaccessible CTCF peaks in the absence or presence of TF motifs. The bar graphs show the distribution of each cluster by genomic context. **D**: Profiles comparing the nucleosome occupancy and insulation score between CTCF-alone and cobound sites stratified by chromatin accessibility.

so at inaccessible regions, and at these sites there is a stronger nucleosome presence surrounding the CBS, obscuring the CTCF footprint in most places. That the 12-hour time point effect occurs in the background of unaltered CTCF binding, indicates that cohesin plays a dominant role in nucleosome repositioning and phasing. However, CTCF alone may exert an

425 additional weaker effect, as the red clusters are not reduced to the same extent as when CTCF
426 is depleted and there is a persistent NFR in the dark blue cluster. We speculate that the delay in
427 loss of nucleosome repositioning at 3 hours could be because nucleosome repositioning needs
428 to be reset during S phase [29].
429
430 In sum, we found that CTCF free CBS, CTCF and SCC1 degradation all give rise to similar
431 clustering profiles with a decreasing NFR over time, indicating that a reduction or absence of
432 CTCF binding can allow binding of TFs that can phase, but are unable to reposition surrounding
433 nucleosomes. Thus, cohesin and loop extrusion could mediate nucleosome positioning and
434 phasing by creating or stabilizing barrier effects at CTCF anchors.
435

CTCF signal strength contributes more to nucleosome repositioning and insulation than
accessibility
When we compare the clustering of strong versus weak CTCF signals and inaccessible versus
accessible sites, we observed that signal strength gives rise to a bigger difference in the
proportion of red clusters (strong CTCF binding in a NFR), compared to accessibility (**Figure 7A**).
Moreover, the differential effects of CTCF signal strength were similar at both accessible and
inaccessible sites, suggesting that CTCF signal strength contributes to nucleosome repositioning
independent of accessibility. We previously performed a stratified analysis showing that CTCF
signal strength, which was associated with cohesin stalling, also contributes more to insulation
than accessibility [9]. Here we highlight that a reduction in nucleosome repositioning at weak versus
strong sites is associated with a corresponding reduction in chromatin insulation compared to that
at inaccessible versus accessible sites (**Figure 7B**).
448

Surrounding motifs affect CTCF-mediated nucleosome repositioning and insulation
independent of signal strength
The presence of TF motifs are associated with differential effects on CTCF binding (**Figure 1C**).
Although CTCF signal strength is increased at accessible sites, we observe a minimal change in
nucleosome repositioning and chromatin insulation (**Figure 7C, D**). The same is true at
inaccessible sites, except when TF motifs are enriched on either side of CTCF. Despite a slight
increase in CTCF signal strength at these sites, both nucleosome repositioning and chromatin
insulation are reduced. Together, these data indicate that CTCF and cohesin signal strength can
be uncoupled from the size of the NFR surrounding CTCF, while the association between
nucleosome repositioning and insulation is maintained. This finding highlights that nucleosome

repositioning is a more reliable predictor of chromatin insulation than CTCF signal strength. The

data also demonstrate that TF binding can interfere with nucleosome repositioning and chromatin

insulation and we speculate that the presence of TFs could be a way to fine tune the effect of

CTCF signal strength on these two processes.

Discussion

Although it is known that a subset of CBSs is altered in a state-specific manner, the mechanisms

underlying this control have not been fully elucidated. Here we used CTCF ChIP-seq in mESCs

combined with CTCF ChIP-seq from ENCODE data in a wide range of human cell types to (i)

characterize accessible versus inaccessible sites, and (ii) determine what factors influence CTCF

binding stability and function within these two chromatin environments. Although CTCF is highly

conserved in mice and humans, we found different features associated with their respective

binding sites. Specifically, mouse cells have proportionately more inaccessible binding sites

compared to human cells. In parallel, TF motifs are enriched at inaccessible sites in mouse, which

we have previously shown generally have weaker CTCF signals and reduced cohesin overlap

compared to accessible sites [9]. Since TF cobinding stabilizes CTCF, we speculate that their

enrichment at inaccessible sites in mouse, restores the balance of strong versus weak

CTCF/cohesin signals in the two species, compensating for the relative reduction of accessible

sites in mouse.

CTCF has been shown by us and others to act as a pioneer factor, binding inaccessible as well

as accessible chromatin [9,30]. To follow up on previously published studies hinting at a role for TFs

in cell state specific binding of CTCF, we performed a comprehensive analysis of enrichment of

TF motifs in the 35bp region of the CBS in mESCs as well as 85 ENCODE cell lines. We identified

a wide range of motifs corresponding to expressed cell state-specific factors that could perform

this function. In human and mouse, the majority of CTCF bound sites are enriched in motifs

located within 35 bps of CTCF. Within a given window, enriched TF motifs show a strong similarity,

suggesting redundancy in the regulation of CTCF binding. Furthermore, motifs are degenerate

which could allow for differential effects of TFs from the same class or motif cluster. Given the

redundancy between factors that can bind, we speculate that at a single CTCF binding site, the

presence of distinct TFs in different cell types could fine-tune CTCF's stability and its ability to

block cohesin in a lineage or state specific manner. Indeed, we demonstrate that CTCF signals

are strengthened at sites enriched with TF motifs and by analyzing the effect of a handful of

cofactor TFs on CTCF binding profiles, we validate their stabilizing effect.

The significance of these findings is underscored by our analysis of human tissues from
ENCODE, all of which have unique transcriptional programs with individual TFs expressed at
different levels. Importantly, increases in the expression level of a TF can both increase or
decrease the enrichment of its motif at CTCF bound sites, depending on whether the TF has a
stabilizing or destabilizing effect. The relationship between TFs and CTCF were validated by allele
specific analysis which demonstrate that the two factors have a reciprocal effect on each other's
binding such that SNPs affecting either the binding of CTCF or the TF will impact enrichment of
the other factor, altering cohesin overlap and accessibility. SNPs in CBSs or TF motifs highlight
both the cooperativity and competitiveness of CTCF-TF cobinding. Our finding that cell type-
specific TFs can have a negative effect on CTCF binding addresses another long-standing
question about why many accessible CBSs remain unbound. Here we show that binding of other
factors can compete with CTCF to prevent its occupation.

AlphaFold3 predictions demonstrate that TF binding to CTCF free CBSs, overlaps the CBS,
providing an explanation for their inability to cobind. In contrast, at cobound sites, binding of the
TF occurs in the upstream or downstream regions, sites that do not interfere with CTCF's
occupation. Although AlphaFold3 cannot predict the impact of cobinding on CTCF stability, our
ChIP-seq analyses confirm that this leads to a stronger CTCF signal, despite TF-mediated
changes in DNA torsion and bending. However, binding of cofactors can alter nucleosome
phasing, and we speculate that changes in DNA torsion and bending could be responsible for
disrupting repositioning, and the outcome could be TF specific as previously shown [22]. Cobinding
of YY1 and CTCF occur at a subset of PAX5 binding sites in the *immunoglobulin heavy chain*
(Igh) locus [31]. This is of interest, because we previously showed that both PAX5 [32,33] and YY1 [34]
are important for mediating chromatin looping and locus contraction in early B cell development
to induce long range V-to-DJ rearrangements which is important for the generation of antibody
diversity. In the absence of either factor, locus contraction and rearrangement are impaired,
consistent with the idea that cofactors are important for CTCF mediated function.

Use of long read, nano-NOME-seq single molecule analysis allowed us to interrogate the impact
of CTCF binding on nucleosome repositioning at a resolution other assays such as MNase could
not achieve. By subsetting the data, we were able to show a continuum of CTCF binding states,
starting with binding to highly accessible chromatin devoid of nucleosomes, which has been
shown to reflect newly replicated chromatin [25]. Binding of CTCF occurs early and helps to organize

nucleosomes around the binding site leading to ordered nucleosome phasing and repositioning,

which likely requires recruitment of chromatin remodellers, such as members of the mSWI/SNF

family [35]. The clustering shows that CTCF can bind with either its N or C terminal abutting a

nucleosome or less frequently, overlapping a nucleosome. This is in contrast to the findings of

Pugacheva *et al*., who posit that CTCF binds only at N terminal nucleosome entry sites [35].

At CTCF free sites, CBSs are occupied either by a nucleosome or TFs that can phase

nucleosomes, but cannot reposition them. The effect of the TF on nucleosomes is more dynamic

as reflected by less stable nucleosome positioning around CTCF, in line with the low residence

time of of these factors compared to CTCF. Furthermore, there is a more robust nucleosome

presence at inaccessible sites, consistent with less TF binding at these regions. Interestingly, we

detect similar overall binding states when we degrade CTCF, suggesting that in the absence of

CTCF, TFs can come in and bind at CBSs. This data further highlights that CTCF and TFs can

bind competitively at CBSs, when their binding sites directly overlap.

While CTCF has been shown to act as a barrier that promotes nucleosome phasing and

positioning at its binding sites [36], SCC1 depletion data suggests that cohesin plays a critical and

dominant role in nucleosome repositioning and phasing at CTCF sites. Notably, the loss of

phasing and repositioning observed following SCC1 degradation - despite unchanged CTCF

binding strength - closely recapitulates the nucleosome configuration seen upon combined loss

of CTCF and SMC3 binding at CBSs after CTCF depletion. This finding suggests that cohesin

and loop extrusion might be responsible for the local nucleosome organization observed at CTCF

anchors.

Clustering demonstrates that the proportion of chromatin with repositioned nucleosomes depends

on accessibility as well as CTCF signal strength, both of which likely influence the recruitment of

nucleosome remodellers. However our studies highlight that signal strength has a greater impact

on nucleosome repositioning, and this is also reflected in a bigger impact on chromatin insulation

than accessibility, confirming our previous findings [9]. However, we also show that CTCF signal

strength and nucleosome repositioning can be uncoupled in the presence of TFs, while the

association between nucleosome repositioning and insulation is still maintained. Thus,

nucleosome repositioning is a more reliable predictor of insulation than CTCF signal strength and

we propose that use of single molecule nucleosome occupancy could be a powerful tool to predict

effects on insulation.

Here we have shown that enrichment of a wide range of TF motifs stabilize CTCF binding and
confirmed this by analyzing the ChIP-seq of a handful of factors at CBSs. Together, these studies
define a mechanism by which transcriptional reprogramming during development and in disease
specific states can determine cell type-specific CTCF profiles, linked to local and long-range
chromatin organization.

METHODS

Cell line generation and culture

The construction of the vector for cloning transgenic, doxycycline-inducible expression of WT and
mutant mouse *Ctcf* cDNA and gene targeting in mouse embryonic stem cells E14Tg2a (karyotype
19, XY; 129/Ola isogenic background) were performed as described in Do *et al.,* [9]. Mouse
embryonic stem cells and all clones derived from these were cultured under feeder-free conditions
in 0.1% gelatin (Sigma ES-006-B) coated dishes (Falcon, 353003) as described in Do *et al.,* [9].
Briefly, the cells were grown in DMEM (Thermo Fisher, 11965-118) supplemented with 15% fetal
bovine serum (Thermo Fisher, SH30071.03), 100 U/ml penicillin - 100 µg/ml streptomycin (Sigma,
P4458), 1 X GlutaMax supplement (Thermo Fisher, 35050-061), 1 mM sodium pyruvate (Thermo
Fisher, 11360-070), 1 X MEM non-essential amino-acids (Thermo Fisher, 11140-50), 50 µM b-
mercaptoethanol (Sigma, 38171), 10^4 U/ml leukemia inhibitory factor (Millipore, ESG1107), 3 µM
CHIR99021 (Sigma, SML1046) and 1 µM MEK inhibitor PD0325901 (Sigma, PZ0162).
SCC1-mEGFP-AID HeLa cell line in which SCC1 is tagged with an auxin induced degron was
previously generated by Wutz et al.,[28] and was kindly provided by Dr. Peters. The cells were
grown in DMEM supplemented with 10% DBS (Thermo Fisher, 10371029), 0.2 M glutamine
(Thermo Fisher, 25030081), penicillin/streptomycin and 1X non-essential amino acids (Thermo
Fisher, 11140050)

Induction of auxin inducible degradation of CTCF, SCC1 and doxycycline induced
expression

For degradation of endogenous CTCF in mESCs, the auxin-inducible degron was induced by
adding 500 µM indole-3-acetic acid (IAA, the chemical analog of auxin) (Sigma, I5148) to the
media. Expression of WT CTCF transgene was achieved by the addition of doxycycline (Dox, 1
µg/ml) (Sigma, D9891) to the media. The cells were treated with IAA and/or Dox for 3h, 18h or 2
days. In Hela, SCC1 were degraded by addition of 500 µM IAA for 3h or 12h.

Nano-NOMe-seq sample and library preparation

Nuclei extractraction, GpC methyl-transferase treatment and DNA extraction

After 2 days of IAA and Dox treatment on WT mESCs, 1 million cells were collected and washed

with cold PBS at 300g at 4°C. As described in Battaglia *et al*., [37], nuclei were extracted using cold

lysis buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 0.5 mM spermidine, 1.5 mM MgCl2, 0.1 mM

EDTA, 0.25% IGEPAL CA-630) for 3-6 minutes after checking nuclei under a light microscope.

The sample was centrifuged for 5 minutes at 800x g at 4°C, washed with 1 ml ice-cold nuclei

wash buffer (10 mM Tris-HCl pH 7.4, 50 mM NaCl, 0.5 mM spermidine, 1.5 mM MgCl2, 0.1 mM

EDTA), and centrifuged again for 5 minutes at 800x g at 4°C. The nuclei were resuspended in

564 μL 1X GpC MTase reaction buffer (NEB, B0227) and equilibrated to 37°C. This was followed

by 6 minutes of enzyme treatment (300 μL 1 M sucrose, 34 μL 10X GpC methyltransferase

reaction buffer, 3 μL 32 mM S-Adenosylmethionine (SAM) (NEB, B9003), 100 μL GpC

methyltransferase (M.CviPI) (NEB, M0227L)), followed by another 6 minutes of enzyme treatment

(50 μL GpC methyltransferase (M.CviPI), 3 μL 32 mM S-Adenosylmethionine (SAM)). An equal

volume of pre-warmed 2X stop buffer (20 mM Tris-HCl pH 7.4, 600 mM NaCl, 10 mM EDTA, 1%

SDS) was added and mixed well. Proteinase K (200 μg/mL) was added to the GpC methylation

reaction and incubated for 4 hours at 56°C, followed by RNase A (100 μg/mL) for 30 minutes at

37°C, followed by an additional round of proteinase K (200 μg/mL) for 30 minutes at 37°C.

HMW DNA was extracted using phenol-chloroform and precipitated by isopropanol. Briefly, an

equal volume of phenol-chloroform was added to the samples and mixed vigorously. The samples

were centrifuged at 13,000 rpm for 10 minutes at room temperature. The aqueous phase was

transferred to a clean tube, and 0.7-1 times the volume of isopropanol was added, mixed well,

and incubated at room temperature for 30 minutes. The samples were then centrifuged at 13,000

rpm for 10 minutes at 4°C. The DNA pellet was washed with 1 ml pre-chilled 70% ethanol. After

removing all the ethanol, the DNA was dissolved in EB buffer and incubated at 37°C for 30

minutes. The final DNA concentration was quantified using Nanodrop 2000, and molecular weight

was measured by genomic DNA screentape (Agilent, 5067-5365).

Nano-NOMe-seq library preparation

For Nano-NOMe-seq, the DNA fragment was sheared to an average size of 20kb with the Covaris

G-Tube (#520079) in an Eppendorf 5415D tabletop microcentrifuge following the manufacture's

protocol. Sample size was confirmed on genomic screentape with the Agilent Tapestation system.

The sheared DNA (1.5ug) was then input into the SQK-LSK114 library prep from Oxford

Nanopore to create the Nanopore library. Components were used from the LSK114 kit along with
several buffers and enzymes from New England Biolab (NEB) following the LSK114 protocol from
Oxford Nanopore. Following library prep, the sample concentration was verified on the Qubit
system using the dsDNA High Sensitivity Kit. The concentration and basepair size were used to
calculate Femtomolar concentration for loading. The sample was resuspended with library loading
beads and loaded at 150-200 femtomolar concentration onto the Promethion R10.4.1 flow cell
according to the LSK protocol. Sample were sequenced on the R10.4.1 Promethion flow cell using
the Oxford Nanopore Promethion P24 system with a sequencing run time of 72 hrs. Standard
settings were applied during the run setup, special settings were applied and fragments less than
1kb rejected and methylation detection was set for both 5mC and 5hmC along with the super
accurate basecalling feature turned on.
640
ChIpmentation
ChIPmentation was performed in the Hela cell lines as described in Do *et al.* [9]. Briefly, 10 millions
cells were double cross linked using 25mM EGS (ethylene glycol bis(succinimidyl succinate);
Thermofisher #21565), followed by addition of 1% formaldehyde (Tousimis #1008A). Chromatin
was sheared using a bioruptor (Diagenode) for 15 minutes. Mouse anti-CTCF (Active Motif,
61311) and Rabbit IgG (abcam, ab37415) antibodies were used for immunoprecipitation.
Tagmentation were performed using 1 µl Tagment DNA Enzyme from the Tagment DNA Enzyme
and Buffer Kit (Illumina #20034198). Purified DNA (20 µl) was amplified and barcoded using
indexed and non-indexed primers and NEBNext High-Fidelity 2X PCR Master Mix (NEB M0541).
DNA was purified using Agencourt AMPure XP beads (Beckman, A63881) to remove fragments
larger than 700 bp as well as the primer dimers. Library quality and quantity were estimated using
Tapestation (Agilent High Sensitivity D1000 ScreenTape #5067-5584 and High Sensitivity D1000
reagents #5067-5585) and quantified by Qubit (Life Technologies Qubit™ 1X dsDNA High
Sensitivity (HS) #Q33230). Libraries were then sequenced with the Novaseq6000 Illumina
technology according to the standard protocols and with around 200 million 150bp paired-end
total per sample.
657
Datasets and Data Processing
ChIP-seq, RNA-seq and ATAC-seq dataset
The CTCF, SMC3 ChIP-seq, RNA-seq and ATAC-seq data for WT mESCs were previously
generated and processed as described in Do *et al.*, [9] using the Seq-N-Slide pipeline [38]. Hela
ChIPmentation data were processed using the same pipeline.

For ENCODE human samples, processed data for samples having CTCF ChIP-seq, RNA-seq

and ATAC-seq performed on the same biosample were downloaded from ENCODE. The

accession numbers of the dataset used in this work are listed in **Table S1A, S1B, S3**. CTCF

peaks were scanned for CTCF consensus motif (JASPAR MA0139.2) [39] using FIMO [40] with a p-

value cutoff of 5e10^{-5}. Only samples with at least 20% of motif containing CTCF peaks were kept

for downstream analyses. Accessible CTCF peaks were identified by overlapping CTCF with

ATAC peaks using bedtools [41].

For the GM12878 cell line, we downloaded an additional set of 145 ENCODE TF ChIP-seq narrow

peak files to assess the percentage of CTCF binding sites with TF co-binding (the accession

numbers are listed in **Table S3**. To analyze allele-specific binding the BAM files for CTCF, RAD21,

TFs and ATAC-seq were also downloaded (the accession numbers are listed in **Table S3**). The

phased genotyping data of GM12878 was retrieved from the Platinium Genomes project [42]

(accession number PRJEB3246).

To validate the effect of TF co-binding, we downloaded KLF4, OCT4, SOX2, MYC, YY1 and input

ChiP-seq data (fastq files) generated in mESCs from GEO [20] [21]. The data were processed using

the Seq-N-Slide pipeline [38] as described in Do *et al.*, [9]. Briefly, the reads were aligned to mm10

genome with Bowtie2 [43] and narrow peaks were called using MACS2 [44] in pair-end mode and with

the input as control.

The CTCF-oriented heatmaps and profiles were generated using the computeMatrix scale-

regions function from deeptools [45] using a 15 bp bin and a region body length of 15 bp which

corresponds to the length of CTCF core motif.

Hi-C data.

The HIC data were previously generated using Arima HiC kit (A510008) and processed using

HiC-Pro and HiCExplorer as described in Do *et al.*, [9]. The HIC matrices were generated at a 10

kb resolution. The insulation scores were computed using GENOVA [46] and the CTCF-oriented

profiles generated using deeptools [45] with a 1 kb bin and a region body length.

Nano-NOMe-seq processing

Base modification calling were performed on pod5 files using dorado basecaller [47] using the "--

min-qscore 10 sup,5mC_5hmC" option, followed by alignment using dorado aligner against the

mm10 reference genome. Reads with less than 10 MAPQ score were filtered out using samtools

[48]. Methylation in HGC context was called using dorado pileup using the "--combine-strands"
option. Single molecule (SM) methylation was extracted using modkit extract with the "--read-
calls-path" option [49]. GpC overlapping the ENCODE blacklisted regions [50] were excluded as well
as sites with more than 100x coverage.

CTCF-oriented GpC methylation profiles were generated using deeptools and single molecule
heatmaps generated using R. Single molecules with occupied or unoccupied CTCF were defined
as molecule containing a unmethylated or methylated GpC in the motif, respectively. No bias was
observed in the nucleosome positioning of molecules with and without GpC containing motifs
(**Figure S4A**). SM Heatmaps were created in R at the single molecule and GpC level using the
GpC methylation status called by modkit extract after centering the reads on the middle of CTCF
consensus motif.

Downstream Analyses

Motif logos and Motif spacing enrichment analysis

Motif logos of the aggregated sequences centered on CTCF motif detected by FIMO [40] were
created using the R package, ggseqlogo [51]. Significantly enriched spacings of TF motifs in the
vicinity of CTCF bound motifs were identified using SpaMo [52]. The motif positional weight matrices
(PWM) were downloaded from JASPAR 2024 (CORE vertebrates non-redundant dataset) [39] . The
PWM for the U and D motifs were derived from Nakahashi *et al.,* [17]. The analyses were performed
using the 500 bp window centered on each bound CTCF motif stratified by chromatin accessibility.
TF motif clustering and trees were performed using the R package, universalmotif [53] for each
window defined as follow: upstream spacer, U window and further upstream, 0 to 5bp, 6-8 bp and
more than 9bp upstream, respectively; downstream spacer, D window and further downstream,
0 to 9 bp, 10-12 bp and more than 13 bp downstream respectively.

Association between TF motif enrichment and TF expression

To ask whether TFs more enriched at accessible sites compared to inaccessible sites were more
expressed in human iPSCs, accessible enriched TFs were defined as TFs with a 2-fold increase
in the percentage of motif found within 35 bp of the CTCF motif identified by the SpaMo analysis.
A multivariate linear regression using a backward-stepwise selection was then performed to test
the association between TF expression level (log2 TPM) and chromatin accessibility controlled
for the epistatic effect of the motif orientation and window of enrichment (relative to the CTCF
motif). The adjusted expression level by accessibility was estimated by computing the model

margins and the p-value by performing the pairwise comparisons of predictive margins using
STATA v17.

To test the overall correlation between TF expression and motif enrichment, we performed an
aggregated analysis across ENCODE human samples (**Table S1A**). The expression level of TFs
and percentage of motif were summed by sample, accessibility, window of enrichment and motif
orientation. A multivariate linear regression using a backward-stepwise selection was then
performed to test the correlation between the aggregated percentage of enriched motifs and TF
expression level (log2 TPM) controlled for the epistatic effect of chromatin accessibility, the motif
orientation and window of enrichment (relative to the CTCF motif). The differential effect of these
covariates on the TF enrichment was estimated using the adjusted slope (interaction term
between the covariate and TF expression) by computing the model margins and the pairwise
comparisons of predictive margins using STATA v17.
Bivariate linear regressions between individual TF motif enrichment and expression level were
performed using R.

AlphaFold3 prediction
For each transcription factor (TF) — KLF4, OCT4, SOX2, MYC, and YY1 — we randomly selected
loci for each category of "CTCF-only," "CTCF-TF cobound," and "TF-only" sites. The DNA
sequences (positive and negative strands) of these loci, centered on the CTCF or TF motif and
including 35 bp flanking sequences, were extracted using Bedtools [41].
TF and CTCF ZF1-11 amino acid sequences were retrieved from UniProt [54]: Q61164 for CTCF,
Q00899 for YY1, Q60793 for KLF4, P20263 for OCT4, P48432 for SOX2, P01108 for MYC, and
P28574 for MAX. The CTCF ZF1-11 positions were obtained from Yang *et al.* [23].
AlphaFold3 predictions [55] were carried out using the DNA sequences, the CTCF sequence, with
or without the TF sequence. The predicted protein-DNA structures were visualized in PyMOL
3.10, and the root mean square deviation (RMSD) between the structures was calculated using
PyMOL 3.10.

Allele-specific binding analyses
Allele-specific binding (ASB) was performed by separating the ChIP-seq or ATAC-seq reads
overlapping phased heterozygous SNPs in the GM12878 cell line using R. Allele-specific BAM
and bigwig files were then generated using samtools [48] and bamCoverage [45]. To identify ASB

SNPs disrupting the TF motif, allele-specific motif affinity was computed using the R package,

atSNP [56].

Nano-NOMe-seq single molecules clustering, trajectory and phasing score

For each condition (untreated, IAA) and genomic context (based on accessibility, CTCF signal

strength, presence of TF motifs), 5000 molecules were randomly sampled. The level of 5mC

methylation in GpC context was binned by 50 bp for each molecule. The missing values were

imputed using k-Nearest Neighbors before performing k-means clustering. UMAP was applied

using the uwot R package with n_neighbors = 15 and min_dist = 0.2 to project the nucleosome

occupancy profiles into two dimensions. For SM trajectories, a principal curve was first fitted on

the UMAP-embedded molecules using the princurve R package [57]. All molecules were then

projected onto this curve to obtain pseudotime estimate and trajectory inferred using the slingshot

R package [24], specifying cluster 1 as the root. Nucleosome phasing scores were calculated by

applying a fast Fourier transform (FFT) to the binarized methylation patterns of each read. Two

separate phasing scores were computed for accessible and inaccessible sites, targeting

periodicities of 190 bp and 170 bp, respectively. For each, the power spectrum was computed

and the phasing score was defined as the proportion of signal intensity within a narrow frequency

band around the expected nucleosome spacing.

Data availability

All raw and processed sequencing data files are deposited at NCBI's Gene Expression

Omnibus (GEO) and will be available to public on publication of the manuscript.

Code availability

No custom code/software was used in the study. The publicly available software used is indicated

in the methods section.

Acknowledgements

This work was supported by 1R35GM122515 (J.S) and NIH P01CA229086 (J.S). GC were

supported by a fellowship from the NCC.

The authors thank Skok lab members for helpful scientific discussions, New York University

School of Medicine High Performance Computing Facility (HPCF) for computing technical

support, Adriana Heguy and the Genome Technology Center (GTC) core for sequencing efforts,

NYU Flow Cytometry and Cell Sorting Center for FACS analysis and sorting. GTC is a shared
resource partially supported by the Cancer Center Support Grant P30CA016087 at the Laura and
Isaac Perlmutter Cancer Center.
801
802 The authors thank Sofia Battaglia from the Bernstein's lab for her advice and protocols for the
803 Nano-NOMe-seq library preparation.
804
805 **Author contributions**
806 These studies were designed by Jane Skok and Catherine Do. The Nano-NOME-seq experiments
807 were performed by Guimei Jiang and Paul Zappile under the supervision of Adriana Heguy. The
808 analysis was performed by Catherine Do The paper was written by Jane Skok and Catherine Do.
809
810 **Declaration of interests**
811 The authors declare no competing interests.
812
813 **SUPPLEMENTARY INFORMATION**
814
Supplemental Tables
Table S1. Tab A: Accession numbers of the CTCF ChIP-seq, ATAC-seq and RNA-seq for human
ENCODE samples used in the motif analyses. Only samples with data in all three assays and
with at least 35% of CTCF motif-containing peaks were included. The percentage of CTCF motif
found using FIMO are listed. **Tab B**: Number and percentage of accessible CTCF peaks in human
and mouse including the CTCF ChIP-seq and ATAC-seq accession numbers for the mouse
ENCODE samples. This data was used in **Figure S1B.** This table relates to **Figure 1.**
822
823 **Table S2. Tab A:** Proportions and counts of the sequences with enriched TF motifs within 35 bp
824 of the CTCF consensus motif at accessible and inaccessible CBSs in WT mESCs. **Tab B:**
825 Proportions and counts of the sequences with enriched TF motifs within 35 bp of the CTCF
826 consensus motif at accessible and inaccessible binding sites in human ENCODE samples with at
827 least 20% of CTCF motif-containing peaks. This table relates to **Figure 1**.
828
829 **Table S3.** ENCODE accession numbers of the TF ChIP-seq narrow peak and BAM files in
830 GM12878 to assess the overall percentage of cobound CTCF and TF sites. This table relates to
831 **Figure 3**.

Table S4. Allele-specific binding of CTCF associated with allele-specific of TF with SNPs
disrupting either CTCF or the TF motifs. This tables relates to **Figure 3**.

Table S5. Significant correlations between the percentage of TF motifs within 35 bp of CTCF
consensus motif and their expression level in ENCODE human samples. This relates to **Figure
4**.

Supplemental Figures

844
845

Figure S1. A: Cumulative percentage of CTCF peaks containing a TF motif within 250 bp at accessible and inaccessible sites. The dashed lines correspond to 35 bp upstream or downstream of the CTCF motif. This figure relates to **Figure 1**. **B:** Dotplot showing the higher percentage of accessible CTCF bound sites in human versus mouse. The red line represents the median. The p-value has been computed using a t-test. This figure is related to **Figure 1.**

Figure S2. A: Motif logos of the enriched TF motifs found in the U motif window at accessible and inaccessible sites in WT mESCs. **B**: Motif logos of the enriched TF motifs found in the U spacer window at accessible and inaccessible sites in WT mESCs. This window is enriched in ETS-family motifs. This figure relates to **Figure 1**.

851
852
853
854

Figure S3. A: Motif logos of the enriched TF motifs found in the D window at accessible sites in WT mESCs. No motifs were significantly enriched at inaccessible sites. **B**: Motif logos of the enriched TF motifs found in the downstream spacer window at accessible and inaccessible sites in WT mESCs. This figure relates to **Figure 1**.

856
857
858
859

Figure S4

Figure S4. A: Motif logos of the enriched TF motifs found in the U motif window at accessible sites in human iPSCs (WCT11)**. B:** Motif logos of the enriched TF motifs found in the U motif window and resembling the U motif at inaccessible sites in human iPSCs (WCT11)**. C**: Motif logos of the enriched TF motifs found in the U motif window and distinct from the U motif at inaccessible sites in human iPSCs (WCT11)**.** This figure relates to **Figure 1**.

867
868

Figure S5. A: Motif logos of the enriched TF motifs found in the D motif window at accessible and inaccessible sites in human iPSCs (WCT11). **B**: Motif logos of the enriched TF motifs found in the upstream spacer at accessible and inaccessible sites in human iPSCs (WCT11)**.** This window is enriched in ETS-family motifs. This figure relates to **Figure 1**.

Figure S6. A: Motif logos of the enriched TF motifs found in the downstream spacer at accessible and inaccessible (**B**) sites in human iPSCs (WCT11)**.** This figure relates to **Figure 1**.

Figure S7: Individual TFs strengthen CTCF signal. Profiles of CTCF (FLAG) and 5 TFs (including 4 pioneer TFs, KLF4, OCT4, MYC and SOX2) showing the effect of TF binding on CTCF signal strength (top) and TF signal strength (bottom) at inaccessible sites where they are co-bound (green) compared to CTCF only (blue) and TF only sites (red). This analysis is restricted to CTCF binding sites with the consensus CTCF motif. The graphs are oriented on the C to N terminal CTCF core motif. This figure relates to **Figure 2**.

Figure S8

A Cobound CBSs: CTCF (colored) with KLF4 (red)
   Potential effect of loss of KLF4 on CTCF binding (grey)

   chr10:128536413-128536483
   RMSD CTCF-DNA: 1.8
   RMSD CTCF: 0.7
   RMSD DNA: 1.8

B KLF4 only CBSs: KLF4 only (blue)

   chr2:104069990-104070075

C Cobound CBSs: CTCF (colored) with MYC-MAX (red)
   Potential effect of loss of MYC-MAX on CTCF binding (grey)

   chr12:111732652-111732722
   RMSD CTCF-DNA: 2.7
   RMSD CTCF: 2.7
   RMSD DNA: 2.0

D MYC only CBSs: MYC-MAX only (blue)

   chr2:143062998-143063083

E Cobound CBSs: CTCF (colored) with OCT4 (red)
   Potential effect of loss of OCT4 on CTCF binding (grey)

   chr11:45008266-5008336
   RMSD CTCF-DNA: 2.2
   RMSD CTCF: 2
   RMSD DNA: 2.3

F OCT4 only CBSs: OCT4 only (blue)

   chr5:120649102-120649187

G Cobound CBSs: CTCF (colored) with SOX2-OCT4 (red)
   Potential effect of loss of SOX2-OCT4 on CTCF binding (grey)

   chr7:30634790-30634860
   RMSD CTCF-DNA: 9.3
   RMSD CTCF: 7.2
   RMSD DNA: 21.8

H SOX2-OCT4 only CBSs: SOX2-OCT4 only (blue)

   chr13:37514368-37514453

Figure S8: TFs can cobind or compete with CTCF at CBSs. AlphaFold3 predictions at cobound sites for KLF4 (**A**), MYC-MAX (**C**), OCT4 (**E**) and SOX2-OCT4 (**G**). To assess the effect of the cobound TF, the structures of CTCF with (colored) or without TF (grey) were aligned and the RMSD (root mean standard deviation) calculated for the CTCF-DNA complex and separately for CTCF and DNA (top). At cobound sites, TFs tend to bind outside CBSs or in the U spacer (between ZF7 and ZF9). SOX2 binding can induce a major bend in the DNA (**G**). AlphaFold3 predictions at TF only bound CBSs for KLF4 (**B**), MYC-MAX (**D**), OCT4 (without SOX2) (**F**) and SOX2-OCT4 (most of the SOX2 peaks overlap with OCT4) (**H**). TFs are shown in in blue. CTCF binding predictions are overlayed to visualize the position of the TF relative to CTCF. At those sites, the TFs tend to bind at CTCF core or extended binding sites and as a result they directly compete with its binding. This figure relates to **Figure 2**.

Figure S9

Figure S9: A: Profiles showing similar nucleosome occupancy phasing of occupied SM for GpC-containing CBSs compared to non GpC-containing CBSs. **B**: Enrichment of each SM cluster for the condition and genomic context under study. The odd ratios and p-values were computed using logistic regressions. **C:** UMAP of SMs colored by the molecule overall nucleosome occupancy. This figure relates to **Figure 5**.

884
885

Figure S10: A: Heatmaps showing the effect of CTCF degradation on nucleosome phasing after 3h and 48h of IAA treatment in mESCs. **B:** Profiles of SMC3 at lost CTCF peaks after CTCF degradation (48h IAA) at accessible and inaccessible sites in mESCs. This figure relates to **Figure 6**.

886
887

889
References

1   Davidson, I. F. *et al.* DNA loop extrusion by human cohesin. *Science* **366**, 1338-1345
(2019). https://doi.org/10.1126/science.aaz3418
2   Fudenberg, G. *et al.* Formation of Chromosomal Domains by Loop Extrusion. *Cell Rep* **15**,
2038-2049 (2016). https://doi.org/10.1016/j.celrep.2016.04.085
3   Dehingia, B., Milewska, M., Janowski, M. & Pekowska, A. CTCF shapes chromatin
structure and gene expression in health and disease. *EMBO Rep* **23**, e55146 (2022).
https://doi.org/10.15252/embr.202255146
4   Guo, C. *et al.* CTCF-binding elements mediate control of V(D)J recombination. *Nature*
477, 424-430 (2011). https://doi.org/10.1038/nature10495
5   Hnisz, D. *et al.* Activation of proto-oncogenes by disruption of chromosome
neighborhoods. *Science* **351**, 1454-1458 (2016). https://doi.org/10.1126/science.aad9024
6   Narendra, V. *et al.* Transcription. CTCF establishes discrete functional chromatin domains
at the Hox clusters during differentiation. *Science* **347**, 1017-1021 (2015).
https://doi.org/10.1126/science.1262088
7   Xiang, Y., Zhou, X., Hewitt, S. L., Skok, J. A. & Garrard, W. T. A multifunctional element in
the mouse Igkappa locus that specifies repertoire and Ig loci subnuclear location. *J*
Immunol **186**, 5356-5366 (2011). https://doi.org/10.4049/jimmunol.1003794
8   Hansen, A. S. CTCF as a boundary factor for cohesin-mediated loop extrusion: evidence
for a multi-step mechanism. *Nucleus* **11**, 132-148 (2020).
https://doi.org/10.1080/19491034.2020.1782024
9   Do, C. *et al.* Binding domain mutations provide insight into CTCF's relationship with
chromatin and its contribution to gene regulation. *Cell Genom* **5**, 100813 (2025).
https://doi.org/10.1016/j.xgen.2025.100813
10   Iurlaro, M. *et al.* Systematic assessment of ISWI subunits shows that NURF creates local
accessibility for CTCF. *Nat Genet* **56**, 1203-1212 (2024). https://doi.org/10.1038/s41588-
024-01767-x
11   Essien, K. *et al.* CTCF binding site classes exhibit distinct evolutionary, genomic,
epigenomic and transcriptomic features. *Genome Biol* **10**, R131 (2009).
https://doi.org/10.1186/gb-2009-10-11-r131
12   Hu, G. *et al.* Systematic screening of CTCF binding partners identifies that BHLHE40
regulates CTCF genome-wide distribution and long-range chromatin interactions. *Nucleic*
Acids Res **48**, 9606-9620 (2020). https://doi.org/10.1093/nar/gkaa705
13   Jerkovic, I. *et al.* Genome-Wide Binding of Posterior HOXA/D Transcription Factors
Reveals Subgrouping and Association with CTCF. *PLoS Genet* **13**, e1006567 (2017).
https://doi.org/10.1371/journal.pgen.1006567
14   Liu, Y. *et al.* CTCF coordinates cell fate specification via orchestrating regulatory hubs with
pioneer transcription factors. *Cell Rep* **42**, 113259 (2023).
https://doi.org/10.1016/j.celrep.2023.113259
15   Ross-Innes, C. S., Brown, G. D. & Carroll, J. S. A co-ordinated interaction between CTCF
and ER in breast cancer cells. *BMC Genomics* **12**, 593 (2011).
https://doi.org/10.1186/1471-2164-12-593
16   Sonmezer, C. *et al.* Molecular Co-occupancy Identifies Transcription Factor Binding
Cooperativity In Vivo. *Mol Cell* **81**, 255-267 e256 (2021).
https://doi.org/10.1016/j.molcel.2020.11.015
17   Nakahashi, H. *et al.* A genome-wide map of CTCF multivalency redefines the CTCF code.
Cell Rep **3**, 1678-1689 (2013). https://doi.org/10.1016/j.celrep.2013.04.024

18   Plasschaert, R. N. *et al.* CTCF binding site sequence differences are associated with
unique regulatory and functional trends during embryonic stem cell differentiation. *Nucleic*
Acids Res **42**, 774-789 (2014). https://doi.org/10.1093/nar/gkt910
19   Soochit, W. *et al.* CTCF chromatin residence time controls three-dimensional genome
organization, gene expression and DNA methylation in pluripotent cells. *Nat Cell Biol* **23**,
881-893 (2021). https://doi.org/10.1038/s41556-021-00722-w
20   O'Dwyer, M. R. *et al.* Nucleosome fibre topology guides transcription factor binding to
enhancers. *Nature* **638**, 251-260 (2025). https://doi.org/10.1038/s41586-024-08333-9
21   Laugwitz, L. *et al.* ZSCAN10 deficiency causes a neurodevelopmental disorder with
characteristic   oto-facial   malformations.   *Brain*   **147**,   2471-2482   (2024).
https://doi.org/10.1093/brain/awae058
22   Malaga Gadea, F. C. & Nikolova, E. N. Structural Plasticity of Pioneer Factor Sox2 and
DNA Bendability Modulate Nucleosome Engagement and Sox2-Oct4 Synergism. *J Mol*
Biol **435**, 167916 (2023). https://doi.org/10.1016/j.jmb.2022.167916
23   Yang, J. *et al.* Structures of CTCF-DNA complexes including all 11 zinc fingers. *Nucleic*
Acids Res **51**, 8447-8462 (2023). https://doi.org/10.1093/nar/gkad594
24   Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genomics* **19**, 477 (2018). https://doi.org/10.1186/s12864-018-
4772-0
25   Ostrowski, M. S. *et al.* The single-molecule accessibility landscape of newly replicated
mammalian   chromatin.   *Cell*   **188**,   237-252   e219   (2025).
https://doi.org/10.1016/j.cell.2024.10.039
26   Grand, R. S. *et al.* Genome access is transcription factor-specific and defined by
nucleosome   position.   *Mol   Cell*   **84**,   3455-3468   e3456   (2024).
https://doi.org/10.1016/j.molcel.2024.08.009
27   Luan, J. *et al.* Distinct properties and functions of CTCF revealed by a rapidly inducible
degron system. *Cell Rep* **34**, 108783 (2021). https://doi.org/10.1016/j.celrep.2021.108783
28   Wutz, G. *et al.* Topologically associating domains and chromatin loops depend on cohesin
and are regulated by CTCF, WAPL, and PDS5 proteins. *EMBO J* **36**, 3573-3599 (2017).
https://doi.org/10.15252/embj.201798004
29   Owens, N. *et al.* CTCF confers local nucleosome resiliency after DNA replication and
during mitosis. *Elife* **8** (2019). https://doi.org/10.7554/eLife.47898
30   Pop, R. T. *et al.* Identification of mammalian transcription factors that bind to inaccessible
chromatin. *Nucleic Acids Res* **51**, 8480-8495 (2023). https://doi.org/10.1093/nar/gkad614
31   Ebert, A. *et al.* The distal V(H) gene cluster of the Igh locus contains distinct regulatory
elements with Pax5 transcription factor-dependent activity in pro-B cells. *Immunity* **34**,
175-187 (2011). https://doi.org/10.1016/j.immuni.2011.02.005
32   Fuxa, M. *et al.* Pax5 induces V-to-DJ rearrangements and locus contraction of the
immunoglobulin   heavy-chain   gene.   *Genes   Dev*   **18**,   411-422   (2004).
https://doi.org/10.1101/gad.291504
33   Roldan, E. *et al.* Locus 'decontraction' and centromeric recruitment contribute to allelic
exclusion of the immunoglobulin heavy-chain gene. *Nat Immunol* **6**, 31-41 (2005).
https://doi.org/10.1038/ni1150
34   Liu, H. *et al.* Yin Yang 1 is a critical regulator of B-cell development. *Genes Dev* **21**, 1179-
1189 (2007). https://doi.org/10.1101/gad.1529307
35   Otto, J. E. *et al.* Structural and functional properties of mSWI/SNF chromatin remodeling
complexes revealed through single-cell perturbation screens. *Mol Cell* **83**, 1350-1367
e1357 (2023). https://doi.org/10.1016/j.molcel.2023.03.013
36   Wiechens, N. *et al.* The Chromatin Remodelling Enzymes SNF2H and SNF2L Position
Nucleosomes adjacent to CTCF and Other Transcription Factors. *PLoS Genet* **12**,
e1005940 (2016). https://doi.org/10.1371/journal.pgen.1005940

37 Battaglia, S. *et al.* Long-range phasing of dynamic, tissue-specific and allele-specific
regulatory elements. *Nat Genet* **54**, 1504-1513 (2022). https://doi.org/10.1038/s41588-
022-01188-8
38 Dolgalev, I. Seq-N-Slide [Computer software]. (2022).
https://doi.org/https://doi.org/10.5281/zenodo.5550459
39 Rauluseviciute, I. *et al.* JASPAR 2024: 20th anniversary of the open-access database of
transcription factor binding profiles. *Nucleic Acids Res* **52**, D174-D182 (2024).
https://doi.org/10.1093/nar/gkad1059
40 Bailey, T. L. *et al.* MEME SUITE: tools for motif discovery and searching. *Nucleic Acids*
Res **37**, W202-208 (2009). https://doi.org/10.1093/nar/gkp335
41 Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841-842 (2010). https://doi.org/10.1093/bioinformatics/btq033
42 Eberle, M. A. *et al.* A reference data set of 5.4 million phased human variants validated by
genetic inheritance from sequencing a three-generation 17-member pedigree. *Genome*
Res **27**, 157-164 (2017). https://doi.org/10.1101/gr.210500.116
43 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9, 357-359 (2012). https://doi.org/10.1038/nmeth.1923
44 Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).
https://doi.org/10.1186/gb-2008-9-9-r137
45 Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A. & Manke, T. deepTools: a flexible platform
for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187-191 (2014).
https://doi.org/10.1093/nar/gku365
46 van der Weide, R. H. *et al.* Hi-C analyses with GENOVA: a case study with cohesin
variants. *NAR Genom Bioinform* **3**, lqab040 (2021).
https://doi.org/10.1093/nargab/lqab040
47 nanoporetech. Dorado. *https://github.com/nanoporetech/dorado* (2024).
48 Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-
2079 (2009). https://doi.org/10.1093/bioinformatics/btp352
49 nanoporetech. Modkit. *https://github.com/nanoporetech/modkit* (2023).
50 Amemiya, H. M., Kundaje, A. & Boyle, A. P. The ENCODE Blacklist: Identification of
Problematic Regions of the Genome. *Sci Rep* **9**, 9354 (2019).
https://doi.org/10.1038/s41598-019-45839-z
51 Wagih, O. ggseqlogo: a versatile R package for drawing sequence logos. *Bioinformatics*
33, 3645-3647 (2017). https://doi.org/10.1093/bioinformatics/btx469
52 Whitington, T., Frith, M. C., Johnson, J. & Bailey, T. L. Inferring transcription factor
complexes from ChIP-seq data. *Nucleic Acids Res* **39**, e98 (2011).
https://doi.org/10.1093/nar/gkr341
53 Tremblay, B. J. universalmotif: An R package for biological motif
analysis. *The Journal of Open Source Software* **9(100):7012** (2024).
https://doi.org/10.21105/joss.07012
54 Apweiler, R. *et al.* UniProt: the Universal Protein knowledgebase. *Nucleic Acids Res* **32**,
D115-119 (2004). https://doi.org/10.1093/nar/gkh131
55 Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493-500 (2024). https://doi.org/10.1038/s41586-024-07487-w
56 Zuo, C., Shin, S. & Keles, S. atSNP: transcription factor binding affinity testing for
regulatory SNP detection. *Bioinformatics* **31**, 3353-3355 (2015).
https://doi.org/10.1093/bioinformatics/btv328
57 Hastie, T. & Stuetzle, W. Principal Curves. *ournal of the American Statistical Association,*
84(406), *502–516.* (1989).
https://doi.org/https://doi.org/10.1080/01621459.1989.10478797
1039