Strip cropping shows promising increases in ground beetle community diversity compared to monocultures

Luuk Croijmans[*a,b], Fogelina Cuperus[*b,c], Dirk F. van Apeldoorn[b,c], Felix J.J.A. Bianchi[b], Walter A.H. Rossing[b], Erik H. Poelman[a]

    a) Laboratory of Entomology, Wageningen University & Research, Wageningen, the Netherlands

    b) Farming Systems Ecology, Wageningen University & Research, Wageningen, the Netherlands

    c) Field Crops, Wageningen University & Research, Lelystad, the Netherlands

[*] = These authors contributed equally to this work.

Email:

luuk.croijmans@wur.nl; felix.bianchi@wur.nl (corresponding authors)

Keywords: Carabidae, intercropping, organic agriculture, crop diversification, agroecosystem, community ecology

Abstract

Global biodiversity is declining at an unprecedented rate, with agriculture as one of the major

drivers. There is mounting evidence that intercropping can increase insect biodiversity while

maintaining or increasing yield. Yet, intercropping is often considered impractical for

mechanized farming systems. Strip cropping is a type of intercropping that is compatible with

standard farm machinery and has been pioneered by Dutch farmers since 2014. Here, we present

ground beetle data from four organically managed experimental farms across four years. Ground

beetles are sensitive to changes in habitats and disturbances, and hold keystone positions in

agroecosystem food webs. We show that strip cropping systems can enhance ground beetle

biodiversity, while other studies showed that these increases have been achieved without

incurring major yield loss. Strip cropped fields had on average 15% more ground beetle species

and 30% more individuals than monocultural fields. The higher ground beetle richness in strip

crops was explained by the merger of crop-related ground beetle communities, rather than by

ground beetle species unique to strip cropping systems. The increase in field-level beetle species

richness in organic agriculture through strip cropping approached increases found for other

readily deployed biodiversity conservation methods, like shifting from conventional to organic

agriculture (+19% - +23%). This indicates that strip cropping is a potentially useful tool

supporting ground beetle biodiversity in agricultural fields without compromising food

production.

Introduction

Insects account for 80% of the animal species in the world and, therefore, the recently reported

unprecedented rate of insect decline is cause for alarm about the state of biodiversity on Earth

(Dirzo et al., 2014; Hallmann et al., 2017; Van Klink et al., 2020). An array of biodiversity

metrics provide strong evidence for declines of especially terrestrial insects across all continents

(Van Klink et al., 2020). This includes declines in abundance of both common and rare species

(Hallmann et al., 2017; Pilotto et al., 2020; Seibold et al., 2019; van Klink et al., 2024) and in

total species richness and changes in assembly composition (Blowes et al., 2022; Wagner et al.,

2021). Insects are essential for crop production through their role in decomposition, pollination,

pest control and sustaining food webs. Therefore, erosion of insect communities can have

potentially devastating effects on ecosystem functioning, provision of ecosystem services and

ultimately on human civilization (Dirzo et al., 2014). The main drivers of insect biodiversity

decline are habitat loss due to conversion to agriculture, pollution, invasive species and climate

change (Díaz et al., 2019; Müller et al., 2024; Wagner et al., 2021). Strategies for biodiversity

conservation in conjunction with adequate food production require understanding of how

biodiversity responds to agricultural management (Cozim-Melges et al., 2024; Mupepele et al.,

2021; Saunders, 2020). Ideally, sustainable agricultural practices retain yield and enhance

biodiversity, preventing the need to convert natural habitats to agricultural land to maintain food

production (Tscharntke et al., 2021).

Increasing crop heterogeneity can facilitate biodiversity conservation in highly productive

agricultural landscapes without compromising yield (Martin-Guay et al., 2018; Sirami et al.,

2019). Crop diversification can enhance niche complementarity by creating heterogeneous

habitats and increasing availability and diversity of resources (Lichtenberg et al., 2017;

Tamburini et al., 2020). A promising crop diversification strategy is strip cropping, where crops

are grown in alternating strips, wide enough for using standard agricultural machines yet narrow

enough to facilitate ecological interactions among crops (Ditzler et al., 2021). Crops that are

grown in strips may benefit from increased resource use efficiency, and the suppression of pests

and diseases (Croijmans et al., 2024; Karssemeijer et al., 2024; Rakotomalala et al., 2023)

without major yield compromises (Campanelli et al., 2023; Juventia and van Apeldoorn, 2024;

Van Oort et al., 2020). Growing multiple crops on a field may foster a larger diversity of

organisms than monocultures through greater plant species richness that cascades into richer

herbivore and predator communities (Crutsinger et al., 2006; Cuperus et al., 2023). Moreover, the

expected increase in available and potentially complementary niches within the agricultural field

due to higher spatial diversity in crops can result in admixture of communities related to

individual crops (Hummel et al., 2012), or the creation of completely new communities by

enhanced richness of agriculture-related species, and/or the occurrence of species rarely found in

agricultural fields (Rischen et al., 2021). So far, it is not well understood if and how insect

communities respond to strip cropping across distinct crops.

Here, we present data of four years of pitfall trapping of ground beetles at several moments

during the growing season in 14 crops at four organic experimental farms across the Netherlands

where strip cropping was compared to monocultures. Ground beetle communities are sensitive to

changes in farming practices and are frequently used to examine agricultural sustainability

(Holland and Luff, 2000; Makwela et al., 2023; Turin, 2022). Furthermore, ground beetle species

are important for maintaining ecological functions as they comprise scavengers and predators of

(weed) seeds, detritivores (e.g., collembolas and earthworms) and herbivores (e.g., aphids and

caterpillars). We first examine whether strip cropping fields have greater ground beetle activity

density, species richness, evenness and diversity than monocultural fields. We also test for

twelve abundant ground beetle genera whether their activity density is higher in strip cropping

than in monocultures. Lastly, we evaluate whether ground beetle community changes are caused

by admixture of communities, whether these assemblages promote species associated with

agricultural or with natural ecosystems, and whether they contain rare species.

☐

Results

A total of 48,108 ground beetles belonging to 71 species were caught using pitfall traps over four

years at four different organically managed experimental farms in The Netherlands: 40,153 at

Almere; 3,777 at Lelystad; 1,126 at Valthermond; and 3,052 at Wageningen (Table S1, S2).

Strip cropping enhances ground beetle richness

Strip cropping fields had on average 15% higher ground beetle taxonomic richness than

monoculture fields after rarefaction to the number of samples of the least-sampled crop

configuration ($\beta_0 = 0.151$, SE = 0.044, p < 0.001; Fig. 1c,d). However, strip cropping fields did

not harbour more species than monocultural fields with the highest ground beetle richness ($\beta_0 = -$

0.008, SE = 0.037, p = 0.821; Fig. 1d). The difference in field-level taxonomic richness could not

be explained by an increase in the number of ground beetle species per crop in strip cropping

compared to monocultures. At crop-level, the 5% increase in ground beetle taxonomic richness in

strip cropping was not statistically significant (Fig. 1f). Similarly, crop-level absolute evenness,

inverse Simpson index and Shannon entropy did not differ significantly among crop

configurations (Fig. 1h-j, S1c,d, S2). The effect of crop configuration on crop-level taxonomic

richness was variable and was not associated with location or crop species. The effect ranged

from 56% more species in potato in monoculture at Wageningen in 2022, to 136% more species

in barley in strip cropping at Wageningen in 2020 (Fig. 1k).

Strip cropping enhances ground beetle activity density

Ground beetle activity density was on average 30% higher in strip cropping fields than in

monoculture fields ($\beta_0 = 0.303$, SE = 0.121, p = 0.012; Fig. 1e), based on rarefaction. However,

there was no significant difference in activity density between the strip cropping fields and

monocultures that harboured the richest beetle communities ($\beta_0 = -0.110$, SE = 0.088, p = 0.215).

Crop-level activity density of ground beetles was not affected by crop configuration (Fig. 1g,

S1a,b).

Crop configuration alters abundance of abundant genera

We tested how crop configuration affected the abundance of twelve abundant ground beetle

genera (Table S3). We analysed this separately per location as some genera only occurred at

specific locations. Four genera were more abundant in strip cropped fields in at least one location

(*Anchomenus, Bembidion, Harpalus* and *Nebria*), whereas four genera were more abundant in

monocultures in at least one location (*Amara, Calathus, Pterostichus* and *Trechus*) (Fig 2). The

other four common genera (*Blemus, Clivina, Loricera* and *Poecilus*) were not significantly

influenced by crop configuration. Furthermore, no ground beetle genus showed significantly

contrasting responses to crop configuration among locations.

Crop configuration alters ground beetle community composition

Ground beetle communities were significantly influenced by crop configuration, but the effects

were highly dependent on the specific context created by the combination of location, year and

crop (Fig. S3-S8, Table 1, S4). The context-dependency of configuration effects on ground beetle

communities is illustrated in redundancy analyses of ground beetle assemblages per crop

combination at Wageningen in 2021 and 2022 (Fig. S9). Here, we found distinct ground beetle

communities among crop configurations for pumpkin, barley and potato in 2021, and for cabbage

and oat in 2022. In the other cases the difference between crop configurations was not significant

(Fig. S9, Table S5). Moreover, in all crop combinations except for potato-grass in 2021, the

difference in ground beetle communities between the monocultures of the constituent crops were

significant, while this was never the case for ground beetle communities of crops in strip

cropping (Table S5). This indicates that strip cropping might lead to overlapping crop-related

communities. However, these results could be spatially autocorrelated as samples from different

crops were in closer proximity of each other in strip cropping than among monocultures (Fig

S10d).

Crop configuration does not increase the number of rare ground beetle species

Among the 461 year series, we only found two rare species (following waarneming.nl): one

individual of *Microlestes minutulus*, and five individuals of *Harpalus signaticornis*. This latter

species was found most often in a wheat monoculture, but this was likely due to a failed crop

which created a very open habitat. *Harpalus signaticornis* is known to inhabit recently disturbed,

dry, open habitats with limited vegetation (Turin et al., 2012), a situation similar to this sparsely-

covered monoculture. All other species were common or relatively common species (Table S2).

Furthermore, most ground beetle species were either ruderal habitat specialists or eurytopic

species that occur in many different habitats (Table S2).

Discussion

It is well established that different crop types have distinct ground beetle communities (Eyre et

al., 2009; Holland and Luff, 2000) and that increasing habitat diversity by including multiple

crops in a field can enhance ground beetle diversity (Cuperus et al., 2024; Puliga et al., 2023).

Our study shows that strip cropping increased field-level ground beetle richness by 15%.

However, ground beetle communities within the same crop in a strip or in monoculture were

mostly similar. This indicates that the 15% increase in richness at the field level can be mostly

attributed by the higher number of crops in strip cropped fields that harboured crop-related

ground beetle communities, and that there was only limited mixing of ground beetle species

among crops in strip crops. This is in line with earlier findings that ground beetle movement is

reduced by crop edges (Allema et al., 2014; Anderson et al., 2024). Further research on

movement behaviours of ground beetles at crop edges might help explain how ground beetles

distribute themselves within a strip cropping field, and whether they utilize the different

resources provided by a more diverse cropping system.

The 30% higher ground beetle activity density in strip cropping fields compared to monoculture

fields may be explained by a more stable and diverse habitat with refuges and alternative

resources in strip crops (Ratnadass et al., 2012). Crop diversification enhances prey biomass for

ground beetles (Lichtenberg et al., 2017) and increases weed seed richness as compared to

monocultures (Ditzler et al., 2023), both of which reduce bottom-up control by increased food

provision (Carbonne et al., 2022). Alternatively, the increase in activity density might be caused

by higher movement of ground beetles, which in turn can be the consequence of food starvation

(Wallin and Ekbom, 1994). Indeed, several papers show reduced abundances of herbivorous

insects in strip- and intercropping (Alarcón☐Segura et al., 2022; Cuperus et al., 2023;

Rakotomalala et al., 2023), which may be potential prey for ground beetles (Turin, 2000).

Furthermore, strip cropping systems may support high herbivore reproduction in combination

with a high predation pressure resulting in herbivore populations dominated by early life stages

(Karssemeijer et al., 2024). Therefore, strip cropping may favour ground beetles that predate on,

for instance, herbivore eggs, such as *Bembidion spp.* and *Anchomenus dorsalis*, which were more

abundant in strip cropping (Finch, 1996), and disadvantaging ground beetles that feed on larger

prey, such as *Pterostichus melanarius,* which was less abundant in strip cropping. Multi-taxa

evaluation of future strip cropping studies will be a valuable approach to increase understanding

of biomass flows and trophic interactions within diverse agricultural systems.

The 15% increase in ground beetle species richness through strip cropping is mostly caused by an

increase in species (relatively) common to agricultural fields, rather than an increase in rare

species or species otherwise common in other habitat types. A keystone species here might be

Pterostichus melanarius, the most common *Pterostichus* species in our samples. *Pterostichus*

melanarius can be very dominant in pitfall traps, as we regularly found hundreds of individuals

per trap. This relatively large ground beetle might compete with or predate on other ground

beetle species (Roubinet et al., 2018). *Pterostichus melanarius* is especially well-adapted to

highly intensive agriculture (Turin, 2022, 2000), which explains why it was more abundant in

monocultures previously (Ditzler et al., 2021). In turn, the reduced *P. melanarius* populations in

strip cropping might allow other species to persist, such as *Harpalus spp.* and *Anchomenus*

dorsalis. Therefore, strip cropping might facilitate ground beetle biodiversity by reducing the

dominance of a common, competitive species.

Biodiversity gains in our study did in most cases not coincide with productivity losses (Table

S6). Earlier studies on crop yield in the strip cropping fields in Lelystad and Wageningen during

the years of pitfall trapping show a yield decrease when strip cropping cabbage (Carrillo-Reche

et al., 2023) and wheat (Ditzler et al., 2023), whereas potato yield was unaffected by crop

configuration (Ditzler et al., 2023). In Almere bean and parsnip yields were higher in strip

cropping, whereas oat and onion yield were lower (Juventia and van Apeldoorn, 2024). In

Valthermond crop productivity was similar for monocultures and strip cropping (Table S6).

Therefore, biodiversity gains do not compromise agricultural production.

We show that changing crop configuration from monoculture to strip cropping, on average,

enhances ground beetle richness by 15% and activity density by 30% within agricultural fields.

These results show that strip cropping can lead to increases in biodiversity that approach those

achieved by shifting from conventional to organic farming practices (+19% richness, Lichtenberg

et al., 2017; +23% richness, Gong et al., 2022) and by other in-field diversification measures, like

hedgerows and flower strips (+23% richness, Lichtenberg et al., 2017; +24% richness, Beillouin

et al., 2021). While organic management or in-field diversification measures generally lead to

lower productivity (Gong et al., 2022), strip cropping can maintain crop productivity without

taking land out of production (Campanelli et al., 2023; Juventia and van Apeldoorn, 2024; Van

Oort et al., 2020), although occasional yield reductions have been reported (Carrillo-Reche et al.,

2023; Ditzler et al., 2023). The biodiversity gain in our study was achieved without additional

crop-level diversification strategies designed for biodiversity conservation, such as cover

cropping or (flowering) companion plants. This increase in biodiversity stacks on top of already

higher biodiversity achieved through organic management (Gong et al., 2022; Lichtenberg et al.,

2017). Furthermore, the estimated richness effects of strip cropping, i.e., no change at crop level

and 15% increase at field level, are conservative because they only consider the pairwise

comparison of biodiversity of two or three crops in a strip cropping configuration to

monocultures, whereas the inclusion of more crops within strip cropping could further enhance

ground beetle biodiversity. Also, the effect of strip cropping on ground beetle biodiversity might

be more pronounced in conventional agriculture and at larger field sizes, as the relatively small-

scale, organic fields that we studied might have already had a relatively high ground beetle

biodiversity (Tscharntke et al., 2021). The ground beetle communities in our study were

dominated by farmland and eurytopic species and contained only two rare species (Turin, 2022).

Future research could test whether the inclusion of other in-field diversification measures within

strip-cropped fields, such as the establishment of perennial semi-natural habitats (Rischen et al.,

2021; Sirami et al., 2019) or uptake at larger spatial extents (Tscharntke et al., 2021) would allow

ground beetles with other habitat preferences to establish in agricultural fields.

Materials and methods

Study area

A multi-location study was conducted on four organic farms across the Netherlands (Fig. 1a).

Three experimental farms were managed by Wageningen University & Research (Lelystad,

Valthermond, Wageningen) and one commercial farm was managed by *Exploitatie*

Reservegronden Flevoland (ERF B.V.) located in Almere. All four locations contained both strip

cropping and monocultural crop fields, but differed in soil type, establishment year of the strip

cropping experiment, number of crops grown, length of the crop rotation, number of sampled

crops and sampling years, and farm and landscape characteristics such as percentage of on-farm

semi-natural habitat (SNH), mean field size, and landscape configuration (Fig. 1b, S10). The

locations Almere and Lelystad were located in a homogeneous, open polder landscape

characterized by intensive arable crop production and non-crop habitats consisting of grass

margins, tree lines and watercourses. Valthermond was located in an open, reclaimed peat

landscape with intensive arable crop production characterized by long and narrow fields

separated by grassy margins and ditches and limited areas of woody elements. The site at

Wageningen was located in a more complex landscape with smaller field sizes, and non-crop

habitat consisting of woodlots, hedgerows, tree lines, ditches and farmyards.

Experimental lay-out

At Almere and Valthermond the crops in strip cropping were all grown alongside each other,

whereas at Lelystad and Wageningen two alternating crops (crop pairs) were grown alongside

each other (Fig. 1b, S10). At each location, strip cropping and monoculture fields were always

paired on the same experimental field (Fig. S10). At Almere eight different crops were grown in

alternating strips of 6 m width, including celeriac (*Apium graveolens* var. *rapaceum*), broccoli

(*Brassica oleracea* var. *italic*), oat (*Avena sativa*), onion (*Allium cepa*), parsnip (*Pastinaca*

sativa), faba bean (*Vicia faba*), potato (*Solanum tuberosum*) and a mix of ryegrass and white

clover referred to as grass-clover (*Lolium perenne/Trifolium repens*) (Fig. S10a). At Lelystad

four different crop pairs were grown in alternating strips of 3 m width, including carrot (*Daucus*

carota subsp. sativus) and onion, white cabbage (*Brassica oleracea var. capitata*) and wheat

(*Triticum aestivum*), sugar beet (*Beta vulgaris*) and barley (*Hordeum vulgare)*, and potato and

ryegrass (Fig. S10b). At Valthermond eight different crops were grown in alternating strips of 6

m width, including potato, barley, barley mixed with broad bean (*Vicia faba*) in 2020, barley

mixed with pea (*Pisum sativum*) in 2021, sweet corn (*Zea mays convar. saccharata var. rugosa*),

sugar beet, common bean (*Phaseolus vulgaris*), and grass-clover (Fig. S10c). At Wageningen

three different crop pairs were grown in alternating strips of 3 m width, including white cabbage

and wheat (2019 – 2021) or oat (2022), barley and pumpkin (*Cucurbita maxima*, 2020 - 2022) or

bare soil due to crop failure (2019), and potato and ryegrass (Fig. S10d). The crop combinations

and neighbors were selected based on literature, expert knowledge and experience of

functionality in terms of expected advantages for yield and pest and disease control. Large scale

monoculture plots (0.25 ha to 2.30 ha) served as reference, hereafter referred to as monoculture

(Fig. S10). At Lelystad, Valthermond and Wageningen not each crop grown in strips was present

as monoculture in each year, but only those crops for which a monoculture was present were

sampled (Fig. 1k, Table S7). All fields were managed according to organic regulations, yet at

each location fertilization and weed management reflected regional practices and were adjusted

to local soil conditions. Flower strips were sown within the experimental fields at Almere and

Valthermond (Fig. S10, Table S8).

Sampling

The ground beetle community was sampled using pitfall traps in all crops for which both a

monoculture and strip cropping field were present, at each location and in multiple rounds per

year between March and September. The sampled crops, number of rounds and number of

pitfalls differed per year and per location (Table S7). Pitfall traps consisted of a transparent

plastic cup (9.2 cm diameter, 14 cm height) placed in the soil so the top of the cup was flush with

the soil surface. We did not use funnels inside the cups. Pitfalls were filled with approximately

100 ml water mixed with non-perfumed soap (around 3 cm heigh water level) and covered with a

black plastic rain guard (12.5 cm diameter) around 2 to 5 cm above the soil surface. Pitfall traps

were placed in the center of a strip (1.5 – 3 meters from the edges of the strip), and at least 10

meters from field edges at fixed locations at different moments in a year. Furthermore, traps were

mostly placed at equal distances from the field edges in strip-cropped and monocultural fields. At

Almere and Valthermond, the number of pitfalls in monocultures and strip cropping was the

same. At Wageningen and Lelystad, the number of pitfalls in strip cropped fields was usually

lower than in the monoculture because the area of the strips was only half of that of the

monocultures.

Ground beetle identification

Ground beetles were identified up to species level for Almere in 2021 and 2022, Lelystad and

Wageningen. At Almere in 2020, identification was kept at the genus level due to an extremely

high abundance of *Pterostichus melanarius/niger* and *Poecilus cupreus/versicolor*, and the

associated high time investment for identification. At Valthermond, due to labour constraints, we

chose to consider species that are complicated to distinguish only up to genus level (like *Poecilus*

cupreus/versicolor, *Harpalus spp.* other than *H. rufipes* and *H. affinis*, *Bembidion spp.*). For all

analyses, year series were made, in which all ground beetle catches from the same pitfall trap

were pooled per year. We examined habitat preferences of ground beetle using Turin et al. (2022)

and ground beetle rarity using www.waarneming.nl (checked on 15-01-2025).

Statistical analyses

We used R, version 4.2.2 for all statistical analyses.

Effect of crop configuration on field-level richness and activity density

To analyse the difference in species richness and activity density between monocultures and strip

cropping configurations at field level, we used rarefaction of samples within the same field. To

rarify to an equal sampling intensity, we calculated the average cumulative number of species or

individuals within x year series, where x is the largest number of year series available for the

crop configuration comparison (correcting for unequal sampling between monocultures and strip

cropping, or for missing samples). Next, we calculated the relative change due to strip cropping

by subtracting the number of species or individuals found in the monoculture field from the

number found in the strip cropping field and then dividing the result by the number of species or

individuals in the monoculture (Zou et al., 2020). This gave the relative change centered around

zero, where negative values indicated higher richness in monocultures and positive values higher

richness in strip cropping. We then analysed this data using Generalized Linear Mixed Models

(GLMM) with a Gaussian distribution and assessed whether the intercept deviated significantly

from zero. As random variables, we used location and year, with year nested in location. We ran

these analyses using a dataset that included all comparisons among monocultural fields and strip

cropping fields of all locations. However, as in this case the strip cropped field was compared

with both monocultural fields of the corresponding crop pair, we also tested the effect of field-

level richness using only one monoculture per strip cropped field. To obtain a conservative

estimate of the effect of strip cropping, we chose the monocultural fields with the highest

taxonomic richness or activity density among the constitutive crops of the strip cropping fields.

Generalized linear models were run using the glmmTMB package (Brooks et al., 2017) and

tested for model fit using the DHARMa package (Hartig, 2018).

Effect of crop configuration on crop-level biodiversity

To quantify biodiversity we used five variables: (1) activity density, the total number of ground

beetles found per year series; (2) taxonomic richness, the total number of species or genera

(lowest taxonomic level available) found per year series; (3) the inverse Simpson index, the

inverse of the sum of proportions of different species over the total abundance (Simpson, 1949);

(4) absolute evenness, the number of effective species calculated by dividing the inverse

Simpson index by the taxonomic richness (Williams, 1964); and (5) Shannon entropy (Shannon

and Weaver, 1949). We chose absolute evenness as our measure for evenness, as this method

removes the richness component from the inverse Simpson index and adheres to all requirements

for an evenness index (Smith and Wilson, 1996; Tuomisto, 2012). We included both the inverse

Simpson index and Shannon entropy as the former is more sensitive to changes in evenness and

the latter to species richness (DeJong, 1975).

To analyse the effect of crop configuration on total ground beetle activity density, taxonomic

richness, evenness, inverse Simpson index and Shannon entropy we used GLMM. We

constructed models for each response variable, using data from all four locations. In these models

we included crop configuration (monoculture or strip cropping) as a fixed factor. We included

location, year and crop as nested random variables in these models, with crop nested in year and

year in location. To quantify and visualize the variation in responses between locations, years

and crops, we ran generalized linear models (GLM) with a variable that combined these three

variables into one, which was also included as a fixed factor. Here, we also included the

interaction between crop configuration and the combined variable for crop, location and year. For

the model on activity density we used the negative binomial distribution (log link function), as

this was count data; for richness, evenness and Shannon entropy we used Gaussian distribution;

and for the inverse Simpson index we used a gamma distribution (inverse link function) as a

Gaussian distribution did not fit well. We used the 'DHARMa' package to validate model

assumptions (Hartig, 2018). We used estimated marginal means to assess differences between

monoculture and strip cropping (Hothorn et al., 2016; Lenth, 2018).

To analyse the effect of crop configuration on the activity density of the twelve most common

ground beetle genera, we used another GLMM with negative binomial distribution. Fixed effects

were crop configuration, location and their interactions, and crop nested in year was added as a

random effect. As certain genera only occurred at specific locations, we only included locations

where the genus was found. All models were tested for model fit using the DHARMa package.

We used estimated marginal means to assess differences between monoculture and strip

cropping.

Community composition of crop configurations and crops

To assess whether crop configurations have distinct ground beetle communities we used

permanova with a Hellinger transformation (with 999 permutations) using the "vegan" package

(Oksanen et al., 2013). We used a Hellinger transformation to give more weight to rarer species,

thus accounting for highly abundant species in some samples (such as *P. melanarius* and *H.*

rufipes). We only included species that occurred in at least 3% of the pitfall samples to avoid

strong influence of very rare species. We used data from locations and years where pitfall catches

had been identified to species level (Almere 2021/22, Lelystad and Wageningen). We considered

four models, one for all locations combined and one for each location separately. In all models

we included crop configuration and the nested variables of location (whenever applicable), year

and crop as fixed factors. We also analysed the interaction between crop configuration and this

nested structure of location, year and crop. To visualize these results, we used non-metric

multidimensional scaling (NMDS) on Hellinger transformed data. To show whether crop

configurations have distinct ground beetle communities, we used redundancy analysis (RDA) on

Hellinger transformed data. Here, we again conducted four analyses, one for all locations

combined and one for each of the three considered locations. We only used crop configuration as

a predictor to force RDA to show any change in ground beetle community associated with crop

configuration. As such, only one RDA axis was created per model, which was plotted against the

first principal component describing the residual variation.

Due to the large influence of location and year on ground beetle communities, visualizing any

general effects of strip cropping on these communities using all available data was challenging.

To address this, we conducted RDA and visualized the effect of crop configuration on a subset of

the data from one location and one year. We chose the data from Wageningen in 2021 and 2022

for this analysis because it provided a set-up where strip cropping of two crops could be

compared with their constituent monocultures within the same experimental fields. Here, we

used both crop configuration, crop and their interaction term as explanatory variables for each

crop pair separately. To analyse whether ground beetle communities significantly differed among

combinations of crops and crop configurations we ran pairwise permanova on all three fields and

two years separately using the "pairwiseAdonis" package (Martinez Arbizu, 2020).

Fig. 1. Effect of crop configuration (monoculture versus strip cropping) on ground beetle

biodiversity. (a) Location of experimental sites in the Netherlands. **(b)** Field set-up of the two crop

configurations: monoculture and strip cropping. At Lelystad and Wageningen, strip cropping consisted of

3-m wide crop strips of two crops (pairs), and multiple crop pairs were assessed. At Almere and

Valthermond, strip cropping consisted of 6-m wide crop strips of eight crops combined (Fig. S10). **(c)**

Sample-based species accumulation curves of all year series from monocultures (brown) and strip

cropping (green), in Almere from 2021 and 2022. This was the only location where an equal number of

samples were taken in the monocultures and in strip cropping on a similar area. Ground beetle species

include *Poecilus cupreus* (left) and *Pterostichus melanarius* (right). Photo credit: Ortwin Bleich, retrieved

from: www.eurocarabidae.de. **(d-e)** Overall relative change in field-level ground beetle **(d)** taxonomic

richness and **(e)** activity density. Positive values indicate higher richness or activity density in strip

cropping, negative values in monocultures. **(f-j)** Overall relative effect of crop configuration on ground

beetle **(f)** taxonomic richness, **(g)** activity density, **(h)** absolute evenness, **(i)** inverse Simpson index, and

(j) Shannon entropy. **(k)** Effect of crop configuration on ground beetle taxonomic richness for each

combination of location, year and crop. Barley-mixture consists of a mixture of barley-bean (2020) or

barley-pea (2021). Squares indicate estimated means, the bar indicates the 95% confidence interval.

Asterisks indicate significant differences among the crop configurations. Empty panels indicate

combinations of years and locations that were not sampled. When no estimated mean and confidence

interval are shown, crops were not grown or sampled in that year. Open circles indicate individual year

series to visualize sample size (Table S7).

Fig. 2. Ground beetle species associated with crop configuration (monoculture (a) versus strip cropping (b)). Results obtained by generalized linear mixed models on the twelve most common genera of ground beetles with the four locations analysed separately (Table S5). Only those genera are given for which cropping system significantly influence activity density within at least one location ($\alpha = 0.05$). Locations were the genus had a higher activity density in one of the cropping systems are indicated between brackets (Al=Almere; Le = Lelystad; Va = Valthermond; Wa = Wageningen). For Pterostichus, data from Almere in 2020 was analysed separately, as models did not fit elsewise. The asterisk "*" next to Pterostichus indicates that only in 2020 the difference between monoculture and strip cropping was significant. Photo credit: Ortwin Bleich, retrieved from: www.eurocarabidae.de.

437 **Table 1. Effect of crop configuration on ground beetle community composition.** Results from

438 permanova analysis using Hellinger's transformation for data from the three locations with species level

439 data (see Table S3 for analyses per location). Crops were a nested variable within years, as these differed

440 among years. Years were nested in locations, as the years that were studied differed among locations.

441 Bold letters indicate significant effects ($\alpha = 0.05$).

Location	Predictor	Df	Sum sq	R2	F	P
All locations	Crop configuration	1	1.68	0.01	7.32	**0.001**
	Location	2	47.7	0.29	103.8	**0.001**
	Location : Year	6	10.7	0.06	7.72	**0.001**
	Location : Year : Crop species	31	33.3	0.20	4.67	**0.001**
	Crop configuration : Location	2	0.63	0.00	1.37	0.149
	Crop configuration : Location : Year	6	2.31	0.01	1.67	**0.012**
	Crop configuration : Location : Year : Crop species	31	14.8	0.09	2.08	**0.001**
	Residual	*245*	*56.3*	*0.34*		
	Total	*324*	*167.4*	*1.00*		

Acknowledgements

First, we are grateful to Ron Anbergen, Martine Arkema, Bart Burger, Jonas Driessen, Michiel van de Glind, Angelo Grievink, Roelof Gruppen, Rolinde de Haan, Willem Hendriks, Nashita Maniran, Marina Martino, Sara Michellin, Ciska Nienhuis, Ralph Rustom, Simone Verdonschot, Pim Vrehen, Rik Waenink, and Xiaoshen Wang for their contribution to data collection. This work could not have been completed without the occasional help of many colleagues and students, for this we thank Zhaoqi Bin, Lenora Ditzler, Hilde Faber, Merel Hofmeijer, Gabriel Joachim, Stella Juventia, Peter Karssemeijer, Nelson Ríos Hernández, and any other people that helped. Thanks go to Roy Michielsen and Dirk van de Weert, for maintaining the Almere fields; the team at Wageningen Field crops, and in particular Joost Rijk and Laurens van Run, for maintaining the Lelystad fields; Gerard Hoekzema for maintaining the Valthermond fields; Olivia Elsenpeter, Esther Hofkamp, Titouan le Noc and the Unifarm staff, and in particular Andries Siepel and Peter van der Zee, for maintaining the Wageningen fields. We thank Ortwin Bleich for allowing us to use his ground beetle pictures. We are grateful to Daan Mertens, Marcel Dicke, Liesje Mommer and Thijs Fijen for constructive feedback on earlier versions of this manuscript. Lastly, we thank editors Bernhard Schmid and Sergio Rasmann, and three anonymous reviewers for their extensive and constructive feedback.

Statements

Funding This study has received funding from the European Union's Horizon 2020 research and innovation program under grant agreements No 727482 (DiverIMPACTS) and No 727672 (LegValue), from the Dutch Public Private Partnership research program under grant agreement

No LWV19129 (Crop diversification), from regional funds provided by provinces Groningen and

Drenthe, and through internal Wageningen University and Research funds financed by the Dutch

Ministry of Agriculture, Nature and Food Quality under grant agreement No KB36003003

(Nature Based Solutions in Field Crops). This publication is part of the project CropMix (with

project number NWA.1389.20.160) of the Dutch Research Agenda (NWA-ORC) which is

(partly) financed by the Dutch Research Council (NWO).

Author contributions *Luuk Croijmans:* Conceptualization, Methodology, Formal analysis,

Investigation, Data curation, Writing – original draft, Writing – review & editing, Visualization,

Supervision. *Fogelina Cuperus:* Conceptualization, Methodology, Formal analysis,

Investigation, Data curation, Writing – original draft, Writing – review & editing, Visualization,

Supervision. *Dirk F. van Apeldoorn:* Conceptualization, Writing – review & editing,

Visualization, Supervision, Project administration, Funding acquisition. *Felix J.J.A. Bianchi:*

Conceptualization, Writing – original draft, Writing – review & editing. *Walter A.H. Rossing:*

Conceptualization, Writing – original draft, Writing – review & editing, Project administration,

Funding acquisition. *Erik H. Poelman:* Conceptualization, Methodology, Writing – original draft,

Writing – review & editing, Supervision, Project administration, Funding acquisition.

Conflict of Interest The authors declare that they have no known competing financial interests

or personal relationships that could have influenced the work reported in this paper.

Ethics approval This research was not reviewed by an institutional or governmental regulatory

body as the work was performed on invertebrates.

Availability of data and materials Data and scripts are publicly available from

4TU.ResearchData via the following DOI: https://doi.org/10.4121/bcf78320-aaa6-428f-acf6-

2eb436baa13e (Croijmans et al., 2025).

References

Alarcón-Segura V, Grass I, Breustedt G, Rohlfs M, Tscharntke T. 2022. Strip intercropping of wheat and
oilseed rape enhances biodiversity and biological pest control in a conventionally managed farm
scenario. *Journal of Applied Ecology* **59**:1513–1523. doi:10.1111/1365-2664.14161

Allema B, Van Der Werf W, Van Lenteren JC, Hemerik L, Rossing WAH. 2014. Movement Behaviour of the
Carabid Beetle Pterostichus melanarius in Crops and at a Habitat Interface Explains Patterns of
Population Redistribution in the Field. *PLoS ONE* **9**:e115751. doi:10.1371/journal.pone.0115751

Anderson ME, Harman RR, Kim TN. 2024. Ground beetle movement is deterred by habitat edges: a mark-
release-recapture study on the effectiveness of border crops in an agricultural landscape. *Journal of
Insect Science* **24**:24. doi:10.1093/jisesa/ieae062

Beillouin D, Ben-Ari T, Malézieux E, Seufert V, Makowski D. 2021. Positive but variable effects of crop
diversification on biodiversity and ecosystem services. *Global Change Biology* **27**:4697–4710.
doi:10.1111/gcb.15747

Blowes SA, Daskalova GN, Dornelas M, Engel T, Gotelli NJ, Magurran AE, Martins IS, McGill B, McGlinn
DJ, Sagouis A, Shimadzu H, Supp SR, Chase JM. 2022. Local biodiversity change reflects interactions
among changing abundance, evenness, and richness. *Ecology* **103**:e3820. doi:10.1002/ecy.3820

Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Mächler M,
Bolker BM. 2017. glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated
Generalized Linear Mixed Modeling. *The R Journal* **9**:378–400.

Campanelli G, Iocola I, Leteo F, Montemurro F, Platani C, Testani E, Canali S. 2023. Strip cropping in
organically managed vegetable systems: agronomic and environmental effects. *Renewable Agriculture
and Food Systems* **38**:e31. doi:10.1017/S1742170523000248

Carbonne B, Bohan DA, Foffová H, Daouti E, Frei B, Neidel V, Saska P, Skuhrovec J, Petit S. 2022. Direct
and indirect effects of landscape and field management intensity on carabids through trophic resources
and weeds. *Journal of Applied Ecology* **59**:176–187. doi:10.1111/1365-2664.14043

Carrillo-Reche J, Le Noc T, Van Apeldoorn DF, Juventia SD, Westhoek A, Shanmugam S, Kristensen HL,
Hondebrink M, Himanen SJ, Kivijärvi P, Lepse L, Dane S, Rossing WAH. 2023. Finding guidelines for
cabbage intercropping systems design as a first step in a meta-analysis relay for vegetables. *Agriculture,
Ecosystems & Environment* **354**:108564. doi:10.1016/j.agee.2023.108564

Cozim-Melges F, Ripoll-Bosch R, Veen GF (Ciska), Oggiano P, Bianchi FJJA, van der Putten WH, van Zanten
HHE. 2024. Farming practices to enhance biodiversity across biomes: a systematic review. *npj biodivers*
3:1–11. doi:10.1038/s44185-023-00034-2

Croijmans L, van Apeldoorn DF, Sanfilippo F, Zangpo T, Poelman EH. 2024. Crop species diversity levels
with attract and reward strategies to enhance Pieris brassicae parasitism rate by Cotesia glomerata in
strip intercropping. *Functional Ecology* **38**:654–667. doi:10.1111/1365-2435.14502

Crutsinger GM, Collins MD, Fordyce JA, Gompert Z, Nice CC, Sanders NJ. 2006. Plant Genotypic Diversity
Predicts Community Structure and Governs an Ecosystem Process. *Science* **313**:966–968.
doi:10.1126/science.1128326

Cuperus F, Allema B, Bianchi FJJA, Rossing WAH, van Apeldoorn DF. 2024. Cauliflower strip cropping
promotes ground-dwelling arthropod richness and spider abundance. *Agriculture, Ecosystems &*
Environment **374**:109137. doi:10.1016/j.agee.2024.109137

Cuperus F, Ozinga WA, Bianchi FJJA, Croijmans L, Rossing WAH, Van Apeldoorn DF. 2023. Effects of field-
level strip and mixed cropping on aerial arthropod and arable flora communities. *Agriculture, Ecosystems*
& Environment **354**:108568. doi:10.1016/j.agee.2023.108568

Díaz S, Settele J, Brondízio ES, Ngo HT, Agard J, Arneth A, Balvanera P, Brauman KA, Butchart SHM, Chan
KMA, Garibaldi LA, Ichii K, Liu J, Subramanian SM, Midgley GF, Miloslavich P, Molnár Z, Obura D, Pfaff A,
Polasky S, Purvis A, Razzaque J, Reyers B, Chowdhury RR, Shin Y-J, Visseren-Hamakers I, Willis KJ, Zayas
CN. 2019. Pervasive human-driven decline of life on Earth points to the need for transformative change.
Science **366**:eaax3100. doi:10.1126/science.aax3100

Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B. 2014. Defaunation in the Anthropocene.
Science **345**:401–406. doi:10.1126/science.1251817

Ditzler L, Apeldoorn DF van, Schulte RPO, Tittonell P, Rossing WAH. 2021. Redefining the field to mobilize
three-dimensional diversity and ecosystem services on the arable farm. *European Journal of Agronomy*
122:126197. doi:10.1016/j.eja.2020.126197

Ditzler L, Rossing WAH, Schulte RPO, Hageman J, Van Apeldoorn DF. 2023. Prospects for increasing the
resolution of crop diversity for agroecosystem service delivery in a Dutch arable system. *Agriculture,*
Ecosystems & Environment **351**:108472. doi:10.1016/j.agee.2023.108472

Eyre MD, Labanowska-Bury D, Avayanos JG, White R, Leifert C. 2009. Ground beetles (Coleoptera,
Carabidae) in an intensively managed vegetable crop landscape in eastern England. *Agriculture,*
Ecosystems & Environment **131**:340–346. doi:10.1016/j.agee.2009.02.006

Finch S. 1996. Effect of beetle size on predation of cabbage root fly eggs by ground beetles. *Entomologia*
Experimentalis et Applicata **81**:199–206. doi:10.1111/j.1570-7458.1996.tb02032.x

Gong S, Hodgson JA, Tscharntke T, Liu Y, Van Der Werf W, Batáry P, Knops JMH, Zou Y. 2022. Biodiversity
and yield trade-offs for organic farming. *Ecology Letters* **25**:1699–1710. doi:10.1111/ele.14017

Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, Stenmans W, Müller A, Sumser H,
Hörren T, Goulson D, De Kroon H. 2017. More than 75 percent decline over 27 years in total flying insect
biomass in protected areas. *PLoS ONE* **12**:e0185809. doi:10.1371/journal.pone.0185809

Hartig F. 2018. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models.
R Packag version 020.

Holland JM, Luff ML. 2000. The Effects of Agricultural Practices on Carabidae in Temperate
Agroecosystems. *Integrated Pest Management Reviews* **5**:109–129.

Hothorn T, Bretz F, Westfall P, Heiberger RM, Scheutzenmeister A, Schreibe S. 2016. Package
"multcomp."

Hummel JD, Dosdall LM, Clayton GW, Harker KN, O'Donovan JT. 2012. Ground Beetle (Coleoptera:
Carabidae) Diversity, Activity Density, and Community Structure in a Diversified Agroecosystem.
Environmental Entomology **41**:72–80. doi:10.1603/EN11072

Juventia SD, van Apeldoorn DF. 2024. Strip cropping increases yield and revenue: multi-year analysis of
an organic system in the Netherlands. *Frontiers in Sustainable Food Systems* **8**.
doi:10.3389/fsufs.2024.1452779

Karssemeijer PN, Croijmans L, Gajendiran K, Gols R, Van Apeldoorn DF, Van Loon JJA, Dicke M, Poelman
EH. 2024. Diverse cropping systems lead to higher larval mortality of the cabbage root fly (Delia
radicum). *Journal of Pest Science* **97**:337–353. doi:10.1007/s10340-023-01629-1

Lenth R. 2018. Package "lsmeans." *The American Statistician* **34.4**:216–221.

Lichtenberg EM, Kennedy CM, Kremen C, Batáry P, Berendse F, Bommarco R, Bosque-Pérez NA,
Carvalheiro LG, Snyder WE, Williams NM, Winfree R, Klatt BK, Åström S, Benjamin F, Brittain C,
Chaplin-Kramer R, Clough Y, Danforth B, Diekötter T, Eigenbrode SD, Ekroos J, Elle E, Freitas BM, Fukuda
Y, Gaines-Day HR, Grab H, Gratton C, Holzschuh A, Isaacs R, Isaia M, Jha S, Jonason D, Jones VP, Klein A,
Krauss J, Letourneau DK, Macfadyen S, Mallinger RE, Martin EA, Martinez E, Memmott J, Morandin L,
Neame L, Otieno M, Park MG, Pfiffner L, Pocock MJO, Ponce C, Potts SG, Poveda K, Ramos M, Rosenheim
JA, Rundlöf M, Sardiñas H, Saunders ME, Schon NL, Sciligo AR, Sidhu CS, Steffan-Dewenter I, Tscharntke
T, Veselý M, Weisser WW, Wilson JK, Crowder DW. 2017. A global synthesis of the effects of diversified
farming systems on arthropod diversity within fields and across agricultural landscapes. *Global Change*
Biology **23**:4946–4957. doi:10.1111/gcb.13714

Makwela MM, Slotow R, Munyai TC. 2023. Carabid Beetles (Coleoptera) as Indicators of Sustainability in
Agroecosystems: A Systematic Review. *Sustainability* **15**:3936. doi:10.3390/su15053936

Martinez Arbizu P. 2020. pairwiseAdonis: Pairwise multilevel comparison using adonis.

Martin-Guay M-O, Paquette A, Dupras J, Rivest D. 2018. The new Green Revolution: Sustainable
intensification of agriculture by intercropping. *Science of The Total Environment* **615**:767–772.
doi:10.1016/j.scitotenv.2017.10.024

Müller J, Hothorn T, Yuan Y, Seibold S, Mitesser O, Rothacher J, Freund J, Wild C, Wolz M, Menzel A.
2024. Weather explains the decline and rise of insect biomass over 34 years. *Nature* **628**:349–354.
doi:10.1038/s41586-023-06402-z

Mupepele A-C, Bruelheide H, Brühl C, Dauber J, Fenske M, Freibauer A, Gerowitt B, Krüß A, Lakner S,
Plieninger T, Potthast T, Schlacke S, Seppelt R, Stützel H, Weisser W, Wägele W, Böhning-Gaese K, Klein
A-M. 2021. Biodiversity in European agricultural landscapes: transformative societal changes needed.
Trends in Ecology & Evolution **36**:1067–1070. doi:10.1016/j.tree.2021.08.014

Oksanen J, Simpson GL, Guillaume Blanchet F, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos P,
Stevens MHH, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G, Chirico M,
de Caceres M, Durand S, Antoniazi Evangelista HB, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill

MO, Lahti L, McGlinn D, Ouellette M-H, Ribeiro Cunha E, Smith T, Stier A, ter Braak CJF, Weedon J,
Borman T. 2013. Package "Vegan."

Pilotto F, Kühn I, Adrian R, Alber R, Alignier A, Andrews C, Bäck J, Barbaro L, Beaumont D, Beenaerts N,
Benham S, Boukal DS, Bretagnolle V, Camatti E, Canullo R, Cardoso PG, Ens BJ, Everaert G, Evtimova V,
Feuchtmayr H, García-González R, Gómez García D, Grandin U, Gutowski JM, Hadar L, Halada L, Halassy
M, Hummel H, Huttunen K-L, Jaroszewicz B, Jensen TC, Kalivoda H, Schmidt IK, Kröncke I, Leinonen R,
Martinho F, Meesenburg H, Meyer J, Minerbi S, Monteith D, Nikolov BP, Oro D, Ozoliņš D, Padedda BM,
Pallett D, Pansera M, Pardal MÂ, Petriccione B, Pipan T, Pöyry J, Schäfer SM, Schaub M, Schneider SC,
Skuja A, Soetaert K, Spriņģe G, Stanchev R, Stockan JA, Stoll S, Sundqvist L, Thimonier A, Van Hoey G, Van
Ryckegem G, Visser ME, Vorhauser S, Haase P. 2020. Meta-analysis of multidecadal biodiversity trends in
Europe. *Nature Communications* **11**:3486. doi:10.1038/s41467-020-17171-y

Puliga GA, Arlotti D, Dauber J. 2023. The effects of wheat-pea mixed intercropping on biocontrol
potential of generalist predators in a long-term experimental trial. *Annals of Applied Biology* **182**:37–47.
doi:10.1111/aab.12792

Rakotomalala AANA, Ficiciyan AM, Tscharntke T. 2023. Intercropping enhances beneficial arthropods
and controls pests: A systematic review and meta-analysis. *Agriculture, Ecosystems & Environment*
356:108617. doi:10.1016/j.agee.2023.108617

Ratnadass A, Fernandes P, Avelino J, Habib R. 2012. Plant species diversity for sustainable management
of crop pests and diseases in agroecosystems: a review. *Agronomy for Sustainable Development* **32**:273–
303. doi:10.1007/s13593-011-0022-4

Rischen T, Frenzel T, Fischer K. 2021. Biodiversity in agricultural landscapes: different non-crop habitats
increase diversity of ground-dwelling beetles (Coleoptera) but support different communities.
Biodiversity and Conservation **30**:3965–3981. doi:10.1007/s10531-021-02284-7

Roubinet E, Jonsson T, Malsher G, Staudacher K, Traugott M, Ekbom B, Jonsson M. 2018. High
Redundancy as well as Complementary Prey Choice Characterize Generalist Predator Food Webs in
Agroecosystems. *Scientific Reports* **8**:8054. doi:10.1038/s41598-018-26191-0

Saunders ME. 2020. Conceptual ambiguity hinders measurement and management of ecosystem
disservices. *Journal of Applied Ecology* **57**:1840–1846. doi:10.1111/1365-2664.13665

Seibold S, Gossner MM, Simons NK, Blüthgen N, Müller J, Ambarlı D, Ammer C, Bauhus J, Fischer M,
Habel JC, Linsenmair KE, Nauss T, Penone C, Prati D, Schall P, Schulze E-D, Vogt J, Wöllauer S, Weisser
WW. 2019. Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature*
574:671–674. doi:10.1038/s41586-019-1684-3

Sirami C, Gross N, Baillod AB, Bertrand C, Carrié R, Hass A, Henckel L, Miguet P, Vuillot C, Alignier A,
Girard J, Batáry P, Clough Y, Violle C, Giralt D, Bota G, Badenhausser I, Lefebvre G, Gauffre B, Vialatte A,
Calatayud F, Gil-Tena A, Tischendorf L, Mitchell S, Lindsay K, Georges R, Hilaire S, Recasens J, Solé-Senan
XO, Robleño I, Bosch J, Barrientos JA, Ricarte A, Marcos-Garcia MÁ, Miñano J, Mathevet R, Gibon A,
Baudry J, Balent G, Poulin B, Burel F, Tscharntke T, Bretagnolle V, Siriwardena G, Ouin A, Brotons L,
Martin J-L, Fahrig L. 2019. Increasing crop heterogeneity enhances multitrophic diversity across

agricultural regions. *Proceedings of the National Academy of Sciences* **116**:16442–16447.
doi:10.1073/pnas.1906419116

Tamburini G, Bommarco R, Wanger TC, Kremen C, Van Der Heijden MGA, Liebman M, Hallin S. 2020.
Agricultural diversification promotes multiple ecosystem services without compromising yield. *Sci Adv*
6:eaba1715. doi:10.1126/sciadv.aba1715

Tscharntke T, Grass I, Wanger TC, Westphal C, Batáry P. 2021. Beyond organic farming – harnessing
biodiversity-friendly landscapes. *Trends in Ecology & Evolution* **36**:919–930.
doi:10.1016/j.tree.2021.06.010

Turin H. 2022. Ecology and conservation of the Dutch ground beetle fauna: lessons from 66 years of
pitfall trapping. Wageningen: Wageningen Academic Publishers.

Turin H. 2000. De Nederlandse loopkevers: verspreiding en oecologie (Coleoptera: Carabidae),
Nederlandse fauna. Leiden: Nationaal Natuurhistorisch Museum.

Turin H, Heijerman T, Noordijk J, Trautner J. 2012. Het recente voorkomen van de loopkever Harpalus
signaticornis in Nederland (Coleoptera: Carabidae). *Nederlandse Faunistische Mededelingen* **38**:9–16.

van Klink R, Bowler DE, Gongalsky KB, Shen M, Swengel SR, Chase JM. 2024. Disproportionate declines of
formerly abundant species underlie insect loss. *Nature* **628**:359–364. doi:10.1038/s41586-023-06861-4

Van Klink R, Bowler DE, Gongalsky KB, Swengel AB, Gentile A, Chase JM. 2020. Meta-analysis reveals
declines in terrestrial but increases in freshwater insect abundances. *Science* **368**:417–420.
doi:10.1126/science.aax9931

Van Oort PAJ, Gou F, Stomph TJ, Van Der Werf W. 2020. Effects of strip width on yields in relay-strip
intercropping: A simulation study. *European Journal of Agronomy* **112**:125936.
doi:10.1016/j.eja.2019.125936

Wagner DL, Grames EM, Forister ML, Berenbaum MR, Stopak D. 2021. Insect decline in the
Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of Sciences*
118:e2023989118. doi:10.1073/pnas.2023989118

Wallin H, Ekbom B. 1994. Influence of hunger level and prey densities on movement patterns in three
species of Pterostichus beetles (Coleoptera: Carabidae). *Environmental Entomology* **23**:1171–1181.

Zou Y, de Kraker J, Bianchi FJJA, Xiao H, Huang J, Deng X, Hou L, van der Werf W. 2020. Do diverse
landscapes provide for effective natural pest control in subtropical rice? *Journal of Applied Ecology*
57:170–180. doi:10.1111/1365-2664.13520

665