PGR5 is needed for redox-dependent regulation of ATP synthase both in chloroplasts and in

cyanobacteria

Lauri Nikkanen[1]*, Laura T. Wey[1], Russell Woodford[2], Henna Mustila[1], Darius Kosmützky[3], Maria

Ermakova[2], Eevi Rintamäki[1], and Yagut Allahverdiyeva[1]*

[1]Molecular Plant Biology, Department of Life Technologies, University of Turku, Finland

[2]School of Biological Sciences, Monash University, Australia

[3]Department of Biochemistry, University of Cambridge, United Kingdom

*Co-corresponding authors

Authors for Correspondence:

Lauri Nikkanen, Molecular Plant Biology, Department of Life Technologies, University of Turku, Turku 11 FI- 20014,

Finland. E-mail: lenikk@utu.fi

Yagut Allahverdiyeva, Molecular Plant Biology, Department of Life Technologies, University of Turku, Turku 11 FI-

20014, Finland. E-mail: allahve@utu.fi

ORCID identifiers:

Lauri Nikkanen: 0000-0002-7192-9322

Laura T. Wey: 0000-0003-2345-0699

Russell Woodford: 0000-0002-6766-2274

Henna Mustila: 0000-0002-1939-7197

Darius Kosmützky: 0000-0002-6678-7550

Maria Ermakova: 0000-0001-8466-4186

Eevi Rintamäki: 0000-0002-1353-3995

Yagut Allahverdiyeva: 0000-0002-9262-1757

Running title: PGR5-dependent regulation of ATP synthase

Number of Figures: 8

Supplementary Data: 18 Figures and 10 Tables

Word count: 9598

Highlight

PGR5 is involved in regulation of ATP synthase in fluctuating light conditions across photosynthetic
organisms in a manner that depends on the thiol redox state of the chloroplast or cell.

ABSTRACT

The regulation of proton motive force (*pmf*) via ATP synthase activity is a critical mechanism by
which photosynthetic organisms maintain redox homeostasis and activate photoprotective responses
under fluctuating light conditions. Here, we used time-resolved electrochromic shift (ECS)
measurements to investigate *pmf* dynamics across a range of photosynthetic organisms. Our results
reveal that ATP synthase is dynamically regulated during light fluctuations, but in *Arabidopsis*
thaliana could this regulation could not be explained by the known light-induced reduction of the
CF□γ subunit, suggesting alternative control mechanisms. The PROTON GRADIENT
REGULATION 5 (PGR5) protein, previously proposed to facilitate cyclic electron transport (CET),
also has a potential role in ATP synthase regulation. The physiological role of cyanobacterial Pgr5 has
remained elusive. We characterised a Δ*pgr5* mutant of *Synechocystis* sp. PCC 6803 and investigated
pmf dynamics in *pgr5* mutants of *Chlamydomonas reinhardtii*, Arabidopsis, and the C4 grass *Setaria*
viridis. While PGR5 was not required for CET in *Synechocystis*, it was needed for downregulating
ATP synthase under high irradiance in all tested organisms via a thiol redox state dependent
mechanism. Furthermore, interaction between AtPGR5 and AtCF□γ suggests that PGR5 functions as
a conserved inhibitor of ATP synthase under high light, contributing to *pmf* retention and
photoprotection.

51
52
53 KEYWORDS: Photosynthesis, ATP synthase, proton motive force, chloroplast, cyanobacteria,
54 Arabidopsis, electrochromic shift

INTRODUCTION

Oxygenic photosynthesis by plants, algae, and cyanobacteria uses light energy to drive oxidation of water to produce ATP and NADPH for CO_2 fixation and other cellular processes. In their natural habitats, oxygenic photosynthetic organisms undergo constant fluctuation of irradiance. Mechanisms to adjust the redox homeostasis in chloroplasts and cyanobacterial cells have evolved to prevent energy-consuming futile reactions and damage to the photosynthetic machinery and other biomolecules via generation of reactive oxygen species (ROS) by the photosynthetic apparatus. ROS can be generated as a result of excitation imbalance between the photosystems, or when electron transport rate exceeds the sink capacity of the reactions downstream of the photosynthetic electron transport chain (PETC). Resulting accumulation of reduced electron carriers in the PETC triggers charge recombination reactions within the photosystems and direct reduction of O_2, generating highly reactive ROS such as superoxide ($O_2^{\bullet-}$) and hydrogen peroxide (H_2O_2) at PSI and singlet oxygen ($^1O_2$) primarily at PSII (Foyer & Hanke, 2022). PSI and its iron sulphur clusters are particularly vulnerable to over-reduction-induced damage, and plants, algae, and cyanobacteria invest heavily in their protection (Tiwari *et al.*, 2016, 2024). These protective mechanisms involve cyclic electron transport around PSI (CET), redistribution of excitation energy between the photosystems via state transitions, non-photochemical quenching (NPQ), and inhibition of PQH_2 oxidation at the cytochrome b_6f complex, known as photosynthetic control (Tikkanen & Aro, 2014; Shimakawa *et al.*, 2016; Nikkanen *et al.*, 2021; Eckardt *et al.*, 2024; Degen & Johnson, 2024). Additionally, in photosynthetic organisms apart from angiosperms, flavodiiron proteins provide a photoprotective electron sink on the acceptor side of PSI by catalysing controlled reduction of O_2 to water (the Mehler-like reaction) (Allahverdiyeva *et al.*, 2013; Ilik *et al.*, 2017).

As photosynthetic control and NPQ (in plants and algae) are induced by acidic pH of the thylakoid lumen, an important mechanism to maintain photosynthetic redox balance involves control of the proton motive force *(pmf)* via regulation of the thylakoid ATP synthase (Kanazawa *et al.*, 2017; Takagi *et al.*, 2017). The pH-dependent component of *pmf* (ΔpH), is generated by release of protons to the lumenal side of the thylakoid membrane by water oxidation at PSII and the Q cycle, and proton pumping from the stromal/cytosolic side by cyclic electron transport, as well as consumption of protons on the stromal side. The proton gradient is released mainly via efflux through the ATP synthase, providing the driving force for ATP synthesis. Proton-conductivity of the ATP synthase is therefore an important regulatory factor for adjusting the magnitude of the *pmf*.

In addition to ΔpH, the *pmf* comprises an electric field component ($\Delta\Psi$). $\Delta\Psi$ derives from charge separation within the photosystems and in the Q cycle, the positive charge of protons, and exchange of cations (e.g. K^+, Ca^{2+} and Mg^{2+}) in and anions (e.g. Cl^-) out of the lumen by ion channels and transporters in the thylakoid membrane (Armbruster *et al.*, 2017). ΔpH and $\Delta\psi$ are

thermodynamically equivalent and both drive ATP synthesis (Hangarter & Good, 1982). However,

build-up of sufficient ΔpH over the thylakoid membrane is required for induction of photoprotective

mechanisms upon sudden increases in light intensity, namely dissipation of excessive excitation

energy as heat in NPQ (Niyogi & Truong, 2013; Krishnan-Schmieden *et al.*, 2021) and photosynthetic

control (Tikhonov *et al.*, 1981; Degen & Johnson, 2024). These mechanisms prevent over-reduction of

the electron transport chain and photodamage to PSI. Conversely, it is also important to down-

regulate ΔpH (and consequently, NPQ and photosynthetic control) upon sudden decreases in

irradiance in order to maintain sufficient supply of electrons on the acceptor side of the electron

transfer chain for carbon fixation in the Calvin–Benson cycle (CBC).

ATP synthase is regulated via several mechanisms. The $F_1\epsilon$ (epsilon) subunits of the chloroplast

(CF_0CF_1) ATP synthase (Feniouk *et al.*, 2006) and the cyanobacterial ATP synthase (Imashimizu *et*

al., 2011; Murakami *et al.*, 2018) have inhibitory roles preventing ATP hydrolysis. $F_1\epsilon$ undergoes

conformational changes triggered by *pmf* generation and changes in ATP/ADP ratio, allowing

inhibition of ATP hydrolysis (Richter & McCarty, 1987; Feniouk *et al.*, 2006). Moreover, upon dark-

to-light transitions, the chloroplast ATP synthase is activated by thioredoxin (TRX)-mediated

reduction of a disulphide in the γ subunit ($CF_1\gamma$) (McKinney *et al.*, 1978; Nalin & McCarty, 1984;

Akiyama *et al.*, 2023). $CF_1\gamma$ is kept in an inactive oxidised state in darkness to avoid hydrolysis of

ATP by reverse activity or loss of the proton gradient (Hisabori *et al.*, 2013; Kohzuma *et al.*, 2017;

Hahn *et al.*, 2018). Interestingly, it was recently reported that the *Chlamydomonas reinhardtii* $CF_1\gamma$,

although the conserved disulphide is present, contains a modified hairpin loop in the redox regulatory

domain that allows $CrCF_0CF_1$ to remain active also in darkness to facilitate heterotrophic metabolism

(Lebok & Buchert, 2024). In plants, maintenance of adequate dark-*pmf* via ATP synthase dark-light

regulation is also needed for powering the insertion of new proteins into the thylakoid membrane via

the Tat- and Sec-pathways to maintain the integrity of the photosynthetic apparatus (Kohzuma *et al.*,

2017). $CF_1\gamma$ is reduced in seconds upon illumination, which lowers the threshold *pmf* for activation of

the ATP synthase (Kramer *et al.*, 1990; Konno *et al.*, 2012; Kohzuma *et al.*, 2017; Nikkanen *et al.*,

2018; Buchert *et al.*, 2021). In turn, ΔpH also facilitates reduction of the $CF_1\gamma$ disulphide bond

(Sekiguchi *et al.*, 2024), while dissipation of ΔpH promotes oxidation of $CF_1\gamma$ (Sekiguchi *et al.*, 2022).

F- and m-type chloroplast TRX isoforms have been understood to be mainly responsible for light-

dependent reduction of $CF_1\gamma$ (Schwarz *et al.*, 1997; Sekiguchi *et al.*, 2020), and chloroplast TRX-like

proteins (TrxL2) and 2-cysteine peroxiredoxins (2-CysPrxs) catalyse the oxidation of the subunit in

the dark (Yoshida *et al.*, 2018; Sekiguchi *et al.*, 2022). Additionally, *in vivo* studies have revealed the

involvement of the NADPH-dependent TRX system (NTRC) in $CF_1\gamma$ reduction in low light

conditions (Carrillo *et al.*, 2016; Nikkanen *et al.*, 2016), while the atypical TRX CDSP32 was recently

suggested to play a role in reduction of $CF_1\gamma$ in potato (Rey *et al.*, 2024).

Reduction of $CF_1\gamma$ constitutes an on/off switch controlling the activation of the ATP synthase upon

the onset of illumination in chloroplasts of plants and algae, and does not occur in mitochondria or in

cyanobacteria (Hisabori *et al.*, 2013). The cyanobacterial γ subunit still has a role in preventing ATP

hydrolysis however, via an ADP-dependent mechanism (Sunamura *et al.*, 2010). Moreover, the small

AtpΘ protein (Song *et al.*, 2022) has been reported to function as an inhibitor of the cyanobacterial

ATP synthase in the dark and in low *pmf* conditions, preventing ATP hydrolysis and collapse of the

proton gradient. Interestingly, the $F_1\alpha$ (AtpA) and $F_1\beta$ (AtpB) subunits of the ATP synthase also

contain conserved Cys residues, and $F_1\alpha$ has been identified as a prospective TRX target in

cyanobacteria (Lindahl & Florencio, 2003; Guo *et al.*, 2014), and $F_1\beta$ in plants (Balmer *et al.*, 2006).

Moreover, the mitochondrial ATP synthase in *Xenopus laeviis* oocytes is regulated by reversible

oxidation of thiols in the $F_1\alpha$ subunit (Cobley *et al.*, 2020).

There are also other regulatory mechanisms that adjust the activation state of the ATP synthase

according to the metabolic state of the stroma or cytosol, such as substrate (ADP and P_i) availability; a

lack of Pi due to low consumption of ATP by an inactive CBC would lead to a decrease ATP synthase

activity, increasing ΔpH and activating photosynthetic control and NPQ to protect PSI from over-

reduction (Kanazawa & Kramer, 2002; Avenson *et al.*, 2005; Takagi *et al.*, 2017). High CO_2 supply

results in increase of chloroplast ATP synthase activity by a mechanism that is independent of $CF_1\gamma$

redox state (Kiirats *et al.*, 2009; Kohzuma *et al.*, 2013). Another potential regulatory mechanism has

been suggested to involve phosphorylation of the $CF_1\beta$ subunit (del Riego *et al.*, 2006), although a

more recent study reports that the phosphorylation is unlikely to have a role in controlling ATP

synthase activity and is instead involved in CF_1 assembly (Strand *et al.*, 2023).

Additionally, thylakoid conductivity and the *pmf* can be adjusted through the activities of ion channels

and transporters in the thylakoid membrane (Spetea *et al.*, 2017). A lack of the K^+/H^+ exchanger

KEA3 or the Cl^- channels CLCe and VCCN1 has been shown to affect the *pmf* and/or induction and

relaxation of NPQ in plants (Armbruster *et al.*, 2014; Herdean *et al.*, 2016b,a; Wang & Shikanai,

2019; von Bismarck *et al.*, 2023). Similarly, lack of the cyanobacterial K^+ channel SynK resulted in

decreased ΔpH generation and impairment of photosynthetic control in the model cyanobacterium

Synechocystis sp. PCC 6803 (*Synechocystis*) (Checchetto *et al.*, 2012). Controlling the ionic

environment of the thylakoid membrane is particularly important to avoid drastic increases in ΔΨ,

which result in production of damaging $^1O_2$ via recombination reactions in PSII (Davis *et al.*, 2016).

Plants deficient in the PROTON GRADIENT REGULATION 5 (PGR5) protein suffer from impaired

pmf generation (Munekage *et al.*, 2002). In plant chloroplasts, PGR5 forms redox-dependent

heterodimers with PGRL1 (PGR5-LIKE 1), which have been proposed to mediate or regulate cyclic

electron transport (CET) from ferredoxin to the plastoquinone pool (DalCorso *et al.*, 2008; Hertle *et*

al., 2013; Yamamoto & Shikanai, 2019), contributing to *pmf* formation (Wang *et al.*, 2015). More

recently, a functional PGRL1-analogue was identified also in cyanobacteria (Dann & Leister, 2019).

The *pgr5* knockout plants are unable to induce photosynthetic control and NPQ during transitions

from low to high light (HL) (Munekage *et al.*, 2002; Suorsa *et al.*, 2012; Yamamoto & Shikanai,

2019). Consequently, exposure of *pgr5* mutant plants to HL results in over-reduction of the electron

transport chain and photodamage to the iron sulphur clusters of PSI (Tiwari *et al.*, 2016, 2024).

Accordingly, the lack of PGR5 is lethal in fluctuating light conditions in *Arabidopsis thaliana*

(Arabidopsis) (Tikkanen *et al.*, 2010; Suorsa *et al.*, 2012) and severely inhibits growth of the model C_4

plant *Setaria viridis* (Woodford *et al.*, 2024) as well as of the model green alga *Chlamydomonas*

reinhardtii (Chlamydomonas) (Jokel *et al.*, 2018). *Synechocystis* cells deficient in a PGR5-orthologue

(SynPgr5) suffer from slightly increased HL-sensitivity (Yeremenko *et al.*, 2005), but no growth

impairment was observed in fluctuating light conditions (Allahverdiyeva *et al.*, 2013). Interestingly,

however, there is strong selection pressure against overexpression of SynPgr5 (Margulis *et al.*, 2020).

The physiological functions of PGR5 and PGRL1 remain controversial, with some reports linking the

proteins to regulation of linear instead of cyclic electron transport from PSI (Nandha *et al.*, 2007;

Joliot & Johnson, 2011; Johnson *et al.*, 2014; Tikkanen *et al.*, 2015; Takagi & Miyake, 2018; Rantala *et*

al., 2020; Maekawa *et al.*, 2024), or regulation of the Q-cycle at the cytochrome b_6f complex (Buchert

et al., 2020). Impaired *pmf* generation and induction of photoprotection in the *pgr5* mutant could also

derive from the inability of the mutant plants to limit the conductivity of the thylakoid membrane

under HL (Avenson *et al.*, 2005; Nikkanen *et al.*, 2018; Degen *et al.*, 2023). Interestingly, elevated

thylakoid (ATP synthase) conductivity upon HL exposure was not observed in isolated and ruptured

pgr5 mutant chloroplasts (Wang *et al.*, 2018). This suggests that a stromal component that is lost in

ruptured chloroplasts plays a role in the effect of PGR5 on thylakoid conductivity.

In the current study, we elucidated the dynamics of *pmf* regulation under light in a variety of

photosynthetic organisms focusing on the potential role of PGR5. We performed *in vivo*

characterisation of photosynthetic electron transport and *pmf* formation in *Synechocystis,*

Chlamydonomas, *Arabidopsis,* and *S. viridis.* Our results reveal that in *Arabidopsis*, dynamic

regulation of ATP synthase conductivity during fluctuations in light intensity is independent of the

thiol redox state of the $CF_1\gamma$ subunit. We show that downregulation of the ATP synthase activity upon

transitions to HL depends on PGR5 as well as the thiol redox state of the stroma or cytosol in both

chloroplasts and cyanobacteria. Unlike in chloroplasts, PGR5 is not essential for oxidation of the

PETC in cyanobacteria. Moreover, we demonstrate that cyanobacterial SynPgr5 is not needed for

CET and suggest that the observed effect of plant PGR5 on cyclic electron transport may be an

indirect consequence of ATP synthase misregulation and increased photoinhibition of PSI.

MATERIALS AND METHODS

Strains, plant lines and growth conditions

Arabidopsis thaliana WT-gl1 or WT-Col-0, *pgr5-1* (At2g05620 knockout) (Munekage *et al.*, 2002), and *pgr5-Cas* (At2g05620 CRISPR-Cas9 mutant line) (Penzler *et al.*, 2022) plants were grown for 4 weeks under 200 µmol photons in a 12/12h photoperiod before collecting individual leaves for experiments.

For the BiFC experiments, WT *Nicotiana benthamiana* plants were grown under 130 µmol photons $m^{-2}\,s^{-1}$ at 23°C in a 16/8 hr light/dark photoperiod.

Chlamydomonas reinhardtii WT 137c (progenitor strain of the CC-124 and CC-125 WT strains), *pgr5* knockout and PGR5C complementation strains (Johnson *et al.*, 2014; Jokel *et al.*, 2018) were grown photoautotrophically under continuous 50 µmol photons $m^{-2}\,s^{-1}$ at 25 °C and 120 rpm in TP medium for four days.

Setaria viridis cv MEO V34-1 *pgr5* knock-out mutant was generated using CRISPR-Cas9 gene-editing as described in detail in (Ermakova *et al.*, 2024). WT plants and plants with *pgr5-3* null alleles described in (Woodford *et al.*, 2024) were analysed. Plants were grown in a controlled environment room with a 16 h : 8 h, light : dark photoperiod, 28°C day and 22°C night, 60% humidity and ambient CO_2 at a light intensity of 300 µmol $m^{-2}\,s^{-1}$ supplied by 1000 W red sunrise 3200K lamps (Sunmaster Growlamps, Solon, OH, USA). Youngest fully expanded leaves of 3–4-week-old plants were used in analysis.

WT, Δpgr5, Δflv3 (Helman *et al.*, 2003) and *Δpgr5 flv3 Synechocystis* sp. PCC 6803 strains as well as WT *Synechococcus elongatus* sp. PCC 7942 cells were grown under continuous 50 µmol photons $m^{-2}\,s^{-1}$ at 30 °C and air-level $[CO_2]$ in pH 7.5 BG-11 medium. Unless stated otherwise, cultures were grown for 4 days in 30 ml flasks, then pelleted and resuspended in fresh BG-11 at an appropriate chlorophyll concentration for each experiment. For the growth curves under fluctuating light, WT, *Δpgr5* and *Δpgr5 flv3* cells were grown for 7 days in conditions where light intensity was held at 20 µmol photons $m^{-2}\,s^{-1}$ and increased to 500 µmol photons $m^{-2}\,s^{-1}$ for 30 sec every 5 min. Samples were taken and $OD_{750\ nm}$ measured with a spectrophotometer every 24h. For the experiment in Figure 5, cells were grown in a regime in which light intensity fluctuated between 1 min periods of 25 and 500 µmol photons $m^{-2}\,s^{-1}$.

Construction of Δ*pgr5* and Δ*pgr5 flv3* knockout mutants

The upstream and downstream flanking sequences of the *pgr5* (*ssr2016*) gene and the *aphI* (Km^R) gene replacing the *ssr2016* open reading frame were amplified from genomic DNA of Δ*pgr5* mutant generated earlier in a different *Synechocystis* WT background as described in (Allahverdiyeva *et al.*, 2013). After transformation into *Synechocystis* sp. PCC 6803 WT strain and the Δ*flv3* strain (Helman *et al.*, 2003), the transformants were selected on BG-11 plates with 50 µg/ml kanamycin. Complete segregation of the mutation was confirmed by PCR with specific forward primer 5′-AGCCAAACCGCATCTAACG-3′ and reverse primer 5′-TTCATAACGCCTCCGGTC-3′. The expected PCR fragment length was 0.7 kb for the WT and 1.7 kb for the mutant (Supplementary Fig. S1). The mutation in the *flv3* open reading frame was confirmed by using forward primer 5′-GTACGGCATGTTCACTACC-3′ and reverse primer 5′- GATCAGCGTTAGCCTGTAA-3′, and the expected PCR fragment length was 1.0 kb for the WT and 2.6 kb for the mutant (Supplementary Fig. S1). Two separate clones of Δ*pgr5* were used interchangeably for the experiments, as no difference was detected in their phenotypes.

Electrochromic shift (ECS) measurements

Individual *Arabidopsis* leaves were measured with the Dual-PAM 100 spectrophotometer (Walz) and its P515/535 accessory module (Schreiber & Klughammer, 2008) as described earlier (Nikkanen *et al.*, 2018; Guinea Diaz *et al.*, 2020). For the experiment in Figure 1, dark-adapted (30 min) leaves were exposed to a light regime fluctuating between 1 min periods of 50 and 825 µmol photons m^{-2} s^{-1}, with 250 ms dark intervals administered every 5 seconds for the first 30 seconds and every 10 seconds thereafter. For the methyl viologen (MV) experiment, detached leaves were dark-incubated overnight on water with 0.05% Tween-20 ± 1 µM MV. In the ECS measurement, dark intervals of 450 ms were administered every 4 s until 28 s during 95 µmol photons m^{-2}s^{-1} of actinic illumination, every 10 s between 28 and 58 s, every 30 s between 58 and 298 s, every 4 s after transition to 825 µmol photons m^{-2}s^{-1} between 298 and 318 s, and every 30 s thereafter. 2,000 Hz pulse frequency of the measuring light was used. ΔECS_T values were normalised to the ΔECS measured upon a 20 µs single-turnover saturating flash administered with the Dual-PAM 100 on dark-adapted leaves (ΔECS_{ST}) (Mathiot & Alric, 2021). For the NEM experiment, detached leaves were incubated in dark on water with 0.05% Tween-20 ±0.1 mM NEM for 2 hours before the experiment. *Chlamydomonas* cultures and *S. viridis* leaves were measured using the same protocol as *Arabidopsis*, with *Chlamydomonas* chlorophyll concentration set to 20 Chl µg ml^{-1}.

ECS changes were measured from *Synechocystis* and *Synechococcus* cells by monitoring the 500-480

nm absorbance difference signal (Viola *et al.*, 2019) using a JTS-10 spectrophotometer (BioLogic) as

described in (Nikkanen *et al.*, 2020). Cells were pelleted and resuspended in fresh BG-11 at 7.5 µg

Chl ml^{-1}. When appropriate, 0.1 mM NEM was added to the samples before a 5 min dark-adaption

prior to the measurement. Each measurement was also conducted with the cells illuminated only by

the measuring light flashes, and the ECS changes recorded were then subtracted from the

measurements under green actinic light, in order to control for any actinic effects of the measuring

light. A moderate (60 µmol photons m^{-2} s^{-1}, relative to growth light of 50 µmol photons m^{-2} s^{-1}) rather

than a low actinic light intensity was used in fluctuating light to allow a sufficient signal-to-noise ratio

for dark-interval relaxation kinetics (DIRK) analysis. The cyanobacterial ECS signals were not

normalised to ECS$_{ST}$ due to a low signal-to-noise ratio of single turnover traces. Instead, a constant

chlorophyll concentration was used.

The *pmf* and g$_{H}$+ parameters were determined as the difference between magnitude of the ECS signal

in light – y0 value of a first-order fit to the decay kinetics of the signal during a dark interval, and the

inverse of the time constant of the fit, respectively, while v$_{H}$+ was derived as *pmf* x g$_{H}$+ (Cruz *et al.*,

2005). Fitting and data processing was done with the OriginPro 2024b (OriginLab) software.

Saturating pulse analysis of chlorophyll fluorescence and P700^{+}

The extents of the donor side and acceptor side limitations of PSI, Y(ND) and Y(NA), respectively,

were determined under fluctuating light conditions by a saturating pulse analysis of the P700^{+} signal

using the Dual-PAM 100 (Klughammer & Schreiber, 2008). Chlorophyll autofluorescence was

simultaneously recorded to determine the Y(NPQ) parameter (in *Arabidopsis*) as described by

(Schreiber *et al.*, 2008). Saturating pulses (8000 µmol photons m^{-2} s^{-1} for *Arabidopsis* and 3000

µmol photons m^{-2} s^{-1} for *Synechocystis*) were administered every 15 seconds during illumination.

Deconvoluted P700, plastocyanin, and ferredoxin redox changes

The deconvoluted plastocyanin (PC), P700, and ferredoxin redox changes were determined by NIR

absorbance difference spectroscopy using the DUAL-KLAS-NIR spectrophotometer (Walz)

(Klughammer & Schreiber, 2016). CET and LET at PSI in *Synechocystis* were quantified using DIRK

analysis according to (Theune *et al.*, 2021) with modifications. *Synechocystis* cells were pelleted and

adjusted to 15 µg Chl ml^{-1} with fresh BG-11. Cells were pre-illuminated for 2 min with 500 µmol

photons m^{-2} s^{-1}, followed by administration of 100 dark intervals of 20 ms. The 100 decay traces

were averaged and initial slopes of the PC and P700 decay curves were used to calculate J_{PSI}. J_{PSI}-CET

was determined by repeating the same experiment in the presence of 10µM DCMU to block electron

transport from PSII. The NIRMAX script (Klughammer & Schreiber, 2016) modified for

cyanobacteria was used to determine the maximal oxidation of PC and P700 and maximal reduction

of ferredoxin (Fed). The model spectra for the deconvolution of PC, P700, and Fed redox changes in

Synechocystis were measured as described in (Nikkanen *et al.*, 2020). Arabidopsis NIRMAX

measurements of dark-adapted leaves were conducted as described in (Nikkanen *et al.*, 2019).

Reduction kinetics of Fed under far red light were measured after running the NIRMAX script,

illuminating detached leaves for 1 second with intensity 20 far-red light.

Cytochrome *f* redox changes

Detached leaves from 4-weeks-old WT-gl1 and *pgr5-1* Arabidopsis were dark-adapted for 5 min,

after which the cytochrome *f* redox kinetics were measured as time-resolved absorbance changes at

554 nm with a baseline drawn between 546 and 573 nm during and after five seconds of illumination

with 500 µmol photons m^{-2} s^{-1} of green light, using the JTS-10 spectrophotometer and sequential

measurements with appropriate interference filters (10 nm FWHM, Edmund Optics). Scattering

artifacts were minimised by BG39 filters (Schott). The cytochrome *f* post-illumination re-reduction

rates ($k_{f\,red}$ s^{-1}) were obtained by fitting the post-illumination kinetics over 300 ms to a first-order

exponential function.

Protein thiol alkylation and SDS-PAGE

4-week-old *Arabidopsis* WT-Col-0 and *pgr5*-Cas plants were dark-adapted, exposed to a light regime

fluctuating between 1-minute periods of 50 and 800 µmol photons $m^{-2}\square s^{-1}$. Samples (individual

leaves) were taken every 20 seconds and quickly placed in 10 % TCA solution to freeze the thiol

redox state of proteins. TCA precipitation, alkylation of thiols with MALPEG-5000, and SDS-PAGE

was then performed as described in Nikkanen et al., (2016). Briefly, free thiol groups were first

alkylated with NEM. Following alkylation, disulphide bonds in proteins were reduced with

dithiothreitol (DTT), and the resulting thiols were then alkylated using methoxypolyethylene glycol

maleimide Mn 5000 (MAL-PEG, Sigma-Aldrich). This treatment makes the in-vivo oxidised forms of

proteins migrate more slowly than reduced ones in SDS-PAGE. MV-treated WT-Col-0 leaves (see

above) were illuminated in growth conditions for 0, 30, 90, or 180 min before sample collection and

placement in 10% TCA.

Total protein extractions from *Synechocystis* cells were done as described in (Zhang *et al.*, 2009) and

from *N. benthamiana* leaves by grinding the tissue in a buffer with 100 mM Tris-HCl pH 8, 50 mM

EDTA, 250 mM NaCl, and 0.75% SDS, heating for 10 min at 68°C, and centrifuging for 10 min at

14,000 rpm. Protein concentration was determined by the DC Protein Assay Kit (Bio-Rad, based on

the Lowry assay). Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and

immunoblotting were performed as previously described (Nikkanen *et al.*, 2016). 4–15% Mini-

PROTEAN TGX precast gels (Bio-Rad) were used for electrophoresis except for Figure 5, for which

12% polyacrylamide gels with 6M urea were cast. PVDF membranes were probed with antibodies

raised against $CF_1\gamma$ (Agrisera, AS08 312), FBPase (courtesy of Dr M. Sahrawy), AtpB (Agrisera, AS),

Flv3 (Genscript), NTRC (Lepistö *et al.*, 2009), or 2-CysPrx (courtesy of Prof. F. J. Cejudo).

Membrane Inlet Mass Spectrometry (MIMS)

Synechocystis in vivo O_2 and CO_2 gas fluxes were monitored in real-time by membrane inlet mass

spectrometry (MIMS) as described earlier (Mustila *et al.*, 2016), with O_2 uptake distinguished from O_2

evolution by enrichment of the $^{18}O_2$ isotope in the medium. Chlorophyll concentration was set to 10

µg Chl ml^{-1}, 1.5 mM $NaHCO_3$ was added, and cells were dark-adapted for 15 min prior to the

measurements. Gas exchange was monitored under 500 µmol photons m^{-2}s^{-1} illumination for 5 min.

See legend for Fig. 2i for experimental details.

Bimolecular Fluorescence Complementation (BiFC) tests

In planta protein–protein interaction tests by bimolecular fluorescence complementation (BiFC) were

conducted as described earlier (Nikkanen *et al.*, 2016, 2018). *Nicotiana benthamiana* plants were used

for transient expression of fusion constructs via *Agrobacterium tumefaciens* strain GV3101

infiltration. Imaging was done 3 days after infiltration using a Zeiss LSM780 confocal microscope.

Construction of the AtPGR5:YFP-C and AtNdhS:YFP-N expression vectors for YFP-fusion proteins

is described in (Nikkanen *et al.*, 2018) and construction of the $AtCF_1\gamma$:YFP-N and AtTRXx:YFP-C

vectors in (Nikkanen *et al.*, 2016).

In silico modelling

The ClusPro 2.0 protein-protein docking service (Kozakov *et al.*, 2017) was used to perform molecular

docking of the AtPGR5 (Uniprot identifier Q9SL05) / $SoCF_1\gamma$ (P05435) / SoCF1 (P00833)

interaction complex, as well as the AtPGR5 / AtPGRL1a (Q8H112)/ AtTRXm4 (Q9SEU6) complex.

The predicted chloroplast transit peptides (amino acid residues 1–60, 1–60, and 1–82) for AtPGR5,

AtPGRL1a, and AtTRXm4, respectively) were removed from the structures before docking.

The amino acid sequences for *Synechocystis* genes *sll1327, slr1330* and *ssr2016* were retrieved from

the Uniprot database (entries P17253, P26533, and P73358, respectively). AlphaFold2 Multimer

Google Colab notebook (Mirdita *et al.*, 2022) was used to model the interaction complex using 6

recycles and pdb100 template mode, with Amber relaxation of the best model. The model with the

best pTM score was selected for visualisation using the Mol* 3D viewer (Sehnal *et al.*, 2021). The

Spinacia oleracea chloroplast ATP synthase structure (Hahn *et al.*, 2018) was retrieved from the

RCSB PDB database (entry 6FKF), while overlaying with the SoCF1γ/AtPGR5 interaction model as

well as modelling of electrostatic surface charges was performed with ChimeraX (Meng *et al.*, 2023).

Statistical tests

372   ANOVA, Tukey's tests for significance, Student's T-tests, and linear mixed effects modelling

373   (LMEM) were performed using the OriginPro 2024 software and its LMEM App. The LMEM models

374   included $gH\square$ as the dependent variable, *pmf* as a continuous covariate, genotype and light intensity

375   as fixed factors, and biological replicate as a random factor. Significance was assessed using the F-

376   tests provided by the software.

RESULTS

Dynamic changes in thylakoid conductivity are independent of the thiol redox state of $CF_1\gamma$ in plants

In plants and algae, measurement of dark-interval relaxation kinetics (DIRK) of the electrochromic

shift (ECS) signal has been widely used for *in vivo* determination of the magnitude of the *pmf*,

partitioning of the *pmf* between ΔpH and $\Delta\Psi$, as well as the conductivity of the thylakoid membrane,

which is mainly determined by the activity of ATP synthase (Sacksteder & Kramer, 2000; Cruz *et al.*,

2001, 2005; Schreiber & Klughammer, 2008; Bailleul *et al.*, 2010). Recently, a reliable method to

measure ECS has also been developed for cyanobacteria (Viola *et al.*, 2019; Nikkanen *et al.*, 2020;

Santana-Sánchez *et al.*, 2023).

Time-resolved ECS analysis in wild type (WT) *Arabidopsis* revealed that upon the onset of

illumination of dark-adapted leaves, the conductivity of the ATP synthase (g_{H+}) initially decreased for

c.a. 30 s (Fig. 1a), as observed previously (Nikkanen *et al.*, 2018; Guinea Diaz *et al.*, 2020). This initial

decrease in g_{H+} allowed transient build-up of *pmf* (Fig. 1b) and induction of NPQ and photosynthetic

control (as indicated by donor-side limitation of PSI) (Fig. 1d–e) in order to avoid over-reduction of

the electron transport chain when redox-regulated CBC enzymes are not yet activated (i.e. reduced).

After the initial decrease, at c.a. 30 s of illumination, g_{H+} started to increase gradually, resulting in

decrease of *pmf* (Fig. 1a–d). Importantly, these changes in g_{H+} did not correlate with the thiol redox

state of $CF_1\gamma$. $CF_1\gamma$ was rapidly reduced at the onset of illumination, becoming almost fully reduced

after 20 s, and remained reduced thereafter (Fig. 1f). In contrast, the CBC enzyme Fructose-1,6-

bisphosphatase (FBPase) remained mostly oxidised (inactive) for a full minute in low light, and only

became partially reduced when leaves were exposed to HL (Supplementary Fig. S2).

In a light regime where light intensity alternates between periods of low light (LL) and HL,

mimicking natural fluctuations, intricate regulatory dynamics of the *pmf* and g_{H+} could be observed.

During LL phases, g_{H+} increased substantially, which contributed to a rapid decline in *pmf* (Fig. 1a–

b). This upregulation of g_{H+} allowed relaxation of photosynthetic control (as inferred from donor-side

limitation of PSI) and NPQ (Fig. 1e–f), allowing sufficient supply of electrons to stromal reactions.

Similarly to downregulation of g_{H+} during dark-to-light transitions, the upregulation of g_{H+} during LL

happened independently of the thiol redox state of $CF_1\gamma$, which remained fully reduced throughout the

change in light intensity (Fig. 1e).

In contrast to LL phases, g_{H+} was rapidly downregulated in WT leaves during HL phases (Fig. 1a),

enabling sufficient *pmf* build-up (Fig. 1b) for induction of photosynthetic control and NPQ (Fig. 1d–

e). Again, the decrease in g_{H+} did not coincide with oxidation of $CF_1\gamma$, which remained fully reduced

(Fig. 1f).

These results demonstrate that ATP synthase activity undergoes dynamic regulation under fluctuating

light in Arabidopsis, and that this regulation is independent of $CF_1\gamma$ redox state and must occur via

another mechanism.

Thylakoid conductivity is dynamically regulated under fluctuating light also in cyanobacteria

and unicellular algae

When WT *Synechocystis* cells were subjected to sudden fluctuations between moderate light (ML)

and HL, time-resolved ECS-DIRK measurements revealed downregulation of g_{H+} during HL phases

(Fig. 2a), resembling that observed in *Arabidopsis*. As in plants, this downregulation likely occurs to

maintain an adequate *pmf* (Fig. 2b) for induction of photosynthetic control. Accordingly, strong donor

side limitation of PSI (Y(ND)) was measured during HL phases (Fig. 2e) indicative of limitation of

electron transport from Cyt b_6f. In contrast to *Arabidopsis*, g_{H+} remained low during the ML phases of

the experiment (Fig. 2a).

We also probed the *pmf* and g_{H+} dynamics under fluctuating light in the unicellular green alga

Chlamydomonas. While *pmf* underwent similar changes as in plants and in cyanobacteria (Fig. 3b), no

clear downregulation of g_H+ was observed during HL, and during the LL phases g_H+ remained low

(Fig. 3a).

430   PGR5 is needed for downregulation of thylakoid conductivity under strong illumination across

431   photosynthetic organisms

During dark-to-LL transitions, the *Arabidopsis pgr5* mutant plants had a lower *pmf* than WT (Fig.

1B), as shown previously and attributed to lower linear electron transport rates (Nikkanen *et al.,*

2018). In fluctuating light, the *pgr5* mutants maintained a significantly higher gH+ than WT during

HL phases (Fig. 1a), resulting in partial loss of *pmf* (Fig. 1b), in agreement with previous studies

(Avenson *et al.,* 2005; Nikkanen *et al.,* 2018). The mutant plants also exhibited impaired induction of

NPQ (Fig. 1d) and photosynthetic control, resulting in an inability to oxidise P700 (Y(ND), Fig. 1e)

and low PSI quantum yield (Supplementary Fig. S3a) due to high acceptor side limitation

(Supplementary Fig. S3b). Interestingly, and in contrast to their behaviour during HL phases, *pgr5*

mutant plants showed a lower g_H+ and slightly lower *pmf* than WT during LL phases (Fig. 1a–b),

which was due to diminished proton flux (vH+) (Fig. 1c). A small amount of reduced $CF_1\gamma$ was

detected in the dark in *pgr5*, but under the fluctuating light treatment the redox state of $CF_1\gamma$ was

indistinguishable from WT, as $CF_1\gamma$ was rapidly reduced upon illumination and maintained as such

throughout the fluctuating light treatment also in *pgr5* (Fig. 1g).

As it has been proposed that PGR5 deficiency results in impaired Q cycle activity in *Chlamydomonas*

(Buchert *et al.,* 2020), we confirmed that the elevated gH+ values in Arabidopsis *pgr5* are not caused

by diminished charge separation in the Q cycle during the dark intervals. No difference in post-

illumination re-reduction rates of cytochrome *f* was detected between WT and *pgr5* (Supplementary

Fig. S4), indicating that the elevated gH+ in *pgr5* does indeed derive from increased proton

conductivity of the thylakoid membrane.

Next, we sought to elucidate the bottleneck in the electron transport chain that impairs PSI oxidation

in *pgr5*. It was proposed recently that Fed binding to PSI is impaired due to damage of F_A and F_B iron-

sulphur clusters in *pgr5* already under moderate light (Tiwari *et al.,* 2024). Since it has also been

suggested that the rate of Fed oxidation is primarily determined by the linear electron transport rate

(Takagi & Miyake, 2018), we compared the Fed reduction and oxidation kinetics between WT and

pgr5 during and after a 1 s pulse of far-red light to minimise the contribution of linear electron flow

and examine just the ability of PSI to reduce, and acceptors to oxidise Fed. Our results show that both

reduction and re-oxidation kinetics of Fed are slowed down in *pgr5,* while oxidation of PC was

increased and oxidation of P700 remained similar to WT (Supplementary Fig. S5), in agreement with

the proposal by Tiwari et al. (2024).

The *Arabidopsis pgr5-1* mutant used in our experiments has been reported to also contain an

additional mutation in the At2g1724 gene, which slightly exacerbates PSI photoinhibition in the

mutant (Wada *et al.*, 2021). Mutants where an intact At2g1724 has been re-introduced (*pgr5*[hope1] and

pgr5-Cas) are, however, also unable to induce photosynthetic control to oxidise PSI (Wada *et al.*,

2021), and have similarly elevated g_{H+} and diminished *pmf* as the *pgr5-1* mutant (Penzler *et al.*, 2022;

Maekawa *et al.*, 2024). While it is therefore unlikely that the At2g1724 mutation in *pgr5-1* plants

would have a significant effect on the phenotypes examined in the current study, we confirmed this by

repeating the ECS experiment (Fig. 1a–c) also with the *pgr5-Cas* mutant (Penzler *et al.*, 2022).

Results with *pgr5-Cas* plants were similar to the ones with *pgr5-1,* apart from the decrease in *pmf*

during initial dark-to-LL transitions being less prominent in *pgr5-Cas* plants than in *pgr5-1* in

comparison to WT (Supplementary Fig. S6). Moreover, while a diminished proton flux (v_{H+}) in LL

and immediately upon shifts to HL was measured in *pgr5-1* mutants in comparison to WT (Fig. 1c),

this decrease was absent in *pgr5-Cas* plants, which instead exhibited elevated v_{H+} during the first HL

phase (Supplementary Fig. S6c). In order to examine if this elevation of vH+ or changes to gH+ were

linked to altered mitochondrial respiration, we repeated the ECS experiment in the presence of

potassium cyanide (KCN) and salicylhydroxamic acid (SHAM), inhibitors of Complex IV and

alternative oxidase (AOX) in mitochondrial respiration, respectively. Compared to WT, *pgr5*-Cas

plants maintained elevated gH+ also in these conditions, while vH+ and *pmf* are lower than WT in LL

but higher than WT in HL (Supplementary Fig. S7). These results indicate that the elevated gH+ and

vH+ observed in HL in *pgr5* mutants do not derive from increased mitochondrial respiration.

We proceeded to investigate whether lack of SynPgr5 affects regulation of g_{H+} in *Synechocystis*.

Elevated g_{H+} was indeed detected upon exposure of Δ*pgr5* cells to HL after illumination in moderate

light (ML) in a fluctuating light regime (Fig. 2a, d), concomitant with lowered (albeit statistically

insignificantly) *pmf* towards the end of the second HL cycle (Fig. 2b). The increased g_{H+} was not due

to an elevated amount of ATP synthase, as the protein level of AtpB remained unchanged in the

mutant (Supplementary Fig. S8). Despite the elevated g_{H+} and in contrast to AtPGR5-deficient

Arabidopsis, no decrease in PSI quantum yield (Supplementary Fig. S3c), donor-side limitation

(indicative of impairment of photosynthetic control) (Fig. 2e) or increase in acceptor side limitation of

PSI (Fig. 2f) was detected during transitions to HL in *Synechocystis* Δ*pgr5* cells. Interestingly, a

slight, albeit statistically non-significant decrease in donor side limitation of PSI was detected in

Δ*pgr5* upon transitions from HL to LL (Fig. 2e).

The ability of the *Synechocystis* Δ*pgr5* strain to maintain sufficient PSI oxidation for unimpaired

photosynthetic activity may be due to the strong electron sink capacity of the Mehler-like reaction

catalysed by flavodiiron proteins (FDPs) under light (Allahverdiyeva *et al.*, 2013). Therefore, we

tested the PSI oxidation capacity of a Δ*flv3 pgr5* double mutant strain, which is deficient in both

SynPgr5 as well as the Flv1/3 hetero-oligomers (Supplementary Fig. S1) which are mainly

responsible for the strong, rapidly induced Mehler-like reaction (Santana-Sanchez *et al.*, 2019). We

detected slightly but non-significantly aggravated acceptor side limitation of in the double mutant in

comparison to the $\Delta flv3$ single mutant during initial LL-to-HL transitions (Fig. 2f). The additional

effect of Pgr5 deficiency was minor compared to lack of Flv3, and no additional aggravation of

ferredoxin redox kinetics, CO_2 fixation, or O_2 evolution was observed in $\Delta flv3\ pgr5$ in comparison to

$\Delta flv3$ (Supplementary Fig. S9). These results indicate that SynPgr5 may affect PSI oxidation also in

cyanobacteria, but its importance is minor compared to plants, while flavodiiron proteins have a

crucial role. This was also reflected in the growth phenotypes of the strains. In agreement with

previous results (Allahverdiyeva *et al.*, 2013), no impairment of growth was detected between WT and

$\Delta pgr5$ under fluctuating light (1 min LL / 1 min HL), with $\Delta pgr5$ even growing slightly faster than

WT, while additional deficiency of Flv3 in $\Delta pgr5\ flv3$ and $\Delta flv3$ resulted in severe impairment of

growth (Fig. 2g).

The $\Delta flv3$ mutant showed a decreased $g_{H}+$ in the HL phases of fluctuating light (Fig. 2a), which

allowed it to compensate for diminished $v_{H}+$ (Fig. 2c) and maintain WT-level *pmf* (Fig. 2b). The

$\Delta pgr5\ flv3$ double mutant, however, exhibited WT-like $g_{H}+$, $v_{H}+$, and *pmf* (Fig. 2a–c), indicating that

the downregulation of $g_{H}+$ observed in $\Delta flv3$ is also at least partly dependent on SynPgr5.

To clarify whether SynPgr5 is involved in CET in *Synechocystis* as previously concluded (Yeremenko

et al., 2005; Dann & Leister, 2019), we quantified the rates of CET and linear electron transport (LET)

in WT and $\Delta pgr5$ cells using DIRK analysis of near-infrared absorbance difference signals, as

described by (Theune *et al.*, 2021). Our results showed that the CET rate was not decreased in the

deletion mutant (Fig. 2h), indicating that SynPgr5 is not involved in CET, at least in steady state

conditions. It has been previously reported that a *Synechocystis* $\Delta pgr5$ mutant has an accelerated rate

of P700 oxidation under far-red light as measured with the Dual-PAM 100 spectrophotometer (Dann

& Leister, 2019), which can be an indication of decreased CET, although results from such assays

should be interpreted with caution (Fan *et al.*, 2007). In contrast to the previous report, we detected no

difference between WT and $\Delta pgr5$ in far-red-induced P700 oxidation rate when measured as the

deconvoluted P700 signal using the Dual-KLAS-NIR instrument (Supplementary Fig. S10). There

results suggest that SynPgr5 is not involved in CET.

Near-infrared spectroscopy revealed no significant differences in the deconvoluted redox signals for

ferredoxin (Fed), plastocyanin (PC) or P700 between WT and $\Delta pgr5$ cells upon onset of strong

illumination (Supplementary Fig. S11a). Moreover, analysis of *in vivo* O_2 and CO_2 gas exchange by

membrane inlet mass spectrometry (MIMS) showed that oxygen evolution (Supplementary Fig.

S11b), dark respiration rate (Supplementary Fig. S11c), light-induced O_2 uptake rate indicative of

FDP activity (Supplementary Fig. S11b, d), as well as CO_2 fixation (Supplementary Fig. S11e) were

all unchanged in Δ*pgr5* in comparison to WT cells. When cells were grown under fluctuating light

and O_2 gas exchange was measured under similar conditions, however, a slightly elevated light-

induced O_2 uptake rate was observed in Δ*pgr5,* although the difference was statistically significant

only during the first HL phase (Fig. 2i).

There is no PGR5 orthologue identified in the genome of another cyanobacterial model species,

Synechococcus elongatus PCC 7942 (*Synechococcus*) (Shimakawa *et al.*, 2016). We therefore

proceeded to investigate whether the regulation of g_{H+} in response to fluctuating light conditions in

Synechococcus differs from *Synechocystis*. Interestingly, no major difference was observed in

comparison to the g_{H+} kinetics in *Synechocystis*, with down-regulation of g_{H+} occurring during the

HL phases after an initial increase (Supplementary Fig. S12). This suggests that the downregulation of

g_{H+} in HL is achieved via a different mechanism in *Synechococcus*.

Elevated g_{H+} under HL was also observed in the *Chlamydomonas pgr5* mutant, in comparison to WT

or a CrPGR5 complementation strain (Fig. 3a). Interestingly, this did not result in lower *pmf* under HL

due to elevated proton flux (v_{H+}) (Fig. 3b, c). We confirmed that the potential presence of damaged

PSI centres in *pgr5* (Tiwari et al., 2024) did not result in major overestimation of the *pmf* and vH+

parameters by also plotting the *Chlamydomonas* ECS data normalised only by Chl concentration

(Supplementary Fig. S13). Finally, the increase in g_{H+} during the HL phases of a fluctuating light

regime was also detected in a *pgr5* mutant of the model C_4 grass *S. viridis* (Supplementary Fig. S12).

Similarly to Arabidopsis but unlike *Synechocystis*, both *Chlamydomonas* (Jokel *et al.*, 2018) and *S.*

viridis (Woodford *et al.*, 2024) *pgr5* mutants suffer from an inability to oxidise PSI in HL.

It has been suggested that the elevated gH+ in *pgr5* mutants is simply a consequence of lower *pmf*,

and that when *pmf* is the same, there would be no difference in gH+ between WT and *pgr5* (Kappel *et*

al., 2023). However, this correlation was made based on steady state measurements. When we plotted

the gH+ values from our time-resolved ECS measurements in fluctuating light as a function of *pmf*,

the correlation between the two parameters differed dramatically between WT and *pgr5* mutants in all

species examined in the current study (Fig. 4). Linear mixed effects models (Supplementary Tables

S1–S7) revealed that Arabidopsis *pgr5-1* mutants exhibited a significantly elevated gH□ in HL

compared to WT independently of *pmf* (p=0.001), with an estimated difference of +5.6 gH+ units

across all *pmf* values. Notably, *pmf* was not a significant predictor of gH□ in this condition (p=0.74).

Similarly, gH+ values increased independently of *pmf* in HL also in Arabidopsis *pgr5*-Cas (+6.89,

p=0.001), and in the full dataset in *Synechocystis* Δ*pgr5* (+12.3, p=0.0154), *Chlamydomonas pgr5*

(+5.72, p=0.003, and *S. Viridis pgr5* (+10.7, p=0.010). These results indicate that the effect of PGR5

on gH+ is not an indirect consequence of diminished *pmf* in any of the studied organisms, and

highlight a conserved function of PGR5 in regulation of the proton conductivity of ATP synthase

from cyanobacteria to chloroplasts in algae, C_3, and C_4 plants.

Elucidating the mechanism of redox-dependent inhibition of thylakoid conductivity

We have previously shown that increasing the reducing capacity of the chloroplast thiol redox regulation systems through overexpression of NTRC in *Arabidopsis* intensifies the downregulation of g_H+ upon increases in light intensity (Nikkanen *et al.*, 2018). We therefore proceeded to investigate potential thiol regulation of thylakoid conductivity or ATP synthase activity, as well as the role of PGR5 in it.

Addition of methyl viologen (MV) results in lowered conductivity of the thylakoid membrane, which has been interpreted as prevention of TRX-mediated reduction of the $CF_1\gamma$ disulphide (Kramer & Crofts, 1989). MV facilitates electron transfer from PSI or reduced ferredoxin to O_2 (the Mehler reaction), resulting in generation of ROS (Fujii *et al.*, 1990; Sétif, 2015; Shapiguzov *et al.*, 2019, 2020). However, MV is also able to donate electrons to redox-regulated enzymes such as nitrite reductase in spinach (Hirasawa & Tamura, 1980) and glutathione reductase in yeast (Llobell *et al.*, 1986). To further elucidate the relationship between stromal/cytosolic thiol redox state and PGR5-dependent downregulation of thylakoid conductivity, we investigated the effects of MV on chloroplastic thiol-regulated enzymes and *pmf* in *Arabidopsis.* We first determined the *in vivo* thiol redox states of the chloroplast redox enzymes as well as $AtCF_1\gamma$ in presence or absence of 1 µM MV. Although electron supply to the TRX systems from PSI is limited by MV, addition of MV caused strong reduction of the 2-cysteine peroxiredoxin (2-CysPrx), FBPase, and $CF_1\gamma$ pools even in darkness (Fig. 5a). NTRC, whose redox state is mostly unaffected by light conditions, thereby acting as a buffer system for plastidial redox state (Nikkanen *et al.*, 2018; Zimmer *et al.*, 2021), also became strongly reduced after 90 min of illumination when treated with MV (Fig. 5a).

Next, we investigated the effect of MV on regulation of *pmf* and g_H+ in WT *Arabidopsis* leaves. Upon illumination of dark-adapted leaves, g_H+ was initially significantly higher in MV-treated WT leaves than in control leaves (Fig. 5b), most likely due to dark-reduction of $AtCF_1\gamma$ by MV (Fig. 5a). As illumination continued, g_H+ diminished in MV-treated leaves and remained at a very low level throughout the rest of the experiment (Fig. 5c). This results in very high *pmf* (Fig. 5d) and NPQ (Shapiguzov *et al.*, 2020). In contrast, MV-treatment of *pgr5* mutant leaves caused only a minor decrease in g_H+ and increase in *pmf* during illumination, and in HL g_H+ was in fact higher in MV-treated *pgr5* mutant than in WT control leaves (Fig. 5c–d), suggesting that the MV-mediated g_H+ decrease is dependent on AtPGR5.

We also investigated the effect of perturbing the thiol redox state of the cell on thylakoid conductivity in *Synechocystis*. As moderate concentrations of MV have only minor effects on cyanobacterial

electron transport (Sétif, 2015), with no significant impact on g_H+ observed after addition of 0.5 mM

MV (Supplementary Fig. S15), we opted to test the effect of supplementation with the thiol-alkylating

agent N-ethylmaleimide (NEM). Addition of 0.1 mM NEM to WT *Synechocystis* cultures resulted in

a significant increase in g_H+ after 5 seconds of illumination (Fig. 6a), similarly to results recently

reported for another maleimide compound 2-methylmaleimide (Hubáček *et al.*, 2024). In Δ*pgr5* cells,

only a slight, statistically non-significant NEM-induced increase in g_H+ was observed. In

Synechococcus, which does not have PGR5, NEM did not have a significant effect on g_H+ (Fig. 6b).

Finally, we also tested the effect of NEM on g_H+ in *Arabidopsis* WT and *pgr5* mutant leaves,

repeating the fluctuating light experiment from Figure 1. NEM decreased g_H+ throughout the

experiment in WT (Fig. 6c), albeit not as dramatically as MV treatment. In contrast, *pgr5* mutant

leaves did not respond to NEM treatment during the first LL-HL cycle (Fig. 6c). During subsequent

cycles g_H+ in NEM-treated *pgr5* mutant leaves decreased slightly in comparison to untreated *pgr5*

mutant leaves, while still remaining significantly higher than untreated WT leaves (Supplementary

Fig. S16) in HL.

These results provided evidence that PGR5 is involved in a redox-dependent regulatory mechanism of

ATP synthase activity that is distinct from $F_1\gamma$ redox state. Interestingly, it has been reported that

PGR5 from cucumber (*Cucumis sativus L.*) can interact with the AtpB subunit of the chloroplast ATP

synthase in yeast-two-hybrid tests (Wu *et al.*, 2021). To investigate the ability of PGR5 to directly

interact with ATP synthase *in planta*, we performed bimolecular fluorescence complementation

(BiFC) tests between AtPGR5 and $AtCF_1\gamma$. Indeed, strong YFP fluorescence co-localising with

chlorophyll autofluorescence was detected, indicating interaction between the two proteins in

chloroplast thylakoid membranes (Fig. 7a). As negative controls we performed BiFC tests between

AtPGR5 and the NdhS subunit of the chloroplast NAD(P)H dehydrogenase-like complex, another

major thylakoid protein complex, as well as $AtCF_1\gamma$ and Thioredoxin x, a chloroplast protein similar

in size to AtPGR5 (Supplementary Fig. S17). $AtCF1\gamma$ +AtTRXf1 and AtPGRL1+AtPGR5

interactions were used as positive controls (Supplementary Fig. S18a–b), as both interactions are well

established in previous studies (Nikkanen *et al.*, 2016, 2018). No cleaved YFP fragments were

detected in immunoblots with an antibody against YFP using in total protein extracts from *N.*

benthamiana leaves transiently expressing the $AtCF_1\gamma$:YFP-N fusion protein (Supplementary Fig.

S18c).

We also performed molecular docking of *Spinacia oleracea* $CF_1\gamma$ and $CF_1\square$ (Hahn *et al.*, 2018) with

an available AtPGR5 structural model using the ClusPro 2.0 protein-protein docking service

(Kozakov *et al.*, 2017). As the $CF_1\varepsilon$-subunit likely remains bound to $CF_1\gamma$ even in its non-inhibitory

conformation (Kato *et al.*, 1997), we also included $CF_1\varepsilon$ in our model. The model predicts that PGR5

may slot between the $CF_1\gamma$, $CF_1\alpha$, and $CF_1\beta$ subunits in a way that possibly hinders the rotation of

$CF_1\gamma$ (Fig. 7b). Moreover, an Alphafold2 Multimer model predicts that SynPgr5 can bind to the $F_1\gamma$

subunit of the *Synechocystis* ATP synthase (Fig. 7c). We analysed the SoCF$_1\gamma$ / AtPGR5

(Supplementary Table S8) as well as SynF1γ / SynPgr5 (Supplementary Table S9) interaction

interfaces using PDBePISA. Finally, we aligned the *Synechocystis* F$_1$y sequences against the sequence

from *Synechococcus elongatus* sp. PCC 7942, which does not have an PGR5 ortholog. Out of the

residues mediating predicted hydrogen bonds with SynPgr5, only Pro 2 in the *Synechocystis* F1γ

sequence is not conserved in the *Synechococcus* sequence (Supplementary Table S10). Therefore,

further work is required to fully elucidate the molecular details of the mechanism of PGR5 interaction

with ATP synthase.

DISCUSSION

Substantial adjustments of thylakoid conductivity take place during sudden changes in light intensity

in both chloroplasts and cyanobacteria, in order to control induction of the photoprotective

mechanisms. Results shown herein indicate that this regulation is independent of the thiol redox state

of $CF_1\gamma$ (Fig. 1), but dependent on another redox-based mechanism that involves the PGR5 protein in

a manner that has been conserved in evolution from free-living cyanobacteria to green algae and C_3

plants, as well as C_4 plants.

Does PGR5 have a role in CET?

The PGR5 protein was originally understood to either directly catalyse or facilitate CET around PSI in

plants (Munekage *et al.*, 2002) and cyanobacteria (Yeremenko *et al.*, 2005). Lack of PGR5 caused a

distinct inability to oxidise PSI in chloroplasts, which was interpreted as being caused by failure to

induce photosynthetic control and NPQ due to insufficient acidification of the thylakoid lumen in the

absence of PGR5-mediated CET (Yamamoto & Shikanai, 2019). However, contrasting evidence was

already provided early on showing that ferredoxin-dependent CET does function in the absence of

PGR5 (Nandha *et al.*, 2007), suggesting that the phenotype of the *pgr5* mutant may be caused by

another mechanism.

We showed previously that in *pgr5 Arabidopsis* plants overexpressing the chloroplast NADPH-

dependent thioredoxin reductase (OE-NTRC *pgr5*), *pmf* was restored to a WT-level during LL-to-HL

transitions (Nikkanen *et al.*, 2018). This was insufficient however, to induce photosynthetic control,

resulting in the OE-NTRC *pgr5* plants suffering from almost as severe over-reduction of the electron

transport chain as *pgr5* plants. Concurring results were more recently reported from *Arabidopsis pgr5*

cgl160 double mutants as well as plants expressing Plastid terminal oxidase 2 from *Chlamydomonas*

reinhardtii (CrPTOX2) in the *pgr5* mutant background. Additional deletion of the ATP synthase

assembly factor CGL160 resulted in a diminished amount of ATP synthase, limiting proton

conductivity of the thylakoid membrane and allowing the *pgr5* mutant to maintain a WT-like *pmf*

while still suffering from extreme acceptor side limitation of PSI (Kappel *et al.*, 2023). Similarly, the

introduction of CrPTOX2 in *pgr5* background resulted in restoration of WT-like (ΔpH-dependent)

NPQ, but oxidation of PSI was still as impaired as in *pgr5* mutants (Zhou *et al.*, 2023). These results

suggest that the inability of *pgr5* mutants to induce photosynthetic control in HL is not simply caused

by loss of ΔpH, but the AtPGR5 protein is directly required. It has been suggested that reduction of a

disulphide in the Rieske subunit of Cyt b_6f could upregulate photosynthetic control by causing a

conformational change that allows protonation of the His128 residue at a higher lumenal pH (Hald *et*

al., 2008; Degen & Johnson, 2024). Moreover, ΔpH-independent metabolic induction of

photosynthetic control in *Chlamydomonas* was recently proposed by (Saroussi *et al.*, 2023). It remains

to be investigated how these potential mechanisms may be affected in PGR5-deficient lines, but as

NTRC overexpression results in stronger reduction of chloroplast TRX systems and redox regulated

proteins (Nikkanen *et al.*, 2016), one could expect the activation of photosynthetic control in OE-

NTRC *pgr5* to occur already at a lower ΔpH instead of being impaired.

Instead, the inability of *pgr5* mutants to oxidise PSI despite restoration of *pmf* generation may be

explained by the presence of a partially photoinhibited PSI population with damaged F_A and F_B FeS

clusters, resulting in repulsion of ferredoxin from the thylakoid membrane (Tiwari et al., 2024). This

means that the bottleneck of electron transport in *pgr5* likely forms already at P700 or alternatively, as

suggested by Buchert et al., (2020) for *Chlamydomonas*, at the Cyt b_6f complex, and not at ferredoxin

as would be expected if PGR5 was directly involved in ferredoxin-dependent CET. Accordingly, our

near-infrared spectroscopy measurements showed that both reduction and re-oxidation kinetics of Fed

are delayed in *pgr5* under far-red light while PC was more strongly oxidised (Supplementary Fig. S5).

This indicated that the bottleneck is not in electron transfer onwards from Fed, but the accumulation

of damaged PSI centres would make it appear as if the mechanism of ferredoxin-dependent CET is

impaired. These results suggest that the phenotype of *pgr5* mutants may not be caused by direct

impairment of CET, but it must be noted that we cannot fully exclude that possibility since

impairment of CET and ATP synthase deregulation could both theoretically result in the original FeS

cluster damage. Interestingly, the PSI oxidation phenotype of *pgr5* could be rescued via expression of

Physcomitrium patens flavodiiron proteins in Arabidopsis *pgr5* background (Yamamoto *et al.*, 2016),

which likely provided a strong enough exogenous sink for electrons from ferredoxin to prevent PSI

photodamage.

Finally, as it has been reported that a lack of CrPGR5 resulted in inefficient operation of the Q cycle

(Buchert et al., 2020), it could be that the elevated gH+ values in *pgr5* mutants derive from

diminished electrogenic activity of the Q cycle during the dark intervals where the rate of ECS signal

decay is monitored. However, the post-illumination re-reduction rate of Cyt *f* was unimpaired in

Arabidopsis *pgr5* (Supplementary Fig. S4). Moreover, Buchert et al. only detected a lower Cyt *f* re-

reduction rate in anoxic *Chlamydomonas pgr5* samples. Therefore, it is unlikely that the elevated gH+

values in *pgr5* mutants are caused by impaired Q cycle activity.

The *Synechocystis* Δ*pgr5* mutant showed no impairment of CET (Fig. 2h). Our data is in line with a

study by Miller et al. showing that the antimycin a-sensitive pathway (=PGR5-dependent) does not

contribute to *pmf* formation during dark/light transitions in *Synechocystis*. Instead, the CET-dependent

fraction of *pmf* generation was entirely attributable to the NDH-1 complex (Miller *et al.*, 2021). As the

extra ATP produced by CET is understood to enable more efficient operation of the CBC by adjusting

the ATP:NADPH ratio, the fact that no impairment of CO_2 fixation rate was observed in Δ*pgr5*

(Supplementary Fig. S11e) further indicates that SynPgr5 is not needed for CET in *Synechocystis*. We

propose it functions instead in reversible redox-dependent inhibition of ATP synthase.

Our ECS results both with the *Chlamydomonas pgr5* mutant (Fig. 3c) as well as the *pgr5-Cas*

Arabidopsis mutant (Supplementary Fig. S6c) showed elevated proton flux (v_H+) in HL, which is also

incompatible with a significant role for PGR5 in CET, and may rather indicate enhancement of the

CET rate. Accordingly, CrPGR5 was not detected in CET supercomplexes (Iwai *et al.*, 2010).

Recruitment of FNR to thylakoid membranes was reported to be diminished in the absence of

CrPGR5 (Mosebach *et al.*, 2017), but a later study concluded that FNR remains associated with the

thylakoids at all times (Kramer *et al.*, 2021). It was also recently speculated, based on simultaneous

near-infrared and chlorophyll fluorescence spectroscopy, that the absence of AtPGR5 may in fact

enhance CET activity in *Arabidopsis* (Maekawa *et al.*, 2024). A lower v_H+ in LL that cannot be fully

explained by a lower linear electron transport rate has, however, been reported in *Arabidopsis* (Degen

et al., 2023) and *S. viridis* (Woodford *et al.*, 2024) *pgr5* mutants. Interestingly though, the

Arabidopsis pgr5-Cas mutant showed WT-like v_H+ in LL (Supplementary Fig. S6c), suggesting that

the low v_H+ in the *pgr5-1* mutant (Fig. 1c) may be affected by the additional mutation of the

At2g1724 gene (Wada *et al.*, 2021). Moreover, in the fluctuating light conditions of the current study,

no impairment of vH+ was detected in the *S. viridis pgr5* mutant (Supplementary Fig. S14c). It should

be noted, however, that the presence of damaged PSI centres in *pgr5* mutants would result in a

decrease in the ECS signal induced by a single-turnover flash (ECS_{ST}), which was used to normalise

the ECS values in plant and algal samples. If this decrease in ECS_{ST} is not proportional to a decrease

in the actinic light-induced ECS signal, it may result in overestimation of the *pmf* and vH+

parameters. However, normalisation of the *Chlamydomonas* ECS data (Fig. 3) to [Chl] only did not

eliminate the differences between WT and *pgr5* (Supplementary Fig. S13), suggesting that no major

artefact is caused by potential PSI damage in *pgr5*.

Conserved PGR5-mediated regulatory mechanism of ATP synthase depends on thiol redox

conditions

The $g_{H}+$ values elevated independently of *pmf* in PGR5-deficient *Synechocystis*, *Chlamydomonas*,

Arabidopsis, and *S. viridis* during high-light phases of fluctuating light (Fig. 1–4, Supplementary Fig.

S14) are consistent with PGR5 functioning as a direct or indirect down-regulator of ATP synthase

activity (Tikkanen *et al.*, 2015; Kanazawa *et al.*, 2017; Nikkanen *et al.*, 2018). *In planta* interaction of

AtPGR5 with $AtCF_1\gamma$ in BiFC tests (Fig. 7a) and *in silico* modelling of PGR5-$CF_1\gamma$ interaction

complexes (Fig. 7b–c) suggest that PGR5 may inhibit ATP synthase by directly binding to the $F_1\gamma$

subunit. $F_1\varepsilon$-inhibition likely prevents ATP hydrolysis in dark and LL conditions, when *pmf* is low,

while PGR5 would limit ATP synthase activity in HL, possibly allowed by the non-inhibitory "down"

conformation of $F_1\varepsilon$ (Feniouk *et al.*, 2010), to allow build-up of sufficient ΔpH for induction of

photoprotection.

The mechanism PGR5-dependent inhibition of ATP synthase activity may be regulated by the thiol

redox state of the chloroplast stroma or cell. PGR5 contains a conserved cysteine residue (Munekage

et al., 2002; Yeremenko *et al.*, 2005), which could be target for regulatory thiol modifications. In line

with this hypothesis, AtPGR5 was found able to interact with AtNTRC in an *in planta* BiFC test, and

overexpression of AtNTRC resulted in intensified downregulation of thylakoid proton conductivity

during transitions to HL, but only in the presence of AtPGR5 (Nikkanen *et al.*, 2018). Moreover, using

a *pgr5 ntrc* double mutant of *Arabidopsis*, Naranjo et al. showed that the high NPQ observed in *ntrc*

mutant plants is dependent on the presence of AtPGR5 (Naranjo *et al.*, 2021). *Ntrc* mutant plants have

a diminished AtPGR5 protein level (Nikkanen *et al.*, 2018), most likely to compensate for the

impaired activation of ATP synthase via reduction of the $AtCF_1\gamma$ disulphide (Carrillo *et al.*, 2016;

Nikkanen *et al.*, 2016).

The *pgr5* mutation could, hypothetically, result in the chloroplast thioredoxin systems themselves

becoming more oxidised or reduced, which would affect the redox state and activity of some other

inhibitory factor of ATP synthase. However, no differences were reported between Arabidopsis WT

and *pgr5* in the vivo redox states of redox regulated CBC enzymes FBPase and SBPase, of NADP-

malate dehydrogenase, or of $CF_1\gamma$ (Okegawa & Motohashi, 2020) (Fig 1g). These findings indicate

that altered activity of the thioredoxin systems is not a major factor in the gH+ phenotype of *pgr5*. A

small population of reduced $CF_1\gamma$ was detected in *pgr5* in the dark (Fig. 1g), but this would not affect

the gH+ dynamics under fluctuating light apart from the very first seconds of the dark-to-light

transition.

It has been suggested that a disulphide bridge between AtPGR5 and AtPGRL1 is required for

PGR5/PGRL1-dependent CET in *Arabidopsis* (Hertle *et al.*, 2013). More recently, however, a model

was proposed where the function of AtPGRL1 is to stabilise AtPGR5 (Rühle *et al.*, 2021). Moreover,

Chlamydomonas PGR5 with the conserved cysteine residue mutated to a serine was able to

complement the impairment of CET in the *Chlamydomonas pgr5* mutant (Buchert *et al.*, 2020),

suggesting that the conserved cysteine residue in PGR5 may be involved in a non-CET related

regulatory mechanism.

Transient reduction of AtPGRL1 occurs upon onset of illumination, coinciding with increase in *pmf*,

PSI oxidation, and induction of NPQ (Hertle *et al.*, 2013; Nikkanen *et al.*, 2018; Wolf *et al.*, 2020;

Chaturvedi *et al.*, 2024). TRXm4 forms a disulphide bridge with AtPGRL1, and cleavage of that

disulphide (most likely by another thioredoxin isoform) and transient release of TRXm4 also occur at

dark-to-light transitions (Okegawa & Motohashi, 2020). This has been suggested to indicate activation

of CET (Hertle *et al.*, 2013; Wolf *et al.*, 2020; Okegawa & Motohashi, 2020; Chaturvedi *et al.*, 2024),

but the same results could also be interpreted as reduction of AtPGRL1 affecting the association of

AtPGR5 with AtPGRL1, releasing AtPGR5 to interact with ATP synthase and to downregulate its

activity (Fig. 8). This would limit H^+ efflux from the lumen and cause build-up of *pmf*, and

consequently, induction of NPQ and photosynthetic control. As recently proposed for *S. viridis* (Nix *et*

al., 2024), PGR5 may be bound to PGRL1 homodimers that are held together by an intermolecular

disulphide between the Cys 116 residues of the monomers, possibly maintained by a bound TRXm4.

Once stromal redox regulatory systems become sufficiently reduced, the disulphide would be reduced,

potentially by another TRX isoform, dissolving the PGRL1 dimer and releasing PGR5.

We therefore suggest that the phenotypes of *pgr5* mutants may have been misinterpreted as being

caused by directly impaired CET. For example, it was recently shown that point mutations of

disulphide-forming cysteine residues in AtPGRL1a results in elevated *pmf*, NPQ, and donor-side

limitation of PSI (Chaturvedi *et al.*, 2024). While the authors did not show data on g_{H+}, it is possible

that this phenotype was caused not by increased CET, but by constitutive inhibition of ATP synthase

activity by AtPGR5. Similarly, it has been reported that AtPGR5 can enhance CET if both AtPGRL1

and AtPGRL2, a protein causing degradation of AtPGR5 in the absence of AtPGRL1, are missing

(Rühle et al., 2021). This conclusion was based on higher *pmf* and NPQ in the *pgrl1a pgrl1b pgrl2*

triple mutant compared to *pgrl1* or *pgr5* single mutants. We suggest, however, that the increased *pmf*

was, again, probably mediated or at least contributed to by lowered ATP synthase conductivity.

In the current study, we explored the effects of perturbing the chloroplast or cellular redox state on the

conductivity of ATP synthase. Interestingly, we found that addition of methyl viologen (MV) results,

either directly or e.g. via affecting the TrxL2-dependent oxidative pathway (Yoshida *et al.*, 2018), in

strong reduction of chloroplast enzymes in *Arabidopsis*, including $AtCF_1\gamma$ (Fig. 5a). While we cannot

fully exclude the possibility of the apparent reduction of chloroplast enzymes in presence of MV

being caused by overoxidation of thiols to sulphenic acids (Huang *et al.*, 2019), the elevation of g_{H+} at

the onset of illumination in MV-treated leaves (Fig. 5b) suggests otherwise, at least in the case of

$CF_1\gamma$ in darkness. Moreover, overoxidation would inactivate 2-CysPrxs, preventing their oxidising

effect on other thiol proteins in the chloroplast, (Pérez-Ruiz *et al.*, 2017; Vaseghi *et al.*, 2018). Thus,

we conclude that MV treatment indeed results in strong reduction of chloroplast redox enzymes,

including $AtCF_1\gamma$.

Accordingly, g_{H+} was elevated in dark-adapted MV-treated leaves (Fig. 5b). However, strong

downregulation of g_{H+} during illumination was measured in MV-leaves, but only in the presence of

AtPGR5 (Fig. 5c). Thiol-alkylating agent NEM had a similar albeit less dramatic effect (Fig. 6c).

Interestingly, lack of AtPGR5 made the plants immune to NEM-induced g_{H+} decrease under HL, but

did not prevent the decrease observed under LL phases of the fluctuating light experiment. These

results provide further evidence that AtPGR5 is required for redox-dependent downregulation of g_{H+}

under high irradiance, although it should be noted that we cannot exclude indirect effects of

MV/NEM treatment interfering with metabolic regulation of ATP synthase (Kohzuma *et al.*, 2013).

We hypothesise that in WT *Arabidopsis*, addition of MV likely causes strong reduction of either

AtPGR5/AtPGRL1/AtTRXm4 directly or some associated factor that would, without an MV

treatment, function as an inhibitor of the ATP synthase during sudden increases in light intensity.

Importantly, this inhibitory effect overrides the activating effect of a fully reduced $AtCF_1\gamma$. In the

absence of AtPGR5, the inhibitory mechanism does not function even in presence of MV.

In *Synechocystis*, addition of NEM caused an opposite phenotype to *Arabidopsis*, with g_{H+} being

elevated rather than diminished (Fig. 6a). The NEM-induced g_{H+} increase was diminished in

Synechocystis Δpgr5 and absent in WT *Synechococcus*, a species that does not have a PGR5

orthologue (Shimakawa *et al.*, 2016). Untreated *Synechococcus* did, however, exhibit similar

downregulation of g_{H+} under strong illumination as *Synechocystis* or *Arabidopsis* (Supplementary

Fig. S12). These results suggest that the regulatory mechanisms involved in this response may differ

between organisms in terms of directionality of redox-regulation and involvement of PGR5, with

Synechococcus possibly using an unrelated, convergently evolved analogous mechanism. The

molecular details of these differing responses remain to be elucidated in future studies.

Interestingly, it has been reported that alkylation of cysteine 89 (Cys 130 in the full sequence

containing the transit peptide) in the (*Spinacia oleracea*) $SoCF_1\gamma$ by NEM or crosslinking Cys89 with

Cys322 inhibits photophosphorylation and induces loss of the proton gradient (McCarty & Fagan,

1973; Soteropoulos *et al.*, 1994; Evron & Pick, 1997). This effect was not observed in the dark,

suggesting Cys89 only becomes accessible to modifications in light upon generation of ΔpH (McCarty

839 & Fagan, 1973). It is therefore possible that Cys89 is responsible for the middle band in the $AtCF_1\gamma$

840 mobility shift assays that is also present in the control sample (Fig. 1e), where all thiols accessible in

841 the dark are alkylated with NEM before addition of MAL-PEG. MAL-PEG likely gains access to

842 Cys89 after the subsequent DTT treatment, resulting in the observed mobility shift. The $F_1\gamma$ Cys89

843 residue is conserved in the green lineage, including cyanobacteria (Hisabori *et al.*, 2013), suggesting a

844 functional role. Our computational models of $AtCF_1\gamma$-AtPGR5 and $SynF_1\gamma$-SynPgr5 interaction

845 complexes predict, however, that neither the conserved Cys30 in SynPgr5, or Cys90 in $SynF_1\gamma$ are

846 involved in mediating the protein–protein interactions (Supplementary Tables S1, S2). This is also

847 consistent with the lack of changes in $AtCF_1\gamma$ redox state in fluctuating light (Fig. 1f–g), suggesting

848 that thiol/disulphide exchanges of $F_1\gamma$ are not involved in the mechanism of adjusting g_{H+} in

849 fluctuating light. The predicted binding of SynPgr5 to $SynF_1\gamma$ in our model somewhat resembles the

850 binding of the inhibitory IF_1 protein to mitochondrial ATP synthase, which slots between the $F_1\gamma$, $F_1\alpha$,

851 and $F_1\beta$ subunits (García-Bermúdez & Cuezva, 2016). This could also explain the observed interaction

852 between cucumber PGR5 and $F_1\beta$ (Wu et al., 2021).

853 An alternative or additional mechanism for the regulation of g_{H+} in fluctuating light in plants could

854 function through changes in intercellular inorganic carbon concentration (C_i). C_i increases upon HL to

855 LL transitions as carbon fixation rate suddenly drops (Vialet-Chabrand *et al.*, 2017), and increase in C_i

856 is known to increase g_{H+} via an unknown regulatory pathway (Kohzuma *et al.*, 2013). It is unlikely,

857 however, that the inability of *pgr5* to down-regulate g_{H+} in HL would be caused by an elevated

858 concentration of C_i, because WT-like levels of C_i as well as CO_2 fixation rates were measured under

859 HL in PGR5-deficient rice plants (Nishikawa *et al.*, 2012; Wada *et al.*, 2018). The carbon fixation

860 capacity of *pgr5* plants only goes down as a consequence of prolonged HL exposure and ensuing PSI

861 photoinhibition (Gollan *et al.*, 2017; Lima-Melo *et al.*, 2018). Similarly, we detected no impairment of

862 CO_2 fixation in the *Synechocystis* $\Delta pgr5$ mutant (Supplementary Fig. S11e). Thus, it is also unlikely

863 that the elevated gH+ in PGR5-deficient plants or cyanobacteria would be caused by an excess of P_i

864 due to a hyperactivated CBC. Interestingly, expression of FlvA and FlvB from the moss

865 *Physcomitrium patens* in the Arabidopsis *pgr5* mutant background alleviated the elevated g_{H+}

866 phenotype (Yamamoto & Shikanai, 2020), highlighting the complex relationship between oxidation of

867 PSI and regulation of ATP synthase.

869 Physiological significance of PGR5-mediated downregulation of ATP synthase in *Synechocystis*

870 Lack of SynPgr5 caused no impairment of PSI oxidation or growth despite lowered *pmf* in

871 *Synechocystis* under fluctuating light (Fig. 2). This may be due to prevalence of FDPs as a protective

872 electron sink on the acceptor side of PSI in cyanobacteria (Allahverdiyeva *et al.*, 2013; Santana-

Sanchez *et al.*, 2019). Slightly elevated FDP activity was indeed measured in fluctuating light grown

Δ*pgr5* under fluctuating light (Fig. 2i), which may allow the mutant to maintain sufficient PSI

oxidation in HL despite diminished *pmf* and to grow similarly to WT in fluctuating light (Fig. 2g). We

have recently proposed that dissipation of *pmf* (lower cytosolic pH) causes FDP hetero-oligomers to

associate more strongly with the thylakoid membrane, increasing photoprotective O_2 photoreduction

activity (Eckardt *et al.*, 2024; Nikkanen *et al.*, 2025). In Δ*pgr5*, the decreased *pmf* due to elevated g_H+

in HL (Fig. 2a–b) may therefore cause the slightly increased FDP activity during the HL phases of

fluctuating light conditions (Fig. 2i), at least partly compensating for the loss of photoprotective

capacity via ΔpH-induced photosynthetic control. Even in the absence of Flv1/3-mediated O_2

photoreduction in the Δ*pgr5 flv3* double mutant, however, no significant additional impairment of PSI

oxidation in comparison to Δ*flv3* single mutant was observed (Fig. 2e–f). The lack of any clear effect

on PSI oxidation in Δ*pgr5* is thus most likely explained by PSI/PSII stoichiometry, as *Synechocystis*

harbours 2-to-5-fold more PSI than PSII complexes (Moore & Vermaas, 2024).

Interestingly, there appears to be a regulatory link between FDPs, ATP synthase, and SynPgr5 in

cyanobacteria. During transitions to HL, lack of Flv1/3 results in a loss up to 75% of the proton flux

during the first minute of illumination in *Synechocystis* (Nikkanen *et al.*, 2020) (Fig. 2c) as well as in

the filamentous model cyanobacterium *Anabaena* sp. PCC 7120 (Santana-Sánchez *et al.*, 2023).

However, cyanobacterial cells compensate for this by downregulating g_H+ (Fig. 2a), achieving WT-

like *pmf* levels after the initial stages of HL transitions (Fig. 2b). In in the absence of both Flv1/3 and

SynPgr5 in the Δ*pgr5 flv3* double mutant, however, this downregulation of g_H+ did not occur,

indicating that SynPgr5-mediated inhibition of ATP-synthase is required for the compensatory

mechanism. Interestingly, proton flux also recovered to WT-levels in Δ*pgr5 flv3* (Fig. 2c), suggesting

e.g. increased CET via NDH-1.

It remains to be investigated what the environmental conditions are where SynPgr5 has a

physiologically significant role. Future studies should focus for example on mixotrophic conditions,

where overexpression of SynPgr5 has been shown to inhibit growth (Margulis *et al.*, 2020) and diurnal

rhythms, which present additional challenges for coordination of the bioenergetic pathways and

metabolism in cyanobacteria (Solymosi *et al.*, 2020; Ortega-Martínez *et al.*, 2024).

Conclusion

The activity of chloroplast or cyanobacterial ATP synthase is dynamically regulated during

fluctuations in light intensity. These adjustments are not dependent on the thiol redox state of the $F_1\gamma$

subunit, but instead depend on the PGR5 protein as well as the thiol redox state of the stroma or

cytosol. This mechanism has been conserved in evolution from free-living cyanobacteria to
chloroplasts in algae, C_3 plants as well as C_4 plants. The inability to downregulate ATP synthase
activity under strong illumination is the only phenotype of PGR5-deficient knockout mutants that can
be observed in all representatives of the groups of photosynthetic organisms studied here;
cyanobacterium *Synechocystis* sp. PCC 6803, green alga *Chlamydomonas reinhardtii*, C_3 plant
Arabidopsis thaliana, and C_4 plant *Setaria viridis*. The role of PGR5 is vital for oxidation of PSI in
plants and algae, but not in cyanobacteria. We also demonstrated that the cyanobacterial Pgr5 is not
needed for cyclic electron transport. We suggest that PGR5 directly binds to ATP synthase and
functions as an inhibitory factor in a manner that is dependent on the thiol redox state of the cell or
chloroplast. The observed effect of plant PGR5 on CET is likely an indirect consequence of
misregulation of ATP synthase and, as shown by (Tiwari *et al.*, 2024), increased photoinhibition of
PSI, although a role for PGR5 in regulating CET via Cyt b_6f cannot be excluded (Buchert *et al.*,
2020). Our study lays important groundwork for understanding the regulatory mechanisms of the *pmf*
and their evolution, possibly enabling biotechnologies to modulate *pmf* regulation for enhancement of
photosynthetic efficiency in crop plants or microbial bioproduction platforms.

Supplementary Data

Fig. S1. PCR analysis to verify full segregation of the Δ*pgr5* and Δ*pgr5 flv3* strains.,

Fig. S2. Mobility shift assay of the in vivo thiol redox state of CBC enzyme fructose 1,6
bisphosphatase (FBPase).

Fig. S3. PSI quantum yield in PGR5-mutants of Arabidopsis and *Synechocystis*.

Fig. S4. Cytochrome *f* post-illumination re-reduction rates in Arabidopsis WT and *pgr5-1* mutant
leaves.

Fig. S5. Redox kinetics of PC, P700, and Fed upon a pulse of far-red light in Arabidopsis WT and
pgr5.

Fig. S6. *Pmf* dynamics in the *Arabidopsis thaliana* WT and *pgr5-Cas* mutant in fluctuating light.

Fig. S7. *Pmf* dynamics in the *Arabidopsis thaliana* WT and *pgr5*-Cas mutant in fluctuating light in the
presence of inhibitors of mitochondrial respiration

Fig. S8. Immunodetection of AtpB and Flv3 levels in *Synechocystis* strains.

Fig. S9. *In vivo* redox kinetics of P700, PC, and Fed and O_2 and CO_2 gas exchange in Δ*pgr5 flv3* and
Δ*flv3* strains of *Synechocystis*.

Fig. S10. P700 oxidation under far-red light in WT and Δ*pgr5* strains of *Synechocystis*.

Fig. S11. Photosynthetic phenotype of the *Synechocystis Δpgr5* strain.

Fig. S12. *Pmf* dynamics in *Synechococcus elongatus* sp. PCC 7942 under fluctuating light.

Fig, S13. *Pmf* and vH+ dynamics in *Chlamydomonas reinhardtii* under fluctuating light without
ECS$_{ST}$ normalisation.

Fig. S14. *Pmf* dynamics in the C$_4$ model grass *Setaria viridis*.

Fig. S15. Effect of methyl viologen on thylakoid conductivity in *Synechocystis*.

Fig. S16. Effect of N-ethylmaleimide (NEM) on thylakoid conductivity under fluctuating light in
Arabidopsis.

Fig. S17. Negative controls for the BiFC tests.

Fig. S18. Positive controls for bimolecular fluorescence com

plementation (BiFC) tests.

Table S1. LMEM report table for Arabidopsis WT vs *pgr5-1* (full data).

Table S2. LMEM report table for Arabidopsis WT vs *pgr5-1* (high light data).

Table S3. LMEM report table for Arabidopsis WT vs *pgr5*-Cas (full data).

Table S4. LMEM report table for Arabidopsis WT vs *pgr5*-Cas (high light only).

Table S5. LMEM report table for *Chlamydomonas* WT vs *pgr5* (full data).

Table S6. LMEM report table for *Setaria viridis* WT vs *pgr5-1* (full data).

Table S7. LMEM report table for *Synechocystis* WT vs Δ*pgr5* (full data).

Table S8. Hydrogen Bond Interactions and salt bridges between AtPGR5 and SoCF$_1$γ in the ClusPro
2.0 docking model.

Table S9. Hydrogen Bond Interactions between SynCF$_1$γ and SynPgr5 in the AlphaFold2 model.

Table S10. Alignment of the ATP synthase F1γ amino acid sequences from Synechocystis sp. PCC
6803 and Synechococcus elongatus sp. PCC 7942

Acknowledgements

We thank Drs Martina Jokel and Arjun Tiwari for providing *Chlamydomonas* strains and the
Arabidopsis *pgr5*-Cas line, respectively. Linda Nevala, Mahfuzur Rahman, Iida Turpeinen, and Tapio
Ronkainen are thanked for technical assistance, and the Turku Bioscience CIC Imaging Core for
imaging infrastructure and assistance.

Author Contributions

Conflicts of interest

977 The authors declare no conflicts of interest.

Funding

This work was supported by the Research Council of Finland grants #354876 (to L.N.), #315119 (to
Y.A.), and # 276392 (to E.R.), funding from Jane and Aatos Erkko Foundation (to E.R.), Novo
Nordisk Foundation grants NNF20OC0064371 (PhotoCat, to Y.A.) and NNF22OCOO79717 (Photo-
e-Microbes, to L.T.W.) as well as Australian Research Council 's Discovery Project (DP230100175)
and the Thomas Davies Research Grant for Marine, Soil and Plant Biology from the Australian
Academy of Sciences (to M.E.).

Data Availability

988 The SynPgr5/SynCF1γ AlphaFold2 multimer model is available in the ModelArchive database at
989 https://www.modelarchive.org/doi/10.5452/ma-v9v6q. All relevant data generated in this study are
990 included in this paper and its Supplementary Data.

References

Akiyama K, Ozawa S-I, Takahashi Y, Yoshida K, Suzuki T, Kondo K, Wakabayashi K, Hisabori T. 2023. Two specific domains of the γ subunit of chloroplast F_oF_1 provide redox regulation of the ATP synthesis through conformational changes. *Proceedings of the National Academy of Sciences* **120**.

Allahverdiyeva Y, Mustila H, Ermakova M, Bersanini L, Richaud P, Ajlani G, Battchikova N, Cournac L, Aro E-M. 2013. Flavodiiron proteins Flv1 and Flv3 enable cyanobacterial growth and photosynthesis under fluctuating light. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 4111–4116.

Armbruster U, Carrillo LR, Venema K, Pavlovic L, Schmidtmann E, Kornfeld A, Jahns P, Berry JA, Kramer DM, Jonikas MC. 2014. Ion antiport accelerates photosynthetic acclimation in fluctuating light environments. *Nature Communications* **5**: 5439.

Armbruster U, Correa Galvis V, Kunz HH, Strand DD. 2017. The regulation of the chloroplast proton motive force plays a key role for photosynthesis in fluctuating light. *Current opinion in plant biology* **37**: 56–62.

Avenson TJ, Cruz JA, Kanazawa A, Kramer DM. 2005. Regulating the proton budget of higher plant photosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* **102**: 9709–9713.

Bailleul B, Cardol P, Breyton C, Finazzi G. 2010. Electrochromism: a useful probe to study algal photosynthesis. *Photosynthesis Research* **106**: 179–189.

Balmer Y, Vensel WH, Hurkman WJ, Buchanan BB. 2006. Thioredoxin target proteins in chloroplast thylakoid membranes. *Antioxidants & Redox Signaling* **8**: 1829–1834.

von Bismarck T, Korkmaz K, Ruß J, Skurk K, Kaiser E, Correa Galvis V, Cruz JA, Strand DD, Köhl K, Eirich J, *et al.* 2023. Light acclimation interacts with thylakoid ion transport to govern the dynamics of photosynthesis in Arabidopsis. *New Phytologist* **237**: 160–176.

Buchert F, Bailleul B, Joliot P. 2021. Disentangling chloroplast ATP synthase regulation by proton motive force and thiol modulation in Arabidopsis leaves. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1862**: 148434.

Buchert F, Mosebach L, Gäbelein P, Hippler M. 2020. PGR5 is required for efficient Q cycle in the cytochrome b6f complex during cyclic electron flow. *The Biochemical journal* **477**: 1631–1650.

Carrillo LR, Froehlich JE, Cruz JA, Savage LJ, Kramer DM. 2016. Multi-level regulation of the chloroplast ATP synthase: the chloroplast NADPH thioredoxin reductase C (NTRC) is required for redox modulation specifically under low irradiance. *Plant Journal* **87**: 654–663.

Chaturvedi AK, Dym O, Levin Y, Fluhr R. 2024. PGR5-LIKE PHOTOSYNTHETIC PHENOTYPE1A redox states alleviate photoinhibition during changes in light intensity. *Plant Physiology* **194**: 1059–1074.

Checchetto V, Segalla A, Allorent G, La Rocca N, Leanza L, Giacometti GM, Uozumi N, Finazzi G,
Bergantino E, Szabò I. 2012. Thylakoid potassium channel is required for efficient photosynthesis in
cyanobacteria. *Proceedings of the National Academy of Sciences* **109**: 11043–11048.

Cobley J, Noble A, Bessell R, Guille M, Husi H. 2020. Reversible Thiol Oxidation Inhibits the
Mitochondrial ATP Synthase in Xenopus laevis Oocytes. *Antioxidants* **9**: 215.

Cruz JA, Avenson TJ, Kanazawa A, Takizawa K, Edwards GE, Kramer DM. 2005. Plasticity in light
reactions of photosynthesis for energy production and photoprotection. *Journal of experimental*
botany **56**: 395–406.

Cruz JA, Sacksteder CA, Kanazawa A, Kramer DM. 2001. Contribution of electric field (Delta psi) to
steady-state transthylakoid proton motive force (pmf) in vitro and in vivo. Control of pmf parsing
into Delta psi and Delta pH by ionic strength. *Biochemistry* **40**: 1226–1237.

DalCorso G, Pesaresi P, Masiero S, Aseeva E, Schuenemann D, Finazzi G, Joliot P, Barbato R, Leister
D. 2008. A complex containing PGRL1 and PGR5 is involved in the switch between linear and cyclic
electron flow in Arabidopsis. *Cell* **132**: 273–285.

Dann M, Leister D. 2019. Evidence that cyanobacterial Sll1217 functions analogously to PGRL1 in
enhancing PGR5-dependent cyclic electron flow. *Nature Communications* **10**: 5299.

Davis GA, Kanazawa A, Schoettler MA, Kohzuma K, Froehlich JE, Rutherford AW, Satoh-Cruz M,
Minhas D, Tietz S, Dhingra A, *et al.* 2016. Limitations to photosynthesis by proton motive force-
induced photosystem II photodamage. *Elife* **5**: e16921.

Degen GE, Jackson PJ, Proctor MS, Zoulias N, Casson SA, Johnson MP. 2023. High cyclic electron
transfer via the PGR5 pathway in the absence of photosynthetic control. *Plant Physiology* **192**: 370–
386.

Degen GE, Johnson MP. 2024. Photosynthetic control at the cytochrome *b* 6 *f* complex. *The Plant*
Cell **36**: 4065–4079.

Eckardt NA, Allahverdiyeva Y, Alvarez CE, Büchel C, Burlacot A, Cardona T, Chaloner E, Engel BD,
Grossman AR, Harris D, *et al.* 2024. Lighting the way: Compelling open questions in photosynthesis
research. *The Plant Cell*.

Ermakova M, Woodford R, Fitzpatrick D, Nix SJ, Zwahlen SM, Farquhar GD, von Caemmerer S,
Furbank RT. 2024. Chloroplast NADH dehydrogenase-like complex-mediated cyclic electron flow is
the main electron transport route in <scp>C</scp> $_4$ bundle sheath cells. *New Phytologist* **243**: 2187–
2200.

Evron Y, Pick U. 1997. Modification of Sulfhydryl Groups in the [gamma]-Subunit of Chloroplast-
Coupling Factor 1 Affects the Proton Slip through the ATP Synthase. *Plant Physiology* **115**: 1549–
1555.

Fan D-Y, Nie Q, Hope AB, Hillier W, Pogson BJ, Chow WS. 2007. Quantification of cyclic electron
flow around Photosystem I in spinach leaves during photosynthetic induction. *Photosynthesis*
Research **94**: 347–357.

Feniouk BA, Kato-Yamada Y, Yoshida M, Suzuki T. 2010. Conformational Transitions of Subunit ⬚ in
ATP Synthase from Thermophilic Bacillus PS3. *Biophysical Journal* **98**: 434–442.

Feniouk BA, Suzuki T, Yoshida M. 2006. The role of subunit epsilon in the catalysis and regulation of
FOF1-ATP synthase. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1757**: 326–338.

Foyer CH, Hanke G. 2022. ROS production and signalling in chloroplasts: cornerstones and evolving
concepts. *The Plant Journal* **111**: 642–661.

Fujii T, Yokoyama E, Inoue K, Sakurai H. 1990. The sites of electron donation of Photosystem I to
methyl viologen. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1015**: 41–48.

García-Bermúdez J, Cuezva JM. 2016. The ATPase Inhibitory Factor 1 (IF1): A master regulator of
energy metabolism and of cell survival. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1857**:
1167–1182.

Gollan PJ, Lima-Melo Y, Tiwari A, Tikkanen M, Aro E-M. 2017. Interaction between photosynthetic
electron transport and chloroplast sinks triggers protection and signalling important for plant
productivity. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**.

Guinea Diaz M, Nikkanen L, Himanen K, Toivola J, Rintamäki E. 2020. Two chloroplast thioredoxin
systems differentially modulate photosynthesis in Arabidopsis depending on light intensity and leaf
age. *Plant Journal* **104**.

**Guo J, Nguyen AY, Dai Z, Su D, Gaffrey MJ, Moore RJ, Jacobs JM, Monroe ME, Smith RD, Koppenaal
DW, *et al.* 2014**. Proteome-wide light/dark modulation of thiol oxidation in cyanobacteria revealed
by quantitative site-specific redox proteomics. *Molecular & cellular proteomics⬚: MCP* **13**: 3270–
3285.

Hahn A, Vonck J, Mills DJ, Meier T, Kühlbrandt W. 2018. Structure, mechanism, and regulation of
the chloroplast ATP synthase. *Science* **360**.

Hangarter RP, Good NE. 1982. Energy Thresholds for Atp Synthesis in Chloroplasts. *Biochimica et
biophysica acta* **681**: 397–404.

Helman Y, Tchernov D, Reinhold L, Shibata M, Ogawa T, Schwarz R, Ohad I, Kaplan A. 2003. Genes
encoding a-type flavoproteins are essential for photoreduction of O-2 in cyanobacteria. *Current
Biology* **13**: 230–235.

Herdean A, Nziengui H, Zsiros O, Solymosi K, Garab G, Lundin B, Spetea C. 2016a. The Arabidopsis
Thylakoid Chloride Channel AtCLCe Functions in Chloride Homeostasis and Regulation of
Photosynthetic Electron Transport. *Frontiers in Plant Science* **7**: 115.

**Herdean A, Teardo E, Nilsson AK, Pfeil BE, Johansson ON, Unnep R, Nagy G, Zsiros O, Dana S,
Solymosi K, *et al.* 2016b**. A voltage-dependent chloride channel fine-tunes photosynthesis in plants.
Nature Communications **7**: 11654.

Hertle AP, Blunder T, Wunder T, Pesaresi P, Pribil M, Armbruster U, Leister D. **2013**. PGRL1 Is the
Elusive Ferredoxin-Plastoquinone Reductase in Photosynthetic Cyclic Electron Flow. *Molecular cell*
49: 511–523.

Hirasawa M, Tamura G. **1980**. Ferredoxin-Dependent Nitrite Reductase from Spinach Leaves.
Agricultural and Biological Chemistry **44**: 749–758.

Hisabori T, Sunamura E-I, Kim Y, Konno H. **2013**. The Chloroplast ATP Synthase Features the
Characteristic Redox Regulation Machinery. *Antioxidants & Redox Signaling* **19**: 1846–1854.

Huang J, Willems P, Wei B, Tian C, Ferreira RB, Bodra N, Gache SAM, Wahni K, Liu K, Vertommen D,
et al. **2019**. Mining for protein S-sulfenylation in Arabidopsis uncovers redox-sensitive sites.
Proceedings of the National Academy of Sciences of the United States of America **116**: 21256–21261.

Hubáček M, Wey LT, Kourist R, Malihan-Yap L, Nikkanen L, Allahverdiyeva Y. **2024**. Strong
heterologous electron sink outcompetes alternative electron transport pathways in photosynthesis.
The Plant Journal.

Ilik P, Pavlovic A, Kouril R, Alboresi A, Morosinotto T, Allahverdiyeva Y, Aro E-M, Yamamoto H,
Shikanai T. **2017**. Alternative electron transport mediated by flavodiiron proteins is operational in
organisms from cyanobacteria up to gymnosperms. *New Phytologist* **214**: 967–972.

Imashimizu M, Bernat G, Sunamura E-I, Broekmans M, Konno H, Isato K, Roegner M, Hisabori T.
2011. Regulation of F0F1-ATPase from Synechocystis sp PCC 6803 by gamma and epsilon Subunits Is
Significant for Light/Dark Adaptation. *Journal of Biological Chemistry* **286**: 26595–26602.

Iwai M, Takizawa K, Tokutsu R, Okamuro A, Takahashi Y, Minagawa J. **2010**. Isolation of the elusive
supercomplex that drives cyclic electron flow in photosynthesis. *Nature* **464**: 1210–1213.

Johnson X, Steinbeck J, Dent RM, Takahashi H, Richaud P, Ozawa S-I, Houille-Vernes L, Petroutsos
D, Rappaport F, Grossman AR, *et al.* **2014**. Proton Gradient Regulation 5-Mediated Cyclic Electron
Flow under ATP- or Redox-Limited Conditions: A Study of Δ *ATPase pgr5* and Δ *rbcL pgr5* Mutants in
the Green Alga *Chlamydomonas reinhardtii*   . *Plant Physiology* **165**: 438–452.

Jokel M, Johnson X, Peltier G, Aro E-M, Allahverdiyeva Y. **2018**. Hunting the main player enabling
Chlamydomonas reinhardtii growth under fluctuating light. *Plant Journal* **94**: 822–835.

Joliot P, Johnson GN. **2011**. Regulation of cyclic and linear electron flow in higher plants.
Proceedings of the National Academy of Sciences of the United States of America **108**: 13317–13322.

Kanazawa A, Kramer DM. **2002**. In vivo modulation of nonphotochemical exciton quenching (NPQ)
by regulation of the chloroplast ATP synthase. *Proceedings of the National Academy of Sciences of*
the United States of America **99**: 12789–12794.

Kanazawa A, Ostendorf E, Kohzuma K, Hoh D, Strand DD, Sato-Cruz M, Savage L, Cruz JA, Fisher N,
Froehlich JE, *et al.* **2017**. Chloroplast ATP Synthase Modulation of the Thylakoid Proton Motive
Force: Implications for Photosystem I and Photosystem II Photoprotection. *Frontiers in Plant Science*
8: 719.

Kappel S, Thiele W, Gefen-Treves S, Henze A, Armbruster U, Schöttler MA. 2023. A congestion
downstream of PSI causes the over-reduction of the electron transport chain in *pgr5* independent of
membrane energization. *bioRxiv*.

Kato Y, Matsui T, Tanaka N, Muneyuki E, Hisabori T, Yoshida M. 1997. Thermophilic F1-ATPase Is
Activated without Dissociation of an Endogenous Inhibitor, ε Subunit. *Journal of Biological Chemistry*
272: 24906–24912.

Kiirats O, Cruz JA, Edwards GE, Kramer DM. 2009. Feedback limitation of photosynthesis at high
CO2 acts by modulating the activity of the chloroplast ATP synthase. *Functional Plant Biology* **36**:
893–901.

Klughammer C, Schreiber U. 2008. Saturation Pulse method for assessment of energy conversion in
PSI. *PAM Application Notes* **1**: 11–14.

Klughammer C, Schreiber U. 2016. Deconvolution of ferredoxin, plastocyanin, and P700
transmittance changes in intact leaves with a new type of kinetic LED array spectrophotometer.
Photosynthesis Research **128**: 195–214.

Kohzuma K, Dal Bosco C, Meurer J, Kramer DM. 2013. Light- and Metabolism-related Regulation of
the Chloroplast ATP Synthase Has Distinct Mechanisms and Functions. *Journal of Biological*
Chemistry **288**: 13156–13163.

Kohzuma K, Froehlich JE, Davis GA, Temple JA, Minhas D, Dhingra A, Cruz JA, Kramer DM. 2017.
The Role of Light-Dark Regulation of the Chloroplast ATP Synthase. *Frontiers in Plant Science* **8**: 1248.

Konno H, Nakane T, Yoshida M, Ueoka-Nakanishi H, Hara S, Hisabori T. 2012. Thiol Modulation of
the Chloroplast ATP Synthase is Dependent on the Energization of Thylakoid Membranes. *Plant and*
Cell Physiology **53**: 626–634.

Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, Beglov D, Vajda S. 2017. The ClusPro web
server for protein–protein docking. *Nature Protocols* **12**: 255–278.

Kramer DM, Crofts AR. 1989. Activation of the Chloroplast Atpase Measured by the Electrochromic
Change in Leaves of Intact Plants. *Biochimica et biophysica acta* **976**: 28–41.

Kramer M, Rodriguez-Heredia M, Saccon F, Mosebach L, Twachtmann M, Krieger-Liszkay A, Duffy
C, Knell RJ, Finazzi G, Hanke GT. 2021. Regulation of photosynthetic electron flow on dark to light
transition by ferredoxin:NADP(H) oxidoreductase interactions. *eLife* **10**.

Kramer DM, Wise RR, Frederick JR, Alm DM, Hesketh JD, Ort DR, Crofts AR. 1990. Regulation of
Coupling Factor in Field-Grown Sunflower - a Redox Model Relating Coupling Factor Activity to the
Activities of Other Thioredoxin-Dependent Chloroplast Enzymes. *Photosynthesis Research* **26**: 213–
222.

Krishnan-Schmieden M, Konold PE, Kennis JTM, Pandit A. 2021. The molecular pH-response
mechanism of the plant light-stress sensor PsbS. *Nature Communications* **12**: 2291.

Lepistö A, Kangasjärvi S, Luomala E-M, Brader G, Sipari N, Keränen M, Keinänen M, Rintamäki E.
2009. Chloroplast NADPH-Thioredoxin Reductase Interacts with Photoperiodic Development in
Arabidopsis. *Plant Physiology* **149**: 1261–1276.

Lima-Melo Y, Gollan PJ, Tikkanen M, Silveira JAG, Aro E-M. 2018. Consequences of photosystem-I
damage and repair on photosynthesis and carbon use in Arabidopsis thaliana. *The Plant Journal* **0**.

Lindahl M, Florencio FJ. 2003. Thioredoxin-linked processes in cyanobacteria are as numerous as in
chloroplasts, but targets are different. *Proceedings of the National Academy of Sciences* **100**: 16107–
16112.

Llobell A, Fernandez VM, Lopezbarea J. 1986. Electron-Transfer between Reduced Methyl Viologen
and Oxidized Glutathione - a New Assay of Saccharomyces-Cerevisiae Glutathione-Reductase.
Archives of Biochemistry and Biophysics **250**: 373–381.

Maekawa S, Ohnishi M, Wada S, Ifuku K, Miyake C. 2024. Enhanced Reduction of Ferredoxin in
PGR5-Deficient Mutant of Arabidopsis thaliana Stimulated Ferredoxin-Dependent Cyclic Electron
Flow around Photosystem I. *International Journal of Molecular Sciences* **25**: 2677.

Margulis K, Zer H, Lis H, Schoffman H, Murik O, Shimakawa G, Krieger-Liszkay A, Keren N. 2020.
Over Expression of the Cyanobacterial Pgr5-Homologue Leads to Pseudoreversion in a Gene Coding
for a Putative Esterase in Synechocystis 6803. *Life* **10**: 174.

Mathiot C, Alric J. 2021. Standard units for ElectroChromic Shift measurements in plant biology.
Journal of Experimental Botany **72**: 6467–6473.

McCarty RE, Fagan J. 1973. Light-stimulated incorporation of N-ethylmaleimide into coupling factor
1 in spinach chloroplasts. *Biochemistry* **12**: 1503–1507.

McKinney DW, Buchanan BB, Wolosiuk RA. 1978. Activation of Chloroplast Atpase by Reduced
Thioredoxin. *Phytochemistry* **17**: 794–795.

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023. <scp>UCSF
ChimeraX</scp>⬚: Tools for structure building and analysis. *Protein Science* **32**.

Miller NT, Vaughn MD, Burnap RL. 2021. Electron flow through NDH-1 complexes is the major driver
of cyclic electron flow-dependent proton pumping in cyanobacteria. *Biochimica et Biophysica Acta
(BBA) - Bioenergetics* **1862**: 148354.

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022. ColabFold: making
protein folding accessible to all. *Nature Methods* **19**: 679–682.

Moore V, Vermaas W. 2024. Functional consequences of modification of the photosystem
I/photosystem II ratio in the cyanobacterium *Synechocystis* sp. PCC 6803. *Journal of Bacteriology*
206.

Mosebach L, Heilmann C, Mutoh R, Gäbelein P, Steinbeck J, Happe T, Ikegami T, Hanke G, Kurisu G,
Hippler M. 2017. Association of Ferredoxin:NADP(+) oxidoreductase with the photosynthetic

apparatus modulates electron transfer in Chlamydomonas reinhardtii. *Photosynthesis Research* **134**:
291–306.

Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T. 2002. PGR5 is involved in cyclic
electron flow around photosystem I and is essential for photoprotection in Arabidopsis. *Cell* **110**:
361–371.

Murakami S, Kondo K, Katayama S, Hara S, Sunamura E, Yamashita E, Groth G, Hisabori T. 2018.
Structure of the γ–ε complex of cyanobacterial F1-ATPase reveals a suppression mechanism of the γ
subunit on ATP hydrolysis in phototrophs. *Biochemical Journal* **475**: 2925–2939.

**Mustila H, Paananen P, Battchikova N, Santana-Sanchez A, Muth-Pawlak D, Hagemann M, Aro E-
M, Allahverdiyeva Y. 2016**. The Flavodiiron Protein Flv3 Functions as a Homo-Oligomer During Stress
Acclimation and is Distinct from the Flv1/Flv3 Hetero-Oligomer Specific to the O-2 Photoreduction
Pathway. *Plant and Cell Physiology* **57**: 1468–1483.

Nalin CM, McCarty RE. 1984. Role of a Disulfide Bond in the Gamma-Subunit in Activation of the
Atpase of Chloroplast Coupling Factor-i. *Journal of Biological Chemistry* **259**: 7275–7280.

Nandha B, Finazzi G, Joliot P, Hald S, Johnson GN. 2007. The role of PGR5 in the redox poising of
photosynthetic electron transport. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1767**: 1252–
1259.

Naranjo B, Penzler J-F, Rühle T, Leister D. 2021. NTRC Effects on Non-Photochemical Quenching
Depends on PGR5. *Antioxidants* **10**: 900.

Nikkanen L, Guinea Diaz M, Toivola J, Tiwari A, Rintamäki E. 2019. Multilevel regulation of non-
photochemical quenching and state transitions by chloroplast NADPH-dependent thioredoxin
reductase. *Physiologia Plantarum* **166**.

Nikkanen L, Santana Sánchez A, Ermakova M, Rögner M, Cournac L, Allahverdiyeva Y. 2020.
Functional redundancy between flavodiiron proteins and NDH-1 in Synechocystis sp. PCC 6803. *The
Plant Journal⬚: for cell and molecular biology* **103**: 1460–1476.

Nikkanen L, Solymosi D, Jokel M, Allahverdiyeva Y. 2021. Regulatory electron transport pathways of
photosynthesis in cyanobacteria and microalgae: Recent advances and biotechnological prospects.
Physiologia Plantarum **173**: 514–525.

Nikkanen L, Toivola J, Rintamäki E. 2016. Crosstalk between chloroplast thioredoxin systems in
regulation of photosynthesis. *Plant Cell and Environment* **39**: 1691–1705.

Nikkanen L, Toivola J, Trotta A, Guinea Diaz M, Tikkanen M, Aro E-M, Rintamäki E. 2018.
Regulation of cyclic electron flow by chloroplast NADPH-dependent thioredoxin system. *Plant Direct*
2: 1–24.

**Nikkanen L, Vakal S, Hubáček M, Santana-Sánchez A, Konert G, Wang Y, Boehm M, Gutekunst K,
Salminen TA, Allahverdiyeva Y. 2025**. Flavodiiron proteins associate pH-dependently with the
thylakoid membrane for ferredoxin-1-powered O2 photoreduction. *New Phytologist*.

Nishikawa Y, Yamamoto H, Okegawa Y, Wada S, Sato N, Taira Y, Sugimoto K, Makino A, Shikanai T.
2012. PGR5-Dependent Cyclic Electron Transport Around PSI Contributes to the Redox Homeostasis
in Chloroplasts Rather Than CO2 Fixation and Biomass Production in Rice. *Plant and Cell Physiology*
53: 2117–2126.

Nix SJ, Morton CJ, Furbank RT, Ermakova M. 2024. A novel *PGRL1* paralog refined the control of
photoprotection in grasses and facilitated cell specialisation in C$_4$ photosynthesis.

Niyogi KK, Truong TB. **2013**. Evolution of flexible non-photochemical quenching mechanisms that
regulate light harvesting in oxygenic photosynthesis. *Current Opinion in Plant Biology* **16**: 307–314.

Okegawa Y, Motohashi K. **2020**. M-type thioredoxins regulate the PGR5/PGRL1-dependent pathway
by forming a disulfide-linked complex with PGRL1. *Plant Cell* **32**: 3866–3883.

Ortega-Martínez P, Nikkanen L, Wey LT, Florencio FJ, Allahverdiyeva Y, Díaz-Troya S. 2024.
Glycogen synthesis prevents metabolic imbalance and disruption of photosynthetic electron
transport from photosystem <scp>II</scp> during transition to photomixotrophy in *Synechocystis* sp.
<scp>PCC</scp> 6803. *New Phytologist* **243**: 162–179.

Penzler J-F, Marino G, Reiter B, Kleine T, Naranjo B, Leister D. 2022. Commonalities and specialties
in photosynthetic functions of PROTON GRADIENT REGULATION5 variants in Arabidopsis. *Plant*
Physiology **190**: 1866–1882.

Pérez-Ruiz JM, Naranjo B, Ojeda V, Guinea M, Cejudo FJ. 2017. NTRC-dependent redox balance of
2-Cys peroxiredoxins is needed for optimal function of the photosynthetic apparatus. *Proceedings of*
the National Academy of Sciences **114**: 12069–12074.

Rantala S, Lempiäinen T, Gerotto C, Tiwari A, Aro EM, Tikkanen M. 2020. PGR5 and NDH-1 systems
do not function as protective electron acceptors but mitigate the consequences of PSI inhibition.
Biochimica et Biophysica Acta - Bioenergetics **1861**.

Rey P, Henri P, Alric J, Blanchard L, Viola S. 2024. Participation of the stress-responsive CDSP32
thioredoxin in the modulation of chloroplast ATP-synthase activity in *Solanum tuberosum*. *Plant, Cell*
& Environment.

Richter ML, McCarty RE. **1987**. Energy-dependent changes in the conformation of the epsilon
subunit of the chloroplast ATP synthase. *Journal of Biological Chemistry* **262**: 15037–15040.

del Riego G, Casano LM, Martin M, Sabater B. 2006. Multiple phosphorylation sites in the beta
subunit of thylakoid ATP synthase. *Photosynthesis Research* **89**: 11–18.

Rühle T, Dann M, Reiter B, Schünemann D, Naranjo B, Penzler J-F, Kleine T, Leister D. 2021. PGRL2
triggers degradation of PGR5 in the absence of PGRL1. *Nature Communications* **12**: 3941.

Sacksteder CA, Kramer DM. 2000. Dark-interval relaxation kinetics (DIRK) of absorbance changes as
a quantitative probe of steady-state electron transfer. *Photosynthesis Research* **66**: 145–158.

Santana-Sánchez A, Nikkanen L, Werner E, Tóth G, Ermakova M, Kosourov S, Walter J, He M, Aro E,
Allahverdiyeva Y. 2023. Flv3A facilitates O2 photoreduction and affects H2 photoproduction
independently of Flv1A in diazotrophic Anabaena filaments. *New Phytologist* **237**: 126–139.

Santana-Sanchez A, Solymosi D, Mustila H, Bersanini L, Aro E-M, Allahverdiyeva Y. 2019.
Flavodiiron proteins 1-to-4 function in versatile combinations in O-2 photoreduction in
cyanobacteria. *Elife* **8**: e45766.

Schreiber U, Klughammer C. 2008. New accessory for the DUAL-PAM-100: The P515/535 module
and examples of its application. *PAM Application Notes* **1**: 1–10.

Schreiber U, Klughammer C, Walz H. 2008. *Complementary PS II quantum yields calculated from*
simple fluorescence parameters measured by PAM fluorometry and the Saturation Pulse method.

Schwarz O, Schürmann P, Strotmann H. 1997. Kinetics and thioredoxin specificity of thiol
modulation of the chloroplast H+-ATPase. *Journal of Biological Chemistry* **272**: 16924–16927.

Sehnal D, Bittrich S, Deshpande M, Svobodová R, Berka K, Bazgier V, Velankar S, Burley SK, Koča J,
Rose AS. 2021. Mol* Viewer: modern web app for 3D visualization and analysis of large biomolecular
structures. *Nucleic Acids Research* **49**: W431–W437.

Sekiguchi T, Yoshida K, Okegawa Y, Motohashi K, Wakabayashi K, Hisabori T. 2020. Chloroplast ATP
synthase is reduced by both f-type and m-type thioredoxins. *Biochimica et Biophysica Acta (BBA) -*
Bioenergetics **1861**: 148261.

Sekiguchi T, Yoshida K, Wakabayashi K-I, Hisabori T. 2022. Dissipation of the proton electrochemical
gradient in chloroplasts promotes the oxidation of ATP synthase by thioredoxin-like proteins. *Journal*
of Biological Chemistry **298**: 102541.

Sekiguchi T, Yoshida K, Wakabayashi K, Hisabori T. 2024. Proton gradient across the chloroplast
thylakoid membrane governs the redox regulatory function of ATP synthase. *Journal of Biological*
Chemistry **300**: 107659.

Sétif P. 2015. Electron-transfer kinetics in cyanobacterial cells: Methyl viologen is a poor inhibitor of
linear electron flow. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1847**: 212–222.

Shapiguzov A, Nikkanen L, Fitzpatrick D, Vainonen JP, Gossens R, Alseekh S, Aarabi F, Tiwari A,
Blokhina O, Panzarová K, *et al*. 2020. Dissecting the interaction of photosynthetic electron transfer
with mitochondrial signalling and hypoxic response in the Arabidopsis rcd1 mutant. *Philosophical*
Transactions of the Royal Society B: Biological Sciences **375**.

Shapiguzov A, Vainonen JP, Hunter K, Tossavainen H, Tiwari A, Järvi S, Hellman M, Aarabi F,
Alseekh S, Wybouw B, *et al*. 2019. Arabidopsis RCD1 coordinates chloroplast and mitochondrial
functions through interaction with ANAC transcription factors. *eLife* **8**.

Shimakawa G, Shaku K, Miyake C. 2016. Oxidation of P700 in Photosystem I Is Essential for the
Growth of Cyanobacteria. *Plant Physiology* **172**: 1443–1450.

Solymosi D, Nikkanen L, Muth-Pawlak D, Fitzpatrick D, Vasudevan R, Howe CJ, Lea-Smith DJ,
Allahverdiyeva Y. **2020**. Cytochrome cM decreases photosynthesis under photomixotrophy in
synechocystis sp. PCC 6803. *Plant Physiology* **183**.

Song K, Baumgartner D, Hagemann M, Muro-Pastor AM, Maaß S, Becher D, Hess WR. **2022**. AtpΘ is
an inhibitor of F0F1 ATP synthase to arrest ATP hydrolysis during low-energy conditions in
cyanobacteria. *Current Biology* **32**: 136-148.e5.

Soteropoulos P, Ong AM, McCarty RE. **1994**. Alkylation of cysteine 89 of the gamma subunit of
chloroplast coupling factor 1 with N-ethylmaleimide alters nucleotide interactions. *Journal of*
Biological Chemistry **269**: 19810–19816.

Spetea C, Herdean A, Allorent G, Carraretto L, Finazzi G, Szabo I. **2017**. An update on the regulation
of photosynthesis by thylakoid ion channels and transporters in Arabidopsis. *Physiologia Plantarum*
161: 16–27.

Sunamura E-I, Konno H, Imashimizu-Kobayashi M, Sugano Y, Hisabori T. **2010**. Physiological Impact
of Intrinsic ADP Inhibition of Cyanobacterial FoF1 Conferred by the Inherent Sequence Inserted into
the γ Subunit. *Plant and Cell Physiology* **51**: 855–865.

Suorsa M, Jarvi S, Grieco M, Nurmi M, Pietrzykowska M, Rantala M, Kangasjarvi S, Paakkarinen V,
Tikkanen M, Jansson S, *et al.* **2012**. PROTON GRADIENT REGULATION5 Is Essential for Proper
Acclimation of Arabidopsis Photosystem I to Naturally and Artificially Fluctuating Light Conditions.
Plant Cell **24**: 2934–2948.

Takagi D, Amako K, Hashiguchi M, Fukaki H, Ishizaki K, Goh T, Fukao Y, Sano R, Kurata T, Demura T,
et al. **2017**. Chloroplastic ATP synthase builds up a proton motive force preventing production of
reactive oxygen species in photosystem I. *Plant Journal* **91**: 306–324.

Takagi D, Miyake C. **2018**. PROTON GRADIENT REGULATION 5 supports linear electron flow to
oxidize photosystem I. *Physiologia Plantarum* **164**: 337–348.

Theune ML, Hildebrandt S, Steffen-Heins A, Bilger W, Gutekunst K, Appel J. **2021**. In-vivo
quantification of electron flow through photosystem I – Cyclic electron transport makes up about
35% in a cyanobacterium. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1862**: 148353.

Tikhonov AN, Khomutov GB, Ruuge EK, Blumenfeld LA. **1981**. Electron transport control in
chloroplasts. Effects of photosynthetic control monitored by the intrathylakoid pH. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics **637**: 321–333.

Tikkanen M, Aro EM. **2014**. Integrative regulatory network of plant thylakoid energy transduction.
Trends in plant science **19**: 10–17.

Tikkanen M, Grieco M, Kangasjarvi S, Aro E-M. **2010**. Thylakoid Protein Phosphorylation in Higher
Plant Chloroplasts Optimizes Electron Transfer under Fluctuating Light. *Plant Physiology* **152**: 723–
735.

Tikkanen M, Rantala S, Aro E-M. **2015**. Electron flow from PSII to PSI under high light is controlled by
PGR5 but not by PSBS. *Frontiers in Plant Science* **6**: 521.

Tiwari A, Mamedov F, Fitzpatrick D, Gunell S, Tikkanen M, Aro E-M. 2024. Differential FeS cluster
photodamage plays a critical role in regulating excess electron flow through photosystem I. *Nature*
Plants.

Tiwari A, Mamedov F, Grieco M, Suorsa M, Jajoo A, Styring S, Tikkanen M, Aro E-M. 2016.
Photodamage of iron-sulphur clusters in photosystem I induces non-photochemical energy
dissipation. *Nature Plants* **2**: 16035.

Vaseghi M-J, Chibani K, Telman W, Liebthal MF, Gerken M, Schnitzer H, Mueller SM, Dietz K-J.
2018. The chloroplast 2-cysteine peroxiredoxin functions as thioredoxin oxidase in redox regulation
of chloroplast metabolism (IT Baldwin, C Foyer, and J-P Reichheld, Eds.). *eLife* **7**: e38194.

Vialet-Chabrand S, Matthews JSA, Simkin AJ, Raines CA, Lawson T. 2017. Importance of
Fluctuations in Light on Plant Photosynthetic Acclimation. *Plant Physiology* **173**: 2163–2179.

Viola S, Bailleul B, Yu J, Nixon P, Selles J, Joliot P, Wollman F-A. 2019. Probing the electric field
across thylakoid membranes in cyanobacteria. *Proceedings of the National Academy of Sciences of*
the United States of America **116**: 21900–21906.

Wada S, Amako K, Miyake C. 2021. Identification of a Novel Mutation Exacerbated the PSI
Photoinhibition in pgr5/pgrl1 Mutants; Caution for Overestimation of the Phenotypes in Arabidopsis
pgr5-1 Mutant. *Cells* **10**: 2884.

Wada S, Yamamoto H, Suzuki Y, Yamori W, Shikanai T, Makino A. 2018. Flavodiiron Protein
Substitutes for Cyclic Electron Flow without Competing CO2 Assimilation in Rice. *Plant Physiology*
176: 1509–1518.

Wang C, Shikanai T. 2019. Modification of Activity of the Thylakoid H+/K+ Antiporter KEA3 Disturbs
Delta pH-Dependent Regulation of Photosynthesis. *Plant Physiology* **181**: 762–773.

Wang C, Takahashi H, Shikanai T. 2018. PROTON GRADIENT REGULATION 5 contributes to
ferredoxin-dependent cyclic phosphorylation in ruptured chloroplasts. *Biochimica Et Biophysica*
Acta-Bioenergetics **1859**: 1173–1179.

Wang C, Yamamoto H, Shikanai T. 2015. Role of cyclic electron transport around photosystem I in
regulating proton motive force. *Biochimica Et Biophysica Acta-Bioenergetics* **1847**: 931–938.

Wolf B, Isaacson T, Tiwari V, Dangoor I, Mufkadi S, Danon A. 2020. Redox regulation of PGRL1 at
the onset of low light intensity. *The Plant Journal* **103**: 715–725.

Woodford R, Watkins J, Moore M, Nix SJ, Yee S, Chan KX, Pogson BJ, von Caemmerer S, Furbank
RT, Ermakova M. 2024. PGR5 promotes energy-dependent non-photochemical quenching to enable
efficient C4 photosynthesis under fluctuating light. *bioRxiv*: 2024.09.15.613160.

Wu X, Wu J, Wang Y, He M, He M, Liu W, Shu S, Sun J, Guo S. 2021. The key cyclic electron flow
protein PGR5 associates with cytochrome b6f, and its function is partially influenced by the LHCII
state transition. *Horticulture Research* **8**: 55.

Yamamoto H, Shikanai T. **2019**. PGR5-Dependent Cyclic Electron Flow Protects Photosystem I under
Fluctuating Light at Donor and Acceptor Sides. *Plant Physiology*: pp.01343.2018.

Yamamoto H, Shikanai T. **2020**. Does the Arabidopsis *proton gradient regulation5* Mutant Leak
Protons from the Thylakoid Membrane? *Plant Physiology* **184**: 421–427.

Yamamoto H, Takahashi S, Badger MR, Shikanai T. **2016**. Artificial remodelling of alternative
electron flow by flavodiiron proteins in Arabidopsis. *Nature Plants* **2**: 16012.

Yeremenko N, Jeanjean R, Prommeenate P, Krasikov V, Nixon PJ, Vermaas WFJ, Havaux M,
Matthijs HCP. **2005**. Open reading frame ssr2016 is required for antimycin A-sensitive photosystem
I-driven cyclic electron flow in the cyanobacterium Synechocystis sp PCC 6803. *Plant and Cell*
Physiology **46**: 1433–1436.

Yoshida K, Hara A, Sugiura K, Fukaya Y, Hisabori T. **2018**. Thioredoxin-like2/2-Cys peroxiredoxin
redox cascade supports oxidative thiol modulation in chloroplasts. *Proceedings of the National*
Academy of Sciences **115**: E8296–E8304.

Zhang P, Allahverdiyeva Y, Eisenhut M, Aro E-M. **2009**. Flavodiiron Proteins in Oxygenic
Photosynthetic Organisms: Photoprotection of Photosystem II by Flv2 and Flv4 in Synechocystis sp
PCC 6803. *Plos One* **4**: e5331.

Zhou Q, Yamamoto H, Shikanai T. **2023**. Distinct contribution of two cyclic electron transport
pathways to P700 oxidation. *Plant Physiology* **192**: 326–341.

Zimmer D, Swart C, Graf A, Arrivault S, Tillich M, Proost S, Nikoloski Z, Stitt M, Bock R, Mühlhaus T,
***et al.* 2021**. Topology of the redox network during induction of photosynthesis as revealed by time-
resolved proteomics in tobacco. *Science Advances* **7**.

1401

Figure Legends

Figure 1. Effect of PGR5 and CF$_1\gamma$ redox state on *pmf*, thylakoid conductivity, and
photoprotection in fluctuating light in *Arabidopsis*. **(a)** Proton conductivity of the thylakoid
membrane (g$_{H+}$ s^{-1}), **(b)** magnitude of the *pmf*, and **(c)** Thylakoid proton flux (v$_{H+}$ = g$_{H+}$ x *pmf*) in
fluctuating light, as determined from dark-interval relaxation kinetics (DIRK) of the ECS signal. ECS
was measured from dark-adapted WT-gl1 and *pgr5-1* mutant *Arabidopsis* leaves under a light regime
alternating between 1 min periods of low (50 µmol photons m^{-2}s^{-1}) and high light (825 µmol photons
m^{-2}s^{-1}). The values in (a) and (c) are averages from 9 and in (b) 11 biological replicates ± SEM. **(d)**
Quantum yield of non-photochemical quenching (Y(NPQ)) and **(e)** fraction of oxidised P700 (donor
side limitation of PSI, Y(ND)) in a fluctuating light regime supplemented with saturating pulses at 15
s intervals. The values in (d–e) are averages from 3 biological replicates ± SEM. In (a–e) * indicates
statistically significant difference between WT and *pgr5-1* according to a two-sample Student's t-test
(P<0.05). **(f-g)** Mobility shift assays of the *in vivo* thiol redox state of the CF$_1$y subunit of the

chloroplast ATP synthase in WT (f) and in *pgr5-1* (g). Samples were taken from fluctuating light (1

min periods of 50 µmol photons $m^{-2}s^{-1}$, LL, and 825 µmol photons $m^{-2}s^{-1}$, HL. The break between the

140 and 160 sec samples in (f) represents division between two gels. –DTT indicates control sample

where DTT was not added after NEM incubation and no alkylation of thiols by MALPEG should

occur.

1420

Figure 2. *Pmf* **dynamics, PSI oxidation, O₂ fluxes, and growth under fluctuating light in**

Synechocystis. Thylakoid conductivity (g_{H+}) **(a)**, *pmf* **(b),** and proton flux (vH+) **(c)** were determined

by the ECS DIRK method from WT, *Δpgr5, Δpgr5 flv3,* and *Δflv3* cells. The values in a–c are

averages from 3–6 (depending on the quality of the first-order DIRK fit) biological replicates ± SEM,

apart from the second data points of WT and *Δpgr5*, where signal-to-noise ratio allowed only single

replicates to be fitted successfully. * indicates statistically significant differences between WT and

Δpgr5 according to a two-sample Student's t-test (P<0.05). **(d)** Representative ECS decay kinetics in

WT and *Δpgr5* cells during the second-to-last 600 ms dark interval in the experiment. First order

exponential fits are shown in cyan and turquoise. **(e)** Donor (Y(ND)) and **(f)** acceptor side limitation

(Y(NA)) of PSI in WT, *Δpgr5, Δpgr5 flv3*, and *Δflv3* strains of *Synechocystis* during transitions from

low light (25 µmol photons $m^{-2}s^{-1}$, LL) to high light (525 µmol photons $m^{-2}s^{-1}$, HL). Y(NA) and

Y(ND) were determined from P700 redox kinetics using the saturating pulse method on a DUAL-

PAM 100 spectrophotometer. Traces are averages of 3–4 biological replicates ± SEM (ribbons). *

indicates statistically significant difference to WT according to a two-sample Student's t-test

(P<0.05). **(g)** Growth curves (OD 750 nm) of WT, *Δpgr5* and *Δpgr5 flv3* cells under fluctuating light

(5 min at 20 µmol photons $m^{-2}s^{-1}$ to 30 sec at 500 µmol photons $m^{-2}s^{-1}$). The values are averages from

3 biological replicates ± SEM. (**h**) Quantification of LET and CET rates in *Synechocystis* WT and

Δ*pgr5*. Chlorophyll was adjusted to 20 µg Chl ml^{-1} in BG-11 pH 7.5. Using a Dual KLAS-NIR

spectrophotometer, cells were illuminated for 2 min under 500 µmol photons $m^{-2}s^{-1}$ before

administering 100x 40 ms dark intervals 10 s apart to monitor the relaxation kinetics of the P700 and

PC signals. The measurement was then repeated in the presence of 10µM DCMU to inhibit LET.

Values means of four biological replicates ± SD. (**i**) O₂ gas fluxes in fluctuating light in WT and

Δpgr5 as measured by membrane inlet mass spectrometry (MIMS). Cells were grown in fluctuating

light (1 min 25 µmol photons $m^{-2}s^{-1}$ / 1 min 500 µmol photons $m^{-2}s^{-1}$), adjusted to 10 µg Chl ml^{-1}, and

$^{18}O_2$ was added in equilibrium with $^{16}O_2$ in order to distinguish O₂ uptake from O₂ evolution. Cells

were dark-adapted for 10 min, after which real-time gas fluxes were measured by MIMS mimicking

the fluctuating growth light conditions, with a 5 min dark period on either side of three LL-HL cycles.

The values are averages of four biological replicates ± SD. The bar chart shows the maximum O₂

uptake rates + SD during the first HL phase of the experiment (dotted arrow). * Indicates statistically

significant difference between WT and *Δpgr5* according to a two-sample Student's t-test (P<0.05).

1451

Figure 3. *Pmf* dynamics in *Chlamydomonas reinhardtii* under fluctuating light. (a) Thylakoid conductivity (g_{H+}). **(b)** *pmf* ($\Delta ECS/ECS_{ST}$). **(c)** Proton flux (v_{H+}). The ECS signal was measured from WT137c, *pgr5* knockout mutant, and PGR5 complementation line (PGR5C) cells under a fluctuating light regime, alternating between 1 min periods of low light (50 µmol photons $m^{-2}s^{-1}$) and high light 825 (µmol photons $m^{-2}s^{-1}$). Values are means ± SEM from three biological replicates. The asterisks indicate statistically significant differences between WT and *pgr5* according to a two-sample Student's t-tests (P<0.05).

1459

Figure 4. Correlation of *pmf* with gH+ in WT and PGR5-deficient mutants across photosynthetic organisms. Time-resolved ECS datasets from fluctuating light conditions. **(a)** *Arabidopsis thaliana* dataset from Fig. 1a–c. **(b)** *Arabidopsis thaliana* dataset from Supplementary Fig. S4. **(c)** *Setaria viridis* dataset from Supplementary Fig. S12. **(d)** *Chlamydomonas reinhardtii* dataset from Fig. 3. **(e)** *Synechocystis* sp. PCC 6804 dataset from Fig. 2a–c. (f) *Synechococcus elongatus* sp. PCC 7942 dataset from Supplementary Fig. S11.

1466

Figure 5. Effect of methyl viologen (MV) on thiol redox states of chloroplast enzymes and on regulation of *pmf* in *Arabidopsis*. (a) Mobility shift assays of AtCF$_1$y, 2-CysPrxs, NTRC, and FBPase thiol redox state in vivo. WT Arabidopsis leaves were incubated in H_2O with (+) or without (-) 1 µM MV in dark and then illuminated in GL conditions for 180 min, with samples taken from darkness (0) and after 30, 90, and 180 min of illumination. Total protein extracts were labelled with MALPEG as described in Materials and Methods, separated by SDS-PAGE and probed with appropriate antibodies. MALPEG-labelled (in vivo oxidised) forms migrate more slowly than unlabelled forms (in vivo reduced). –DTT indicates control sample where no alkylation of thiols by MALPEG should occur, and * indicates dimeric 2CysPrx. We cannot fully exclude the possibility of the apparent reduction of chloroplast enzymes in presence of MV being caused by overoxidation of thiols to sulphenic acids (Huang et al., 2019), especially in the case of 2-CysPrxs (Puerto-Galan et al., 2013; Cerveau et al., 2016). S-sulphenylated proteins cannot form disulphides and thus cannot be labelled by the alkylating agent in our protocol, migrating similarly to fully reduced proteins in mobility shift assays. Apart from 2-CysPrxs, where the thiols are required for the catalytic activity of the enzyme, overoxidation may in fact result in in a conformation that is functionally similar to the reduced state of the protein. **(b)** Conductivity of the thylakoid membrane (g_{H+}) after 1 s of illumination in growth light (GL, 95 µmol photons $m^{-2}s^{-1}$) in dark-adapted WT and *pgr5* mutant leaves incubated in H_2O with or without 1 µM MV. The values are mean + SD from 4–7 biological replicates. Statistical significance of differences was tested by one-way ANOVA and Tukey's tests

1486 (p<0.05). **(c)** Thylakoid conductivity and **(d)** *pmf* in dark-adapted WT and *pgr5* mutant leaves
1487 incubated in H_2O with or without 1 μM MV during 5 min illumination in GL (95 μmol photons $m^{-2}s^{-1}$)
1488 and consequent increase in light intensity to 825 μmol photons $m^{-2}s^{-1}$ (HL), as determined from dark-
1489 interval relaxation kinetics of the ECS signal. The values in C-D are averages from 4–7 biological
1490 replicates ± SEM (ribbon).

1491

Figure 6. Effect of NEM on thylakoid conductivity under illumination. (a) The effect of 0.1 mM
NEM on gH+ in *Synechocystis* WT and Δ*pgr5* cells. ECS DIRK were measured 5 seconds after the
onset of illumination at 500 μmol photons m^{-2} s^{-1}. The values are averages of 7–11 biological
replicates + SD. Statistical significance of differences was tested by one-wayANOVA and Tukey's
tests for comparison of means (P<0.05**). (b)** The effect of 0.1 mM NEM on gH+ in *Synechococcus*
elongatus sp. PCC 7942 WT cells. ECS DIRK were measured as averaged decay kinetics from 5 dark
intervals spaced 5 seconds apart, with the first DIRK being administered 5 seconds after the onset of
illumination at 500 μmol photons m^{-2} s^{-1}. The values are averages of 3 biological replicates + SD. n.s.
= statistically non-significant difference according to a paired *t*-test (P=0.20). **(c)** gH+ was measured
in *Arabidopsis* WT-gl1 and *pgr5* mutant leaves incubated on water or water + 0.1 mM NEM. The
water samples are the same as in Figure 1, while the +NEM values are averages of 4 biological
replicates ± SEM.

1504

Figure 7. Protein–protein interaction between ATP synthase and PGR5. (a) Bimolecular
fluorescence complementation (BiFC) test for protein-protein interaction between Arabidopsis $CF_1\gamma$
fused to an N-terminal fragment of yellow fluorescent protein ($AtCF_1\gamma$:YFP-N) and Arabidopsis
PGR5 fused to a C-terminal YFP fragment (AtPGR5:YFP-C) in tobacco leaves. The left panel shows
Chlorophyll *a* autofluorescence in purple, the middle panel YFP fluorescence in yellow, and the right
panel a merged image of YFP and chlorophyll images. The scale bar is 10 μm. **(b)** Predicted
interaction complex of *Spinacia oleracea* $CF_1\gamma$ and $CF_1\square$ (Hahn *et al.,* 2018), and AtPGR5 (PDB
identifier Q9SL05) docked using the ClusPro 2.0 protein-protein docking service. The $SoCF_1\gamma$ /
$SoCF_1\square$ / AtPGR5 with the of the Spinach ATP synthase complex ($SoCF_0CF_1$, PDB identifier 6FKF)
superimposed (Hahn *et al.,* 2018). Modelling of electrostatic surface charges of AtPGR5 and the
surrounding parts of $SoCF_1$ is shown in the smaller inlet on the right. **(c)** An Alphafold2 Multimer
prediction of an interaction complex between *Synechocystis* ATP synthase γ subunit (*sll1327*, green),
ATP synthase ε subunit (*slr1330,* yellow*)*, and *Synechocystis* Pgr5 (*ssr2016*, magenta). The top-
ranked model with a pTM score of 0.797 and an ipTM score of 0.702 is shown.

1519

Figure 8. Hypothetical model of PGR5-mediated downregulation of ATP synthase activity in plant chloroplasts. A model of an *Arabidopsis thaliana* 2xPGRL1a/2xPGR5/TRXm4 complex was created with the protein–protein docking service ClusPro 2.0, docking monomeric AlphaFold models with the PDB identifiers Q8H112, Q9SL05, and Q9SEU6. An interaction complex of AtPGR5, $AtCF_1\gamma$, and $AtCF_1\square$ was modelled with AlphaFold3 and overlaid with the from the *Spinacia oleracea* CF_0CF_1 crystal structure (6FKF). **(a)** Under low irradiance when stromal redox regulatory systems are largely oxidised, PGR5 is bound to the PGRL1a/PGR5/TRXm4 complex, possibly held by an intermolecular disulphide bond between the Cys116 residues in the PGRL1a monomers. This allows ATP synthase to maintain high activity, effectively dissipating the proton gradient over the thylakoid membrane**. (b)** Under high irradiance, stromal redox regulatory systems become reduced, resulting in reduction of the PGRL1a-dimer disulphide and release of PGR5 to bind to ATP synthase. This inhibits the activity of ATP synthase, lowering thylakoid conductivity to protons, buildup of high *pmf* and induction of photoprotective mechanisms.

(a)

CF$_1$γ

	MV	min

2-CysPrx

NTRC

FBPase

Ox. 2
*
Ox. 1
Red.

Ox. 3
Ox. 2
Ox. 1
Red.

(b)

(c)

(d)

(a)

AtCF$_1\gamma$:YFP-N + AtPGR5:YFP-C

Chl *a*

YFP

Merged

10 μm

10 μm

10 μm

(b)

SoCF$_1\gamma$

AtPgr5

SoCF$_1\varepsilon$

– +

(c)

pTM=0.797

SynPgr5

SynF$_1\gamma$

SynF$_1\varepsilon$

(a)

Thylakoid conductivity / Light intensity

TRXm4
PGRL1a
PGRL1a
PGR5
PGR5

Stroma
Thylakoid
Lumen

H+ H+ H+
H+ H+ H+ H+

(b)

Thylakoid conductivity / Light intensity

TRX?
TRXm4
PGRL1a
PGRL1a
PGR5

Stroma
Thylakoid
Lumen

H+ H+
H+ H+ H+ H+ H+ H+