Title: The Polygenic Score - Rare Variant Causal Pivot: A Conditional Approach to Discovery in Complex Disease Genetics

Authors and Affiliations:

Chad Shaw[1,2,4]*, Christopher Williams[3], Tao Tao Tan[1], Nicholas Di[2], Joshua Shulman[1,4] and John W. Belmont[1,3]

1. Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas
2. Department of Statistics, Rice University, Houston, Texas
3. Genetics & Genomics Services, Inc., Houston, Texas
4. Department of Neurology, Baylor College of Medicine, Houston, Texas

*Correspondence: cashaw@bcm.edu

Abstract

We develop a structural causal methodology to analyze genetic heterogeneity in complex diseases. Our method, the Causal Pivot (CP), exploits one known cause to detect the contribution of a second suspected cause. We employ polygenic scores (PRS) as a known cause; we consider selected rare variation (RV) and RV ensembles as candidate causes. We condition on disease to determine a method that incorporates outcome induced association. We derive a conditional maximum likelihood procedure for both binary and quantitative traits, and we develop a likelihood ratio test (CP-LRT). We demonstrate the CP-LRT's strong power and robustness to misspecification of the RV rate compared to alternatives.

We then use data from the UK Biobank (UKBB) to examine performance. As demonstrations, we consider hypercholesterolemia (HC, LDL-c direct $\geq$ 4.9 mmol/L; n_c=24656), breast cancer (BC, female, ICD10 C50; n_c=12479) and Parkinson's Disease (PD, ICD10 G20, n_c=2940). For PRS we use UKBB data; for RV we select ClinVar pathogenic and likely pathogenic variants in known disease genes. For HC we focus on *LDLR,* for BC we focus on *BRCA1*, and in PD we consider *GBA*. Using CP-LRT we detect RV signals for all three demonstration diseases. Cross-disease analyses serve as comparators. We extend our CP-LRT approach using matching and inverse probability weighting to address potential confounding driven by ancestry. Finally, we present a CP oligogenic load analysis in PD using lysosomal storage. Taken together our work demonstrates the versatility of the PRS-RV CP as a method of inference and discovery in complex disease genetics.

Introduction

Causal heterogeneity, where root causes of disease vary among affected individuals, is a signature characteristic of complex disease. Affected individuals may reach the disease phenotype through distinct etiologies which may be influenced by alternative genetic factors. Although the causes of disease vary, they ultimately lead to a shared pathophysiology or disease state. Moreover, complex disease genetic architectures are known to vary from monogenic to highly polygenic.[1,2] Oligogenic inheritance, environmental phenocopies, and gene X environment interactions are also thought to contribute to complex disease etiologies. This causal heterogeneity impacts both prevention and management because distinct causes may require different approaches for effective interventions or risk mitigation.[3]

Genome wide statistical association studies (GWAS) in complex disease are well developed for both rare and common DNA variants[4,5], but integration across the allele frequency spectrum remains challenging. Moreover, non-familial association approaches are typically case-control designs and may not be applicable to all settings.[6,7] Although GWAS has tools for both rare and common variants, almost all studies are marginal analyses that treat each genetic locus as a separate testing unit; results are summarized with Manhattan plots. The marginal approach avoids confronting heterogeneity with respect to alternative genetic factors differentially present or absent in individuals. Consequently, these methods do not classify individuals into causal groups or subtypes. Resolution of heterogeneity is desirable to avoid overgeneralizing results across different subpopulations and to identify which specific causal mechanism or pathway is potentially responsible in individual cases.[8]

Genetic variants -- while exhibiting allelic heterogeneity, pleiotropy and potential confounding with ancestry and environment -- are explicitly considered to be causal factors that drive downstream disease relevant processes. Consequently, methods developed for causal inference and structural causal modeling (SCM) in fields such as epidemiology and econometrics [9,10] have been adapted to problems in genetics. The most successful example is

Mendelian randomization (MR)[11-14], which uses genetic variants as instrumental variables to

evaluate potential causal biomarkers and endophenotypes for common diseases. We reasoned

causal analysis methods could have broader applications in genetics[15-19] and could offer a

potential approach to reframe inquiry into heterogeneity in complex disease genetics.

We hypothesized that we could adopt SCM as a framework to consider multiple genetic

factors differentially present or absent in individuals to examine the underlying mechanistic

heterogeneity of complex disease. This approach has the goal of deconvolving patient

populations into causal groups or subtypes; it has the additional goal of providing individual level

insights following the diagnostic process where evaluating and eliminating alternative potential

causes is fundamental to personalized medicine.

We develop a statistical approach which we call the Causal Pivot (CP) which can be

applied in cases only or a case-control context. Our CP is based on a structural model of

disease as well as the well-known collider bias concept[20-22]   in which conditioning on an

outcome induces a synthetic correlation between otherwise independent causes. Under these

conditions, if we observe the disease status of individuals and condition on their outcome state

– either case or control —then, within those outcome groups, independent causal factors will

have induced correlation in mathematically well defined patterns. Importantly, model factors will

show no induced correlation if the factors are unconditionally independent and not mutually

causal. We view outcome induced association between distinct causal factors as a feature to

resolve heterogeneity and to drive discovery instead of as a source of bias in the research

process.[23-25]

To develop CP we reasoned that in most research and diagnostic scenarios

investigators have some prior knowledge --   although incomplete -- about causal factors and

the strength of their relationship to outcomes. By exploiting this known information together with

the induced correlation between a known factor and a candidate cause conditional on outcomes

we reasoned it might be possible to pivot from known causal factors to discover novel causes.

Moreover, this approach could also enable discovery in cases-only analysis because the

induced correlation would be present without the need for comparative control individuals.

We proceed from the premise that GWAS has uncovered strong evidence for genetic

loci driving disease and that in aggregate the observed genetic associations represent causal

factors. We hypothesized that we could recruit polygenic risk scores (PRS) as a genome wide

representation of genetic causal contribution to complex disease outcomes. Moreover, because

variants identified in GWAS are frequently common and because PRS are frequently restricted

to common variants above a given minor allele frequency cutoff (i.e. MAF >0.01), we can take

MAF restricted PRS as representative of common genetic variation's contribution to disease

which excludes rare variation (RV, MAF < 0.01). Therefore, we chose to treat the PRS as a CP

to search for new causes by incorporating induced correlation between alternative disease

drivers and PRS conditional on outcomes. We focus attention on RV or RV ensembles as

candidate causes (Figure 1). The general concept that low PRS can be used to prioritize

individuals for RV sequencing or to classify them as having monogenic disorders has been

noted[26,27], and more generally, the use of colliders or v-structures to infer causal graphs is well

known.[28-30] However, more detailed statistical analysis is still needed to elevate and generalize

the procedure, and recognition of the overarching paradigm for PRS - RV causal opposition with

respect to complex traits has been lacking. Importantly, prior methods have been non-

parametric, lacking explicit parameter estimation and lacking distinction between main effects

and interaction effects between PRS and RV. Moreover, non-parametric methods lack interval

estimation to quantify uncertainty in results. Moreover, the lack of statistical formalism has

hampered clarity on performance of the method and its extension to address confounding by

ancestry as well as alternative models such as pathway load.

We developed a highly general statistical framework for the CP using SCM and Bayes

rule. This approach leads naturally to a likelihood framework, maximum likelihood estimation,

confidence intervals, and to a CP likelihood ratio test (CP-LRT). The same mathematical

structure applies in the context of both binary and quantitative traits. We performed power

analyses and compared CP-LRT approach to alternatives [26,27] . Importantly, the CP-LRT

derives power from both the disease rate change in RV as well as the conditionally induced

dependency between RV and PRS. This insight leads to demonstration of the robust properties

of the CP-LRT in terms of sensitivity and specificity. We then demonstrate the performance of

CP-LRT in complex disease examples of breast cancer (BC), hypercholesterolemia (HC) and

Parkinson's disease (PD using the UK Biobank (UKBB). We then extend CP to address

confounded structures for PRS-RV association potentially driven by ancestry. We demonstrate

two approaches to address ancestry confounding using both matching and inverse probability

weighting. Finally, we develop an RV pathway burden test to demonstrate the possible utility of

cases-only CP to investigate oligogenic models for pathway contribution to complex disease

using lysosomal storage in PD as an example. Our results demonstrate the generality and

potential of CP as an approach to causal discovery and to address heterogeneity in complex

disease genetics.

Methods

Causal Models and Graphs

When structural modeling is equipped with a counterfactual interpretation it is also called

causal analysis or structural causal modeling. Our methods are developed through structural

causal modeling, and we adopt this counterfactual interpretation. In our context the

counterfactual assertion is that **but for** the presence or absence of genetic exposures the

potential health outcome of an individual could have been different. In the structural framework

directed acyclic graphs are used to express model assumptions where directed arrows depict

stochastic dependencies and the lack of a directed path between nodes indicates unconditional

independence.

We consider simple models with three or four nodes and two or four edges, respectively

(**Figure 1 a, d** and **Figure 4 a,b,c**). The disease outcome variable is denoted **Y**. This outcome

represents the phenotypic status, for example as recorded in the UKBB metadata by ICD-10

codes or biomarker measurement information. This outcome variable can be binary or

continuous. In our work we focs on two genetic exposures: a PRS -- a continuous variable --

which we will denote as **X** representing the combined causal contribution of common variation

(MAF >0.01), and a binary RV variable denoted as **G** that represents the presence of a strongly

acting RV (**Figure 1**). This binary RV represents the state that an individual is positive for any of

an ensemble of strongly acting RV that in themselves may be vanishingly rare or singleton but

in aggregate can have frequency approaching 0.001 or larger in a population or study sample.

These **X** and **G** are depicted as nodes with edges incident to the **Y** outcome (**Figure 1 a**). Later

we extend this three-node model to a four-node graph (**Figure 4 a**) and we let the PRS and RV

variables be coordinately influenced by ancestry denoted **A** with arrows incident to **X** and **G;**

thereby **A** acts as a common cause in the four node, four edge model.

Causal Pivot

When two causes are incident to the same effect that effect is called a collider for the

two causes. Conditioning on the collider outcome induces an observational – but not causal –

correlation between the distinct causes. This situation is a well-known source of bias in

epidemiologic studies because researchers may unknowingly condition on a collider structure

and report the induced association between variables **X** and **G** as causal.

The CP exploits collider induced correlation as a source of signal rather than noise.

When one causal variable incident to a collider has a **known effect on the outcome** – such as

the effect of the **PRS** on **Y**, then the outcome-conditionally-induced **RV-PRS** correlation can be

used to estimate and to test the causal **Y~RV** relationship. The Null Hypothesis is an absence of

the **Y~RV** relationship, and in this situation no induced correlation will appear between **X** and **G**

conditional on **Y**. As we show in the Mathematical Supplement, the CP is generally applicable

and not limited to genetics.

Conditional Analysis

Here we present an overview of the mathematical derivation of the CP approach; more

details are in the Mathematical Supplement. In what follows we refer to the variables PRS as **X**,

RV as **G** and disease outcome **Y.** The CP derives from manipulation of the probabilistic

factorization determined by the graphical model (**Figure 1A**). The joint density under the model

is:

(1) $$f_\theta(x, g, y) = f_{\theta_R}(y \mid x, g) f_{\theta_X}(x) f_{\theta_G}(g)$$

In the expression $f_\theta(\cdot)$ represents the respective probability density functions for

their arguments, and the bar symbol expresses conditioning on variables to the right-hand side

of the bar. The parameter vector θ represents the unknown parameters governing the

probabilistic structure of the system; for clarity we emphasize that the parameter θ is comprised

of the parameters θ_R that determine the forward conditional relationship between **Y** and **X**,**G** and

the separate parameters θ_{XG} determine the distributions of **X** and **G** – which in the simple model

are exogenous and independent.

The Causal Pivot procedure follows from application of Bayes' rule to the probabilistic

factorization implied by the graphical model. In notation applying Bayes rule to (1):

(2) $$f_\theta(g \mid x, y) = \frac{f_\theta(y \mid x, g)}{E_G(f_\theta(y \mid x, G))} f_\theta(g)$$

For **G** representing RV status, the expression (2) states that the conditional probability

for a sampled individual to bear an RV – denoted by **G=1** – or to not have an RV -- denoted **G**=0

-- is given by the ratio of the forward conditional probability of their phenotype **Y** given PRS **X**

and **G=b** divided by the average probability of **Y** integrating out **G** multiplied against the prior

probability of **G=b**. This factorization is general and does not depend on the model specification

of **f(y | x,g)** or **f(x)** or **f(g)**; moreover, the factorizations in (1) and (2) do not depend on a

univariate restriction on **X**, **G** or **Y**, nor does it depend on the binary character of **G** or the

distributional character of **X**.

Model Specification

It is useful to provide a parametric model in order to perform analyses, make

calculations, enable parametric testing and to perform power comparisons. Non-parametric

approaches also apply, but parametric modeling provides both mathematical insight and yields

practical results such as confidence intervals. We adopt a generalized linear model framework

for the forward distribution of **Y** given **X** and **G.** The general linear model framework is

encompassing of both logistic regression in the case of binary outcomes as well as linear

regression for quantitative traits. The general expression for the forward conditional expectation

of **Y** given **X** and **G** is:

(3) $$h\left(E\left(Y \mid X, G\right)\right) = \alpha + \beta \cdot X + \gamma \cdot G + \eta \cdot X \cdot G$$

Examples for the function h() are the logit or log-odds for binary outcomes, such as for a

binary trait like BC; the identity function can be used for continuous **Y**, such as in the case of

LDL measurements. We assume the parameters α and β are known or estimable from

population data; we also assume knowledge of a baseline frequency of RV, set at ω. In many

cases the estimate of the **X**, **Y** relationship is reported. If not, the easiest way to estimate it is to

apply a forward model to unconditional **Y** including **X** and exclude **G.** Although the α parameter

depends on **G** we argue that it is negligible because **G** is rare. We adopt this approach to

determine α and β in our demonstration work. The frequency of rare variation is estimable from

population tabulation such as GNOMAD or from UKBB summary information.

194   Likelihood Analysis Conditional on Y

195   To examine the properties of the CP we constructed a test procedure for the possible

196   effect of RV using the factorization in (2) to determine a likelihood function for observed data

197   conditional on **Y=a.** We executed this for both binary **Y** and for Y>δ for continuous **Y**. In the

198   case of discrete Y, we have:

$$
\begin{aligned}
L(\gamma, \eta \mid X, G, Y = a) &= \prod_{i=1}^{n} f_\theta(g_i \mid x_i, y_i = a) \\
&= \prod_{i=1}^{n} \frac{f_\theta(y_i = a \mid x_i, g_i)}{E_G(f_\theta(y_i \mid x_i, G))} f(g_i)
\end{aligned}
$$

199   (4)

200   Taking logarithms and emphasizing that the unknown elements of θ τηατ concern the impact of

201   the RV on outcome are γ,η:

$$
\begin{aligned}
l(\gamma, \eta \mid X, G, Y = a) &= \sum_{i=1}^{n} l(\gamma, \eta \mid g_i, x_i, Y_i = a) \\
&= \sum_{i=1}^{n} l(Y_i = a, \gamma, \eta \mid x_i, g_i) - log\left(E_G(f(Y_i = a, \gamma, \eta \mid x_i, G))\right) + l(g_i)
\end{aligned}
$$

(5)

As shown in the Mathematical Supplement this likelihood function can be maximized in

the variables γ and η to produce maximum likelihood estimates; this analysis is performed by

differentiating the log-likelihood with respect to γ and η and solving for the root where the

derivative is zero. Confidence intervals are obtained using Fisher Information as using the

observed means of the squared derivatives of the log-likelihood. We performed this analysis for

both logistic and a liability model for a continuous trait (see Supplemental Materials). For the

continuous trait we consider the circumstance where the observable sample comprises only

individuals with a sufficiently extreme quantitative trait Y>δ where individuals in this class are

deemed cases and individuals not in the class could be deemed controls; we suppose we have

access to the trait values for the cases. Importantly, we note that the CP conditional analysis

can be performed on cases only, controls only, or on both cases and controls taken together but

where the analysis is performed conditionally in both groups. The availability of appropriate

research subjects determines which is pursued. In our investigation we focus on cases only

analyses.

Likelihood Ratio Test

As mentioned, the likelihood approach determines maximum likelihood estimates (MLE)

by solving for the roots of the derivatives of log-likelihood equations treating these as functions

of the unknown parameters. A likelihood ratio test can be constructed by plugging in the

maximum likelihood estimates into the log-likelihood function to compute a test statistic. The

Null Hypothesis is that the parameters γ and η are 0. Under the Null Hypothesis the asymptotic

distribution of : $-2 * l(0,0\ |X,Y,G) - l(\gamma, \eta|X,Y,G)$ is a chi-square on two degrees of freedom. The

procedure considers the influence of the RV – both individually and in its potential interaction

with the PRS. Additional details are provided in the Mathematical Supplement.

Logistic Model

The logistic model considers the disease outcome to be binary where the log-odds of

disease is linear in the explanatory variable. We additionally treat RV as a binary variable. In the

initial treatment there is no additional stochasticity in the conditional distribution of the outcome

Y given **X** and **G** except for the coin-flipping variability of a Bernoulli trial; such extensions are

possible but out of scope in this paper. The log-likelihood analysis conditional on **Y=a** produces

four different sums: the sum of contributions for **Y=0** where **RV=1**, **Y=0** where **RV=0**, **Y=1** where

RV=1, and **Y=1** where **RV=0**. Breaking the likelihood calculation into these four parts is both

computationally practical and useful for analysis. In cases only analysis there is no contribuition

from groups where **Y=0.** The details of the log-likelihood analysis and derivative derivation

appear in the R markdown code and in the Supplement.

Liability Model

The Causal Pivot liability model considers conditioning on the outcome Y>δ for a

quantitative trait that is considered to be conditionally distributed with a mean as in (3) and

where additional exogenous noise is Gaussian. The approach generalizes to non-Guassian

noise, but the Gaussian situation is both canonical and illustrative. Here the model considers the

outcome variable Y | I(Y>δ), x, g. The forward model has a distribution:

$$f(y \mid I_{Y>\delta} = 1, x, g) = \begin{cases} \frac{f(y|x,g)}{1-F(\delta|x,g)} & Y > \delta \\ 0 & Y \le \delta \end{cases}$$

245   (6)

246   For situations where the outcome value is less than δ we have:

$$f(y \mid I_{Y>\delta} = 0, x, g) = \begin{cases} \frac{f(y|x,g)}{F(\delta|x,g)} & y < \delta \\ 0 & y \ge \delta \end{cases}$$

(7)

To perform analysis with this liability model we use the general expression in (2) that

becomes expression (5) after applying the log-likelihood transformation. As with the logistic

model there are four configurations: where the outcome Y< δ and RV=1, where Y<δ and RV=0,

where the Y>δ and RV=1, and where Y>δ and RV=0.

Power Analysis

To perform power analysis, we built a forward simulation procedure to generate outcome

data under both the logistic and liability framework. For each scenario, we simulated RV as a

binary variate with frequency 0.001. For PRS we generated unit normal outcomes. We fixed β

according to values consistent with real data. We let γ and η range across many possible

values. For logistic regression we set α=-2.2 and β =0.5, we set γ and η to values collection of

alternatives. We generated **Y** outcomes according to the inverse transformed log-odds. We then

applied the likelihood expression in (5) to obtain the MLE, and we compute the likelihood ratio

test statistic. We performed 1000 simulations at each model parameterization, and we recorded

outcomes. To determine power, we calculated the proportion of simulation runs where the

observed p-value was less than 0.05. For the liability model we generated outcome variables in

a parallel fashion, first generating X ~N(0,1) and G~ber(0.001) and fixing the parameters α, β

and a grid of alternative **RV** effect sizes. We used the traditional unconditional case-control

forward regression as a comparator for CP power. This analysis considers a logistic regression

for **Y** given **X** and **G** and a linear model for **Y** given **X** and **G** in the quantitative outcome

situation.

Cases Only Analysis and Case Control Conditional Analysis

Our CP procedure can be applied to cases-only data, to controls only data or to case-

control data taken together. In all of these situations, data analysis proceeds according to a

conditional analysis using the fundamental factorization in (2) and to application of the maximum

likelihood procedure in (5) followed by a likelihood ratio test. We focused on comparison of

cases only CP analysis against the traditional forward regression approach and to alternative

cases-only analyses.

Comparator Models and Robustness Analyses

The CP-LRT benefits from the full information in the stochastic system as described by

the structural causal graph and as encapsulated in the conditional likelihood function. This

conditional likelihood gives the expected rate change in G given both X and Y.   We considered

two comparator non-parametric models to our CP-LRT . First, we consider a Z-test procedure

where we assume the frequency of RV status in the source population –deemed ω -- is exactly

known. Under this scenario if the RV status has no effect on outcome, the expected number of

RV in the outcome group **Y=a** would be binomial with expectation $N_{Y=a}\omega$ and variance

$N_{Y=a}\omega(1-\omega)$. We take **Z** to be the difference between the observed RV+ disease individuals

($N_{Y=a,RV+}$) and the expected under H_0 standardizing by square root of variance. This comparator

focuses on the rate change of a functional variant state in the disease population. Alternatively,

prior work proposed a Wilcoxon test on the PRS values among RV+ and RV- individuals in Y=a.

The Wilcoxon is a non-parametric test for a shift in means between two populations, in this case

RV+ and RV- conditional on Y=a.

Robustness Analyses

We consider the impact of the precision with which ω is known. We examine the type 1 error

when **G** has no effect and scrutinize the impact of underspecification of the rate ω in the general

population. The type 1 error is the rate at which the CP-LRT and Z-test reject H_0 and falsely

identify an effect of **G** when there is none. These analyses are performed across a range of

underspecification of the RV rate which we deem ω^*; the true simulation parameter values in

these analyses are α=-2.2, β=0.6,ω=0.001, γ=0,η=0.

We then consider the power impact of overspecifying the RV rate. We examine this

situation in the context where α=-2.2, β=0.6, ω=0.001,γ=2.2, η=0. In this case there is an RV effect, but

oversetting the value of ω^* makes it more difficult to detect. We examine a range of values ω^* up to

and beyond the conditional frequency of RV of 0.0045.

UKBB Data

We analyzed human data from cohorts and controls drawn from the UKB under an

approved application (Project 98786) titled "Genetic Heterogeneity in Diseases with Complex

Inheritance". We used hypercholesterolemia (LDL direct | Instance 0 ≥ 4.9 mmol/L; Data-Field

30780), breast cancer (ICD-10 C50; Data-Field 40006), and Parkinson's Disease (ICD-10 G20;

Data-Field 131022) as disease models with expected positive findings. Cases were filtered

based on inferred European ancestry, availability of exome and genotyping data, female-only for

breast cancer, and availability of a phenotype-specific polygenic risk score. The final samples

included: hypercholesterolemia (HC cases 24656; controls 347889); breast cancer (BC, cases

12479; controls 198311); and Parkinson's (PD, cases 2949; controls 388198). As this study

exclusively utilized de-identified data from the UKBB, it qualifies for exemption from ethical

review under the guidelines for research involving non-identifiable human data.

PRS

We used PRS made available through the UKBB (Field ID 26220 Standard PRS for BC;

26250 Standard PRS for low density lipoprotein cholesterol; and 26260 Standard PRS for PD).[31]

Exome Variant Extraction and Classification

We used PLINK2 to read genotypes from the UKBB exome BGEN files (Data Field

23159) and extract variants within our target gene lists (see bedfile

https://github.com/chadashaw/causal-pivot/) for the disease-defined cohort samples. The

extracted genotype files were converted to Matrix Market format using a custom python script.

Variants were, then, annotated with OpenCRAVAT; annotators included those pulled from the

OpenCRAVAT respository (gnomad3, revel, cadd_exome) as well as custom annotators

(tcg_clinvar, AlphaMissense, ESM1b) produced from publicly available data. We exclude

variants that were absent (homozygous-ref) across all samples.

We focused on three example genes for the demonstration diseases: *BRCA1* for BC,

LDLR for HC and *GBA* for PD. We used the OpenCRAVAT annotations to classify rare

deleterious and/or pathogenic variants within these 3 genes. Rare deleterious variants were

those with a gnomAD v3 Non-Finnish European allele frequency of less than 0.1% and were,

functionally, either stop-gain or frameshift. Pathogenic variants were ClinVar "Pathogenic" or

"Likely Pathogenic" without conflicting "Benign" annotations. Variants not meeting either of these

criteria were considered negatives.

We evaluated as RV positive those individuals with at least one alternate allele for any

rare deleterious or pathogenic variant within each gene. To ensure independence between our

two causal factors (RV status and PRS) we further excluded all variants from each gene

strongly associated with the PRS for that gene's demonstration disease. We used logistic

regression to estimate the statistical significance of any correlation between individual variants

and their associated PRS and removed those with an effect p-value of less than 0.05.

We extracted relevant cohort metadata (age, sex, disease indicators, LDL, and ancestry

components) from the UKB cohort browser.

Controlling for Ancestry

We consider potential confounding between PRS and RV arising from shared Ancestry,

denoted A (**Figure 1B**). We adopt two approaches to address this situation: matching and

inverse probability weighting. With both methods we focus on the conditional outcome where

Y=1, cases only analysis. We expect that if RV influences Y then a negative association will

appear between RV and PRS such that affected individuals with RV=1 will have an average

lower PRS than affected individuals with RV=0.

To perform the matching, samples in each of the three cohort groups were differentiated based

on positive disease (case) status. A distance matrix between all samples was calculated using

the `distances` package in R. In the absence of eigenvalues for the available PCs in UKBB an

exponentially decreasing weighting function (scaled to unity) [decay = e^{r*5}, weights = decay /

sum(decay) **r = 0.637**] was used to weight distances for each PC in descending order. RV+

cases were matched to the closest (k=2) RV- cases using the calculated distances. For each

RV+ case, the difference between the observed PRS value and the mean PRS value for the

matched RV- cases was calculated. An overall test statistic was calculated by taking the

average of these values over all RV+ cases. Permutation testing was performed using the RV+

status as the treatment variable. A 1-tailed permutation test of means was then performed using the observed test-statistic against the permutation distribution.

We also performed an analysis using inverse probability weighting. Briefly, inverse

probability weighting is a general method for normalizing outcome data in causal analysis to

account for unequal representation of the causal exposure that correlates with an observable

common cause. In this situation the outcome variable is taken to be the PRS among the

affected individuals and the exposure is the binary RV; the common cause is ancestry. We use

the random forest method to perform a machine learning predictive analysis for RV positivity

using the ancestry principal components as predictors in each disease cohort. The outcome of

random forest is a probability of RV=1 for each observation. We then computed the inverse

probability weighted sum of PRS among RV+ and subtracted the sum of one minus the

probability of RV of the respective in the RV- cases. We again use permutation assignment of

the RV status to determine a null distribution and a one sided p-value.

Test of RV Load in Individuals

We tabulated a count of ClinVar pathogenic variants in a pathway and considered a

count variable as the candidate cause for the CP against PRS; in this approach the count is

total RV variants in a pathway using the same rule as for our binary RV analyses. For this

oligogenic analysis we focus on the lysosomal storage pathway in PD. We enumerated variants

in 54 lysosomal storage genes that were also present in UKBB. We then counted the number of

variants in each individual. To examine association of this count variable with PRS conditional

on disease we use Poisson regression to consider the average RV count as a function of PRS.

Briefly, in Poisson regression the logarithm of the rate of RV is modeled as a linear function of

the PRS. We used the LRT for Poisson regression as implemented in R to determine p-values.

Results

Comparative power analyses

We hypothesized our Causal Pivot LRT (CP-LRT) would have promising statistical properties to discover RV causal candidates by exploiting a PRS as a known cause. The model assumes the known factor is previously established and has a known effect size. The candidate cause is a binary variable representing the presence of at least one functional RV. This variable is assumed to have a known sparse rate of occurrence in the general population, and we use the value 0.001. Simulation-based power analyses presented in **Figure 2a** show the performance of the CP-LRT in the context of a binary outcome trait considering the possibility of both a main effect of RV as well as a PRS-RV interaction. In all simulations the forward logistic regression using the full data of both cases and controls has the highest power; this test is using approximately an order of magnitude more subjects because n=500,000 for case-control, vs n_c of approximately 50,000. Importantly, using only cases – approximately 10% of the population – achieves 90% of the power of the case-control analysis for the CP-LRT when the effect of the RV is at least as large as the effect of the PRS. In the situation of cases-only analyses with no interaction effect, the Z-test looking for overrepresentation of RV+ individuals has the highest power. When the RV main effect is sufficient and there is an interaction between RV and PRS the CP-LRT has superior power over the Z-test. The CP-LRT has superior power over the previously proposed Wilcox test (purple curve) when there is no interaction effect.

For the liability model (**Figure 2b**) we simulated according to a linear model for continuous quantitative outcomes. We modeled our analysis on the example with parameters $\alpha=0$, $\beta=0.3$, $\omega=0.001$, and we used a cutoff value of $\delta=1.13$,corresponding to the 92%-ile of the trait distrubtion, consistent with the LDL cholesterol cutoffs. Conditional cases-only analysis corresponds to the situation where observed samples are restricted to those with trait values over a cutoff. In this situation the CP-LRT has power that is stronger than the Wilcox test when

there is no interaction effect. When the interaction effect is strong the Wilcox test has power of

unity. The forward linear regression model using both case and control data has the strongest

power, and the Z-test also has high power.

We also perform robustness analysis focusing on the binary trait situation in the cases-

only circumstance. In this situation analyses for the Z-test and the CP-LRT rely on prior

information concerning RV variant frequency. Therefore, we explore the impact of errors in the

specification of this frequency. **Figure 2c** considers the situation where there is no RV effect,

with both $\gamma=0$ and $\eta=0$. Results show that the CP-LRT hast lower Type 1 error rate compared to

the Z-test, as indicated by the cyan Z-test curve dominating the red CP-LRT curve. The CP-

LRT retains a type 1 error of 0.05 when the RV rate is misspecified by 10%, from 0.001 down to

0.0009.

Conversely, the rate parameter for the RV might be falsely specified above its true value.

Figure **2d** shows this situation when the RV effect is set to $\alpha=-2.2$, with $\gamma=2.2$ and the PRS

parameter is $\beta=0.6$ – values driven from the UKBB data for BC. In these cases, the CP-LRT

has higher power to detect an effect while the Z-test power is sharply reduced, as indicated by

the red CP-LRT curve dominating the Z-test cyan curve. Interestingly, the CP-LRT retains a

power of 75% when the RV frequency is set to the expected expected RV frequency among

cases of 0.0045; in this circumstance the Z-test power is only 0.05.

Application to UKB data

To assess relevance of CP in real data we used UKB exomes and PRS. We used well

studied complex traits with both valid standard PRS as well as established single gene

contributors. The latter allowed us to recruit ClinVar annotations to select disease causing rare

variants as positive controls. Disease cohorts and variant selection are as described in the

Methods.

Cross Cohort Comparisons

We employ a cross-cohort comparative approach. We extract affected individuals in

each disease population and their respective PRS value as well as an RV determination mining

exome data for qualifying RV (see Methods). For visualization of the relative rate of RV against

PRS we split the respective cohorts into tertiles by their PRS. For each PRS tertile in each

cohort we counted the rate of RV in each PRS tertile bin; this analysis is repeated for each of

the three example genes: *BRCA1*, *LDLR* and *GBA*, respectively. We perform statistical testing

by application of our CP-LRT treating PRS quantitatively (**Figure 3a**). For each gene we

perform a CP-LRT for the rate of RV vs PRS by cohort. The diagonal of the plot are highly

significant (p< 1×10^{-6}) indicating that the CP-LRT found a significant negative association where

the matching gene RV state is tabulated against its disease specific PRS. Cross-cohort

analyses are mostly not significant, indicating that RV status for *BRCA1* and *GBA* are not

associated with either PD state or BC state, respectively. The gene *LDLR* shows weak but

significant CP-LRT association with both BC and PD. Supplemental Figure 2 shows the MLE

estimates of the RV and interaction parameters; the point estimates and confidence intervals

are provided. The point estimates using the CP MLE strongly agree with the results from case-

control analyses.

Controlling for the effect of Ancestry

Ancestry may confound the PRS-RV relationship by acting as a common cause for an

individual to have a functional RV as well as a high or low PRS. We call this scenario the

diamond graph (**Figure 4 a-c**). We use both matching and inverse probability weighting to

examine for signal while controlling for this possible confounding effect.

For matching we use the ancestry principal components to identify neighborhoods of

affected individuals. In each neighborhood we take the difference between the PRS of the RV+

and RV- affected individuals, and then average the PRS differences. The results show that

cases the RV+ individuals have on average significantly lower PRS scores than RV- individuals

after controlling for ancestry (**Figure 4d**).

Propensity scoring and inverse probability weighting is an alternative approach to control

for confounding. We treat RV status as a binary exposure, and we construct a model to predict

the chance of this exposure using Ancestry information from UKB population PC values. We

use random forests to predict the probability of RV+ status using the first 10 principal

components. We then compute the IP-weighted differences in PRS between RV+ and RV-

individuals. The results again show that PRS values are significantly lower in RV+ compared to

RV- individuals (**Supplemental Figure 1**).

Generalization to pathway burden scores

We considered the lysosomal storage pathway in the context of PD as an example of

pathway load analysis in the context of the CP. The lysosomal storage pathway is the system

that supports recycling of macromolecules in the brain. There are many lines of evidence that

point to lysosomal storage as a central pathway in neurodegeneration, and it plays a key role in

PD. We established a set of 54 human lysosomal storage pathway genes (**Supplemental**

table). We tabulate for each individual the number of qualifying RV in the lysosomal storage

pathway for each individual. We then compute the mean PRS of individuals with disease

stratifying on the number of pathway RV. We then use a Poisson Regression to examine the

association of the PRS with pathway burden of RV. The results are presented in **Figure 5a**.

Among the affected individuals there is a downward trend in PRS with increasing pathway

burden. To emphasize the combinatorial sparsity of the data we present the count of two-

variant combinations among cases **Figure 5b**.

Discussion

To date, the general principles of causal inference and structural causal models (SCM)

have been applied in complex trait genetics using Mendelian Randomization (MR)[14,16] which

exploits DNA variants as instrumental variables. Importantly, MR methods are typically used to

investigate single mechanisms that flow through a candidate biomarker and are not used to

differentiate individuals based on multiple independent causes or causal classes. Given the

success of MR, it is likely that SCM can be used even more broadly to model genetic effects,

suggesting several new avenues for addressing common diseases and other complex traits.[32]

The study of independent causes, in classical genetics called locus and allelic

heterogeneity, is central to understanding the genetic architecture of monogenic disorders.[33]

Parallel concepts of causal heterogeneity are needed for complex traits where current methods

focus on the average causal contribution of variants rather than their operation in potentially

heterogeneous subgroups.[34] There are already well-known examples in certain complex traits in

which the total population of affected individuals includes both those with high polygenic risk

along with individuals affected by monogenic forms. In this work, we have developed and

evaluated an approach to address this form of causal heterogeneity using a general SCM

framework. By examining the possible conditional relationship between a known cause and a

distinct candidate cause conditional on the phenotypic outcome we demonstrate that one can

establish their independent contributions. Moreover, we show that this approach applies in

cases-only, controls-only or combined case-control analyses. We call this conditional, induced

correlation approach the Causal Pivot (CP) to indicate that the known cause is used to leverage

the conclusion.

Simulation based power analyses enable controlled comparsion of the CP approach and

alternatives. The CP applies in cases-only analyses; the Wilcox and Z-test are chosen as non-

parametric comparators that also apply in cases-only analysis. The Wilcox test has been

proposed previously [26,27], and focuses exclusively on the induced correlation between PRS and

RV state conditional on disease. Alternatively, the Z-test focuses exclusively on the rate change

in RV in the disease population compared to its expected frequency in the general population.

Thus the Z-test ignores or fails to incorpoate the PRS information. The CP approach

incorporates both the frequency information from the general population and also the PRS-RV

conditional dependency. The results show that in the absence of a strong interaction the CP-LRT has stronger power than the Wilcox test, but it is weaker than the Z-test. When there is an interaction effect, the power of the CP-LRT exceeds the Z-test, but the Wilcox test surpasses it. Importantly, the CP likelihood approach results in parameter estimates and confidence intervals (see Supplemental materials); these interval estimates are an important advantage of the CP procedure over the non-parametric methods. Results show that the parameter estimates from the CP MLE in the logistic model framework are almost exactly equivalent to results from case-control logistic regression (see Supplmental Figure 2). Importantly, all of the tests have good power in the effect size range of interest for strong RV – where the RV effect exceeds that of the PRS.

The robustness analyses in Figure 2c and 2d are perhaps most revealing about the properties of the CP method. In the cases-only context, prior information concerning RV frequency is a precondition to apply either the Z-test or the CP-LRT. Importantly, the CP-LRT is informed by the structural model and includes the PRS as part of its procedure, while the Z-test does not. In terms of specificity, Figure 2c shows that CP-LRT is more stable against false positives when the population rate of RV is underestimated. Conversely, the Figure 2d shows that CP-LRT is also more sensitive when the RV rate is overestimated. Taken together Figure 2c and 2d show a key advantage of the CP-LRT. Because CP procedure is drawing information from the PRS it is effective when the Z-test fails.

The causal pivot is not a foreign concept in medical research.[6] Differential diagnosis is

an example of causal pivot thinking. Given that an individual is observed to have some

phenotype, when certain causes are ruled out, other causes become more likely. In genomic

medicine, the causal pivot has been observed in somatic cancer, where driver mutations are

observed to exclude each other.[35-37] This exclusion principle is also at the heart of

complementation group analysis in experimental genetics.

We chose polygenic risk scores (PRS)[38-43] as our pivot variable. PRS are quantitative

metrics for individual subjects derived from the marginal associations of higher frequency

markers with disease. To determine a PRS the individual's allelic counts are multiplied by

weights derived fromby genome wide association studies (GWAS) – these weights themselves

are commonly corrected for population structure and other covariates. PRS quantitatively

represents the aggregate genetic risk from a large number of GWAS signals - summing a large

number of diffuse or small effect size contributions. No single marker drives the PRS. Rather the

PRS aggregates the additive genetic effects by including tens, hundreds, thousands or even

millions of contributing variants. The advantage of the PRS is that it is encompassing and

produces a straightforward causal interpretation. The disadvantage of the PRS is the

disadvantage of any marker-based association analysis – the markers it employs are mostly

common and have individually modest effects, while the effect of rare, private and *de-novo*

variation is discounted or excluded.

RV are predominantly newer variants in a population and can have large functional

effects. Studies of rare and Mendelian diseases are rooted in the large effects of rare variation

as drivers of disease. RV are also observed in complex disease, where familial forms of many

complex diseases – including the example diseases of Hypercholesterolemia (HC), Breast

Cancer (BC)and Parkinson's Disease (PD). These diseases have familial and sporadic forms

caused by large-effect RV.

Our simple three node causal pivot model makes a strong assumption that there are no

common causes of the pivot variable and the causal candidate; in this case meaning that there

are not common causes of the the PRS value and the RV state. We demonstrated that the

assumptions can be relaxed and the causal pivot method can be extended to cases where the

independence assumption may not hold. We considered global ancestry as an obvious

possible confounder illustrated in the 'diamond' graph (Figure 1B). The relationship between

ancestry and the PRS may be complex and non-linear; the RV-ancestry relationship can be

equally cryptic; as such linear approaches to control for the relationship may fail. We used the

principal component vectors from genome wide ancestry analysis to implement non-parametric

approaches to control for the possible role of ancestry confounding. We used both matching and

inverse probability weighting where the procedure was determined by machine learning from the

ancestry principle components. In both cases our results indiciated a highly significant induced

negative correlation between PRS and RV after controlling for ancestry in the demonstration

diseases (**Figure 4).**

The Causal Pivot approach extends to the context of non-binary candidate causes. We

examined its use in the context of a pathway burden effect using the lysosomal storage pathway

in PD. Extensive literature has demonstrated the importance of the lysosome in

neurodegenerative disease, and many examples of Mendelian PD involve variants in lysosomal

storage genes. We modeled the count of ClinVar pathogenic variants in the pathway as a count

variable and we show that rate of the rare pathogenic variation negatively covaries with PRS

among affected individuals. We also present a graphical view of the PRS as a function of the

variant count and see that its mean behavior is decreasing in the variant burden.

We used simulation and real data to consider CP analyses in a cases-only context. The

cases-only test has slightly less power than a comparator case-control analysis, indicating that

most of the information about the causal effect of the RV is carried in the cases. It is important

to note that such a cases-only analysis is not possible using traditional forward association

tests. The new strategy is therefore ideal when selection on the phenotype is unavoidable.

There are many such scenarios relevant to complex diseases. Genotype X environment

interaction studies investigate whether the genetic variant and environmental exposure jointly

influence the risk, which cannot be tested in those who do not have the outcome. Case-

crossover designs the study is restricted to individuals who have already experienced the

outcome, and the focus is on understanding the exposure timing in relation to the event within

the same individual. Cases only analysis also applies in the context of somatic mutation

outcomes in cancer where analysis is confined to individuals who already have cancer, and the

focus is on the correlation between mutations and exposures or germline genetic variation.

Another cases only context occurs in drug safety studies that investigate the association

between drug exposure specific adverse outcomes; the design also applies to pharmacogenetic

studies that assess the drug response among patients who have already experienced a

particular outcome, such as an adverse drug reaction. In all these circumstances cases-only

genetic analyses are required.

The study of causal heterogeneity in complex disease genetics is a fertile area for

research. Significant progress has been made in identifying genes and mechanistic processes

that drive complex disease. In this context comparably less effort has been assigned to

deconvolving the disease population into subgroups. Our CP and CP-LRT method exploits prior

research from GWAS as well as surveys of genetic variation to drive discovery and assignment

of causal types. Additional research in the area of causal heterogeneity using SCM is likely to

prove productive.

Appendices

Derivation of Causal Pivot - .pdf

Supplementary Figures - .pdf

Table of Annotated Variants - .xlsx

Declarations of Interests

JWB and CAS are co-owners of Texas Genomics Consulting ~~which to date has worked in the development and implementation of clinical genetic testing in rare disease and cancer~~.

Acknowledgments

Grant from Genetics & Genomics Services, Inc.

Author Contributions

CAS developed the concept of the Causal Pivot and derived the SCM representation. Working together with TTT he developed the likelihood representation and likelihood ratio test. CAS derived the likelihood score equations for the likelihood analysis as well as the ratio representation for simplifying the CP-LRT; he developed the approach to Fisher Information and interval calculation for the method. He also developed the matching and inverse probability weighting method for Ancestry correction as well as the pathway burden modeling. He supervised the implementation of software and he wrote the manuscript and supplement. *Conceptualization, Formal analysis, Funding acquisition, Methodology, Project administration, Software, Resources, Supervision, Writing – original draft, Writing – review & editing*

CJW participated heavily in data curation as well as refinement and execution in software implementation of both power simulations and UKBB validations. CJW also executed the implementation of matching and random forest analysis to control for ancestry. He generated the results and figures for the manuscript. *Data curation, Software, Validation, Visualization, Writing – original draft*

JWB worked together with CAS to develop the concept of CP for complex disease and its context in genetic epidemiology. He guided the choice of example diseases and the selection of the proof of concept genes. He contributed to the derivation of the log-likelihood score equations. He provided funding support, and he was strongly involved in writing the manuscript. *Conceptualization, Data curation, Funding acquisition, Project administration, Software, Resources, Writing – original draft, Writing – review & editing*

JS contributed to development of the CP concept in neurodegenerative disease and identified *GBA* as a candidate gene for CP analysis in PD. He reviewed and assisted with the manuscript and provided funding support. *Conceptualization, Data curation, Funding acquisition, Writing – review & editing*

TTT worked with CAS on development of the LRT for CP and on initial simulation studies. He participated in development of the manuscript. *Formal analysis, Methodology , Writing – review & editing*

ND assisted with implementation of the log-likelihood score equations for both the logistic and liability model. He performed simulation studies for power analysis. *Formal analysis, Methodology , Writing – review & editing*

Web Resources

N/A

Data and code availability

Github ....

References

1.   Selvaraj, M.S., Li, X., Li, Z., Pampana, A., Zhang, D.Y., Park, J., Aslibekyan, S., Bis,
J.C., Brody, J.A., Cade, B.E., et al. (2022). Whole genome sequence analysis of
blood lipid levels in >66,000 individuals. Nat Commun *13*, 5995. 10.1038/s41467-
022-33510-7.

2.   Goodrich, J.K., Singer-Berk, M., Son, R., Sveden, A., Wood, J., England, E., Cole, J.B.,
Weisburd, B., Watts, N., Caulkins, L., et al. (2021). Determinants of penetrance and
variable expressivity in monogenic metabolic conditions across 77,184 exomes.
Nat Commun *12*, 3505. 10.1038/s41467-021-23556-4.

3.   Medeiros, A.M., and Bourbon, M. (2021). Polygenic contribution for familial
hypercholesterolemia (FH). Curr Opin Lipidol *32*, 392-395.
10.1097/MOL.0000000000000787.

4.   Sun, N., and Zhao, H. (2020). Statistical Methods in Genome-Wide Association
Studies. Annual Review of Biomedical Data Science *3*, 265-288. 10.1146/annurev-
biodatasci-030320-041026.

5.   Chen, W., Coombes, B.J., and Larson, N.B. (2022). Recent advances and challenges
of rare variant association analysis in the biobank sequencing era. Front Genet *13*,
1014947. 10.3389/fgene.2022.1014947.

6.   Pierce, B.L., and Ahsan, H. (2010). Case-only genome-wide interaction study of
disease risk, prognosis and treatment. Genet Epidemiol *34*, 7-15.
10.1002/gepi.20427.

7.   Wu, L., Schaid, D.J., Sicotte, H., Wieben, E.D., Li, H., and Petersen, G.M. (2015).
Case-only exome sequencing and complex disease susceptibility gene discovery:
study design considerations. J Med Genet *52*, 10-16. 10.1136/jmedgenet-2014-
102697.

8.   Webster, A.J. (2022). Causal attribution fractions, and the attribution of smoking
and BMI to the landscape of disease incidence in UK Biobank. Sci Rep *12*, 19678.
10.1038/s41598-022-23877-4.

9.   Glass, T.A., Goodman, S.N., Hernan, M.A., and Samet, J.M. (2013). Causal inference
in public health. Annu Rev Public Health *34*, 61-75. 10.1146/annurev-publhealth-
031811-124606.

10.   Rothman, K.J., and Greenland, S. (2005). Causation and causal inference in
epidemiology. Am J Public Health *95 Suppl 1*, S144-150.
10.2105/AJPH.2004.059204.

11.   Smith, G.D., and Ebrahim, S. (2004). Mendelian randomization: prospects,
potentials, and limitations. Int J Epidemiol *33*, 30-42. 10.1093/ije/dyh132.

12.   Zheng, J., Baird, D., Borges, M.C., Bowden, J., Hemani, G., Haycock, P., Evans, D.M.,
and Smith, G.D. (2017). Recent Developments in Mendelian Randomization
Studies. Curr Epidemiol Rep *4*, 330-345. 10.1007/s40471-017-0128-6.

13.   Sanderson, E., Glymour, M.M., Holmes, M.V., Kang, H., Morrison, J., Munafo, M.R.,
Palmer, T., Schooling, C.M., Wallace, C., Zhao, Q., and Smith, G.D. (2022).
Mendelian randomization. Nat Rev Methods Primers *2*. 10.1038/s43586-021-00092-
5.

14. Hu, X., Cai, M., Xiao, J., Wan, X., Wang, Z., Zhao, H., and Yang, C. (2024).
Benchmarking Mendelian randomization methods for causal inference using
genome-wide association study summary statistics. Am J Hum Genet *111*, 1717-
1735. 10.1016/j.ajhg.2024.06.016.
15. Dudbridge, F. (2021). Polygenic Mendelian Randomization. Cold Spring Harb
Perspect Med *11*. 10.1101/cshperspect.a039586.
16. Pingault, J.B., Richmond, R., and Davey Smith, G. (2022). Causal Inference with
Genetic Data: Past, Present, and Future. Cold Spring Harb Perspect Med *12*.
10.1101/cshperspect.a041271.
17. Pearl, J. (2010). An introduction to causal inference. Int J Biostat *6*, Article 7.
10.2202/1557-4679.1203.
18. Petersen, M.L. (2011). Compound treatments, transportability, and the structural
causal model: the power and simplicity of causal graphs. Epidemiology *22*, 378-
381. 10.1097/EDE.0b013e3182126127.
19. Lipsky, A.M., and Greenland, S. (2022). Causal Directed Acyclic Graphs. JAMA *327*,
1083-1084. 10.1001/jama.2022.1816.
20. Holmberg, M.J., and Andersen, L.W. (2022). Collider Bias. JAMA *327*, 1282-1283.
10.1001/jama.2022.1820.
21. Cai, S., Hartley, A., Mahmoud, O., Tilling, K., and Dudbridge, F. (2022). Adjusting for
collider bias in genetic association studies using instrumental variable methods.
Genet Epidemiol *46*, 303-316. 10.1002/gepi.22455.
22. Coscia, C., Gill, D., Benitez, R., Perez, T., Malats, N., and Burgess, S. (2022).
Avoiding collider bias in Mendelian randomization when performing stratified
analyses. Eur J Epidemiol *37*, 671-682. 10.1007/s10654-022-00879-0.
23. Mahmoud, O., Dudbridge, F., Davey Smith, G., Munafo, M., and Tilling, K. (2022). A
robust method for collider bias correction in conditional genome-wide association
studies. Nat Commun *13*, 619. 10.1038/s41467-022-28119-9.
24. Mitchell, R.E., Hartley, A.E., Walker, V.M., Gkatzionis, A., Yarmolinsky, J., Bell, J.A.,
Chong, A.H.W., Paternoster, L., Tilling, K., and Smith, G.D. (2023). Strategies to
investigate and mitigate collider bias in genetic and Mendelian randomisation
studies of disease progression. PLoS Genet *19*, e1010596.
10.1371/journal.pgen.1010596.
25. Swanson, S.A., Robins, J.M., Miller, M., and Hernan, M.A. (2015). Selecting on
treatment: a pervasive form of bias in instrumental variable analyses. Am J
Epidemiol *181*, 191-197. 10.1093/aje/kwu284.
26. Zhou, D., Yu, D., Scharf, J.M., Mathews, C.A., McGrath, L., Cook, E., Lee, S.H.,
Davis, L.K., and Gamazon, E.R. (2021). Contextualizing genetic risk score for
disease screening and rare variant discovery. Nat Commun *12*, 4418.
10.1038/s41467-021-24387-z.
27. Lu, T., Forgetta, V., Richards, J.B., and Greenwood, C.M.T. (2022). Polygenic risk
score as a possible tool for identifying familial monogenic causes of complex
diseases. Genet Med *24*, 1545-1555. 10.1016/j.gim.2022.03.022.
28. Spirtes, P., Glymour, C., and Scheines, R. (2001). Causation, Prediction, and
Search, second edition (The MIT Press).

29. Kalisch, M., Mächler, M., Colombo, D., Maathuis, M.H., and Bühlmann, P. (2012).
Causal Inference Using Graphical Models with the R Package pcalg. Journal of
Statistical Software *47*, 1 - 26. 10.18637/jss.v047.i11.
30. Badsha, M.B., Martin, E.A., and Fu, A.Q. (2021). MRPC: An R Package for Inference
of Causal Graphs. Front Genet *12*, 651812. 10.3389/fgene.2021.651812.
31. Thompson, D.J., Wells, D., Selzam, S., Peneva, I., Moore, R., Sharp, K., Tarran, W.A.,
Beard, E.J., Riveros-Mckay, F., Palmer, D., et al. (2022). UK Biobank release and
systematic evaluation of optimised polygenic risk scores for 53 diseases and
quantitative traits. medRxiv, 2022.2006.2016.22276246.
10.1101/2022.06.16.22276246.
32. Madsen, A.M., Ottman, R., and Hodge, S.E. (2011). Causal models for investigating
complex genetic disease: II. what causal models can tell us about penetrance for
additive, heterogeneity, and multiplicative two-locus models. Hum Hered *72*, 63-
72. 10.1159/000330780.
33. Stern, C. (1973). Principles of Human Genetics, 3rd Edition (W. H. Freeman and
Company).
34. Woodward, A.A., Urbanowicz, R.J., Naj, A.C., and Moore, J.H. (2022). Genetic
heterogeneity: Challenges, impacts, and methods through an associative lens.
Genet Epidemiol *46*, 555-571. 10.1002/gepi.22497.
35. Babur, O., Gonen, M., Aksoy, B.A., Schultz, N., Ciriello, G., Sander, C., and Demir, E.
(2015). Systematic identification of cancer driving signaling pathways based on
mutual exclusivity of genomic alterations. Genome Biol *16*, 45. 10.1186/s13059-
015-0612-6.
36. Dao, P., Kim, Y.A., Wojtowicz, D., Madan, S., Sharan, R., and Przytycka, T.M. (2017).
BeWith: A Between-Within method to discover relationships between cancer
modules via integrated analysis of mutual exclusivity, co-occurrence and functional
interactions. PLoS Comput Biol *13*, e1005695. 10.1371/journal.pcbi.1005695.
37. Zhang, W., Zeng, Y., Wang, L., Liu, Y., and Cheng, Y.N. (2020). An Effective Graph
Clustering Method to Identify Cancer Driver Modules. Front Bioeng Biotechnol *8*,
271. 10.3389/fbioe.2020.00271.
38. Abraham, G., Rutten-Jacobs, L., and Inouye, M. (2021). Risk Prediction Using
Polygenic Risk Scores for Prevention of Stroke and Other Cardiovascular Diseases.
Stroke *52*, 2983-2991. 10.1161/STROKEAHA.120.032619.
39. Lakeman, I.M.M., Rodriguez-Girondo, M.D.M., Lee, A., Celosse, N., Braspenning,
M.E., van Engelen, K., van de Beek, I., van der Hout, A.H., Gomez Garcia, E.B.,
Mensenkamp, A.R., et al. (2023). Clinical applicability of the Polygenic Risk Score
for breast cancer risk prediction in familial cases. J Med Genet *60*, 327-336.
10.1136/jmg-2022-108502.
40. O'Sullivan, J.W., Raghavan, S., Marquez-Luna, C., Luzum, J.A., Damrauer, S.M.,
Ashley, E.A., O'Donnell, C.J., Willer, C.J., Natarajan, P., American Heart Association
Council on, G., et al. (2022). Polygenic Risk Scores for Cardiovascular Disease: A
Scientific Statement From the American Heart Association. Circulation *146*, e93-
e118. 10.1161/CIR.0000000000001077.

41.   Rao, A.S., and Knowles, J.W. (2019). Polygenic risk scores in coronary artery
disease. Curr Opin Cardiol *34*, 435-440. 10.1097/HCO.0000000000000629.
42.   Riveros-Mckay, F., Weale, M.E., Moore, R., Selzam, S., Krapohl, E., Sivley, R.M.,
Tarran, W.A., Sorensen, P., Lachapelle, A.S., Griffiths, J.A., et al. (2021). Integrated
Polygenic Tool Substantially Enhances Coronary Artery Disease Prediction. Circ
Genom Precis Med *14*, e003304. 10.1161/CIRCGEN.120.003304.
43.   Roberts, E., Howell, S., and Evans, D.G. (2023). Polygenic risk scores and breast
cancer risk prediction. Breast *67*, 71-77. 10.1016/j.breast.2023.01.003.

Figure Titles and Legends

Figure 1 Core Logic of the Causal Pivot. **Panel a)** depicts the basic model with PRS as a known causal factor indicated by the solid line and arrow terminating at Y; RV is an independent factor whose effect on Y is unknown and being evaluated. **Panel b)** depicts the impact of the solid arrow from PRS to Y; the solid line indicates a correlative relationship between PRS and Y which manifests as a higher PRS in cases where Y=1 than in controls where Y=0. **Panel c)** depicts the model of unconditional independence between PRS and RV. The unconditional independence results in a lack of pattern between PRS and RV state, which can also be viewed as a lack of correlation in the rate of the RV+ as a function of the tertiles of the quantitative PRS. In **panel d)** we depict the circumstance where RV influences phenotype as indicated by the solid arrow from RV to Y; the red box around Y indicates conditioning on the phenotypic state. In the circumstance where RV drives Y and where the value of Y is conditioned upon (red box), then an induced correlation appears between PRS and RV. This collider-induced correlation is depicted by the dashed red line connecting PRS and RV. In **panel e)** we depict the rate change in RV+ status according to the value of Y as would result from a solid arrow connecting RV and Y. In **panel f)** we depict the collider-induced correlation between PRS and RV conditional on Y; this correlation manifests as a Y-conditional shift in PRS between RV+ and RV-; alternatively, the rate of the RV+ state covaries with PRS conditional on Y. This collider induced correlation between PRS and RV juxtaposes with the unconditional independence of the PRS and RV as shown in **panel c)**. The Causal Pivot recruits the collider induced correlation together with the rate change in the RV – as shown in **panel d --** as signal to infer the RV-Y relationship.

Figure 2. Simulation Analyses The figure presents simulation analyses comparing the CP-LRT to alternatives. Monte Carlo methods are used to generate the results in panels a-d; these simulations generate PRS and RV status which are then used to generate trait outcomes under

the corresponding probabilistic model – either logisitic or quantitative. Alternative analyses are then performed to detect the effect of the RV factor and to compare alternative approaches on the same simulated data. Detailed markdown documentation and code to produce these analyses are provided (https://github.com/chadashaw/pivotr). Panel a) focuses on binary traits; the outcome is generated by a logistic model where PRS and RV are causal factors. Total population size is n=500,000, comparable to a UKBB scale. We set the intercept α=-2.2 corresponding to a disease prevalence of 10%, and the PRS effect β=0.5, consistent with UKBB data for our three demonstration diseases; the RV state is generated with frequency ω=0.001. The x-axis represents the main effect of the rare variant -- parameterized as γ. The y-axis represents the statistical power as determined by the proportion of simulation runs that reject the null hypothesis of no PRS effect under a nominal Type 1 error rate of 0.05. The curves present power comparisons to detect the RV effect as the main effect of the RV varies. The cases-only CP-LRT is shown in red. Cases-only alternatives are the Wilcox test (purple) that considers non-parametric mean shift in PRS values in cases between RV+ vs RV-; the Z-test (cyan) considers the increase in the frequency of RV+ among cases compared to what would be expected from the background frequency ω=0.001. Also included is the forward logistic regression using both cases and controls (green) that represents a typical regression approach. For the dashed line curves the PRS-RV interaction is set to η= -.4. In the absence of an interaction effect the Wilcox test has sharply lower power than CP-LRT and the Z-test. The Z-test is more sensitive to detect RV effects than the CP-LRT; the Z-test is also non-parametric and does not distinguish main effects and interaction effects between PRS and RV state. The Panel b) presents analyses when the phenotypic outcome is quantitative; again, a biobank scale

population is simulated, but here the trait is generated according to a linear model; we set

$\alpha=0$ and set $\beta=0.3$; RV occurs at a rate of 0.001. To construct a conditional outcome we

identify individuals who have a trait value larger than a quantile cutoff, and we use an empirical

percentile of 0.93 consistent with the HC cutoff from UKBB corresponding to LDL> 190. We

compare CP-LRT to cases-only alternatives of the Wilcox test (purple) and a Z-test (cyan) as well

as an unconditional linear regression model (green) for the trait using the full sample. The CP-

LRT has superior power to the Wilcox test in the absence of an interaction term. The Z-test still

outperforms; the data linear regression has the best power. When the interaction effect is

present all tests have higher power. The Wilcox test is notably higher. In panel c) we scrutinize

fold increase in Type-1 error (y-axis, ratio of simulated type 1 error to 0.05) as a measure of the

comparative stability of the CP-LRT compared to the Z-test. In c) simulations use the same

parameters $\alpha,$ β,ω as in panel a) but holding γ,η at 0. Both the CP-LRT and Z-test require

specification of ω the population rate of RV. The x-axis in c) considers ω^* where the analysis

underspecifies the RV rate. This under specification results in a sharp increase in the type 1

error of the Z-test compared to the CP-LRT; the Z-test error immediately exceeds 0.05 as ω^* is

decreased from the true value of 0.001. Differently, the CP-LRT – which leverages the collider

induced correlation – is more robust and retains a type 1 error of close to 0.05 beyond a 10%

reduction in ω. The red curve is uniformly below the Z-test indicating the CP-LRT retains better

type 1 error robustness. In **panel d)** we again focus on the logistic model as in **panel a).** Here we

compare the sensitivity of CP-LRT to Z-test when the RV has an effect but where the analysis

false misspecifies the RV rate ω^* (x-axis) above its true value of 0.001 (dahsed vertical line).

Incorrect overspecification of RV rate diminishes the power of the Z-test. In these simulations

we again set $\alpha=-2.2$; we set $\beta=0.6$ and $\gamma=2.2$, consistent with the parameterization for the

BRCA1 gene in BC (Supplemental Table 1). The vertical dashed line near $\omega^*=0.045$ indicates

performance when the RV rate is set to the expected frequency of the RV in the Y=1 population.

In this case the rejection rate of Z-test is about 0.05, while the CP-LRT has about 75% power.

The retains CP-LRT power because it recruits information from the istructurally nduced

correlation between RV and PRS.

Figure 3. **Demonstration of the Causal Pivot in UKBB data.** Panel a) shows a 3x3 collection of

barplots where the vertical bars represent the count of RV+ Samples within each PRS-tertile

grouping which defines the X-axes. Each row represents a different disease cohort paired with

its disease specific PRS; HC in row one, BC in row two and PD in row three. Each column

presents a different disease gene as indicated by the column headers, with *LDLR* variation

tabulated in column one, *BRCA1* variation in column two and *GBA* in column three. The

diagonal of panel a) shows the negative correlation between the number of RV+ samples and

PRS scores within each PRS-Disease-Gene grouping and a lack of association characterizes the

barplots off the diagonal; the CP-LRT was used to determine p-values for all 9 analyses, and the

p-values for all tests are p<10^-5 along the diagonal. Off diagonal entries are not significant

except for the gene *LDLR* which shows modest association pattern with both BC and PD with

p<0.05. More detailed results including confidence intervals for γ and η appear in the

Supplemental materials. In panel cb the graphic presents the multiplicity and typology of the RV

represented in the panel a) data. Most RV occur in only a single sample. A variety of different

variant types are present. A detailed listing of the variants that inform the analysis is provided

in a Supplemental Variant Table.

Figure 4 Controlling for Ancestry In any genetic association analysis ancestry is a potential common cause. In Panel a) we depict the scenario where ancestry represented by node A drives both PRS and RV state. In panel b) RV does not drive Y, and conditioning on both A and Y there is no induced or residual correlation between PRS and RV. In panel c) RV drives Y, and conditioning on both A and Y yields an induced correlation between PRS and RV. In panel d) we present three distinct rows of analyses; HC in row one, BC in row two and PD in row three. Each row shows three results. The scatter plots show RV$^-$ (black points) and RV$^+$ (red points) cases as points in an ancestry visualization space. The scatter plot representation is determined by the first 5 PCs reduced to two dimensions using t-distributed stochastic neighbor embedding (t-SNE). A zoom in view of each row is indicated by the excerpted rectangular region. Each RV$^+$ case is matched to the nearest (k=2) RV$^-$ cases in the 5 dimensional ancestry space. The mean difference in the PRS of each RV+ and its ancestry matched neighbors is determined. The average of these PRS differences is then calculated and shown in the dashed vertical red line in each histrogram plot. The histogram plots show both these real data calculations as well as the results of permutation testing depicted in the histrograms. The permutation distributions are generated by randomly permuting the RV states of each sample in the cohort while fixing the ancestry data; the procedure of identifying neighbors and calculing the mean difference in PRS is repeated. In addition to the histograms that present the permutation distributions, the 5% threshold value is shown by the dashed black lines. The observed value from the true RV+ ancestry matched data is shown by the dashed red line is significantly below the 0.05 quantile in all cases, consistent with the pattern that RV+ cases have lower PRS than their RV- ancestry-matched neighbors.

Figure 5. Pathway burden analysis. Panel a) shows the the PRS values (y-axis) against the count of pathogenic/likely pathogenic RV in the lysosomal storage pathway among PD+ cases from UKBB. Each boxplot presents cases with the tabulated count of RV+ per case. There is a decreasing trend in PRS, and a Poisson regression analysis which models the rate of RV as a function of PRS demonstrates a statisticallly significant negative association between the rate of RV+ and the PRS value (p=0.03). Panel b) depicts the sparse occupancy of positivity for 2 variants. No single pair accounts for the majority of instances.

Tables

Rare Variant Table from UKBB

Figure 1: Core Logic of the Causal Pivot

Figure 2: Simulation Analyses

Figure 3: UKBB Results

Figure 4: Controlling for Ancestry

* p < 0.05

Figure 5: Pathway Burden