Longer Internal Exons Tend to Have More Tandem Repeats and Experience Insertions

and Deletions More Frequently

Keiichi Homma[* 1], Hiroto Anbo[1], Motonori Ota[2], and Satoshi Fukuchi[1]

[1] Program for Information Systems, Division of Informatics, Bioengineering and Bioscience,

Maebashi Institute of Technology, 460-1 Kamisadori-machi, Maebashi-shi, 371-0816 Japan

[2] Graduate Schol of Informatics, Nagoya University, Nagoya, 464-8601 Japan

[*]**Corresponding author:** khomma@maebashi-it.ac.jp

Abstract

Insertions and deletions (indels) within the coding sequences of eukaryotic proteins are known

to preferentially encode intrinsically disordered regions (IDRs), protein regions that by

themselves do not form unique three-dimensional structures. As a previous investigation of

our group showed that long internal exons have a tendency to encode IDRs in eukaryotes

(Fukuchi et al. 2023), it was thought worthwhile to analyse how indels alter internal exons and

affect IDRs of the encoded proteins. For consideration of evolutionary roles indels play, it is

important to focus on indels commonly observed in all variants ("fixed" indels) since indels in

minor variants may represent transient aberrations in splicing. Here, by comparison of

orthologous variants of closely related species together with those of outgroups, fixed indels in

the internal exons were identified in four mammals and two flies. Deletion tends to occur more

frequently than insertion does and indels are generally more prevalent in long internal exons.

On average 51% of inserted and 40% of deleted residues are attributable to alterations in

tandem repeats. Nearly all indel lengths are multiples of three nucleotides (nt), shorter than 19

nt and mostly encode IDRs. Intriguingly, it was discovered that tandem repeats preferentially

occur in long internal exons. As tandem repeats mostly encode IDRs, this finding at least

partially explains the high incidence of IDRs in long internal exons. We propose that long

internal exons had been produced in early eukaryotes mainly by repeat expansion but are

experiencing frequent indels by alterations in tandem repeats.

Key words: intrinsically disordered region, evolution, indel, exon, FECA, LECA

Significance

Besides structural domains, many proteins have regions that by themselves do not form unique three-dimensional structures, called intrinsically disordered regions (IDRs), which constitute approximately a third of residues in eukaryotic proteins, but a much smaller fraction in prokaryotic proteins. As insertions and deletions (indels) frequently encode IDRs, we here rigorously identified indels to investigate how eukaryotic proteins had accumulated IDRs in evolution and discovered that alterations in tandem repeats constitute a major generation mechanism of indels and that indels occur more often in longer internal exons. Since tandem repeats were found to be more prevalent in long internal exons, we propose that expansion of tandem repeats had elongated some internal exons and added IDRs to primordial eukaryotic proteins.

1   Introduction

Only a small fraction of prokaryotic proteins (residue-wise 2.0% of archaean and 4.2% of

eubacterial proteins) consist of intrinsically disordered regions (IDRs) that by themselves do

not form unique three-dimensional structures, whereas 33.0% of residues in eukaryotic

proteins are comprised of IDRs (Ward et al. 2004). IDRs in eukaryotic proteins participate in

binding to other molecules and are thereby involved in important functions such as gene

transcription and signaling (Anbo et al. 2019). Two mechanisms of *de novo* IDR generation

are expansion of repetitive DNA sequences (Tompa 2003) and exonisation of introns

(Kondrashov et al. 2003; Sorek 2007; Marquez et al. 2015), but to our knowledge their

proportional shares have not been quantified. It is intriguing to uncover how IDRs had been

acquired in the evolutionary process from the first eukaryotic common ancestor (FECA) to the

last eukaryotic common ancestor (LECA).

We previously reported the general eukaryotic tendency of long internal exons to encode

IDRs and proposed that long internal exons in the LECA had resulted from short exons by

addition of IDR-encoding nucleotides (Fukuchi et al. 2023). As indels play important roles in

exon length alterations and mostly encode IDRs (Light et al. 2013A, B; Khan et al. 2015),

indel analyses may reveal mechanisms of exon length alterations and thereby explain how the

preferential encoding of IDRs by long internal exons came about.

Indels in human genomes cause as much variation as small nucleotide polymorphisms

(Mullaney et al. 2010) and one way to identify them in coding sequences is to find indels in

genomes and select those in coding sequences (Anzai et al. 2003; Mills et al. 2006; Wetterbom

et al. 2006; Fan et al. 2007; Khan et al. 2015; Lin et al. 2017). As coding sequences vary from

one alternative splicing variant to another, however, different sets of coding sequences result

in different indel identifications. Another method to select indels in proteins is to compare the

amino acids sequences of orthologous proteins either by sequence alignments (Benner et al.

1993; Light et al. 2013A, B) or by structural alignments (Pascarella et al. 1992; Qian and

Goldstein 2001). Since indels thereby identified are dependent on the selection of orthologous

proteins, splicing variants need to be considered, and structural alignments cannot be made to

stretches containing IDRs, which is a relevant issue because, as stated above, indels frequently

encode IDRs (Light et al. 2013A, B; Khan et al. 2015). As the increasing availability of variant sequences makes it possible to identify indels shared by all variants, we conducted sequence alignments of all available variants, selected commonly observed indels, and called them "fixed" indels.

Fixed indels can result from constitutive changes in splicing and genomic mutations. Changes in splicing that give rise to insertions can be regarded as intron to exon conversions (exonisations), while those resulting in deletions can be considered as exon to intron conversions (intronisations). Indels by genomic mutations have three plausible causes: DNA slippage, DNA damage followed by imperfect repair (Redelings et al. 2024), and transposable elements, which constitute approximately 4% of human protein-coding genes (Nekrutenko and Li 2001). DNA slippage frequently occurs in short tandem repeats that in most cases encode IDRs (Simon and Hancock 2009, Levinson and Gutman 1987), while DNA damage is mostly repaired by homologous repair pathways.

Since splicing of exons longer than 300 nt is inhibited unless flanked by short introns (Robberson et al. 1990; Chen and Chasin 1994; Sterner et al. 1996), many exons that became excessively long may not be efficiently spliced. On the other hand, analyses of RNA-Seq data demonstrated that introns shorter than 70 nt are not completely spliced out (Abebrese et al. 2017). Exon definition was proposed to explain the splicing mechanism in vertebrates in which short exons separated by long (>250 bp) introns predominate (Robberson et al. 1990; Berget 1994; De Conti et al. 2012), while intron definition was postulated to account for the splicing of long exons with short introns prevalent in lower eukaryotes (Lang and Spritz 1983; Berget 1994; De Conti et al. 2012). Recent research disclosed that the same spliceosome assembled on introns provides the mechanisms for both intron and exon definition in *Saccharomyces cerevisiae* (Li et al. 2019).

In this investigation, we identified fixed indels that are uniformly present in all variants and attempted to determine their generation mechanisms and found a considerable number of indels generated by alteration in the number of tandem repeats, which frequently encode IDRs. Our analyses also revealed that longer internal exons tend to experience more indels and generally have a higher prevalence of tandem repeats, suggesting that long internal exons were

generated chiefly by repeat expansions. We propose that the expansion of tandem repeats encoding DRs was a crucial evolutionary mechanism by which the LECA arose from the FECA.

Results

Selection of Fixed Indels in Internal Exons

Exploiting a wealth of sequences in the Ensembl database (Martin et al. 2023), we first aligned

all variants of orthologous genes in closely related sp. 1 and sp. 2 and selected sections

missing in sp. 2 as indel candidates (Figure 1a, c). We then carried out BLASTN alignments

(Altschul et al. 1990) of all variants of orthologous genes in spp. 1 and 2 and an outgroup to

classify them into insertion and deletion candidates; those present in an outgroup variant were

considered as insertion candidates because the segments were probably absent before the

bifurcation of sp. 1 and sp. 2 (Figure 1b), while those missing in an outgroup variant were

regarded as deletion candidates as the segments were likely to be present in the immediate

ancestor of sp. 1 and sp. 2 (Figure 1d). To filter out candidates that correspond to splicing

variants, we applied uniformity tests; we selected the insertion candidates whose segments are

present in all sp. 1 variants and are unexceptionally absent in sp. 2 and outgroup variants and

regarded them as fixed insertions (Figure 1b); we likewise chose the deletion candidates

whose segments are invariably absent in sp. 2, but omnipresent in both sp. 1 and outgroup

variants and considered them as fixed deletions (Figure 1d). Finally, only those in internal

exons are selected and are called fixed indels. We present the actual steps followed to select

fixed indels in internal exons and the number of cases in supplementary figure S1 and table S1

(Supplementary Material online). (For explanations of step 5, see below.)

Fig. 1.—Selection of fixed indels. Rectangles represent sections of coding sequences and

are arranged so that aligned segments match vertically. a, b) The criteria for selecting fixed

insertions and c, d) fixed deletions are schematically shown.

Insertions are probably generated either by intron to exon conversion (exonisation)

(Figure 2a-h) or by genomic modifications, the latter of which may be caused by expansion of

tandem repeats (Figure 2i), homologous recombination (Figure 2j), or insertion of exogenous

segments (Figure 2k). Inserted segments that are presumably generated by exonisation but

correspond to introns with lengths less than 70 nt (Figure 2b, f) were discarded (step 5 in

supplementary figure S1, Supplementary Material online) because such introns may be

imperfectly removed (Abebrese et al., 2017), making them unqualified as "fixed" insertions.

Similarly, the three generating mechanisms of deletions are exon to intron conversion

(intronisation) (Figure 2i-s), contraction of tandem repeats (Figure 2t), homologous

recombination (Figure 2u), and genomic deletion (Figure 2v). Deleted segments that

corresponded to short (<70 nt) introns (Figure 2m, q) were removed for the aforementioned

reason. (In the following, we additionally carried out analyses of the permissive sets of fixed

indels to demonstrate that the removal does not affect conclusions.) Insertions with

corresponding introns in sp. 2 possessing no significant homology were considered as possibly

generated by exonisation (Figure 2a, c, d), while those alignable to intron segments in sp. 2 were classified as probably generated by exonisation (Figure 2e, g, h). As introns generally have a higher mutation rate than exons do (Li 1997), exons created by exonisation may no longer possess detectable sequence homology with the introns. Likewise, we consider deletions as possibly generated by intronisation if the corresponding intron segment in sp. 2 has no discernable similarity to the exons in sp. 1 (Figure 2l, n, o), while sorting those as probably generated by intronisation if the intron segments were alignable to the exons (Figure 2p, r, s). We distinguish fixed insertions of entire exon(s) (Figure 2c, g) from those within exons (Figure 2a, e) and regard fixed deletions of entire exon(s) (Figure 2n, r) separately from those within exons (Figure 2l, p). The total number of indels within internal exons as well as their breakdown by indel location within exons are shown in supplementary tables S2 and S3 (Supplementary Material online), the latter of which list numbers without removing those corresponding to short (<70 nt) introns (designated the "permissive" sets of fixed indels). Fixed indels of entire exon(s) are rare and almost all the fixed indels in internal exons are located in the middle of rather than at 5' and 3' ends of exons (supplementary figure S2, Supplementary Material online).

Fig. 2.--Generation mechanisms of fixed indels. Presumed generating mechanisms of

fixed indels are pictorially explained. Grey rectangles and horizontal bars stand for exons and

introns, respectively, while parallel vertical bars signify tandem repeats.

Six examples of fixed indels are provided (Figure 3) and two examples of indels that were

removed due to their correspondence to excessively short (<70 nt) introns are shown

(supplementary figure S3, Supplementary Material online). The human protein in Figure 3a is

a protein enabled homologue involved in a range of processes dependent on cytoskeleton

remodeling and cell polarity, the *D. melanogaster* protein in Figure 3b is E2F transcription

factor, isoform D, whereas the rat protein presented as Figure 3c is nuclear factor erythroid 2-

related factor 2, which is a transcription factor that plays a key role in the response to

oxidative stress. The chimpanzee protein depicted in Figure 3d is a trinucleotide repeat-

containing protein that plays a role in RNA-mediated gene silencing, the human protein shown

in Figure 3e is GRIP and coiled-cold domain-containing protein isoform 2, while the human

protein in Figure 3f is SMG1 phosphatidylinositol 3-kinase-related kinase partaking in both

mRNA surveillance and genotoxic stress response pathways.

Fig 3.--Actual examples of fixed indels. Examples of fixed insertions a-c) and deletions d-f) depicted as in Fig. 2 in the scale indicated with tandem repeats represented by parallel vertical bars. The encoded proteins are shown as orange rectangles in arbitrary scales with horizontal parallel bars representing IDRs and grey rectangles corresponding to the depicted genomic sections.

Generation Mechanisms of Fixed Indels

The residue-wise distributions of generation mechanisms of fixed indels are graphically

presented (Figure 4) with the indels whose generating mechanisms remain unidentified labeled

"unknown". No indel cases involving transposable elements were detected even if the

expectation value of the BLASTN search was relaxed to 10 and no insertion cases whose

generation mechanisms are unknown were long enough to be assessed for involvement of

other exogenous elements. The corresponding figure of the permissive sets of fixed indels is

presented as supplementary figure S4, Supplementary Material online. The numbers of fixed

indel events and those of residues in insertion and deletion are tabulated (supplementary table

S4, Supplementary Material online), while the corresponding numbers of the permissive sets

of indels are presented as supplementary table S5, Supplementary Material online. Although

interspecies variations are considerable, on average 51% of inserted and 40% of deleted

residues were attributable to alterations in tandem repeats and the generation mechanisms of a

substantial number of fixed indels remain unidentified (see Discussion). Fixed indels

generated by intronisation and exonisation exist but have small shares except in human

insertions, chimpanzee deletions, and rat indels, whereas those attributable to homologous

recombination are universally rare.

Fig. 4.--Generating mechanisms of fixed indels.
21

Long Internal Exons Tend to Have a High Incidence of Indels

As we previously found that long internal exons tend to encode IDRs (Fukuchi et al. 2023), we sought to search for evolutionary generation mechanisms of long internal exons and thus determined the dependence of indels on exon length. We first calculated the abundance of internal exons in 30 nt length bins (supplementary figure S5, Supplementary Material online) and found that, while long internal exons are rare in the four mammals, the two *Drosophila* species showed distributions shifted to longer length, confirming a previous report (Hawkins 1988). We then examined the frequencies of indels in each length bin of internal exons, using the length of the corresponding sp. 1 exons for the analyses of insertion in sp. 1 and deletion in sp. 2. The length of the internal exon with an insertion utilized was that of the exon minus the insertion length because that is the presumed exon length before insertion. In the example presented as Figure 3c in which a 21 nt insertion exists in rat exon 4 of 189 nt, we used168 nt as the length of the primordial exon in which the insertion occurred. (Although 168 nt happens to be the length of the corresponding sp. 2 exon in this case, the lengths sometimes disagree as some sp. 2 exons have been altered). The indels of entire exon(s) (Figure 2c, g, n, r), which constitute small fractions (supplementary figure S2, tables S2 and S3, Supplementary Material online), were not included in this and subsequent analyses because our aim was to analyse incremental length changes in internal exons. Interestingly, we found higher frequencies of indels in long internal exons without exception (Figure 5 and supplementary figure S6, Supplementary Material online). The correlation coefficients significantly differ from 0 ($P<0.01$) for insertions in *Drosophila melanogaster* and all deletions except for those in human and chimpanzee. Also, deletion frequency tends to be higher than insertion frequency in each species except in short intron length bins of human insertions.

Fig. 5.--Long internal exons tend to have a high incidence of indels in each species. One,
two, and three asterisks signify that the correlation coefficient is significantly different from
zero at $P<10^{-2}$, 10^{-3}, and 10^{-4}, respectively. a) The normalized averages of insertion (left scale)
and deletion (right scale) frequencies are shown with the scale of deletion frequencies
multiplied by the average ratio of deletion to insertion frequency, which is the arithmetic
average of the ratios of deletion to insertion frequency of the six species. The dotted lines are
regression lines for insertion (yellow) and deletion (blue) frequencies. b-g) The indel
frequencies in each internal length bin are graphed for each species.

As the smallness of indel residues seemed to cause fluctuations in indel frequency, we calculated the average indel frequency of the six species in each internal exon length bin. The combined dependence of indel frequency on internal exon length (Figure 5a and supplementary figure S6a, Supplementary Material online) shows that positive, statistically significant correlations exist, and that deletion frequency is on average higher than insertion frequency with the difference more pronounced in long internal exons. The latter observation reflects the fact that insertion frequency mostly levels off in the long length range, while deletion frequency does not.

Most Fixed Indels Encode Intrinsically Disordered Regions, Conserve Reading Frames, and Are Short

As indels reportedly occur preferentially in IDRs (Light et al. 2013A, B; Khan et al. 2015), we calculated the fractions of IDRs in the fixed indels. For deleted segments in sp. 2, we used the corresponding segments in sp. 1 (Figure 1d) as a proxy. In agreement with the literature, most indels encode IDRs and the fractions are all much higher than those of all coding exons (Figure 6a and supplementary figures S7 and S8, Supplementary Material online). We also examined indel lengths and verified reports (Wetterbom et al. 2006, Mills et al. 2006; Khan et al. 2015) that those of nearly all indels in coding sequences are multiples of 3 nt (Figure 6b and supplementary figures S9 and S10, Supplementary Material online). Frame preserving insertions except for those containing stop codons merely add residues to the encoding proteins, while nonframeshifting deletions excluding those containing start or stop codons simply remove residues. Since most indels encode IDRs, alterations in the number of amino acids caused by fixed indels are likely to be tolerated. The length distributions (Figure 6c and supplementary figures S11 and S12, Supplementary Material online) demonstrate that shorter indels predominate, in agreement with reported findings (Benner et al. 1993; Qian and Goldstein 2001; Khan et al. 2015) and almost all fixed indels are shorter than 19 nt.

Fig. 6.—Most indels encode IDRs, are multiples of 3 nt in length, and are short, and

amino acid compositions of indels resemble those of IDRs. The bars and line graphs represent

arithmetic averages of the six species. a-c) The label unknown signifies that the generation

mechanisms are unknown. d) The frequency of each amino acid encoded by all internal exons

("All"), that of fixed indels, and that of IDRs are represented by bars (left scale). The fold

changes in frequency relative to that of all internal exons are shown by line graphs (right

scale).

Amino Acid Compositions of Indels Resemble That of IDRs

As most indels encode IDRs and IDRs have a characteristic amino acid composition (Dunker et al. 2008), we calculated the amino acid compositions of indels (Figure 6d, supplementary figure S12, Supplementary Material online). Indels have amino acid compositions similar to that of IDRs in that order-promoting residues are depleted, while disorder-promoting residues appear more frequently than the average. For the data presented, the correlation coefficient between log fold changes of insertion and those of IDR is 0.776 (statistically significant at $P<10^{-4}$), while that between log fold changes of deletion and those of IDR is 0.826 (statistically significant at $P<10^{-5}$).

Long Internal Exons Tend to Have More Tandem Repeats

What accounts for the higher incidences of indels in longer internal exons? We found no grounds to suppose that intronisation, exonization, and homologous recombination occur more frequently in longer internal exons. Since tandem repeats account for a large fraction of indels (Figure 4) and longer internal exons generally experience more indels (Figure 5), we thought it possible that tandem repeats are more prevalent in long internal exons. We thus calculated the fraction of each internal exon occupied by tandem repeats and investigated its dependence on internal exon length. The results (Figure 7) verified the conjecture; the longer internal exons are, the higher the repeat frequency with all the correlation coefficients statistically significant at $P<10^{-2}$ and the conclusion remains unchanged in a longer length range (1-840 nt instead of 1-450 nt) (supplementary figure S14, Supplementary Material online).

To examine how much of the higher prevalence of IDRs in longer internal exons is explained by tandem repeats, we calculated the fractions of IDRs, tandem repeats within IDRs, and others within IDRs (i.e., IDRs that are not in tandem repeat segments) in each length bin in the six species in the 1-450 nt (Figure 7) and 1-840 nt length ranges (supplementary figure S14, Supplementary Material online). Note that the previously reported dependence of the fraction of IDRs on internal exon length (Fukuchi et al. 2023) was more pronounced in the longer exon length range. Since the IDRs are divided into tandem repeats (within IDRs) and others, the sum of the fractions of the two constituents is equal to that of IDRs and, consequently, the sum of the slopes of the regression lines of the two is identical to that of

IDRs. If the positive correlation of IDRs are entirely accounted for by tandem repeats, the slope of the regression line of tandem repeats within IDRs will be the same as that of IDRs, whereas the slope of others will be zero, that is, the fraction of others will be constant irrespective of intron length. The observed slope of the regression line of the fraction of tandem repeats on IDRs was 0.438 per cent/unit, where 1 unit equals 30 nt, and that of others in IDRs was 0.478, while that of the corresponding slope of the fraction of IDRs was 0.916. Thus, only 48% of the dependence of IDRs on internal exon length is contributed by tandem repeats. As tandem repeats constitute a mere 28.3% of IDRs on average, however, they make a disproportionate contribution to IDRs. On the other hand, in the longer range of exon length (supplementary figure S14a, Supplementary Material online), the contribution ratio of tandem repeats in IDRs was 36%, lower than 48%.

Fig. 7.--Long internal exons tend to have tandem repeats as well as IDRs. The fractions of tandem repeats, IDRs, tandem repeats in IDRs, and others in IDRs are shown with the correlation coefficients of the fractions with exon length. Asterisks signify statistical significances as in Fig. 5 legend. a) Arithmetic averages are plotted with regression lines (dotted lines).

The Frequency of Indels Correlates with that of Tandem Repeats

Since both the frequency of indel and that of tandem repeats show positive correlation with internal exon length, the two frequencies are likely to be correlated. The correlations between the normalized insertion/deletion frequency and the normalized repeat frequency are shown by scatter plots (Figure 8a, supplementary figure S15, Supplementary Material online). The correlation coefficient between insertion and repeat frequencies is 0.597 (statistically significant at $P<0.05$), while that between deletion and repeat frequencies is 0.885 (statistically significant at $P<10^{-5}$). In confirmation of the expectation, the indel frequencies are significantly correlated with repeat frequency, with the deletion frequency more strongly so.

Fig. 8.--Indel frequencies are correlated with tandem repeat frequency and a proposed model. a) Scatter plots of the normalized repeat frequency of each length bin and the corresponding normalized insertion (yellow dots) and deletion (blue dots) frequencies are shown. Dotted lines (yellow: insertion, blue: deletion) represent regression lines. b-d) Proposed evolutionary steps that had led to the last eukaryotic common ancestor that have long internal exons that frequently encode IDRs. e) An ongoing process.

Discussion

We unexpectedly discovered that fixed indels occur preferentially in longer internal exons and large fractions of indels were generated by alterations in tandem repeats. We also found a high prevalence of tandem repeats in long internal exons, which accounts for the previously observed tendency of long internal exons to encode IDRs (Fukuchi et al. 2022) as tandem repeats mostly encode IDRs (Simon and Hancock 2009). As the coding sequences in prokaryotes rarely have tandem repeats (Albà, Tompa, and Veitia 2007), the first eukaryotic common ancestor (FECA) probably had a low repeat frequency in the coding sequences. After the acquisition of introns, we speculate that many exons had elongated by repeat expansion, as the countervailing repeat contraction initially occurred less often, and that the resultant increase in IDRs in proteins had facilitated further evolution. In view of the findings of this research, we prose that long internal exons had resulted from primordial short exons mainly by repeat expansion but are currently subject to frequent contractions and, to a lesser extent, expansions due to alterations in tandem repeats (Figure 8b-e).

That deletion generally exceeds insertion (Figure 5) is in agreement with previous reports (de Jong and Rydén 1981; Redelings et al. 2024). The number of human indels in ALL exons identified in this investigation is the sum of the number of human insertions and that of chimpanzee deletions at step 5, and is 863 (supplementary table S2, Supplementary Material online), which is larger than the reported number, 262, of human indels in all coding exons (Mills et al. 2006). The increase in number is attributable not only to revised sequence data we had the good fortune to use, but also to the methodology that identifies changed alternative splicing in addition to genomic alterations in constitutively spliced exons.

As nearly all fixed indels are nonframeshifting (Figure 6b), indels generated by tandem repeats mostly result in tandem repeats in proteins. Since protein tandem repeats play crucial roles in ligand binding and transcriptional regulation (Kobe and Kajava 2001; Yagi et al. 2014), indels generated by tandem repeats may have important biological significance. We consider the fractions of fixed indels generated by tandem repeats in exons represent an underestimate; slight sequence changes after repeat expansion and contraction make tandem repeats imperfect, making them undetectable by the repeat detection program. It is thus conceivable that indels generated by unknown mechanisms contain those attributable to tandem repeats, although they

may well include some cases generated by inverted and mirror repeats (Burssed et al. 2022), which are undetectable by the program we used. In fact, the fractions of IDRs in indels generated by alterations in tandem repeats are comparable to those produced by unknown mechanisms (Figure 6a), their length distributions are nearly identical (Figure 6c), and amino acid compositions of the two groups are similar (supplementary figure S16, Supplementary Material online). It is thus plausible that many indels whose generation mechanisms remain unidentified were really generated by alterations in tandem repeats. An underestimation of tandem repeats can explain why the tandem repeats explain only 48% (in the internal exon length range 1-450 nt, Figure 7a) or even less in the longer (1-840 nt) exon length range of the dependence of IDRs on the length of internal exons; unidentified tandem repeats may account for some of the remaining dependence.

Notwithstanding all the emphasis on tandem repeats, we would like to reiterate that intronisation and exonisation also generate deletions and insertions, respectively, and they account for considerable fractions of fixed insertions in *Homo sapiens*, fixed deletions in chimpanzee, and fixed indels in rat (Figure 4). The fact that many indels were "probably'" generated by intronisation and exonisation makes it quite likely that they really account for some indels. The wide variation in the fractions of intronisation and exonisation is largely explainable by disparity in location distribution of indels within internal exons; most insertions at 5' and 3' ends of exons and those involving entire exons were judged to be generated by exonisation (70.1, 77.2 and 58.6%, respectively), and nearly all deletions at 5' and 3' ends of exons and those of entire exon(s) were attributable to exonisation (85.0, 91.1 and 97.9%, respectively). By contrast, intronisation and exonisation were seldom assigned as the generation mechanisms of indels within exons (0.7 and 1.0%, respectively). The insertions in human and rat and the deletions in chimpanzee and rat have high total proportions of those at the ends of exon and those involving entire exons (supplementary figure S2, Supplementary Material online), and these have comparatively high fractions of cases generated by intronisation/exonisation. Indel analyses of other species are needed to judge whether intronisation and exonisation generally generate less indels than tandem repeats do.

Although the frequencies of the detected fixed indels are small (less than 0.06%), they may well be an underestimate as our methods of selecting fixed indels probably capture only a

fraction of actual fixed indels. This is both because transcripts with corresponding segments must be present in all three species in our identification method and because BLASTN alignments can be made only for highly conserved nucleotide sequences. Additionally, the outgroup requirement means that the capture rate of fixed indels varies from one species group from another and thus makes it inappropriate to compare indel frequencies of different groups; we for instance cannot meaningfully compare the indel frequencies of human and mouse. Moreover, the availability of more variant data may disqualify some fixed indels due to the presence of counterexamples in the newly identified variants. Since the methodology has no bias on exon length, however, the finding that higher indel fluency is observed in longer internal exons is unaffected by these shortcomings.

As the general preponderance of deletion rates over insertion rates is probably unaffected by methodological limitations, either, what accounts for the phenomenon? We consider it possible that insertion is evolutionarily disfavored especially in long internal exons because splicing of many excessively long exons (>300 nt) is inhibited (Robberson et al. 1990; Chen and Chasin 1994; Sterner et al. 1996). This interpretation dovetails with the observed tendencies that insertion frequency of internal exons initially rises but levels off at around 300 nt, while deletion frequency shows a monotonically increasing trend (Figure 5a). However, we excluded fixed insertions that create entire exons (Figure 2c, g) and fixed deletions that eliminate whole exons (Figure 2n, r) from our analyses of exon length dependence and our analysis cannot capture cases of exon fusion and fission that do not affect coding sequences. Although analysis of exon evolution including these neglected cases is expected to elucidate exon evolution dynamics in its entirety, it is beyond the scope of the current study.

The existence of a positive correlation between tandem repeat frequency and internal exon length was an interesting finding. Since four other eukaryotes, *Oryza sativa*, *Arabidopsis thaliana*, *Caenorhabditis elegans*, and *Schizosaccharomyces pombe* were shown to have a positive correlation of the fraction of IDRs and internal exon length (Fukuchi et al. 2023), we checked if repeat frequency and internal exon length are correlated in the 1-450 nt range in these model eukaryotes, too. We found the first three model eukaryotes have statistically significant ($P<0.01$) positive correlations, but *S. pombe* does not (supplementary figure S17, Supplementary Material online). The case of *S. pombe* is apparently inconsistent with the

notion that long internal exons were produced mainly by the expansion of tandem repeats. However, as stated above, the fraction of tandem repeats is likely to be underestimated, and the underestimation can explain the lack of correlation in the yeast species.

As far as we are aware, this represents the first systematic identification of fixed indels including those generated by constitutive alterations in alternative splicing and examination of the dependence of their frequencies on internal exon length. As our observations are limited to indels in six species only, the generality of our findings needs to be tested by indels in other species. Since the current selection method requires the presence of a trio of closely related species with completely sequenced genomes and an abundance of sequenced variants, however, it is not easy to apply the current methodology to many other species. For instance, the application of our methods to *C. elegans*, *C. briggsae* with *C. japonica* as the outgroup identified less than 70 cases of fixed insertions in *C. elegans*, a number judged too small for further analyses. Hopefully, widespread availability of variant and genome sequences will make extensive testing possible.

Materials and Methods

Sequence and Other Data Used

For the selection of *Homo sapiens* and *Pan troglodytes* (chimpanzee) indels, *Gorilla gorilla* was chosen as the outgroup. As the outgroup of *Mus musculus* (mouse) and *Rattus norvegicus* (rat), *Peromyscus maniculatus* was used, while *Drosophila yakuba* served as the outgroup of *Drosophila melanogaster* and *Drosophila simulans.* All the sequence and exon data, and the lists of orthologous genes except for that between *D. melanogaster* and *D. simulans* were downloaded from the Ensembl database (Martin et al. 2023). As a list of orthologous genes between the two *Drosophila* species was unavailable from the Ensemble database, we selected pairs of orthologous genes by mutual best hit of BLASTN alignments of the longest variants. The list of transposable elements used was in release 19 of the TREP Transposable Element Platform (Wicker et al. 2002).

Selection of Fixed Indels and Their Classification by Location

The actual selection steps followed are schematically shown (supplementary figure S1, Supplementary Material online). BLASTN alignments (Altschul et al. 1990) were performed with the default parameters except that the gap-opening penalty was set at 10 and the expectation value was changed to 0.001, unless otherwise stated. We identified indel candidates in BLASTN alignments with special considerations for sections of multiple alignments; we regarded two aligned sections continuous if the last query residue of one section coincides with the first query residue of the other within 3 nt. (The 3 nt allowance was made to cope with the uncertainty in alignments.) We judged indel sections identical if their genomic addresses agreed within 3 nt. In the uniformity tests, variants that do not have sections covering indels were disregarded. An indel was regarded as coinciding with the 5' or 3' end of an exon of sp. 1 if the start or the end of the segment coincides with the start or the end of the exon within 3 nt. An indel without coincidence with exon border(s) is judged to be located inside an exon.

Generation Mechanisms of Fixed Indels

An insertion is considered to be "possibly" generated by exonisation if an intron whose length is more than or equal to two thirds of the length of the insertion exists in sp. 2 at the corresponding location but is upgraded to "probably" generated by exonisation if at least two thirds of the residues were aligned (Figure 2a, c-e, g, h). Likewise, a deletion is regarded as "possibly" generated by intronisation if the corresponding segment in sp. 2 has an intron of at least two thirds of the length of the deleted segment but is reclassified as "probably" generated by intronisation if the aligned segment is more than or equal to two thirds of the length of the deleted segment (Figure 2l, n-p, r, s).

The generation mechanisms of the rest of the fixed indels were identified as follows. Homologous recombination is chosen as the generation mechanism of an insertion in sp. 1 based on the following conjecture (supplementary figure S18, Supplementary Material online); sequence A with insertion generated by homologous recombination in sp. 1 is probably more like sequence A' in sp. 1 that replaced the sequence than to the orthologous sequence B in sp. 2. Based on this idea, we set two criteria for an insertion to be a product of homologous recombination. Firstly, there must be at least one sequence A' of a different gene in sp. 1 aligned to the sp. 2 orthologue B, and secondly the fraction of the identity between the sequences of the insertion-containing exon and A' is not significantly lower than that of the exon and B sequences (chi-square test). Similarly, a deletion in sp. 2 is considered to have been generated by homologous recombination if 1) sequence B in sp. 2 with an exon containing the deletion has at least one sequence B' mapped to a different gene in sp. 2 that is aligned to the sp. 1 orthologue A and 2) the fraction of sequence identity between the exon and B' is not significantly lower than that between the exon and A (chi-square test).

Tandem repeats were identified in coding sequences by Tandem Repeats Finder (Benson 1999) with the mismatch and indel penalties and maximum period size set at 5, 3, and 2000, respectively, while the other parameters were at default settings. If there are segments of tandem repeats in the fixed indels, the tandem repeats were regarded as responsible for generating the indels.

Assignment of IDRs and statistical analyses

IUPred3 (Erdős et al. 2021) was run and amino acids with a score more than or equal to 0.502 were judged to be in IDRs. All the tests of statistical significance were carried out by t-test unless otherwise stated. For the data shown in Figures 5a, 8a and supplementary figures S6a and S15a, Supplementary Material online, each frequency of a length bin was first divided by the average frequency of the species and the normalized values of the six species were arithmetically averaged.

Supplementary Material

Supplementary data are available online.

Acknowledgments

This work was supported in part by Grant-in-Aid for Transformative Research Areas 'Multifaceted proteins: Expanding and transformative protein world', from the Ministry of Education, Culture, Sports, Science and Technology in Japan, 20H05932.

Data Availability

All data and code necessary to reproduce these anayses are available at Figshare; the data as well as a program list, the programs for primates, rodents, *Drosophila* species, and the remaining four eukaryotes have been respectively deposited at

https://figshare.com/account/articles/27366162,

https://figshare.com/account/articles/27410016,

https://figshare.com/account/articles/27411264,

https://figshare.com/account/articles/27411279, and

https://figshare.com/account/articles/27411291.

References

Abebrese EL, Ali SH, Arnold ZR, Andrews VM, Armstrong K, Burns L, Crowder HR, Day RT Jr, Hsu DG, Jarrell K, et al. Identification of human short introns. PLoS One 2017: 12(5):e0175393. https://doi: 10.1371/journal.pone.0175393

Albà MM, Tompa P, Veitia RA. Amino acid repeats and the structure and evolution of proteins. Genome Dyn. 2007:3:119-130. https://doi.org/10.1159/000107607

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990:215(3):403-410. https://doi.org/10.1016/S0022-2836(05)80360-2

Anbo H, Sato M, Okoshi A, Fukuchi S. Functional segments on intrinsically disordered regions in disease-related proteins. Biomolecules 2019:9(3):88. https://doi.org/10.3390/biom9030088

Anzai T, Shiina T, Kimura N, Yanagiya K, Kohara S, Shigenari A, Yamagata T, Kulski JK, Naruse TK, Fujimori Y, et al. Comparative sequencing of human and chimpanzee MHC class I regions unveils insertions/deletions as the major path to genomic divergence. Proc Natl Acad Sci U S A. 2003:100(13):7708-7713. https://doi.org/10.1073/pnas.1230533100

Basile W, Sachenkova O, Light S, Elofsson A. High GC content causes orphan proteins to be intrinsically disordered. PLoS Comput Biol. 2017:13(3):e1005375. https://doi.org/10.1371/journal.pcbi.1005375

Benson G. Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids Res. 1999:27(2):573-580. https://doi.org/10.1093/nar/27.2.573

Benner SA, Cohen MA, Gonnet GH. Empirical and structural models for insertions and deletions in the divergent evolution of proteins. J Mol Biol. 1993:229(4):1065-1082. https://doi.org/10.1006/jmbi.1993.1105

Berget SM. Exon recognition in vertebrate splicing. J Biol Chem. 1995:270(6):2411-2414. doi: https://doi.org/10.1074/jbc.270.6.2411

Burssed B, Zamariolli M, Bellucco FT, Melaragno MI. Mechanisms of structural chromosomal rearrangement formation. Mol Cytogenet. 2022:15(1)23. https://doi.org/10.1186/s13039-022-00600-6

Chen IT, Chasin LA. Large exon size does not limit splicing in vivo. Mol Cell Biol. 1994:14(3):2140-2146. https://doi.org/110.1128/mcb.14.3.2140-2146.1994

De Conti L, Baralle M, Buratti E. Exon and intron definition in pre-mRNA splicing. Wiley Interdiscip Rev RNA. 2013:4(1):49-60. doi: https://doi.org/10.1002/wrna.1140*

de Jong WW, Rydén L. Causes of more frequent deletions than insertions in mutations and protein evolution. Nature 1981:290(5802):157-159. doi: 10.1038/290157a0

Dunker AK, Oldfield CJ, Meng J, Romero P, Yang JY, Chen JW, Vacic V, Obradovic Z, Uversky VN. The unfoldomics decade: an update on intrinsically disordered proteins. BMC Genomics. 2008:9(Suppl. 2): S1. https://doi.org/10.1186/1471-2164-9-S2-S1

Erdős G, Pajkos M, Dosztányi Z. IUPred3: prediction of protein disorder enhanced with unambiguous experimental annotation and visualization of evolutionary conservation. Nucleic Acids Res. 2021: 49(W1):W297–W303. https://doi.org/10.1093/nar/gkab408

Fan Y, Wang W, Ma G, Liang L, Shi Q, Tao S. Patterns of insertion and deletion in mammalian genomes. Curr Genomics. 2007:8(6):370-378. https://doi.org/10.2174/138920207783406479

Fukuchi S, Noguchi T, Anbo H, Homma K. Exon elongation added intrinsically disordered regions to the encoded proteins and facilitated the emergence of the last eukaryotic common ancestor. Mol Biol Evol. 2023:40(1):msac272. https://doi.org/10.1093/molbev/msac272

Hawkins JD. A survey on intron and exon lengths. Nucleic Acids Res. 1988:16(21):9893-9908. https://doi.org/10.1093/nar/16.21.9893

Khan T, Douglas GM, Patel P, Nguyen BA, Moses AM. Polymorphism analysis reveals reduced negative selection and elevated rate of insertions and deletions in intrinsically disordered protein regions. Genome Biol Evol. 2015 :7(6):1815-1826. https://doi.org/10.1093/gbe/evv105

Kobe B, Kajava AV. 2001. The leucine-rich repeat as a protein recognition motif. Curr Opin Struct Biol. 2001:11(6):725-732. https://doi.org/10.1016/s0959-440x(01)00266-4

Kondrashov FA, Koonin EV. 2003. Evolution of alternative splicing: deletions, insertions and origin of functional parts of proteins from intron sequences. Trends Genet. 2003:19(3):115-119. https://doi.org/10.1016/S0168-9525(02)00029-X

Lang KM, Spritz RA. RNA splice site selection: evidence for a 5' leads to 3' scanning model. Science. 1983:220(4604):1351-1355. https://doi.org/10.1126/science.6304877

Levinson G, Gutman GA. Slipped-strand mispairing: a major mechanism for DNA sequence evolution. Mol Biol Evol. 1987:4(3):203-221. https://doi.org/10.1093/oxfordjournals.molbev.a040442

Li WH. 1997. Molecular Evolution. Sinauer Associates.

Li X, Liu S, Zhang L, Issaian A, Hill RC, Espinosa S, Shi S, Cui Y, Kappel K, Das R, et al. A unified mechanism for intron and exon definition and back-splicing. Nature. 2019:573(7774):375-380. https://doi.org/10.1038/s41586-019-1523-6

Light S, Sagit R, Ekman D, Elofsson A. Long indels are disordered: a study of disorder and indels in homologous eukaryotic proteins. Biochim Biophys Acta. 2013A:1834(5):890-897. https://doi.org/10.1016/j.bbapap.2013.01.002

Light S, Sagit R, Sachenkova O, Ekman D, Elofsson A. Protein expansion is primarily due to indels in intrinsically disordered regions. Mol Biol Evol. 2013B:30(12):2645-2653. https://doi.org/10.1093/molbev/mst157

Lin L, Shen S, Tye A, Cai JJ, Jiang P, Davidson BL, Xing Y. Diverse splicing patterns of exonized Alu elements in human tissues. PLoS Genet. 2008:4(10):e1000225. https://doi.org/10.1371/journal.pgen.1000225

Lin M, Whitmire S, Chen J, Farrel A, Shi X, Guo JT. Effects of short indels on protein structure and function in human genomes. Sci Rep. 2017:7(1):9313. https://doi.org/10.1038/s41598-017-09287-x.

Marquez Y, Höpfler M, Ayatollahi Z, Barta Z, Kalyna M. Unmasking alternative splicing inside protein-coding exons defines exitrons and their role in proteome plasticity. Genome Res. 2015:25(7):995-1007. https://doi.org/10.1101/gr.186585.1147

Martin FJ, Amode MR, Aneja A, Austine-Orimoloye O, Azov AG et al. Ensembl 2023. Nucleic Acids Res. 2023:51(D1):D933-D941. https://doi.org/10.1093/nar/gkac958

Mills RE, Luttig CT, Larkins CE, Beauchamp A, Tsui C, Pittard WS, Devine SE. An initial map of insertion and deletion (INDEL) variation in the human genome. Genome Res. 2006:16(9):1182-1190. https://doi.org/10.1101/gr.4565806

Mullaney JM, Mills RE, Pittard WS, Devine SE. Small insertions and deletions (INDELs) in human genomes. Hum Mol Genet. 2010:19(R2):R131-136. https://doi.org/10.1093/hmg/ddq400

Nekrutenko A, Li WH. Transposable elements are found in a large number of human protein-coding genes. Trends Genet. 2001:17(11):619-621. https://doi.org/10.1016/s0168-9525(01)02445-3

Pascarella S, Argos P. Analysis of insertions/deletions in protein structures. J Mol Biol. 1992:224(2):461-471. https://doi.org/10.1016/0022-2836(92)91008-d

Qian B, Goldstein RA. Distribution of Indel lengths. Proteins. 2001:45(1):102-104. https://doi.org/10.1002/prot.1129

Redelings BD, Holmes I, Lunter G, Pupko T, Anisimova M. Insertions and deletions: Computational methods, evolutionary dynamics, and biological applications. Mol Biol Evol. 2024:41(9):msae177. https://doi.org/ 10.1093/molbev/msae177

Robberson BL, Cote GJ, Berget SM. Exon definition may facilitate splice site selection in RNAs with multiple exons. Mol Cell Biol. 1990:10(1):84-94. https://doi.org/10.1128/mcb.10.1.84-94.1990

Simon M, Hancock JM. Tandem and cryptic amino acid repeats accumulate in disordered regions of proteins. Genome Biol. 2009:10(6):R59. https://doi.org/10.1186/gb-2009-10-6-r59

Sorek R, Ast G, Graur D. Alu-containing exons are alternatively spliced. Genome Res. 2002:12(7):1060-1067. https://doi.org/10.1101/gr.229302

Sorek R. The birth of new exons: mechanisms and evoutionary consequences. RNA 2007:13(10):1603-1608. https://doi.org/10.1261/rna.682507

Sterner DA, Carlo T, Berget SM. Architectural limits on split genes. Proc Natl Acad Sci U S A. 1996:93(26):15081-15085. https://doi.org/10.1073/pnas.93.26.15081

Tompa P. Intrinsically unstructured proteins evolve by repeat expansion. BioEssays 2003:25(9):847-855. https://doi.org/10.1002/bies.10324.

Ward JJ, Sodhi JS, McGuffin LJ, Buxton BH, Jones DT. Prediction and functional analysis of native disorder in proteins from the three kingdoms of life. J Mol Biol. 2004: 337(3): 635-645. https://doi.org/10.1016/j.jmb.2004.02.002

Wetterbom A, Sevov M, Cavelier L, Bergström TF. Comparative genomic analysis of human and chimpanzee indicates a key role for indels in primate evolution. J Mol Evol. 2006:63(5):682-690. https://doi.org/10.1007/s00239-006-0045-7

1   Wicker T, Matthews DE, Keller B. TREP: a database for Triticeae repetitive elements. Trends

2   Plant Sci. 2022:7(12):561-562. https://doi.org/10.1016/S1360-1385(02)02372-5

3   Yagi Y, Nakamura T, Small I. The potential for manipulating RNA with pentatricopeptide

4   repeat proteins. Plant J. 2014:78(5):772-782. https://10.1111/tpj.12377