HER2-driven mammary tumorigenesis enhances bioenergetics despite reductions in mitochondrial content

Sara M Frangos[1], Henver S Brunetta[1], Dongdong Wang[2,3], Maria Joy Therese Jabile[2,3], Leslie Jeffries[4], Grace Mencfeld[4], David W L Ma[1], William J Muller[5], Cezar M Khursigara[6], Kelsey H Fisher-Wellman[7], Jim Petrik[4], Gregory R Steinberg[2,3], Graham P Holloway[1]

[1]Department of Human Health Sciences, University of Guelph, ON, Canada
[2]Centre for Metabolism, Obesity and Diabetes Research, McMaster University, Hamilton, ON, Canada
[3]Division of Endocrinology and Metabolism, Department of Medicine, McMaster University, Hamilton, ON, Canada
[4]Department of Biomedical Sciences, University of Guelph, ON, Canada
[5]Department of Biochemistry, McGill University, Rosalind and Morris Goodman Cancer Research, Montreal, QC, Canada
[6]Department of Molecular and Cellular Biology, University of Guelph, ON, Canada
[7]Department of Cancer Biology, Medical Center Boulevard, Wake Forest University School of Medicine, Winston-Salem, NC, USA
Corresponding authors: Sara M Frangos (sfrangos@uoguelph.ca) and Graham P Holloway (ghollowa@uoguelph.ca)

ABSTRACT

It is now recognized that mitochondria play a crucial role in tumorigenesis, however, it has become clear that tumor metabolism varies significantly between cancer types. The failure of recent clinical trials aimed at directly targeting tumor respiration through oxidative phosphorylation inhibitors underscores the critical need for further studies providing an in-depth evaluation of mitochondrial bioenergetics. Accordingly, we comprehensively assessed the bulk tumor and mitochondrial metabolic phenotype in murine HER2-driven mammary cancer tumors and benign mammary tissue. Transcriptomic and proteomic profiling revealed a broad downregulation of mitochondrial genes/proteins in tumors, including OXPHOS subunits comprising Complexes I-IV. Despite reductions in tumor mitochondrial proteins, mitochondrial respiration was several-fold higher compared to benign mammary tissue, which persisted regardless of normalization method (wet weight, total protein content and when corrected for mitochondrial content). This upregulated respiratory capacity could not be explained by OXPHOS uncoupling, suggesting HER2 signaling regulates intrinsic mitochondrial bioenergetics. In further support, lapatinib, an EGFR/HER2 tyrosine kinase inhibitor, attenuated mitochondrial respiration in NF639 murine mammary tumor epithelial cells. Together, this data highlights that the typical correlation between mitochondrial content and respiratory capacity may not apply to all tumor types and implicates HER2-linked activation of mitochondrial respiration supporting tumorigenesis in this model.

INTRODUCTION

The field of cancer metabolism has rapidly evolved in the last decade. A specific focus of many researchers has been on understanding the role of mitochondria in tumorigenesis with the goal to potentially target aspects of mitochondrial metabolism to reduce proliferation[1,2]. It was initially hypothesized that a universal characteristic of cancer cells was dysfunctional oxidative phosphorylation (OXPHOS), as proposed by Otto Warburg in the 1920s[3,4]. This theory spurred interest in studying mitochondria in various cancers, highlighting the diverse mitochondrial phenotypes in this context. While instrumental for OXPHOS, mitochondria are now recognized for their multifaceted roles in regulating cellular homeostasis, including supporting macromolecule biosynthesis[5], maintaining redox balance[6] and generating metabolites that regulate gene transcription[7]. Given their pleiotropic effects within the cell, there is growing recognition that mitochondria play an essential role in cancer initiation, progression, and metastasis[1], and an appreciation that interrogating mitochondrial bioenergetics in diverse cancers may identify novel therapeutic strategies.

Reductions in the content/activity of mitochondrial enzymes within the tricarboxylic acid (TCA) cycle
and electron transport chain (ETC) often correlate with impaired mitochondrial respiratory capacity and
coincides with activating transcriptional programs driving tumorigenesis[8–10]. Indeed, inhibition of
succinate dehydrogenase (Complex II) in osteosarcoma cells[8], and renal cell carcinoma[9] enhances cancer
growth despite OXPHOS inhibition. Furthermore, in colorectal cancer, age-related Complex I deficiency
coincides with reduced respiratory capacity but enhanced serine synthesis and subsequent growth rates[10].
Despite this evidence suggesting that disruptions in mitochondrial bioenergetics contribute to tumor
progression, several elegant studies using genetic and pharmacological approaches have solidified that a
functional ETC is required for cancer cell growth [6,11,12]. Furthermore, the use of whole-genome sequencing
specifically interrogating mtDNA phenotypes across several human cancers has revealed that loss of
function mutations in mtDNA genes are often under negative selection[13], suggesting the maintenance of
mitochondrial function may support tumorigenesis in some cancers and impairments in mitochondrial
metabolism are not a ubiquitous finding. However, mitochondrial content represents one dimension of
bioenergetic regulation and does not alone determine respiratory capacity. Therefore, a comprehensive
assessment of mitochondrial function, particularly within the context of the mitochondrial proteome, is
warranted to provide deeper insights into the relationship between mitochondrial content and
bioenergetics in different cancers.

Within this framework, there has been considerable interest in investigating mitochondrial
bioenergetics and dynamics in breast cancer subtypes due to the limited therapeutic options for triple-
negative breast cancer (TNBC) and therapeutic resistance in breast cancers driven by estrogen (ER),
progesterone (PR) and human epidermal growth factor (HER2) receptor amplification[14–16]. Indeed,
increases in mitochondrial respiration or "OXPHOS reliance" is associated with poor outcomes in
treatment naïve TNBC[17], metastatic ER+ tumors[18], and HER2-driven breast cancers[19]. While inhibiting
OXPHOS in this context[17,18] (i.e. with Complex I inhibitor IACS-010759) has shown some pre-clinical
success, systemic toxicity has prevented the translation of this strategy to the clinic[20]. Furthermore, it is
unclear whether increases in mitochondrial respiration arise from increases in mitochondrial content or
intrinsic enhancements in mitochondrial function (e.g. coupling efficiency, substrate delivery/sensitivity,
oncogenic signaling events). This information is critical to develop successful mitochondrial-targeted
therapies with a wider therapeutic window. Thus, in the current study, we comprehensively assessed the
bulk tumor and mitochondrial phenotype between murine HER2-driven mammary cancer tumors and
benign mammary tissue in the context of the underlying transcriptome and proteome. Respiratory
capacity supported by both pyruvate and lipids was enhanced in HER2-driven tumors despite lower
mitochondrial content compared to benign mammary tissue. Mitochondrial bioenergetics were supressed
by the dual tyrosine kinase inhibitor lapatinib, suggesting events linked to HER2-mediated oncogenic
signaling may drive increases in OXPHOS during tumorigenesis.

RESULTS

Mammary tumors from MMTV/neu[ndl]-YD5 mice display canonical HER2 pathway activation

Cell lines are commonly used to study isolated mechanisms driving tumorigenesis, however, they neglect
the variety of cell types in the tissue of origin and the influence of the tumor microenvironment, which is
a key regulator of cancer growth[21,22]. To this end, we have utilized tumor and benign mammary tissues
from MMTV/neu[ndl]-YD5 transgenic mice[23] to investigate HER2-driven tumor biology **(Fig.1A).** Signaling
events linked to HER2 activation rely on the autophosphorylation of the tyrosine kinase domain[23–25] and
MMTV/neu[ndl]-YD5 mice possess an add-back mutation at the Y1226/7 (YD) site, activating downstream
signaling events linked to SHC/GRB2 with similar transforming potential as the wild-type protein[25,26]
(Fig.1B). While the proximal signaling events linked to HER2 activation in this model have been described[25],
to validate our bulk tissue-based model, we assessed the mRNA (transcriptomics) and protein
(quantitative label-free proteomics and western blotting) expression of canonical HER2 pathway members
(RAS/MAPK and PI3K/AKT/mTOR pathways) that are well-characterized in various HER2-overexpressing
mammary cancer cell lines **(Fig.1B)**[27–29]. *Her2* transcript levels were increased ~175-fold (**Fig.1C**) and
transcripts belonging to the Ras/MAPK (*Hras, Mapk1, Mapk3*) and PI3K/AKT/mTOR (*Pik3cg, Pik3c2a,*

Pik3c2b, Pikrcg, Pik3c3, Akt3) pathways were also upregulated in the tumors compared to the benign
mammary tissue **(Fig.1D)**. At the protein level, HER2 was readily detectable in mammary tumors by
quantitative label-free proteomics but undetectable in paired benign tissues **(Fig.1E)** suggesting enhanced
HER2 transcription also manifests as an increase in receptor content in the tumors. Downstream of HER2
activation, the total protein content of p38 MAPK, ERK1/2 and eEF2 were significantly increased in tumors
compared to benign tissue detected by western blot **(Fig.1F,G)**. When normalized to total protein, p38
MAPK and eEF2 phosphorylation were not higher in tumors due to proportional increases in total protein
abundance, while in contrast, phosphorylated ERK1/2 and mTOR were significantly upregulated suggesting
both Ras/MAPK and the PI3k/AKT/mTOR pathways were activated in tumors **(Fig.1H, S-Fig.1)**. Overall,
these data demonstrate that mammary tumors display classical hallmarks of HER2-driven cancer cell
signaling, allowing for the subsequent investigation of HER2-driven tumor biology and metabolism in this
model.

Assessment of the HER2-driven mammary tumor transcriptome

To understand the biology and metabolism of HER2-driven mammary tumors, we assessed their
transcriptomes using RNA sequencing. Principal component analysis (PCA) using the entire gene
expression profile showed distinct separation between groups **(Fig.2A)**. Visualizing all differentially
expressed genes (DEGs) using a volcano plot **(Fig.2B)** and the top 3000 DEGs using a heatmap **(Fig.2C)**
revealed several differentially upregulated and downregulated genes between groups where genes
involved in MAPK- and AKT-linked signaling, ECM remodelling and cellular stress responses *(Etv1, Dusp6,*
Papln, Ier3) were among the most upregulated genes in the tumors and genes involved in lipid metabolism
(*Dgat2, Lipe, Cd36*) were among the most downregulated. Enrichment analysis of the top 3000
upregulated DEGs **(Fig.2D)** identified pathways characteristic of HER2-driven signaling (e.g. tyrosine kinase
signaling, GTPase signaling) and gene ontology (GO) analysis identified processes indicative of active
transcription, translation, and protein turnover characteristic of a rapidly proliferating tissue **(Fig.2E)**.
Interestingly, enrichment analysis of the top 3000 downregulated genes in the tumors consistently
identified that mitochondrial metabolism, and in particular components of the oxidative phosphorylation
(OXPHOS) system, were downregulated in tumors compared to benign mammary tissue **(Fig.2F,G)**.

Assessment of HER2-driven mammary tumor proteome

We further validated our bulk-tissue transcriptomic findings at the protein level using label-free
quantitative proteomics and detected 2589 proteins in the benign mammary tissue and 2448 proteins in
the HER2-driven tumors; 1848 of which were found in both groups **(Fig.3A)**. This initial analysis revealed
741 and 600 uniquely expressed (i.e. only detected in one tissue) proteins in benign mammary tissue and
tumors respectively **(S-Fig.2A)**. Interestingly, OXPHOS and aerobic respiration were identified among the
top pathways in benign tissue, but not in tumors **(S-Fig.2B-G)**, further supporting the notion of
downregulated mitochondrial content within tumors. Reflecting the PCA for the transcriptomic data,
analysis of all overlapping proteins between benign and tumor samples revealed distinct separation
(Fig.3B). Of the 1848 proteins found in both the tumors and benign tissue, 420 were significantly different
(p-adj. <0.1, **Fig 3A,C,D**). However, the majority of the differentially expressed proteins were
downregulated (n=411) in tumors compared to benign mammary tissue. Upregulated proteins (n=9)
included those involved in canonical HER2 signaling, protein synthesis and processing, and glycolytic
metabolism **(S-Table 1)**. Supporting transcriptomic data, which indicated a reduction in mitochondrial
content and OXPHOS, enrichment analysis revealed that 8 of the top 10 downregulated pathways **(Fig. 3E)**
and all top 10 downregulated GO **(Fig. 3F)** were related to mitochondrial biogenesis and/or metabolism,
and OXPHOS was identified in the top 10 downregulated KEGG pathways **(Fig.3G)**. Together, these
proteomic data support our transcriptomic analysis suggesting HER2-driven tumors are characterized by
lower mitochondrial content than benign mammary tissue.

Integrated assessment of the HER2-driven tumor transcriptome and proteome

The concordant nature of our transcriptomic and proteomic datasets was highlighted when correlating the
log2FC between overlapping gene and protein targets **(Fig.4A)**. Sorting these data by mitochondrial

proteins highlighted that the majority of overlapping mitochondrial proteins detected in both datasets
were downregulated **(Fig.4B)**. Thus, we further analyzed the 144 overlapping downregulated targets
between these analyses **(Fig.4C)**. KEGG enrichment analysis of the downregulated targets identified
OXPHOS among the top 10 pathways, as well as carbon metabolism, BCAA degradation and fatty acid
metabolism which encompass numerous mitochondrial targets **(Fig.4D)**. The top 10 common
downregulated pathways **(Fig.4E)** and GO **(Fig.4F)** all pertained to mitochondrial substrate
metabolism/catabolism and OXPHOS. Together, transcriptomic and proteomic analyses strongly support
that HER2-driven tumors are characterized by downregulated mitochondrial content.

Assessment of the mitochondrial transcriptome and proteome

Our initial transcriptomic and proteomic analyses encompassed all differentially expressed genes/proteins
between tissues (including mitochondrial targets) and clearly identified reductions in mitochondrial
content in the tumors. This emphasized the robust nature of mitochondrial adaptations in this model,
considering mitochondrial proteins (identified using the mouse MitoCarta 3.0 database) represented a
small fraction of total cellular protein in both tumors and benign mammary tissue **(S-Fig.3A)**. By filtering
our proteomic dataset using MitoCarta, we identified 125 mitochondrial proteins that were commonly
expressed between the tumors and benign mammary tissue **(Fig.5A)**. When visualizing the 125 proteins
with a heatmap, the majority of the mitochondrial proteome was downregulated in the tumors compared
to the benign mammary tissue **(Fig.5B)**. The summed absolute abundance of all detected mitochondrial
proteins in each group was also lower in tumors **(Fig.5C)**, again supporting our transcriptomic and
proteomic analyses suggesting a reduction in mitochondrial content exists in the tumors.

While mitochondrial biogenesis is a coordinated process simultaneously upregulating several
mitochondrial proteins, the total mitochondrial transcriptome/proteome does not necessarily change
proportionally with OXPHOS complex and/or TCA cycle/dehydrogenase content, especially in cancerous
tissues. Indeed, reductions in the content of certain respiratory complexes seem to be cancer specific,
which can have functional repercussions on mitochondrial respiration rates and/or efficiency[10,31]. Within
this context, we assessed the gene expression profile of all OXPHOS complex subunits between groups,
which revealed relatively ubiquitous reductions in transcript count (transcripts per million) across subunits
of Complexes I-IV in the tumors **(Fig.5D-H)**. Interestingly, while the majority of Complex V subunits were
also downregulated, this complex had the highest proportion of subunits displaying no differences
between groups **(Fig.5D,I)**. The TCA cycle transcriptome also reflected a general downregulation of
transcript abundance, apart from alpha-ketoglutarate dehydrogenase (*Ogdhl*) and succinyl-CoA
synthetase GDP forming subunit-β (*Suclg2*), which had a higher transcript abundance in some tumor
samples **(Fig.5J)**. While we focused on transcriptomic data for this analysis due to lower detection of
certain OXPHOS subunits using label-free quantitative proteomics **(S-Fig.4A-E)**, the summed abundance of
all OXPHOS subunits detected per complex expressed relative to the underlying mitochondrial proteome
(S-Fig.4F), and relative to the total proteome **(S-Fig.4G)** displayed a similar pattern. This interpretation was
further validated by western blot, where tumor CII and CIV content were lower than benign mammary
tissue **(S-Fig.4H).** Overall, these data further suggest that the mitochondrial proteome and OXPHOS
Complexes I-IV are downregulated in HER2-driven tumors.

While these findings strongly support reduced mitochondrial gene expression in tumors, differences could
theoretically result from shifts in cell-type during tumorigenesis. To address this, we first assessed OXPHOS
expression in a publicly available single-cell RNA sequencing (scRNA-seq) dataset from human breast
tumors[31]. In the scRNA-seq dataset, tumor epithelial cells expressed the highest proportion of CI-CIV
subunits when grouped by cell type **(Fig.6A,B).** We next assessed cell-type composition in our tumor and
benign mammary tissue samples by immunofluorescence **(Fig.6C)**. Although the immunopositive area was
lower across all cell types in benign samples**,** mesenchymal and epithelial cells remained the dominant
populations in both groups, indicating the absence of major shifts in the most abundant cell types **(Fig.6D).**
To integrate cell-type-specific OXPHOS mRNA expression (scRNA-seq) with cell-type abundance in our
samples, we calculated a cell-type-weighted OXPHOS mRNA Index, reflecting each population's estimated
contribution to Complex I–V transcript levels in our tissues. The OXPHOS mRNA Index **(Fig.6E)** indicated

that epithelial cells were likely the dominant contributors to OXPHOS transcripts in the present tumor
samples. When these values were summed across complexes to generate a Total OXPHOS mRNA Score,
epithelial cells remained the primary contributors **(Fig.6F).** Altogether, these analyses suggest that our
whole-tissue RNA-seq signatures are consistent with canonical HER2-driven signaling and indicate that the
reduction in OXPHOS gene expression observed by bulk-tissue RNA-seq likely cannot be explained by a
loss of high-OXPHOS epithelial cells or a gain in low-OXPHOS cell populations. Instead, these findings
support a cell-intrinsic suppression of mitochondrial gene expression in HER2-driven tumors.

Mitochondrial respiratory capacity is increased in HER2-driven mammary tumors

Considering our transcriptomic and proteomic data strongly suggest HER2-driven mammary tumors
display reductions in mitochondrial content, we next assessed whether this would manifest as a reduction
in respiratory capacity by measuring mitochondrial respiratory flux rates. In permeabilized tumors and
benign mammary tissue, we assessed basal/endogenous respiration and respiration with saturating
carbohydrate (pyruvate/malate) and lipid (palmitoyl-carnitine) substrates. For each protocol, several
biological replicates (individual tumors) were utilized, spanning a continuum of whole tumor wet
weights/volumes **(S-Fig.5A,B)**. Since no strong correlations were found between tumor wet weight or
volume and basal or maximal respiration supported by carbohydrate- or lipid-linked substrates **(S-Fig.5C-**
J), all data points were pooled. When normalizing respiration to tissue wet weight **(Fig. 7A)**, tumors
displayed significantly higher basal/endogenous respiration and higher respiration with subsequent
additions of pyruvate, malate (CI-linked respiration), ADP (State III respiration), glutamate (CI-linked
respiration also supported by alpha-ketoglutarate anaplerosis), and succinate (to maximize CII-linked
respiration) **(Fig.7B)**. Equally, tumors displayed significantly higher respiration rates supported by lipids
(palmitoyl-carnitine, **Fig.7C**), suggesting that HER2-driven tumors are capable of relying on both substrate
sources, perhaps to maximize aerobic metabolism. This initial analysis of respiratory rates was normalized
to tissue wet weight, which can be influenced by differences in total and mitochondrial protein content in
a given tissue volume. To address this, we first normalized respiratory flux rates to total cellular protein
(Fig.7D,E), which minimizes differences in factors such as tissue density and hydration that are not directly
contributing to mitochondrial oxygen consumption. The third normalization method employed was
normalizing respiratory flux to a mitochondrial enrichment factor (MEF) (**Fig.7F,G**) which is the ratio of
mitochondrial proteins to the underlying total proteome **(S-Fig.3B)**. This approach controls for differences
in mitochondrial content between tissues to determine whether increases in respiration are due to
increases in mitochondrial content or intrinsic changes in mitochondrial function. Interestingly, the
fundamental observation of increased respiratory flux rates in tumors compared to benign tissues was
conserved independent of the normalization method, suggesting that HER2-driven mammary tumors are
characterized by high respiratory flux rates independent of total cellular protein or reduced mitochondrial
protein content. While greater intrinsic OXPHOS was observed with all substrates, lipid supported
respiration was stimulated to a greater extent in the tumors, resulting in comparable carbohydrate
(pyruvate, malate, ADP) and lipid (palmitoyl-carnitine, malate, ADP) supported respiration **(Fig.8A)**. This
effect occurred despite many of the genes and proteins involved in lipid metabolism at the plasma
membrane and in the cytosol (e.g. *Acsl1, Adipoq, Dgat1/2, Lipe, Fabp4, Cd36, Pparg*) and mitochondria
(*e.g. Etfa, Etfb, Etfdh, Hadh, Hadhb, Cpt1b, Cpt2*) being downregulated in tumors compared to benign
mammary tissue **(Fig.8B-E)**. The intrinsic activation of respiration was not a ubiquitous finding for all
processes linked to OXPHOS, as reactive oxygen species (**S-Fig.6A**) emission rates were higher in tumors
compared to benign tissue when normalized to wet weight **(S-Fig.6B)**, lower when normalized to total
protein **(S-Fig.6C)** and not different when normalized to mitochondrial content **(S-Fig.6D)**.

High respiratory flux rates cannot be explained by OXPHOS uncoupling

Typically, respiratory capacity correlates with mitochondrial content in fully differentiated tissues[31].
However, given the apparent discrepancy between mitochondrial/OXPHOS protein content and
respiratory flux rates in HER2-driven tumors, we investigated alternative explanations for this observation.
One possible explanation for the observed enhanced respiratory rates could be due to the uncoupling of
OXPHOS from ATP synthesis, especially when ROS emission is not proportional to respiratory flux. To

address this possibility, we utilized oligomycin (OMY) to inhibit the F_0 subunit of ATP synthase (Complex V)
and examine its inhibitory potency between groups **(Fig.8F)**. While OMY inhibited respiration in benign
mammary tissue by ~55%, respiration in tumors was inhibited by ~80% **(Fig.8G)**. Subsequent additions of
antimycin A (AMA) to inhibit Complex III and rotenone (ROT) to inhibit Complex I, fully inhibited respiration
in both tissues **(Fig.8G)**. Since respiration was inhibited to a greater extent in the tumors in response to
OMY, this protocol strongly suggests that HER2-driven tumor mitochondria are highly coupled.

Tumor mitochondria display altered morphology

When assessing the subcellular ultrastructure of HER2-driven mammary tumors using transmission
electron microscopy (TEM), we observed smaller, more punctate mitochondria in tumors compared to
benign tissue **(Fig.9A)**. To determine whether ultrastructural observations aligned with the expression of
genes and proteins involved in mitochondrial turnover, we filtered our transcriptomic and proteomic
dataset using MitoCarta, specifically investigating mitochondrial proteins involved in mitochondrial
dynamics and mitophagy. This approach revealed the upregulation of several genes and proteins involved
in mitochondrial fission and generally a downregulation of genes and proteins involved in mitochondrial
fusion (**Fig.9B-F**). Together, these data show morphological differences in tumor versus benign tissue
mitochondria, which could potentially influence OXPHOS efficiency.

HER2 inhibition supresses tumor respiration

Although the exact mechanism(s) driving OXPHOS activation despite reductions in mitochondrial content
in our model remain unclear, we next asked whether HER2 activity directly regulates mitochondrial
bioenergetics. To address this, we used NF639 cells—an epithelial line derived from mammary tumors
from MMTV-c-neu transgenic mice, which, like our MMTV/neu[ndl]-YD5 model, display upregulated murine
HER2/neu signaling[32–35] serving as a highly relevant cell-line analog to our intact tumor measurements.
NF639 cells were treated with lapatinib, a dual HER2/EGFR tyrosine kinase inhibitor **(Fig.10A)** a well-
described and validated inhibitor of HER2-mediated signaling[29,36,37]. Cells were exposed to 250nM lapatinib
for 24 hours, a dose and time regimen that was validated and did not notably compromise viability
(Fig.10B). Using the same CI+CII linked respirometry protocol applied to the murine tumors **(Fig.7B),**
lapatinib significantly reduced leak respiration, CI-linked respiration, and CI+CII-supported respiration
relative to DMSO-treated cells **(Fig.10C)**. Importantly, lapatinib did not induce a cytochrome c (CC)
response (% change in respiration with CC, CON=-0.83%, LAP=-2.9%) and did not alter respiratory
inhibition with OMY, AMA, or ROT **(Fig.10C)**, indicating that mitochondrial integrity and coupling were
maintained. Collectively, these findings demonstrate that upregulated HER2-signaling directly stimulates
mitochondrial bioenergetics.

DISCUSSION

The aim of the current study was to investigate the metabolic characteristics of a mouse model of HER2-
driven mammary cancer compared to benign mammary tissue. Specifically, we focused on the
bioenergetic adaptations that have occurred with HER2-driven tumorigenesis both with respect to
mitochondrial function (respiration and ROS) and mitochondrial content (total mitochondrial proteome
and OXPHOS complexes). This investigation revealed several findings: 1) HER2-driven mammary tumors
exhibit high respiratory rates compared to benign mammary tissue, 2) tumors are capable of oxidizing
lipids and pyruvate to a similar extent despite the downregulation of several genes involved in lipid
catabolism, 3) high tumor respiration rates cannot be explained by higher mitochondrial content, and, 4)
HER2 signaling regulates the upregulation of tumor respiration. These data highlight that HER2-driven
mammary tumors exhibit high OXPHOS rates despite lower mitochondrial content, challenging the typical
bioenergetic relationship in fully differentiated tissues.

Relationship between mitochondrial content and function in breast cancer

Studies simultaneously assessing mitochondrial content and mitochondrial respiration, and the
contribution of these factors to breast cancer malignant transformation are scarce. While reductions in
ETC subunit content/activity have been described in mitochondria isolated from human mammary
carcinoma cultures compared to mammary epithelial cells[38], it has been assumed that a reduction in ETC
subunit content would manifest as a reduction in respiratory capacity. However, this is at odds with several
reports that classify breast cancer cells as OXPHOS reliant due to increases in mitochondrial respiration
correlating with advanced disease[19,39,40]. Indeed, across many cancers it is generally assumed that
mitochondrial content and respiratory function increase or decrease concomitantly, however, recent work
has challenged this dogma, demonstrating that high rates of respiration normalized to tissue weight can
be negated when corrected for mitochondrial mass,[41,42] revealing intrinsic alterations in cancer cell
bioenergetics. In the current study, utilizing this same approach to control for differences in mitochondrial
content, we found that respiratory capacity supported by pyruvate and lipids in HER2-driven mammary
tumors was significantly higher (~2-4 fold) compared to benign mammary tissue despite lower
mitochondrial enrichment (total mitochondrial proteome) and lower abundance of several respiratory
complex subunits, suggesting mitochondrial content independent increases in mitochondrial respiratory
capacity are acquired during HER2-driven tumorigenesis. This is a key finding independent of the specific
mechanisms underlying the enhanced bioenergetics.

While the exact mechanisms underlying the activation of OXPHOS in the current work remain unclear, our
findings allow us to speculate on several possibilities that could enhance respiration in a mitochondrial-
content–independent manner. It is unlikely these factors are mutually exclusive and include increased
coupling of electron transfer to ATP synthesis (coupling efficiency), changes in the delivery of substrates
across the mitochondrial membranes, intrinsic changes in the activity of ETC complexes and/or ATP
synthase, alterations in mitochondrial morphology/dynamics and signaling events linked to HER2
activation.

Possible mitochondrial content-independent mechanisms supporting enhanced respiration rates in
HER2-driven mammary cancer

Coupling Efficiency: Several studies have suggested that a key function of mitochondrial respiration in
cancer cells is the regeneration of NAD^+, FAD and ubiquinone to provide the necessary metabolites for
biomass production, rather than ATP synthesis[5,6,43]. From a bioenergetic perspective, this could be
achieved by inducing uncoupled respiration, which could manifest in an apparent increase in intrinsic
mitochondrial respiration, given that measures of respiratory flux quantify oxygen consumption rather
than ATP synthesis. However, in the present study the Complex V inhibitor oligomycin inhibited respiration
to a greater extent in HER2-driven tumors, suggesting that uncoupling cannot explain the enhanced
respiration rates observed, and if anything, tumors are more coupled than benign mammary tissue. This
interpretation aligns with work in MCF7 cells where mitochondrial ATP represents the majority of cellular
ATP production[44] suggesting the ATP synthetic function of OXPHOS in breast cancer is likely intact. Indeed,
enhanced mitochondrial ATP provision and OXPHOS-dependent biomass expansion are not mutually
exclusive processes and increased coupled respiration would satisfy both the bioenergetic and
biosynthetic requirements of rapid cell proliferation during tumorigenesis.

Substrate Delivery and Preference: Recent work has highlighted the dependency of HER2+ tumor cells on
both lipid (long chain fatty acids) and carbohydrate-derived (pyruvate) substrate for optimal proliferative
capacity[45]. In addition to pyruvate, HER2-driven tumors may be particularly poised to utilize lipids, given
the proximity of cancer cells to adipocytes in the mammary tumor microenvironment[46], which has spurred
interest in inhibiting lipid use/fat oxidation as an adjunct therapy in HER2-driven cancer[45]. Indeed, the use
of endogenous substrate may be preferred as solid tumors expand and factors such as increased interstitial
fluid pressure and extracellular matrix remodelling impact tumor vessel quality[47], possibly limiting
exogenous substrate delivery. However, at the transcript and protein level, enzymes involved in
mitochondrial lipid uptake (CPT1, CPT2) and pyruvate uptake (MPC) were largely downregulated
compared to benign tissue which would limit rather than enhance mitochondrial substrate delivery.
Despite this lower abundance of enzymes supporting mitochondrial substrate delivery, mitochondrial

respiration supported by pyruvate and lipids were significantly higher in tumors. Considering lipid and carbohydrate oxidation converge within the mitochondrial matrix, both providing acetyl-CoA to the TCA cycle, it is possible that enhanced OXPHOS efficiency lies at a distal site, implicating intrinsic OXPHOS adaptations as a possible explanation for enhanced respiration in HER2-driven tumors.

Intrinsic OXPHOS Adaptations: Several cancers are characterized by the altered expression of individual ETC subunits and/or ATP synthase which can have functional repercussions on respiratory function[10,41]. In the current study, despite a consistent downregulation of subunits comprising Complexes I-IV in HER2-driven tumors, a subset of Complex V genes were not downregulated (specifically two subunits of the c-ring of F_0 domain (*Atp5g1, Atp5g2*), two subunits of the F6 domain stalk connecting the F_0 and F_1 domains (*Atp5j1, Atp5j2*), and regulatory proteins ATPase inhibitory factor 1 (*Atpif1*) and distal membrane assembly component 2 (*Dmac2l*)). This shift essentially increases the ratio of ATP synthase/ETC and raises the possibility that a maintenance of these subunits may enhance OXPHOS rates in the face of downregulated CI-IV subunit content since maximal coupled respiration is usually limited by proton flux through ATP synthase. Interestingly this observation was further supported by our assessment of scRNA-seq data in human breast tumors, where CV expression was generally uniformly and highly expressed across major cell types, further supporting the notion that the maintenance of CV may be important for tumorigenesis. Indeed, despite mtDNA lesions in proton pumping complexes (CI, CIII and CIV) being common in many cancers, presumably reducing the content/function of these complexes, ATP synthase mutations are less common, suggesting positive selection may exist to maintain coupled respiration and ATP production[10,48,49]. Thus, while speculative, it is possible that higher respiration rates in HER2-driven tumors may partially reflect maintained ATP synthase content/function.

While bioenergetic efficiency is typically related to intrinsic changes in OXPHOS stoichiometry, mitochondrial morphology can indirectly affect efficiency. We observed smaller, punctate mitochondria in HER2-driven mammary tumors when visualized by TEM, generally aligning with reductions in mitochondrial fusion and increases in mitochondrial fission-related targets. While the direct relationship between mitochondrial morphology and bioenergetics remains contentious, evidence suggests smaller mitochondria (due to enhanced fission or reduced fusion) may be associated with enhanced bioenergetics. Smaller mitochondria tend to have a higher surface-area-to-volume ratio which has been correlated with upregulated OXPHOS in the liver[50], and low rates of respiration is associated with the inhibition of mitochondrial fission[51]. Furthermore, rare genetic mitochondrial diseases are often characterized by hyper-fused mitochondria which is associated with OXPHOS limitations[52]. Indeed, morphological differences in breast cancer mitochondria are well described[16,53–55] and targeting mitochondrial dynamics for therapy is an active area of study[56–58], however, there is a clear need to ascertain the relationship between mitochondrial dynamics and bioenergetics in tumorigenesis.

Intrinsic activation of OXPHOS mediated by HER2

No matter the mechanisms underlying our observed bioenergetic activation in HER2-driven tumors, our experiments in NF639 cells demonstrate that HER2 activation appears to regulate respiratory function. Although findings in the literature regarding how HER2 activation or suppression influences mitochondrial content and bioenergetics remain mixed, several lines of evidence support the idea that HER2 is important for tumorigenesis and respiratory competence[19,59–61]. Importantly, these studies collectively reinforce that changes in mitochondrial content alone cannot fully account for alterations in respiratory function, suggesting that HER2 may regulate intrinsic mitochondrial activity. Independent groups have shown that HER2 may localize to the inner mitochondrial membrane and directly interact with respiratory complexes (CI,CIV,CV) influencing their activity[19,59,60]. Alternatively, indirect activation of bioenergetics via HER2 signaling is another possibility – which is supported by work showing mitochondrial ADP availability may be enhanced via mitochondrial creatine kinase (MtCK) phosphorylation by HER2 downstream effectors[61]. Both mechanisms would alter bioenergetics without necessarily altering mitochondrial content, which aligns with our observation that content markers do not scale with respiratory capacity. Nonetheless, further research is required to more clearly delineate the relationship between HER2 signaling, OXPHOS

activation and possible mitochondrial content independent mechanisms conferring respiratory
competence.

CONCLUSION

We provide evidence that HER2-driven mammary tumors display enhanced respiratory capacity despite
lower mitochondrial content. This increased respiratory capacity could not be explained by changes in
OXPHOS coupling, however, differences in mitochondrial morphology and the maintained expression of
several subunits of ATP synthase suggest HER2-driven mammary tumors may undergo intrinsic
adaptations in mitochondrial function during tumorigenesis. No matter the mechanism driving OXPHOS
activation in our model, upregulated HER2 signaling is essential for the observed increases in
mitochondrial respiratory capacity.

MATERIALS AND METHODS

Animals and Breeding

All rodent experiments were conducted in accordance with institutional guidelines approved by the
Animal Care Committee at the University of Guelph (AUP#5071). MMTV/neu^{ndl}-YD5 mice on an FVB
background were obtained from Dr. William Muller (McGill University) and Dr. David Ma (University of
Guelph). Female MMTV/neu^{ndl}-YD5 mice were obtained by breeding male heterozygous MMTV/neu^{ndl}-YD5
mice with female FVB mice to yield WT or heterozygous transgenics. Mice were group housed in ventilated
cages at room temperature (22ºC) and kept on a 12:12 light-dark cycle with *ad libitum* access to a chow
diet and water. Starting at 10 weeks, mice were monitored for tumor development on a biweekly basis for
~10 additional weeks. Mice with tumors exceeding 17 mm in length/width or a volume over 5000 m^3 were
euthanized before the 20-week time point.

Tumor and Benign Mammary Tissue Collection

Mice were anesthetized using sodium pentobarbital (60 mg/kg body weight) and checked for depth of
anaesthesia by leg retraction after tail pinch. Once under the surgical plane, a ventral incision was made
up the midline and the skin separated bilaterally from the underlying fascia to allow for the excision of the
intact mammary tumors. Depending on the number of tumors present (typically 1-5), tumors were
allocated for mitochondrial function (placed fresh in BIOPS preservation buffer), histology (TEM: fixed in
2.5% glutaraldehyde/1% paraformaldehyde, IF: fixed in 10% formalin) or snap frozen in liquid nitrogen for
further analysis (western blotting, transcriptomics, quantitative label-free proteomics) immediately after
whole tumor wet weight was recorded. Paired mammary tissue with no visible tumors, furthest from the
site of tumor excision (i.e. if tumors localized in cervical region benign tissue was harvested from inguinal
region) were used as an internal comparison, denoted as benign mammary tissue. While mitochondrial
respiration experiments were performed in paired tissue (benign and tumor from same mouse),
transcriptomic, proteomic and histological analyses were performed on tumors and benign samples from
different mice due to tissue limitations.

RNA Sequencing and Transcriptomic Analyses

Frozen tissues were homogenized and lysed in TRIzol reagent. After centrifuging, the supernatant
(aqueous phase) was applied to the RNeasy kit (Qiagen, 74106) for subsequent total RNA extraction and
purification in accordance with the manufacturer's protocol. RNA samples were sent to the McMaster
Genomics Facility where an RNA quality check, poly A enrichment, library prep and quality check were
performed as previously described[62]. These samples were then run in an Illumina Nextseq P2, 2x50bp
sequencing run. FastQC and MultiQC were used for quality control of raw data. Salmon's transcript-level
quantification[63] and DESeq2[64] was used to detect DEGs with the threshold of adjusted p-value<0.1. PCA

analysis was performed by using variance stabilizing transformation (VST) data through DESeq2 using the
pcaExplorer package[65] in R (version 4.2.1). Functional enrichment analysis was performed by GO
enrichment analysis[66] and Kyoto Encyclopedia of Genes and Genomes (KEGG) mapping using the GOstats,
KEGG.db, and Bioconductor packages. Heatmaps and volcano plots were generated with ggplot2 package
using all detected genes or genes sorted by an adjusted p-value<0.1 by false discovery rate method.
Detailed descriptions for each analysis are included in the figure descriptions.

Western Blotting

Tumor and benign mammary tissues were homogenized in lysis buffer, diluted to equal protein
concentration (1 μg/uL), separated by SDS-PAGE, transferred onto PVDF membranes and detected using
enhanced chemiluminescence (ChemiGenius2 Bioimaging System, SynGene, Cambridge, UK) as previously
described[67,68]. Commercially available primary antibodies were used to detect total and phosphorylated
targets (p38 MAPK (CS9212S, 1:1000), ERK1/2 (CS9102S, 1:1000), Akt (CS4691S, 1:1000), mTOR (CS2972S,
1:1000), p70s6k1 (CS9202S, 1:1000), eEf2 (CS2332S, 1:1000), p-p38 MAPK (CS9211S, 1:1000), p-ERK1/2
(CS9101S, 1:1000), p-AKT (CS9271S, 1:1000), p-mTOR (CS2971S, 1:1000), p-p70s6k1 (CS9234S, 1:1000), p-
eEF2 (CS2331S, 1:1000), OXPHOS complexes CII and CIV (Ab110413, 1:1000)). Due to rapid tissue/protein
turnover in tumor tissues, Ponceau staining was an inappropriate loading control, however, trends in total
protein targets between groups aligned with quantitative label-free proteomic analyses as a secondary
method measuring protein content, validating targets detected by western blotting.

Protein Extraction and Purification for LC-MS

A Bradford assay was used to determine the protein concentration in the samples. Twenty-five μg of
protein per lysate was resolubilized with denaturation buffer (6M urea/2M thiourea) and concurrently
reduced with 10 mM DTT and alkylated with 20 mM iodoacetamide at RT for 60 min. Samples were
precipitated by adding 6:1 v/v cold acetone and kept at -80°C for 60 minutes and the pellet was collected
after centrifugation at 10,000g for 10 min at 4°C. Samples were resuspended with 50 mM ammonium
bicarbonate and MD-grade trypsin (Thermo Fisher Scientific, Cat#90057) was added at a ratio of 1:50
protease to protein. Digestion occurred overnight at 37°C. After digestion, samples were dried by vacuum
centrifugation and purified using Pierce C18 Spin columns (Thermo Fisher Scientific, cat#89873).

LC-MS

The Vanquish Neo UHPLC system was coupled with Orbitrap Exploris 240 mass-spectrometer using the
Easy-Spray source for nanoLC-MS protein identification. The Vanquish Neo UHPLC system was configured
for trap and elute analysis. Peptides were first trapped and washed on a Pepmap Neo C18 trap column (5
micron, 300 micron x5 mm) then separated on EASY-Spray columns 75 μm I.D. × 50 cm with the maximum
pressure of 1200 bar. The nanoLC-MS system was controlled with Standard Instrument Integration (SII)
for Xcalibur software. All hardware and data acquisition software were from Thermo Fisher Scientific. The
mobile phase A and weak wash liquid was water with 0.1% FA, and the mobile phase B and strong wash
liquid was 80% acetonitrile with 0.1% FA. The gradient was as follows 4-19% B over 72 min, 19-29% B over
28 min, 29-45% B over 20 min and a 14.5 min wash at 100% B with a flow rate of 300 nL/min. The
autosampler temperature was 7 °C, and the column temperature was 45 °C. The sample was injected with
Fast Loading set to 'Enabled' with Pressure Control at 500 bar. The column Fast Equilibration function was
set to 'Enabled' with Pressure control at 800 bar, and the equilibration factor was set to 3. Vial bottom
detection was set to 'Enabled'.

DDA MS Method

The Orbitrap Exploris 240 MS was operated in DDA mode using a full scan with m/z range 375–1500,
Orbitrap resolution of 60 000, normalized AGC target value 300%, and maximum injection time set to
Auto. The intensity threshold for precursor was set to 1×10^4. MS/MS spectra starting from 120 m/z were

acquired in data-dependent acquisition (DDA) mode with a cycle time of 2 s, where the precursors were
isolated in a window of 1.6 Da and subsequently fragmented with HCD using a normalized collision energy
of 30%. Orbitrap resolution was set to 15 000 for MS2. The normalized AGC target was standard, and the
maximum injection time was set to Auto.

Proteomics data analysis

Analysis of the tissue proteome was performed in R (version 4.2.1) and R Studio (version 2024.04.2+764).
Principal component analysis was performed using the Vegan package. Volcano plots were created using
log2 transformed fold change comparing cancer/benign tissue and -log10 adjusted p-value. P-values of
differentially regulated proteins between each group were corrected with p.adjust (method = "fdr")
within each comparison. Depending on the analysis, heatmaps and volcano plots were generated using
all detected proteins or proteins sorted by p.adjust <0.1, detailed descriptions for each analysis are
included in the figure descriptions. Gene set enrichment was performed with the Kyoto Encyclopedia of
genes and genomes (KEGG)[69] and EnrichR[70].

Cell Culture

NF639 murine epithelial mammary tumor cells (ATCC: CRL-3090) were cultured in DMEM supplemented
with 10% fetal bovine serum (FBS; Gibco), 4 mM L-glutamine, 1 mM sodium pyruvate, 50 U/mL penicillin
and 50 ug/mL streptomycin (Wisent), maintained by passaging in 100 mm plates, and incubated at 37°C
and 5% CO_2. Cells were seeded in 96-well plates (10^4 cells/well; n=3 technical replicates) for microplate
viability assays or in 100 mm plates for mitochondrial respiration experiments. At 60% confluency, cells
were serum-starved for 6 hours, followed by lapatinib treatment (Cayman, 11493) in complete culture
medium containing 2% FBS for 24 hours. Cell viability in the presence of lapatinib was tested using a crystal
violet assay. Cells were treated with lapatinib at 0 nM, 100 nM, 250 nM, 500 nM, 1000 nM, or 5000 nM
for 24 hours. Following treatment incubation, cells were washed and stained with crystal violet dye
(Abcam) and absorbance was measured at 570 nm as per the manufacturer's instructions. Assays were
repeated in triplicate. For mitochondrial respiration experiments, cells were treated with 250 nM lapatinib
or DMSO (control) in complete culture medium containing 2% FBS for 24 hours. Cells were washed,
trypsinized (1x trypsin-EDTA, Wisent), and centrifuged at 300g for 5 min before counting. Cells were
washed and resuspended in 1x PBS and centrifuged at 300g for 5 minutes. Cells were resuspended in
Mir05 respiration buffer (0.5 mM EGTA, 3 mM $MgCl_2 \cdot H_2O$, 60 mM potassium lactobionate, 10 mM KH_2PO_4,
20 mM HEPES, 110 mM sucrose, 20 mM taurine, and 1 g/L fatty acid free BSA; pH 7.1) for mitochondrial
respiration.

Mitochondrial Respiration

Mitochondrial respiration experiments were performed using the Oroboros Oxygraph-2k systems at 37ºC
in MiR05 respiration with constant stirring at room air saturation (starting at 180 µM O_2). For respiration
experiments in intact tumors and benign tissue, 2-5 mg of permeabilized tumor tissue or 15-20 mg of
permeabilized (both permeabilized with 20 µg/µL saponin) mammary tissue were used per 2 mL chamber
and each protocol was conducted as at least two technical replicates (duplicate) per tumor or benign
sample. Prior to the addition of substrates for any protocol, endogenous (basal) respiration was assessed
with no substrates added. Complex I and Complex II supported respiration was assessed with the addition
of 5 mM pyruvate and 2 mM malate, 5 mM ADP, 10 mM glutamate and 10 mM succinate. This respiratory
capacity measurement represents supraphysiological substrate concentrations delivered in the absence of
diffusion limitations. In a subset of these experiments, subsequent additions of oligomycin (OMY, 1 µM),
antimycin A (AMA, 2.5 µM) and rotenone (ROT, 1 µM) were added to assess OXPHOS coupling. In a
separate protocol, lipid-supported respiration was assessed with 20 µM palmitoyl-carnitine in the
presence of 2 mM malate and 5 mM ADP. For respiration experiments in NF639 cells, the same respiration
buffers and chamber conditions were utilized. One million cells were added per technical replicate and
Complex I and II supported respiration was assessed using the same protocol described in intact tumors

and benign tissue in the presence of 20 µg digitonin to ensure permeabilization. Respiration was expressed
per viable cell.

Reactive Oxygen Species Emission

Mitochondrial hydrogen peroxide (H_2O_2) emission was determined fluorometrically (Lumina, Thermo
Scientific, Waltham, MA) in permeabilized mammary tissue and mammary tumors as previously
described[67] with minor modifications. In brief, 2-5 mg of minced tissue was loaded into a cuvette
containing Amplex Red (Invitrogen, Waltham, MA, USA), horseradish peroxidase (5U/mL), saponin (20
µg/µL), superoxide dismutase (40 U/mL) in Buffer Z (105 mM K-MES, 30 mM KCl, 1 mM EGTA, 10 mM
K_2HPO_4, 5 mM $MgCl_2$, 5 µM glutamate, 5 µM malate, 0.5% BSA, pH 7.4). Emission rates were assessed at
37ºC in the presence of 10 mM succinate and calculated using a standard curve generated with known
concentrations of H_2O_2.

Data Normalization

Mitochondrial respiration rates and H_2O_2 emission were normalized to individual sample wet weight,
group total protein content and group mitochondrial content (mitochondrial enrichment factor, MEF).
Total protein content was calculated as a group average (n=6/group) based on a Bradford assay and back-
calculated total protein in a given wet weight of starting tissue. Data was normalized to mitochondrial
protein by calculating the MEF for each group (n=g/group) based on all detected mitochondrial proteins
by quantitative label-free proteomics (Mouse MitoCarta 3.0 database) normalized to total protein
abundance per sample as previously described by others[31,42].

Histology

Transmission Electron Microscopy: Immediately following tissue dissection, tumor and benign mammary
tissues were fixed in 2.5% glutaraldehyde/1% paraformaldehyde overnight at 4ºC. The following day,
tissues were washed 3x in 100 mM HEPES buffer in preparation for embedding. Tissue was stained with
1% osmium tetroxide for 3h, washed 3x in HEPES buffer and further stained with 1% uranyl acetate for 3h
before washing 2x in HEPES buffer and once in ddH2O. Stained tissue was dehydrated with a graded
ethanol series followed by infiltration with LR White resin. Blocks were polymerized at 60ºC for 18 hours
before ultramicrotomy. Sections were post-stained with uranyl acetate and lead citrate. Sections were
allowed to fully dry before imaging on the FEI Tecnai G2 F20 transmission electron microscope operated
under normal conditions at 120 kVe.

Immunofluorescence: Tumor and benign samples (n=4/group) were fixed in 10% formalin for 24 hours
and stored in 70% ethanol at 4ºC. Samples were blocked in paraffin and sectioned, deparaffinized in
xylene, and rehydrated through a graded ethanol series. Samples were washed in PBS and incubated for
20 min in 0.02% sodium borohydride in PBS to quench endogenous phosphatase activity. Antigen retrieval
was performed in citrate buffer (90ºC) for 12 min, followed by a 20-min cooling period. Sections were then
blocked in 5% BSA in PBS for 1h to prevent nonspecific antibody binding. Primary antibodies (anti-vimentin
(1:100, Novus Bio NBP1-31327), anti-cytokeratin (1:100, Novus Bio NBP2-29429), anti-CD4 (1:200, Abcam
AB183685), anti-CD138 (1:100, StemCell Technologies 60035), anti-CD31 (1:150, Abcam AB28364), or anti-
CD68 (1:100, Novus Bio NB100-683)) were diluted in 1% BSA and applied overnight at 4 °C in a humidified
chamber. The following day, sections were incubated for 2 hours at room temperature with the
appropriate fluorophore-conjugated secondary antibodies (Invitrogen): Alexa Fluor 594 anti-rabbit (1:100
or 1:200), Alexa Fluor 488 anti-mouse (1:100), Alexa Fluor 488 anti-rat (1:200), or Alexa Fluor 488 anti-
rabbit (1:100). Slides were washed and mounted with ProLong™ Gold Antifade Mountant containing DAPI
(Thermo Fisher) and protected from light for 24 hours before being transferred to -20°C for storage. Images
were acquired at 40× magnification using an Olympus BX-61 epifluorescence microscope equipped with
Metamorph imaging software. Monochromatic images captured at 594 nm excitation were pseudocolored
red, while those captured at 488 nm were pseudocolored green. Staining intensity was quantified in

ImageJ using manual region-of-interest selection, and immunopositive area was expressed as a percentage
of total tissue area (expressed as % immunopositive area).

OXPHOS Index and OXPHOS Score Calculations

A publicly available single-cell RNA-seq dataset from of 26 human breast tumors[31] was combined with cell-
type proportion analyses from our tumor and benign tissues (see Immunofluorescence methods) to
estimate cell-type-specific contribution to OXPHOS.

Using human MitoCarta 3.0, we filtered for all detected OXPHOS subunit genes in the scRNA-seq dataset.
For each major cell type described in the dataset (B-, T-, plasmablasts, CAFs, PVL, endothelial, myeloid,
normal epithelial, and cancer epithelial cells), we extracted the percent of cells expressing each subunit
gene. These percent expression values were averaged per complex to generate a mean CI-V expression
value for each cell type. To account for differences in cell-type abundance within the scRNA-seq dataset,
this percent expression was multiplied by the cell-type proportion reported in the same study. In
subsequent calculations, this is reported as $\%Expression_{CT}$.

To estimate cell-type-specific OXPHOS expression in our benign and tumor samples, we calculated an
OXPHOS mRNA Index for all cell types present in both datasets (myeloid, T-, plasmablasts, endothelial,
mesenchymal, and epithelial cells):

$$OXPHOS\ mRNA\ Index = (\%Expression_{CT}) * (\%Immunopositive_{CT})$$

Where, $\%Expression_{CT}$ is the average gene expression data per respiratory complex in each complementary
major cell type (CT), and $\%Immunopositive_{CT}$ is cell type proportion measured by IF in our samples. This
index represents how much each cell type is expected to contribute to CI-V expression in our tissues,
integrating both expression and abundance.

Next, to estimate the contribution of each cell type to the OXPHOS transcriptome in the current study, the
OXPHOS mRNA index in each cell type were summed to calculate a Total OXPHOS mRNA Score.

$$Total\ OXPHOS\ mRNA\ Score = \sum_{i=I,II,III,IV,V} OXPHOS\ RNA\ Index\ Ci$$

Statistical Analysis

Statistical analysis and data visualization was performed using Microsoft Excel, GraphPad Prism (version
10.2.3), R (version 4.2.1) and R Studio (version 2024.04.2+764). All data are presented as mean ± SD and
data was considered statistically significant if p<0.05. For transcriptomic and proteomic analyses, an
adjusted p-value of <0.1 was utilized. Details regarding specific statistical analyses are included in the
Results section and within each figure legend.

DATA AVAILABILITY

Raw RNA sequencing and proteomics data were deposited in GEO (accession number: GSE315552) and
PRIDE databases (accession number: PXD071897) respectively.

ACKNOWLEDGEMENTS

This work is funded by a National Sciences and Engineering Research Council (NSERC) grant by GPH
(400362). SMF is funded by an NSERC-CGS doctoral scholarship.

FIGURE LEGENDS

Figure 1. Experimental design and canonical HER2 signaling. A) Schematic of mammary tumors from the mammary fat pad of
female MMTV-neu[ndl]-YD5 mice and normal (benign) mammary tissue, representative images of tumor morphology (tumor
capsule, peripheral tumor tissue and necrotic core), tissue processing workflow for transcriptomics, proteomics, western blotting,
mitochondrial respiration, ROS emission, and histology (transmission electron microscopy (TEM) and immunofluorescence (IF)).
B) Schematic of classical signaling pathways (Ras/MAPK and PI3K/AKT/mTOR) downstream of HER2 receptor stimulation in the

YD5 model. **C)** *Her2* gene expression (transcripts per million, TPM), **D)** Gene expression for canonical HER2 signaling. **E)** Absolute abundance of HER2 protein detected by quantitative label-free proteomics (n.d. = HER2 not detected in benign). **F)** Representative western blots of proteins involved in canonical HER2 signaling. **G,H)** Quantified western blots (% of benign) for **G)** total protein content and **H)** phosphorylated targets normalized to the total protein content of each target. P-values in panels **D** and **E** listed as adjusted p values from transcriptomic (DEG) data (p-adj. <0.1 considered statistically significant). In panels **G,H** data is presented as means ± SD and statistical significance was determined using an unpaired, two tailed Student's t-test (*=p<0.05, n=5-6 per group/protein target). ***Abbrev.*** *AKT - protein kinase B; eEf2 - eukaryotic elongation factor 2; eIF4E – eukaryotic translation initiation factor 4E; ERK – extracellular signal-regulated kinase; GRB2 – growth factor receptor-bound protein-2, MEK – mitogen-activated protein kinase kinase; mTORC1/2 – mechanistic target of rapamycin complex 1 and 2; PDK – 3-phosphoinositide-dependent protein kinase-1; PI3K – phosphoinositide 3-kinase; PIP2/3 – phosphatidylinositol 4,5 bisphosphate, PIP3 – phosphatidylinositol 3,4,5-triphosphate; p70S6k1 – ribosomal protein S6 kinase beta-1; RAF – rapidly accelerated fibrosarcoma; RAS – rate sarcoma virus; SHC – Src homolog and collagen homolog; 4EBP1 – eukaryotic translation initiation factor 4E-binding protein 1.*

Figure 2. Assessment of HER2-driven mammary tumor and benign mammary tissue transcriptomes. A) PCA of benign mammary tissue and mammary tumors, **B)** Volcano plot showing differentially expressed genes (DEGs) identified between mammary tissue and mammary tumors. **C)** Heatmap of top 3000 DEGs, **D)** Top 10 upregulated pathways (Reactome 2022) and **E)** gene ontologies (GO Biological Process 2023) in tumors compared to benign mammary tissue, **F)** Top 10 downregulated pathways (Reactome 2022) and **G)** gene ontologies (GO Biological Process 2023) in tumors compared to benign mammary tissue. Panels **D,E** assess the top 3000 upregulated DEGs (log2fc>1), panels **F,G** assess the top 3000 downregulated DEGs (log2fc<1). An adjusted p-value of <0.1 was considered statistically significant across all analyses. The Enrichr open access analysis web tool was used to generate panels **D-G** which were sorted by p-value.

Figure 3. Assessment of HER2-driven mammary tumor and benign mammary tissue proteomes. A) Venn diagram showing all detected proteins. Of the 1848 overlapping targets, 420 were significantly different based on a p-adj. <0.1 and 411 of those proteins were downregulated in tumors. **B)** PCA of benign mammary samples and mammary tumors, **C)** Volcano plot showing proteins identified between mammary tumors and benign mammary tissue, those in blue (downregulated) or red (upregulated) were significantly different (n=420 proteins, p-adj. <0.1). **D)** Heatmap of 420 differentially expressed proteins in benign and tumor samples. **E)** Top 10 downregulated pathways (Reactome 2022) and **F)** gene ontologies (GO Biological Process, 2023), **G)** KEGG analysis of downregulated proteins. The Enrichr open access analysis web tool was used to generate panels **E,F** which were sorted by p-value.

Figure 4. Overlapping genes and proteins between HER2-driven mammary tumor transcriptome and proteome. Correlation of log2FC for overlapping targets between the HER2-driven tumor and benign mammary tissue transcriptomes and proteomes grouped by up/downregulation **A),** and **B)** protein localization (mitochondrial vs other). **C)** Venn diagram showing 144 overlapping downregulated targets between the assessed HER2-driven tumor and benign mammary tissue transcriptomes and proteomes. **D)** KEGG analysis for common downregulated targets. Top 10 downregulated **E)** pathways (Reactome 2022) and **F)** gene ontologies (GO Biological Process 2023). An adjusted p value of <0.1 was utilized for all analyses. The Enrichr open access analysis web tool was used to generate panels **E,F** which were sorted by p-value.

Figure 5. Assessment of the mitochondrial transcriptome and proteome. A) Venn diagram showing mitochondrial proteins detected in the current dataset when cross-referenced with the Mouse MitoCarta 3.0 database. **B)** Heatmap of 125 mitochondrial proteins identified in benign mammary tissue and HER2-driven tumors (n=5/group). **C)** Absolute abundance of all mitochondrial proteins detected in benign and tumor tissues. **D)** Gene expression (detected by transcriptomics; transcripts per million) of all detected OXPHOS subunits separated per respiratory complex, black and white bars represent subunits with no significant differences between groups. **E-J)** Heatmaps of **E)** Complex I, **F)** Complex II, **G)** Complex III, **H)** Complex IV, **I)** Complex V (ATP synthase) and **J)** TCA cycle transcriptomes. All proteomic analyses **(panels A-C)** were conducted on n=5/group. All transcriptomic analyses **(panels D-J)** were conducted on n=6/group. **Panels B,C** were analyzed using an unpaired two tailed Student's t-test (*p<0.05). In **Panel D**, * and coloured (blue and green bars) represent significant differences between benign and tumor samples based on an adjusted p value of <0.1. OXPHOS complex schematics in panel **D** were obtained from the open access RCSB Protein Data Bank *(Complex I = TW1V, Complex II=2A06, Complex III= 3SFD Complex IV=6NMP Complex V = 5FIL). Bars in panel **D** are listed in the following order: Complex I: Ndufa1, Ndufa10, Ndufa11, Ndufa12, Ndufa13, Ndufa2, Ndufa3, Ndufa5, Ndufa6, Ndufa7, Ndufa8, Ndufa9, Ndufab1, Ndufb10, Ndufb11, Ndufb2, Ndufb3, Ndufb4, Ndufb5, Ndufb6, Ndufb7, Ndufb8, Ndufb9, Ndufc1, Ndufc2, Ndufs1, Ndufs2, Ndufs3, Ndufs4, Ndufs5, Ndufs6, Ndufs7, Ndufs8, Ndufv1, Ndufv2, Ndufv3; Complex II: Sdha, Sdhb, Sdhc, Sdhd; Complex III: Cyc1, Uqcr10, Uqcr11, Uqcrb, Uqcrc1, Uqcrc2, Uqcrfs1, Uqcrh, Uqcrq; Complex IV: Cox4i1, Cox4i2, Cox5a, Cox5b, Cox6a1, Cox6b1, Cox6b2, Cox6c, Cox7a1, Cox7a2, Cox7a2l, Cox7b, Cox7c, Cox8a, Cox8b, Ndufa4; Complex V: Atp5a1, Atp5b, Atp5c1, Atp5d, Atp5e, Atp5g1, Atp5g2, Atp5g3, Atp5h, Atp5j, Atp5j2, Atp5k, Atp5l, Atp5md, Atp5mpl, Atp5o, Atp5pb, Atpif1, Dmac2l.*

Figure 6. Cell type specific OXPHOS mRNA analysis in benign and tumor samples. A) OXPHOS expression per cell type from a publicly available single-cell RNA sequencing dataset[31] shown as a dot plot for each respiratory complex (Complexes I-V) subunit that was detected. Dot size indicates the percentage of cells of a given major type expressing each subunit gene (% cells expressing). Dot color represents gene expression scaled relative to each gene's expression distribution across all major cell types. **B)** Percentage of cells expressing OXPHOS subunits, arranged in ascending order per cell type for Complexes I-V. **C)** Representative immunofluorescence images of benign mammary tissue and tumor samples stained for vimentin, cytokeratin, CD68, CD31, CD4 and CD138. **D)** Quantification of cell-type proportion (% immunopositive area) in benign and tumor samples. **E)** OXPHOS mRNA

Index calculated for Complexes I-V. $OXPHOS\ mRNA\ Index = (\%Expression_{CT}) * (\%Immunopositive_{CT})$ where CT represents cell type. **F)** Total OXPHOS mRNA Score for each cell type. $TOTAL\ OXPHOS\ mRNA\ Score = \sum_{i=I,II,III,IV,V} OXPHOS\ mRNA\ Index\ Ci$. Where i represents mitochondrial respiratory complexes.

Figure 7. Assessment of lipid and carbohydrate linked mitochondrial respiration and OXPHOS coupling in benign mammary tissue and HER2-driven mammary tumors. A) Schematic depicting methods of normalization for respiration rates (*per wet weight, total protein, mitochondrial protein/enrichment*). **B)** Complex I and complex II- supported respiration normalized to tissue wet weight, **C)** lipid-supported respiration normalized to tissue wet weight, **D)** Complex I and complex II-supported respiration normalized to total protein content, **E)** lipid-supported respiration normalized to total protein content, **F)** Complex I and complex II-supported respiration normalized to mitochondrial protein (mitochondrial enrichment factor, MEF). **G)** Lipid supported respiration normalized to mitochondrial protein. All individual values represent biological replicates (n=9-22/group depending on the protocol). Data are presented as means ± SD, statistical analyses were conducted using an unpaired, two-tailed Student's t-test (*p<0.05). ***Abbrev:*** *E = Endogenous, PM = pyruvate (5 mM), malate (2 mM), D = ADP (5 mM), G = glutamate (10 mM), S = succinate (10 mM), PC = palmitoyl-carnitine (20 µM), CI+CII = maximal CI+CII-supported respiration (PMDGS).*

Figure 8. Lipid and carbohydrate linked respiration and lipid metabolism transcriptome and proteome. A) Rates of carbohydrate (pyruvate, malate, ADP) and lipid (palmitoyl-carnitine, malate, ADP) supported respiration in tumors and benign mammary tissue. **B)** Gene expression (transcripts per million) of mitochondrial pyruvate carrier and carnitine palmitoyltransferase isoforms mediating mitochondrial pyruvate and long chain fatty acid transport respectively. **C)** Network interaction showing downregulated (unsupervised analysis) gene clustering in the HER2-driven tumor transcriptome, **D)** Heatmap of differentially expressed genes involved in lipid oxidation. **E)** Heatmap of differentially expressed proteins involved in lipid metabolism. **F)** Representative trace of protocol used to assess mitochondrial coupling. Following maximal rates of complex I and II- supported respiration, 1 µM oligomycin (OMY) was added to inhibit Complex V (ATP synthase), followed by 2.5 µM antimycin-A (AMA) and 1 µM rotenone (ROT) to inhibit complexes III and I respectively. **G)** Group data for experiment depicted in **(F)** depicted as relative inhibition from maximal complex I and II supported respiration (PMDGS) in each group. In panels **A, B, F** and **G** data are presented as means ± SD and statistical analyses were conducted using an unpaired, two-tailed Student's t-test (*p<0.05).

Figure 9. Mitochondrial morphology and assessment of mitochondrial fusion, mitochondrial fission and mitophagy transcriptomes and proteomes in benign mammary tissue and HER2-driven tumors. A) Transmission electron microscopy (TEM) images of tumor and benign samples showing punctate (tumor) versus elongated (benign) mitochondria. **B)** Gene expression (detected by transcriptomics; transcripts per million) of fusion, fission and mitophagy related genes separated per process filtered by MitoCarta, coloured bars and * represent genes with an adjusted p value of <0.1. Black and white bars represent genes with no significant differences between groups. **C)** Heatmap of fusion transcriptome, **D)** Heatmap of fusion proteome, **E)** Heatmap of fission transcriptome, **F)** Heatmap of fission proteome, **G)** Heatmap of mitophagy transcriptome, **H)** Heatmap of mitophagy proteome. TEM images are from independent tumor and benign tissue samples. Heatmaps depict all detected proteins (n=5/group) and transcripts (n=6/group).

Figure 10. Mitochondrial respiration is supressed by tyrosine kinase inhibitor lapatinib. A) Schematic showing mechanism of lapatinib mediated inhibition of HER2 tyrosine kinase activity. **B)** NF639 cell viability assessed using a crystal violet assay expressed as a percentage of untreated viability (0nM lapatinib). **C)** Complex I and II supported respiration in the presence of 20µg digitonin in NF639 cells untreated (grey) or treated with 250 nM lapatinib for 24h. Left inset: Schematic of substrates supporting Complex I-II respiration, schematic of lapatinib or DMSO treatment. Right inset: Percent inhibition of maximal CI+II supported respiration (PMDGSC) with OMY, AMA and ROT. Following maximal rates of complex I and II- supported respiration, 1 µM oligomycin (OMY) was added to inhibit Complex V (ATP synthase), followed by 2.5 µM antimycin-A (AMA) and 1 µM rotenone (ROT) to inhibit complexes III and I respectively. Each datapoint represents a technical replicate (duplicate) and data are presented as means ± SD. Statistical analyses were conducted using an unpaired, two-tailed Student's t-test (*p<0.05). ***Abbrev:*** *E = Endogenous, PMD = pyruvate (5 mM), malate (2 mM), D= ADP (5 mM), G = glutamate (10 mM), S = succinate (10 mM), CC = cytochrome C (10 µM).*

Supplementary Figure 1. Absolute phosphorylated protein targets downstream of HER2 activation. A) Quantified western blots for phosphorylated canonical HER2 signaling proteins (n=5-6/group). Phosphorylation sites are indicated for each target. **B)** Representative blot for each phosphorylated target. Data are presented as mean ± SD and analyzed with an unpaired, two tailed Student's t-test (*p<0.05).

Supplementary Figure 2. Unique HER2-driven tumor and benign mammary tissue proteins. A) Venn diagram depicting 741 and 600 unique proteins identified in benign mammary tissue and HER2-driven tumors respectively, **B)** KEGG analysis of all detected unique benign mammary tissue proteins, **C)** Top 10 pathways (Reactome 2022), **D)** Top 10 gene ontologies (GO Biological Process) in benign mammary tissue, **E)** KEGG analysis of unique HER2-driven tumor proteins, **F)** Top 10 pathways (Reactome 2022), **G)** Top 10 gene ontologies (GO Biological Process) in HER2-driven tumors. KEGG, pathway and GO analyses were conducted on all mammary and tumor proteins detected by quantitative label-free proteomics. The Enrichr open access analysis web tool was used to generate panels **C,D,F and G** which were sorted by p-value.

Supplementary Figure 3. Mitochondrial enrichment in benign mammary tissue and HER2-driven mammary tumors. A) Total and mitochondrial (Mouse MitoCarta 3.0) absolute protein abundance assessed by quantitative label-free proteomics. **B)** Mitochondrial enrichment factor (MEF) calculated as the ratio of all detected MitoCarta proteins relative to the underlying total

proteome per sample (n=5/group). Statistical analysis was conducted using an unpaired two tailed Student's t-test (*p<0.05). In panel **A**, *# denotes significant differences between both total and MitoCarta proteomes between groups.

Supplementary Figure 4. Proteomic analysis of detected OXPHOS subunits. Heatmaps showing subunits detected for **A)** Complex I, **B)** Complex II, **C)** Complex III, **D)** Complex IV, **E)** Complex V (ATP synthase) using quantified label-free proteomics. **F)** OXPHOS Complex I-V absolute abundance as a percentage of the mitochondrial proteome (n=5/group). **G)** OXPHOS complex I-V absolute abundance as a percentage of the total detected proteome (n=5/group). **H)** Representative and quantified western blots for Complex II and Complex IV expressed as percentage of benign. Data in panels **F,G,H** are presented as means ± SD and analyzed with an unpaired, two tailed Student's t-test (*p<0.05).

Supplementary Figure 5. Correlation of tumor size and volume with maximal carbohydrate and lipid-supported respiration rates. A) Tumor wet weight (whole tumor post excision), **B)** Tumor volume calculated as a modified ellipse (V =1/2(length*width^2)). **C-F)** Endogenous (no substrate, CI+CII protocol), Endogenous (no substrate, lipid protocol), State 3 (5 mM pyruvate, 2 mM malate, 5 mM ADP) and maximal lipid-supported (2 mM malate, 5 mM ADP, 20 μM palmitoyl-carnitine) respiration rates correlated with tissue wet weight. **G-J)** Same respiration data as **(C-F)** but correlated with tumor volume. The Pearson correlation coefficient (R) values are listed for each correlation.

Supplementary Figure 6. Reactive oxygen species emission in benign mammary tissue and HER2-driven mammary tumors. A) Schematic depicting major sites of superoxide (O_2^{*-}) emission from respiratory complexes I and III. Superoxide dismutase 2 (SOD2) subsequently catalyzes the dismutation of O_2^{*-} into hydrogen peroxide (H_2O_2) which can be measured fluorometrically. **B)** H_2O_2 emission per wet tissue weight, **C)** H_2O_2 emission per total protein, **D)** H_2O_2 emission per mitochondrial protein (MEF, mitochondrial enrichment factor). Statistical analyses were conducted using an unpaired, two-tailed Student's t-test (*p<0.05).

Supplementary Table 1. List of 9 significantly upregulated proteins between tumor and benign samples, sorted in order of fold change from benign (smallest to largest fold change). General pathways or functions are listed per protein based on ID mapping from the Uniprot protein sequence database. Adjusted p-values are listed per target.

REFERENCES

1. Vasan, K., Werner, M., and Chandel, N.S. (2020). Mitochondrial Metabolism as a Target for Cancer Therapy. Cell Metab. *32*, 341–352. https://doi.org/10.1016/j.cmet.2020.06.019.

2. DeBerardinis, R.J., and Chandel, N.S. (2020). We need to talk about the Warburg effect. Nat. Metab. *2*, 127–129. https://doi.org/10.1038/s42255-020-0172-2.

3. Warburg, O. (1924). Über den Stoffwechsel der Carcinomzelle. Naturwissenschaften *12*, 1131–1137. https://doi.org/10.1007/BF01504608.

4. Warburg, O. (1956). On respiratory impairment in cancer cells. Science *124*, 269–270.

5. Sullivan, L.B., Gui, D.Y., Hosios, A.M., Bush, L.N., Freinkman, E., and Vander Heiden, M.G. (2015). Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferating Cells. Cell *162*, 552–563. https://doi.org/10.1016/j.cell.2015.07.017.

6. Martínez-Reyes, I., Cardona, L.R., Kong, H., Vasan, K., McElroy, G.S., Werner, M., Kihshen, H., Reczek, C.R., Weinberg, S.E., Gao, P., et al. (2020). Mitochondrial ubiquinol oxidation is necessary for tumour growth. Nature *585*, 288–292. https://doi.org/10.1038/s41586-020-2475-6.

7. Morris, J.P., Yashinskie, J.J., Koche, R., Chandwani, R., Tian, S., Chen, C.-C., Baslan, T., Marinkovic, Z.S., Sánchez-Rivera, F.J., Leach, S.D., et al. (2019). α-Ketoglutarate links p53 to cell fate during tumour suppression. Nature *573*, 595–599. https://doi.org/10.1038/s41586-019-1577-5.

8. Sciacovelli, M., Guzzo, G., Morello, V., Frezza, C., Zheng, L., Nannini, N., Calabrese, F., Laudiero, G., Esposito, F., Landriscina, M., et al. (2013). The mitochondrial chaperone TRAP1

promotes neoplastic growth by inhibiting succinate dehydrogenase. Cell Metab. *17*, 988–
999. https://doi.org/10.1016/j.cmet.2013.04.019.

9.   Cardaci, S., Zheng, L., MacKay, G., van den Broek, N.J.F., MacKenzie, E.D., Nixon, C.,
Stevenson, D., Tumanov, S., Bulusu, V., Kamphorst, J.J., et al. (2015). Pyruvate carboxylation
enables growth of SDH-deficient cells by supporting aspartate biosynthesis. Nat. Cell Biol.
17, 1317–1326. https://doi.org/10.1038/ncb3233.

10. Smith, A.L., Whitehall, J.C., Bradshaw, C., Gay, D., Robertson, F., Blain, A.P., Hudson, G., Pyle,
A., Houghton, D., Hunt, M., et al. (2020). Age-associated mitochondrial DNA mutations
cause metabolic remodelling that contributes to accelerated intestinal tumorigenesis. Nat.
Cancer *1*, 976–989. https://doi.org/10.1038/s43018-020-00112-5.

11. Tan, A.S., Baty, J.W., Dong, L.-F., Bezawork-Geleta, A., Endaya, B., Goodwin, J., Bajzikova, M.,
Kovarova, J., Peterka, M., Yan, B., et al. (2015). Mitochondrial genome acquisition restores
respiratory function and tumorigenic potential of cancer cells without mitochondrial DNA.
Cell Metab. *21*, 81–94. https://doi.org/10.1016/j.cmet.2014.12.003.

12. Weinberg, F., Hamanaka, R., Wheaton, W.W., Weinberg, S., Joseph, J., Lopez, M.,
Kalyanaraman, B., Mutlu, G.M., Budinger, G.R.S., and Chandel, N.S. (2010). Mitochondrial
metabolism and ROS generation are essential for Kras-mediated tumorigenicity. Proc. Natl.
Acad. Sci. U. S. A. *107*, 8788–8793. https://doi.org/10.1073/pnas.1003428107.

13. Yuan, Y., Ju, Y.S., Kim, Y., Li, J., Wang, Y., Yoon, C.J., Yang, Y., Martincorena, I., Creighton, C.J.,
Weinstein, J.N., et al. (2020). Comprehensive molecular characterization of mitochondrial
genomes in human cancers. Nat. Genet. *52*, 342–352. https://doi.org/10.1038/s41588-019-
0557-x.

14. Wang, Z., Zheng, Z., Jia, S., Liu, S., Xiao, X., Chen, G., Liang, W., and Lu, X. (2022).
Trastuzumab resistance in HER2-positive breast cancer: Mechanisms, emerging biomarkers
and targeting agents. Front. Oncol. *12*.

15. Avagliano, A., Ruocco, M.R., Aliotta, F., Belviso, I., Accurso, A., Masone, S., Montagnani, S.,
and Arcucci, A. (2019). Mitochondrial Flexibility of Breast Cancers: A Growth Advantage and
a Therapeutic Opportunity. Cells *8*, 401. https://doi.org/10.3390/cells8050401.

16. Baek, M.L., Lee, J., Pendleton, K.E., Berner, M.J., Goff, E.B., Tan, L., Martinez, S.A., Mahmud,
I., Wang, T., Meyer, M.D., et al. (2023). Mitochondrial structure and function adaptation in
residual triple negative breast cancer cells surviving chemotherapy treatment. Oncogene *42*,
1117–1131. https://doi.org/10.1038/s41388-023-02596-8.

17. Evans, K.W., Yuca, E., Scott, S.S., Zhao, M., Paez Arango, N., Cruz Pico, C.X., Saridogan, T.,
Shariati, M., Class, C.A., Bristow, C.A., et al. (2021). Oxidative Phosphorylation Is a Metabolic
Vulnerability in Chemotherapy-Resistant Triple-Negative Breast Cancer. Cancer Res. *81*,
5572–5581. https://doi.org/10.1158/0008-5472.CAN-20-3242.

18. El-Botty, R., Morriset, L., Montaudon, E., Tariq, Z., Schnitzler, A., Bacci, M., Lorito, N., Sourd,
L., Huguet, L., Dahmani, A., et al. (2023). Oxidative phosphorylation is a metabolic
vulnerability of endocrine therapy and palbociclib resistant metastatic breast cancers. Nat.
Commun. *14*, 4221. https://doi.org/10.1038/s41467-023-40022-5.

19. Rohlenova, K., Sachaphibulkij, K., Stursa, J., Bezawork-Geleta, A., Blecha, J., Endaya, B.,
Werner, L., Cerny, J., Zobalova, R., Goodwin, J., et al. (2017). Selective Disruption of
Respiratory Supercomplexes as a New Strategy to Suppress Her2high Breast Cancer.
Antioxid. Redox Signal. *26*, 84–103. https://doi.org/10.1089/ars.2016.6677.

20. Yap, T.A., Daver, N., Mahendra, M., Zhang, J., Kamiya-Matsuoka, C., Meric-Bernstam, F.,
Kantarjian, H.M., Ravandi, F., Collins, M.E., Francesco, M.E.D., et al. (2023). Complex I
inhibitor of oxidative phosphorylation in advanced solid tumors and acute myeloid
leukemia: phase I trials. Nat. Med. *29*, 115–126. https://doi.org/10.1038/s41591-022-
02103-8.

21. Mun, S., Lee, H.J., and Kim, P. (2024). Rebuilding the microenvironment of primary tumors in
humans: a focus on stroma. Exp. Mol. Med. *56*, 527–548. https://doi.org/10.1038/s12276-
024-01191-5.

22. de Visser, K.E., and Joyce, J.A. (2023). The evolving tumor microenvironment: From cancer
initiation to metastatic outgrowth. Cancer Cell *41*, 374–403.
https://doi.org/10.1016/j.ccell.2023.02.016.

23. Ursini-Siegel, J., Schade, B., Cardiff, R.D., and Muller, W.J. (2007). Insights from transgenic
mouse models of ERBB2-induced breast cancer. Nat. Rev. Cancer *7*, 389–397.
https://doi.org/10.1038/nrc2127.

24. Iqbal, N., and Iqbal, N. (2014). Human Epidermal Growth Factor Receptor 2 (HER2) in
Cancers: Overexpression and Therapeutic Implications. Mol. Biol. Int. *2014*, 1–9.
https://doi.org/10.1155/2014/852748.

25. Dankort, D.L., Wang, Z., Blackmore, V., Moran, M.F., and Muller, W.J. (1997). Distinct
Tyrosine Autophosphorylation Sites Negatively and Positively Modulate Neu-Mediated
Transformation. Mol. Cell. Biol. *17*, 5410–5425. https://doi.org/10.1128/MCB.17.9.5410.

26. Dankort, D., Jeyabalan, N., Jones, N., Dumont, D.J., and Muller, W.J. (2001). Multiple ErbB-
2/Neu Phosphorylation Sites Mediate Transformation through Distinct Effector Proteins. J.
Biol. Chem. *276*, 38921–38928. https://doi.org/10.1074/jbc.M106239200.

27. Ruiz-Saenz, A., Dreyer, C., Campbell, M.R., Steri, V., Gulizia, N., and Moasser, M.M. (2018).
HER2 amplification in tumors activates PI3K/Akt signaling independent of HER3. Cancer Res.
78, 3645–3658. https://doi.org/10.1158/0008-5472.CAN-18-0430.

28. Kirouac, D.C., Du, J., Lahdenranta, J., Onsum, M.D., Nielsen, U.B., Schoeberl, B., and
McDonagh, C.F. (2016). HER2+ Cancer Cell Dependence on PI3K vs. MAPK Signaling Axes Is
Determined by Expression of EGFR, ERBB3 and CDKN1B. PLoS Comput. Biol. *12*, e1004827.
https://doi.org/10.1371/journal.pcbi.1004827.

29. Serra, V., Scaltriti, M., Prudkin, L., Eichhorn, P.J.A., Ibrahim, Y.H., Chandarlapaty, S.,
Markman, B., Rodriguez, O., Guzman, M., Rodriguez, S., et al. (2011). PI3K inhibition results
in enhanced HER signaling and acquired ERK dependency in HER2-overexpressing breast
cancer. Oncogene *30*, 2547–2557. https://doi.org/10.1038/onc.2010.626.

30. Boykov, I.N., Montgomery, M.M., Hagen, J.T., Aruleba, R.T., McLaughlin, K.L., Coalson, H.S.,
Nelson, M.A., Pereyra, A.S., Ellis, J.M., Zeczycki, T.N., et al. (2023). Pan-tissue mitochondrial
phenotyping reveals lower OXPHOS expression and function across cancer types. Sci. Rep.
13, 16742. https://doi.org/10.1038/s41598-023-43963-5.

31. Wu, S.Z., Al-Eryani, G., Roden, D.L., Junankar, S., Harvey, K., Andersson, A., Thennavan, A.,
Wang, C., Torpy, J.R., Bartonicek, N., et al. (2021). A single-cell and spatially resolved atlas of
human breast cancers. Nat. Genet. *53*, 1334–1347. https://doi.org/10.1038/s41588-021-
00911-1.

32. Amundadottir, L.T., and Leder, P. (1998). Signal transduction pathways activated and
required for mammary carcinogenesis in response to specific oncogenes. Oncogene *16*,
737–746. https://doi.org/10.1038/sj.onc.1201829.

33. Muller, W.J., Sinn, E., Pattengale, P.K., Wallace, R., and Leder, P. (1988). Single-step induction
of mammary adenocarcinoma in transgenic mice bearing the activated c-neu oncogene. Cell
54, 105–115. https://doi.org/10.1016/0092-8674(88)90184-5.

34. Min, C., Kirsch, K.H., Zhao, Y., Jeay, S., Palamakumbura, A.H., Trackman, P.C., and
Sonenshein, G.E. (2007). The Tumor Suppressor Activity of the Lysyl Oxidase Propeptide
Reverses the Invasive Phenotype of Her-2/neu–Driven Breast Cancer. Cancer Res. *67*, 1105–
1112. https://doi.org/10.1158/0008-5472.CAN-06-3867.

35. Romieu-Mourez, R., Landesman-Bollag, E., Seldin, D.C., and Sonenshein, G.E. (2002). Protein
kinase CK2 promotes aberrant activation of nuclear factor-kappaB, transformed phenotype,
and survival of breast cancer cells. Cancer Res. *62*, 6770–6778.

36. Xiang, Z., Song, S., Zhu, Z., Sun, W., Gifts, J.E., Sun, S., Li, Q.S., Yu, Y., and Li, K.K. (2019).
LncRNAs GIHCG and SPINT1-AS1 Are Crucial Factors for Pan-Cancer Cells Sensitivity to
Lapatinib. Front. Genet. *10*, 25. https://doi.org/10.3389/fgene.2019.00025.

37. Leung, W., Roxanis, I., Sheldon, H., Buffa, F.M., Li, J., Harris, A.L., and Kong, A. (2015).
Combining lapatinib and pertuzumab to overcome lapatinib resistance due to NRG1-
mediated signalling in HER2-amplified breast cancer. Oncotarget *6*, 5678–5694.
https://doi.org/10.18632/oncotarget.3296.

38. Putignani, L., Raffa, S., Pescosolido, R., Aimati, L., Signore, F., Torrisi, M.R., and Grammatico,
P. (2008). Alteration of expression levels of the oxidative phosphorylation system (OXPHOS)
in breast cancer cell mitochondria. Breast Cancer Res. Treat. *110*, 439–452.
https://doi.org/10.1007/s10549-007-9738-x.

39. Hu, Y., Xu, W., Zeng, H., He, Z., Lu, X., Zuo, D., Qin, G., and Chen, W. (2020). OXPHOS-
dependent metabolic reprogramming prompts metastatic potential of breast cancer cells
under osteogenic differentiation. Br. J. Cancer *123*, 1644–1655.
https://doi.org/10.1038/s41416-020-01040-y.

40. Pacheco-Velázquez, S.C., Robledo-Cadena, D.X., Hernández-Reséndiz, I., Gallardo-Pérez, J.C.,
Moreno-Sánchez, R., and Rodríguez-Enríquez, S. (2018). Energy Metabolism Drugs Block
Triple Negative Breast Metastatic Cancer Cell Phenotype. Mol. Pharm. *15*, 2151–2164.
https://doi.org/10.1021/acs.molpharmaceut.8b00015.

41. Nelson, M.A., McLaughlin, K.L., Hagen, J.T., Coalson, H.S., Schmidt, C., Kassai, M., Kew, K.A.,
McClung, J.M., Neufer, P.D., Brophy, P., et al. (2021). Intrinsic OXPHOS limitations underlie
cellular bioenergetics in leukemia. eLife *10*, e63104. https://doi.org/10.7554/eLife.63104.

42. McLaughlin, K.L., Hagen, J.T., Coalson, H.S., Nelson, M.A.M., Kew, K.A., Wooten, A.R., and
Fisher-Wellman, K.H. (2020). Novel approach to quantify mitochondrial content and intrinsic
bioenergetic efficiency across organs. Sci. Rep. *10*, 17599. https://doi.org/10.1038/s41598-
020-74718-1.

43. Luengo, A., Li, Z., Gui, D.Y., Sullivan, L.B., Zagorulya, M., Do, B.T., Ferreira, R., Naamati, A., Ali,
A., Lewis, C.A., et al. (2021). Increased demand for NAD+ relative to ATP drives aerobic
glycolysis. Mol. Cell *81*, 691-707.e6. https://doi.org/10.1016/j.molcel.2020.12.012.

44. Giddings, E.L., Champagne, D.P., Wu, M.-H., Laffin, J.M., Thornton, T.M., Valenca-Pereira, F.,
Culp-Hill, R., Fortner, K.A., Romero, N., East, J., et al. (2021). Mitochondrial ATP fuels ABC
transporter-mediated drug efflux in cancer chemoresistance. Nat. Commun. *12*, 2804.
https://doi.org/10.1038/s41467-021-23071-6.

45. Nandi, I., Ji, L., Smith, H.W., Avizonis, D., Papavasiliou, V., Lavoie, C., Pacis, A., Attalla, S.,
Sanguin-Gendreau, V., and Muller, W.J. (2024). Targeting fatty acid oxidation enhances
response to HER2-targeted therapy. Nat. Commun. *15*, 6587.
https://doi.org/10.1038/s41467-024-50998-3.

46. Wang, Y.Y., Attané, C., Milhas, D., Dirat, B., Dauvillier, S., Guerard, A., Gilhodes, J., Lazar, I.,
Alet, N., Laurent, V., et al. Mammary adipocytes stimulate breast cancer invasion through
metabolic remodeling of tumor cells. JCI Insight *2*, e87489.
https://doi.org/10.1172/jci.insight.87489.

47. Matuszewska, K., Pereira, M., Petrik, D., Lawler, J., and Petrik, J. (2021). Normalizing Tumor
Vasculature to Reduce Hypoxia, Enhance Perfusion, and Optimize Therapy Uptake. Cancers
13, 4444. https://doi.org/10.3390/cancers13174444.

48. Wu, S., Akhtari, M., and Alachkar, H. (2018). Characterization of Mutations in the
Mitochondrial Encoded Electron Transport Chain Complexes in Acute Myeloid Leukemia. Sci.
Rep. *8*, 13301. https://doi.org/10.1038/s41598-018-31489-0.

49. Schöpf, B., Weissensteiner, H., Schäfer, G., Fazzini, F., Charoentong, P., Naschberger, A.,
Rupp, B., Fendt, L., Bukur, V., Giese, I., et al. (2020). OXPHOS remodeling in high-grade
prostate cancer involves mtDNA mutations and increased succinate oxidation. Nat.
Commun. 2020 111 *11*, 1–16. https://doi.org/10.1038/s41467-020-15237-5.

50. Zhou, Y., Long, D., Zhao, Y., Li, S., Liang, Y., Wan, L., Zhang, J., Xue, F., and Feng, L. (2022).
Oxidative stress-mediated mitochondrial fission promotes hepatic stellate cell activation via
stimulating oxidative phosphorylation. Cell Death Dis. *13*, 1–15.
https://doi.org/10.1038/s41419-022-05088-x.

51. Lhuissier, C., Desquiret-Dumas, V., Girona, A., Alban, J., Faure, J., Cassereau, J., Codron, P.,
Lenaers, G., Baris, O.R., Gueguen, N., et al. (2024). Mitochondrial F0F1-ATP synthase
governs the induction of mitochondrial fission. iScience *27*.
https://doi.org/10.1016/j.isci.2024.109808.

52. Zou, W., Chen, Q., Slone, J., Yang, L., Lou, X., Diao, J., and Huang, T. (2021). Nanoscopic
quantification of sub-mitochondrial morphology, mitophagy and mitochondrial dynamics in
living cells derived from patients with mitochondrial diseases. J. Nanobiotechnology *19*,
136. https://doi.org/10.1186/s12951-021-00882-9.

53. Ren, K., Zhou, D., Wang, M., Li, E., Hou, C., Su, Y., Zou, Q., Zhou, P., and Liu, X. (2021).
RACGAP1 modulates ECT2-Dependent mitochondrial quality control to drive breast cancer
metastasis. Exp. Cell Res. *400*, 112493. https://doi.org/10.1016/j.yexcr.2021.112493.

54. Zhao, J., Zhang, J., Yu, M., Xie, Y., Huang, Y., Wolff, D.W., Abel, P.W., and Tu, Y. (2013).
Mitochondrial dynamics regulates migration and invasion of breast cancer cells. Oncogene
32, 4814–4824. https://doi.org/10.1038/onc.2012.494.

55. Han, X.-J., Yang, Z.-J., Jiang, L.-P., Wei, Y.-F., Liao, M.-F., Qian, Y., Li, Y., Huang, X., Wang, J.-B.,
Xin, H.-B., et al. (2015). Mitochondrial dynamics regulates hypoxia-induced migration and
antineoplastic activity of cisplatin in breast cancer cells. Int. J. Oncol. *46*, 691–700.
https://doi.org/10.3892/ijo.2014.2781.

56. Zamberlan, M., Boeckx, A., Muller, F., Vinelli, F., Ek, O., Vianello, C., Coart, E., Shibata, K.,
Christian, A., Grespi, F., et al. (2022). Inhibition of the mitochondrial protein Opa1 curtails
breast cancer growth. J. Exp. Clin. Cancer Res. CR *41*, 95. https://doi.org/10.1186/s13046-
022-02304-6.

57. Yin, M., Lu, Q., Liu, X., Wang, T., Liu, Y., and Chen, L. (2016). Silencing Drp1 inhibits glioma
cells proliferation and invasion by RHOA/ ROCK1 pathway. Biochem. Biophys. Res. Commun.
478, 663–668. https://doi.org/10.1016/j.bbrc.2016.08.003.

58. Tang, Q., Liu, W., Zhang, Q., Huang, J., Hu, C., Liu, Y., Wang, Q., Zhou, M., Lai, W., Sheng, F.,
et al. (2018). Dynamin-related protein 1-mediated mitochondrial fission contributes to IR-
783-induced apoptosis in human breast cancer cells. J. Cell. Mol. Med. *22*, 4474–4485.
https://doi.org/10.1111/jcmm.13749.

59. Novotna, E., Milosevic, M., Prukova, D., Magalhaes-Novais, S., Dvorakova, S., Dmytruk, K.,
Gemperle, J., Zudova, D., Nickl, T., Vrbacky, M., et al. (2024). Mitochondrial HER2 stimulates
respiration and promotes tumorigenicity. Eur. J. Clin. Invest. *54*, e14174.
https://doi.org/10.1111/eci.14174.

60. Ding, Y., Liu, Z., Desai, S., Zhao, Y., Liu, H., Pannell, L.K., Yi, H., Wright, E.R., Owen, L.B., Dean-
Colomb, W., et al. (2012). Receptor tyrosine kinase ErbB2 translocates into mitochondria
and regulates cellular metabolism. Nat. Commun. *3*, 1271.
https://doi.org/10.1038/ncomms2236.

61. Kurmi, K., Hitosugi, S., Yu, J., Boakye-Agyeman, F., Wiese, E.K., Larson, T.R., Dai, Q., Machida,
Y.J., Lou, Z., Wang, L., et al. (2018). Tyrosine Phosphorylation of Mitochondrial Creatine
Kinase 1 Enhances a Druggable Tumor Energy Shuttle Pathway. Cell Metab. *28*, 833-847.e8.
https://doi.org/10.1016/j.cmet.2018.08.008.

62. Wang, D., Townsend, L.K., DesOrmeaux, G.J., Frangos, S.M., Batchuluun, B., Dumont, L.,
Kuhre, R.E., Ahmadi, E., Hu, S., Rebalka, I.A., et al. (2023). GDF15 promotes weight loss by

enhancing energy expenditure in muscle. Nature *619*, 143–150.

https://doi.org/10.1038/s41586-023-06249-4.

63. Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast

and bias-aware quantification of transcript expression. Nat. Methods *14*, 417–419.

https://doi.org/10.1038/nmeth.4197.

64. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.

https://doi.org/10.1186/s13059-014-0550-8.

65. Marini, F., and Binder, H. (2019). pcaExplorer: an R/Bioconductor package for interacting

with RNA-seq principal components. BMC Bioinformatics *20*, 331.

https://doi.org/10.1186/s12859-019-2879-1.

66. Mi, H., Muruganujan, A., Huang, X., Ebert, D., Mills, C., Guo, X., and Thomas, P.D. (2019).

Protocol Update for large-scale genome and gene function analysis with the PANTHER

classification system (v.14.0). Nat. Protoc. *14*, 703–721. https://doi.org/10.1038/s41596-

019-0128-8.

67. Paglialunga, S., Ludzki, A., Root-McCaig, J., and Holloway, G.P. (2015). In adipose tissue,

increased mitochondrial emission of reactive oxygen species is important for short-term

high-fat diet-induced insulin resistance in mice. Diabetologia *58*, 1071–1080.

https://doi.org/10.1007/s00125-015-3531-x.

68. Miotto, P.M., and Holloway, G.P. (2019). Exercise-induced reductions in mitochondrial ADP

sensitivity contribute to the induction of gene expression and mitochondrial biogenesis

through enhanced mitochondrial H2O2 emission. Mitochondrion *46*, 116–122.

https://doi.org/10.1016/j.mito.2018.03.003.

69. Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M. (2016). KEGG as a

reference resource for gene and protein annotation. Nucleic Acids Res. *44*, D457–D462.

https://doi.org/10.1093/nar/gkv1070.

70. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann, A.,

Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene Set Knowledge

Discovery with Enrichr. Curr. Protoc. *1*, e90. https://doi.org/10.1002/cpz1.90.

1007

A

Benign

420 common proteins
(p adj. <0.1)

9 **Upregulated** in cancer
411 **Downregulated** in cancer

1848

Tumors

B

Benign
Tumor

PC2
(9% variance)

PC1
(57% variance)

C

Downregulated
Upregulated

-log₁₀adj-p-value

log₂FC

D

N=411 DOWNREGULATED TUMOR PROTEINS

E REACTOME 2022

Metabolism R-HSA-1430728
Respiratory Electron Transport R-HSA-611105
Citric Acid (TCA) Cycle And Respiratory Electron Transport R-HSA-1428517
Respiratory Electron Transport, ATP Synthesis By Chemiosmotic Coupling, Heat Production By Uncoupling Proteins R-HSA
Complex I Biogenesis R-HSA-6799198
Fatty Acid Metabolism R-HSA-8978868
Mitochondrial Fatty Acid Beta-Oxidation R-HSA-77289
Platelet Activation, Signaling And Aggregation R-HSA-76002
Hemostasis R-HSA-109582
Branched-chain Amino Acid Catabolism R-HSA-70895

F GO BIOLOGICAL PROCESS 2023

Cellular Respiration (GO:0045333)
Mitochondrial ATP Synthesis Coupled Electron Transport (GO:0042775)
Aerobic Electron Transport Chain (GO:0019646)
Oxidative Phosphorylation (GO:0006119)
Aerobic Respiration (GO:0009060)
Proton Motive Force–Driven Mitochondrial ATP Synthesis (GO:0042776)
Mitochondrial Electron Transport, NADH To Ubiquinone (GO:0006120)
Proton Motive Force–Driven ATP Synthesis (GO:0015986)
Fatty Acid Beta-Oxidation (GO:0006635)
Fatty Acid Catabolic Process (GO:0009062)

G KEGG ENRICHMENT ANALYSIS

Amyotrophic lateral sclerosis
Chemical carcinogenesis-reactive oxygen species
Huntington disease
Parkinson disease
Prion disease
Thermogenesis
Diabetic cardiomyopathy
Non-alcoholic fatty liver disease
Oxidative Phosphorylation
Propanoate Metabolism

GeneRatio

p.adjust
8.600880e-24
9.294614e-18
1.858922e-17
2.788383e-17
3.717843e-17

Count
20
30
40
50

BENIGN   TUMOR

Figure 7

A

NORMALIZATION: WET WEIGHT

Benign

Tumor

NORMALIZATION: TOTAL PROTEIN

Benign

Tumor

Protein

NORMALIZATION: MITOCHONDRIAL PROTEIN

Benign

Tumor

B **CI+CII SUPPORTED RESPIRATION**
per wet weight

JO_2 (pmol•s^{-1}•mg ww^{-1})

E   +PM   +D   +G   +S

C **LIPID-SUPPORTED RESPIRATION**
per wet weight

JO_2 (pmol•s^{-1}•mg ww^{-1})

E   +MD   +PC

D **CI+CII SUPPORTED RESPIRATION**
per total protein

JO_2 (pmol•s^{-1}•mg pro^{-1})

E   +PM   +D   +G   +S

E **LIPID-SUPPORTED RESPIRATION**
per total protein

JO_2 (pmol•s^{-1}•mg pro^{-1})

E   +MD   +PC

F **CI+CII SUPPORTED RESPIRATION**
per mitochondrial protein

JO_2 (pmol•s^{-1}•mg pro^{-1})

E   +PM   +D   +G   +S

G **LIPID-SUPPORTED RESPIRATION**
per mitochondrial protein

JO_2 (pmol•s^{-1}•mg pro^{-1})

E   +MD   +PC

CANONICAL HER2 SIGNALING - PROTEIN

N=741 UNIQUE BENIGN MAMMARY TISSUE PROTEINS

N=600 UNIQUE HER2-DRIVEN TUMOR PROTEINS

$$MEF = \frac{MitoCarta\ Protein\ Abundance}{Total\ Protein\ Abundance}$$

A

LFQ

*#

Total
MitoCarta

BEN   TUM

B

MEF
(MitoCarta/Total)

*

BEN   TUM

Abbrev.	Protein	Pathway/Function	Fc (vs BEN)	p.adj
PGK1	Phosphoglycerate kinase	Glycolysis	2.48	0.068
KPYM	Pyruvate kinase	Glycolysis	3.17	0.048
MP2K4	Mitogen activated protein kinase kinase 4	MAPK Pathway	3.65	0.012
RL38	Large ribosomal subunit protein eL38	Protein synthesis/translation	4.11	0.078
PGTA	Geranylgeranyl transferase type-2 subunit alpha	Cysteine prenylation of Rab proteins	4.28	0.067
Rl23	Large ribosomal subunit protein uL14	Protein synthesis/translation	4.41	0.042
CADH1	Cadherin-1	Cell adhesion and motility, epithelial cell marker	12.75	0.084
LMCD1	LIM and cysteine rich domains protein 1	Transcriptional cofactor for GATA6 involved in cell proliferation and metastasis (MMP, EGFR)	15.26	0.077
MPRI	Cation independent mannose 6 phosphate receptor	Lysosomal/Golgi trafficking, associates with IGFR2	22.3	0.093