Decoding Liver Cancer Prognosis: From Multi-omics Subtypes, Prognostic Models to Single Cell

Validation

Yanbin Wang[1], Yuqi Wu[1], Hong Zhang[1], Xiao Zhou[1], Anping Song[1], Hong Qiu[1], Xianglin Yuan[1], Hua

Xiong[1]*, Yanmei Zou[1]*

[1*]Department of Oncology, Tongji Hospital, Tongji Medical College, Huazhong University of Science

and Technology(HUST), Wuhan, 430030, China.

Running title: Prognostic Models Based on Multi-omics Subtypes.

***Corresponding author**.

*These authors contributed equally to this work and should be considered co-corresponding

authors.

E-mail: cnhxiong@tjh.tjmu.edu.cn; whtjzym@tjh.tjmu.edu.cn;

Postal address: 1095 Liberation Avenue, Qiaokou District, Wuhan City, Hubei Province, China;

Contributing authors: ybwang0907@hust.edu.cn; 18049463686@163.com; 1954528173@qq.com;

zhouxiaotj@qq.com; anpingsong@tjh.tjmu.edu.cn; qiuhong@hust.edu.cn; yuanxianglin@hust.edu.cn.

Conflict of interest disclosure statement: The authors declare no potential conflicts of interest.

The main text of the study consists of 4854 words and 8 figures.

Abstract

Purpose: Hepatocellular carcinoma (HCC) is a highly aggressive tumor characterized by significant heterogeneity and invasiveness, leading to a lack of precise individualized treatment strategies and poor patient outcomes. This necessitates the urgent development of accurate patient stratification methods and targeted therapies based on distinct tumor characteristics.

Experimental Design: By integrating gene expression data from The Cancer Genome Atlas (TCGA), International Cancer Genome Consortium (ICGC), and Gene Expression Omnibus (GEO), we identified subtypes through a multi-omics consensus clustering approach amalgamated from 10 clustering techniques. Subsequently, we developed a prognostic model, employing machine learning algorithms, based on subtype classification features. Finally, by analyzing single cell sequencing data, we investigated the mechanisms driving prognostic variations among distinct subtypes.

Results: First, we developed a novel consensus clustering method that categorizes liver cancer patients into two subtypes, CS1 and CS2. Second, we constructed a prognostic prediction model, which demonstrated superior predictive accuracy compared to several models published in the past five years. Finally, we observed differences between CS1 and CS2 in various metabolic pathways, biological processes, and signaling pathways, such as fatty acid metabolism, hypoxia levels, and the PI3K-AKT signaling pathway.

Introduction

Liver cancer is currently one of the most common tumors, and among primary liver cancers, hepatocellular carcinoma (HCC) is the most common pathological type. In 2022, liver cancer saw widespread new cases globally, and its incidence rate has been gradually increasing in recent years, ranking 6th among all tumors worldwide[1]. Meanwhile, liver cancer ranks 3rd among tumors causing patient deaths, with a relative 5-year survival rate of about 18%, highlighting the significantly poor prognosis of this disease[2].

With the advancement of current treatment technologies, there are ongoing attempts to employ

advanced treatment methods for innovative liver cancer therapy. Nevertheless, up to now, the progress

of strategies aimed at treating liver cancer, especially unresectable liver cancer, has been limited. For

instance, targeted therapies and immunotherapies have been the focus of extensive research efforts and

have shown promising results in some patients. However, a significant number of patients still do not

obtain clinical benefits from these approaches. This may be due to the significant heterogeneity of liver

cancer; classifying liver cancer patients according to molecular subtypes might help overcome this

difficulty[3]. This urgently necessitates leveraging large-scale multi-omics data and advanced machine

learning algorithms to identify biomarkers capable of efficiently predicting clinical treatment outcomes

for liver cancer patients.

In this study, we combined mRNA, long non-coding RNA (lncRNA), and microRNA (miRNA)

expression profiles, genomic mutations, and epigenomic DNA methylation data, using a consensus

strategy integrating 10 multi-omics clustering methods to develop comprehensive consensus subtypes

of liver cancer, and conducted preliminary exploration of the molecular characteristics and potential

clinical applications of the two subtypes. Subsequently, we further expanded the clinical application

value of the subtype. Based on the differential gene expression profiles of different subtypes, we used a

combination of 101 machine learning algorithms to construct the best-performing prognostic model.

This prognostic model showed excellent prognostic value in both training and validation cohorts, and

outperformed several other prognostic models in comparisons. Finally, we validated the subtypes at the

single-cell sequencing cell level using the same typing method, and further explored the possible

causes and mechanisms of prognostic differences in patients with different subtypes.

Materials and Methods

Data collection and processing

Download the molecular profiles of the TCGA-LIHC cohort dataset from the TCGA database,

including 432 LIHC cases with complete transcriptome expression, somatic mutations, copy number

alterations (CNA), DNA methylation, and clinical information. The transcriptome profiles of mRNA

and lncRNA were obtained using the "TCGAbiolinks" R package[4], and GENCODE27 was used to

convert Ensemble IDs to gene symbols; Somatic mutation data were also obtained using

"TCGAbiolinks" R package and processed with the "maftools" R package[5]; DNA methylation profiles

and clinical information were downloaded from UCSC Xena (https://xenabrowser.net/); Meanwhile,

we also downloaded 445 LIHC cases with complete transcriptome expression and clinical information

from the LIHC.JP cohort in the ICGC database (https://dcc.icgc.org/); The single-cell sequencing

dataset is sourced from the GEO database (https://www.ncbi.nlm.nih.gov/geo/, accession number

GSE229772).

Construct multi-omics consensus clustering classification and perform validation

In this study, we utilized the "getElites" function from the multi-omics integration and visualization

(MOVICS) package to screen gene features in cancer subtypes[6]. For continuous variables (mRNA,

lncRNA, miRNA, and methylation), we set the "method" parameter of the "getElites" function to

"mad" to screen the top 1000 genes with the greatest variability;   For binary variable gene mutation

data, we conducted screening based on mutation frequency by setting the "method" parameter to "freq"

to identify genes with a mutation frequency exceeding 10. The results from these five dimensions were

incorporated into our study for further analysis. After the initial feature selection, we used the

"getClustNum" function in the MOVICS package to further determine the optimal number of clusters

for our study and ultimately decided to divide them into two subtypes. Next, we applied the

"getMOIC" function for clustering analysis. We used 10 clustering algorithms (CIMLR,

ConsensusClustering, SNF, iClusterBayes, PINSPlus, moCluster, NEMO, IntNMF, COCA, and LRA)

as input for the "methods.list" parameter and employed the default parameters provided by the

MOVICS package. As a result, we obtained the clustering results for each method. After calculating the

clustering results of the 10 methods, we integrated the results of different algorithms based on the

concept of consensus clustering using the "getConsensusMOIC" function, obtaining the final clustering

results. Subsequently, we conducted nearest template prediction(NTP) analysis, performed subtyping in

the external ICGC-JP cohort, and validated the subtyping method.

Molecular Characteristics of Multi-Omics Consensus Clustering Subtypes

We analyzed the mutation status using the "maftools" R package, then used the limma method to

analyze the differentially expressed genes between the two subtypes[7], and performed enrichment

analysis with the clusterProfiler R package[8].

Cell, signaling pathways, stemness, and other phenotypic characteristics of multi-omics

consensus clustering subtypes

Using ssGSEA analysis, we calculated the enrichment scores of different immune cell characteristics

between the two subtypes based on immune cell gene sets reported in the literature[9]; Next, we

analyzed the activation status of cancer pathways and some biological processes in patients using the

GSEA algorithm, and analyzed the sensitivity of the two subtypes to conventional therapeutic drugs.

Both of these steps were performed using the "GSVA" R package[10]. The expression data from CCLE

were sourced from the Broad Institute Cancer Cell Line Encyclopedia

(https://sites.broadinstitute.org/ccle/); The CTRP v.2.0 (https://portals.broadinstitute.org/ctrp) and

PRISM dataset (24Q2; https://depmap.org/portal/prism/) were utilized to acquire drug sensitivity data;

The half-maximal inhibitory concentration (IC50) was used as an indicator of drug sensitivity.

Constructing a prognostic prediction model based on the consensus clustering of multi-omics

features

To build the prognostic model, we randomly split the TCGA-LIHC cohort into training and validation

sets at a 1:1 ratio and incorporated 10 machine learning algorithms, including CoxBoost, stepwise Cox,

Lasso, Ridge, Enet, survival-SVMs, GBMs, SuperPC, plsRcox, and RSF, creating a total of 101

different combinations of these algorithms[11]. Ultimately, these 101 algorithm combinations will be

utilized to develop the most predictive prognostic model with the best C-index performance. We then

scored each sample in the training and validation sets based on the model and classified the samples

into high-risk and low-risk groups according to the optimal cut-off value of scores, and the

"survminer" R package(https://github.com/kassambara/survminer/tree/master) was used to determine

the optimal cut-off value. Cox regression and Kaplan–Meier analyses were performed via the "survival"

R package(https://CRAN.R-project.org/package=survival).

Comparison of Prognostic Prediction Models

We evaluated the prognostic predictive ability of all models included in the study using the C-index in

each cohort. To enhance the clinical utility of the model, we constructed a nomogram based on factors

obtained from multivariable Cox regression. We plotted time-dependent C-index curves and calibration

curves to describe accuracy and used decision curve analysis to calculate the clinical benefit for

patients.

scRNA-seq Data Processing

We utilized the "Seurat" R package to process single-cell sequencing data from the GSE229772

dataset[12]. We removed samples with over 10% mitochondrial genes and over 5% ribosomal genes, and

filtered out genes expressed in fewer than 3 cells. Next, we identified the most variable genes using the

FindVariableFeatures function from the Seurat R package, conducted dimensionality reduction via

principal component analysis (PCA), and identified a total of 13 clusters through t-SNE clustering

analysis. Finally, we annotated the cell populations by combining the expression of characteristic genes

in each cell cluster with the expression of commonly used cell-type-specific genes across the clusters.

Characteristic Analysis of Different Cell Subpopulations

We used the "irGSEA" R package[13], employing 6 methods (AUCell, UCell, singscore, ssGSEA,

JASMINE, and viper) for pathway enrichment analysis among different cell populations, and

performed a comprehensive evaluation of the differential analysis results using robust rank aggregation

(RRA) to identify gene sets significantly enriched in most enrichment analysis methods. Moreover, we

additionally utilized the "scMetabolism" R package to compare the metabolic difference across various

cell types[14]. The metabolic activity of each cell was quantified through integrating the gene sets

contained in the KEGG database. Subsequently, we used the "CellChat" R package to explore the

interaction patterns among cell populations[15], defining ligands and receptors as outgoing and

incoming signals, respectively.

Characteristics of Different Subtypes of Malignant Cells at the Single-Cell Level

We applied the "CopyKAT" R package to analyze gene expression levels across all cells[16], identifying

diploid cells as normal cells, with normal and tumor cells separated due to differences in gene

expression distributions. Subsequently, we classified tumor cells based on the previously obtained

multi-omics consensus clustering approach, and used the irGSEA R package to perform pathway

enrichment analysis on different subtypes, conducted metabolic differential analysis with the

scMetabolism R package, and ultimately explored the interaction patterns of different subtype

malignant cells with other cell populations using the CellChat package.

Results

Construct multi-omics consensus clustering classification and perform validation

We utilized clustering prediction indicators, gap statistical analysis, silhouette scores, and insights from

previous research involving 10 multi-omics integrated clustering algorithms to ascertain the number of

subtypes (Fig.1A). Using a consensus integration method, we linked the clustering results to distinct

molecular expression patterns of the transcriptome (mRNA, lncRNA, and miRNA), epigenetic

methylation, and somatic mutations, thereby identifying two subtypes (CS1 and CS2) independently

(Fig.1B). Significant differences in mRNA, lncRNA, miRNA, DNA CpG methylation sites, and

mutation gene characteristics were observed between the two subtypes (Fig.1C). This classification

method is strongly correlated with overall survival ($p < 0.001$), with patients of the CS2 subtype

exhibiting better prognostic survival compared to those of the CS1 subtype (Fig.1D). These

characteristics suggest that our classification method can effectively distinguish between patients.

Subsequently, employing the NTP method, we selected 200 genes specifically upregulated in each

subtype as classifiers to validate the classification in an independent external validation set, ICGC, and

found a high degree of consistency in the classification results. Patients with CS1 and CS2 subtypes in

the ICGC cohort exhibited similar prognostic differences (Fig.1E-G). This demonstrates the versatility

and practical value of our classification method.

Molecular Characteristics of Multi-Omics Consensus Clustering Subtypes

We analyzed the gene mutation landscape of two subtypes and found that both subtypes exhibited a

high frequency of mutations in TP53, KMT2D, and DNAH6 genes(Fig.2A). Based on the correlation

between tumor mutation burden (TMB) and response to immunotherapy, we initially evaluated the

TMB of patients within the two subtypes. The results indicated no significant difference in TMB

between the two subtypes (Fig.2B-C). Subsequently, we analyzed the bulk RNA sequencing data of

patients within the two subtypes, identified differentially expressed genes, and conducted enrichment

analysis on these genes. The results revealed significant differential enrichment in several biological

pathways, including those related to metabolism and signal transduction, such as lipid peroxidation,

fatty acid metabolism, and amino acid metabolism, suggesting notable biological differences between

the tumors and associated tissues of the two subtypes (Fig.2D-E). Previous studies have reported that

heterogeneity among different tumor tissues of the same tumor type largely stems from metabolic

reprogramming and changes in signal transduction pathway activity[17-18]. This implies that the

heterogeneity between the tumor tissues of the two subtypes may also arise from these factors. Finally,

we compared the consistency of subtyping methods with the pathologic staging of patients(Fig.2F).

Cell, signaling pathways, stemness, and other phenotypic characteristics of multi-omics

consensus clustering subtypes

Given the impact of the tumor immune microenvironment on tumor tissue heterogeneity and patient

treatment responsiveness, we used the ssGSEA algorithm to assess the enrichment status of

immune-related cells in the tumor microenvironment (TME) of two subtypes(Fig.3A). The gene set

utilized for this analysis was compiled and summarized from various literature sources. Then, we

employed drug intervention data to evaluate the sensitivity of the two subtypes to commonly used

cancer therapies. The CS2 subtype exhibited higher sensitivity to 5-fluorouracil, Gefitinib, Afatinib,

and Erlotinib compared to the CS1 subtype (Fig.3B). Among these, 5-fluorouracil is the main drug

used in hepatic arterial infusion chemotherapy for hepatocellular carcinoma (HCC) patients, whereas

Gefitinib, Afatinib, and Erlotinib are tyrosine kinase inhibitors (TKIs) that primarily inhibit epidermal

growth factor receptor (EGFR). We further compared the level of infiltration of each cell taxon in

subtypes in the TME(Fig.3C). Within the TME of the CS2 subtype, associated with a comparatively

more favorable prognosis, nearly all assessed cell distribution levels demonstrated a decrease in

comparison to those in the CS1 subtype. This encompasses traditionally recognized inhibitory immune

cells like MDSCs and Regulatory T cells, alongside pivotal tumor-killing immune cells like Activated

CD8+ T cells and NK cells. These findings may imply reduced levels of tumor-associated antigens

within CS2 tumor tissue, leading to a comparatively subdued activation and induction of immune cells

within the TME. We also evaluated several common biological pathways that may influence the TME

and tumor tissue responsiveness to treatment. We discovered that the CS1 subtype is more active than

the CS2 subtype in multiple pathways associated with tumor development and progression, including

the MAPK, JAK-STAT, NFκB, PI3K-AKT, and VEGF signaling pathways, as well as pathways

involved in angiogenesis and hypoxia. No significant differences were observed in the cAMP signaling

pathway or the mRNAsi stemness score (Fig. 3D).

Constructing a prognostic prediction model based on the consensus clustering of multi-omics

features

In a previous study, we utilized the mRNA expression profiles of TCGA-LIHC from the TCGA

database to compute the differential gene expression profiles between CS1 and CS2 subtypes. We

selected the top 30 significantly different genes as candidate genes for prognostic risk. Subsequently,

we applied 101 algorithmic approaches to determine the correlation and impact weights between these

30 genes and prognosis, leading to the construction of a prognostic prediction model. Simultaneously,

we calculated the average C index of each model to assess the predictive capability across all models

(Fig. 4A). Employing this model to evaluate patients' prognostic risk within the training set

TCGA-LIHC cohort and stratifying them based on the optimal cutoff value, we observed a notably

poorer prognosis among high-risk group patients compared to their low-risk counterparts. The model

comprises 5 central genes, and the Hazard Ratio associated with these genes in the TCGA-LIHC cohort

is depicted in Fig. 4C. Additionally, we illustrated the patients' prognostic outcomes and variations in

central genes as the prognostic risk escalated (Fig. 4D). Subsequently, we validated the model using an

independent external validation set, the ICGC cohort. Our findings demonstrated its efficacy in

stratifying patients into high and low-risk groups within the ICGC cohort. A notable disparity in

prognosis between the two groups was evident, aligning with the outcomes observed in the training set

(Fig. 4E-F).

Comparison of Prognostic Prediction Models

With the rapid advancement and widespread adoption of Next Generation Sequencing (NGS)

technology, numerous prognostic prediction models based on gene expression have been extensively

reported. After developing our model, we reviewed and collected prognostic prediction models

reported in the past five years and included seven of these models, which are based on various

foundations such as immune infiltration, glucose metabolism, and lipid metabolism. We calculated the

C-index for a total of eight models using the training set TCGA-LIHC cohort and the validation set

ICGC cohort. The results indicated that our model demonstrated superior prognostic prediction

performance compared to all other models in both cohorts (Fig. 5A). Subsequently, we incorporated

clinical features related to prognosis into the model to construct a nomogram (Fig. 5B). The calibration

curve demonstrated that the nomogram had predictive accuracy more consistent with actual conditions

(Fig. 5C). Decision curve analysis (DCA) indicated that both the nomogram and the original model

could benefit patients, with the nomogram performing slightly better (Fig. 5D).

scRNA-seq Data Processing

Given the substantial prognostic variances between CS1 and CS2 patients and the notable

heterogeneity in tumor tissues at both cellular and molecular levels, we sought to explore the potential

mechanisms driving these differences through scRNA-seq data analysis. The GSE229772 dataset from

the GEO database was chosen for our analysis. Following quality control filtering, 56,938 cells were

derived from 31 samples. By employing hierarchical and unsupervised clustering methods, we

delineated cell subclusters according to their expression profiles, categorizing the cells into 13

subclusters (Fig. 6A). Subsequently, we employed established marker genes to profile various cell

types and assign annotations, discerning 12 primary cell subgroups: B cells, stromal cells, vascular

endothelial cells, CD8+ T cells, epithelial cells, fibroblasts, germ cell-like cells, hepatic stellate cells,

hepatocytes, macrophages, neutrophils, and T cells (Fig. 6B-D). Within these, the immune cell group

prevailed as the most abundant, succeeded by stromal cells and fibroblasts (Fig. 6E).

Characteristic Analysis of Different Cell Subpopulations

To investigate pathway heterogeneity among different cell subclusters, we identified differentially

expressed characteristic genes within these subclusters. Subsequently, we employed multiple

algorithms, including AUCell, UCell, singscore, ssgsea, JASMINE, and viper, to assess pathway

activity variation. Differential analysis results were comprehensively evaluated using robust rank

aggregation (RRA), and gene sets significantly enriched across multiple enrichment analysis methods

were filtered to obtain consensus results. Statistically significant differentially enriched pathways from

these consensus results were visualized (Fig.7A-B). We observed significant differences in several

metabolic and signaling pathways among different cell subclusters. Given the reported impact of

metabolic changes and signaling pathway variations on the TME and their implications for

heterogeneity, we specifically analyzed these pathways. Utilizing the VISION algorithm, we assessed

metabolic levels across each cell subcluster and visualized the differences in a dot plot (Fig.7C). Our

findings indicated pronounced metabolic variations among cell subclusters, with germ cell-like cells

and hepatocytes notably exhibiting heightened metabolic activity compared to other cell groups.

Subsequently, we utilized the CellChat R package's integrated databases for secreted signaling, ECM

receptor signaling, cell-cell contact, and non-protein signaling to explore cell interactions within the

single-cell dataset. Initially, we examined the cell communication landscape between distinct cell

groups and constructed a cell communication network (Fig.7D). Our analysis revealed robust

communication links between various cell groups, particularly notable between fibroblasts and hepatic

stellate cells, which exhibited signaling connections with nearly all other cells. Then, we visualized the

signaling output and input patterns (Fig.7E-G) to illustrate the contributions of different cell groups at

both ends.

Characteristics of Different Subtypes of Malignant Cells at the Single-Cell Level

To delve deeper into the heterogeneity of the two tumor subtypes, we initially utilized the copykat

algorithm to pinpoint potential malignant cell subclusters, including hepatocytes, fibroblasts, and

pericellular cells, by detecting whole-genome aneuploidy. This approach facilitated the identification of

potential malignant cells, which were subsequently aligned with the prior cell annotation results

(Fig.8A). Subsequently, we employed the previously derived multi-omics consensus clustering features

to classify the malignant cells into two subtypes: CS1 and CS2. This analysis enabled the identification

of differentially expressed genes between the two subtypes. Enrichment analysis of these genes

revealed significant differences in various metabolic and signaling pathways between CS1 and CS2

malignant cells, including purine metabolism, pyrimidine metabolism, glycolysis, the PI3K-AKT

signaling pathway, and the P53 signaling pathway (Fig.8B). A targeted analysis of the metabolic

variances between the CS1 and CS2 subtypes revealed notable differences, indicating significantly

higher metabolic activity in CS1 compared to CS2 (Fig.8C). Then, upon integrating the malignant cells

classified as CS1 and CS2 into the comprehensive cell communication network, we observed that CS1

and CS2 exhibited widespread regulation of all other cell groups in the TME (Fig.8D). Subsequently,

we analyzed the signal output and input patterns of various cellular subgroups and found that the MIF

signaling pathway exhibited the highest frequency and greatest contribution in the signal

communication network, whether at the input or output end(Fig.8E-G). We further analyzed the output

levels of the MIF signaling pathway in various cellular subgroups, focusing on which cells were

outputting the MIF signal. The results showed that the CS1 and CS2 subtypes mainly output MIF, with

CS1 subtype being relatively predominant(Fig.8H). Further investigation into the regulatory effects of

CS1 malignant cells on other cell groups demonstrated that the CS1 subtype exerted extensive

regulatory effects on nearly all immune cells in the TME, including CD8+ T cells, macrophages, B

cells, and neutrophils, primarily through the MIF- (CD74+CXCR4) pathway (Fig.8I).

Discussion

The role of genes in disease development, particularly in tumorigenesis, is well-established. Studying

gene expression processes significantly enhances our understanding of tumors. Gene expression is

intricately regulated by genetic mechanisms such as methylation and mutations. Therefore, integrating

multi-omics data analysis is crucial for gaining deeper insights into disease-specific regulatory

mechanisms. However, most studies to date have primarily focused on single omics, such as mRNA

and miRNA expression research. Moreover, the choice of omics clustering methods in research often

relies on personal preference, exacerbating the limitations of individual approaches in studies. Our

research aims to bridge this gap by integrating ten state-of-the-art clustering algorithms to identify two

prognostic subtypes with distinct characteristics, which could potentially facilitate precise stratification

and treatment of HCC patients. Furthermore, our novel classification method has demonstrated stability

across multiple cohorts.

Significant prognostic differences among patients with the same tumor type may be influenced by

various factors, including tumor tissue heterogeneity, metabolic reprogramming, signaling patterns, and

disparities in the TME. As such, our study undertook initial investigations to elucidate the

underpinnings of these prognostic discrepancies across patient subtypes. Initially, we evaluated the

mutation landscape across subtypes and observed no notable differences in TMB and FGA between the

two subtypes, implying that subtype heterogeneity may not originate from gene mutations. Subsequent

analysis of differentially expressed genes and gene enrichment indicated significant enrichment in

tumor-related pathways, particularly metabolism and signal transduction pathways like lipid

peroxidation, fatty acid metabolism, amino acid metabolism, PI3K-NOX, and EGF-EGFR signaling

pathways. This suggests potential variances in metabolic reprogramming and signaling patterns

between the subtypes, which can influence patient prognosis by governing the TME. Consequently, we

conducted a further analysis of immune cell infiltration in the TME, revealing lower immune cell

infiltration in CS2 compared to CS1, implying a potentially lower malignancy in CS2 with reduced

release of tumor-associated antigens and subsequent lower immune response and infiltration.

Additionally, we thoroughly examined the signaling pathways and biological characteristics closely

associated with tumor development. The findings revealed that the CS1 subtype exhibited greater

activity than the CS2 subtype in multiple pathways linked to tumorigenesis, including the MAPK,

JAK-STAT, NFκB, PI3K-AKT, VEGF, angiogenesis, and hypoxia pathways. These pathways are

widely recognized for their close association with tumor development and their significant roles in

tumor transformation and progression. Furthermore, the hypoxia pathway and angiogenesis pathway

are well-documented to contribute to the formation of an immunosuppressive TME in HCC[19].

Hypoxia directly induces VEGF production[20-21], which can exert inhibitory effects on various immune

cells, such as CD8+ T cells[22]. Additionally, VEGF-induced angiogenesis within the TME exacerbates

hypoxia, leading to a detrimental feedback loop[23]. These factors likely contribute to the observed

heterogeneity and prognostic differences between the two subtypes. Finally, we conducted an in-depth

assessment of the impact of subtyping on clinical treatment and prognosis assessment for patients. Our

findings revealed that CS2 exhibited enhanced sensitivity to the conventional drug 5-fluorouracil

utilized in hepatic arterial infusion chemotherapy, as well as to several tyrosine kinase inhibitors,

including gefitinib, afatinib, and erlotinib, compared to CS1. This discovery holds the potential to

inform tailored treatment strategies for patients based on their subtype, consequently improving

treatment efficacy and prognosis.

The above findings suggest that our classification method reliably distinguishes patients into two

subtypes characterized by significant prognostic differences. These prognostic disparities may arise

from heterogeneity linked to variations in metabolic reprogramming and signaling patterns, thus

providing a theoretical foundation for the classification's validity. Consequently, we developed a

prognostic prediction model based on subtype characteristics to enhance the clinical applicability of the

classification and effectively predict patient prognosis. Machine learning algorithms serve as powerful

tools for analyzing multi-omics data and constructing prognostic models. We utilized the TCGA-LIHC

cohort as the training set and the ICGC-JP cohort as the validation set. After testing 101 algorithm

combinations to identify the optimal model, we employed the C-index to evaluate predictive

performance, ultimately determining the model with the highest C-index, which signifies superior

predictive capability.

The data revealed that the prognostic model, developed using the Random Survival Forest (RSF)

algorithm and incorporating 5 hub genes, demonstrated superior performance. Upon assigning risk

scores and stratifying patients in both the training and validation sets based on this model, it became

evident that individuals in the high-risk category exhibited significantly poorer prognoses compared to

those in the low-risk category, affirming the success of our model construction. Subsequently, we

compared our model with several previously established prognostic prediction models to predict patient

outcomes and assessed their performance in both cohorts. Through comprehensive evaluations, our

model consistently outperformed the other models across both sets. Additionally, we conducted further

assessments using calibration curves and decision curve analysis, yielding consistent results indicating

the robust predictive capabilities of our model for patient prognosis.

After exploring the clinical implications of our classification method, we conducted an analysis of

tumor subtype heterogeneity at the single-cell level to uncover potential mechanisms contributing to

differences in prognosis and heterogeneity between subtypes. Initially, we annotated a single-cell

sequencing dataset comprising tumor tissue samples from 31 HCC patients, effectively categorizing

cells into 12 distinct groups. Subsequently, leveraging the CopyKAT algorithm, we analyzed gene

expression levels to identify non-diploid cells as malignant. Employing the same classification method

used previously, we successfully categorized malignant cells into two subtypes and assessed

differential gene expression and metabolic pathways between them. Our findings aligned with

RNA-bulk sequencing data, highlighting significant distinctions between subtypes across various

tumor-related signaling pathways and metabolic processes.

Our findings indicate significant heterogeneity among tumor cells of the two subtypes. Additionally, we

conducted an analysis of signaling patterns between tumor cells and other cell groups. The results

revealed that CS1 malignant cells exert substantial regulatory effects on immune cells within the TME,

including CD8+ T cells, macrophages, B cells, and neutrophils, mediated by the MIF-(CD74+CXCR4)

pathway. Prior research has elucidated that MIF binds to its receptor CD74, leading to its

phosphorylation and the recruitment of CD44, triggering downstream phosphorylation signals via the

proto-oncogene SRC[24-25]. This cascade encompasses the transcriptional activation of cyclin D1,

phosphorylation of retinoblastoma protein, and the involvement of multiple pro-proliferation and

anti-apoptotic pathways such as the MAPK/ERK and PI3K/AKT pathways[26-27].

The MIF signaling pathway is intricately involved in diverse cellular processes. It stabilizes the

p53-MDM2 complex, promoting cell proliferation and inhibiting apoptosis[28]. There is a synergistic

relationship between MIF and HIF-1α, where HIF-1α binds to the 5'-UTR hypoxia response element of

MIF, thereby enhancing MIF expression regulated by CREB[29]. Additionally, MIF activates a

p53-dependent feedback loop of HIF-1α, crucial for tumor hypoxia adaptation[30]. Moreover, MIF

prevents proteasome-mediated degradation of HIF-1α and forms a ternary complex with CSN5 and

HIF-1α, amplifying the tumor hypoxia response[31]. Our prior findings indicate heightened activity in

MAPK, PI3K-AKT signaling, and the hypoxia pathway in the CS1 subtype compared to CS2.

Integrating RNA-bulk and single-cell sequencing data suggests a pivotal role for the MIF pathway in

subtype heterogeneity.

In comparison to prior studies, our research exhibits several distinctive characteristics. First, we

acknowledged the potential substantial heterogeneity in HCC patients, shaping our research perspective

towards offering more precise stratification and treatment for patients. Second, we incorporated 5 types

of omics data and employed 10 clustering methods for analysis, ultimately amalgamating them to

derive a consensus clustering method, thereby mitigating the impact of clustering method selection bias

on our analysis. Third, our modeling was underpinned by the features of consensus clustering subtypes,

utilizing 10 machine learning algorithms combined into 101 algorithmic configurations for modeling.

We identified the model with the best average C-index performance, surpassing several previously

reported prediction models in terms of predictive efficacy. Fourth, subsequent to identifying potential

heterogeneity between subtypes in RNA-bulk data analysis, we further delved into single-cell data. Our

exploration revealed the involvement of the MIF signaling pathway in tumor heterogeneity and the

resulting prognostic disparities in HCC patients. This comprehensive approach allowed us to conduct

an extensive study around the consensus clustering method, elucidating both its clinical application

value and the mechanisms underpinning the observed heterogeneity and prognostic differences

between subtypes.

Conclusion

In summary, our study is based on a new multi-omics clustering consensus typing method. While

discovering the prognostic value of this typing method through the analysis of RNA-bulk data and

scRNA-seq data, we focused on the tumor heterogeneity caused by metabolic reprogramming and

changes in signaling transduction patterns. We initially explored the reasons for the differences in

prognosis among patients of different types and found that the MIF signaling pathway is involved in

the occurrence and development of HCC. Due to the use of bioinformatics and limited samples and

survival data in our study, it is necessary to further validate potential mechanisms through subsequent

basic experiments and analysis of larger cohort data.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under grant 82272902 and the Beijing Public Health Foudation under grant 2023158.

Data Availability Statement

The data analyzed in this study were obtained from The Cancer Genome Atlas(TCGA) at <u>TCGA-LIHC</u>, International Cancer Genome Consortium(ICGC) at <u>LIHC-JP</u>, and Gene Expression Omnibus (GEO) at <u>GSE229772</u>.

References

[1] Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, Jemal A. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin. 2024;74(3):229-63.

[2] Siegel RL, Giaquinto AN, Jemal A. Cancer statistics, 2024. CA Cancer J Clin. 2024;74(1):12-49.

[3] Zheng H, Pomyen Y, Hernandez MO, Li C, Livak F, Tang W, Dang H, Greten TF, Davis JL, Zhao Y, Mehta M, Levin Y, Shetty J, Tran B, Budhu A, Wang XW. Single-cell analysis reveals cancer stem cell heterogeneity in hepatocellular carcinoma. Hepatology. 2018;68(1):127-40.

[4] Colaprico A, Silva TC, Olsen C, Garofano L, Cava C, Garolini D, Sabedot TS, Malta TM, Pagnotta SM, Castiglioni I, Ceccarelli M, Bontempi G, Noushmehr H. TCGAbiolinks: an R/Bioconductor package for integrative analysis of TCGA data. Nucleic Acids Res. 2016;44(8):e71.

[5] Mayakonda A, Lin DC, Assenov Y, Plass C, Koeffler HP. Maftools: efficient and comprehensive analysis of somatic variants in cancer. Genome Res. 2018;28(11):1747-56.

[6] Lu X, Meng J, Zhou Y, Jiang L, Yan F. MOVICS: an R package for multi-omics integration and visualization in cancer subtyping. Bioinformatics. 2021;36(22-23):5539-41.

[7] Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.

2015;43(7):e47.

[8] Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, Fu X, Liu S, Bo X,

Yu G. clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. Innovation

(Camb). 2021;2(3):100141.

[9] Charoentong P, Finotello F, Angelova M, Mayer C, Efremova M, Rieder D, Hackl H, Trajanoski Z.

Pan-cancer Immunogenomic Analyses Reveal Genotype-Immunophenotype Relationships and

Predictors of Response to Checkpoint Blockade. Cell Rep. 2017;18(1):248-62.

[10] Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and

RNA-seq data. BMC Bioinformatics. 2013;14:7.

[11] Liu Z, Liu L, Weng S, Guo C, Dang Q, Xu H, Wang L, Lu T, Zhang Y, Sun Z, Han X. Machine

learning-based integration develops an immune-derived lncRNA signature for improving

outcomes in colorectal cancer. Nat Commun. 2022;13(1):816.

[12] Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G,

Madad S, Fernandez-Granda C, Satija R. Dictionary learning for integrative, multimodal and

scalable single-cell analysis. Nat Biotechnol. 2024;42(2):293-304.

[13] Fan C, Chen F, Chen Y, Huang L, Wang M, Liu Y, Wang Y, Guo H, Zheng N, Liu Y, Wang H, Ma

L. irGSEA: the integration of single-cell rank-based gene set enrichment analysis. Brief Bioinform.

2024;25(4).

[14] Wu Y, Yang S, Ma J, Chen Z, Song G, Rao D, Cheng Y, Huang S, Liu Y, Jiang S, Liu J, Huang X,

Wang X, Qiu S, Xu J, Xi R, Bai F, Zhou J, Fan J, Zhang X, Gao Q. Spatiotemporal Immune

Landscape of Colorectal Cancer Liver Metastasis at Single-Cell Level. Cancer Discov.

2022;12(1):134-53.

[15] Jin S, Plikus MV, Nie Q. CellChat for systematic analysis of cell-cell communication from

single-cell and spatially resolved transcriptomics. J bioRxiv. 2023.11.05.565674.

[16] Gao R, Bai S, Henderson YC, Lin Y, Schalck A, Yan Y, Kumar T, Hu M, Sei E, Davis A, Wang F,

Shaitelman SF, Wang JR, Chen K, Moulder S, Lai SY, Navin NE. Delineating copy number and

clonal substructure in human tumors from single-cell transcriptomes. Nat Biotechnol.

2021;39(5):599-608.

[17] Martínez-Reyes I, Chandel NS. Cancer metabolism: looking forward. Nat Rev Cancer.

2021;21(10):669-80.

[18] Barkley D, Moncada R, Pour M, Liberman DA, Dryg I, Werba G, Wang W, Baron M, Rao A, Xia

B, França GS, Weil A, Delair DF, Hajdu C, Lund AW, Osman I, Yanai I. Cancer cell states recur

across tumor types and form specific interactions with the tumor microenvironment. Nat Genet.

2022;54(8):1192-201.

[19] Kudo M. Combination immunotherapy with anti-VEGF/TKI for hepatocellular carcinoma: present

and future perspective. Hepatobiliary Surg Nutr. 2021;10(2):241-5.

[20] Li Y, Zhang MZ, Zhang SJ, Sun X, Zhou C, Li J, Liu J, Feng J, Lu SY, Pei-Jun L, Wang JC.

HIF-1α inhibitor YC-1 suppresses triple-negative breast cancer growth and angiogenesis by

targeting PlGF/VEGFR1-induced macrophage polarization. Biomed Pharmacother.

2023;161:114423.

[21] Du R, Lu KV, Petritsch C, Liu P, Ganss R, Passegué E, Song H, Vandenberg S, Johnson RS, Werb

Z, Bergers G. HIF1alpha induces the recruitment of bone marrow-derived vascular modulatory

cells to regulate tumor angiogenesis and invasion. Cancer Cell. 2008;13(3):206-20.

[22] Oura K, Morishita A, Tani J, Masaki T. Tumor Immune Microenvironment and

Immunosuppressive Therapy in Hepatocellular Carcinoma: A Review. Int J Mol Sci. 2021;22(11).

[23] Shigeta K, Datta M, Hato T, Kitahara S, Chen IX, Matsui A, Kikuchi H, Mamessier E, Aoki S,

Ramjiawan RR, Ochiai H, Bardeesy N, Huang P, Cobbold M, Zhu AX, Jain RK, Duda DG. Dual

Programmed Death Receptor-1 and Vascular Endothelial Growth Factor Receptor-2 Blockade

Promotes Vascular Normalization and Enhances Antitumor Immune Responses in Hepatocellular

Carcinoma. Hepatology. 2020;71(4):1247-61.

[24] Bozzi F, Mogavero A, Varinelli L, Belfiore A, Manenti G, Caccia C, Volpi CC, Beznoussenko GV,

Milione M, Leoni V, Gloghini A, Mironov AA, Leo E, Pilotti S, Pierotti MA, Bongarzone I,

Gariboldi M. MIF/CD74 axis is a target for novel therapies in colon carcinomatosis. J Exp Clin

Cancer Res. 2017;36(1):16.

[25] Gil-Yarom N, Radomir L, Sever L, Kramer MP, Lewinsky H, Bornstein C, Blecher-Gonen R,

Barnett-Itzhaki Z, Mirkin V, Friedlander G, Shvidel L, Herishanu Y, Lolis EJ, Becker-Herman S,

Amit I, Shachar I. CD74 is a novel transcription regulator. Proc Natl Acad Sci U S A.

2017;114(3):562-7.

[26] Penticuff JC, Woolbright BL, Sielecki TM, Weir SJ, Taylor JA, 3rd. MIF family proteins in

577 genitourinary cancer: tumorigenic roles and therapeutic potential. Nat Rev Urol.

578 2019;16(5):318-28.

579 [27] Sumaiya K, Langford D, Natarajaseenivasan K, Shanmughapriya S. Macrophage migration

580 inhibitory factor (MIF): A multifaceted cytokine regulated by genetic and physiological strategies.

581 Pharmacol Ther. 2022;233:108024.

582 [28] Fukaya R, Ohta S, Yaguchi T, Matsuzaki Y, Sugihara E, Okano H, Saya H, Kawakami Y, Kawase

583 T, Yoshida K, Toda M. MIF Maintains the Tumorigenic Capacity of Brain Tumor-Initiating Cells

584 by Directly Inhibiting p53. Cancer Res. 2016;76(9):2813-23.

585 [29] Safi W, Kraus A, Grampp S, Schödel J, Buchholz B. Macrophage migration inhibitory factor is

586 regulated by HIF-1α and cAMP and promotes renal cyst cell proliferation in a

587 macrophage-independent manner. J Mol Med (Berl). 2020;98(11):1547-59.

588 [30] Thirusangu P, Vigneshwaran V, Prashanth T, Vijay Avin BR, Malojirao VH, Rakesh H, Khanum

589 SA, Mahmood R, Prabhakar BT. BP-1T, an antiangiogenic benzophenone-thiazole pharmacophore,

590 counteracts HIF-1 signalling through p53/MDM2-mediated HIF-1α proteasomal degradation.

591 Angiogenesis. 2017;20(1):55-71.

592 [31] Winner M, Koong AC, Rendon BE, Zundel W, Mitchell RA. Amplification of tumor hypoxic

593 responses by macrophage migration inhibitory factor-dependent hypoxia-inducible factor

594 stabilization. Cancer Res. 2007;67(1):186-93.

FIGURE 1. Multi-omics consensus clustering classification. (A) Calculation of optimal number of subtypes based on cluster prediction index and gap-statistics. (B) Heatmap of the consistency of the integration of the results of the 10 typing algorithms. (C-D) Significant multiomics characterization and prognostic differences between subtypes. (E-G) Validation of consistency of typing results and prognostic value in an external cohort.

FIGURE.2 Differences in molecular characterization of subtypes. (A) Differences in mutation landscapes of subtypes. (B-C) Genomic characterization from TMB and FGA of subtypes. (D-E) Functional enrichment of up and down-regulated differential genes between subtypes. (F) Concordance between subtypes and patients' pathologic staging.

FIGURE 3. Phenotypic features of subtypes that may be associated with tumor development. (A) Landscape of subtype TME immune infiltration. (B) Differences in subtype responsiveness to different drugs. (C) Comparison of immune cell infiltration levels in subtypes of TME. (D) Comparison of hepatocellular carcinoma-related signaling pathways and biological process activity in subtypes.

FIGURE 4. Building prognostic prediction models through machine learning. (A) 101 Machine Learning Algorithms to Build Screening Optimal Prognostic Models. (B) Calculating risk scores and comparing prognostic differences based on predictive modeling. (C-D) Correlation between 5 prognostic genes and prognosis. (E-F) Validating prognostic model efficacy in an external cohort external cohort.

FIGURE 5. Comparison and refinement of prognostic prediction models. (A) Comparison of predictive performance of prognostic models with some prognostic models published in the last 5 years. (B) Constructing a nomogram incorporating patient clinical characteristics. (C-D) Evaluating model performance based on calibration curve and decision curve analysis.

FIGURE 6. ScRNA-seq cell sorting and annotation. (A-C) Cellular subpopulation delineation and annotation based on cellular characterization genes. (D-E) Visualization of the proportions of each cell subgroup. (F) Differentially expressed genes by cellular subpopulations.

FIGURE 7. Phenotypic characterization of cellular subpopulations. (A-B) Functional enrichment analysis of differentially expressed genes within cell subpopulation. (C) Analysis of the metabolic landscape of cell subpopulations. (D) Cellular polymeric communication network. (E) The contribution of signal strength in both incoming and outgoing signals shapes the landscape of cellular subgroups. (F-G) Analysis of signal transmission into and out of cellular subgroups.

FIGURE 8. Exploring the characteristics of two subtypes of malignant cells at the scRNA-seq level. (A) Mapping of malignant cells in the cell clustering results. (B) Enrichment analysis of differential genes between different subtypes of malignant cells. (C) Metabolic level analysis of different subtypes of malignant cells. (D) Annotation of the cellular subpopulation signaling landscape after malignant cell annotation. (E-G) Analysis of signal input-output patterns in cellular subpopulations. (H) Differential levels of the MIF signaling pathway in cellular subpopulation outputs. (I) Signaling modulation pattern of CS1 subtype malignant cells on other cells.