Weberviruses are gut-associated phages that infect *Klebsiella* spp.

Samuel J. T. Dawson[1], Preetha Shibu[2]*, Sara Garnett[1], Fiona Newberry[1], Thomas C.

Brook[2], Tobi Tijani[2], Magdalena Kujawska[3,4,5], Lindsay J. Hall[3,4,5], Anne L. McCartney[1],

David Negus[1], Lesley Hoyles[1]

[1] Department of Biosciences, Nottingham Trent University, UK

[2] Life Sciences, University of Westminster, UK

[3] Microbes, Infection and Microbiomes, School of Infection, Inflammation and Immunology,

College of Medicine and Health, University of Birmingham, UK

[4] Food, Microbiome and Health, Quadram Institute Bioscience, Norwich Research Park,

Norwich, UK

[5] Norwich Medical School, University of East Anglia, Norwich Research Park, Norwich

Corresponding authors: David Negus, david.negus@ntu.ac.uk; Lesley Hoyles,

lesley.hoyles@ntu.ac.uk

* **Present address:** Berkshire and Surrey Pathology Services, Frimley Health NHS Trust,

Wexham Park Hospital, Slough, UK.

Keywords: microbiota, comparative genomics, environment, metagenome-assembled

genomes, *Klebsiella pneumoniae*.

Abbreviations: MAG, metagenome-assembled genome; MDR, multidrug-resistant; ST,

sequence type; TEM, transmission electron micrograph; UHGG, Unified Human

Gastrointestinal Genome.

Data availability. The sequences for the seven new phage genomes described herein have

been deposited in DDBJ/ENA/GenBank under accession numbers OM065837–OM065843.

The genome sequences of bacteria described herein have been deposited under BioProject

PRJNA917129. All supplementary material is available from

https://figshare.com/projects/Weberviruses_are_gut-

associated_phages_that_infect_Klebsiella_spp_/128516.

ABSTRACT

Weberviruses are bacteriophages (phages) that have been shown to infect and lyse members of the genus *Klebsiella*. Due to their antimicrobial properties, phages and their products have emerged as a potential therapeutic option to help tackle increasing rates of antimicrobial resistance amongst *Klebsiella* spp. Weberviruses are an attractive option due to their high burst sizes, long shelf life and associated depolymerase enzymes. In this study we isolated and characterized seven new weberviruses and compared their genomes with those of their closest relatives to further our understanding of the genus *Webervirus*. We show that the seven phages isolated in this study can lyse a range of clinical multidrug-resistant *Klebsiella* strains, including *K. variicola*. Gene-sharing network-based analysis using vConTACT2 and incorporating all publicly available webervirus genomes (*n*=67) confirmed the seven phages as members of the genus *Webervirus* and that weberviruses are affiliated with the family *Drexlerviridae*. Using the online resource PhageClouds, we demonstrated that weberviruses can be readily detected in metagenome-derived phage datasets from a diverse range of geographical locations. Using a curated database of 67 isolated phage genomes and 60 metagenome-assembled genomes (*n*=127 total), we found that weberviruses are primarily associated with samples originating from the gut, with human faeces (61/127, 48 %), sewage (39/127, 31 %) and wastewater (14/127, 11 %) being the most frequent sources of these phages.

INTRODUCTION

Webervirus KP36 was isolated from treated sewage from a communal wastewater plant in Poland (Kęsik-Szeloch et al., 2013). This lytic phage infected ~12 % of clinical *Klebsiella pneumoniae* strains it was screened against, and exhibited depolymerase activity on its hosts (Kęsik-Szeloch et al., 2013; Majkowska-Skrobek et al., 2016). It represented the type species of a novel genus, *KP36virus*, of *Siphoviridae* belonging to the family *Tunavirinae* (Kęsik-Szeloch et al., 2013). Subsequent changes to the virus taxonomic code led to the renaming of *KP36virus* to *Webervirus* (Walker et al., 2019), with the genus *Webervirus* moved to the family *Drexliviridae* within the order *Caudovirales* in 2020 (Walker et al., 2021). Abolishment of the *Caudovirales* in July 2021 led to the family *Drexliviridae* being classified within the class *Caudoviricetes*, phylum *Uroviricota*. As of 18 January 2023, the genus *Webervirus* encompasses 32 different phage species ([International Committee on Taxonomy of Viruses](#)).

All weberviruses described to date have been isolated on *Klebsiella* hosts, and have proven easy to recover from sewage, wastewater and, occasionally, intestinal contents (Herridge et al., 2020). Although *Webervirus F20* was originally described as being isolated on *Enterobacter aerogenes* (Mishra et al., 2012), this bacterium has subsequently been reclassified as *Klebsiella aerogenes*. Their high burst sizes and long shelf life make weberviruses ideal phages to work with for biotechnological and clinical applications (Fang et al., 2022; Herridge et al., 2020). Weberviruses tend to have narrow host ranges (Hoyles et al., 2015; Pertics et al., 2021). However, our previous (and ongoing) work has suggested that their depolymerases can target the capsules of non-host *Klebsiella* spp. (Hoyles et al., 2015). Depolymerase activity is common to weberviruses, and is being actively investigated as a tool to lyse capsules of *Klebsiella* spp. that often hinder or make treatment with antimicrobials difficult (Cai et al., 2019; Majkowska-Skrobek et al., 2016; Pertics et al., 2021). For example, the webervirus depolymerase Depo32 has been shown to protect mice from otherwise lethal *K. pneumoniae* infections in a mouse model of disease (Cai et al., 2023). In addition, a webervirus (P39) has recently been used in combination with another lytic phage (P24, *Przondovirus*) to decolonize mice of carbapenem-resistant *K. pneumoniae* (Fang et al., 2022).

Previously we isolated *Webervirus KLPN1* from the caecum of a healthy female, along with its host *K. pneumoniae* subsp. *pneumoniae* L4-FAA5 (Hoyles et al., 2015). During studies to isolate additional phages that infect L4-FAA5 and multidrug-resistant (MDR) *Klebsiella* spp., we identified seven new representatives of the genus *Webervirus*. Here we describe these

new phages and compare their genomes with those of their closest relatives. We also

demonstrate that weberviruses are readily detectable within recent metagenome-derived

phage datasets.

METHODS

Bacterial strains. Details of all *Klebsiella* strains included in this study are given in **Table 1**.

The antimicrobial resistance profiles of the isolates, determined according to EUCAST

guidelines as described previously (Shibu et al., 2021), can be found in **Supplementary**

Table 1.

Generation of sequence data for bacterial isolates. Genome data for clinical strains

included in this study were generated as described previously (Shibu et al., 2021). Illumina

and ONT Nanopore data for *K. pneumoniae* L4-FAA5 were generated by microbesNG

(Birmingham, UK) as described previously (Newberry et al., 2023). *De novo* genome

assembly was performed by microbesNG using Unicycler v0.4.0 (Wick et al., 2017).

CheckM2 v0.1.3 (Chklovski et al., 2022) was used to confirm the quality (in terms of

completeness and contamination) of all assembled genomes. PubMLST was used to assign

sequence types (STs) to genomes (Jolley et al., 2018); Kaptive was used to assign capsule

(K) and lipopolysaccharide (O) locus types to genomes (Wick et al., 2018).

Sample collection and processing. Sewage samples were collected from different mixed-

liquor tanks at Mogden Sewage Treatment Works (Thames Water). All sewage samples

were filtered-sterilized with 0.45 µm cellulose acetate filters (Millipore), and kept at 4 °C prior

to use.

Isolation and propagation of phages. Filter-sterilized sewage samples were screened for

phages using *Klebsiella* strains L4-FAA5, PS_misc5 and PS_misc6 (**Table 1**). Aliquots (9

ml) of each sample were inoculated into 1 ml of 10× sterile nutrient broth (Fluka) containing

50 mM $CaCl_2$ and 50 mM $MgCl_2$ and 200 µl of overnight broth culture, and incubated at 37

°C for 6–16 h. Phage-positive enrichments showed lysis (clear appearance) of the culture

after incubation. Dilution series (10^{-1} to 10^{-10}) were prepared in nutrient broth from the filtered

lysates of the phage-positive samples and used in double-agar overlays to obtain individual

plaques as described previously (Hoyles et al., 2015).

Isolation of phage DNA

vB_KpnS-KLPN2, vB_KpnS-KLPN3 and vB_KpnS-KLPN4. Phages were precipitated from

100 ml of each lysate using PEG 8000/6 % NaCl and DNA isolated from the three phages as

described previously (Hoyles et al., 2015). Integrity and quality of DNA were checked using

agarose gel electrophoresis and Nanodrop.

vB_KvaS-KLPN5, vB_KvaS-KLPN6, vB_KvaS-KLPN7 and vB_KpnS-KLPN8. These phages

were concentrated and DNA extracted as described previously (Smith-Zaitlik et al., 2022).

Transmission electron microscopy. Transmission electron micrographs (TEMs) of phages

vB_KpnS-KLPN2, vB_KpnS-KLPN3 and vB_KpnS-KLPN4 were generated as described

previously (Hoyles et al., 2015). TEMs for phages vB_KvaS-KLPN5, vB_KvaS-KLPN6,

vB_KvaS-KLPN7 and vB_KpnS-KLPN8 were generated as described previously (Smith-

Zaitlik et al., 2022), with images analysed using ImageJ (Schindelin et al., 2012).

Phage genome sequencing, assembly and annotation. Illumina sequence data for

phages vB_KpnS-KLPN2, vB_KpnS-KLPN3 and vB_KpnS-KLPN4 were generated by

microbesNG in 2016. Data quality for the provided trimmed sequence data were checked

using FastQC v0.11.9 and did not need further trimming. For phages vB_KvaS-KLPN5,

vB_KvaS-KLPN6, vB_KvaS-KLPN7 and vB_KpnS-KLPN8, sequence data were generated

on an Illumina MiSeq at Nottingham Trent University according to (Smith-Zaitlik et al., 2022)

(Illumina MiSeq Reagent Kit v2; 150-bp paired-end reads). Quality of raw sequence data

was assessed using FastQC v0.11.9. Reads had a mean phred score above 30 and no

adapter contamination, so data were not trimmed.

All genomes were assembled using SPAdes v3.13.0 (default settings) (Bankevich et al.,

2012), and visualized to confirm circularization of genomes using Bandage v0.8.1 (Wick et

al., 2015). CheckV v0.8.1 (checkv-db-v1.0; (Nayfach et al., 2021a)) was used to determine

contamination and completeness of the genomes.

Genes in all phage genomes included in this study (**Table 2**) were predicted using the

PHROG (Terzian et al., 2021) hmm dataset and Prokka v1.14.6 (Seemann, 2014).

Comparison of webervirus genomes. ViPTree v1.9.1 (Nishimura et al., 2017) was used to

determine whether the seven phage genomes were closely related to previously described

double-stranded DNA viruses. Initial analyses showed them to be closely related to

Webervirus KP36 (Kęsik-Szeloch et al., 2013) and other weberviruses. As the online

ViPTree database does not contain genomes of all members of the genus *Webervirus*, we

curated a list of all known webervirus sequences within NCBI GenBank in April 2022 (**Table**

2). The sequence files were concatenated and uploaded to VIRIDIC web

(http://rhea.icbm.uni-oldenburg.de/VIRIDIC/) (Moraru et al., 2020). They were also added to

the vConTACT v2.0 database (Bin Jang et al., 2019; Bolduc et al., 2017). The similarity

matrix created using VIRIDIC was visualized using heatmap.2 in gplots v3.1.1. The gene-

sharing network created using vConTACT v2.0 was imported into Gephi v0.9.2 (Bastian et

al., 2009) for visualization.

Large-subunit terminase genes were extracted from genomes using Biostrings v2.62.0

(Pagès et al., 2021). A multiple-sequence alignment was created for the genes (Clustal

Omega 1.2.2 implemented in Geneious Prime v2020.0.5; options – group sequences by

similarity, 5 representative iterations). This alignment was used to create a bootstrapped

maximum-likelihood tree. Phangorn v2.7.1 and Ape v5.5 (Paradis and Schliep, 2019;

Schliep et al., 2017) were used to generate the trees using the LG model (Le and Gascuel,

2008).

Identification of weberviruses in metagenomic datasets. We used PhageClouds

(Rangel-Pineros et al., 2021) to identify relatives of weberviruses in metagenome-assembled

genome (MAG) datasets. The genome of *Webervirus KLPN1* was searched against the

PhageClouds database with a threshold of 0.15, as we had previously looked for relatives of

this phage in metagenomic datasets and are interested in gut-associated phage

communities (Hoyles et al., 2015). The nucleotide sequences of the relevant phage MAGs

(from (Camarillo-Guerrero et al., 2021; Gregory et al., 2020; Tisza and Buck, 2021)) were

recovered from the relevant datasets. CheckV v0.8.1 (checkv-db-v1.0; (Nayfach et al.,

2021a)) was used to determine contamination and completeness of our curated dataset;

those genomes that were complete and not already included in checkv-db-v1.0 were added

to the database (**Supplementary Table 2**), which was then used to determine the quality

(completeness, contamination) of the MAGs identified as being closely related to *Webervirus*

KLPN1.

The MAG sequences were analysed using ViPTree v2.0 (online) to confirm their affiliation

with the genus *Webervirus*. They were also included in a vConTACT analysis with our

curated set of 67 webervirus genomes.

Distribution of weberviruses. The global distribution of weberviruses was determined by

identifying the source and location information for the 67 GenBank genomes (**Table 2**) and

the 60 MAGs (**Supplementary Table 3**). Data were aggregated based on isolation source or

geographical location, with these latter data visualized using the R package rworldmap
v1.3.6 (South, 2011).
203
Host prediction for MAGs. The CRISPR Spacer Database and Exploration Tool (Dion et
al., 2021) and HostPhinder 1.1 (Villarroel et al., 2016) were used to predict hosts for the
weberviruses recovered from metagenomic datasets. The MAGs were also subject to a
BLASTN search against the Unified Human Gastrointestinal Genome (UHGG) CRISPR
spacer database according to (Nayfach et al., 2021b). For this, a BLASTN database was
created from 1,846,441 spacers from 145,053 CRISPR arrays from 79,735 UHGG genomes
(Nayfach et al., 2021b). Spacers were searched against viral genomes using BLASTN from
the blast+ package v.2.12.0 (options:-dust=no; -word-size=18); a maximum of one mismatch
or gap was allowed over ≥95 % of the spacer length.
213
214 **Identification of potential depolymerases among weberviruses**. A BLASTP database
215 was created using amino acid sequences from experimentally validated *Webervirus*
216 depolymerases and a BLASTP search was ran versus all webervirus genomes. Sequences
217 used to build the BLAST database can be found in **Supplementary Table 4**. MUSCLE
218 alignments were created in Geneious prime (v2023.1.2).
219
220
RESULTS
Initial characterization of the seven new phages
Seven phages were isolated on two different strains of *K. pneumoniae* subsp. *pneumoniae*
(L4-FAA5 – vB_KpnS-KLPN2, vB_KpnS-KLPN3, vB_KpnS-KLPN4; PS_misc6 – vB_KpnS-
KLPN8) and one strain of *K. variicola* subsp. *variicola* (PS_misc5 – vB_KvaS-KLPN5,
vB_KvaS-KLPN6, vB_KvaS-KLPN7). Strain L4-FAA5 (K2:O1v1, ST380) was originally
isolated from human caecal effluent along with *Webervirus KLPN1* (Hoyles et al., 2015),
while strains PS_misc5 (K81:OL101; gapA-16, infB-24, mdh-21, pgi-31, phoE-47, rpoB-22,
tonB-281) and PS_misc6 (K110:O2v1, ST716) were part of a collection (*n*=36) of clinical
MDR and/or carbapenem-resistant *Klebsiella* isolates currently being used in our laboratory
in phage-related and other studies (Shibu, 2019) (**Supplementary Table 1**).
232
233 TEM showed the seven phages had a siphovirus-type morphology (**Supplementary Figure
234 A**). Host-range analysis showed the seven phages had different infection profiles (**Table 1**).
235
Genome-based analyses of the seven new phages

Whole-genome sequences were generated for the seven phages. Bandage (data not
shown) and CheckV (**Supplementary Table 2**) analyses confirmed all the genomes were
circular and complete. None of the genomes was contaminated. An initial online ViPTree
analysis showed the new genomes belonged to the genus *Webervirus* (data not shown). All
publicly available webervirus genomes (available as of April 2022) were downloaded from
GenBank to allow comparison with our newly sequenced phages.
243
A gene-sharing network was created with 67 webervirus genomes derived from isolated
phages (**Table 2**) using vConTACT v2.0 (Bin Jang et al., 2019; Bolduc et al., 2017). The
network had 2,619 nodes and 51,589 edges. It was filtered based on the giant component
topology, removing all nodes that were not connected to the main cluster (**Figure 1**). This
left 1,523 nodes (58.15 % of 2,619) and 33,573 edges (65.08 % of 51,589) visible. The
vConTACT-based analysis confirmed the affiliation of the seven phages with the family
Drexlerviridae (**Figure 1**). The monophyletic nature of the genus *Webervirus* – including the
seven new phages – was confirmed by analysis of large-subunit terminase genes (**Figure
2**).
253
254 VIRIDIC (**Supplementary Figure B**) data suggested there were three species within the
255 genus *Webervirus*, based on the clustering of the 67 genomes.
256
Detection of weberviruses in metagenomic datasets
PhageClouds is an online resource that allows researchers to search a reference dataset of
~640,000 phage genomes for phages with genomes related to query sequences. Of the 67
webervirus genomes in our curated database, 47 were included in the PhageClouds output.
Five PhageClouds hits with the prefix PIGEON (Ter Horst et al., 2021) represented phages
already included in our curated dataset (**Supplementary Table 5**), so were excluded from
further analyses. Fifty-four of the PhageClouds hits represented MAGs derived from the Gut
Phage Database (GPD) (Camarillo-Guerrero et al., 2021), six were from the Cenote Human
Virome Database (CHVD) (Tisza and Buck, 2021) and two were from the Gut Virome
Database (GVD) (Gregory et al., 2020). MAG
Ma_2019_SRR413710_NODE_378_length_50715_cov_48.086538 from the GVD was
identical to uvig_330395 from the GPD (Camarillo-Guerrero et al., 2021) so was removed
from further analyses (PhageClouds scores identical, 100 % pairwise identity; an
unsurprising finding as both MAGs are derived from the same dataset (Ma et al., 2018)).
Two MAGs from the GPD were also found to be identical: uvig_314355 and uvig_315584
were high-quality genomes both derived from the same four samples (SRR1952259,
SRR1162648, SRR1162662, SRR1162654 (Tisza and Buck, 2021)); only uvig_314355 was

retained for further analyses. This left us with 60 MAGs (available as

60_MAG_sequences.fasta in **Supplementary Material**), potentially representing

weberviruses, that were subject to further analyses.

The MAGs ranged from 10,230 to 55,276 nt (mean 41,737 nt) in length (**Supplementary**

Table 6). Thirteen of our 67 curated genomes were determined to be complete (CheckV)

and of high quality (MIUVIG) by CheckV so were added to the CheckV database

(**Supplementary Table 2**). The updated database was then used to assess the quality

(completeness, contamination) of the 60 MAGs. Twenty-seven MAGs were complete/high

quality, 15 were high quality/high quality, eight were medium quality/genome fragments and

10 were low quality/genome fragments (**Supplementary Table 6**).

The 60 MAGs were uploaded to ViPTree v2.0 for analysis. In a comparison of 2,981

sequences, all MAGs clustered with the genus *Webervirus* (**Supplementary Material**).

vConTACT analysis confirmed the association of the MAGs with the genus *Webervirus*, with

only weberviruses represented in the corresponding module (**Figure 3a**).

Distribution and diversity of weberviruses

Our dataset represented a total of 67 isolated phages and 60 MAGs. Weberviruses were

recovered from a variety of sample types, with human faeces (61/127, 48 %), sewage

(39/127, 31 %) and wastewater (14/127, 11 %) the most frequent sources of these phages

(**Figure 3b**, **Supplementary Table 7**). Seven (5.5 %) of the weberviruses were of unknown

origin. Only one genome was recovered from each of the other sample types (activated

sludge, human caecal effluent, different water types).

Of the 127 different webervirus genomes included in this study (*n*=67 isolated phage, *n*=60

MAGs), 126 were of known geographical origin (**Figure 3c**). According to our analysis,

weberviruses have been detected most frequently in samples emanating from: China (*n*=54);

USA (*n*=24); UK (*n*=10); Spain (*n*=8); Fiji (*n*=5); Denmark and Russia (*n*=4 each); South

Korea (*n*=3); India, Hungary and Austria (*n*=2 each); Turkey, Thailand, Sweden, Singapore,

Poland, Pakistan, Cote d'Ivoire and Cameroon (*n*=1 each). The majority (108/127, 85 %) of

webervirus genomes appear to have been detected only once. However, for MAGs

described by the GPD (uvig prefix) (Camarillo-Guerrero et al., 2021), data were available

with respect to the number of metagenomic samples each had been recovered from: 11 had

been recovered from two samples each, one had been recovered from four samples, two

had been recovered from eight samples each, and five had been recovered from 16 samples

each (**Supplementary Table 3**). Recovery of webervirus genomes from one more than

sample was country-specific (i.e. for the five genomes recovered from 16 samples, all
derived from Fiji).

ViPTree analysis of all 127 genomes showed clustering of several phage genomes based on
geographical origin (**Supplementary Figure C**).

Phage–host prediction for the MAGs
Each of the seven new phage genomes (nt sequences) we generated was uploaded to
CRISPR Spacer Database and Exploration Tool (Dion et al., 2021). None of the phages
could be assigned to known hosts using this tool. Using the BLASTN approach of (Nayfach
et al., 2021b) with the MAG sequences, only SAMEA2737751_a1_ct5309 had sufficient
coverage; this MAG had two hits to *Klebsiella* species (*K. pneumoniae* and *K. variicola*).
HostPhinder 1.1 (Villarroel et al., 2016) was able to predict hosts for our KLPN phages, with
all assigned to *Klebsiella pneumoniae*. Consequently, this tool was used to predict hosts for
the 60 *Webervirus* MAGs (**Supplementary Table 8**). All were predicted to infect *Klebsiella*
pneumoniae.

Depolymerases of weberviruses
As our newly isolates phages all displayed apparent depolymerase activity against one or
more hosts, we aimed to identify potential depolymerases encoded within the genomes of
127 high-quality and complete webervirus genomes. Currently, four experimentally validated
depolymerases from weberviruses have been reported in the literature: depoKP36
(Majkowska-Skrobek et al., 2016), Depo32 (Cai et al., 2023), DpK2 (Dunstan et al., 2021)
and B1dep (Pertics et al., 2021). These four depolymerases were used to create a BLASTP
database to interrogate the 127 phage genomes for similar amino acid sequences.

In addition to the four previously reported sequences, our search identified four additional
potential depolymerases encoded within separate webervirus genomes, with two of these
belonging to metagenome-assembled genomes (**Supplementary Table 9**). All four
sequences shared >98 % amino acid identity and 100 % sequence coverage with the K2
capsule-degrading depolymerases Depo32, DpK2 and B1dep, but less than 64 % amino
acid over < 36 % query coverage with depolymerase depoKP36, an enzyme specific for the
K63 capsular type. An amino acid similarity matrix displaying percent amino acid identity
between all eight full-length sequences in included in (**Supplementary Table 10**). Alignment
of the depolymerase amino acid sequences highlighted a high level of N-terminal
conservation (**Supplementary Figure D**).

DISCUSSION

Members of the *Klebsiella pneumoniae* species complex are opportunistic pathogens that can cause serious hospital-acquired infections, and are major contributors to global deaths associated with antimicrobial resistance (Antimicrobial Resistance Collaborators, 2022). Carbapenem-resistant isolates of *K. pneumoniae* are resistant to a range of frontline β-lactam antibiotics. The difficulty of treating such isolates with conventional antibiotics has resulted in the investigation of new therapeutic modalities, including phages and their products. We have previously published an extensive review on *Klebsiella* phage and their potential as therapeutics (Herridge et al., 2020). To realize the potential of phage therapy, it is important to comprehensively characterize phages with clinical utility. Here, we describe the isolation and characterization of seven new phages belonging to the genus *Webervirus* with lytic activity against clinical MDR *Klebsiella* spp. and characterize these in the context of other available *Webervirus* genome sequences.

Isolation and characterization of seven new weberviruses

Studies from a diverse range of geographical locations have reported the isolation or detection of weberviruses from samples associated with the human gut (e.g. wastewater, sewage, faeces, caecal effluent) (Herridge et al., 2020). To date, the majority of weberviruses have been isolated using *K. pneumoniae* as a host. However, weberviruses have been reported to infect other *Klebsiella* spp. including *K. oxytoca* (Brown et al., 2017; Park et al., 2017) and *K. aerogenes* (Hudson et al., 2021). In the present study, we isolated seven new weberviruses from sewage samples, including three phages (vB_KvaS-KLPN5, vB_KvaS-KLPN6, vB_KvaS-KLPN7) that were isolated using a strain of *K. variicola* as the host. To our knowledge, this is the first report of weberviruses infecting *K. variicola*, a recognized emerging human pathogen (Rodríguez-Medina et al., 2019) increasingly associated with carbapenem and colistin resistance (Kim et al., 2023; Li et al., 2024). *K. variicola* PS_misc5 (this study) is a carbapenem-resistant clinical isolate that encodes the class D β-lactamase OXA-48. The remaining four phages were isolated on either the caecal isolate *K. pneumoniae* L4-FAA5 (vB_KpnS-KLPN2, vB_KpnS-KLPN3, vB_KpnS-KLPN4) or the clinical isolate *K. pneumoniae* PS_misc6 (vB_KpnS-KLPN8). Strain PS_misc6 encodes the carbapenem-degrading metallo-β-lactamase NDM, while L4-FAA5 is resistant to cefotaxime, temocillin, ceftoxitin and aztreonam. The lytic phages isolated as part of this study may, therefore, represent attractive future therapeutics for the treatment of drug-resistant isolates belonging to the *K. pneumoniae* species complex.

In agreement with previous work (Hoyles et al., 2015; Pertics et al., 2021), the weberviruses
described herein exhibited relatively narrow host ranges when screened against a panel of
Klebsiella (including 36 clinical MDR) isolates representing a range of STs and capsule (K)
types (**Table 1**, **Supplementary Table 1**). Phage host range is very much related to
isolation host rather than phage phylogeny, with lysis appearing to be restricted based on K
type. Phages *Webervirus KLPN1*, vB_KpnS-KLPN2, vB_KpnS-KLPN3 and vB_KpnS-KLPN4
had the broadest host ranges and were all isolated on the same host – *K. pneumoniae* L4-
FAA5 (K2). Based on our phylogenetic analysis of phage-encoded large-subunit terminase
amino acid sequences *Webervirus KLPN1* belongs to cluster 1 whereas vB_KpnS-KLPN2,
vB_KpnS-KLPN3 and vB_KpnS-KLPN4 belong to cluster 2 (**Figure 2**). Phages *Webervirus*
KLPN1, KpnS-KLPN2 and vB_KpnS-KLPN3 shared identical host ranges, showing lytic
activity against the same 15 strains from the collection of 37 isolates (phage vB_KpnS-
KLPN4 exhibited a near identical host range but lysed one strain fewer) (**Table 1**). All four
phages were restricted to hosts predicted to produce either the K2 (14/15 strains lysed) or
K64 (1/15 strains lysed) capsule. Phages vB_KvaS-KLPN5, vB_KvaS-KLPN6 and vB_KvaS-
KLPN7 were isolated on *K. variicola* PS_misc5 (K81) and shared identical, narrow host
ranges; infecting only a single isolate (PS_Kpn24; K2) other than their original host. Phages
vB_KvaS-KLPN5 and vB_KvaS-KLPN7 were identified as belonging to cluster 1, whereas
vB_KvaS-KLPN6 groups with cluster 2. Phage vB_KpnS-KLPN8 was restricted to lysis of its
original host strain, *K. pneumoniae* PS_misc6 and is a member of cluster 2 (**Figure 2**).
404
Capsular (K) type has been shown to be a major determinant of host tropism in *Klebsiella*
phages (Beamud et al., 2023). Bacterial capsules are known to prevent phage attachment
by masking cell-surface-associated receptor proteins (Dunstan et al., 2021; Scholl et al.,
2005). To overcome this physical barrier, phages encode enzymes – frequently referred to
as depolymerases – that selectively degrade polysaccharides or polypeptides that comprise
the bacterial capsule (Cai et al., 2023; Dunstan et al., 2021; Hoyles et al., 2015; Majkowska-
Skrobek et al., 2016; Pertics et al., 2021). Phage-encoded depolymerases, therefore,
contribute to host tropism and previous studies have identified that weberviruses encode
functionally active depolymerase enzymes (Cai et al., 2023; Dunstan et al., 2021;
Majkowska-Skrobek et al., 2016; Pertics et al., 2021). While performing our host-range
analysis, we observed the presence of haloes indicative of depolymerase activity for a small
number of phage–host combinations (haloes were was associated with 18 % of the plaques;
Table 1). We, therefore, undertook a bioinformatic analysis to identify potential
depolymerase enzymes encoded within webervirus genomes. Our BLASTP search identified
four potential depolymerases that shared significant sequence similarity with the webervirus
depolymerases Depo32, DpK2 and B1dep. These enzymes have been shown to selectively

degrade the *K. pneumoniae* K2 capsule (Cai et al., 2023; Dunstan et al., 2021; Pertics et al.,
2021). As the four sequences shared far less similarity with the third depolymerase included
in our BLAST database, a webervirus enzyme which hydrolyses the K63 capsule, the
identified depolymerases may have specificity towards the K2 capsule. All depolymerases
shared a high level of sequence identity at the N-terminal region (**Supplementary Figure**
D). The N-terminal region of phage-associated depolymerases is typically responsible for
anchoring the baseplate of the phage virion and is often highly conserved (Knecht et al.,
2019; Latka et al., 2019). Structural analysis of Depo32 from phage GH-K3 has revealed
that, in addition to the N-terminal domain, Depo32 contains a short neck helix and
connection domain (residues 186–271), a β-helix domain (residues 272–642), a connection
helix domain (residues 643–666), a carbohydrate-binding module (residues 667–846), and a
C-terminal domain (residues 847–907) (Cai et al., 2023). It is the β-helix domain that is
responsible for hydrolysis of the polysaccharide capsule. Given the high level of amino acid
identity between Depo32 and the four additional sequences identified, it is highly likely that
these potential depolymerases are structurally similar.
436
We were unable to identify any coding sequences in the genomes of our isolated KLPN
phages sharing high similarity to the three experimentally validated webervirus
depolymerase sequences used to create our BLASTP database. Thus, it is likely that any
depolymerase activity associated with the phages isolated in our study is due to enzyme(s)
that remain to be characterized experimentally. As part of our previous analysis of the
genome of phage KLPN1, we hypothesized that ORF34 and/or ORF35 may encode the
depolymerase activity of phage KLPN1 as these sequences include a predicted endo-*N*-
acetylneuraminidase/endosialidase domain (Hoyles et al., 2015). Further experimental work
is required to determine whether these are functionally active depolymerases. As most of the
plaques we observed had no discernible haloes, it may be that alternative mechanisms are
used by weberviruses for penetrating the bacterial capsule. Depolymerase-independent
penetration of the capsule by *Klebsiella* phages has been reported in the literature (Beamud
et al., 2023).
450
Genome-based analyses of weberviruses
Our gene-sharing network analysis of 67 webervirus genomes derived from isolated phages
confirmed their association with the family *Drexlerviridae* (**Figure 1**). Taxonomic assignment
of phages using whole genome gene-sharing profiles has been shown to be highly accurate;
a recent study showed that vConTACT2 produces near-identical replication of existing
genus-level viral taxonomy assignments from the International Committee on Taxonomy of
Viruses (ICTV) (Bin Jang et al., 2019). It has been suggested that genomes comprising a

genus should be evaluated by phylogenetics with the use of conserved 'signature genes'

that are conserved throughout all members (Turner et al., 2021). Such analyses should

always produce trees that are monophyletic. Using the large-subunit terminase as a

'signature gene', we were able to show that the genus *Webervirus* is indeed monophyletic

(**Figure 2**). To assess the number of different species present within the genus, we used

VIRIDIC to determine the intergenomic similarity between phage genomes (**Supplementary**

Figure B). Guidelines suggest any two phages belong to the same species if they are more

than 95 % identical across their entire genome (Turner et al., 2021). Based on these criteria,

our results show that weberviruses cluster into three distinct species.

As phages are among the most abundant biological entities on Earth, it is important to gain

knowledge on their presence within different environments. To better understand the wider

diversity and distribution of weberviruses, we searched the genome sequence of *Webervirus*

KLPN1 against metagenomic datasets using the online resource PhageClouds (Rangel-

Pineros et al., 2021). We used PhageClouds previously to demonstrate the presence of

slopekviruses in human faeces (Smith-Zaitlik et al., 2022). We identified 60 high-quality

webervirus MAGs in a range of samples, thereby doubling the number of webervirus

genomes available to the research community for genomic analyses (**Supplementary Table**

3). Their affiliation with the genus *Webervirus* was confirmed by our vConTACT2 (**Figure 3a**)

and phylogenetic analyses (**Supplementary Figure C**). We also determined that

weberviruses are distributed globally and predominated by phages associated with human

faeces or water supplies contaminated with human faeces (**Figure 3**). Lack of detection in

South America is likely due to absence of metagenomic datasets from this part of the world

rather than weberviruses not being represented in faecal samples from individuals living in

countries within this region. Compared with shotgun metagenomic datasets characterizing

the total microbiota found in faeces, there are very few studies – worldwide – examining

solely the intestinal virome, and PhageClouds is populated with phage genomes derived

from virome datasets.

It was notable when curating our MAG dataset that none of the studies describing these data

was able to predict hosts for the webervirus MAGs we have identified. Our analysis using

HostPhinder predicts that all 60 webervirus MAGs infect *K. pneumoniae*. HostPhinder

predicts the host species of a phage by searching for the most genetically similar phages in

a database of reference phages with known hosts (Villarroel et al., 2016). Although the

authors have shown that this whole-genome similarity-based approach is highly accurate,

host range can be altered by a relatively low number of mutations, especially those localized

to tail fibre proteins which are often determinants of host-cell specificity (Latka et al., 2021;

Taslem Mourosi et al., 2022). In the present study, we used a strain of *K. variicola* to isolate
three weberviruses and phages of this genus have also been isolated on *K. oxytoca* and *K.*
aerogenes. Although it is highly likely that all 127 weberviruses discussed herein are phages
of *Klebsiella* spp., determination of host range via plaque assays is still informative,
especially when determining therapeutic utility.
500
501

SUMMARY

503 We successfully characterized seven novel weberviruses that infect clinically relevant
504 *Klebsiella* spp. We have doubled the number of available webervirus genomes through
505 combining genomic data from isolated phage and MAG datasets. In doing so, we have
506 demonstrated the importance of interrogating MAG datasets to expand the availability of
507 curated phage genome sequences for use in genomic and ecological studies.
508
509

ACKNOWLEDGEMENTS

Consultant microbiologist Dr Frances Davies (Imperial College Healthcare NHS Trust) is
thanked for providing access to clinical strains. We thank Emily Goren for providing mixed-
liquor samples from Mogden Sewage Treatment Works (Thames Water). Dave Baker and
the QIB core sequencing team are thanked for WGS library preparation and sequencing for
the PS-prefixed bacterial genomes. Andrew Millard (University of Leicester) is thanked for
providing us with the PHROG hmm dataset for use with Prokka. Horst Neve is thanked for
generating TEMs of phages vB_KpnS-KLPN2, vB_KpnS-KLPN3 and vB_KpnS-KLPN4.
518

LH, ALM and DN designed the study. PS, ALM, SG, TT and LH isolated and purified the
phages. TCB and SJTD assembled and annotated the phage genomes. ALM and MK
processed clinical isolates for whole-genome sequencing; PS and LH assembled and
annotated the *Klebsiella* genomes. PS determined the antimicrobial profiles for the *Klebsiella*
strains included in this study. FN determined the host ranges of the MAGs. SJTD did all
bioinformatics work associated with the seven new phage genomes and the vConTACT
analyses; LH did all other bioinformatics work associated with the MAGs. DN produced
sequence data and TEM images. LH and ALM supervised PS. LH supervised TT, TCB and
SJTD. DN supervised SG. SJTD, DN and LH drafted the manuscript. All authors read and
approved the final version of the manuscript.

529
530

FUNDING

Imperial Health Charity is thanked for contributing to registration fees for the Professional

Doctorate studies of PS. PS was in receipt of an IBMS Research Grant (project title

"Isolation of lytic bacteriophages active against antibiotic-resistant *Klebsiella pneumoniae*").

This work used computing resources provided by UK Med-Bio (Medical Research Council

grant number MR/L01632X/1) and the Research Contingency Fund of the Department of

Biosciences, Nottingham Trent University. SJTD was funded by Nottingham Trent

University. SG completed this work as part of an MRes degree at NTU. LJH is funded by

Wellcome Trust Investigator Award no. 220876/Z/20/Z.

CONFLICTS OF INTEREST

543   The authors declare that there are no conflicts of interest.

ETHICS

547   The study of anonymised clinical isolates beyond the diagnostic requirement was approved

548   by an NHS research ethics committee (number 06/Q0406/20).

REFERENCES

Antimicrobial Resistance Collaborators, 2022. Global burden of bacterial antimicrobial
resistance in 2019: a systematic analysis. Lancet Lond. Engl. 399, 629–655.
https://doi.org/10.1016/S0140-6736(21)02724-0

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N.,
Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: a new genome assembly
algorithm and its applications to single-cell sequencing. J. Comput. Biol. J. Comput.
Mol. Cell Biol. 19, 455–477. https://doi.org/10.1089/cmb.2012.0021

Bastian, M., Heymann, S., Jacomy, M., 2009. Gephi: an open source software for exploring
and manipulating networks, in: International AAAI Conference on Weblogs and
Social Media.

Beamud, B., García-González, N., Gómez-Ortega, M., González-Candelas, F., Domingo-Calap,
P., Sanjuan, R., 2023. Genetic determinants of host tropism in Klebsiella phages. Cell
Rep. 42, 112048. https://doi.org/10.1016/j.celrep.2023.112048

Bin Jang, H., Bolduc, B., Zablocki, O., Kuhn, J.H., Roux, S., Adriaenssens, E.M., Brister, J.R.,
Kropinski, A.M., Krupovic, M., Lavigne, R., Turner, D., Sullivan, M.B., 2019. Taxonomic
assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing
networks. Nat. Biotechnol. 37, 632–639. https://doi.org/10.1038/s41587-019-0100-8

Bleriot, I., Blasco, L., Pacios, O., Fernández-García, L., Ambroa, A., López, M., Ortiz-
Cartagena, C., Cuenca, F.F., Oteo, J., Pascual, Á., Martínez-Martínez, L., Domingo-
Calap, P., Wood, T.K., Tomás, M., 2021. «PemIK (PemK/PemI) type II TA system from

Klebsiella pneumoniae clinical strains inhibits lytic phage».
https://doi.org/10.21203/rs.3.rs-679460/v1

Bolduc, B., Jang, H.B., Doulcier, G., You, Z.-Q., Roux, S., Sullivan, M.B., 2017. vConTACT: an
iVirus tool to classify double-stranded DNA viruses that infect Archaea and Bacteria.
PeerJ 5, e3243. https://doi.org/10.7717/peerj.3243

Brown, T.L., Petrovski, S., Hoyle, D., Chan, H.T., Lock, P., Tucci, J., 2017. Characterization and
formulation into solid dosage forms of a novel bacteriophage lytic against Klebsiella
oxytoca. PloS One 12, e0183510. https://doi.org/10.1371/journal.pone.0183510

Cai, R., Ren, Z., Zhao, R., Lu, Y., Wang, X., Guo, Z., Song, J., Xiang, W., Du, R., Zhang, X., Han,
W., Ru, H., Gu, J., 2023. Structural biology and functional features of phage-derived
depolymerase Depo32 on Klebsiella pneumoniae with K2 serotype capsular
polysaccharides. Microbiol. Spectr. 11, e0530422.
https://doi.org/10.1128/spectrum.05304-22

Cai, R., Wang, Z., Wang, G., Zhang, H., Cheng, M., Guo, Z., Ji, Y., Xi, H., Wang, X., Xue, Y., Ur
Rahman, S., Sun, C., Feng, X., Lei, L., Tong, Y., Han, W., Gu, J., 2019. Biological
properties and genomics analysis of vB_KpnS_GH-K3, a Klebsiella phage with a
putative depolymerase-like protein. Virus Genes. https://doi.org/10.1007/s11262-
019-01681-z

Camarillo-Guerrero, L.F., Almeida, A., Rangel-Pineros, G., Finn, R.D., Lawley, T.D., 2021.
Massive expansion of human gut bacteriophage diversity. Cell 184, 1098-1109.e9.
https://doi.org/10.1016/j.cell.2021.01.029

Cao, F., Wang, X., Wang, Linhui, Li, Z., Che, J., Wang, Lili, Li, X., Cao, Z., Zhang, J., Jin, L., Xu,
Y., 2015. Evaluation of the efficacy of a bacteriophage in the treatment of
pneumonia induced by multidrug resistance Klebsiella pneumoniae in mice. BioMed
Res. Int. 2015, 752930. https://doi.org/10.1155/2015/752930

Castillo, M., Tran, R., Newkirk, H., Liu, M., Gill, J.J., Ramsey, J., 2019. Complete Genome
Sequence of Sin4, a Siphophage Infecting Carbapenemase-Producing Klebsiella
pneumoniae. Microbiol. Resour. Announc. 8. https://doi.org/10.1128/MRA.01048-19

Chklovski, A., Parks, D.H., Woodcroft, B.J., Tyson, G.W., 2022. CheckM2: a rapid, scalable
and accurate tool for assessing microbial genome quality using machine learning.
https://doi.org/10.1101/2022.07.11.499243

Cranston, A., Danielson, P., Arens, D.K., Barker, A., Birch, E.K., Brown, H., Carr, E., Cero, P.,
Chow, J., Correa, E., Dean, J., Dunn, M., Eberhard, N., Egbert, A., Foster, K., Gaertner,
R., Gleave, A., Gomez, A., Gordon, J.B., Harris, E.B., Heaps, C., Hyer, M., Johnson, A.,
Johnson, L., Kim, M., Kruger, J.L., Leonard, T., LeSueur, A., Lima, S., Marshall, N.,
Moulton, R., Newey, C.R., Owen, D., Packard, A., Rolfson, A., Suorsa, A.R., Rodriguez,
W., Sandoval, C., Sharma, R., Smith, A., Sork, C., Soule, C., Soule, S., Stewart, J.,
Stoker, T., Thompson, D.W., Thurgood, T., Walker, J., Zaugg, E., Casjens, S.R., Grose,
J.H., 2022. Genome Sequences of 22 T1-like Bacteriophages That Infect
Enterobacteriales. Microbiol. Resour. Announc. 11, e0122121.
https://doi.org/10.1128/mra.01221-21

Dion, M.B., Plante, P.-L., Zufferey, E., Shah, S.A., Corbeil, J., Moineau, S., 2021. Streamlining
CRISPR spacer-based bacterial host predictions to decipher the viral dark matter.
Nucleic Acids Res. 49, 3127–3138. https://doi.org/10.1093/nar/gkab133

Dunstan, R.A., Bamert, R.S., Belousoff, M.J., Short, F.L., Barlow, C.K., Pickard, D.J., Wilksch,
J.J., Schittenhelm, R.B., Strugnell, R.A., Dougan, G., Lithgow, T., 2021. Mechanistic
Insights into the Capsule-Targeting Depolymerase from a Klebsiella pneumoniae

Bacteriophage. Microbiol. Spectr. 9, e0102321.
https://doi.org/10.1128/Spectrum.01023-21
Fang, Q., Feng, Y., McNally, A., Zong, Z., 2022. Characterization of phage resistance and
phages capable of intestinal decolonization of carbapenem-resistant Klebsiella
pneumoniae in mice. Commun. Biol. 5, 48. https://doi.org/10.1038/s42003-022-
03001-y
Gao, S., Linden, S.B., Nelson, D.C., 2017. Complete Genome Sequence of Klebsiella
pneumoniae Phages SopranoGao, MezzoGao, and AltoGao. Genome Announc. 5.
https://doi.org/10.1128/genomeA.01009-17
Gramer, J., Kenny, S., Newkirk, H., Liu, M., Gill, J.J., Ramsey, J., 2019. Complete Genome
Sequence of Klebsiella pneumoniae Siphophage Skenny. Microbiol. Resour.
Announc. 8. https://doi.org/10.1128/MRA.01036-19
Gregory, A.C., Zablocki, O., Zayed, A.A., Howell, A., Bolduc, B., Sullivan, M.B., 2020. The Gut
Virome Database Reveals Age-Dependent Patterns of Virome Diversity in the Human
Gut. Cell Host Microbe 28, 724-740.e8. https://doi.org/10.1016/j.chom.2020.08.003
Haines, M.E.K., Hodges, F.E., Nale, J.Y., Mahony, J., van Sinderen, D., Kaczorowska, J.,
Alrashid, B., Akter, M., Brown, N., Sauvageau, D., Sicheritz-Pontén, T., Thanki, A.M.,
Millard, A.D., Galyov, E.E., Clokie, M.R.J., 2021. Analysis of Selection Methods to
Develop Novel Phage Therapy Cocktails Against Antimicrobial Resistant Clinical
Isolates of Bacteria. Front. Microbiol. 12, 613529.
https://doi.org/10.3389/fmicb.2021.613529
Herridge, W.P., Shibu, P., O'Shea, J., Brook, T.C., Hoyles, L., 2020. Bacteriophages of
Klebsiella spp., their diversity and potential therapeutic uses. J. Med. Microbiol. 69,
176–194. https://doi.org/10.1099/jmm.0.001141
Hesse, S., Rajaure, M., Wall, E., Johnson, J., Bliskovsky, V., Gottesman, S., Adhya, S., 2020.
Phage Resistance in Multidrug-Resistant Klebsiella pneumoniae ST258 Evolves via
Diverse Mutations That Culminate in Impaired Adsorption. mBio 11, e02530-19.
https://doi.org/10.1128/mBio.02530-19
Horváth, M., Kovács, T., Koderivalappil, S., Ábrahám, H., Rákhely, G., Schneider, G., 2020.
Identification of a newly isolated lytic bacteriophage against K24 capsular type,
carbapenem resistant Klebsiella pneumoniae isolates. Sci. Rep. 10, 5891.
https://doi.org/10.1038/s41598-020-62691-8
Hoyles, L., Murphy, J., Neve, H., Heller, K.J., Turton, J.F., Mahony, J., Sanderson, J.D.,
Hudspith, B., Gibson, G.R., McCartney, A.L., van Sinderen, D., 2015. Klebsiella
pneumoniae subsp. pneumoniae-bacteriophage combination from the caecal
effluent of a healthy woman. PeerJ 3, e1061. https://doi.org/10.7717/peerj.1061
Hudson, A.T., Clark, J., Gill, J.J., Liu, M., 2021. Complete Genome Sequence of Klebsiella
aerogenes Siphophage Solomon. Microbiol. Resour. Announc. 10, e01419-20.
https://doi.org/10.1128/MRA.01419-20
Jolley, K.A., Bray, J.E., Maiden, M.C.J., 2018. Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. Wellcome Open
Res. 3, 124. https://doi.org/10.12688/wellcomeopenres.14826.1
Kęsik-Szeloch, A., Drulis-Kawa, Z., Weber-Dąbrowska, B., Kassner, J., Majkowska-Skrobek, G.,
Augustyniak, D., Lusiak-Szelachowska, M., Zaczek, M., Górski, A., Kropinski, A.M.,
2013. Characterising the biology of novel lytic bacteriophages infecting multidrug
resistant Klebsiella pneumoniae. Virol. J. 10, 100. https://doi.org/10.1186/1743-
422X-10-100

Kim, S.J., Jo, J., Ko, K.S., 2023. Lipid A modification-induced colistin-resistant Klebsiella
variicola from healthy adults. J. Med. Microbiol. 72.
https://doi.org/10.1099/jmm.0.001680
Knecht, L.E., Veljkovic, M., Fieseler, L., 2019. Diversity and Function of Phage Encoded
Depolymerases. Front. Microbiol. 10, 2949.
https://doi.org/10.3389/fmicb.2019.02949
Latka, A., Leiman, P.G., Drulis-Kawa, Z., Briers, Y., 2019. Modeling the Architecture of
Depolymerase-Containing Receptor Binding Proteins in Klebsiella Phages. Front.
Microbiol. 10, 2649. https://doi.org/10.3389/fmicb.2019.02649
Latka, A., Lemire, S., Grimon, D., Dams, D., Maciejewska, B., Lu, T., Drulis-Kawa, Z., Briers, Y.,
2021. Engineering the Modular Receptor-Binding Proteins of Klebsiella Phages
Switches Their Capsule Serotype Specificity. mBio 12, e00455-21.
https://doi.org/10.1128/mBio.00455-21
Le, S.Q., Gascuel, O., 2008. An improved general amino acid replacement matrix. Mol. Biol.
Evol. 25, 1307–1320. https://doi.org/10.1093/molbev/msn067
Li, L., Zhang, Y., Wang, W., Chen, Y., He, F., Yu, Y., 2024. Global Emergence and Genomic
Epidemiology of blaNDM-Carrying Klebsiella variicola. Infect. Drug Resist. 17, 1893–
1901. https://doi.org/10.2147/IDR.S460569
Ma, Y., You, X., Mai, G., Tokuyasu, T., Liu, C., 2018. A human gut phage catalog correlates
the gut phageome with type 2 diabetes. Microbiome 6, 24.
https://doi.org/10.1186/s40168-018-0410-y
Majkowska-Skrobek, G., Łątka, A., Berisio, R., Maciejewska, B., Squeglia, F., Romano, M.,
Lavigne, R., Struve, C., Drulis-Kawa, Z., 2016. Capsule-Targeting Depolymerase,
Derived from Klebsiella KP36 Phage, as a Tool for the Development of Anti-Virulent
Strategy. Viruses 8, E324. https://doi.org/10.3390/v8120324
Martinez, N., Williams, E., Newkirk, H., Liu, M., Gill, J.J., Ramsey, J., 2019. Complete Genome
Sequence of Klebsiella pneumoniae Phage Sweeny. Microbiol. Resour. Announc. 8.
https://doi.org/10.1128/MRA.01047-19
Mishra, C.K., Choi, T.J., Kang, S.C., 2012. Isolation and characterization of a bacteriophage
F20 virulent to Enterobacter aerogenes. J. Gen. Virol. 93, 2310–2314.
https://doi.org/10.1099/vir.0.043562-0
Moraru, C., Varsani, A., Kropinski, A.M., 2020. VIRIDIC—A Novel Tool to Calculate the
Intergenomic Similarities of Prokaryote-Infecting Viruses. Viruses 12, 1268.
https://doi.org/10.3390/v12111268
Nayfach, S., Camargo, A.P., Schulz, F., Eloe-Fadrosh, E., Roux, S., Kyrpides, N.C., 2021a.
CheckV assesses the quality and completeness of metagenome-assembled viral
genomes. Nat. Biotechnol. 39, 578–585. https://doi.org/10.1038/s41587-020-00774-
7
Nayfach, S., Páez-Espino, D., Call, L., Low, S.J., Sberro, H., Ivanova, N.N., Proal, A.D.,
Fischbach, M.A., Bhatt, A.S., Hugenholtz, P., Kyrpides, N.C., 2021b. Metagenomic
compendium of 189,680 DNA viruses from the human gut microbiome. Nat.
Microbiol. 6, 960–970. https://doi.org/10.1038/s41564-021-00928-6
Nazir, A., Qi, C., Shi, N., Gao, X., Feng, Q., Qing, H., Li, F., Tong, Y., 2022. Characterization and
Genomic Analysis of a Novel Drexlervirial Bacteriophage IME268 with Lytic Activity
Against Klebsiella pneumoniae. Infect. Drug Resist. 15, 1533–1546.
https://doi.org/10.2147/IDR.S347110

Newberry, F., Shibu, P., Smith-Zaitlik, T., Eladawy, M., McCartney, A.L., Hoyles, L., Negus, D.,
2023. Phage vB_KmiS-Kmi2C infects members of the Klebsiella oxytoca complex and
represents a novel genus of lytic bacteriophage. bioRxiv.
https://doi.org/10.1101/2023.01.12.523727
Nguyen, D.T., Lessor, L.E., Cahill, J.L., Rasche, E.S., Kuty Everett, G.F., 2015. Complete
Genome Sequence of Klebsiella pneumoniae Carbapenemase-Producing K.
pneumoniae Siphophage Sushi. Genome Announc. 3.
https://doi.org/10.1128/genomeA.00994-15
Nir-Paz, R., Gelman, D., Khouri, A., Sisson, B.M., Fackler, J., Alkalay-Oren, S., Khalifa, L.,
Rimon, A., Yerushalmy, O., Bader, R., Amit, S., Coppenhagen-Glazer, S., Henry, M.,
Quinones, J., Malagon, F., Biswas, B., Moses, A.E., Merril, G., Schooley, R.T.,
Brownstein, M.J., Weil, Y.A., Hazan, R., 2019. Successful Treatment of Antibiotic-
resistant, Poly-microbial Bone Infection With Bacteriophages and Antibiotics
Combination. Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am. 69, 2015–2018.
https://doi.org/10.1093/cid/ciz222
Nishimura, Y., Yoshida, T., Kuronishi, M., Uehara, H., Ogata, H., Goto, S., 2017. ViPTree: the
viral proteomic tree server. Bioinforma. Oxf. Engl. 33, 2379–2380.
https://doi.org/10.1093/bioinformatics/btx157
Pagès, H., Aboyoun, P., Gentleman, R., DebRoy, S., 2021. Biostrings: Efficient manipulation
of biological strings. R package version 2.62.0.
Paradis, E., Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinforma. Oxf. Engl. 35, 526–528.
https://doi.org/10.1093/bioinformatics/bty633
Park, E.-A., Kim, Y.-T., Cho, J.-H., Ryu, S., Lee, J.-H., 2017. Characterization and genome
analysis of novel bacteriophages infecting the opportunistic human pathogens
Klebsiella oxytoca and K. pneumoniae. Arch. Virol. 162, 1129–1139.
https://doi.org/10.1007/s00705-016-3202-3
Pertics, B.Z., Cox, A., Nyúl, A., Szamek, N., Kovács, T., Schneider, G., 2021. Isolation and
Characterization of a Novel Lytic Bacteriophage against the K2 Capsule-Expressing
Hypervirulent Klebsiella pneumoniae Strain 52145, and Identification of Its
Functional Depolymerase. Microorganisms 9, 650.
https://doi.org/10.3390/microorganisms9030650
Rangel-Pineros, G., Millard, A., Michniewski, S., Scanlan, D., Sirén, K., Reyes, A., Petersen, B.,
Clokie, M.R.J., Sicheritz-Pontén, T., 2021. From Trees to Clouds: PhageClouds for Fast
Comparison of ~640,000 Phage Genomic Sequences and Host-Centric Visualization
Using Genomic Network Graphs. PHAGE 2, 194–203.
https://doi.org/10.1089/phage.2021.0008
Richardson, R.W., Lessor, L., O'Leary, C., Gill, J., Liu, M., 2019. Complete Genome Sequence
of Klebsiella pneumoniae Siphophage Sanco. Microbiol. Resour. Announc. 8.
https://doi.org/10.1128/MRA.01252-19
Rodríguez-Medina, N., Barrios-Camacho, H., Duran-Bedolla, J., Garza-Ramos, U., 2019.
Klebsiella variicola: an emerging pathogen in humans. Emerg. Microbes Infect. 8,
973–988. https://doi.org/10.1080/22221751.2019.1634981
Saldana, R., Newkirk, H., Liu, M., Gill, J.J., Ramsey, J., 2019. Complete Genome Sequence of
Shelby, a Siphophage Infecting Carbapenemase-Producing Klebsiella pneumoniae.
Microbiol. Resour. Announc. 8. https://doi.org/10.1128/MRA.01037-19

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri,
K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-
image analysis. Nat. Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019
Schliep, K., Potts, A.J., Morrison, D.A., Grimm, G.W., 2017. Intertwining phylogenetic trees
and networks. Methods Ecol. Evol. 8, 1212–1220. https://doi.org/10.1111/2041-
210X.12760
Scholl, D., Adhya, S., Merril, C., 2005. Escherichia coli K1's capsule is a barrier to
bacteriophage T7. Appl. Environ. Microbiol. 71, 4872–4874.
https://doi.org/10.1128/AEM.71.8.4872-4874.2005
Seemann, T., 2014. Prokka: rapid prokaryotic genome annotation. Bioinforma. Oxf. Engl. 30,
2068–2069. https://doi.org/10.1093/bioinformatics/btu153
Shibu, P., 2019. Investigations of carbapenem-resistant Klebsiella species and associated
clinical considerations. Westminster, London.
Shibu, P., McCuaig, F., McCartney, A.L., Kujawska, M., Hall, L.J., Hoyles, L., 2021. Improved
molecular characterization of the Klebsiella oxytoca complex reveals the prevalence
of the kleboxymycin biosynthetic gene cluster. Microb. Genomics 7.
https://doi.org/10.1099/mgen.0.000592
Smith-Zaitlik, T., Shibu, P., McCartney, A.L., Foster, G., Hoyles, L., Negus, D., 2022. Extended
genomic analyses of the broad-host-range phages vB_KmiM-2Di and vB_KmiM-4Dii
reveal slopekviruses have highly conserved genomes.
https://doi.org/10.1101/2022.04.06.486684
South, A., 2011. rworldmap: a New R package for mapping global data. R J. 3, 35–43.
Tabassum, R., Shafique, M., Khawaja, K.A., Alvi, I.A., Rehman, Y., Sheik, C.S., Abbas, Z.,
Rehman, S.U., 2018. Complete genome analysis of a Siphoviridae phage TSK1
showing biofilm removal potential against Klebsiella pneumoniae. Sci. Rep. 8, 17904.
https://doi.org/10.1038/s41598-018-36229-y
Taslem Mourosi, J., Awe, A., Guo, W., Batra, H., Ganesh, H., Wu, X., Zhu, J., 2022.
Understanding Bacteriophage Tail Fiber Interaction with Host Surface Receptor: The
Key "Blueprint" for Reprogramming Phage Host Range. Int. J. Mol. Sci. 23, 12146.
https://doi.org/10.3390/ijms232012146
Ter Horst, A.M., Santos-Medellín, C., Sorensen, J.W., Zinke, L.A., Wilson, R.M., Johnston,
E.R., Trubl, G., Pett-Ridge, J., Blazewicz, S.J., Hanson, P.J., Chanton, J.P., Schadt, C.W.,
Kostka, J.E., Emerson, J.B., 2021. Minnesota peat viromes reveal terrestrial and
aquatic niche partitioning for local and global viral populations. Microbiome 9, 233.
https://doi.org/10.1186/s40168-021-01156-0
Terzian, P., Olo Ndela, E., Galiez, C., Lossouarn, J., Pérez Bucio, R.E., Mom, R., Toussaint, A.,
Petit, M.-A., Enault, F., 2021. PHROG: families of prokaryotic virus proteins clustered
using remote homology. NAR Genomics Bioinforma. 3, lqab067.
https://doi.org/10.1093/nargab/lqab067
Thung, T.Y., White, M.E., Dai, W., Wilksch, J.J., Bamert, R.S., Rocker, A., Stubenrauch, C.J.,
Williams, D., Huang, C., Schittelhelm, R., Barr, J.J., Jameson, E., McGowan, S., Zhang,
Y., Wang, J., Dunstan, R.A., Lithgow, T., 2021. Component Parts of Bacteriophage
Virions Accurately Defined by a Machine-Learning Approach Built on Evolutionary
Features. mSystems 6, e0024221. https://doi.org/10.1128/mSystems.00242-21
Thurgood, T.L., Sharma, R., Call, J.J., Chronis, J.D., Dawson, D.D., Finnegan, Z.K., Foster, K.W.,
Meek, T., Potts, E., Sirrine, M.R., Atkinson, A.D., Fairholm, J.D., Handoko, Y.A., Ong,

K.L., Tateoka, O.B., Arens, D.K., Johnson, L., Kruger, J.L., Loertscher, E., Thompson,
D.W., Walker, J.K., Robison, R.A., Casjens, S.R., Grose, J.H., 2020. Genome Sequences
of 12 Phages That Infect Klebsiella pneumoniae. Microbiol. Resour. Announc. 9,
e00024-20. https://doi.org/10.1128/MRA.00024-20

Tisza, M.J., Buck, C.B., 2021. A catalog of tens of thousands of viruses from human
metagenomes reveals hidden associations with chronic diseases. Proc. Natl. Acad.
Sci. U. S. A. 118, e2023202118. https://doi.org/10.1073/pnas.2023202118

Turner, D., Kropinski, A.M., Adriaenssens, E.M., 2021. A Roadmap for Genome-Based Phage
Taxonomy. Viruses 13, 506. https://doi.org/10.3390/v13030506

Villarroel, J., Kleinheinz, K.A., Jurtz, V.I., Zschach, H., Lund, O., Nielsen, M., Larsen, M.V.,
2016. HostPhinder: A Phage Host Prediction Tool. Viruses 8, 116.
https://doi.org/10.3390/v8050116

Walker, P.J., Siddell, S.G., Lefkowitz, E.J., Mushegian, A.R., Adriaenssens, E.M., Alfenas-
Zerbini, P., Davison, A.J., Dempsey, D.M., Dutilh, B.E., García, M.L., Harrach, B.,
Harrison, R.L., Hendrickson, R.C., Junglen, S., Knowles, N.J., Krupovic, M., Kuhn, J.H.,
Lambert, A.J., Łobocka, M., Nibert, M.L., Oksanen, H.M., Orton, R.J., Robertson, D.L.,
Rubino, L., Sabanadzovic, S., Simmonds, P., Smith, D.B., Suzuki, N., Van Dooerslaer,
K., Vandamme, A.-M., Varsani, A., Zerbini, F.M., 2021. Changes to virus taxonomy
and to the International Code of Virus Classification and Nomenclature ratified by
the International Committee on Taxonomy of Viruses (2021). Arch. Virol. 166, 2633–
2648. https://doi.org/10.1007/s00705-021-05156-1

Walker, P.J., Siddell, S.G., Lefkowitz, E.J., Mushegian, A.R., Dempsey, D.M., Dutilh, B.E.,
Harrach, B., Harrison, R.L., Hendrickson, R.C., Junglen, S., Knowles, N.J., Kropinski,
A.M., Krupovic, M., Kuhn, J.H., Nibert, M., Rubino, L., Sabanadzovic, S., Simmonds, P.,
Varsani, A., Zerbini, F.M., Davison, A.J., 2019. Changes to virus taxonomy and the
International Code of Virus Classification and Nomenclature ratified by the
International Committee on Taxonomy of Viruses (2019). Arch. Virol. 164, 2417–
2429. https://doi.org/10.1007/s00705-019-04306-w

Wick, R.R., Heinz, E., Holt, K.E., Wyres, K.L., 2018. Kaptive Web: User-Friendly Capsule and
Lipopolysaccharide Serotype Prediction for Klebsiella Genomes. J. Clin. Microbiol. 56.
https://doi.org/10.1128/JCM.00197-18

Wick, R.R., Judd, L.M., Gorrie, C.L., Holt, K.E., 2017. Unicycler: Resolving bacterial genome
assemblies from short and long sequencing reads. PLoS Comput. Biol. 13, e1005595.
https://doi.org/10.1371/journal.pcbi.1005595

Wick, R.R., Schultz, M.B., Zobel, J., Holt, K.E., 2015. Bandage: interactive visualization of de
novo genome assemblies. Bioinforma. Oxf. Engl. 31, 3350–3352.
https://doi.org/10.1093/bioinformatics/btv383

Wintachai, P., Naknaen, A., Thammaphet, J., Pomwised, R., Phaonakrop, N., Roytrakul, S.,
Smith, D.R., 2020. Characterization of extended-spectrum-β-lactamase producing
Klebsiella pneumoniae phage KP1801 and evaluation of therapeutic efficacy in vitro
and in vivo. Sci. Rep. 10, 11803. https://doi.org/10.1038/s41598-020-68702-y

Zurabov, F., Zhilenkov, E., 2019. Complete Genome Sequences of Lytic Polysaccharide-
Degrading Klebsiella pneumoniae Bacteriophages vB_KpnS_FZ10, vB_KpnP_FZ12,
vB_KpnM_FZ14, and vB_KpnS_FZ41. Microbiol. Resour. Announc. 8.
https://doi.org/10.1128/MRA.00914-19

Table 1. Strains of *Klebsiella* included in this study and their phage infection profiles

Strain *	Source	K type †	O antigen†	MLST ‡	Infection type §							
					KLPN1	KLPN2	KLPN3	KLPN4	KLPN5	KLPN6	KLPN7	KLPN8
L4-FAA5	Human caecum (Hoyles et al., 2015)	K2	O1v1	ST380	++++ d	++++ d	++++ d	++++ d	0	0	0	0
PS_misc2	Groin (Shibu, 2019)	K64	O1v1	ST14	0	0	0	0	0	0	0	0
PS_misc3	Rectum (Shibu, 2019)	K52	OL101	ST45	0	0	0	0	0	0	0	0
PS_misc5	Rectum (Shibu, 2019)	K81	OL101	16-24-21-31-47-22-281	0	0	0	0	++++ d	++++ d	++++ d	0
PS_misc6	Rectum (Shibu, 2019)	K110	O2v1	ST716	0	0	0	0	0	0	0	+++ d
PS_misc7	Rectum (Shibu, 2019)	K30	O1v1	ST294	0	0	0	0	0	0	0	0
PS_misc8	Perineum (Shibu, 2019)	K52	OL101	ST45	0	0	0	0	0	0	0	0
PS_Kpn1	Perineum (Shibu, 2019)	K2	O4	1-X-1-1-1-1	+++	++	++	++	0	0	0	0
PS_Kpn2	Rectum (Shibu, 2019)	K64	O1v1	ST14	+++	+++	+++	++	0	0	0	0
PS_Kpn3	Rectum (Shibu, 2019)	K64	O1v1	1-1-1-1-1-1-X	0	0	0	0	0	0	0	0
PS_Kpn4	Cross-infection (Shibu, 2019)	K64	O1v2	ST15	0	0	0	0	0	0	0	0
PS_Kpn7	Rectum (Shibu, 2019)	K2	O1v1	ST14	++	++	++	++	0	0	0	0
PS_Kpn9	Rectum (Shibu, 2019)	K18	O2v1	2-1-1-X-1-4	0	0	0	0	0	0	0	0
PS_Kpn10	Rectum (Shibu, 2019)	K2	O1v1	ST14	++++	+++	+++	++	0	0	0	0
PS_Kpn11	Urine (Shibu, 2019)	K107	O2v2	ST258	0	0	0	0	0	0	0	0
PS_Kpn12	Rectum (Shibu, 2019)	K15	O4	ST11	0	0	0	0	0	0	0	0
PS_Kpn13	Rectum (Shibu, 2019)	K2	O1v1	ST14	+++	+++	+++	0	0	0	0	0
PS_Kpn14	Rectum (Shibu, 2019)	K2	O1v1	ST14	++++	+++	+++	++	0	0	0	0
PS_Kpn15	Rectum (Shibu, 2019)	K52	OL101	ST45	0	0	0	0	0	0	0	0
PS_Kpn16	Rectum (Shibu, 2019)	K17	O1v1	ST101	0	0	0	0	0	0	0	0
PS_Kpn24	Rectum (Shibu, 2019)	K2	O1v1	ST14	++	++	++	++	++	++	++	0
PS_Kpn25	Rectum (Shibu, 2019)	K2	O1v1	ST14	+++	++	++	++	0	0	0	0
PS_Kpn26	Rectum (Shibu, 2019)	K64	O1v1	ST147	0	0	0	0	0	0	0	0
PS_Kpn27	Urine (Shibu, 2019)	K107	O2v2	ST258	0	0	0	0	0	0	0	0
PS_Kpn28	Urine (Shibu, 2019)	K2	O1v1	ST14	++++	+++	++	++	0	0	0	0
PS_Kpn29	Urine (Shibu, 2019)	K2	O1v1	ST14	++	++	++	++	0	0	0	0
PS_Kpn30	Mouth (Shibu, 2019)	K112	O1v1	ST15	0	0	0	0	0	0	0	0
PS_Kpn31	Perineum (Shibu, 2019)	K2	O1v1	ST14	++++	++	++	++	0	0	0	0
PS_Kpn32	Drain fluid (Shibu, 2019)	K22	O1v1	ST35	0	0	0	0	0	0	0	0
PS_Kpn33	Urine (Shibu, 2019)	K2	O1v1	ST14	+++ (d)	+++ (d)	++	++	0	0	0	0
PS_Kpn35	Urine (Shibu, 2019)	K2	O1v1	ST14	+++ (d)	+++ (d)	++	++	0	0	0	0
PS_Kpn36	Urine (Shibu, 2019)	K24	O2v1	ST11	0	0	0	0	0	0	0	0
PS_Kpn37	Wound (Shibu, 2019)	K51	O3b	ST16	0	0	0	0	0	0	0	0
PS_Kpn38	High vaginal swab (Shibu, 2019)	K2	O1v1	ST14	+++	++	+++	++	0	0	0	0
PS_Kpn39	Wound (Shibu, 2019)	K64	O1v1	ST14	0	0	0	0	0	0	0	0
PS_Kpn40	Wound (Shibu, 2019)	K107	O2v2	ST512	0	0	0	0	0	0	0	0
PS_Kpn41	Leg (Shibu, 2019)	K112	O1v1	ST15	0	0	0	0	0	0	0	0

* All strains with PS prefix identified as *K. pneumoniae* subsp. *pneumoniae* by average nucleotide identity and phylogenetic analyses against

type strains of the genus *Klebsiella*, except for PS_misc5 (*K. variicola* subsp. *variicola*) (Shibu, 2019).

† Determined using Kaptive (Wick et al., 2018).

‡ MLST, multi-locus sequence type determined via PubMLST (Jolley et al., 2018).

857 § The appearance of the spot was graded according to (Haines et al., 2021): ++++, complete lysis; +++, lysis with resistant colonies; ++, hazy
858 lysis; +, visible plaques; 0, no visible plaques. We also noted whether depolymerase activity was detected (i.e. formation of haloes around
859 plaques): d, depolymerase activity; (d), weak depolymerase activity detected.
860

861 **Table 2.** Webervirus genomes associated with *Klebsiella* hosts included in this study

Phage/species	Accession	Isolated from	Location	Reference
Klebsiella phage vB_KpnS-KLPN2	OM065837	Sewage	UK	This study
Klebsiella phage vB_KpnS-KLPN3	OM065838	Sewage	UK	This study
Klebsiella phage vB_KpnS-KLPN4	OM065839	Sewage	UK	This study
Klebsiella phage vB_KvaS-KLPN5	OM065840	Sewage	UK	This study
Klebsiella phage vB_KvaS-KLPN6	OM065841	Sewage	UK	This study
Klebsiella phage vB_KvaS-KLPN7	OM065842	Sewage	UK	This study
Klebsiella phage vB_KpnS-KLPN8	OM065843	Sewage	UK	This study
Klebsiella phage 066039 (P39)	MW042802.1	Sewage	China	(Fang et al., 2022)
Klebsiella phage ABTNL-2	MZ221764.1	Sewage	China	–
Klebsiella phage B1	MW672037.1	Sewage	Hungary	(Pertics et al., 2021)
Klebsiella phage IME268	MZ398242.1	Unknown	China	(Nazir et al., 2022)
Klebsiella phage LF20	MW417503.1	Hospital wastewater	China	–
Klebsiella phage MMBB	MT894005.1	Merri Creek watercourse	Australia	(Thung et al., 2021)
Klebsiella phage NJR15	NC_048044.1	Sewage	China	–
Klebsiella phage NJS2	NC_048043.1	Sewage	China	–
Klebsiella phage NJS3	MH633486.1	Hospital sewage	China	–
Klebsiella phage P1	MZ598515.1	Water	China	–
Klebsiella phage P528	MW021764.1	Sewage	USA	(Cranston et al., 2022)
Klebsiella phage PhiKpNIH-10	MN395285.1	Wastewater treatment plant	USA	(Hesse et al., 2020)
Klebsiella phage RAD2	NC_055956.1	Hospital wastewater	UK	(Dunstan et al., 2021)
Klebsiella phage Sanco	MK618657.1	Wastewater treatment plant	USA	(Richardson et al., 2019)
Klebsiella phage SH-Kp 160016	KY575286.1	Sewage	China	–
Klebsiella phage Solomon	MT701592.1	Wastewater treatment plant	USA	(Hudson et al., 2021)
Klebsiella phage vB_KpnS_2811	LR757892.1	Unknown	UK	(Haines et al., 2021)
Klebsiella phage vB_KpnS_KingDDD	MN013078.1	Sewage	USA	(Thurgood et al., 2020)
Klebsiella phage vB_KpnS_Penguinator	MN013087.1	Sewage	USA	(Thurgood et al., 2020)
Klebsiella phage vB_KpnS_ZX2	MW722081.1	Faeces	China	–
Klebsiella phage vB_kpnS-VAC10	MZ428227.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_KpnS-VAC11	MZ428228.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_kpnS-VAC2	MZ428221.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_KpnS-VAC4	MZ428222.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_KpnS-VAC5	MZ428223.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_KpnS-VAC6	MZ428224.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_KpnS-VAC7	MZ428225.1	Sewage	Spain	(Bleriot et al., 2021)
Klebsiella phage vB_KpnS-VAC8	MZ428226.1	Sewage	Spain	(Bleriot et al., 2021)
Webervirus alina	NC_049837.1	Sewage	USA	(Thurgood et al., 2020)
Webervirus call	NC_049836.1	Sewage	USA	(Thurgood et al., 2020)
Webervirus domnhall	NC_049835.1	Sewage	USA	(Thurgood et al., 2020)
Webervirus F20	NC_043469.1	Lake water	South Korea	(Mishra et al., 2012)
Webervirus FZ10	NC_049840.1	Sewage	Russia	(Zurabov and Zhilenkov, 2019)
Webervirus GHK3	NC_048162.1	Sewage	China	(Cai et al., 2019)
Webervirus IMGroot	NC_049834.1	Sewage	USA	(Thurgood et al., 2020)
Webervirus JY917	NC_049843.1	Unknown	China	–
Webervirus KL	NC_049838.1	Unknown	Unknown	–
Webervirus KLPN1	NC_028760.1	Human caecal effluent	UK	(Hoyles et al., 2015)
Webervirus KLPPOU149	NC_049842.1	Unknown	Cote d'Ivoire	–
Webervirus KOX1	NC_047825.1	Wastewater	Australia	(Brown et al., 2017)
Webervirus KP1801	NC_049848.1	Hospital wastewater	Thailand	(Wintachai et al., 2020)
Webervirus KP36	NC_029099.1	Wastewater	Poland	(Majkowska-Skrobek et al., 2016)
Webervirus KpCol1	NC_047907.1	Wastewater	Turkey	–
Webervirus KpKT21phi1	NC_048143.1	Unknown	USA	(Nir-Paz et al., 2019)
Webervirus KpNIH2	NC_049845.1	Wastewater	USA	(Hesse et al., 2020)
Webervirus KpV522	NC_047784.1	Sewage	Russia	–
Webervirus mezzogao	NC_047850.1	Wastewater	USA	(Gao et al., 2017)
Webervirus N141	NC_047841.1	Unknown	South Korea	–

Phage/species	Accession	Isolated from	Location	Reference
Webervirus NJS1	NC_048024.1	Sewage	China	–
Webervirus PKP126	NC_031053.1	Sewage	South Korea	(Park et al., 2017)
Webervirus segescirculi	NC_049833.1	Sewage	USA	(Thurgood et al., 2020)
Webervirus shelby	NC_049846.1	Pond water	USA	(Saldana et al., 2019)
Webervirus sin4	NC_049847.1	Wastewater	USA	(Castillo et al., 2019)
Webervirus skenny	NC_049841.1	Activated sludge	USA	(Gramer et al., 2019)
Webervirus sushi	NC_028774.1	Sewage	USA	(Nguyen et al., 2015)
Webervirus sweeny	NC_049839.1	Wastewater	USA	(Martinez et al., 2019)
Webervirus TAH8	NC_048042.1	Hospital sewage	China	–
Webervirus TSK1	NC_048126.1	Sewage	Pakistan	(Tabassum et al., 2018)
Webervirus wv13	NC_049844.1	Wastewater	Hungary	(Horváth et al., 2020)
Webervirus wv1513	NC_028786.1	Sewage	China	(Cao et al., 2015)

Figure 1. vConTACT filtered gene-sharing network in which only nodes connected to the

main cluster are shown (filtered based on giant component; Fruchterman Reingold layout).

The modularity of the network was determined (Gephi options selected: randomness, use

weights, resolution 1.0). The network comprised 33 modules. Only modules representing >4

% of all nodes in the network are coloured. The circled module contained all 67 webervirus

genomes (orange nodes) plus some other members of the family *Drexlerviridae* (yellow

nodes). All the *Klebsiella* phage genomes (i.e. weberviruses) clustered together.

Figure 2. Phylogenetic relationships (maximum-likelihood tree) of members of the family

Drexlerviridae based on analysis of large-subunit terminase amino acid sequences encoded in

phage genomes. Bootstrap values are expressed as a percentage of 100 replications; scale bar,

mean number of amino acid substitutions per position; the tree is rooted at the midpoint. Clusters

match those presented in **Supplementary Figure B** for the VIRIDIC analysis.

Figure 3. Distribution of weberviruses. (a) vConTACT filtered gene-sharing network in which only nodes connected to the main cluster are shown (Fruchterman Reingold layout; filtered based on giant component; Gephi modularity options selected – randomness, use weights, resolution 1.0). The network comprises 34 modules. Only modules representing >3 % of all nodes in the network are coloured. Module 0 (represented by the circled orange and yellow nodes in the image) contained only the 127 webervirus genomes (*n*=67 isolated phages; *n*=60 MAGs). (b) Stacked bar graph showing the sources of the 127 webervirus genomes (*n*=67 isolated phages; *n*=60 MAGs). (c) Geographical distribution of 126 of the webervirus genomes included in this study (the location information was not available for one isolated phage, namely *Webervirus KL*).

Supplementary figures for Dawson *et al.*, Weberviruses are gut-associated phages that infect *Klebsiella* spp.

Supplementary Figure A. TEM images of (a) KLPN1, (b) vB_KpnS-KLPN2, (c) vB_KpnS-KLPN3, (d) vB_KpnS-KLPN4, (e) vB_KvaS-KLPN5, (f) vB_KvaS-KLPN6, (g) vB_KvaS-KLPN7 and (h) vB_KpnS-KLPN8.

Supplementary Figure B. Heatmap showing bidirectional clustering of VIRIDIC similarity data for the 67 webervirus genomes. The three numbered clusters at the side of the figure correspond to the three clusters detected by VIRIDIC. NC_028760.1 is the genome sequence for *Webervirus KLPN1*.

Supplementary Figure C. ViPTree analysis of 127 webervirus genomes with information (where known) provided for country of isolation of each phage.

Supplementary Figure D. MUSCLE amino acid alignment of depolymerases identified in this study. A high level of sequence conservation (indicated in green) was identified at the N-terminal region of all the depolymerase sequences (first 100 amino acids).