Type IV pili-associated secretion of a biofilm matrix protein from *Clostridium perfringens* that

forms intermolecular isopeptide bonds

Running title: Isopeptide bonds link *C. perfringens* biofilm proteins

Sarah E. Kivimaki[1*], Samantha Dempsey[1#], Collette Camper[1†], Julia M. Tani[1#], Ian K. Hicklin[2‡],

Crysten E. Blaby-Haas[3,4], Anne M. Brown[2], Stephen B. Melville[1]

1. Department of Biological Sciences, Virginia Tech, Blacksburg, VA 24061, USA

2. University Libraries and Department of Biochemistry, Virginia Tech, Blacksburg, VA 24061,

USA

3. Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

4. DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA, USA

*Correspondence:

Stephen Melville

melville@vt.edu

* Current address: Janelia Research Campus, 19700 Helix Dr., Ashburn VA, 20147

Current address: TechLab Inc., 2001 Kraft Drive, Blacksburg, VA 24060-6358 USA

† Current address: University of South Florida, Tampa Campus, 4202 E. Fowler Avenue, Tampa,

FL 33620

‡ Current address: University of California, Berkeley, 110 Sproul Hall #5800, Berkeley, CA 94720

Key words: *Clostridium perfringens*, biofilm matrix protein, protein secretion, isopeptide bonds

Abstract

Clostridium perfringens is a Gram-positive, anaerobic, spore-forming, bacterial pathogen of

humans and animals. *C. perfringens* also produces type IV pili (T4P) and has two complete sets of

T4P-associated genes, one of which has been shown to produce surface pili needed for cell

adherence. One hypothesis about the second set of T4P genes is that they comprise a system

analogous to the type II secretion systems (TTSS) found in Gram-negative bacteria, which is used

to export folded proteins from the periplasm through the outer membrane to the extracellular

environment. Gram-positive bacteria have a similar secretion barrier in the thick peptidoglycan

(PG) layer, which blocks secretion of folded proteins >25 kD. To determine if the T4P-associated

genes comprise a Gram-positive TTSS, the secretome of mutants lacking type IV pilins were

examined and a single protein, a von Willebrand A domain containing protein BsaC (CPE0517)

was identified as being dependent on PilA3 for secretion. The *bsaC* gene is in an operon with genes

encoding a signal peptidase and two putative biofilm matrix proteins BsaA and BsaB, with remote

homology to *Bacillus subtilis* TasA. We analyzed BsaA monomer interactions with *de novo*

modeling, which projected that the monomers formed isopeptide bonds as part of a donor strand

exchange process. Mutations in residues predicted to form the isopeptide bonds led to loss of

oligomerization, supporting an exchange and lock mechanism. Phylogenetic analysis showed the

BsaA family of proteins are widespread among bacteria and archaea but only a subset are predicted

to form isopeptide bonds.

Importance

For bacteria to secrete folded proteins to the environment, they have to overcome the physical

barriers of an outer membrane in Gram-negative bacteria and the thick peptidoglycan layer in

Gram-positive bacteria. One mechanism to do this is the use of a Type II secretion system in

Gram-negative bacteria, which has a structure similar to type IV pili and is modeled to act as a

piston that pumps folded proteins through the outer membrane to the environment. *Clostridium*

perfringens, like all or most all of the clostridia, has type IV pili and, in fact, has two sets of

pilus-associated genes. Here we present evidence that *C. perfringens* uses one set of pilus genes

to secrete a biofilm associated protein and may be responsible for secreting the main biofilm

protein, BsaA. We show that BsaA monomers are, unlike most other biofilm matrix proteins,

linked by intermolecular isopeptide bonds, enhancing the physical strength of BsaA fibers.

 Introduction

Bacterial protein secretion to the exterior environment involves transporting the protein across

several physical barriers. For Gram-negative bacteria these include the cytoplasmic membrane

(CM), a relatively thin peptidoglycan (PG) cell wall and the outer membrane. Gram-negative

bacteria have evolved several different mechanisms to achieve this, including Type II secretion

systems (TTSS), which have many proteins similar to those found in Type IV pili (T4P) (Fig. S1

and (1, 2)). TTSS are modeled to function as pistons that use a short pseudopilus to pump folded

periplasmic proteins through a protein channel (secretin) in the outer membrane (Fig. S1 and (3,

4)). Protein secretion in Gram-positive bacteria involves transport across the CM and a usually

thick PG layer. There are many protein secretion mechanisms for transiting the CM known for

Gram-positive bacteria, including the Sec system, Twin arginine translocation (Tat), Type IV

secretion systems, Esat-6 or type VII secretion system, flagella export apparatus (FEA), fimbrilin

(pilus) protein exporter (FPE), and holin-dependent translocation. The Type IV, Type VII and FEA

are able to translocate proteins across the PG layer (5, 6). However, proteins secreted through the

CM via the Sec system, used for the majority of proteins, need to fold and, once they do, the PG

layer of Gram-positive bacteria acts as a barrier to secretion to the environment because the mesh

size in the PG layer is too small to allow diffusion of globular proteins >25 kD, as determined by

measurements of the PG from the Gram-positive bacterium *Bacillus subtilis* (7).

One possible mechanism for secretion of folded proteins from the space between the CM and the

PG layer in Gram-positive bacteria would be a system analogous to TTSS in Gram-negative

bacteria. Our discovery that T4P systems are ubiquitous and perhaps universal in all of the

Clostridia (8) provided a means for a potential TTSS to exist in the Clostridia. We hypothesized in

a review (9) that in Clostridial species with two separate T4P systems, one of them could function

as a TTSS to transport proteins through the PG layer. Multiple species of Clostridia do carry

multiple copies of complete T4P systems, including *C. perfringens* (Fig. 1) and *Clostridioides*

difficile (9). The core proteins needed for functional T4P usually include pili, an assembly ATPase

(PilB), an inner membrane core protein (PilC), and associated assembly proteins PilM, PilN, and

PilO (9). *C. perfringens* carries two homologs of PilB, two homologs of PilC and single copies of

PilM, PilN, and PilO (Fig.1), making it plausible that one of these systems could function as a

TTSS. To test this hypothesis, we examined the secretome of strains with in-frame deletions in

genes encoding Type 4 pilins; in *C. perfringens* these are designated as PilA1-PilA4 (Fig. 1). We

noticed a protein of ~70 kD was present in the WT secretome but absent in the secretome of one

deletion strain, *pilA3* (Fig. 2A). Mass spectrometry identified the 70 kD protein as CPE0517,

which was encoded by a gene in an operon with three other genes, including a SipW signal

peptidase homolog and two biofilm matrix proteins, CPR0515 (BsaA) and CPE0516 (BsaB) (Fig.

2C). We expressed the entire operon in *C. perfringens* and found that it produced large amounts of

biofilm matrix proteins (Fig. S2B), and that the biofilm matrix protein (CPE0515) was present as

a long oligomer and was resistant to boiling in SDS-PAGE buffer (Fig. S2D). We also noted that

CPE0515 had some predicted structural homology to the TasA protein of *B. subtilis*, which is also

a biofilm matrix protein that uses a SipW signal peptidase for secretion (10-12). At around this

same time, a paper by Obana et al (13) was published in which they examined many features of

the same operon, which they designated as *sipW-bsaA-bsaB-bsaC*. While the majority of their

work concerned the regulation of matrix protein synthesis, they discovered some important

physical features of the biofilm matrix protein (BsaA, Fig. 2C): it is the main component of the

biofilm matrix; it required SipW for signal processing and secretion; expression in *E. coli* of a

form of BsaA lacking a signal sequence led to spontaneous oligomerization of BsaA, which was

resistant to denaturation by SDS or formic acid, indicting the oligomers were held together by very

strong intermolecular interactions (13).

In this report we describe the process of identifying T4P proteins involved in secreting CPE0517

(BsaC), the 70 kD protein described above, as well as pilin proteins involved in secreting the BsaA

oligomers. We also propose an "exchange and lock" mechanism involving isopeptide bond

formation that is responsible for the stability of the BsaA oligomer. We characterized the nature

of these bonds and conservation of isopeptide bond formation based on phylogeny of the

BsaA/TasA family of proteins. Identifying a potential TTSS-like system in a Gram-positive

bacterium and the novel finding of isopeptide bond formation in a well-studied family of proteins

will lead to significant advances in both of these emerging fields of microbial cell biology.

137   **Results**

Secretome analysis of a *pilA3* mutant indicates a 70 kD protein with a Von Willebrand

domain A (CPE0517, BsaC) is absent. We screened the secretome of in-frame deletion mutants

in four pilin-encoding genes, *pilA1*, *pilA2*, *pilA3*, *pilA4* for missing proteins, indicating they may

be dependent on the pilin for secretion. Only the *pilA3* mutant showed evidence of a distinct

missing band in the secretome and this band was restored by complementation of a WT copy of

the *pilA3* gene (Fig. 2A). The band was excised from a gel and subjected to trypsin digestion and

analysis by mass spectrometry. Peptide fragments matched those of the CPE0517 (BsaC) protein

from the *C. perfringens* strain 13 genome. The protein has an unknown function but does contain

a von Willebrand A domain, which can potentially function in protein-protein interactions with

host cells. This gene is the fourth gene in a four gene operon containing a type I signal peptidase

(CPE0514, SipW) and two genes (*CPE0515-CPE0516*, *bsaA-bsaB*) encoding proteins with

similarity to the TasA protein in *B. subtilis* (Fig. 2C). Western blots using anti-BsaC antibodies

showed a similar loss of CPE0517 secretion by the *pilA3* mutant and complementation by

expression of a WT copy of the *pilA3* gene (Fig. 2B).

To determine if other Type IV pilus (TFP) related genes were needed for secretion of BsaC, we

expressed the gene in a bank of in-frame deletion mutants encompassing each gene related to

synthesis and assembly of TFP (Fig. 1) using a vector we developed containing a lactose-inducible

promoter (14). We noted variation in day-to-day results with the WT strain and, since these were

western blots on TCA precipitated protein from culture supernatants, we normalized all of the data

from mutant strains to that of the WT strain on the same blot (Fig. 3). Using quantitative

densitometry on western blots of the secreted oligomers, we measured statistically significant

decreased secretion of BsaC in the following mutants: *pilA3*, *pilB2*, *pilC1*, *pilD*, *pilM*, *pilN* and

pilO (Fig. 3). With the exception of *pilC1*, these results suggest efficient secretion of BsaC requires

components of the TFP system present in the large *pilD-CPE2277* operon in *C. perfringens* (Fig.

1 and (8, 9)). Whether these form a type IV pilus with PilA3 as the pilin or a Gram-positive TTSS

remains to be determined. However, a major difference in the structures of T4P and TTSS is the

type IV pilus extends beyond the outer membrane (Gram-negative) or PG layer (Gram-positive,

(8)) but in the TTSS in Gram negative bacteria the pilus does not (2). Our model for a TTSS in

Gram-positive bacteria is that the pilus (i.e., PilA3) does not extend past the PG layer (Fig. S1).

To test this model, we exposed *C. perfringens* cells to rabbit anti-PilA3 antibodies and a

fluorescently tagged goat anti-rabbit secondary antibody and found that we could detect binding

localized to the poles of the cells (Fig. S2). This refined the location of the pili, and that they extend

far enough out past the PGY layer to be exposed to antibodies.

To determine if there are generalized secretion defects in TFP mutants, we examined

whether each mutant in the TFP-associated genes were able to secrete another protein, the

phospholipase-sphingomyelinase, PLC. Secretion of PLC can be easily detected by a ring of

precipitation around a colony using egg yolk agar, and we found that none of the TFP mutants

showed a detectable decrease in PLC secretion (Fig. S3), suggesting the secretion defects seen

with BsaC were not due to generalized secretion defects.

Expression of the *bsa* operon leads to synthesis of extracellular biofilm matrix material. Since

the *bsa* operon encodes proteins with similar synteny and predicted structural homology to the

sipW-tasA-tapA biofilm matrix operon in *B. subtilis* (12), we placed the entire operon (*sipW-bsaA-*

bsaB-bsaC) under control of the lactose inducible promoter for high level expression. The cells

produced large quantities of an extracellular gel like material (Fig. S4B), which, after staining with

crystal violet, appeared as amorphous material between the bacterial cells (Fig. S4C). A His_6 tag

was added to the C-terminus of the BsaA protein and, when the operon was expressed, a distinct

high molecular weight ladder of BsaA was visible in anti-His_6 antibody western blots after running

on SDS-PAGE gels using samples boiled in SDS-PAGE buffer (Fig. S4D). This suggested BsaA

was in very heat-stable oligomers. This was confirmed by the results of Obana et al who found

that BsaA was the major component of the biofilm matrix and that BsaA oligomers were stable

after boiling in SDS-PAGE buffer and resistant to dissociation by high levels of SDS and formic

acid (13). These results suggested BsaA oligomers were in an extremely stable conformation which

may include covalent bonds. However, these stable oligomers were not due to the activity of the

sortase-encoding gene *srtB* (Fig. 2B), since a mutant lacking that gene still produced highly stable

oligomers of BsaA (13).

Efficient secretion of the BsaA oligomer is dependent on some TFP-associated genes. Since it was possible that, like BsaC, BsaA in its oligomer form was secreted with the aid of TFP-associated proteins, we expressed a clone containing just the *sipW* and *bsaA* genes in which a FLAG-tag was added to the C-terminus of BsaA for quantification in western blots using densitometry. An example of a typical blot is shown in Fig. S5A. The WT strain showed significant levels of secretion of the BsaA oligomer (Fig. S5B). Using this method, we found that the *pilA1*, *pilB1*, *pilB1/pilB2*, *pilN* and *pilO* mutants exhibited decreased levels of BsaA oligomers in the supernatants (Fig. S5B). However, since this profile of T4P-associated genes lacked an inner membrane core protein (*pilC*) component, which is essential for TFP functions (15) the results were not definitive. Therefore, we used a quantitative slot blot method as an alternative.   In this experiment, culture supernatants were serially diluted in medium and passed through a membrane, and probed with an anti-FLAG antibody (Fig. S6A). The densitometry readings were then plotted to identify an amount of signal that was in the linear range for quantification (Fig. S6B). However, using this method only the *pilA1*, *pilA2* (higher), and *cpe1841* (higher) mutants showed a statistical difference from the WT strain (Fig. S6C), which does not comprise a full T4P system.

Models of BsaA oligomers predicts an isopeptide bond is formed linking adjacent monomers. BsaA forms very heat and SDS resistant oligomers ((13) and Figs. S4D and S5A). To investigate the structural nature of the interactions between monomers of BsaA that leads to this stability, a structure would need to be developed of a homomultimer. Due to the difficulty of solving the structure of long, thin chains by experimental methods, computational modelling was used. Models were initially created using Phyre2 (16), with the top model being an x-ray crystal structure of TasA from *B. subtilis* (17). That structure, and homology models from that structure, were monomeric, and not useful for determining intermolecular interactions between monomers.

AlphaFold Multimer was then used to create a multimeric model to determine inter-chain

interactions. Using ColabFold 1.3.0 (18), which is based on AlphaFold 2.2.0 (19), a homotrimeric

structure of BsaA was created (Fig. 4A). The SipW signal sequences (residues 1-27) were included

in the initial model, but only formed disordered loops which do not have any notable interactions.

Removing the signal sequences did not change the overall structure of the monomers so this was

done for subsequent analyses.

In monomeric structures, residues 28-38 would extend out as disordered loops (Fig. 4A,

monomer 1). In the multimeric structures, these residues incorporated themselves into the beta

sheet structure of adjacent monomers (Fig. 4A). Being fully incorporated into a nearby structure

indicates that residues 28-38 provide specific interactions with the receiving monomer, providing

the basis of a mechanism for which oligomers of BsaA can be created. For example, V33 and F37

of the interdigitated strand become part of the hydrophobic core of the receiving monomer (Fig.

4B). The interactions between specific side chains of residues from the interdigitated strand and

the receiving monomer were investigated further for less standard interactions.

The *de novo* created structures indicate a bond between two side chains: N35 of the

interdigitated strand and K74 of the receiving monomer (Fig. 4C). At first, this was thought to be

an error in AlphaFold's model generation, since the chemistry would not be possible. The bond

was created between the oxygen of N35 and the nitrogen of K74, resulting in three bonds in the

oxygen. However, this bond was created in nearly every model generated, so it was investigated

further. A crosslink between asparagine and lysine is a type of isopeptide bond, which has been

characterized through a few different mechanisms and can be found as an intermolecular crosslink,

as in ubiquitylation (20), or as an intramolecular crosslink found in Gram-positive pili (21). In

Gram-positive pili, this bond is formed between an asparagine and a lysine, autocatalyzed by a

glutamate within the hydrophobic core of the protein. Residues equivalent to these are found

surrounding the bond in BsaA (Fig. 4C). The hydrophobic region surrounding the N35-K74 pair

(Fig. 4B, brown colored residues) is believed to be necessary to help the catalytic cross-linking

reaction (22).

It is unclear why this bond was created by AlphaFold. In the documentation of AlphaFold Multimer

(23), there are no mentions of predicting crosslinks. One potential reason is that it generated the

side chains so close to each other there was a steric clash, but the rest of the model was so favorable

the structure was created regardless. Models created in ColabFold have an option to perform

energy minimization using Amber. For these models, this was not performed. Recreating this

model while enabling energy minimization eliminates the bond every time. This is expected given

the incorrect chemistry of the bond. AlphaFold3 (24) does not include a minimization step, and,

like AlphaFold2, creates the isopeptide bond.

This amyloid-like fold and interdigitation is seen in other Gram-positive bacteria, such as

TasA in *B. subtilis*. A cryo-EM structure has been created for a chain of TasA (10), which

demonstrated how hydrophobic interactions between donor and acceptor monomers drive

interdigitation. No crosslinks were found or mentioned, and the required triad of N-E-K residues

is not present in the structure (PDB 8aur). Therefore, the isopeptide bonds in BsaA oligomers

would be the first example of an intermolecular bond of this mechanism found in this family of

proteins.

Mutations in isopeptide bond forming residues abolish oligomerization of BsaA. The

AlphaFold Multimer models predicted an isopeptide bond was formed between N35 and K74 from

adjacent residues in a BsaA oligomer. To test this computational prediction, we made point mutant

substitutions of each residue to alanine, forming N35A and K74A mutants, along with a double

N35A/K74A mutant. His-tagged versions of these mutant forms were expressed along with *sipW*

and the cell pellets and culture supernatants were examined for the presence of the mutant proteins

in oligomeric form. Mutating N35 or K74 to alanine residues resulted in a loss of oligomers in the

supernatants and the appearance of monomers of the K74A mutant but not of the N35A mutant,

suggesting the N35A version was not effectively released by the cell (Fig. 5A). Monomeric forms

of both mutants appeared in the cell pellet of some samples (Fig. 5A). In some experiments, small

amounts of dimers and higher mol wt forms of BsaA appeared in the cell pellet, but the large

majority were in monomeric form, which was absent in the WT form of BsaA (Fig. 5A), perhaps

due to the stability of oligomers even in the absence of an isopeptide bond. To test this, we

suspended cell pellet extracts from BsaA-His$_6$ and N35A-His$_6$ in SDS-PAGE buffer and heated at

95 C for 0, 10, 20 and 40 min before running the samples on SDS-PAGE gels and performing

western blots. The N35A-His mutant maintained stable oligomers in SDS-PAGE buffer and after

being run through an SDS-PAGE gel in the absence of heating, but heating the samples for 10-40

min led to dissociation of the oligomers, in contrast to the WT BsaA protein, which was not

affected by heating (Fig. S5). This suggests that N35, and therefore the isopeptide bond, is essential

for forming detergent and heat resistant oligomers, and that the monomer-monomer interactions

are stable even in the presence of 2% SDS (the concentration in the SDS-PAGE buffer), which is

in contrast to oligomers formed by the *B. subtilis* TasA protein, which were dissociated by SDS

(11).

We also tested the N35A/K74A mutant and observed that it showed a similar loss of

oligomerization as did the N35A mutant (Fig. 5B, left four lanes). These experiments were all done

in strain HN13 (WT) background where there was an intact *bsaA* gene. To determine if this

potential source of BsaA, even though it was not His-tagged, affected the results we observed with

the mutants, we expressed the N35A and N35A/K74A mutants in a strain with an in-frame deletion

of the *bsaA* gene, but did not observe any differences to the pattern seen in the strain with an intact

bsaA gene (Fig. 7B, right four lanes), suggesting the chromosomal source of BsaA was not forming

oligomers in combination with the N35A and N35A/K74A mutant forms of BsaA.

The Glu56 residue of BsaA is essential for efficient isopeptide bond formation. Spontaneous

or self-catalyzed isopeptide bond formation is catalyzed by an acidic Glu or Asp residue located

within the hydrophobic pocket where the Asn and Lys residues are located (22). In AlphaFold 2

BsaA models of dimer interfaces, Glu56 is predicted to be in the precise location to catalyze bond

formation (Fig. 4B). To test this was actually occurring, we constructed an E56A substitution in

the BsaA protein and examined its ability to oligomerize by using western blotting on cell extracts

and concentrated supernatants from the mutant and WT strain (Fig. 5C). The E56A mutant showed

a clear lack of oligomer formation and an accumulation of the monomeric form of the protein

which was not observed with the WT form of BsaA (Fig. 5C), providing strong evidence that

Glu56 is responsible for catalyzing isopeptide bond formation in BsaA.

A FAST fusion to BsaA displays surface exposed localization. To investigate the cellular

location of the BsaA oligomers, a FAST gene fusion to the *bsaA* gene was constructed and, along

with the *sipW* gene, was placed under control of the lactose-inducible promoter. The FAST protein

binds a soluble dye which, upon binding, exhibits a large increase in fluorescence (25-27) allowing

localization of the fusion proteins. We expressed the *sipW-bsaA*-FAST fusion genes and exposed

the cells to two different dyes, Coral, which is membrane permeable, and Amber-NP which is

impermeable to membranes (both obtained from The Twinkle Factory). To test if the Coral and

Amber-NP permeability profiles were working in *C. perfringens* membranes, we expressed a *pilT*-

FAST gene fusion, which should be located only in the cytoplasm, and exposed them to the Coral

and Amber-NP dyes. Consistent with the predicted cytoplasmic location of the *pilT*-FAST fusion,

only the Coral dye showed significant fluorescence after treatment (Fig. 6A). However, when the

bsaA-FAST fusion was expressed along with *sipW*, both dyes were able to bind and fluoresce (Fig.

6B), indicating the BsaA-FAST fusion was competent for secretion in *C. perfringens* and was

located on the surface of the bacteria.

BsaA oligomers are anchored to the membrane and not the peptidoglycan in strain HN13.

Because the BsaA-FAST fusion experiments indicated BsaA oligomers were located on the cell

surface and SEM images of the BsaA matrix proteins suggested they were in the form of fibrils

associated with the bacterial surface (13), we wanted to determine if the oligomers were embedded

in the cytoplasmic membrane or covalently linked to the PG layer. To test this, we expressed *sipW-*

bsaA-His$_6$ constructs for the WT and each mutant in our lactose-inducible promoter system and

then boiled the cells in 10% SDS for 30 min, followed by digestion of the PG matrix by lysozyme.

Our experimental approach was based on the hypothesis that if BsaA was anchored in the

membrane, boiling in SDS would release it from the cell but if it was covalently anchored to the

PG, it would be freed only after digestion with lysozyme. We tested the WT, N35A, K74A and

N35/K74 mutants and found that all of the long wild-type oligomers were released after the SDS

boiling step, while the cells appeared to retain a small number of monomers and dimers of BsaA

(Fig. S4). However, after treatment with lysozyme, the monomers and dimers were no longer

detected. For the mutants that can't oligomerize, the monomer/dimers were retained after boiling

in SDS, but not after lysozyme treatment. We interpret these findings as indicating the oligomers

are anchored to the membrane but after SDS boiling they are released. A small number of

monomers and dimers remain trapped inside the sacculi produced by boiling in SDS but lysozyme

treatment releases these from the sacculi. However, they are then too dilute to detect on western

blots. This is consistent with an analysis of the *bsaA* and *bsaB* protein sequences, both of which

lack a recognizable LPXTG motif, despite the presence of a sortase gene, *srtB*, adjacent to the

operon (Fig. 1C).

BsaB, an ortholog of BsaA, forms dimers but not long oligomers. Previous results by Obana et

al (13) showed a *bsaB* mutant still made significant amounts of the BsaA oligomers but they did

not test to see if BsaB was incorporated into BsaA oligomers. Therefore, we added an HA-tag to

the BsaB protein and expressed the *sipW-bsaA-bsaB*-HA-*bsaC* genes in our inducible system and

examined the culture supernatants for BsaB-HA using western blots. Based on the predicted

molecular weight of the BsaB protein, we detected BsaB-HA only in a monomeric and a higher

molecular weight form which could be either a homodimer or a BsaA-BsaB heterodimer (Fig. S6),

but it is not covalently attached to the longer BsaA oligomers.

Phylogeny of isopeptide bond formation in the TasA family. Since a BlastP search with BsaA

only identifies homologs in closely related Clostridial species, an iterative search with profile

hidden Markov models was employed. This analysis resulted in the identification of 2,172 proteins,

including the previously described TasA from *B. subtilis* and CalY from *Bacillus cereus* (Fig. 7).

Of these, 632 hits belong to the "peptidase M73, camelysin" (InterPro: IPR022121) family; so-

called because of the likely mis-identification of CalY as a peptidase. In addition to the use of

probabilistic models, the relatedness of these proteins is evident at the structural level. Like other

family members, BsaA is predicted to adopt a β-sheet–rich Ig-fold-like structure and is predicted

to participate in donor-strand exchange (Fig. 4), as described for TasA (10). These analyses suggest

that while the structures of these proteins and homopolymer assembly are highly conserved,

sequences are highly divergent.

In addition to BsaA, we identified three other uncharacterized proteins in *C. perfringens*

(BsaB, CPE2262 and CPE0554), but these paralogs are not expected to form an intermolecular

isopeptide bond, because the N-E-K triad is missing. Nevertheless, over 100 BsaA-like proteins in

other species were identified with the N-E-K triad, suggesting conservation of the donor-strand-

exchange-and-lock mechanism. These proteins are largely from Clostridia with some homologs

from *Coprobacillaceae*, *Enterococcaceae*, and *Thermococcus* (Fig. 7).

Based on the putative functions of gene neighbors, the BsaA/TasA family members appear

to be involved in the assembly of surface filament structures (Fig. 7C). They are often in putative

operons with genes encoding proteins containing the Gram-positive cell wall anchor motif LPxTG,

fibronectin domains, SpaA domains, choice-of-anchor A domains, as well as signal peptidase and

sortase domains. Multiple BsaA/TasA paralogs can be found in a single putative operon (Fig. 7C),

suggesting the possibility of donor-strand exchange assembly of heteropolymers.

366   **Discussion**

The original goal of this work to determine if Gram-positive bacteria have the functional

equivalent of TTSS that are found in Gram-negative bacteria. Since TTSS have a similar overall

architecture and mechanism as T4P (Fig. S1), it made sense to test a Gram-positive bacterium

known to have T4P. With a few notable exceptions (e.g., *Streptococcus sanguinis* (28-30))

amongst Gram-positive bacteria, T4P are confined within the Clostridia (9) so *C. perfringens* was

chosen as representing a typical clostridial species. The overall concept was that the thick Gram-

positive PG layer represents a similar barrier to protein secretion to the environment as the Gram-

negative outer membrane. Since TTSS depend upon a pseudopilus to function, we tested mutants

with deletions in each of the four main pilin-encoding genes in *C. perfringens* for changes in the

secretome compared to a WT strain. We identified one protein BsaC (CPE0517) that was depleted

in the secretome of a *pilA3* deletion strain (Fig. 2A-B). We then examined a bank of in-frame

deletion mutants in all of pilin-associated genes (Fig. 1) and found that, for BsaC secretion, a

homolog of each of the essential proteins needed for a type IV pilus assembly, PilA3 (pilin), PilB2

(assembly ATPase), PilC1 (inner membrane core protein), PilD (prepilin peptidase), and PilM-N-

O, (inner membrane accessory proteins) (Fig. 1) were required for efficient secretion (Fig. 3). This

appears to be specific for BsaC secretion, since the secretion of the toxin PLC was not affected by

mutations in any of the T4P-associated genes (Fig. S2). While this may indicate the presence of a

TTSS like apparatus in *C. perfringens*, the lines separating T4P- and T2SS-dependent secretion

have been blurred by the discoveries that *Pseudomonas aeruginosa* and *Dichelobacter nodosus*,

Gram-negative bacteria, have T4P systems capable of mediating protein secretion (31, 32).

However, further analysis of the structure and components of these Gram-negative pilus-dependent

secretion systems have not been published. PilA3 pilin was detected on the surface of *C.*

perfringens, so it is possible the PilA3-dependent secretion system acts more like a T4P.

BsaC monomer secretion may appear to be independent of BsaA oligomers, but it is

possible that they are being secreted simultaneously, since we did not track BsaA and BsaC

secretion at the same time. The biological role that BsaC plays in biofilm matrix functions is

unknown, but it is intriguing that the protein contains a VWA domain, which is widespread in

eukaryotes, bacteria and archaea and is associated with protein-protein interactions (33).

Therefore, the role BsaC plays in biofilm functions is also currently under investigation in the

laboratory.

BsaC is in an operon with genes encoding a signal peptidase, SipW and two proteins (BsaA

and BsaB) with homology to the main biofilm protein of *B. subtilis*, TasA (Fig. 2C). Because this

gene synteny is similar to the *sipW-tasA-tasB* biofilm operon in *B. subtilis*, we expressed the entire

operon to see if a biofilm matrix was produced and noted that high levels of a gel like matrix

material was formed in the culture supernatant (Fig. S3B-C). A high mol wt oligomer of BsaA

was seen in SDS-PAGE gels (Fig. S3D), which was similar to that seen by Obana et al in their

analysis of the same operon (13). Because BsaC was dependent on T4P-associated proteins for

secretion, it was logical to determine if secretion of the BsaA oligomer was as well. Two different

methods were tried, using densitometry of western blots of the BsaA oligomers in culture

supernatants and densitometry to measure the amount of BsaA in the culture supernatants using a

slot blot device. Although we saw significantly decreased levels of secretion in mutants using both

methods (Fig. S4 and S5), neither one gave conclusive evidence that a full T4P system was needed

for secretion. Both experimental approaches were hindered by relatively high amounts of

experimental variability in trying to measure an oligomer in solution; this can be seen by the high

range of the SEM determined for most of the samples (Fig. S4 and S5) despite multiple repetitions

of each experiments. However, the fact that several genes in each type of experiment was shown

to be necessary for efficient secretion, does strongly suggest that T4P proteins play a role in the

process.

Although we observed that a *pilA1* deletion strain did not secrete the biofilm matrix

oligomers efficiently (Fig. S4 and S5), we did not observe this phenotype in our initial screening

for proteins absent in the secretome, as shown for PilA3 in Fig. 2A-B. We think this is likely due

to the fact that our growth conditions for the experiment, overnight culture in BHI liquid medium

at 37 ºC, is not optimum for production of the biofilm matrix proteins, as described by Obana et al

(13). These findings point out the limitations of screening for T4P-dependent secretion: using only

a single type of environmental condition may not be the one in which a protein is made and secreted

so it would not be discovered. Even 2-D gels would be unlikely to have uncovered the presence of

the BsaA oligomers due to their extremely high mol weight. Therefore, it seems protein-based

mass spectrometry on samples from a wide variety of environmental conditions would be optimal

for detecting T4P-dependent secretion substrates.

Since our results and that of Obana et al both showed that oligomers of BsaA were

extremely resistant to heat and other denaturants (Fig, S3D and (13)), this suggested that covalent

bonding might be involved. Subsequent analysis using AlphaFold 3 indicated there were likely

isopeptide bonds being formed by donor strand exchange involving an N35 of one monomer and

a K74 of another, catalyzed by an E56 residue in the recipient monomer (Fig. 4). This model was

confirmed by subsequent substitution of each residue involved with an Ala, which led to a

complete lack of oligomerization (Fig. 5). However, isopeptide bonds are formed relatively slowly

(22) so we hypothesized that the donor strand exchange resulted in formation of a temporally stable

interaction that gave the isopeptide bonds time to form. However, based on experiments with the

N35A mutant, the oligomers are stable even without the isopeptide bond, as the mutant version

was resistant to denaturation in SDS-PAGE buffer and being run through an SDS-PAGE gel. Only

after heating in the SDS-PAGE buffer at 95 ºC did the N35 mutant depolymerize (Fig. S6), while

the WT oligomer withstood this treatment. We further characterized the formation of the oligomer

and showed, using FAST fluorescent protein fusions, that the oligomers were anchored to the

surface of the bacteria (Fig. 6) but via the cytoplasmic membrane and not directly to the PG layer

(Fig. S7).

BsaB, the other biofilm protein encoded in the operon, did not form oligomers, did not

become covalently linked to BsaA oligomers (Fig. S8) and both BsaA and BsaB have no apparent

LPXTG motif present in their protein sequence (data not shown). These structural features seem

somewhat at odds with the presence of a sortase gene (*srtB*) adjacent to, but oriented in the opposite

direction, of the *sipW* gene (Fig. 2C). Obana et al (13) tested the premise that SrtB was responsible

for oligomerization of BsaA, but found that a *srtB* mutant had normal levels of oligomer present

(13). However, examination of the *bsa* operon in other strains of *C. perfringens* reveals that a

different gene synteny is in place, where *sipW* is followed by a *bsaA* homolog and two *bsaB*-like

genes, the first of which carries an identifiable LPXTG motif (Fig. 2D). The BsaB sequence from

strain 13, when subjected to a BLAST analysis (34) against other *C. perfringens* strains, shows

proteins with high levels of homology to either the N-terminal half or C-terminal half of the strain

13 BsaB, but not both. This suggests strain 13 underwent a genetic rearrangement event in which

the two *bsaB* homologs were recombined into a single gene with the loss of the LPXTG motif

found in the first gene (Fig. 2D). This brings up the distinct possibility that the SrtB sortase could

catalyze the cross-linking of the first BsaB ortholog to the PG layer in strains other than strain 13.

What role BsaB plays in BsaA oligomerization is still unknown, since a *bsaB* mutant still formed

similar amounts of oligomers of BsaA as the wild-type strain (13). However, based on the

differences in composition of BsaB homologs between strain 13 and other strains (Fig. 2C-D), it

is possible that BsaB is involved in anchoring the BsaA oligomers to the PG layer via the activity

of SrtB and the BsaB homolog with an LPXTG motif, but this is speculative.

Taking all of these results into consideration, we have constructed a model showing the

mechanisms underlying the assembly of BsaA into cross-linked oligomers (Fig. 8). In this model,

BsaA monomers are secreted via the Sec pathway, fold on the outer surface of the CM but are

anchored in the membrane by their N-terminal signal sequence. An oligomer of BsaA with the

most recent monomer having its N-terminal disordered region exposed, contacts the newly

secreted monomer and undergoes a donor strand exchange in which the N-terminal disordered

region binds to the exposed beta-sheet fold and interdigitates as a newly formed beta sheet (Fig.

4A). This is followed by cleavage of the signal sequence by the signal peptidase SipW. We believe

the SipW acts after oligomer binding to the newly secreted monomer because the BsaA oligomers

appear to be anchored to the membrane (Fig. S7) and the oligomers are in a relatively stable

assembly even before the monomers are "locked" together by isopeptide bond formation (Fig. S6).

After SipW cleavage, the oligomer is released and can then be used to add another monomer or be

secreted through the PG layer to the external medium. The presence of significant amounts of

oligomers attached to the bacterial surface (Fig. 6) suggests the monomers assemble in the space

between the CM and PG rather than being secreted as monomers for oligomerization in the

environment. The mechanism for export of the oligomers through the PG layer is not complete but

there is evidence that T4P play at least some role in the process (Fig. S4 and S5).

The phylogeny of BsaA-like proteins shows they are widespread amongst Gram-positive

bacteria but also present in some Gram-negative bacteria and archaea (Fig. 7). However, the

homologs that are predicted to form isopeptide bonds are more narrowly focused in the Clostridia

and Enterococcus genera (Fig. 7). This brings up the question as to what advantage isopeptide

bond formation bestows on the biofilm matrix properties of the bacteria that form them. Formation

of isopeptide bonds can greatly increase the mechanical resistance to distension and separation of

the BsaA fibers. As an example, the intramolecular isopeptide bonds formed in sortase-dependent

pili made by Gram-positive bacteria have been shown to greatly increase the amount of force that

needs to be applied to distend the individual subunits, thereby strengthening the entire fiber (35).

To trap macrophages, neutrophils or other phagocytic cells, *C. perfringens* and other species may

have evolved a mechanism to form isopeptide bonds to increase the mechanical strength of the

fibers, since macrophages, neutrophils and other phagocytic cells can impart a large locomotive

force (1.9 to 10.7 nN) during their motility (36). It is also worth considering that *C. perfringens* is

an extremely ubiquitous bacterium in nature, found in soil, freshwater sediments and even extreme

environments like Antarctica (37, 38) and likely forms biofilms in these natural environments. The

BsaA biofilm matrix fibers could protect them from killing by amoeba, protozoa and other

predators found in these environments. Analyzing the relative mechanical strength of fibers with

and without isopeptide bonds will help answer some of these questions.

498   **Materials and Methods**

Bacterial strains and growth conditions. Bacterial strains, plasmids, and primers used in this

study are listed in Table 1. *Escherichia coli* strain DH10B was grown in Luria Bertani broth at

37°C for all transformations. When necessary, kanamycin and chloramphenicol were added to the

media at a concentration of 100 μg/ml and 20 μg/ml, respectively. *C. perfringens* strain HN13, a

Δ*galKT* derivative of strain 13, was used as the wild type strain in this study. *C. perfringens* strains

were grown anaerobically in PGY (30 g proteose peptone #3, 20 g glucose, 10 g yeast extract, 1 g

sodium thioglycolate per liter) or brain-heart infusion (BHI) (Thermo Fisher) in an anaerobic

chamber (Coy Laboratory Products, Inc.). Anaerobic egg yolk agar medium was prepared as

previously   described   (https://www.fda.gov/food/laboratory-methods-food/bam-media-m12-

anaerobic-egg-yolk-agar#).

Construction of in-frame deletions of TFP-associated genes. In-frame deletions of TFP-

associated genes were made using the method of Nariya et al. (39), modified as described in

Hendrick et al (40). The primers used to amplify the flanking DNA for each gene are listed in

Table S1. All deletions were confirmed by PCR across the deleted region. Complementation of

the in-frame *pilA3* mutant (Fig. 2A-B) was done using the lactose-inducible promoter vector

pKRAH1 (14) to express the *pilA3* gene. The construct was made by PCR using the primers listed

in Table S7.

SDS-PAGE and staining of secretome proteins. Cultures of strain HN13 were grown for 16 h

in BHI medium (i.e., into stationary phase) and the bacteria were removed by centrifugation in a

table top centrifuge. The culture supernatants were removed and passed through a 0.45 μM filter.

Proteins in the culture medium were precipitated using 10% trichloroacetic acid (TCA)

precipitation and washed twice in acetone. After drying, sample pellets were resuspended in water

and then 4X SDS-PAGE sample buffer (41) was added to a final concentration of 1X. Secretome

samples were separated on an SDS-PAGE gel and then stained with the fluorescent protein stain,

Sypro-Ruby according to the manufacturer's instructions (Thermo Fisher). Stained gels were

imaged with a Gel-Doc Imaging system (Bio-Rad, Inc).

Cloning the *bsa* operon into the lactose-inducible expression vector, pKRAH1. Because the

SipW signal peptidase is likely needed for processing the secretion signal on BsaC (13), to analyze

secretion of BsaC in TFP gene associated mutants, the entire *sipW-basA-bsaB-bcaC* operon was

expressed under control of the lactose inducible promoter in plasmid pKRAH1 (14) using primers

OHL145 (PstI site) and OHL146 (BamHI site) by PCR (Table 1 and Table S3). The PCR fragment

was ligated to the PCR cloning vector pGEM-T to make pHLL64.   pHLL64 and pKRAH1 were

digested with PstI and BamHI and ligated to make pHLL65.

Construction of a FAST protein fusion to BsaA. The gene fusion was constructed using

overlapping PCR of the FAST coding region on plasmid pAG104 FAST (25) to the *bsaA* gene

linked to *sipW* (i.e., the *bsaB* and *bsaC* genes were absent). The FAST protein was fused to the C-

terminal end of BsaA because of the processing of the signal sequence on the N-terminal end. The

primers used to construct the gene fusion are listed in Table S5.

Production of antibodies to BsaC, PilA3 and immunofluorescence microscopy. Rabbit polyclonal antibodies against the BsaC peptide sequence N-CNIEGDKISWDRGEN-C were produced by GenScript USA, Inc. along with pre-immune serum as a negative control. Rabbit polyclonal antibodies against the PilA3 peptide sequence N-CIKIDNSKDSIDLTR-C were produced by Yenzyme Antibodies, LLC. along with pre-immune serum as a negative control. To test for surface exposure of PilA3 pilins, the bacteria were fixed with 2.5% paraformaldehyde in Dulbecco's phosphate buffer saline (DPBS), incubated with rabbit serum with anti-PilA3 antibodies, washed with DPBS and incubated with goat anti-rabbit antibodies fluorescently tagged with Alexafluor 647 (Invitrogen). Stained cells were imaged on an Olympus IX71 fluorescent inverted microscope equipped with a CoolSnap HQ2 CCD camera and DeltaVision deconvolution and image analysis software.

Quantitative immunoblotting of secreted BsaC. Overnight cultures were grown in BHI supplemented with chloramphenicol, and diluted 1:50 into fresh BHI medium pre-incubated at 37°C. The cultures were grown for one hour then induced by adding lactose, to a final concentration of 10 mM, to the growth media every 30 min for 3 h. Following induction, 1 mL of culture supernatant was collected by centrifugation, and proteins precipitated using TCA, as described above. Resulting precipitates were resuspended in 50 uL of SDS-PAGE loading buffer with 100 mM DTT, the pH was adjusted using one μL of 1.5 M Tris-HCl, pH 9, if necessary. Five μL of sample was loaded per well, and SDS-PAGE was performed. Following SDS-PAGE, proteins were transferred to a polyvinylidene difluoride (PVDF) membrane utilizing the iBlot system (Bio-Rad, Inc), and developed with serum containing anti-BsaC antibodies at 1:250 dilution, followed by a DyLight 550 labeled fluorescent secondary goat anti-rabbit antibody

(Thermo-Fisher). Imaging was performed on a Typhoon gel scanner, and mean pixel intensity

was measured for each band utilizing the Amersham Typhoon densitometry control software.

Purification of BsaC. The *bsaC* gene with a C-terminal His$_6$ tag was cloned into the expression

vector pET24a by PCR using the primers listed in Table S4. An overnight culture of *E. coli*

LOBSTR pET24a::BsaC-His$_6$ was used to inoculate 2 L of 37^{o}C pre-warmed terrific broth in a 4

L beveled flask. The culture was allowed to grow to 0.6 OD$_{600}$ where it was induced with 1 mM

IPTG. After 6 hours of induction, the culture was centrifuged to remove cells (BsaC-His$_6$ is

secreted by *E. coli*), and the resulting supernatant was precipitated with 90% ammonium sulfate.

Precipitated protein was solubilized in PBS, and used to load on to an ÄKTA purifier FPLC for

Ni-NTA chromatography in PBS with 10 mM imidazole as loading buffer, and PBS with 1M

imidazole as elution buffer. Eluted fractions were concentrated with an Amicon Ultra 2 mL 10 K

cutoff centrifugation columns. Concentrated elution fractions were then further processed using a

Superose 6 gel filtration column linked to an ÄKTA purifier FPLC.

Immunoblotting of BsaA oligomers. BsaA oligomer secretion by *C. perfringens* was analyzed

using western blots and slot blots. For the immunoblot experiments, the proteins were blotted onto

a PVDF membrane while submerged in cold buffer which contained 3 mM Na$_2$CO$_3$, 10 mM

NaHCO$_3$ and 20% methanol.

Immunoblotting Method One. The membranes were placed in a Snap i.d. 2.0 system

(Millipore Sigma) which uses a vacuum system to pull solutions through the membrane (referred

to a swashing steps below). Once set up, a solution containing Tris-buffered saline (TBS) (pH 7.4),

0.1% Tween 20 (TBST), and 1% bovine serum albumin (BSA) was added as a blocker. Both

primary and secondary antibodies were added to separate aliquots of the solution containing TBST

and 1% BSA. In the first step, the primary antibody solution was incubated for 10 minutes on the

membrane, followed by three washings with TBST. Then the secondary antibody solution was left to sit for 10 minutes on the membrane, followed by three washings with TBST. The primary antibody utilized was OctA-probe H-5 (1:250 dilution) used to detect the FLAG epitope (Santa Cruz Biotechnology). The secondary antibody utilized for these experiments was StarBright700 goat anti-mouse (Bio-Rad) (1:10,000 dilution). The fluorescently labeled immunoblots were imaged using a ChemiDoc MP imaging system (Bio-Rad), and the intensity of the fluorescence was measured using the ChemiDoc MP imaging software for densitometry measurements.

Immunoblotting method 2. Membranes were transferred to a container and blocked with EveryBlot blocking Buffer (Bio-Rad). Antibodies were diluted with EveryBlot blocking buffer and applied to membranes, followed by a minimum of five washes with TBST.

The primary antibodies utilized included OctA-probe H-5 (1:227 dilution) used to detect FLAG, and HA-Tag F-7 (1:227 dilution) used to detect HA (both from Santa Cruz Biotechnology, Santa Cruz, CA). Membranes submerged in EveryBlot blocking buffer with primary antibody were incubated at 4°C overnight. The secondary antibody utilized for these experiments was StarBright700 goat anti-mouse (Bio-Rad) (1:13,333 dilution), incubated for one hour at room temperature. Images of the immunoblots were taken using a ChemiDoc MP imaging system (Bio-Rad), to measure the intensity of the fluorescence.

Slot blots to detect secretion of BsaA oligomers. For slot blots, supernatants samples were prepared by diluting supernatants with sterile BHI. For initial experiments, samples of 100% pure culture supernatant, 50% culture supernatant and 50% blank media, 25% culture supernatant and 75% blank media, 10% culture supernatant and 90% blank media, 1% culture supernatant and 99% blank media, and 100% blank media were prepared. For later experiments, samples of 50% culture supernatant and 50% blank media, 20% culture supernatant and 80% blank media, 10% culture

supernatant and 90% blank media, 5% culture supernatant and 95% blank media, 1% culture

supernatant and 99% blank media, and 100% blank media were prepared. The 10% culture

supernatant samples were most often used to collect the raw intensity data for quantitative results.

A PVDF membrane was soaked in TBST prior to being loaded into the slot blot device

while still damp. The slot blot was loaded with TBST in each slot for 10 minutes. The TBST was

pulled through by vacuum, and then the diluted supernatants for the slot blots were loaded into the

individual slots of the device, and allowed to sit for 10 minutes. The supernatants were pulled

through by vacuum, and then the slot blot were loaded with TBST in each slot for another 10

minutes. The TBST was vacuumed off as well, and the membrane was developed using

immunoblotting method 2.

Western blots on *bsaA* mutants. Western blots to detect oligomerization of BsaA mutants were

performed by expressing the *sipW-bsaA-His₆* genes under control of the lactose-inducible

promoter. Supernatants were separated form cell pellets by centrifugation. The proteins in the

supernatant were concentrated by TCA precipitation (as described above), suspended in SDS-

PAGE loading buffer with 100 mM DTT and heated at 95 °C for 20 min. The bacteria in the cell

pellet were suspended in PBS and disrupted by bead beating with 0.1 mm zirconia-silica beads.

The beads were then removed by centrifugation and the cell extract brought to 1 X SDS-PAGE

buffer with 100 mM DTT and heated at 95 °C for 20 min. After running on SDS-PAGE gels, the

proteins were transferred to a PVDF membrane and developed using immunoblotting method 2

above, except the primary antibody was mouse monoclonal anti-His6 antibody His.H8 (Santa Cruz

Biotechnology).

Modeling BsaA Structure. BsaA from *C. perfringens* (accession number WP_011009871) was

created as a homotrimer using ColabFold v1.3.0 (18) based on AlphaFold v2.2.0 (19). Amber

minimization was not enabled. In the top ranked model based on pLDDT PyMOL v2.5.0 was used

to analyze side chains of interdigitated beta strands for potential sidechain crosslinks. Images were

created using ChimeraX v1.2. (42, 43) The same trimer of BsaA was created in AlphaFold 3 (24)

to observe if the new version creates crosslinks, and all trimer models made did contain the

crosslinks. Structures of BsaA were created with AlphaFold 3 (24).

Phylogeny of BsaA-like proteins. BsaA-like protein sequences were identified with 3 iterations

of jackhmmer (44) (HmmerWeb version 2.41.2) (45) using BsaA (UniProt:Q8XN23) from *C.*

perfringens as the seed. The protein sequence similarity network of proteins identified either with

jackhmmer or as a match to IPR022121 was constructed using the EFI-EST tool

(https://efi.igb.illinois.edu/efi-est/) (46, 47) with an alignment score of 10. Nodes were collapsed

at a sequence identity of 100%. The network was visualized with Cytoscape

(https://www.cytoscape.org) (48) using the Prefuse Force Directed OpenCL Layout. Gene

neighborhoods were retrieved using the EFI-GNT tool (https://efi.igb.illinois.edu/efi-gnt/) (47).

The presence of a N-E-K triad was determined by manual inspection of the multiple sequence

alignment. For phylogenetic reconstruction, sequences were aligned with MAFFT on XSEDE

using CIPRES (49, 50), edited in Jalview (51), and analyzed with IQ-Tree (52-54). Maximum

likelihood trees were constructed using the LG+F+G4 substitution model and ultrafast bootstrap

(1000 replicates). Consensus trees were visualized, rooted at the midpoint, and annotated with

iTOL (55). Data files are available in Supplemental File Data Set 1.

Acknowledgments

We thank W. Keith Ray, Richard Helm and the Virginia Tech Proteomics Incubator for assistance

with mass spectrometry and Megan Rochford and Hualan Liu for construction of in-frame

deletions mutants. Research reported in this publication was supported by the National Institute of

Allergy and Infectious Disease of the National Institutes of Health under award

number R21AI109391 to S. B. M. The content is solely the responsibility of the authors and does

not necessarily represent the official views of the National Institutes of Health. Work at the

Molecular Foundry was supported by the Office of Science, Office of Basic Energy Sciences, of

the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. The work conducted

by the U.S. Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE

Office of Science User Facility, is supported by the Office of Science of the U.S. Department of

Energy operated under Contract No. DE-AC02-05CH11231.

Author Contributions

Sarah E. Kivimaki: Investigation, Methodology, Visualization, and Writing – Review and Editing.

Samantha Dempsey: Investigation, Methodology, Visualization, and Writing.

Gary J. Camper: Investigation, Methodology, Visualization, and Writing.

Julia M. Tani: Investigation, Methodology, Visualization.

Ian K. Hicklin: Investigation, Methodology, Visualization, and Writing.

Crysten E. Blaby-Haas: Methodology, Visualization, and Writing – review and editing.

Anne M. Brown: Methodology, Visualization, and Writing – review and editing.

Stephen B. Melville: Conceptualization, Funding acquisition, Supervision, Investigation,

Methodology, Project administration, Writing, Review and editing.

References

1.   **Singh, P. K., J. Little, and M. S. Donnenberg.** 2022. Landmark Discoveries and Recent
Advances in Type IV Pilus Research. Microbiol Mol Biol Rev **86:**e0007622.
2.   **Naskar, S., M. Hohl, M. Tassinari, and H. H. Low.** 2021. The structure and mechanism of
the bacterial type II secretion system. Mol Microbiol **115:**412-424.
3.   **Korotkov, K. V., M. Sandkvist, and W. G. Hol.** 2012. The type II secretion system:
biogenesis, molecular architecture and mechanism. Nature reviews. Microbiology
10:336-351.
4.   **Ayers, M., P. L. Howell, and L. L. Burrows.** 2010. Architecture of the type II secretion and
type IV pilus machineries. Future microbiology **5:**1203-1218.
5.   **Desvaux, M., M. Hebraud, R. Talon, and I. R. Henderson.** 2009. Secretion and
subcellular localizations of bacterial proteins: a semantic awareness issue. Trends
Microbiol **17:**139-145.
6.   **Anne, J., A. Economou, and K. Bernaerts.** 2017. Protein Secretion in Gram-Positive
Bacteria: From Multiple Pathways to Biotechnology. Curr Top Microbiol Immunol
404:267-308.
7.   **Demchick, P., and A. L. Koch.** 1996. The permeability of the wall fabric of Escherichia coli
and Bacillus subtilis. Journal of bacteriology **178:**768-773.
8.   **Varga, J. J., V. Nguyen, D. K. O'brien, K. Rodgers, R. A. Walker, and S. B. Melville.** 2006.
Type IV pili-dependent gliding motility in the Gram-positive pathogen *Clostridium*
perfringens and other Clostridia. Mol Microbiol **62:**680-694.
9.   **Melville, S., and L. Craig.** 2013. Type IV pili in Gram-positive bacteria. Microbiol Mol Biol
Rev **77:**323-341.
10.   **Bohning, J., M. Ghrayeb, C. Pedebos, D. K. Abbas, S. Khalid, L. Chai, and T. a. M. Bharat.**
2022. Donor-strand exchange drives assembly of the TasA scaffold in Bacillus subtilis
biofilms. Nat Commun **13:**7082.
11.   **Erskine, E., R. J. Morris, M. Schor, C. Earl, R. M. C. Gillespie, K. M. Bromley, T.**
Sukhodub, L. Clark, P. K. Fyfe, L. C. Serpell, N. R. Stanley-Wall, and C. E. Macphee. 2018.
Formation of functional, non-amyloidogenic fibres by recombinant Bacillus subtilis TasA.
Mol Microbiol **110:**897-913.
12.   **Stover, A. G., and A. Driks.** 1999. Secretion, localization, and antibacterial activity of
TasA, a Bacillus subtilis spore-associated protein. J Bacteriol **181:**1664-1672.
13.   **Obana, N., K. Nakamura, and N. Nomura.** 2020. Temperature-regulated heterogeneous
extracellular matrix gene expression defines biofilm morphology in *Clostridium*
perfringens. NPJ Biofilms Microbiomes **6:**29.
14.   **Hartman, A. H., H. Liu, and S. B. Melville.** 2011. Construction and characterization of a
lactose-inducible promoter system for controlled gene expression in *Clostridium*
perfringens. Appl Environ Microbiol **77:**471-478.
15.   **Nunn, D., S. Bergman, and S. Lory.** 1990. Products of three accessory genes, pilB, pilC,
and pilD, are required for biogenesis of Pseudomonas aeruginosa pili. J Bacteriol
172:2911-2919.
16.   **Kelley, L. A., S. Mezulis, C. M. Yates, M. N. Wass, and M. J. Sternberg.** 2015. The Phyre2
web portal for protein modeling, prediction and analysis. Nat Protoc **10:**845-858.

17.   **Diehl, A., Y. Roske, L. Ball, A. Chowdhury, M. Hiller, N. Moliere, R. Kramer, D. Stoppler,**
718   **C. L. Worth, B. Schlegel, M. Leidert, N. Cremer, N. Erdmann, D. Lopez, H. Stephanowitz,**
719   **E. Krause, B. J. Van Rossum, P. Schmieder, U. Heinemann, K. Turgay, U. Akbey, and H.**
720   **Oschkinat.** 2018. Structural changes of TasA in biofilm formation of Bacillus subtilis. Proc
721   Natl Acad Sci U S A **115:**3237-3242.
722   18.   **Mirdita, M., K. Schutze, Y. Moriwaki, L. Heo, S. Ovchinnikov, and M. Steinegger.** 2022.
723   ColabFold: making protein folding accessible to all. Nat Methods **19:**679-682.
724   19.   **Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K.**
725   **Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S. a. A.**
726   **Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S.**
727   **Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T.**
728   **Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli,**
729   **and D. Hassabis.** 2021. Highly accurate protein structure prediction with AlphaFold.
730   Nature **596:**583-589.
731   20.   **Kang, H. J., and E. N. Baker.** 2011. Intramolecular isopeptide bonds: protein crosslinks
732   built for stress? Trends Biochem Sci **36:**229-237.
733   21.   **Kang, H. J., F. Coulibaly, F. Clow, T. Proft, and E. N. Baker.** 2007. Stabilizing isopeptide
734   bonds revealed in gram-positive bacterial pilus structure. Science **318:**1625-1628.
735   22.   **Baker, E. N., C. J. Squire, and P. G. Young.** 2015. Self-generated covalent cross-links in
736   the cell-surface adhesins of Gram-positive bacteria. Biochem Soc Trans **43:**787-794.
737   23.   **Evans, R. O. N., M.; Pritzel, A.; Antropova, N.; Senior, A.; Green, T.; Žídek, A.; Bates, R.;**
738   **Blackwell, S.; Yim, J.; Et Al.** 2022. Protein Complex Prediction with AlphaFold-Multimer.
739   bioRxiv **2022**.
740   24.   **Abramson, J., J. Adler, J. Dunger, R. Evans, T. Green, A. Pritzel, O. Ronneberger, L.**
741   **Willmore, A. J. Ballard, J. Bambrick, S. W. Bodenstein, D. A. Evans, C. C. Hung, M.**
742   **O'neill, D. Reiman, K. Tunyasuvunakool, Z. Wu, A. Zemgulyte, E. Arvaniti, C. Beattie, O.**
743   **Bertolli, A. Bridgland, A. Cherepanov, M. Congreve, A. I. Cowen-Rivers, A. Cowie, M.**
744   **Figurnov, F. B. Fuchs, H. Gladman, R. Jain, Y. A. Khan, C. M. R. Low, K. Perlin, A.**
745   **Potapenko, P. Savy, S. Singh, A. Stecula, A. Thillaisundaram, C. Tong, S. Yakneen, E. D.**
746   **Zhong, M. Zielinski, A. Zidek, V. Bapst, P. Kohli, M. Jaderberg, D. Hassabis, and J. M.**
747   **Jumper.** 2024. Accurate structure prediction of biomolecular interactions with AlphaFold
748   3. Nature **630:**493-500.
749   25.   **Plamont, M. A., E. Billon-Denis, S. Maurin, C. Gauron, F. M. Pimenta, C. G. Specht, J.**
750   **Shi, J. Querard, B. Pan, J. Rossignol, K. Moncoq, N. Morellet, M. Volovitch, E. Lescop, Y.**
751   **Chen, A. Triller, S. Vriz, T. Le Saux, L. Jullien, and A. Gautier.** 2016. Small fluorescence-
752   activating and absorption-shifting tag for tunable protein imaging in vivo. Proc Natl Acad
753   Sci U S A **113:**497-502.
754   26.   **Tebo, A. G., F. M. Pimenta, Y. Zhang, and A. Gautier.** 2018. Improved Chemical-Genetic
755   Fluorescent Markers for Live Cell Microscopy. Biochemistry **57:**5648-5653.
756   27.   **Gautier, A., L. Jullien, C. Li, M. A. Plamont, A. G. Tebo, M. Thauvin, M. Volovitch, and S.**
757   **Vriz.** 2021. Versatile On-Demand Fluorescent Labeling of Fusion Proteins Using
758   Fluorescence-Activating and Absorption-Shifting Tag (FAST). Methods Mol Biol **2350:**253-
759   265.

28.   **Fives-Taylor, P. M., and D. W. Thompson.** 1985. Surface properties of *Streptococcus*
sanguis FW213 mutants nonadherent to saliva-coated hydroxyapatite. Infect Immun
47:752-759.
29.   **Berry, J. L., I. Gurung, J. H. Anonsen, I. Spielman, E. Harper, A. M. J. Hall, V. J. Goosens,**
C. Raynaud, M. Koomey, N. Biais, S. Matthews, and V. Pelicic. 2019. Global biochemical
and structural analysis of the type IV pilus from the Gram-positive bacterium
Streptococcus sanguinis. J Biol Chem **294:**6796-6808.
30.   **Gurung, I., I. Spielman, M. R. Davies, R. Lala, P. Gaustad, N. Biais, and V. Pelicic.** 2016.
Functional analysis of an unusual type IV pilus in the Gram-positive *Streptococcus*
sanguinis. Mol Microbiol **99:**380-392.
31.   **Han, X., R. M. Kennan, D. Parker, J. K. Davies, and J. I. Rood.** 2007. Type IV fimbrial
biogenesis is required for protease secretion and natural transformation in *Dichelobacter*
nodosus. J Bacteriol **189:**5022-5033.
32.   **Kirn, T. J., N. Bose, and R. K. Taylor.** 2003. Secretion of a soluble colonization factor by
the TCP type 4 pilus biogenesis pathway in *Vibrio cholerae*. Molecular microbiology
49:81-92.
33.   **Whittaker, C. A., and R. O. Hynes.** 2002. Distribution and evolution of von
Willebrand/integrin A domains: widely dispersed domains with roles in cell adhesion and
elsewhere. Mol Biol Cell **13:**3369-3387.
34.   **Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J.**
Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. Nucleic Acids Res **25:**3389-3402.
35.   **Alegre-Cebollada, J., C. L. Badilla, and J. M. Fernandez.** 2010. Isopeptide bonds block
the mechanical extension of pili in pathogenic Streptococcus pyogenes. J Biol Chem
285:11235-11242.
36.   **Guilford, W. H., R. C. Lantz, and R. W. Gore.** 1995. Locomotive forces produced by single
leukocytes in vivo and in vitro. Am J Physiol **268:**C1308-1312.
37.   **Rood, J. I., and S. T. Cole.** 1991. Molecular genetics and pathogenesis of *Clostridium*
perfringens. Microbiol Rev **55:**621-648.
38.   **Lisle, J. T., J. J. Smith, D. D. Edwards, and G. A. Mcfeters.** 2004. Occurrence of microbial
indicators and *Clostridium perfringens* in wastewater, water column samples, sediments,
drinking water, and Weddell seal feces collected at McMurdo Station, Antarctica. Appl
Environ Microbiol **70:**7269-7276.
39.   **Nariya, H., S. Miyata, M. Suzuki, E. Tamai, and A. Okabe.** 2011. Development and
application of a method for counterselectable in-frame deletion in *Clostridium*
perfringens. Applied and environmental microbiology **77:**1375-1382.
40.   **Hendrick, W. A., M. W. Orr, S. R. Murray, V. T. Lee, and S. B. Melville.** 2017. Cyclic Di-
GMP Binding by an Assembly ATPase (PilB2) and Control of Type IV Pilin Polymerization
in the Gram-Positive Pathogen *Clostridium perfringens*. J Bacteriol **199**.
41.   **Sambrook, J., and D. W. Russel.** 2001. Molecular Cloning, A Laboratory Manual, 3rd
Edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
42.   **Goddard, T. D., C. C. Huang, E. C. Meng, E. F. Pettersen, G. S. Couch, J. H. Morris, and T.**
E. Ferrin. 2018. UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. Protein Sci **27:**14-25.

43.   **Pettersen, E. F., T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, T. I. Croll, J. H.**
Morris, and T. E. Ferrin. 2021. UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. Protein Sci **30:**70-82.

44.   **Johnson, L. S., S. R. Eddy, and E. Portugaly.** 2010. Hidden Markov model speed heuristic
and iterative HMM search procedure. BMC Bioinformatics **11:**431.

45.   **Potter, S. C., A. Luciani, S. R. Eddy, Y. Park, R. Lopez, and R. D. Finn.** 2018. HMMER web
server: 2018 update. Nucleic Acids Res **46:**W200-W204.

46.   **Gerlt, J. A., J. T. Bouvier, D. B. Davidson, H. J. Imker, B. Sadkhin, D. R. Slater, and K. L.**
Whalen. 2015. Enzyme Function Initiative-Enzyme Similarity Tool (EFI-EST): A web tool
for generating protein sequence similarity networks. Biochim Biophys Acta **1854:**1019-
1037.

47.   **Oberg, N., R. Zallot, and J. A. Gerlt.** 2023. EFI-EST, EFI-GNT, and EFI-CGFP: Enzyme
Function Initiative (EFI) Web Resource for Genomic Enzymology Tools. J Mol Biol
435:168018.

48.   **Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.**
Schwikowski, and T. Ideker. 2003. Cytoscape: a software environment for integrated
models of biomolecular interaction networks. Genome Res **13:**2498-2504.

49.   **Katoh, K., K. Kuma, H. Toh, and T. Miyata.** 2005. MAFFT version 5: improvement in
accuracy of multiple sequence alignment. Nucleic Acids Res **33:**511-518.

50.   **M. A. Miller, W. P. a. T. S.** Creating the CIPRES Science Gateway for inference of large
phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE)**:**1-8.

51.   **Clamp, M., J. Cuff, S. M. Searle, and G. J. Barton.** 2004. The Jalview Java alignment
editor. Bioinformatics **20:**426-427.

52.   **Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. Von Haeseler, and L. S. Jermiin.**
2017. ModelFinder: fast model selection for accurate phylogenetic estimates. Nat
Methods **14:**587-589.

53.   **Nguyen, L. T., H. A. Schmidt, A. Von Haeseler, and B. Q. Minh.** 2015. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol
Evol **32:**268-274.

54.   **Hoang, D. T., O. Chernomor, A. Von Haeseler, B. Q. Minh, and L. S. Vinh.** 2018.
UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol Evol **35:**518-522.

55.   **Letunic, I., and P. Bork.** 2024. Interactive Tree of Life (iTOL) v6: recent updates to the
phylogenetic tree display and annotation tool. Nucleic Acids Res **52:**W78-W82.

56.   **Grant, S. G., J. Jessee, F. R. Bloom, and D. Hanahan.** 1990. Differential plasmid rescue
from transgenic mouse DNAs into Escherichia coli methylation-restriction mutants. Proc
Natl Acad Sci U S A **87:**4645-4649.

57.   **Andersen, K. R., N. C. Leksa, and T. U. Schwartz.** 2013. Optimized E. coli expression
strain LOBSTR eliminates common contaminants from His-tag purification. Proteins
81:1857-1861.

58.   **Guzman, L. M., D. Belin, M. J. Carson, and J. Beckwith.** 1995. Tight regulation,
modulation, and high-level expression by vectors containing the arabinose PBAD
promoter. J Bacteriol **177:**4121-4130.

 Figure legends

Figure 1. Diagram showing the type IV pili-associated genes in *C. perfringens* strain 13. Figure

derived from reference (9).

Figure 2. (A). SDS-PAGE gel showing the proteins in the secretome of the respective strains. The

gel was stained with the fluorescent protein dye Sypro-Ruby, the colors inverted to make the

protein bands in the samples appear dark but the pre-stained molecular weight markers appear as

bright bands. **(B).** Western blot showing the levels of secretion of the BsaC protein in different

strains. The secondary antibody was fluorescently tagged for visualization. Purified BsaC was

added in lane 5 as a positive control for detecting BsaC. **(C).** The gene synteny of the *bsa* operon

and adjacent sortase-encoding gene in *C. perfringens* strain 13. **(D).** The gene synteny of the *bsa*

operon and adjacent sortase-encoding gene in *C. perfringens* strain ATCC 13124 and most other

strains of *C. perfringens* that were observed. The sections of the *bsaB* gene from strain 13 that

were homologous to those in strain ATCC 13124 are labeled, along with the location of an LPXTG

sortase dependent sequence in one of the *bsaB*-encoding genes.

Figure 3. Results from densitometry analysis of western blots from the secretome of strains with

in-frame deletions of T4P-encoding genes. Each strain contained plasmid pGC1 for regulated

expression of the *bsa* operon with a His$_6$ tag added to the *bsaC* gene. The role of each gene product

in pilus function is shown in Fig. 1. For each experiment, the relative fluorescence intensity of the

wild-type strain (HN13) was set at 1.0 and the relative fluorescence intensity for each other strain

in the same experiment was then set relative to that value. Each strain was tested at least three

times with triplicate samples for each experiment. All of the experimental runs were combined and

one sample t and Wilcoxon tests were run to determine if the mutant strains were statistically

different from 1. Asterisks indicate strains that were different ($P < 0.05$) from strain HN13 (i.e.,

1).

Figure 4. (A) AlphaFold model of a homotrimer of BsaA in cartoon representation. Each

independent monomer is colored separately. Residues 28-38 of a monomer incorporate themselves

into the beta strands of an adjacent monomer. **(B)** Close-up view of interdigitated beta strands from

two monomers in cartoon representation, showing relevant side chains as sticks colored by

element. N35 and the giving chain are colored purple, E56, K74, and the receiving monomer are

colored cyan. Side chains forming the hydrophobic pocket are colored brown. The bond created

by AlphaFold between N35 and K74 is between an oxygen and nitrogen due to limitations in

AlphaFold; in reality, an amide bond would form between the side chains with ammonia as the

leaving group. **(C)** Close-up view of the N35-E56-K74 catalytic triad, showing the close proximity

of the catalytic E56 residue to the bond forming N35 and K74. All images were created with

ChimeraX v1.2. (42, 43).

Figure 5. Western blots showing loss of oligomer formation in BsaA with mutations in residues

involved in isopeptide bond formation. **(A)** Oligomer formation seen in TCA precipitated culture

supernatants or in the disrupted whole cell extracts of strains producing BsaA with single

mutations. **(B)** Comparison of BsaA oligomer formation in the N35A and N35A/K74A mutants in

a wild-type background (left side) and one in which the chromosomal copy of the *bsaA* gene was

deleted. Note that the chromosomal *bsaA* mutation did not change the oligomerization levels of

the mutants. **(C)** Western blot showing mutation of the E56 residue to Ala abolished

oligomerization of BsaA. P, disrupted cell pellet; S, TCA precipitate of culture supernatant.

Figure 6. BsaA oligomers are found on the bacterial surface. **(A)** Expression of a FAST protein

fusion to the cytoplasmic PilT protein (pSM413) only shows fluorescence with the membrane

permeable Coral dye. **(B)** A BsaA-FAST fusion protein (pSM412) fluoresces in the presence of

both a membrane permeable dye (Coral) and a membrane impermeable dye (Amber-NP),

indicating it is surface exposed to these dyes.

Figure 7. BsaA phylogeny. (**A**) A consensus tree of representative BsaA-like proteins. Leaf labels

are colored by taxonomy according to the color key. BsaA, BsaB, and two other BsaA-like proteins

identified in *C. perfringens* are labeled, as are TasA from *B. subtilis* and CalY from *B. cereus*.

Bootstrap values are represented as scaled circles according to the inset key. (**B**) Sequence

similarity network of identified BsaA-like proteins. Nodes representing labelled *Clostridium* and

Bacillus proteins in panel A are labeled with an arrow. Nodes are colored by taxonomy according

to the color key. Proteins with conserved residues that would be expected to be involved in

formation of a spontaneous intermolecular isopeptide bond are represented by triangle nodes with

a red outline. (**C**) Example gene neighborhoods of genes encoding BsaA-like proteins and genes

encoding putative surface filament structures. The complete matrix tree data set is included in a

Supplemental Information file.

Figure 8. Model of BsaA oligomerization. BsaA monomers (purple ovals) are oligomerized by

isopeptide bond formation after secretion via the Sec system in *C. perfringens*. SipW is the signal

peptidase for BsaA secretion (Fig. 2C and (13)). Secretion of the oligomers past the PG layer may

be due to the action of T4P-associated proteins (Fig. S4).

Table 1. Strains and plasmids used in this report.

Strain or plasmid	Relevant characteristics	Reference or Source
Strains		
Clostridium perfringens type A strain HN13	*ΔgalKT*	(39)
Escherichia coli strain DH10B	F– *mcr*A Δ(*mrr-hsd*RMS-*mcr*BC) Φ80*lac*ZΔM15 Δ*lac*X74 *rec*A1 *end*A1 *ara*D139 Δ(*ara leu*) 7697 *gal*U *gal*K *rps*L *nup*G λ–	(56)
Escherichia coli strain LOBSTR	Derivative of strain BL21, designed for low background binding to Ni-affinity columns	(57)
Plasmids		
pKRAH1	Cm^r; Lactose-inducible expression system plasmid for use in *C. perfringens*	(14)
pET24a	Protein expression vector	Novagen
pCRBluntIITOPO	PCR cloning vector	Invitrogen
pBAD30	Amp^r; Arabinose-inducible expression system plasmid for use in *E. coli*	(58)
pKRAH1-*pilA3*	Expression of *pilA3* for complementation	This study
pHLL65	The *bsa* operon in pKRAH1	This study
pGC1	pHLL65 with a His$_6$ tag added to the C-terminus of *bsaC*	This Study
pGC8	Vector based on pET24-a; used to express and purify VWA-His6 in *E. coli*	This Study
pSK1	The *bsa* operon in pKRAH1 with a FLAG-tag added to the C-terminus of *bsaA*	This Study

pSK2	Same as pSK1 with an HA tag at the C-terminus of *bsaB*	This Study
pSK4	The *sipW* and *bsaA* genes in pKRAH1 with a FLAG tag at the C-terminus of *bsaA*.	This Study
pSM407	The *sipW* and *bsaA* genes in pKRAH1 with a His$_6$ tag at the C-terminus of *bsaA*.	This Study
pSM408	pSM407 with a N35A mutation in *bsaA*	This Study
pSM409	pSM407 with a K74A mutation in *bsaA*	This Study
pSM410	pSM407 with N35A/K74A mutations in *bsaA*	This Study
pSM411	pSM407 with E56A mutations in *bsaA*	This Study
pSM412	pSM407 with a FAST gene fusion at the C-terminus of *bsaA* in place of the His$_6$ tag	This Study
pSM413	A *pilT*-FAST gene fusion in pKRAH1	This Study

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

Coral
(permeable)

Amber-NP
(impermeable)

B

Coral
(permeable)

Amber-NP
(impermeable)

Figure 7

Figure 8

Peptidoglycan layer

BsaA

Disordered N-terminus

SipW

BsaA secretion

BsaA secretion

Cytoplasmic Membrane