Microbes display broad diversity in cobamide preferences

Kenny C. Mok[a], Olga M. Sokolovskaya[a,b], Adam M. Deutschbauer[a,c], Hans K. Carlson[c], Michiko
E. Taga[a]#

[a]Department of Plant & Microbial Biology, University of California, Berkeley, Berkeley, CA
94720, U.S.A.

[b]Present address: Department of Biological Engineering, Massachusetts Institute of Technology,
Cambridge, MA 02139, U.S.A.

[c]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National
Laboratory, Berkeley, CA 94720, U.S.A.

#Correspondence to: taga@berkeley.edu

Abstract

Cobamides, the vitamin B_{12} (cobalamin) family of cofactors, are used by most organisms but produced by only a fraction of prokaryotes, and are thus considered key shared nutrients among microbes. Cobamides are structurally diverse, with multiple different cobamides found in most microbial communities. The ability to use different cobamides has been tested for several bacteria and microalgae, and nearly all show preferences for certain cobamides. This approach is limited by the commercial unavailability of cobamides other than cobalamin. Here, we have extracted and purified seven commercially unavailable cobamides to characterize bacterial cobamide preferences based on growth in specific cobamide-dependent conditions. The tested bacteria include engineered strains of *Escherichia coli*, *Sinorhizobium meliloti*, and *Bacillus subtilis* expressing native or heterologous cobamide-dependent enzymes, cultured under conditions that functionally isolate specific cobamide-dependent processes such as methionine synthesis. Comparison of these results to previous studies of diverse bacteria and microalgae revealed that a broad diversity of cobamide preferences exists not only across different organisms, but also between different cobamide-dependent metabolic pathways within the same organism. The microbes differed in the cobamides that support growth most efficiently, cobamides that do not support growth, and the minimum cobamide concentrations required for growth. The latter differ by up to four orders of magnitude across organisms from different environments and by up to 20-fold between cobamide-dependent enzymes within the same organism. Given that cobamides are shared, required for use of specific growth substrates, and essential for central metabolism in certain organisms, cobamide preferences likely impact community structure and function.

Importance

Nearly all bacteria are found in microbial communities with tens to thousands of other species. Molecular interactions such as metabolic cooperation and competition are key factors underlying community assembly and structure. Cobamides, the vitamin B_{12} family of enzyme cofactors, are one such class of nutrients, produced by only a minority of prokaryotes but required by most microbes. A unique aspect of cobamides is their broad diversity, with nearly 20 structural forms identified in nature. Importantly, this structural diversity impacts growth, as most bacteria that have been tested show preferences for specific cobamide forms. We measured cobamide-dependent growth in several model bacteria and compared the results to previous analyses of cobamide preference. We found that cobamide preferences vary widely across bacteria, showing the importance of characterizing these aspects of cobamide biology to understand the impact of cobamides on microbial communities.

B$_{12}$ and other cobamide cofactors are required by organisms in all domains of life (1). Only certain
bacteria and archaea synthesize cobamides (2), while others must acquire them exogenously. Thus,
cobamides are considered shared nutrients within microbial communities. A unique aspect of
cobamides is their structural diversity. While B$_{12}$ (cobalamin, Cbl) is a well-known vitamin
important for human health, nearly 20 cobamides with alternative lower ligands exist in nature
(Fig. 1A) (3-5). Though research with alternate cobamides is limited because they are
commercially unavailable, cobamide-dependent enzymes and organisms are known to have
distinct preferences for different cobamides (6-16). Further, addition of different cobamides to soil
or soil-derived enrichment cultures elicited distinct shifts in bacterial abundances, suggesting that
cobamide structure influences bacterial growth at the community level (17). Thus, cobamide
preference in bacteria is likely important for microbial community structure.

Here, we address four questions about cobamide use and preference: 1) Do different cobamide-
dependent processes in the same bacterium have the same cobamide requirements? 2) How many,
and which, cobamides can support growth? 3) Which cobamides are preferred, and do preferences
differ across organisms? 4) How much cobamide do microbes need for growth, and do these
requirements vary by enzyme, taxonomy, or environment? We addressed these questions by
measuring growth of wild-type and engineered bacteria in different cobamide-dependent
conditions with up to eight cobamides at a range of concentrations. Cobamide preference was
defined based on the concentration of added cobamide that elicits half-maximal growth (EC$_{50}$),
with lower values corresponding to more preferred cobamides. This *in vivo* assay encompasses the
different aspects of cobamide utilization, including uptake, adenosylation, riboswitch-based
regulation, and the cobamide-dependent enzymes themselves. We compared these measurements
with those from published literature to gain a comprehensive view of *in vivo* cobamide preferences
across taxa and environments.

Different cobamide-dependent enzymes in a single organism can have distinct cobamide
preferences

First we compared the cobamide requirements of three cobamide-dependent processes in a
Sinorhizobium meliloti strain engineered to lack corrinoid biosynthesis capability. Wild type *S.*
meliloti produces Cbl for methionine synthase (MetH), methylmalonyl-CoA mutase (MCM), and
ribonucleotide reductase (NrdJ), each of which can be tested separately for cobamide preference
by altering the genetic background and growth substrates (see Methods). Although MetH-
dependent growth was supported by all cobamides tested, the benzimidazolyl cobamides Cbl and
[Bza]Cba were strongly preferred, with EC$_{50}$ values two to three orders of magnitude lower than
the purinyl and phenolyl cobamides (Fig. 1B, Table S1). The concentration requirements and
preferences are similar to what we previously observed for MCM-dependent growth (8), except
that [2-MeAde]Cba better supported MCM-dependent growth (Fig. 2). NrdJ-dependent growth
required 25- and 15-fold higher Cbl and [Bza]Cba concentrations, respectively (Fig. 1B-C, Table
S1), which suggests either more cobamide is required for NrdJ function or higher levels of NrdJ
enzyme are necessary to support growth. The relative cobamide preferences for MetH- and NrdJ-
dependent growth are similar, though phenolyl cobamides did not support NrdJ-dependent growth,
consistent with NrdJ being a base-on enzyme (18) and phenolyl cobamides being unable to adopt

the base-on conformation (Fig. 1A-C, Fig. 2). Cbl was the preferred cobamide for all three cobamide-dependent conditions, consistent with adaptation to endogenously produced Cbl (Fig. 2).

We used *Escherichia coli*, a bacterium unable to synthesize cobamides *de novo*, to compare cobamide preferences of ethanolamine ammonia-lyase (EAL) and MetH using engineered strains cultured under different conditions. *E. coli* requires EAL for growth when ethanolamine is the nitrogen source and requires MetH for methionine synthesis when *metE*, encoding the cobamide-independent methionine synthase, is deleted. EAL-dependent growth required 5- to 78-fold higher cobamide concentrations than MetH-dependent growth and showed different cobamide preferences (Fig. 1D-E, Table S1). [2-MeAde]Cba, produced by *E. coli* when provided the precursor cobinamide (19), was most preferred for EAL-dependent growth, but less preferred by MetH. In contrast, Cbl was the least preferred cobamide that supports EAL-dependent growth, but most preferred for MetH-dependent growth (Fig. 1D-E). [*p*-Cre]Cba did not support EAL-dependent growth, consistent with EAL being a base-on enzyme (10). Together, these results show that cobamide-dependent pathways in the same organism can have distinct cobamide preferences and require different cobamide concentrations.

Cobamide use in a single process varies across bacteria

We next examined cobamide preferences for MetH-dependent growth in several additional bacteria. These included two cobamide non-producers that inhabit the human gut, *Bacteroides thetaiotaomicron* and *Ruminococcus gnavus*, which have MetH but lack MetE, as well as engineered *E. coli* and *Bacillus subtilis* strains heterologously expressing MetH orthologs from *Vibrio cholerae* and *Priestia megaterium*, respectively. In each strain, growth was supported by all or most of the cobamides tested, suggesting there is promiscuity in cobamide uptake, adenosylation, regulation, and use by MetH (Fig. 1E-J). However, EC_{50} values spanned a minimum of two orders of magnitude for each individual strain among cobamides supporting growth (Fig. 1, 2). While in all cases, either [Ade]Cba or a phenolyl cobamide was least preferred, and Cbl was most preferred except by *B. thetaiotaomicron*, variability in cobamide preference was observed across organisms (Fig. 1F-J). This variability is most apparent when contrasting *B. thetaiotaomicron* and *B. subtilis* expressing *P. megaterium metH*, which did not grow with phenolyl or purinyl cobamides, respectively, at any tested concentration (Fig. 1H-I). The results of the latter strain largely mirror prior growth measurements at a single cobamide concentration (20), and the preference for Cbl is consistent with *P. megaterium* being a Cbl producer. In contrast, we previously showed that the human gut microbe *Akkermansia muciniphila* has no cobamide preference (Fig. 2) due to its ability to remodel diverse cobamides to [Ade]Cba (21).

Expressing *V. cholerae metH* in *E. coli* afforded the opportunity to compare the cobamide preferences of *E. coli* and *V. cholerae* MetH orthologs in the same intracellular environment. Unlike *E. coli*, wild-type *V. cholerae* cannot use [Ade]Cba for MetH-dependent growth (12). Our results suggest this is due to its poor use by MetH, as the EC_{50} for [Ade]Cba was nearly 300-fold higher for *E. coli* expressing *V. cholerae metH* compared to its native *metH* (52 versus 0.18 nM) (Fig. 1F-G, Table S1). Overexpression of *E. coli metH* led to improved growth with the least preferred cobamides (Fig. 1E-F). This suggests MetH is limiting for *E. coli* growth with certain

cobamides including [Ade]Cba, and is consistent with our previous observation that mutation of the regulator *metR* or the *metH* 5′ untranslated region improved growth of *E. coli* with [Ade]Cba (22). Overall, even with a limited number of taxa, we observed considerable variability in cobamide use with MetH-dependent growth in different bacteria. While generally, benzimidazolyl cobamides were most preferred and phenolyl cobamides the least, growth with specific cobamides varied greatly between organisms, particularly the purinyl cobamides.

Cobamide requirements in bacteria and microalgae span orders of magnitude and correspond to environment

Comparison of these results to other studies that evaluated cobamide-dependent growth demonstrated that taxonomically diverse microbes from soil are also variable in their relative preferences for different cobamides, and that Cbl requirements of eukaryotic microalgae vary by nearly four orders of magnitude (Fig. 2). Further, the lowest EC_{50} values for most soil bacteria and microalgae are 1-2 orders of magnitude lower than those for human gut commensal bacteria (Fig. 2). This suggests the former are adapted to survive at much lower cobamide concentrations, which could be due to more efficient use of cobamides through improvements to processes such as uptake or binding by cobamide-dependent enzymes. If EC_{50} values are indicative of cobamide concentrations in these environments, the bioavailable cobamide concentration can be estimated as 0.1 to 10 pM in aquatic environments, 1-10 pM in soil, and 10-100 pM in the human gut. Consistent with these values, Cbl has been detected at pM concentrations in aquatic environments (23). 41 nM cobamide has been detected in soil, though soil's robust ability to adsorb cobamides suggests not all may be bioavailable (17). Cobamide concentrations in the gut regions that harbor the bacteria in this study have not been measured. It is worth noting that while the cobamides we tested supported growth in nearly all cases (Fig. 1), the less preferred cobamides for certain bacteria are unlikely to be present in sufficient quantities to support growth of these bacteria in their natural environments.

Together, these results demonstrate that cobamide preferences of enzymes in the same organism and of cobamide-dependent growth across taxonomically diverse microbes are variable. Furthermore, cobamide concentrations required for growth vary by orders of magnitude across environments, suggesting microbes are adapted to cobamide levels in their environment, likely by tuning the sensitivity of cobamide uptake, adenosylation, regulation, and use by cobamide-dependent enzymes. The characterization of cobamide abundances, bioavailability, requirements, and preferences are therefore necessary to understand the role of cobamide metabolism within microbial communities.

Figure 1. Cobamide structures and cobamide-dependent growth. A) Structure of B_{12} (cobalamin), which contains the lower ligand 5,6-dimethylbenzimidazole, is shown in the base-on conformation in which the ring nitrogen is coordinated to the cobalt ion (dashed line) (left). Alternative lower ligands of cobamides in this study (right) are shown with the names of the cobamides given below each structure. (B-J) Cobamide dose-dependent growth assays showing OD_{600} measured at the indicated times for (B) MetH-dependent growth of *S. meliloti*, (C) NrdJ-dependent growth of *S. meliloti*, (D) EAL-dependent growth of *E. coli*, (E) MetH-dependent growth of *E. coli*, (F) MetH-dependent growth of *E. coli* expressing *E. coli metH* on a plasmid, (G) MetH-dependent growth of *E. coli* expressing *V. cholerae metH* on a plasmid, (H) MetH-dependent growth of *B. subtilis* expressing *P. megaterium metH*, (I) MetH-dependent growth of *B. thetaiotaomicron*, and (J) MetH-dependent growth of *R. gnavus*. The EC_{50} values calculated from the curves in B-J and genotypes of the engineered strains in B-H are shown in Table S1. OD_{600} values for [Bza]Cba-supplemented cultures of *B. subtilis* were recorded after 72 hr because growth was not observed until after 45 hr. Points represent the means of 3-6 biological replicates; error bars represent standard deviation.

Figure 2. Comparison of EC$_{50}$ values for cobamide-dependent growth. Organisms examined in the current study (bold) are compared with those from previous studies (8, 13, 21, 24, 25). *S. meliloti* Rm1021 Δ*nrdJ cobD*::*gus* Gm^R *metH*::Tn*5* pMSO3-*nrdAB*(*E. coli*) was used for *S. meliloti* MCM-dependent growth (8). EC$_{50}$ values for *S. meliloti* are higher than for the other tested microbes likely because wild-type *S. meliloti* synthesizes Cbl *de novo* and lacks a high-affinity cobamide uptake system. MetH-dependent growth of *C. reinhardtii* was tested in a *metE* mutant (13). Symbols show the mean EC$_{50}$ values. Capped bars represent 95% confidence intervals, except with organisms labeled with *, which indicates error as standard deviation. Error of *C. reinhardtii* and *P. lutheri* EC$_{50}$ values were not reported (13). Bars are uncapped on the left or right when lower or upper bounds for 95% confidence intervals could not be determined, respectively. The lower bound for *Blastococcus* sp. B12_004 grown with [5-OHBza]Cba is 10^{-7} nM (25). The base of the leftward and rightward arrows represents maximal and minimal concentrations for EC$_{50}$ from dose-response assays in which lack of growth or saturating growth was not reached, respectively. For *O. tauri* and *A. carterae*, EC$_{50}$ values could not be calculated, but the capped bars for [Ade]Cba show the upper and lower bounds (13). Symbols in the shaded region on the right represent cobamides that were unable to support growth at any concentration tested.

Acknowledgements

We thank members of the Taga lab for helpful discussions and Zoila Alvarez-Aponte and Rebecca Procknow for critical reading of the manuscript. We thank Kristen LeGault and Kim Seed for genomic DNA of *Vibrio cholerae* O1 biovar El Tor str. N16961. This research was supported by NIH grant R35GM139633 to M.E.T. A.M.D. and H.K.C. were supported by ENIGMA-Ecosystems and Networks Integrated with Genes and Molecular Assemblies (http://enigma.lbl.gov), a Science Focus Area Program at Lawrence Berkeley National Laboratory, and is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under contract number DE-AC02-05CH11231.

References

1. Sokolovskaya OM, Shelton AN, Taga ME. 2020. Sharing vitamins: Cobamides unveil microbial interactions. Science 369:eaba0165. doi:10.1126/science.aba0165.
2. Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, Haft DR, Taga ME. 2019. Uneven distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative genomics. ISME J 13:789-804. doi:10.1038/s41396-018-0304-9.
3. Allen RH, Stabler SP. 2008. Identification and quantitation of cobalamin and cobalamin analogues in human feces. Am J Clin Nutr 87:1324-35. doi:10.1093/ajcn/87.5.1324.
4. Renz P. 1999. Biosynthesis of the 5,6-dimethylbenzimidazole moiety of cobalamin and of the other bases found in natural corrinoids, p 557-566. *In* Banerjee R (ed), Chemistry and Biochemistry of B12. John Wiley & Sons, Inc., New York.
5. Yan J, Bi M, Bourdon AK, Farmer AT, Wang PH, Molenda O, Quaile AT, Jiang N, Yang Y, Yin Y, Simsir B, Campagna SR, Edwards EA, Loffler FE. 2018. Purinyl-cobamide is a native prosthetic group of reductive dehalogenases. Nat Chem Biol 14:8-14. doi:10.1038/nchembio.2512.
6. Yi S, Seth EC, Men YJ, Stabler SP, Allen RH, Alvarez-Cohen L, Taga ME. 2012. Versatility in corrinoid salvaging and remodeling pathways supports corrinoid-dependent metabolism in *Dehalococcoides mccartyi*. Appl Environ Microbiol 78:7745-52. doi:10.1128/AEM.02150-12.
7. Tanioka Y, Miyamoto E, Yabuta Y, Ohnishi K, Fujita T, Yamaji R, Misono H, Shigeoka S, Nakano Y, Inui H, Watanabe F. 2010. Methyladeninylcobamide functions as the cofactor of methionine synthase in a cyanobacterium, *Spirulina platensis* NIES-39. FEBS Lett 584:3223-6. doi:10.1016/j.febslet.2010.06.013.
8. Sokolovskaya OM, Mok KC, Park JD, Tran JLA, Quanstrom KA, Taga ME. 2019. Cofactor Selectivity in Methylmalonyl Coenzyme A Mutase, a Model Cobamide-Dependent Enzyme. mBio 10:e01303-19. doi:10.1128/mBio.01303-19.
9. Shelton AN, Lyu X, Taga ME. 2020. Flexible Cobamide Metabolism in *Clostridioides* (*Clostridium*) *difficile* 630 Δ*erm*. J Bacteriol 202:e00584-19. doi:10.1128/JB.00584-19.
10. Poppe L, Bothe H, Bröker G, Buckel W, Stupperich E, Rétey J. 2000. Elucidation of the coenzyme binding mode of further B_{12}-dependent enzymes using a base-off analogue of coenzyme B_{12}. J Mol Catal 10:345-350. doi:10.1016/S1381-1177(00)00136-3.
11. Poppe L, Stupperich E, Hull WE, Buckel T, Rétey J. 1997. A base-off analogue of coenzyme-B_{12} with a modified nucleotide loop. [1]H-NMR structure analysis and kinetic studies with (*R*)-methylmalonyl-CoA mutase, glycerol dehydratase, and diol dehydratase. Eur J Biochem 250:303-307. doi:10.1111/j.1432-1033.1997.0303a.x.
12. Ma AT, Tyrell B, Beld J. 2020. Specificity of cobamide remodeling, uptake and utilization in *Vibrio cholerae*. Mol Microbiol 113:89-102. doi:10.1111/mmi.14402.
13. Helliwell KE, Lawrence AD, Holzer A, Kudahl UJ, Sasso S, Kräutler B, Scanlan DJ, Warren MJ, Smith AG. 2016. Cyanobacteria and Eukaryotic Algae Use Different Chemical Variants of Vitamin B_{12}. Curr Biol 26:999-1008. doi:10.1016/j.cub.2016.02.041.
14. Watanabe F, Nakano Y, Stupperich E. 1992. Different corrinoid specificities for cell growth and cobalamin uptake in *Euglena gracilis* Z. J Gen Microbiol 138:1807-1813. doi:10.1099/00221287-138-9-1807.
15. Barker HA, Smyth RD, Weissbach H, Toohey JI, Ladd JN, Volcani BE. 1960. Isolation and properties of crystalline cobamide coenzymes containing benzimidazole or 5,6-dimethylbenzimidazole. J Biol Chem 235:480-8. doi:10.1016/S0021-9258(18)69550-X.
16. Keller S, Kunze C, Bommer M, Paetz C, Menezes RC, Svatoš A, Dobbek H, Schubert T. 2018. Selective Utilization of Benzimidazolyl-Norcobamides as Cofactors by the Tetrachloroethene Reductive Dehalogenase of *Sulfurospirillum multivorans*. J Bacteriol 200. doi:10.1128/JB.00584-17.

17. Hallberg ZF, Nicolas AM, Alvarez-Aponte ZI, Mok KC, Sieradzki ET, Pett-Ridge J, Banfield JF,
Carlson HK, Firestone MK, Taga ME. 2024. Soil microbial community response to corrinoids is
shaped by a natural reservoir of vitamin B_{12}. ISME J 18. doi:10.1093/ismejo/wrae094.
18. Lawrence CC, Gerfen GJ, Samano V, Nitsche R, Robins MJ, Rétey J, Stubbe J. 1999. Binding of
Cob(II)alamin to the Adenosylcobalamin-dependent Ribonucleotide Reductase from
Lactobacillus leichmannii. Identification of Dimethylbenzimidazole as the Axial Ligand. J Biol
Chem 274:7039-7042. doi:10.1074/jbc.274.11.7039.
19. Hazra AB, Han AW, Mehta AP, Mok KC, Osadchiy V, Begley TP, Taga ME. 2015. Anaerobic
biosynthesis of the lower ligand of vitamin B_{12}. Proc Natl Acad Sci U S A 112:10792-7.
doi:10.1073/pnas.1509132112.
20. Kennedy KJ, Widner FJ, Sokolovskaya OM, Innocent LV, Procknow RR, Mok KC, Taga ME.
2022. Cobalamin Riboswitches Are Broadly Sensitive to Corrinoid Cofactors to Enable an
Efficient Gene Regulatory Strategy. mBio 13:e0112122. doi:10.1128/mbio.01121-22.
21. Mok KC, Sokolovskaya OM, Nicolas AM, Hallberg ZF, Deutschbauer A, Carlson HK, Taga ME.
2020. Identification of a Novel Cobamide Remodeling Enzyme in the Beneficial Human Gut
Bacterium *Akkermansia muciniphila*. mBio 11. doi:10.1128/mBio.02507-20.
22. Mok KC, Hallberg ZF, Procknow RR, Taga ME. 2025. Laboratory evolution of *E. coli* with a
natural vitamin B_{12} analog reveals roles for cobamide uptake and adenosylation in methionine
synthase-dependent growth. J Bacteriol doi:10.1128/jb.00284-24:e0028424.
23. Bannon CC, Mudge EM, Bertrand EM. 2024. Shedding light on cobalamin photodegradation in
the ocean. Limnol Oceanograph Lett 9:135-44. doi:10.1002/lol2.10371.
24. Tang YZ, Koch F, Gobler CJ. 2010. Most harmful algal bloom species are vitamin B_1 and B_{12}
auxotrophs. Proc Natl Acad Sci U S A 107:20756-61. doi:10.1073/pnas.1009566107.
25. Alvarez-Aponte ZI, Govindaraju AM, Hallberg ZF, Nicolas AM, Green MA, Mok KC, Fonseca-
García C, Coleman-Derr D, Brodie EL, Carlson HK, Taga ME. 2024. Phylogenetic distribution
and experimental characterization of corrinoid production and dependence in soil bacterial
isolates. ISME J 18. doi:10.1093/ismejo/wrae068.

272