Microbes display broad diversity in cobamide preferences

Kenny C. Mok[a], Olga M. Sokolovskaya[a,b], Adam M. Deutschbauer[a,c], Hans K. Carlson[c], Michiko E. Taga[a]#

[a]Department of Plant & Microbial Biology, University of California, Berkeley, Berkeley, CA 94720, U.S.A.

[b]Present address: Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, MA 02139, U.S.A.

[c]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, U.S.A.

#Correspondence to: taga@berkeley.edu

Abstract

Cobamides, the vitamin B_{12} (cobalamin) family of cofactors, are used by most organisms but produced by only a fraction of prokaryotes, and are thus considered key shared nutrients among microbes. Cobamides are structurally diverse, with multiple different cobamides found in most microbial communities. The ability to use different cobamides has been tested for several bacteria and microalgae, and nearly all show preferences for certain cobamides. This approach is limited by the commercial unavailability of cobamides other than cobalamin. Here, we have extracted and purified seven commercially unavailable cobamides to characterize bacterial cobamide preferences based on growth in specific cobamide-dependent conditions. The tested bacteria include engineered strains of *Escherichia coli*, *Sinorhizobium meliloti*, and *Bacillus subtilis* expressing native or heterologous cobamide-dependent enzymes, cultured under conditions that functionally isolate specific cobamide-dependent processes such as methionine synthesis. Comparison of these results to previous studies of diverse bacteria and microalgae revealed that a broad diversity of cobamide preferences exists not only across different organisms, but also between different cobamide-dependent metabolic pathways within the same organism. The microbes differed in the cobamides that support growth most efficiently, those that do not support growth, and the minimum cobamide concentrations required for growth. The latter differ by up to four orders of magnitude across organisms from different environments and by up to 20-fold between cobamide-dependent enzymes within the same organism. Given that cobamides are shared, required for use of specific growth substrates, and essential for central metabolism in certain organisms, cobamide preferences likely impact community structure and function.

Importance

Nearly all bacteria are found in microbial communities with tens to thousands of other species. Molecular interactions such as metabolic cooperation and competition are key factors underlying community assembly and structure. Cobamides, the vitamin B_{12} family of enzyme cofactors, are one such class of nutrients, produced by only a minority of prokaryotes but required by most microbes. A unique aspect of cobamides is their broad diversity, with nearly 20 structural forms identified in nature. Importantly, this structural diversity impacts growth, as most bacteria that have been tested show preferences for specific cobamide forms. We measured cobamide-dependent growth in several model bacteria and compared the results to previous analyses of cobamide preference. We found that cobamide preferences vary widely across bacteria, showing the importance of characterizing these aspects of cobamide biology to understand the impact of cobamides on microbial communities.

B[12] and other cobamide cofactors are required by organisms in all domains of life (1). Only certain bacteria and archaea synthesize cobamides (2), while others must acquire them exogenously. Thus, cobamides are considered shared nutrients within microbial communities. A unique aspect of cobamides is their structural diversity. While B[12] (cobalamin, Cbl) is a well-known vitamin important for human health, nearly 20 cobamides with alternative lower ligands exist in nature (Fig. 1A) (3-5). Though research with alternate cobamides is limited because they are commercially unavailable, cobamide-dependent enzymes and organisms have distinct preferences for different cobamides (6-16). Further, addition of different cobamides to soil or soil-derived enrichment cultures elicited distinct shifts in bacterial abundances, suggesting that cobamide structure influences bacterial growth at the community level (17). Thus, cobamide preference in bacteria is likely important for microbial community structure.

Here, we address four questions about cobamide use and preference: 1) Do different cobamide-dependent enzymes in the same bacterium have the same cobamide requirements? 2) How many, and which, cobamides can support growth? 3) Which cobamides are preferred, and do preferences differ across organisms? 4) How much cobamide do microbes need for growth, and do these requirements vary by enzyme, taxonomy, or environment? We addressed these questions by measuring growth of wild-type and engineered bacteria in different cobamide-dependent conditions with up to eight cobamides at a range of concentrations. Cobamide preference was defined based on the concentration that elicits half-maximal growth (EC_{50}), with lower values corresponding to more preferred cobamides. This *in vivo* assay encompasses the different aspects of cobamide utilization, including uptake, adenosylation, riboswitch-based regulation, and the cobamide-dependent enzymes themselves. We compared these measurements with those from published literature to gain a comprehensive view of cobamide preferences across taxa and environments.

Different cobamide-dependent enzymes in a single organism can have distinct cobamide preferences

First we compared the cobamide requirements of three cobamide-dependent processes in *Sinorhizobium meliloti*. *S. meliloti* produces Cbl for methionine synthase (MetH), methylmalonyl-CoA mutase (MCM), and ribonucleotide reductase (NrdJ), each of which can be tested separately for cobamide preference by altering the genetic background and growth substrates. MetH-dependent growth was supported by all cobamides tested, though EC_{50} values spanned three orders of magnitude (Fig. 1B, Table S1). The concentration requirements and preferences are similar to what we previously observed for MCM-dependent growth (8), except that [2-MeAde]Cba better supported MCM-dependent growth (Fig. 2). NrdJ-dependent growth required 25- and 15-fold higher Cbl and [Bza]Cba concentrations, respectively (Fig. 1B-C, Table S1), which suggests either more cobamide is required for NrdJ function or higher levels of NrdJ enzyme are necessary to support growth. The relative cobamide preferences for MetH- and NrdJ-dependent growth are similar, though phenolyl cobamides did not support NrdJ-dependent growth, consistent with NrdJ being a base-on enzyme (18) and phenolyl cobamides being unable to adopt the base-on conformation (Fig. 1A-C, Fig. 2). Cbl was the preferred cobamide for all three cobamide-dependent conditions, consistent with adaptation to endogenously produced Cbl (Fig. 2).

We used *Escherichia coli* to compare cobamide preferences of ethanolamine ammonia-lyase
(EAL) and MetH using engineered strains cultured under different conditions. EAL-dependent
growth required 5- to 78-fold higher cobamide concentrations than MetH-dependent growth and
showed different cobamide preferences (Fig. 1D-E, Table S1). [2-MeAde]Cba, produced by *E. coli*
when provided the precursor cobinamide (19), was most preferred for EAL-dependent growth but
less preferred by MetH. In contrast, Cbl was the least preferred cobamide that supports EAL-
dependent growth, but most preferred for MetH-dependent growth (Fig. 1D-E). [*p*-Cre]Cba did
not support EAL-dependent growth, consistent with EAL being a base-on enzyme (10). Together,
these results show that cobamide-dependent enzymes in the same organism can have distinct
cobamide preferences and requirements.

Cobamide use in a single process varies across bacteria

We next examined cobamide preferences for MetH-dependent growth in several additional
bacteria, including *Bacteroides thetaiotaomicron* and *Ruminococcus gnavus*, which have MetH
but lack cobamide-independent methionine synthase MetE, as well as *E. coli* and *Bacillus subtilis*
strains heterologously expressing MetH orthologs from *Vibrio cholerae* and *Priestia megaterium*,
respectively. In each organism, growth was supported by all or most of the cobamides tested,
suggesting there is promiscuity in cobamide uptake, adenosylation, regulation, and use by MetH
(Fig. 1E-J). However, EC_{50} values spanned two orders of magnitude among cobamides supporting
growth (Fig. 1, 2). While in all cases, either [Ade]Cba or a phenolyl cobamide was least preferred,
and Cbl was most preferred except by *B. thetaiotaomicron*, variability in cobamide preference was
observed across organisms (Fig. 1F-J). This variability is most apparent when contrasting *B.*
thetaiotaomicron and *B. subtilis* expressing *P. megaterium metH*, which did not grow with
phenolyl or purinyl cobamides, respectively, at any tested concentration (Fig. 1H-I). The cobamide
preferences of the latter strain are largely consistent with prior growth measurements at a single
cobamide concentration (20). In contrast, *Akkermansia muciniphila* showed no cobamide
preference due to its ability to remodel diverse cobamides to [Ade]Cba (Fig. 2) (21).

Expressing *V. cholerae metH* in *E. coli* afforded the opportunity to compare the cobamide
preferences of *E. coli* and *V. cholerae* MetH orthologs in the same intracellular environment.
Unlike *E. coli*, wild-type *V. cholerae* cannot use [Ade]Cba for MetH-dependent growth (12). Our
results suggest this is due to its poor use by MetH, as the EC_{50} for [Ade]Cba was nearly 300-fold
higher for *E. coli* expressing *V. cholerae metH* compared to its native *metH* (Fig. 1F-G, Table S1).
Overexpression of *E. coli metH* led to improved growth with the least preferred cobamides (Fig.
1E-F). This suggests MetH is limiting for *E. coli* growth with certain cobamides including
[Ade]Cba, and is consistent with our previous observation that mutation of the regulator *metR* or
the *metH* 5′ untranslated region improved growth of *E. coli* with [Ade]Cba (22).

Cobamide requirements in bacteria and microalgae span orders of magnitude and correspond to
environment

Comparison of these results to other studies that evaluated cobamide-dependent growth
demonstrated that taxonomically diverse microbes from different environments are variable in
their relative cobamide preferences (Fig. 2). Further, the lowest EC_{50} values for most soil bacteria

and microalgae are 1-2 orders of magnitude lower than those for human gut commensal bacteria
(Fig. 2), suggesting the former are adapted to survive at much lower cobamide concentrations. If
EC_{50} values are indicative of cobamide concentrations in these environments, the bioavailable
cobamide concentration can be estimated as 0.1 to 10 pM in aquatic environments, 1-10 pM in
soil, and 10-100 pM in the human gut. Consistent with these values, Cbl has been detected at pM
concentrations in aquatic environments (23). 41 nM cobamide has been detected in soil, though
soil's robust ability to adsorb cobamides suggests not all may be bioavailable (17). Cobamide
concentrations in the gut regions that harbor the bacteria in this study have not been measured.

Together, these results demonstrate that cobamide preferences of enzymes in the same organism
and of cobamide-dependent growth across taxonomically diverse microbes are variable.
Furthermore, cobamide concentrations required for growth vary by orders of magnitude across
environments, suggesting microbes are adapted to cobamide levels in their environment, likely by
tuning the sensitivity of cobamide uptake, adenosylation, regulation, and use by cobamide-
dependent enzymes. The characterization of cobamide abundances, bioavailability, requirements,
and preferences are therefore necessary to understand the role of cobamide metabolism within
microbial communities.

Figure 1. Cobamide structures and cobamide-dependent growth. A) Structure of B_{12} (cobalamin), which contains the lower ligand 5,6-dimethylbenzimidazole, is shown in the base-on conformation in which the ring nitrogen is coordinated to the cobalt ion (dashed line) (left). Alternative lower ligands of cobamides in this study (right) are shown with the names of the cobamides given below each structure. (B-J) Cobamide dose-dependent growth assays showing OD600 measured at the indicated times for (B) MetH-dependent growth of *S. meliloti*, (C) NrdJ-dependent growth of *S. meliloti*, (D) EAL-dependent growth of *E. coli*, (E) MetH-dependent growth of *E. coli*, (F) MetH-dependent growth of *E. coli* expressing *E. coli metH* on a plasmid, (G) MetH-dependent growth of *E. coli* expressing *V. cholerae metH* on a plasmid, (H) MetH-dependent growth of *B. subtilis* expressing *P. megaterium metH*, (I) MetH-dependent growth of *B. thetaiotaomicron*, and (J) MetH-dependent growth of *R. gnavus.* The EC_{50} values calculated from the curves in B-J and genotypes of the engineered strains in B-H are shown in Table S1. OD_{600} values for [Bza]Cba-supplemented cultures of *B. subtilis* were recorded after 72 hr because growth was not observed until after 45 hr. Points represent the means of 3-6 biological replicates; error bars represent standard deviation.

Figure 2. Comparison of EC$_{50}$ values for cobamide-dependent growth. Organisms examined in the current study (bold) are compared with those from previous studies (8, 13, 21, 24, 25). *S. meliloti* Rm1021 Δ*nrdJ cobD*::*gus* Gm^R *metH*::Tn*5* pMSO3-*nrdAB*(*E. coli*) was used for *S. meliloti* MCM-dependent growth (8). EC$_{50}$ values for *S. meliloti* are higher than for the other tested microbes likely because wild-type *S. meliloti* synthesizes Cbl *de novo* and lacks a high-affinity cobamide uptake system. MetH-dependent growth of *C. reinhardtii* was tested in a *metE* mutant (13). Symbols show the mean EC$_{50}$ values. Capped bars represent 95% confidence intervals, except with organisms labeled with *, which indicates error as standard deviation. Error of *C. reinhardtii* and *P. lutheri* EC$_{50}$ values were not reported (13). Bars are uncapped on the left or right when lower or upper bounds for 95% confidence intervals could not be determined, respectively. The lower bound for *Blastococcus* sp. B12_004 grown with [5-OHBza]Cba is 10^{-7} nM (25). The base of the leftward and rightward arrows represents maximal and minimal concentrations for EC$_{50}$ from dose-response assays in which lack of growth or saturating growth was not reached, respectively. For *O. tauri* and *A. carterae*, EC$_{50}$ values could not be calculated, but the capped bars for [Ade]Cba show the upper and lower bounds (13). Symbols in the shaded region on the right represent cobamides that were unable to support growth at any concentration tested.

Acknowledgements

We thank members of the Taga lab for helpful discussions and Zoila Alvarez-Aponte and Rebecca Procknow for critical reading of the manuscript. We thank Kristen LeGault and Kim Seed for genomic DNA of *Vibrio cholerae* O1 biovar El Tor str. N16961. This research was supported by NIH grant R35GM139633 to M.E.T. A.M.D. and H.K.C. were supported by ENIGMA-Ecosystems and Networks Integrated with Genes and Molecular Assemblies (http://enigma.lbl.gov), a Science Focus Area Program at Lawrence Berkeley National Laboratory, and is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, under contract number DE-AC02-05CH11231.

References

1.   Sokolovskaya OM, Shelton AN, Taga ME. 2020. Sharing vitamins: Cobamides unveil microbial
interactions. Science 369:eaba0165. doi:10.1126/science.aba0165.
2.   Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, Haft DR, Taga ME. 2019. Uneven
distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative
genomics. ISME J 13:789-804. doi:10.1038/s41396-018-0304-9.
3.   Allen RH, Stabler SP. 2008. Identification and quantitation of cobalamin and cobalamin
analogues in human feces. Am J Clin Nutr 87:1324-35. doi:10.1093/ajcn/87.5.1324.
4.   Renz P. 1999. Biosynthesis of the 5,6-dimethylbenzimidazole moiety of cobalamin and of the
other bases found in natural corrinoids, p 557-566. *In* Banerjee R (ed), Chemistry and
Biochemistry of B12. John Wiley & Sons, Inc., New York.
5.   Yan J, Bi M, Bourdon AK, Farmer AT, Wang PH, Molenda O, Quaile AT, Jiang N, Yang Y, Yin Y,
Simsir B, Campagna SR, Edwards EA, Loffler FE. 2018. Purinyl-cobamide is a native prosthetic
group of reductive dehalogenases. Nat Chem Biol 14:8-14. doi:10.1038/nchembio.2512.
6.   Yi S, Seth EC, Men YJ, Stabler SP, Allen RH, Alvarez-Cohen L, Taga ME. 2012. Versatility in
corrinoid salvaging and remodeling pathways supports corrinoid-dependent metabolism in
Dehalococcoides mccartyi. Appl Environ Microbiol 78:7745-52. doi:10.1128/AEM.02150-12.
7.   Tanioka Y, Miyamoto E, Yabuta Y, Ohnishi K, Fujita T, Yamaji R, Misono H, Shigeoka S, Nakano
Y, Inui H, Watanabe F. 2010. Methyladeninylcobamide functions as the cofactor of methionine
synthase in a cyanobacterium, *Spirulina platensis* NIES-39. FEBS Lett 584:3223-6.
doi:10.1016/j.febslet.2010.06.013.
8.   Sokolovskaya OM, Mok KC, Park JD, Tran JLA, Quanstrom KA, Taga ME. 2019. Cofactor
Selectivity in Methylmalonyl Coenzyme A Mutase, a Model Cobamide-Dependent Enzyme.
mBio 10:e01303-19. doi:10.1128/mBio.01303-19.
9.   Shelton AN, Lyu X, Taga ME. 2020. Flexible Cobamide Metabolism in *Clostridioides*
(*Clostridium*) *difficile* 630 Δ*erm*. J Bacteriol 202:e00584-19. doi:10.1128/JB.00584-19.
10.   Poppe L, Bothe H, Bröker G, Buckel W, Stupperich E, Rétey J. 2000. Elucidation of the
coenzyme binding mode of further B_{12}-dependent enzymes using a base-off analogue of
coenzyme B_{12}. J Mol Catal 10:345-350. doi:10.1016/S1381-1177(00)00136-3.
11.   Poppe L, Stupperich E, Hull WE, Buckel T, Rétey J. 1997. A base-off analogue of coenzyme-B_{12}
with a modified nucleotide loop. ^{1}H-NMR structure analysis and kinetic studies with (*R*)-
methylmalonyl-CoA mutase, glycerol dehydratase, and diol dehydratase. Eur J Biochem 250:303-
307. doi:10.1111/j.1432-1033.1997.0303a.x.
12.   Ma AT, Tyrell B, Beld J. 2020. Specificity of cobamide remodeling, uptake and utilization in
Vibrio cholerae. Mol Microbiol 113:89-102. doi:10.1111/mmi.14402.
13.   Helliwell KE, Lawrence AD, Holzer A, Kudahl UJ, Sasso S, Kräutler B, Scanlan DJ, Warren MJ,
Smith AG. 2016. Cyanobacteria and Eukaryotic Algae Use Different Chemical Variants of
Vitamin B_{12}. Curr Biol 26:999-1008. doi:10.1016/j.cub.2016.02.041.
14.   Watanabe F, Nakano Y, Stupperich E. 1992. Different corrinoid specificities for cell growth and
cobalamin uptake in *Euglena gracilis* Z. J Gen Microbiol 138:1807-1813. doi:10.1099/00221287-
138-9-1807.
15.   Barker HA, Smyth RD, Weissbach H, Toohey JI, Ladd JN, Volcani BE. 1960. Isolation and
properties of crystalline cobamide coenzymes containing benzimidazole or 5,6-
dimethylbenzimidazole. J Biol Chem 235:480-8. doi:10.1016/S0021-9258(18)69550-X.
16.   Keller S, Kunze C, Bommer M, Paetz C, Menezes RC, Svatoš A, Dobbek H, Schubert T. 2018.
Selective Utilization of Benzimidazolyl-Norcobamides as Cofactors by the Tetrachloroethene
Reductive Dehalogenase of *Sulfurospirillum multivorans*. J Bacteriol 200. doi:10.1128/JB.00584-
17.

17. Hallberg ZF, Nicolas AM, Alvarez-Aponte ZI, Mok KC, Sieradzki ET, Pett-Ridge J, Banfield JF,
Carlson HK, Firestone MK, Taga ME. 2024. Soil microbial community response to corrinoids is
shaped by a natural reservoir of vitamin B_{12}. ISME J 18. doi:10.1093/ismejo/wrae094.
18. Lawrence CC, Gerfen GJ, Samano V, Nitsche R, Robins MJ, Rétey J, Stubbe J. 1999. Binding of
Cob(II)alamin to the Adenosylcobalamin-dependent Ribonucleotide Reductase from
Lactobacillus leichmannii. Identification of Dimethylbenzimidazole as the Axial Ligand. J Biol
Chem 274:7039-7042. doi:10.1074/jbc.274.11.7039.
19. Hazra AB, Han AW, Mehta AP, Mok KC, Osadchiy V, Begley TP, Taga ME. 2015. Anaerobic
biosynthesis of the lower ligand of vitamin B_{12}. Proc Natl Acad Sci U S A 112:10792-7.
doi:10.1073/pnas.1509132112.
20. Kennedy KJ, Widner FJ, Sokolovskaya OM, Innocent LV, Procknow RR, Mok KC, Taga ME.
2022. Cobalamin Riboswitches Are Broadly Sensitive to Corrinoid Cofactors to Enable an
Efficient Gene Regulatory Strategy. mBio 13:e0112122. doi:10.1128/mbio.01121-22.
21. Mok KC, Sokolovskaya OM, Nicolas AM, Hallberg ZF, Deutschbauer A, Carlson HK, Taga ME.
2020. Identification of a Novel Cobamide Remodeling Enzyme in the Beneficial Human Gut
Bacterium *Akkermansia muciniphila*. mBio 11. doi:10.1128/mBio.02507-20.
22. Mok KC, Hallberg ZF, Procknow RR, Taga ME. 2024. Laboratory evolution of *E. coli* with a
natural vitamin B_{12} analog reveals roles for cobamide uptake and adenosylation in methionine
synthase-dependent growth. bioRxiv doi:10.1101/2024.01.04.574217.
doi:10.1101/2024.01.04.574217.
23. Bannon CC, Mudge EM, Bertrand EM. 2024. Shedding light on cobalamin photodegradation in
the ocean. Limnol Oceanograph Lett 9:135-44. doi:10.1002/lol2.10371.
24. Tang YZ, Koch F, Gobler CJ. 2010. Most harmful algal bloom species are vitamin B_1 and B_{12}
auxotrophs. Proc Natl Acad Sci U S A 107:20756-61. doi:10.1073/pnas.1009566107.
25. Alvarez-Aponte ZI, Govindaraju AM, Hallberg ZF, Nicolas AM, Green MA, Mok KC, Fonseca-
García C, Coleman-Derr D, Brodie EL, Carlson HK, Taga ME. 2024. Phylogenetic distribution
and experimental characterization of corrinoid production and dependence in soil bacterial
isolates. ISME J 18. doi:10.1093/ismejo/wrae068.

	Cbl	[Bza]Cba	[5-MeBza]Cba	[5-OMeBza]Cba	[Ade]Cba	[2-MeAde]Cba	[p-Cre]Cba	[Phe]Cba
S. meliloti (MetH) (ΔnrdJ cobD::gus bhbA::Tn5 pMSO3-nrdAB(E. coli))	3.481 (2.548 - 4.672)	14.47 (12.76 - 16.32)			1077 (949 - 1216)	740.4 (< 808.5)	2159 (> 1971)	> 5000
S. meliloti (NrdJ) (ΔmetH cobD::gus, bhbA::Tn5)	88.69 (68.58 - 116.2)	223.2 (142.3 - 564.3)			> 2500	1830 (1223 - 7914)	NG	NG
E. coli (EAL) (ΔmetH)	3.676 (ND)	1.069 (0.9621 - 1.182)	2.176 (> 2.006)	1.442 (< 1.634)	1.042 (> 0.937)	0.946 (> 0.847)	NG	
E. coli (native MetH) (ΔmetE)	0.047 (0.037 - 0.059)	0.069 (0.060 - 0.080)	0.048 (0.040 - 0.056)	0.058 (0.050 - 0.066)	0.222 (0.208 - 0.237)	0.094 (0.085 - 0.105)	10.87 (> 10.01)	> 25
E. coli (E. coli MetH plasmid) (ΔmetE ΔmetH pETmini-metH(E. coli))	0.042 (0.035 - 0.049)	0.064 (0.057 - 0.073)	0.044 (0.037 - 0.052)	0.052 (0.046 - 0.059)	0.180 (0.168 - 0.193)	0.094 (0.086 - 0.102)	4.816 (4.414 - 5.270)	5.382 (4.723 - 6.113)
E. coli (V. cholerae MetH plasmid) (ΔmetE ΔmetH pETmini-metH(V. cholerae))	0.047 (0.041 - 0.053)	0.074 (0.063 - 0.085)	0.053 (0.048 - 0.059)	0.072 (0.065 - 0.080)	51.75 (47.11 - 56.92)	0.171 (0.154 - 0.188)	7.715 (7.020 - 8.568)	5.672 (5.206 - 6.177)
B. subtilis (P. megaterium MetH) (ΔqueG::Pveg-btuFCDR ΔmetE::loxP amyE::Pveg-metH(P. megaterium))	0.161 (0.141 - 0.184)	1.273 (< 6.495)	0.187 (0.169 - 0.208)	0.496 (0.441 - 0.557)	NG	NG	17.49 (15.99 - 19.21)	19.18 (16.52 - 22.35)
B. thetaiotaomicron (MetH)	0.219 (0.205 - 0.234)	0.788 (0.755 - 0.823)	0.049 (0.046 - 0.053)	0.195 (0.184 - 0.207)	1.728 (1.604 - 1.861)	0.030 (0.027 - 0.033)	NG	NG
R. gnavus (MetH)	0.039 (0.032 - 0.048)	0.056 (0.045 - 0.072)	0.049 (0.039 - 0.063)	0.139 (0.112 - 0.186)	0.121 (0.101 - 0.147)	0.078 (0.065 - 0.093)	> 2.5	> 5

Table S1. EC$_{50}$ values calculated from data in Figure 1 B-J. EC$_{50}$ values (nM) and 95% confidence intervals (in parentheses) are given for each growth condition shown in Figure 1B-J (four-parameter non-linear fit in GraphPad Prism (v9.5.1)). Genotypes are given for engineered strains. NG, no growth; empty, not measured. A greater than (>) value for EC$_{50}$ is an estimate for cultures that failed to reach saturation at any cobamide concentration tested. Greater than (>) and less than (<) symbols in confidence intervals were used when upper or lower bounds could not be determined, respectively; ND represents a confidence interval in which both upper and lower bounds could not be determined.