Spatiotemporal transcriptomic niches of complement pathway and serine protease inhibitor activation in aging and infection

Fabian Kern[1,2,&,*], Viktoria Wagner[2,&], Vanessa Wahl[3], Nicole Ludwig[2,4], Simon Graf[1,2,5], Jeremy Amand[2], Aulden Foltz[6], Micaiah Atkins[6], Blen Kedir[6], Alexandros Hadjilaou[7,8,9], Philipp Donate[1], Matthias Flotho[1,2], Friederike Grandke[2], Shusruto Rishik[2], Philipp Wartenberg[3], Monika I. Hollenhorst[10,11], Daniel Krug[1], Hanna Lotter[12], Rolf Müller[1], Ulrich Boehm[3,+], Oliver Hahn[6,13,+], Thomas Jacobs[7,+], Gabriela Krasteva-Christ[10,11,+], Tony Wyss-Coray[6,14], Andreas Keller[1,2,*]

Helmholtz-Institute for Pharmaceutical Research Saarland (HIPS), Saarland University Campus, Saarbrücken, Germany
Chair for Clinical Bioinformatics, Center for Bioinformatics, Saarland University, Saarbrücken, Germany
Pharmacology & Toxicology, Saarland University, Homburg, Germany
Core Facility Molecular Single Cell and Particle Analysis, Medical Faculty, Saarland University, Homburg, Germany
Saarbrücken Graduate School of Computer Science, Saarland University, Saarbrücken, Germany
Department of Neurology and Neurological Sciences, Stanford University School of Medicine, Stanford, CA, USA
Protozoa Immunology, Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany
Department of Neurology, Medical Center Hamburg-Eppendorf (UKE), Hamburg, Germany
Institute for Neuroimmunology and Multiple Sclerosis, Medical Center Hamburg-Eppendorf (UKE), Hamburg, Germany
Institute of Anatomy and Cell Biology, Saarland University, Homburg, Germany
Center for Gender-Specific Biology and Medicine (CGBM), Saarland University, Homburg, Germany
Molecular Infection Immunology, Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany
Calico Life Sciences LLC, San Francisco, CA, USA
The Phil and Penny Knight Initiative for Brain Resilience, Stanford University, Stanford, CA, USA
& These authors contributed equally
+ These authors contributed equally, sorted in alphabetical order
* Correspondence should be addressed to Andreas Keller (andreas.keller@ccb.uni-saarland.de) and Fabian Kern (fabianmichael.kern@helmholtz-hips.de).

Abstract

Aging is a multifactorial and complex physiological process, affecting every organ with

characteristic manifestations. Understanding the molecular mechanisms that drive

aging processes is crucial to targeting age-related disorders. Recent reports suggest

that severe post-infection syndromes can partially accelerate aging. However, the

underlying gene-encoded regulatory interplay, whether being shared or distinct

between aging and infection biology are poorly understood. Here, we employed spatial

transcriptomics to establish a multi-organ atlas (brain, heart, kidney, liver, lung, and

spleen) across the mouse lifespan (4, 17, and 26 months). Dissecting high-quality fresh-

frozen tissue samples at unbiased molecular resolution, we found both organ-specific

and cross-organ gene dysregulation upon aging. We identified age-related trajectories

in gene expression and cell state, some only detectable within their spatial context, and

provide validation at subcellular resolution. The most prominent effect was organ-wide

immune system activation with spatially variable severity. We therefore evaluated how

aging mimics the expression signatures observed in systemic infection, using spatial

transcriptomics slices from young mice infected with Plasmodium berghei ANKA. While

on the gene level the effect sizes caused by the infection outweighed those of aging, we

reveal a shared activation of the early complement pathway (C4b) and serine protease

inhibitors (Serpin gene family) within by phenotype distinct spatial niches. We show

that this common RNA signature is driven by tissue-specific cell types and eventually
affects protein levels in the aged brain, rendering them a target for future mechanistic
and drug discovery studies. Taken together, our study provides a coherent in-depth and
cross-organ transcriptomics atlas to systematically study aging and infection in the
mouse at spatiotemporal resolution.
25
Key highlights

- Large-scale and high-resolution atlas of spatial transcriptomics from six organs to study
aging and systemic infection across two mouse cohorts.
- Strong transcriptional alterations found in distinct organ-specific niches for aging and
acute malaria, with organ- and cell type-associated immune responses.
- Dysregulation of early complement proteases (C4b) and serine protease inhibitors
(Serpina3n) as common theme across central nervous system and peripheral organs.

Main

The population-wide risk of cardiovascular, neurodegenerative, and metabolic syndromes is inextricably connected to societal aging, which involves an organism-wide deterioration of tissue structure and function[1]. Previous genomics studies across multiple organs have examined molecular changes associated with aging and provided insights into unique trajectories occurring in both organs and individual cell types[2-5]. Single-cell RNA-seq approaches revolutionized our understanding of cell type-specific gene regulation in health and disease. The human and the murine brain have been of particular interest for the application of -omics approaches to identify molecular mechanisms in age-related cognitive decline and progressive neurodegeneration[6-9]. However, a detailed and spatially-resolved understanding of the molecular programs underlying the functional consequences of aging in the hundreds of mammalian cell types within and between organs is still missing. Recent advances in image- and sequencing-based spatial transcriptomics approaches have enabled us to profile the expression of thousands of genes from tissue slices in two- or three-dimensional space[10]. These approaches facilitate the discovery of fine-grained structural changes across tissue sections. Thereby, they overcome the existing limitations of single-cell RNA-seq approaches that typically miss the microenvironment, imposed by the inevitable tissue dissociation process leading to the loss of spatial information[11]. Among those, unbiased genome-wide approaches operating at the (super-)cellular level together with targeted, gene panel-based approaches of sub-cellular resolution have been developed. In the context of aging, spatial transcriptomics hold the promise of narrowing down the cell etiology, specifically by identifying early cellular hot spots of gene dysregulation[9].

The mechanistic interplay between the innate and adaptive immune systems in aging, for instance the observed chronic low-grade sterile inflammation ("inflammaging"), has been a long withstanding aspect in the field[12]. But, the task to discern cause (driver) from effect (passenger) of the aging process is still one of the major ongoing challenges, in particular the aspect how acute events might eventually promote chronic symptoms. In fact, an entire spectrum of inflammatory phenotypes of increasing severity exists, although their relationship to human aging is mostly unclear. The wealth of COVID-19 accelerated interdisciplinary research has pinpointed a potential causal connection between severe infectious diseases and aging-associated disease development on a global scale[13]. This innumerable data indicates common molecular immune pathways across the human body underlying both systemic aging and infection. Still, when and where such a potentially conserved biological mechanism is at play has remained unclear. Therefore, we performed 168 spatial transcriptomics experiments at an unprecedented scale utilizing two independent mouse cohorts, reporting 100 manually annotated spatial compartments based on genome-wide expression clustering of 383,040

high-quality barcoded sequencing spots. For in-depth validation of brain aging signatures, we used STOmics Stereo-seq, an unbiased genome-wide spatial transcriptomics technique, providing sub-cellular views on gene expression. We comprehensively compared organismal aging against systemic infection on the RNA level, both in the periphery and the central nervous system, and identified a shared mechanism of complement dysregulation occurring in specific local niches.

Spatial transcriptomics of aging mouse organs

We first performed spatial transcriptomics experiments using the spot-based 10x Visium platform on a wildtype mouse aging cohort, consisting of up to five mice per timepoint (young: 4 months; middle-aged: 17 months; aged: 26 months) followed by deep sequencing (**Fig. 1a**). For each individual mouse, we processed single sections for the peripheral organs (heart, kidney, liver, lung and spleen) and five sections from the brain (Bregma: –1.7, -2.06, -2.7, -3.52 and -4.72). Following a stringent image-based and sequencing metrics-based quality control, we focused our downstream analysis on 111 high-quality samples (**Extended Data Fig. 1-2**, **Supplementary Table 1**). With a tissue-specific average number, out of a total of 4992 spots available per capture area, we maximized the number of spots located under tissue (mean of 2522 before QC, 2441 after QC), while retaining a high level of reproducibility between replicates of the same organ and across all ages (**Fig. 1b**). The data set covered around 22.3 billion reads (~201 Mio. per sample) for an average of 3783 median genes detected at each spot (12,048 average of median UMIs). To this end, we selected histologically motivated cutting angles, trying to optimize capture of the inherent cellular heterogeneity of each peripheral organ (**Fig. 1c**). After filtering individual spot locations of lower quality (cf. Methods), we found specific properties of the spot clustering in our cleaned data set. While gene expression programs in spots of the structurally more defined organs like the kidney primarily clustered according to their spatial domain, the spot clustering in more homogenous tissues was overly driven by cell type-specific markers (**Fig. 1d, Supplementary Table 2**). In fact, the size and number of distinct expression clusters correlated with histological tissue complexity. We thus followed a hierarchical annotation approach, defining either a local region of gene expression (e.g. "Atria" in the heart), or if possible, naming each cluster after the most enriched cell type (e.g. "Periportal hepatocytes" in the liver) based on known marker genes.

For the peripheral organs, markers for most of the expected cell types were identified, and no major deviations in size and location of each cluster were observed between the aging groups (**Extended Data Fig. 3a-e**). Nevertheless, comparing the expression for each annotated cluster between young and old mice returned a differentiated picture, with the kidney showing more than ten times the number of significant differentially expressed genes (DEGs) than the

heart (**Fig. 1e**). Thereby, the peripheral organs each retained an almost disjunct aging-driven signature of dysregulation with notably varying effect sizes (**Fig. 1f, Extended Data Fig. 3f-j, Supplementary Table 3**). Interestingly, the peripheral gene expression clusters with the most DEGs were those known to harbor many immune cells (**Fig. 1g**). Yet, we found strong aging DEGs in each organ directly related to tissue function (**Fig. 1h**), such as genes of the myosin heavy chain in the heart (Myh7), members of the cytochrome P450 family in the kidney (Cyp2d12), or major urinary protein coding genes in the liver (Mup7). We wondered however, whether tissue-specific DEGs exhibited a spatially variable domain, independent of our supervised spot annotation. Thus, we determined the list of highly significant and reproducible spatially variable genes (SVGs) for each organ without making use of the spot cluster labeling (cf. Methods, **Extended Data Fig. 4a, b**, **Supplementary Table 4**). Similarly to the DEGs, the number of SVGs was organ-specific and we found that per tissue between 5% (spleen) and 24% (kidney) of the genes were both differentially expressed and spatially variable (**Extended Data Fig. 4c**). Still, except for liver and spleen, the number of SVGs clearly outweighed the number of aging DEGs. As one example, the prolactin hormone receptor (Prlr), a type I cytokine receptor, was induced upon aging specifically in the inner and outer stripe of the outer kidney medulla (**Fig. 1i**). Prolactin accumulation was previously reported as a sign of renal failure[14]. Another way to leverage the spatial resolution of the data is by investigating whether the local neighborhood, i.e. tissue microenvironment depends on the age (cf. Methods), since we found only minor changes in cluster proportion along the aging trajectory. Indeed, we again found tissue-specific neighborhood alterations, such as a progressive depletion of fibroblasts in the myocardium of the aging heart (**Fig. 1j**), or the loss of Kupffer cells in adjacency to periportal hepatocytes in the aging liver (**Extended Data Fig. 4d-g**). The immune system appeared to be implicated in all peripheral organs with increasing age, suggesting a more systemic relationship, especially considering the autocrine effects of parts of the dysregulated transcriptome such as the Prlr gene, which is typically also mediated through the central nervous system.

Spatial aging signatures across the mouse brain

To better capture the structural and functional complexity of the murine brain, we decided to profile each sample at five different bregmata, using expert annotations from the Allen Mouse Brain Atlas for guidance (**Fig. 2a**). We observed well reproducible cluster gene signatures for each brain section (e.g. cortical layers I-Vi) with highly preserved marker expression between adjacent slides and used a previously established nomenclature to annotate the spot clusters[9] (**Fig. 2b**, **Supplementary Table 5**). Similar to the peripheral organs, differences in observed cluster proportions were driven by the bregma and not by age. There was however, a trend for an overall decreasing clustering complexity for lower bregma

(**Extended Data Fig. 5a-e**), due to the increasing area assigned to the midbrain structure. As
expected, the different brain sections shared most of the DEGs due to similar constituting
clusters, with a notable peak in Brain5, the only slice already covering parts of the cerebellum
(**Fig. 2c, d**). Overall, the global effect sizes of aging DEGs in the brain were comparable to the
peripheral organs, although much lower in their absolute number (**Extended Data Fig. 5f-j**,
Supplementary Table 6). Interestingly, across all bregmata the number of SVGs clearly
outweighed the number of DEGs by a factor of almost 100 (**Extended Data Fig. 6**,
Supplementary Table 7). Still, the 17 aging DEGs shared between all five brain sections
showed an enrichment of immune-system-related and disease-associated functions, such as
C1qa-c and Trem2, while also exhibiting the strongest dysregulation in the white matter tract
(**Fig. 2e**). Among those, 16 genes are also SVGs in the brain, demonstrating the need for
transcriptomics in spatial dimensions to better understand the directions of aging-driven
dysregulation. In particular, Egr1, Gfap and Trem2 are all known markers for neurons,
astrocytes and microglia, respectively, triggering the question whether certain cell types
specifically contribute on their own to the immune activation commonly observed during brain
aging.
160
To follow up on this observation, we validated our analysis of the super-cellular 10x Visium
brain experiments with the sub-cellularly-resolved Stereo-seq[15] technology developed by BGI
STOmics, sparing one mouse brain from each age group (**Fig. 2f**). To evaluate concordance
after quality control, we first aggregated the Stereo-seq spots into bins, artificially replicating
the size of Visium spots. This allowed us to reproduce most of the gene expression clusters
for each bregma and age but with slightly varying marker genes (**Fig. 2g, Extended Data Fig.**
7, Supplementary Table 8). All of the 17 shared brain aging genes from the Visium data set
were successfully reported as DEG in at least four of the five matching brain layers in the
Stereo-seq dataset, corroborating our findings (**Supplementary Table 9**). Next, we utilized the
specific capabilities of Stereo-seq to perform an image-based cell binning of all spots near
DNA fluorescence signals followed by cell type clustering and annotation. This enabled us to
differentiate several types of neurons from common glia cells in the spatial domain (**Fig. 2h**).
With our close-to-cell resolution (bin20), we confirmed the elevated expression of Trem2, a
well-known microglia activation marker[16], throughout the whole aged mouse brain (**Fig. 2i**).
Taken together our findings further suggest a progressive dysregulation of the innate immune
system that co-occurs with an increased spatially-dependent prevalence of major brain cell
types in proinflammatory states.
178
Because normal brain function relies on numerous neurochemical pathways in different
regions, presumable affected by aging through vascular-related oxidative stress, dysregulated

glycolysis, or neurotransmitter dysfunction[17-19], we prepared high-throughput spatial

metabolomics experiments for the aging cohort using the other brain hemispheres for a set of

tissue slices at Bregma 2.06 (cf. Methods). Each sample was then imaged in a 20 x 20 μm

matrix grid to obtain per-pixel peaks. Obtaining four replicates per age group, we detected

2713 unique features across the 12 slices after computationally aligning the transformed

MALDI timsTOF fleX mass readouts between samples, of which 97 peaks were reference-

annotated to known metabolites (**Supplementary Table 10**). After a stringent quality filtering,

909 mass peak features remained for down-stream analysis. Among the strongest fold-

changes (log(FC)=-9.7) between aged and young mouse brains we found N-Acetyl-L-aspartic

Acid, located in the regions of white matter, striatum as well as the deeper cortical layers, even

though statistical significance diminished after FDR correction (**Extended Data Fig. 8**). Thus,

we argue that existing spatial metabolomics platforms are not yet sensitive enough to

unbiasedly discover metabolome-wide changes in the context of brain aging, where often only

subtle changes occur over larger time differences, but are better suited to perform targeted

validation at cellular or sub-cellular levels[20]. The N-Acetyl-L-aspartic Acid molecule is a

required metabolic factor for enabling lipid and myelin synthesis in oligodendrocytes[21]. We

confirmed this finding in a comprehensive aging mouse brain atlas describing differential

metabolomics in individually dissected anatomical regions, where N-Acetyl-L-aspartic Acid

was also found be upregulated with aging, but specifically in the olfactory bulb[22].

Bulk and spatial transcriptomics of mouse organs in acute malaria

Severe infectious diseases (e.g. long COVID, tuberculosis, hepatitis, malaria) are

associated with accelerated biological aging of the immune system in humans[23-25]. To dissect

how organismal aging is (dis-)similar to systemic infection at the molecular level and which

parts of the immune system activation are shared at the gene and pathway level, we repeated

the 10x Visium experiments with a separate cohort of mice (cf. Methods). The infection cohort

consisted of five young mice infected with the mouse parasite *Plasmodium berghei*, an animal

model of cerebral malaria[26] and five age- and gender-matched young controls (**Fig. 3a**).

Because heart and spleen showed rather small effect sizes in the aging cohort, sections from

the infection cohort were processed for three peripheral organs (liver, kidney and lung), in

addition to two preselected brain regions at bregma –2.06 and bregma 3.89 (the olfactory bulb),

the latter of which is the very first affected brain structure in this model[26]. After QC, we retained

42 high-quality capture areas with comparable spot statistics as mentioned above (**Fig. 3b**,

Extended Data Fig. 9, **Supplementary Table 11**). The angles of histological slices were

analogously chosen to best capture the tissue complexity, yielding a highly similar number,

type and spatial distribution of spot clusters as for the aging cohort (**Fig. 3c, d**, **Supplementary**

Table 12). However, here we noticed significant differences in the spot cluster proportions

between the infected mice and controls. For example, the loss of blood vessels in kidney and
liver parallel to a massive increase of immune cells in the liver and lung (**Extended Data Fig.**
10a-e). In line with this observation, the total number of DEGs in the infection cohort was much
higher than for the aging cohort, with most being strongly upregulated after infection in a tissue-
specific manner, meeting our expectations of an acute inflammation resembling a much more
severe phenotype than aging (**Fig. 3e, f**, **Supplementary Table 13**). In particular the leukocyte,
blood and lymphatic vessels as well as endothelial spot clusters exhibited the most DEGs and
highest effect sizes across tissues (**Extended Data Fig. 10f-k**). In stark contrast to the aging
cohort, the reproducible number of SVGs exceeded the number of DEGs exclusively in the
brain, but with similar percentages of overlap per tissue, reaching from 7% in the olfactory bulb
to 19% in the kidney (**Extended Data Fig. 11a-c**, **Supplementary Table 14**). The substantial
expansion of connective tissue around the lung pulmonary veins together with a massive
leukocyte invasion represented an intriguing phenotype alteration of the infected lung, which
is immediately visible from the spatially annotated gene expression data alone (**Fig. 3g**). All
three peripheral organs underwent remarkable changes in their spatial microenvironment, like
the increased proximity between Kupffer and other immune cells in the infected liver, as
revealed through a differential spot neighborhood analysis (**Extended Data Fig. 11d-f**).

Since *P. berghei* causes cerebral malaria and strongly affects the olfactory bulbs, we
investigated the intersection of SVGs and DEGs more closely to understand the underlying
mechanism. In general, many immune cell surface genes from the clusters of differentiation
were significantly upregulated at specific locations in the infected olfactory bulbs, including
Cd151, Cd164, Cd1d, Cd200, Cd24a, Cd44, Cd52, Cd55, Cd63, Cd74, Cd82, Cd9, and
Cd99l2. Moreover, it appeared that antimicrobial factors such as Cystatin C (Cst3) were highly
upregulated in the center of the infected olfactory bulb. This dysregulation pattern completely
diminished toward the outer layers, thus showing an intriguing spatial expression pattern (**Fig.**
3h). Because *P. berghei* is a eukaryotic pathogen, we exploited the fact that both host and
pathogen transcribe polyadenylated messenger-RNA, such that in principle both should be
captured through the reverse poly-T tag in 10x Visium 3' gene expression libraries. Indeed, we
were able to map a certain percentage of high-quality reads from each infected sample against
the *P. berghei* ANKA genome, with olfactory bulb, liver, and lung exhibiting the highest
percentage of cells with pathogenic reads (**Fig. 3i**). We were thus able to narrow down the
position of individual parasites within the tissue by following the distribution of pathogenic reads
into the different spot expression clusters (**Extended Data Fig. 12**). Our analysis further
revealed that they preferentially reside in the outer layers of the olfactory bulb, rendering the
expression of Cst3 reciprocal to the distance between center and parasite location. We then
identified the spots in the olfactory bulb clearly enriched for parasite reads and compared the

gene expression of each such center spot with its six direct neighbors in the hexagonal spot
array on 10x Visium capture areas. Neighbors of a parasite-enriched center spot exhibited a
significant activation of the cytokine CC-chemokine ligand 5 (Ccl5) as well as the cluster of
differentiation 74 (Cd74), indicating an acute T cell response around the pathogens. Such a
severe immune system activation is a necessary means to fight infections.

Alteration of mRNA expression is modulated by several independent mechanisms, but partially
explained by post-transcriptional gene regulation. We previously showed that both tissue-
specific and global microRNA (miRNA) regulators are either activated or deactivated upon
aging in the mouse[4]. We therefore wondered whether such a regulatory interplay also explains
the spatial immune activation observed during infection and performed small non-coding RNA
sequencing for the infection cohort (cf. Materials and Methods, **Supplementary Table 15**). We
revealed that variability of miRNA expression is strongly driven by tissue origin and not by
technical bias (**Extended Data Fig. 13a, b**). In total, 542 miRNAs were well expressed in at
least one tissue and within one of the experimental groups (**Extended Data Fig. 13c**). The
observed expression was highly correlated among samples from the same tissue (**Extended
Data Fig. 13d**). Surprisingly, none as well as only one of the miRNAs remained significantly
differentially expressed for brain and and bulb (mmu-miR-3473b, up), respectively (**Extended
Data Fig. 13e, f**). For the peripheral organs however, we obtained 46 up-regulated miRNAs in
kidney, 188 up-regulated and 12 down-regulated miRNAs in liver and again none for lung
(**Extended Data Fig. 13g-i**), suggesting once more a tissue-specific gene regulatory
mechanism to be involved in the systemic immune response to malaria disease. Looking into
their potential function, we found the target gene Cd28 (adj. p=0.0021318), an essential factor
for T-cell proliferation and cytokine production and the GO terms "positive regulation of
inflammatory response to antigenic stimulus" (adj. p=0.0029171), and "positive regulation of
peptidyl-serine phosphorylation" (adj. p=0.0029986). For the up-regulated liver miRNAs we
found an enrichment of miRNA targets controlling "angiogenesis" (adj. p=4.79e-7) and
"macrophage differentiation" (adj. p = 1.58e-6). All of these observations are in line with our
hypothesis that miRNAs mediate peripheral immunity specifically and locally, triggering distant
cellular interaction in the central nervous system and further trans-dysregulation of brain or
other spatially restricted mRNAs. Still, it raises questions as to whether such an inflammatory
process causes permanent organ damage and how in contrast the aging-induced immune
activation may initially cause more attenuated but, in the end, similar disease phenotypes.

Shared spatial dysregulation in aging and systemic infection
To better understand similarities and differences between organismal aging and
systemic infection using our high-resolution spatiotemporal transcriptomics readouts, we

correlated the effect sizes between aging and infection cohorts for the same tissues and expression clusters. Here, the brain showed a remarkable resemblance between aging and infection, with the complement component C4b being one of the top upregulated genes (**Fig. 4a**). Similar trends were observed in the peripheral organs (**Supplementary Table 16**). C4b is part of the early complement pathway, specifically upregulated in the white matter tract of the aging brain, also upregulated in the infected brain but beyond the white matter area, thus showing a different spatial presence (**Fig. 4b**).

We then performed an over-representation analysis for all the cluster-specific DEGs shared between infection and aging cohorts, and obtained "Neutrophil degranulation" and "Regulation of complement cascade" as top significantly enriched categories for brain, kidney, liver and lung (**Fig. 4c**, **Supplementary Table 16**). To validate this finding, we first created a gene signature feature for the complement pathway to measure its activity through gene module scores. Indeed, the spatially restricted activity of the early complement pathway was a common molecular feature of dysregulation between organismal aging and systemic infection across multiple organs (**Fig. 4d**). Since the enrichment analysis yielded the "Regulation of complement cascade", we next wondered which gene regulatory components known from the literature are centered around C4b, revealing with STRING[27] a dense network of other complement genes but also two serine protease inhibitors (Serpina3n, Serping1) (**Fig. 4e**). Surprisingly, Serpina3n was one of the top dysregulated genes in the aging brain and also upregulated in the infected brain. Leveraging aggregated pseudobulk expression, we confirmed the almost perfect intra-sample correlation between C4b and Serpina3n across both cohorts and all organs in common (**Fig. 4f**). These findings suggested a central gene regulatory role between serin proteases involved in complement activation and serine protease inhibitors, mediating immune responses in two physiologically different processes that each depend on extrinsic and intrinsic stimuli. We then looked into the whole family of Serpin genes and discovered a tissue-specific dysregulation of its members in aging and infection (**Fig. 4g**). Subsequently, we made use of our sub-cellularly resolved Stereo-seq data sets to closely compare the complement activity against the activity of reactive astrocytes that drive parts of the brain aging signature and obtained a high degree of spatial correlation (**Fig. 4h**). Utilizing our respective marker-based cell bin annotations, we deconvolved the spatial pattern of correlated C4b and Serpina3n expression to primarily originate from both from Astrocytes and Oligodendrocytes. We confirmed this for the cell-marker enriched spots in the aging and infected brain Visium samples (**Extended Data Fig. 14a, b**). Moreover, we compared our findings with a previous study[4] describing in-depth cross-organ bulk miRNA and mRNA expression in aged mice, finding indeed multiple members of the Serpin gene family (Serpina1f, -3f, -3g, -3k) as anticorrelated targets of aging-associated miRNAs in several

tissues. Finally, we externally validated a significant aging-associated upregulation of C4b and

Serpina3n on the protein level in the mouse brain cortex and hippocampus, using a previously

published proteomics data set[28] (**Extended Data Fig. 14c**).

In sum, our findings indicated a cross-organ gene regulatory mechanism as part of the immune

system, being centered around serine proteases (**Extended Data Fig. 15**). This dysregulated

pathway appeared to be orchestrated in each organ in a cell type-specific and spatiotemporal

manner and translated from the transcriptional level into differential protein abundances,

increasing the level of confidence for functional relevance.

Discussion

Understanding the heterogeneous and physiological consequences of mammalian

aging is a challenging task. Our large-scale spatiotemporal cross-organ study provides new

molecular insights into the underlying mechanisms. Several spatially and temporally resolved

genomic data sets on the mammalian brain[9,29-33], olfactory bulb[34-36], heart[37-46], kidney[47-51],

liver[52-54], lung[55-61], and spleen[62-65] have been published, each describing their molecular

architecture in health and disease. However, most of them lacked a sample size sufficient to

compare multiple experimental groups and organs in parallel, especially lacking from the same

donor[66]. Large sequencing studies are inherently challenging to perform and thus carry certain

risks of producing unwanted batch effects[67]. We tackled these challenges by utilizing two

separate breeds of wildtype mice, one from the United States for the aging cohort and one

from Germany for the infection cohort, while ensuring a minimum number of three replicates

per experimental group. The aging cohort was processed first and fully completed before

continuing with the infection cohort to prevent potential sample cross-contamination.

Our findings suggest shared physiological processes between aging and infectious diseases,

at least in malaria, that are progressively dysregulated and are centered around complement

activation and serine protease inhibitors. C4b and Serpina3n were previously reported to be

upregulated in the subventricular zone of the aged mouse brain and as being indicative of a

reactive type of oligodendrocyte[68]. Yet, Hajdarovic et al. recently reported an activation of C4b

in reactive Astrocytes from the female hypothalamus and hippocampus, suggesting a more

general, sex-specific role of glia cell types in mediating a neuroinflammatory or autoimmune

phenotype[6,69]. Moreover, Allen et al. narrowed down C4b upregulation in the aged mouse brain

in both astrocytes and oligodendrocytes that was further escalated by LPS treatment [7], also

corroborating our findings. Others reasoned that such a phenotype could solely arise from the

body-wide accumulation of senescent cells or an organ-dependent dysregulation of the

circadian clock, two other hallmarks of aging[12,70]. In humans, C3 and C4 levels were reported

to be anti-correlated with centenarian longevity[71]. It therefore remains intriguing, having

perceived an almost perfect correlation between C4b and Serpina3n across the aging and

infection cohorts, to raise further questions about the mutual gene regulation of complement

proteases and protease inhibitors, and conceivable downstream effects. Proteases play an

important role in innate immune defense mechanisms, but need to be tightly regulated in order

to prevent autoimmunity and self-attack, and thus have already been brought in connection

with age-related decline and neurodegeneration[72]. Within the complement system, C4 is

cleaved through the serin protease activity of C1 in the classical pathway and through MBL-

associated serine proteases (MASPs) within the lectin pathway[73]. Both proteases can be

inhibited by Serping1, a serin protease inhibitor[74], which is upregulated in the aging mouse

heart and kidney as well as during malaria infection of kidney, brain, olfactory bulb, liver and

lung. We were able to confirm for cortex and hippocampus that indeed the upregulation of C4b

and Serpina3n RNA transcripts results in increased protein abundance. Besides, Serpina3n is

another dysregulated serine protease inhibitor of chymotrypsin, like Serping1, but is mostly

known for the regulation of Cathepsin G[75]. After glycosylation, Serpina3n can also bind double-

stranded DNA in the nucleus and cause chromatin condensation[76]. Whether the observed

correlation of C4b and Serpina3n is explainable through its regulatory capacity within the

nucleus can only be determined in further proteomics studies with functional validation assays.

In general, looking beyond the complement system, the distribution of lymphoid and myeloid

cells in aging tissues, as well as their contribution to sterile inflammation during the aging

process is being increasingly recognized as highly complex and tissue-dependent[77]. Our data

as well as previous reports[78] point toward a cross-communication of native tissue parenchymal

cells with lymphoid cells that is implicated in the different tissue-specific aging niches, for

instance the white matter compartment in the brain or the lung alveoli. A more targeted and

deeper approach using single-cell or spot-free spatial *in situ* mapping techniques to quantify

the contribution of myeloid and lymphoid cells toward the development of shared aging- and

infection-driven niches and how this is mediated through the peripheral systems, is needed.

And even though the high-throughput analysis of the aging brain metabolome has neither

directly confirmed nor invalidated our findings on C4b, we have detected almost 1000 yet

uncharacterized mass feature peaks exhibiting strong fold-changes in spatially restricted

zones. Further research is needed to decode their individual structure and function, leaving

high potential for a comprehensive understanding of the metabolome and its role in brain aging.

Moreover, it remains to show whether the effects described here occur in the same or similar

way in human tissue, given the expression of cell markers obtained with single-cell

experiments was already shown to be partially discordant between species[79]. In fact, an

evolutionary discrepancy determines in any case the baseline chances of success to carry

over our findings into new therapeutic strategies targeting aging and infectious diseases. With

the establishment of improved spatial transcriptomics platforms capable of resolving RNA

expression at the sub-cellular level, we will be able to better narrow down the origins of

individual RNA classes and molecules, for instance whether they are enriched in the nucleus

or the cytosol. This will help to resolve missing details about the so far hidden cell to cell

crosstalk involved in complement biology as just recently suggested[80]. Our findings position

individual members of the complement system and protease inhibitor families as potential

broad therapeutic targets in progressive aging and infectious disease research. Such targets

are desperately needed to tackle pathogenicity, as for example cerebral malaria is still lethal

for approximately 20% of affected patients, with many survivors being affected by prolonged

neurocognitive impairment after primary therapy[81]. We therefore reason that a complement

activation-based pathomechanism plays an important role in autoimmunity-related tissue

damage that in turn contributes to diverse aging- and infectious disease-related phenotypes.

Beyond aging research, single-cell and spatial transcriptomics are pivotal techniques to study

the host-pathogen interface and reveal specific biological mechanisms such as cell invasion

or clearance[82,83]. While most bacteria and virus specific protocols still have to be developed in

order to dual-detect both host and pathogen from the same sequencing sample, multicellular

parasite species can be directly sequenced due to the same type of polyadenylated

messenger-RNA as in the host[84]. This compatibility was already broadly used to describe the

organismal development of *P. berghei*, mechanisms behind its transmission as well as specific

effects on gene expression induced in individual organs and cell types[65,85-91].

We initially combined the existing literature knowledge on individual tissues and cell types as

well as several independent single-cell and spatial studies to carefully annotate the spot gene

expression clusters needed for the downstream analysis. Unfortunately, there exists no

common data annotation standard in the community, even though a few interesting

nomenclatures and solutions to single-cell reference atlases were already proposed[92-94]. Our

data set is specifically designed to foster spatiotemporal driven research into specific organs

and cell types that are of particular interest in the aging and infection research communities,

but also beyond to serve as a reference data set for any other biomedical domain. However,

analyzing almost 200 Visium samples in an integrative approach brings also significant

computational challenges. Many tools are not ready to scale up to large samples sizes,

requiring extensive compute infrastructure and workflow optimizations. Therefore, we

anticipate that our data set is also beneficial to the computational methods community,

fostering the development of novel and scalable data analysis methods in the spatial

transcriptomics domain.

Figures

Figure 1: Spatial transcriptomics of the aging mouse.

a, Illustration of the experimental workflow conducted for the mouse aging cohort.

b, Distribution of the number of cleaned spots under tissue per organ and age. Average number of cleaned spots located under tissue varied between organs (Brain1: 2825, Brain2: 2839, Brain3: 2764, Brain4: 2534, Brain5: 1939, Heart: 3093, Kidney: 3342, Liver: 2824, Lung: 1926, Spleen: 759), out of the 4992 spots possible per capture area. In general, the number of spots per sample was influenced by slight inter-individual variances in organ size. Colored boxes span the first to the third quartile with the line inside the box representing the median value. The whiskers show the minimum and maximum values or values up to 1.5-times the interquartile range below or above the first or third quartile if outliers are present (shown as separate, colored dots).

c, Illustration of cut angle plane and one representative H&E-stained capture area per peripheral organ.

d, From left to right: Two-dimensional UMAP representation of colored spot clusters computationally integrated by organ (top to bottom), pie chart showing the proportion of annotated clusters across all samples, one representative annotated Visium sample with the spot cluster identities plotted over the H&E-stained tissue image.

e, Number of differentially expressed genes per aging peripheral organ (old vs. young) and direction of dysregulation. Total DEG counts were derived across all organ clusters, without removing duplicates.

f, UpSet plot comparing the unique gene sets across all organ cluster DEGs shown in (**e**).

g, Bar plot showing the number of DEGs per annotated spot cluster. Tissues sharing the same cluster names are stacked on top of each other.

h, Heatmap showing scaled expression of representative aging DEGs (rows) across all pseudobulk samples (columns).

i, Normalized expression of Prolactin receptor (Prlr) in the kidney across two spot clusters and three aging groups (left) and representative spatial transcriptomics samples from each aging group with Prlr expression overlaid (right).

j, Clustered heatmap showing differential neighborhood distance analysis for clusters of the aging heart (old vs. young).

Figure 1: Spatial transcriptomics of the aging mouse

a
5 wildtype mice per age group (N=15) ♂
X
6 organs (10 cutting planes)
Brain (5 Bregmata)
Heart
Kidney
Liver
Lung
Spleen
= 150 capture areas
10x Visium protocol (FF)
111 areas after QC
4 (Y)
17 (M)
26 (O)
months

b
Anterior → Posterior   Organ size →
Brain1 Brain2 Brain3 Brain4 Brain5 Heart Kidney Liver Lung Spleen
Number of spots
3000
2000
1000
Young Middle Old
Age

c
Organ | Cutting plane | Example
Heart
Kidney
Liver
Lung
Spleen

d
Spot clustering | Cell type / region annotation | Example

Heart
UMAP 2 / UMAP 1
Annotated cluster (N=14)
Myocardium
Blood cells
Ventricular cardiomyocytes
Vascular endothelium
Atria
Tissue border
Fibroblasts (Connective tissue)
Immune cells
Endocardium
Smooth muscle cells
Lymphatic endothelium
S100A8/A9+ Cardiomyocytes
Connective tissue
Responding immune cells

Kidney
Annotated cluster (N=15)
Inner medulla
Cortex
Inner stripe of outer medulla
Papilla
Proximal tubuli S2/S3 (Cortex)
Blood / immune cells
Outer stripe of outer medulla
Proximal tubuli S2 (Cortex)
Proximal tubuli S1 (Cortex)
Blood vessels
Podocyte in renal corpuscle (Cortex)
SMCs and FiBs (interstitium and vasculature)
Distal tubular cells (Cortex)
Border adipose tissue
Medullary rays (Cortex)

Liver
Annotated cluster (N=7)
Periportal hepatocytes
Midlobular hepatocytes
Pericentral hepatocytes (Glu+)
Blood vessels / Connective tissue
Kupffer cells
Immune cells
Rsad2+ Cmpk2+ cells (Sinusoide)

Lung
Annotated cluster (N=13)
Alveolar region
Secretory cells (Epithelial layer)
AT1 / AT2 cells (Epithelial layer)
Pulmonary vascular endothelium
Smooth muscle cells
Pulmonary lymphatic vessels
Blood / immune cells
Plasma cells
Deuterosomal and ciliated cells (Respiratory epithelium)
AT1 cells (Epithelial layer)
Mesothelial cells
Platelets
Tissue border

Spleen
Annotated cluster (N=6)
Red pulp zone
B cell zone
Erythrocytes
T cell zone
Plasma cells
Megakaryocytes

e
Number of DEGs
Heart Kidney Liver Lung Spleen
162 30 | 2028 417 | 220 572 | 329 643 | 122 856
Age dysregulation (Old vs. Young)
Up
Down

f
614 420 121 105 69 62 54 51 36 26 24 21 18 17 11
Kidney Up 821
Spleen Down 517
Lung Down 221
Liver Down 143
Lung Up 108
Kidney Down 95
Heart Up 81
Liver Up 67
Spleen Up 56
Heart Down 10

g
Number of DEGs
600
400
200

h
Heart 17 / Heart 73 / Heart 112 / Kidney 17 / Kidney 73 / Kidney 112 / Liver 17 / Liver 73 / Liver 112 / Lung 17 / Lung 73 / Lung 112 / Spleen 17 / Spleen 73 / Spleen 112
Myh7
Cyp2d12
Mup7
Bpifa1
Acap1
Scaled pseudobulk expression
2
1
0
-1

i
Prolactin receptor (Prlr)
Normalized expression (Kidney)
Inner stripe of outer medulla | Outer stripe of outer medulla
Age
17
73
112
Normalized expression
2.5
2.0
1.5
1.0
0.5
0.0

j
Fibroblasts (Connective tissue)
Atria
Myocardium
Connective tissue
Immune cells
Tissue border
Blood cells
Lymphatic endothelium
Vascular endothelium
Smooth muscle cells
Responding immune cells
Ventricular cardiomyocytes
Endocardium
S100A8/A9+ Cardiomyocytes
40
20
0
-20
-40

Figure 2: Spatial transcriptomics across the aging mouse brain.

a, Visualization of the five Bregmata selected to study different regions of the aging brain.

b, From left to right: Two-dimensional UMAP representation of colored spot clusters computationally integrated by brain slice (top to bottom), pie chart showing the proportion of annotated clusters across all brain samples, one representative annotated Visium sample with the spot cluster identities plotted over the H&E-stained tissue image.

c, Number of differentially expressed genes per aging brain bregma (old vs. young) and direction of dysregulation. Total DEG counts were derived across all organ clusters, without removing duplicates.

d, Five-dimensional Venn diagram comparing the brain DEG sets from (**c**).

e, Heatmap showing scaled expression of the 17 aging DEGs (rows) shared between all five brain slices, using the Brain1 pseudobulk samples and spot clusters for visualization (columns). All genes except Rbm3 are also significant SVGs in at least one of the five brain bregmata.

f, Sketch of 10x Visium and STOmics Stereo-seq examples comparing the features of both spatial transcriptomics technology platforms.

g, Examples for binned (bin200) and annotated spot clusters of the aging brain (top to bottom; young, middle, old) at Bregma#1 sequenced with Stereo-seq.

h, Dot plot showing the top 3 most significant marker genes per cell type annotated spot cluster using the Stereo-seq cellbin resolution of Brain1 samples.

i, Normalized spatial expression of Trem2 across all 15 STOmics Stereo-seq brain samples using the near-cellular resolution bin20 (from left to right: young, middle, old; from top to bottom: Brain1-5). For visualization spot sizes were rescaled into the point interval [0.1, 1.5] according to their expression of Trem2.

Figure 2: Spatial transcriptomics across the aging mouse brain

Figure 3: Spatial transcriptomics of the acute malaria-infected mouse.

a, Illustration of the experimental workflow conducted for the malaria infection cohort.

b, Distribution of the number of cleaned spots under tissue per organ and experimental group (controls, *P. berghei* infection). Average number of cleaned spots located under tissue varied between organs (Brain: 2760, Olfactory bulb: 722, Kidney: 3154, Liver: 3435, Lung: 3024), out of the 4992 spots possible per capture area. In general, the number of spots per sample was influenced by slight inter-individual variances in organ size. Colored boxes span the first to the third quartile with the line inside the box representing the median value. The whiskers show the minimum and maximum values or values up to 1.5-times the interquartile range below or above the first or third quartile if outliers are present (shown as separate, colored dots).

c, Illustration of cut angle plane and one representative H&E-stained capture area per organ.

d, From left to right: Two-dimensional UMAP representation of colored spot clusters computationally integrated by organ (top to bottom), pie chart showing the proportion of annotated clusters across all samples, one representative annotated Visium sample with the spot cluster identities plotted over the H&E-stained tissue image.

e, Number of differentially expressed genes per organ (day 6 post *P. berghei* infection vs. controls) and direction of dysregulation. Total DEG counts were derived across all organ clusters, without removing duplicates.

f, UpSet plot comparing the gene sets shown in (**e**).

g, In-depth zoom into a lung pulmonary vein from a control mouse (top) and an infected mouse (bottom), demonstrating the expansion of connective tissue and leukocyte invasion caused by the malaria disease. Spots are colored according to cluster identity, in particular Leukocytes in red, Smooth muscle cells in purple and Blood / immune cells in light purple.

h, Normalized expression of Cystatin C (Cst3) in all spot clusters of the olfactory bulb split by infection and controls with clusters ordered by their ascending distance to the center of the tissue (left). Trend lines were computed with loess regression of y ~ log(x). Shown on the right is the normalized expression of Cst3 in the spatial domain of one representative control (top) and infected mouse (bottom).

i, Percentage of reads per spot mapped to the *P. berghei* ANKA genome split by experimental group and organ.

Figure 3: Spatial transcriptomics of the acute malaria-infected mouse

Figure 4: Molecular niches shared between aging and systemic infection.

a, Scatter plot showing the average log2-scaled fold-change between aging and infection for all brain DEGs matched and colored by spot cluster.

b, Normalized spatial expression of C4b in four representative brain samples (Bregma: -2.06), two from aging (top left: young, top right: old) and infection (bottom left: control, bottom right: infected) cohort.

c, Dot plot showing adjusted and log-scaled Hypergeometric test p-values and the number of gene hits for the enriched categories from the Reactome pathway database across the four organs shared between aging and infection cohort. For each row of results a different list of genes was used as input (cf. Methods).

d, Relative (scored) complement pathway activity in pseudobulk samples across all aging and infection cohort organs, split and colored by the five experimental groups (young, middle, old, healthy controls, infected).

e, STRING network for Serpina3n in Mus musculus after performing one level of network expansion and removing edges from text mining and gene fusion. Edges are colored according to the type of association or interaction: curated databases (light blue), experientially determined (purple), gene neighborhood (green), gene co-occurrence (dark blue), co-expression (black), or protein homology (light purple). Nodes are colored according to their shell of interactions towards Serpina3n.

f, Scatter plot showing normalized pseudobulk expression of C4b (x-axis) against Serpina3n (y-axis) across all samples and colored by the five experimental groups (young, middle, old, healthy controls, infected).

g, Row and column clustered heatmap showing the scaled pseudobulk expression for all members of the serine protease inhibitor (Serpin*) gene family.

h, Demonstration of spatially connected (sub-)cellular activity of the complement pathway (C4b+) and activated Astrocytes (Gfap+) in the aging brain (Bregma -2.06). Normalized expression values from one young representative on the top and one old representative on the bottom originating from the Stereo-seq samples at bin20 resolution are displayed.

Figure 4: Molecular niches shared between aging and systemic infection

Methods

Mouse tissue processing and screening

For the aging cohort, male C57BL/6JN mice were shipped from the National Institute on Aging colony (Charles River) to the Stanford ChEM-H animal facility, where they were housed at 12/12h light/dark cycle in cages of 2-3 mice. Water and food were provided ad libitum. The aging cohort consisted of 5 mice per age group (age groups: 17, 73 and 112 weeks; 3.91, 16.8 and 25.78 months). Tissues from brain, heart, kidney, liver, lung and spleen were collected after anaesthetization with 2.5% v/v Avertin and transcardial perfusion with 20 mL cold PBS and flash frozen in isopentane cooled with liquid nitrogen, all between 10:15 AM- 12:00 PM on a single day. For brain samples, the right and left hemisphere were separated before freezing and stored separately. Samples were stored at -80°C until embedding in OCT. Samples were embedded for sectioning in precooled OCT on dry ice. All animal care and procedures complied with the Animal Welfare Act and were in accordance with institutional guidelines and approved by the institutional administrative panel of laboratory animal care at Stanford University.

For the infection cohort, experimental procedures were conducted as described previously[95]. In brief, C57BL/6 mice were bred at the animal facility and *P. berghei* ANKA parasites were obtained from the blood of sporozoite-infected C57BL/6 mice provided by the Parasitology Section of the Bernhard Nocht Institute. Mice in the control and experimental groups were age-matched (6 to 8 weeks old) and sex-matched (male & female). For the experimental group, C57BL/6 mice were infected i.p. with 1×10^5 parasitized red blood cells. Positive parasitemia was determined in Giemsa-stained blood smears from tail blood. Body weights were determined at different time points. All required organs were collected on two consecutive days, from 9:30 AM (Ctrl.) and 10:30 AM (day 6 post-infection) until 12 PM. To prevent lung deflation, the tracheae were sutured, and the lungs were inflated with a 1:1 mixture of PBS and OCT. The small lung lobe was separated and frozen separately from the residual lung tissue. Brains were frozen without separating the hemispheres. All tissue samples were simultaneously frozen while embedded in OCT, in an isopentane bath cooled with liquid nitrogen and kept stored at -80°C. All animal care and procedures complied with institutional guidelines and were approved by the administrative panel of laboratory animal care at the Bernhard Nocht Institute under the admission TVA N118/2020, as well as the Hamburg Animal Welfare agency.

Cryosectioning and section mounting were performed according to the manufacturer's protocol, in a Leica cryostat (LEICA CM3050 S) set to -15°C with a slice thickness of 16 μm (except for the spleen of the infection cohort; 10 μm). For the aging cohort, five brain sections were mounted using visual landmarks and the Allen Mouse Brain Atlas, resulting in sections from bregma –1.7, -2.06, -2.7, -3.52 and -4.72. For the infection cohort, two brain sections were mounted, namely from bregma –2.06 and bregma 3.89 (Olfactory bulb). Sections were mounted on designated capture areas and stored at -80°C until further processing. After tissue mounting on the slide, an additional 10-15 sections were collected for RNA Quality assessment. All samples used in this study had a RIN value between 4.4 and 9.0 (**Supplementary Table 1, 11**).

Spatial gene expression assay (10x Visium)

Following the Tissue Optimization protocol, we determined the permeabilization times to 18 minutes for the brain, 24 minutes for the olfactory bulb, 15 minutes for heart, 15 minutes for kidney, 16 minutes for liver, 20 minutes for lung, and 30 minutes for the spleen samples using the Tissue Optimization (TO) Slide & Reagent Kit (10x Genomics, PN-1000184).

We followed the standard protocol for fresh frozen tissues (compare document CG000239 - Rev F) using the Visium Spatial Gene Expression (GEX) Slide & Reagent Kits with recommended reagents (10x Genomics, PN-1000193). In brief, after H&E staining and imaging (10x magnification, Axio Imager KMAT - ZEISS), tissues were permeabilized with the previously defined permeabilization times. After reverse transcription on the slide and

denaturation, second strand synthesis is performed. For further processing, cDNA is released
via denaturation. To determine cycle number for amplification the KAPA SYBER FAST qPCR
Master Mix Kit (KAPA Biosystems) is used in a qPCR. All samples in this study were amplified
using 14 – 19 PCR cycles. After bead purification using SPRIselect beads (Beckman Coulter,
Product No. B233318), cDNA is quantified via 2100 Bioanalyzer using the Bioanalyzer High
Sensitivity DNA kit (Agilent, cat.no.:5067-4626).
653
The library construction was either performed manually or using the MGISP-100 liquid handling
system according to the 10x Visium user guide. In short, the following steps were performed:
fragmentation, end repair, A-tailing, double sided size selection, adaptor ligation, post ligation
cleanup, sample index PCR, double sided size selection; followed by post library construction
QC performed using the Bioanalyzer High Sensitivity DNA kit (Agilent, cat.no.:5067-4626) and
KAPA Library Quantification Kit (Roche, KK4824).
660
661   Libraries were pooled in an equimolar fashion with up to 15 samples in one library, multiplexing
662   only samples that originated from the same organ whenever possible. All libraries were
663   sequenced using NovaSeq 6000 (Illumina).
664

Spatial gene expression assay (STOmics Stereo-seq)
One brain sample of each age was processed at the MGI Tech Co., Ltd. (Riga, Latvia) using
the STOmics Stereo-seq Transcriptomics T Kit (MGI). For each brain, five 10µm sections
(Bregma: –1.7, -2.06, -2.7, -3.52 and -4.72) identical to the Visium cohort were collected and
loaded onto the Stereo-seq Chip T-Slide according to manufacturer´s protocol. In short, after
fixation of the tissue using ice-cold methanol, cell nuclei were stained using the ssDNA Qubit
dye (Thermo Fisher) and imaged in 10x resolution in FITC channel. Subsequently, brain tissue
was permeabilized for 12 min at 37°C, releasing mRNAs from the cells onto the Chip T-Slide.
Reverse transcription of mRNAs was conducted for 3 hours at 42°C generating cDNA with
sequence barcodes indicating their unique location on the T-Slide. Afterwards, the tissue was
removed using tissue removal buffer for 10 min at 55°C and cDNA was collected from the chip
o/n at 55°C. CDNA was then purified, amplified and size distribution was checked using High
Sensitivity DNA Bioanalyser Kit (Agilent). Library Preparation was performed using Stereo-seq
Library Preparation Kit (MGI) using 30ng of cDNA and 13 cycles of amplification following the
manufacturer´s protocol. Final quality check of the size distribution of the library was performed
using High Sensitivity DNA Bioanalyzer Kit. Libraries were sequenced on DNBSEQ-Tx
sequencer (MGI).
682

Small non-coding RNA profiling
Total RNA was isolated from 5-10 10µm tissue sections using the miRNeasy Mini Kit (Qiagen,
Hilden, Germany) according to the manufacturers protocol. RNA concentration and integrity
was determined using Nanodrop (ThermoFisher Scientific, Waltham, MA, USA) and RNA 6000
Nano Bioanalyzer Kit (Agilent, Santa Clara, CA, USA), respectively.
Libraries for miRNA expression profiling were generated with the MGIEasy Small RNA Library
Prep Kit (MGI Tech, Shenzhen, China) according to manufacturer's recommendations with
100ng totalRNA starting material. Sequencing was performed on an DNBSEQ-G400RS
instrument by the Sequencing Unit of the Core Facility Molecular Single Cell and Particle
Analysis of Saarland University using the 50bp single end sequencing strategy.
693

Image and data preprocessing
We stitched the image series from the Axio Imager KMAT brightfield microscope using the
Zeiss ZEN software (v3.6) and cropped and rotated the resulting high-resolution TIFF files
using Fiji. Individual sample images were color corrected where needed to match the other
images. In the next step, we ran Spaceranger (v2.1) with the demultiplexed FASTQ files and
the corresponding TIFF images. In brief, Spaceranger aligns the reads to a reference genome
(refdata-gex-mm10-2020-A), detects tissue under spots and the fiducial frame in the image, to
create a spot x UMI count matrix. For the infection cohort, we additionally built a custom
Spaceranger reference genome combining mouse (GRCm39) and *Plasmodium berghei* -

ANKA genome (PBANKA01). For two samples the alignment of the fiducial frame was
manually corrected after the automatic detection had failed initially.
705
706 In case of the STOmics Stereo-seq samples, images were obtained and stitched with the Zeiss
707 Moticam Pro 500i. Combined QC and primary analysis of sequencing and image data was
708 performed using the STOmics analysis workflow (SAW) v1.0 to obtain quantification matrices
709 for the mouse genome (GRCm38). The SAW pipeline was also used to accumulate DNB spot
710 gene expression counts into the specific area and cell binnings.
711
Bioinformatics and quality control
Based on the Spaceranger reports, we either excluded a sample completely from further
analysis, or using the 10x Loupe browser tool (v6.5.0) manually selected small spot regions to
exclude from downstream analysis where necessary (e.g. removing corners with uneven
permeabilization), and further applied the following criteria-based spot filtering on each sample,
to eventually obtain a high-quality data set. To this end, we inspected the variable distributions
for the number of UMIs, the number of read counts, mitochondrial content, as well as spot
complexity (=log10(Number of features) / log10(Number of counts)) for each tissue separately,
and applied the following filtering criteria on the filtered spot count matrix from Spaceranger
after loading into Seurat objects: (*Maximum percentage of mitochondrial read content;*
Minimum number of UMIs; Maximum number of UMIs; Minimum number of detected features;
Maximum number of detected features; Minimum spot complexity)
Brain: (35; 500; 70,000; 300; 10,000; 0.75), Bulb: (35; 500; 70,000; 300; 10,000; 0.75),
Heart: (60; 1000; 25,000; 300; 5000; 0.75), Kidney: (30; 1000; 50,000; 300; 8000; 0.75)
Liver: (15; 1000; 40,000; 500; 8000; 0.75), Lung: (10; 1000; 30,000; 300; 8000; 0.75)
Spleen: (10; 500; 35,000; 300; 8000; 0.75).
728
We then rotated the images of the remaining samples to best match the desired orientation for
each tissue. We then followed the default Seurat (v4.4) workflow to preprocess each sample
by applying *NormalizeData*, *FindVariableFeatures* with variance stabilizing transformation for
2000 features, followed by *SCTransform* (v1), regressing out the percentage of mitochondrial
reads and subsetting to 3000 variable features. Subsequently, we performed *RunPCA* on the
SCT assay, *FindNeighbors* on up to 30 PCs, *FindClusters* with leiden clustering followed by
RunUMAP with up to 30 dimensions. Using the cleaned list of Visium samples, we created one
Seurat object for each tissue using the SCT-based integration workflow as follows: set the
default assay to SCT, run *SelectIntegrationFeatures* with 3000 features, then
PrepSCTIntegration, and *FindIntegrationAnchors* followed by *IntegrateData*, each with "SCT"
set as normalization method and up to 30 dimensions. Next, we performed the standard
dimension reduction workflow consisting of *RunPCA*, *RunUMAP*, *FindNeighbors*, and
FindClusters again with up to 30 dimensions and fixed random seeds.
742
For the STOmics Stereo-seq samples, we first converted the outputs of the Stereo-Seq
Analysis Workflow (SAW) pipeline[96] (v1.0) into Seurat objects for both bin200 (200x200
Stereo-seq spots) and cell bins. This was accomplished using the *io.stereo_to_anndata*
function in the Stereopy package[97] (v1.1.0), followed by applying the *h5ad2rds.R* script
provided by BGI. Cell bins were obtained using image-based cell segmentation of the
corresponding ssDNA-stained images. Filtering was performed using the following criteria:
(*Maximum percentage of mitochondrial read content; Minimum number of UMIs; Maximum*
number of UMIs; Minimum number of detected features; Maximum number of detected
features; Minimum spot complexity)
Brain (bin200): (35; 500; 100,000; 300; 10,000; 0.75), Brain (cell bin): (35; 100; 10,000; 100;
3,000; 0.75).
754
In line with the Visium 10x samples we applied a default Seurat preprocessing workflow for
each sample. Using the cleaned list of STOmics Stereo-Seq samples, we performed the
Harmony-based integration workflow as follows: set the default assay to SCT, run
SelectIntegrationFeatures with 5000 features, merge the sample list into one Seurat sample,

run *RunPCA* with 50 components, run *RunHarmony* on SCT assay and the PCA reduction
using *Sample_ID* and *Age* as latent variables, run JoinLayers on the RNA assay. Then, we
performed the sample dimension reduction workflow as in the Visium samples with up to 50
dimensions.
763
Spot clustering annotation
For the Visium aging cohort we employed tissue-specific clustering resolutions to match their
individual complexity as follows: Brain1 0.8, Brain2 0.7, Brain3, 0.6, Brain4 0.5, Brain5 0.5,
Heart 0.5, Kidney 0.5, Liver 0.2, Lung 0.5, and Spleen 0.6. For the Visium infection cohort, we
set the clustering resolution for Brain to 1, Olfactory bulb to 0.6, Kidney to 0.5, Liver to 0.2 and
Lung to 0.5, aiming to reproduce about the same number and type of gene expression clusters
as for the aging cohort. In the next step, we computed the enriched cluster marker genes for
each tissue object in a one vs. all approach by using *FindAllMarkers* on the normalized data
slot of the spatial assay in combination with MAST[98], setting the minimal percentage of positive
cells to 0.1, requiring a minimal alpha to 0.01 and log fold-change of 0.1 for Heart, Kidney,
Lung, Spleen as well as 0.25 for brain and using the sample identity of each spot as latent
variable for gene regression. We then applied a hierarchical approach of annotating the
clusters using the top-enriched genes, trying to resolve at best the most-enriched cell type, if
possible, through the expression of clear marker genes, and annotating the spatial
compartment based on histological knowledge and functional properties instead. We made
combined use of transcriptomics literature on each organ, in particular single-cell and single-
nucleus reference data, reference atlases such as the Allen Mouse Brain atlas, as well as an
in-depth analysis of the stained tissue slices, to manually annotate all clusters with a
nomenclature used previously. Thereby, clusters having no distinctive but unclear markers but
showed shared marker expression overall to other clusters, were merged together.
784
For the STOmics Stereo-seq samples with bin 200 (200x200 DNB bins), we employed tissue-
specific clustering resolution aiming to approximately reproduce the number of clusters from
the Visium 10x aging cohort at each bregma as follows: Brain1 3.2, Brain2 3, Brain3 3, Brain4
2.5, Brain5 2.5. For stereo-seq samples with single-cell resolution we utilized a clustering
resolution of 0.5 across all Bregmata. In the next step, we computed the enriched cluster
marker genes for each tissue object in a one vs. all approach by using *FindAllMarkers* on the
normalized data slot of the spatial assay in combination with MAST, setting the minimal
percentage of positive cells to 0.1, requiring a minimal alpha of 0.01 and fold-change of 0.25
for all samples and using the sample identity of each spot as latent variable for gene
regression. This was accomplished equivalently for both bin200 and cell bins. For bin200, we
performed annotation by first calculating the marker overlap of each cluster with every cluster
from the corresponding Visium 10x samples. Here, we always considered the top 50 markers
selected by lowest adjusted p-value. To annotate a cluster, we then identified Visium clusters
with marker overlap and additionally took the anatomical location, via the Allen Mouse Brain
Atlas of the corresponding spots, into account. In most cases, this allowed us to pinpoint one
annotation for each cluster. In cases where multiple annotations we plausible, we split the
clusters based on spatial location and anatomical information. Clusters annotated with the
same label were merged together. For cell bins we aimed for a high-level cell type annotation
matching the limited counts per cell of the stereo-seq cell segmentation. We utilized the current
literature on single-cell and single-nucleus datasets as reference for our annotation together
with the anatomical location of the clusters. Spot clusters with similar markers and anatomical
location were then merged together.
807
Differential gene expression analysis
To compare the gene expression program of each tissue cluster between old and young mice
as well as infected and control mice, we performed a DEG analysis using the *FindMarkers*
function of Seurat in combination with MAST. Thereby, we used the normalized data slot of the
spatial assay, requiring a minimal fold-change of 0.25, minimal percentage expressed of 0.1,
as well as an alpha level of 0.01, while again using the sample identity of each spot as latent

variable for gene regression. A gene was considered as significantly dysregulated for an organ
if it returned as significant in at least one of the spot clusters.
816

Spatially variable gene analysis
We applied SPARK-X[99,100] (v1.1.1) to each Spaceranger-filtered and preprocessed but
uncleaned sample matrix to detect the spatially variable genes in each tissue sample,
independent of our clustering and annotation approach and thus to prevent any artifacts arising
from our cleaning procedures. To this end, we used the SCT normalized counts as input to
SPARK-X and selected the mixture projection kernel. We then only consider genes as spatially
variable, if they are significant in the *mtest* slot to an alpha level of 0.01. For the comparison
to the tissue cluster DEGs we further filtered the SVGs by keeping only those with the lowest
15% of all adjusted p-values and which are then significant in more than 50% of all the samples
in each tissue over all ages.
827

Pseudobulk and spatial neighborhood expression analysis
Expression analysis on the sample-level was performed using pseudobulk aggregation of
normalized counts via the Seurat function AggregateExpression, either per experimental group
or per spot cluster. Resulting count matrices were then scaled to zero mean and unit variance
using the ScaleData function. Clustering of expression matrices was performed using
hierarchical clustering with Euclidean distance metric. To perform a spot neighborhood
expression analysis for parasite invasion hot spots in the infection cohort, we selected all spots
with more than 0.5% of ANKA mapped reads, and pulled all up to six direct neighbors in the
hexagonal spot array. We then compared the group of center spots against their neighbors
using the Seurat *FindMarkers* function with a minimum log fold-change of 0.25, minimum
percentage of expressed spots of 0.1, an alpha level of 0.05 as well as using Spot sample
identity for latent variable regression. To analyze the combined expression of genes from the
complement pathway via an activity score, we used the *AddModuleScore* function from Seurat
with a fixed seed on the data slot of the spatial assay and all other parameters kept at default.
842

Spatial neighborhood enrichment analysis
To perform spatial neighborhood analysis on our Visium 10x samples, we first converted all
Seurat objects to Squidpy[101] (v1.4.1) *AnnData* objects. This was accomplished by first
extracting gene expression matrix, spatial coordinates and required metadata from the Seurat
object, subsequently loading them into Squidpy using the *AnnData* function. To process groups
of samples together, we first manually shifted the spatial coordinates of each sample such that
there is no spatial overlap between samples in the same dimensions. Based on those common
coordinates, we first calculated the spatial connectivity matrix using the function
gr.spatial_neighbours. To validate that there is no inter-sample connectivity we created spatial
scatter plots using the *pl.spatial_scatter* function with connectivity key set to
spatial_connectivities. Then, we used the Squidpy function *gr.nhood_enrichment* with mode
zscore to calculate the neighborhood enrichment matrix for each group of samples. To
compare between two groups (e.g. young vs. old), we computed the pointwise difference
between their neighborhood enrichment matrices, providing us with a differential neighborhood
enrichment matrix per contrast.
858

Pathway enrichment analysis
To perform gene set over-representation analysis on DEGs we used the Transcriptomics
module of GeneTrail3[102], with the target species set to Mus musculus, null hypothesis set to
two-sided, and Benjamini-Hochberg FDR p-value correction at an alpha level of 0.05. We
tested against Gene Ontology (Biological process, Cellular component, Molecular function),
KEGG, Reactome, Wikipathway, and Pfam databases with all supported mouse genes as
background normalization set.
866

Analysis of small non-coding RNA sequencing data
Raw reads were cut, cleaned, filtered and mapped against small non-coding RNA annotations
using the end-to-end pipeline miRMaster2[103]. Genome-wide mature miRNA counts were

generated for each valid sample using the *Mus musculus* annotation of miRBase[104] v22.1.
Overall, three of 48 samples with unexpectedly low total read counts were removed from any
further analysis. We then performed standard QC steps such as dimension reduction, sample
and feature correlation and clustering analysis, count normalization as well as differential
expression using our established pipeline in miRMaster2. Enrichment analysis was conducted
using miEAA[105] at default parameters and with all recommended category source databases
enabled.

Spatial metabolomics data generation
A timsTOF flex MALDI-2 (Bruker Daltonics) operated in positive MALDI (Matrix-Assisted Laser
Desorption/Ionization) mode with Post Ionization laser and enabled ion mobility separation was
used for the data acquisition. The Imaging 20 μm Smart Beam laser intensity was set to 85%,
creating one burst of 200 Shots with a frequency of 1000 Hz per pixel. Trigger delay for the
Post Ionization laser was set to 5 μs and the used laser power for the post ionization laser was
calculated automatically. A scan range of 50-1300 *m/z* was chosen. The offset of the MALDI
plate was set to 50 V, deflection 1 delta to 70 V, Funnel 1 RF to 250 Vpp, Funnel 2 RF and the
multipole RF to 200 Vpp. The collision cell energy was set to 10 eV and the collision RF to
350 Vpp. The quadrupole low mass was set to 50 *m/z* with an ion energy of 5 eV. The transfer
time for the pre TOF focus was set to 75 μs with a pre pulse storage of 5 μs. A mobility range
given in $1/K_0$ was measured from 0.50 Vs cm^{-2} to 1.40 Vs cm^{-2} with a ramp time of 100 ms and
an accumulation time of 193 ms. The duty cycle was set to 100% with a ramp rate of 5.02 Hz.
Mass calibration was conducted with red phosphorus on the respective cluster formed with the
MALDI laser and ion mobility calibration dimension was calibrated linearly using 4 selected
ions from ESI Low Concentration Tuning Mix (Agilent Technologies, USA) [*m/z*, $1/k_0$:
(322.048121, 0.7363 Vs cm^{-2}), (622.028960, 0.9915 Vs cm^{-2}), (922.0098, 0.9915 Vs cm^{-2})] in
positive mode. The mobility values for the mobility calibration were taken from the CCS
compendium.

Sample preparation including the slicing is identical to the transcriptomics sample preparation,
using an embedding in 2% CMC (Carboxymethylcellulose) solution (water) and matched
samples from the other brain hemispheres. The slices were transferred immediately onto ITO
(Indium Tin Oxide) coated glass slides after cutting, dried at room temperature and put in a
desiccator for 24 h. Before matrix application the edges of every tissue were marked with a
solvent resistant pen by making four crosses in a rectangular order onto the ITO-coated slide
in 2 mm distant from the tissue border each. Each glass slides was scanned to obtain a picture
showing the position of the tissue and the marked crosses on the glass slide (EPSON
perfection V30). 15 mg/mL 2,5 - Dihydroxybenzoic acid (DHB), dissolved in a mixture of
acetonitrile and water (90:10) spiked with 0.1% TFA (trifluoroacetic acid) was sprayed onto the
tissue (HTX M5 sprayer). Fourteen passes of matrix were added with a velocity of
1200 mm/min, a track spacing of 3 mm and a flow rate of 0.125 mL/min. The nozzle height
was set to 40 mm and a temperature to 60°C while the nitrogen gas flow was set to 2 L/min
and a pressure of 10 psi.

Spatial metabolomics analysis
The automatic MALDI imaging runs were set up using fleximaging 7.6 (Bruker Daltonics) and
the timsTOF flex MALDI-2 was operated with timscontrol 6.16. Raw data files were then
processed together with mzmine[106] v4.3.0 using batch mode and the following processing
steps in order: 1) "Import MS data", 2) "Mass detection", 3) "Image builder", 4) "Ims expander",
5) "Smoothing", 6) "Isotopic peaks finder", 7) "Image co-localization", 8) "Join aligner", 9)
"Feature list rows filter", 10) "Local compound database search", 11) "Export CSV", 12) "Raw
data export". The resulting mass feature peak list was analyzed with MetaboAnalyst6.0[107] and
individual feature ion images were generated using the Bruker SCiLS lab software.
MetaboAnalyst was configured to remove features with low variance using the interquartile
range filter set to 40% and to remove features with low abundance using the mean intensity
filter set to 40%. Peak intensities were then normalized using quantile normalization, log
scaling and mean centering.

926
Statistics and reproducibility

All scripts were executed in a reproducible environment using a custom Snakemake v7.32.4 workflow with dedicated conda environments, fixing each utilized R and python package to a particular version: R v4.3.2, python v3.12, numpy v1.26.4, pandas v2.2.1, tidyverse v.2.0.0, patchwork v1.2.0, umap v0.2.10.0, pals v1.8, colorspace v2.1_0, ggrepel v.0.9.5, complexupset v1.3.3, bioconductor-complexheatmap v.2.18.0, readxl v1.4.3, openxlsx v4.2.5.2, future v1.33.0, pbapply v1.7_2, bioconductor-mast v1.28.0, seurat v4.4.0, seuratobject v4.1.4, seurat-disk v0.0.0.9021, matrix v1.6_1.1, sctransform v0.4.1, bioconductor-glmgampoi v1.14.0, hdf5r v1.3.10, python leidenalg v0.10.2, leidenalg v1.1.3, reticulate v1.34.0, python-igraph v0.11.4, umap-learn v0.5.5, devtools v2.4.5, remotes v2.4.2, yaml v2.3.8, doparallel v1.0.17, data.table v 1.15.2, ggVennDiagram v1.5.2, viridis v0.6.5, ggdendro v0.2.0, fmsb v0.7.6, Stereopy (v1.1.0), SPARK-X (v1.1.1), and Squidpy (v1.4.1). Statistical tests were performed two-tailed if not stated otherwise. Raw p-values were adjusted for multiple testing bias to an alpha level of 0.05 using either the Bonferroni (DEG analysis) or Benjamini-Hochberg FDR (pathway analysis) correction method. Trend curves were obtained through Local Polynomial Regression Fitting (loess) of $y \sim log(x)$ with a span of 0.75, two degrees of freedom and a standard error confidence interval of 0.95.

Data availability

The sequencing datasets and images generated throughout this study are available in the NCBI Sequence Read Archive (SRA) via accession-ID PRJNA1163328 (reviewer link: https://dataview.ncbi.nlm.nih.gov/object/PRJNA1163328?reviewer=43g3vvl2a3s5o0nnrksc23pu9) as well as Gene Expression Omnibus (GEO) via accession-ID GSE283283 (reviewer password: ybkzqyygdjatvad). Preprocessed Seurat R objects will also be made available upon request.

Code availability

Custom scripts used to produce the results of this study are available from the corresponding authors upon request.

Acknowledgements and funding

We thank our peers from all the contributing labs providing their continued support and feedback on the project. We are grateful for having received advanced user support and guidance from 10x Genomics and BGI on establishing their sequencing-based spatial transcriptomics platforms. This study was funded by the M.J. Fox Foundation (MJFF-021418), the Knight Initiative for Brain Resilience, the Schaller-Nikolich Foundation, the Hans-und-Ruth-Giessen Foundation, and Saarland University. Computational resources used within this study were financed through the DFG project 469073465. G.K-C. received further support by the DFG CRC TRR152 grant P22 and a DFG KR 4338/1-2 grant.

Author contributions

F.K. and Vi.W. wrote the manuscript with input from all other authors; F.K. led the data analysis with support by S.G., J.A., A.H., M.F., F.G., and S.R.; Vi.W., Va.W, and N.L. performed the spatial transcriptomics experiments; A.F., M.A., B.K., A.H., H.L., T.J., and O.H. organized the cohort setup, animal treatment, organ extraction and sample logistics; Vi.W., Va.W, A.H., P.W., M.H., T.J., G.K-C., U.B., and O.H. supported the histological analysis and cluster annotation procedures; P.D., and D.K. performed the spatial metabolomics experiments and supported the data analysis; U.B., O.H., and T.W-C. supported the design of the mouse aging cohort; H.L., R.M., and T.J. supported the design of the mouse infection cohort; T.J., G.K-C., U.B., T.W-C., and A.K. supported data interpretation and supervised the study; F.K., T.W-C. and A.K. provided the study funding. All authors read and approved upon the manuscript. U.B., O.H., T.J., and G.K-C. contributed equally and were listed in alphabetical order of surnames.

Corresponding authors
Correspondence and requests for materials should be addressed to Andreas Keller (andreas.keller@ccb.uni-saarland.de) or Fabian Kern (fabianmichael.kern@helmholtz-hips.de).

Competing interests
The authors declare no conflicts of interests.

Supplementary Information
Supplementary Information is available for this paper.

Supplementary Table 1: Sample metadata and quality control metrics for the mouse aging cohort.
Supplementary Table 2: Spot cluster marker genes for peripheral organs from the aging cohort (10x Visium).
Supplementary Table 3: Aging DEGs per cluster and peripheral organ (10x Visium).
Supplementary Table 4: Aging SVGs per peripheral organ and sample (10x Visium).
Supplementary Table 5: Spot cluster marker genes for brain sections from the aging cohort (10x Visium).
Supplementary Table 6: Aging DEGs per cluster and brain bregma (10x Visium).
Supplementary Table 7: Aging SVGs per brain bregma and sample (10x Visium).
Supplementary Table 8: Spot cluster marker genes for brain sections from the aging cohort (STOmics Stereo-seq).
Supplementary Table 9: Aging DEGs per cluster and brain bregma (STOmics Stereo-seq).
Supplementary Table 10: Computationally aligned and annotated mass feature peaks for the aging brain spatial metabolomics experiments.
Supplementary Table 11: Sample metadata and quality control metrics for the mouse infection cohort.
Supplementary Table 12: Spot cluster marker genes for all organs from the infection cohort (10x Visium).
Supplementary Table 13: Infection DEGs per cluster and organ (10x Visium).
Supplementary Table 14: Infection SVGs per organ and sample (10x Visium).
Supplementary Table 15: Annotation, normalized counts, differential expression and enrichment statistics from miRNA sequencing of the infection cohort.
Supplementary Table 16: Raw GeneTrail3 results for the over-representation analysis of brain, kidney, liver and lung DEGs shared between aging and infection clusters.

1017   **Extended Data Figures**
1018
1019   **Extended Data Figure 1: Aging peripheral organ samples quality control.**
1020
1021   **a-e**, Distribution of percentage of mitochondrial reads (top left), total number of UMIs (top
1022   right), number of detected gene features (bottom left) and spot complexity (bottom right), per
1023   cleaned high-quality sample and experimental group, for heart (**a**), kidney (**b**), liver (**c**), lung
1024   (**d**), and spleen (**e**).

Extended Data Figure 1: Aging peripheral organ samples quality control

1025 **Extended Data Figure 2: Aging brain samples quality control.**
1026
1027 **a-e**, Distribution of percentage of mitochondrial reads (top left), total number of UMIs (top
1028 right), number of detected gene features (bottom left) and spot complexity (bottom right), per
1029 cleaned high-quality sample and experimental group, for brain1 (**a**), brain2 (**b**), brain3 (**c**),
1030 brain4 (**d**), and brain5 (**e**).

Extended Data Figure 2: Aging brain samples quality control

1031　**Extended Data Figure 3: Integrated annotations and DEG analysis for the aging**
1032　**peripheral organs.**

1033
1034　**a-e**, H&E-stained image slices (left) ordered by age (top row: young, middle row: middle,
1035　bottom row: old) with annotated spot clusters overlaid for all high-quality samples from heart
1036　(**a**), kidney (**b**), liver (**c**), lung (**d**), and spleen (**e**). Pie charts (right) showing the proportion of
1037　spot clusters per age group.

1038
1039　**f-j**, Volcano plots of aging DEGs (old vs. young) colored by spot cluster for heart (**f**), kidney
1040　(**g**), liver (**h**), lung (**i**), and spleen (**j**).

Extended Data Figure 3: Integrated annotations and DEG analysis for the aging peripheral organs

1041   **Extended Data Figure 4: SVG analysis and neighborhood enrichments for aging**
1042   **peripheral organs.**
1043
1044   **a**, Bar plots showing the number of significant SVGs (y-axis) by the minimum number of
1045   replicate samples (x-axis) per age group and peripheral organ as determined with SPARK-X.
1046
1047   **b**, Histogram showing the adjusted and -log10 scaled p-value distribution by age and
1048   peripheral organ for all significant spatially variables genes determined with SPARK-X.
1049
1050   **c**, Venn diagram comparing the significant aging DEGs (old vs. young) and significant SVGs
1051   per peripheral organ.
1052
1053   **d-g**, Like in **Fig. 1j** but for kidney (**d**), liver (**e**), lung (**f**), and spleen (**g**).

Extended Data Figure 4: SVG analysis and neighborhood enrichments for aging peripheral organs

Extended Data Figure 5: Integrated annotations and DEG analysis for the aging brain slices.

a-e, As in **Extended Data Fig. 3a-e** but for the aging brain slices.

f-j, As in **Extended Data Fig. 3f-j** but for the aging brain samples.

Extended Data Figure 5: Integrated annotations and DEG analysis for the aging brain slices

1060   **Extended Data Figure 6: SVG analysis for aging brain slices.**

1061

1062   **a**, As in **Extended Data Fig. 4a** but for the aging brain samples.

1063

1064   **b**, As in **Extended Data Fig. 4b** but for the aging brain samples.

1065

1066   **c**, As in **Extended Data Fig. 4c** but for the aging brain samples.

Extended Data Figure 6: SVG analysis for aging brain slices

 Extended Data Figure 7: Quality control and annotations for STOmics aging brain
 slices.

 a, Illustration of the five different brain bregma used for STOmics Stereo-seq in accordance
 with the Visium data set. Representative H&E stains are shown for each Bregma. Since
 Stereo-seq does not support H&E stains directly from the sequenced tissue slices, an
 adjacent (directly before or after) tissue slice was prepared and stained before running the
 spatial transcriptomics experiments.

 b, From left to right and per brain bregma (top to bottom): integrated UMAP representation of
 all cleaned Stereo-seq spot clusters using the bin200 resolution, pie charts and per replicate
 spatial projections of the final annotated spot clusters. Cluster names and colors were
 assigned in accordance with the Visium data set (cf. Methods).

 c, Distribution of four main quality control features across the cleaned spots and per Stereo-
 seq brain replicate at bin200 resolution.

Extended Data Figure 7: Quality control and annotations for STOmics aging brain slices

1083 **Extended Data Figure 8: Spatial metabolomics of the aging brain.**
1084
1085 **a**, MS peak area intensities before and after normalization displayed by feature.
1086
1087 **b**, MS peak area intensities before and after normalization displayed by sample.
1088
1089 **c**, Two-dimensional principal component matrix for the first five dimensions from a PCA of
1090 peak areas after normalization. Point shape and color correspond to a sample age group.
1091
1092 **d**, Volcano plot with raw unpaired t-test p-values against log fold-changes comparing spatial
1093 metabolomics peak features for old (N=4) versus young (N=4) mouse brains. Points are
1094 colored according to fold-change.
1095
1096 **e**, Experimentally measured and computationally integrated peak intensities of N-Acetyl-L-
1097 aspartic Acid (~ 176.0576 *m/z*) in the spatial domain of twelve brain hemispheres from the
1098 aging cohort.
1099
1100 **e**, Peak intensities for N-Acetylaspartic acid across mouse brain regions and four age groups
1101 as reported by Ding et al[22].

Extended Data Figure 8: Spatial metabolomics of the aging brain

1102 **Extended Data Figure 9: Quality control for organ samples of the infection cohort.**
1103
1104 **a-e**, Like in **Extended Data Fig. 1a-e** but for all organs from the infection cohort.

Extended Data Figure 9: Quality control for organ samples of the infection cohort

1105 **Extended Data Figure 10: Integrated organ annotations and DEG analysis for the**
1106 **infection cohort.**
1107
1108 **a-e**, As in **Extended Data Fig. 3a-e** but for all the organs from the infection cohort, split by
1109 experimental group.
1110
1111 **f**, Like in **Fig. 1g**, but for all tissue-cluster DEGs from the infection cohort.
1112
1113 **g-k**, As in **Extended Data Fig. 3f-j** but for all the organs from the infection cohort, comparing
1114 day6 post-infection against healthy controls.

Extended Data Figure 10: Integrated organ annotations and DEG analysis for the infection cohort

 Extended Data Figure 11: SVG analysis and neighborhood enrichments for the
 infection cohort.

 a, As in **Extended Data Fig. 4a** but for all the infection cohort organs, split by experimental
 group.

 b, As in **Extended Data Fig. 4b** but for all the infection cohort organs, split by experimental
 group.

 c, As in **Extended Data Fig. 4c** but for all the infection cohort organs, comparing SVGs
 against day6 post-infection vs. healthy control DEGs.

 d-f, Like in **Fig. 1j** but for kidney (**d**), liver (**e**), and lung (**f**) from the infection cohort.

Extended Data Figure 11: SVG analysis and neighborhood enrichments for the infection cohort

1128 **Extended Data Figure 12: ANKA genome read detection analysis for the infection**
1129 **cohort.**
1130
1131 **a**, Heatmap showing the scaled pseudobulk expression by brain sample and experimental
1132 group of the top 50 most variable genes from the *P. berghei* (ANKA) genome.
1133
1134 **b**, Distribution of the percentage of ANKA-related read counts per spot across all annotated
1135 spot clusters of the brain and split by experimental group.
1136
1137 **c**, **d**, As in (**a**,**b**) but for olfactory bulb.
1138
1139 **e**, **f**, As in (**a**,**b**) but for kidney.
1140
1141 **g**, **h**, As in (**a**,**b**) but for liver.
1142
1143 **i**, **j**, As in (**a**,**b**) but for lung.

Extended Data Figure 12: ANKA genome read detection analysis for the infection cohort

Extended Data Figure 13: MicroRNA expression analysis for the infection cohort.

a, Small non-coding RNA sequencing samples from the infection cohort projected onto the first two principal components of a PCA using the normalized miRNA expression. Each biological sample is a dot colored by tissue of origin. Values at axis labels describe the percentage of variance explained for each component.

b, Results of the Principal Variance Component Analysis (PVCA) for the normalized miRNA expression from the infection cohort. Each bar shows the total percentage of explainable variance for a single covariate or a linear combination of such. The category 'residual' means the total variance left unexplained by all other variables.

c, Heatmap of z-scores for the 542 miRNAs robustly detected in at least one organ for the infection cohort. The order of rows (miRNAs) and columns (samples) were determined using hierarchical clustering.

d, Heatmap of sample-to-sample spearman correlation analysis. The order of rows (miRNAs) and columns (samples) were determined using hierarchical clustering.

e-i, Volcano plots for miRNA differential expression statistics of brain, bulb, kidney, liver, and lung samples, comparing infected samples against controls.

Extended Data Figure 13: Analysis of tissue-specific microRNA expression in the infection cohort

Extended Data Figure 14: Validation of C4b and Serpina3n upregulation in the aging mouse brain.

a, Combined analysis of C4b and Serpina3n by per-spot multiplication of normalized expression values across four groups of samples (young, old, control, infected). Shown are the combined expression values for Astrocyte assigned spots from the cell binning resolved brain1 Stereo-seq samples (left), and the Astrocyte marker enriched spots from the aging (middle) and malaria disease (right) mouse brain Visium samples.

b, As in (**a**) but for the Oligodendrocyte assigned spots (Stereo-seq) and Oligodendrocyte marker enriched spots (Visium).

c, Normalized protein expression intensities of C4b and Serpina3n in young (3 month), adult middle aged (15 month), and old (24 month) mouse brain cortex (left) and hippocampus (right), as originally obtained by Tsumagari *et al*[28].

Extended Data Figure 14: Validation of C4b and Serpina3n upregulation in the aging mouse brain

a   Astrocytes

b   Oligodendrocytes

c

1180 **Extended Data Figure 15: Summarized findings on complement and protease inhibitor**
1181 **activation.**
1182
Members of the Serine protease inhibitor family regulate the expression of the classical and
lectin complement pathways by interacting with individual proteases. We found a wide-spread
and strong upregulation of these genes in the aging and malaria diseased mouse within
different spatial compartments of several peripheral organs and the central nervous system
(e.g. white matter tract), a gene signature which can also be traced down to individual cell
types.

Extended Data Figure 15: Summarized findings on complement and protease inhibitor activation

References

1. Tian, Y. E. *et al.* Heterogeneous aging across multiple organ systems and prediction of chronic disease and mortality. *Nature Medicine* **29**, 1221–1231 (2023). https://doi.org/10.1038/s41591-023-02296-6

2. Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature* **583**, 596–602 (2020). https://doi.org/10.1038/s41586-020-2499-y

3. Almanzar, N. *et al.* A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature* **583**, 590–595 (2020). https://doi.org/10.1038/s41586-020-2496-1

4. Wagner, V. *et al.* Characterizing expression changes in noncoding RNAs during aging and heterochronic parabiosis across mouse tissues. *Nature Biotechnology* (2023). https://doi.org/10.1038/s41587-023-01751-6

5. Kimmel, J. C. *et al.* Murine single-cell RNA-seq reveals cell-identity- and tissue-specific trajectories of aging. *Genome Research* **29**, 2088–2103 (2019). https://doi.org/10.1101/gr.253880.119

6. Hajdarovic, K. H. *et al.* Single-cell analysis of the aging female mouse hypothalamus. *Nature Aging* **2**, 662–678 (2022). https://doi.org/10.1038/s43587-022-00246-4

7. Allen, W. E., Blosser, T. R., Sullivan, Z. A., Dulac, C. & Zhuang, X. Molecular and spatial signatures of mouse brain aging at single-cell resolution. *Cell* **186**, 194–208.e118 (2023). https://doi.org/10.1016/j.cell.2022.12.010

8. Jeffries, A. M. *et al.* Single-cell transcriptomic and genomic changes in the aging human brain. *bioRxiv*, 2023.2011.2007.566050 (2023). https://doi.org/10.1101/2023.11.07.566050

9. Hahn, O. *et al.* Atlas of the aging mouse brain reveals white matter as vulnerable foci. *Cell* **186**, 4117–4133.e4122 (2023). https://doi.org/10.1016/j.cell.2023.07.027

10. Williams, C. G., Lee, H. J., Asatsuma, T., Vento-Tormo, R. & Haque, A. An introduction to spatial transcriptomics for biomedical research. *Genome Medicine* **14**, 68 (2022). https://doi.org/10.1186/s13073-022-01075-1

11. Vandereyken, K., Sifrim, A., Thienpont, B. & Voet, T. Methods and applications for single-cell and spatial multi-omics. *Nature Reviews Genetics* **24**, 494–515 (2023). https://doi.org/10.1038/s41576-023-00580-2

12. Ogrodnik, M. *et al.* Whole-body senescent cell clearance alleviates age-related brain inflammation and cognitive impairment in mice. *Aging Cell* **20**, e13296 (2021). https://doi.org/10.1111/acel.13296

13. Yang, A. C. *et al.* Dysregulation of brain and choroid plexus cell types in severe COVID-19. *Nature* **595**, 565–571 (2021). https://doi.org/10.1038/s41586-021-03710-0

14. Dourado, M., Cavalcanti, F., Vilar, L. & Cantilino, A. Relationship between Prolactin, Chronic Kidney Disease, and Cardiovascular Risk. *International Journal of Endocrinology* **2020**, 9524839 (2020). https://doi.org/10.1155/2020/9524839

15. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays. *Cell* **185**, 1777–1792.e1721 (2022). https://doi.org/10.1016/j.cell.2022.04.003

16. Marschallinger, J. *et al.* Lipid-droplet-accumulating microglia represent a dysfunctional and proinflammatory state in the aging brain. *Nature Neuroscience* **23**, 194–208 (2020). https://doi.org/10.1038/s41593-019-0566-1

17. Shi, S. M. *et al.* Glycocalyx dysregulation impairs blood–brain barrier in ageing and disease. *Nature* **639**, 985–994 (2025). https://doi.org/10.1038/s41586-025-08589-9

18. Pietzner, M. *et al.* Comprehensive metabolic profiling of chronic low-grade inflammation among generally healthy individuals. *BMC Medicine* **15**, 210 (2017). https://doi.org/10.1186/s12916-017-0974-6

19. Vallianatou, T. *et al.* Applying Spatial Metabolomics To Investigate Age- and Drug-Induced Neurochemical Changes. *ACS Chemical Neuroscience* **15**, 2822–2829 (2024). https://doi.org/10.1021/acschemneuro.4c00199

20. Clarke, H. A. *et al.* Spatial mapping of the brain metabolome lipidome and glycome. *Nature Communications* **16**, 4373 (2025). https://doi.org/10.1038/s41467-025-59487-7

21. Kim, S. *et al.* PubChem 2025 update. *Nucleic Acids Research* **53**, D1516–D1525 (2025). https://doi.org/10.1093/nar/gkae1059

22. Ding, J. *et al.* A metabolome atlas of the aging mouse brain. *Nature Communications* **12**, 6021 (2021). https://doi.org/10.1038/s41467-021-26310-y

23. Cao, X. *et al.* Accelerated biological aging in COVID-19 patients. *Nature Communications* **13**, 2135 (2022). https://doi.org/10.1038/s41467-022-29801-8

24   Lord, J. M. *et al.* Accelerated immune ageing is associated with COVID-19 disease severity.
Immunity & Ageing **21**, 6 (2024). https://doi.org/10.1186/s12979-023-00406-z

25   Bobak, C. A. *et al.* Increased DNA methylation, cellular senescence and premature epigenetic
aging in guinea pigs and humans with tuberculosis. *Aging* **14**, 2174–2193 (2022).
https://doi.org/10.18632/aging.203936

26   Rénia, L. & Howland, S. W. Targeting the olfactory bulb during experimental cerebral malaria.
Trends in Parasitology **30**, 375–376 (2014). https://doi.org/10.1016/j.pt.2014.05.002

27   Szklarczyk, D. *et al.* The STRING database in 2023: protein–protein association networks and
functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Research*
51, D638–D646 (2022). https://doi.org/10.1093/nar/gkac1000

28   Tsumagari, K., Sato, Y., Aoyagi, H., Okano, H. & Kuromitsu, J. Proteomic characterization of
aging-driven changes in the mouse brain by co-expression network analysis. *Scientific Reports*
13, 18191 (2023). https://doi.org/10.1038/s41598-023-45570-w

29   Yao, Z. *et al.* A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse
brain. *Nature* **624**, 317–332 (2023). https://doi.org/10.1038/s41586-023-06812-z

30   Langlieb, J. *et al.* The molecular cytoarchitecture of the adult mouse brain. *Nature* **624**, 333–
342 (2023). https://doi.org/10.1038/s41586-023-06818-7

31   Shi, H. *et al.* Spatial atlas of the mouse central nervous system at molecular resolution. *Nature*
622, 552–561 (2023). https://doi.org/10.1038/s41586-023-06569-5

32   Ortiz, C. *et al.* Molecular atlas of the adult mouse brain. *Science Advances* **6**, eabb3446 (2020).
https://doi.org/10.1126/sciadv.abb3446

33   Chen, A. *et al.* Single-cell spatial transcriptome reveals cell-type organization in the macaque
cortex. *Cell* **186**, 3726–3743.e3724 (2023). https://doi.org/10.1016/j.cell.2023.06.009

34   Wang, I. H. *et al.* Spatial transcriptomic reconstruction of the mouse olfactory glomerular map
suggests principles of odor processing. *Nature Neuroscience* **25**, 484–492 (2022).
https://doi.org/10.1038/s41593-022-01030-8

35   Navarro, J. F. *et al.* Spatial Transcriptomics Reveals Genes Associated with Dysregulated
Mitochondrial Functions and Stress Signaling in Alzheimer Disease. *iScience* **23** (2020).
https://doi.org/10.1016/j.isci.2020.101556

36   Grisanti Canozo, F. J., Zuo, Z., Martin, J. F. & Samee, M. A. H. Cell-type modeling in spatial
transcriptomics data elucidates spatially variable colocalization and communication between
cell-types in mouse brain. *Cell Systems* **13**, 58–70.e55 (2022).
https://doi.org/10.1016/j.cels.2021.09.004

37   Jung, S.-H. *et al.* Spatiotemporal dynamics of macrophage heterogeneity and a potential
function of Trem2hi macrophages in infarcted hearts. *Nature Communications* **13**, 4580 (2022).
https://doi.org/10.1038/s41467-022-32284-2

38   Kanemaru, K. *et al.* Spatially resolved multiomics of human cardiac niches. *Nature* **619**, 801–
810 (2023). https://doi.org/10.1038/s41586-023-06311-1

39   Mantri, M. *et al.* Spatiotemporal transcriptomics reveals pathogenesis of viral myocarditis.
Nature Cardiovascular Research **1**, 946–960 (2022). https://doi.org/10.1038/s44161-022-
00138-1

40   Yamada, S. *et al.* Spatiotemporal transcriptome analysis reveals critical roles for mechano-
sensing genes at the border zone in remodeling after myocardial infarction. *Nature*
Cardiovascular Research **1**, 1072–1083 (2022). https://doi.org/10.1038/s44161-022-00140-7

41   Litviňuková, M. *et al.* Cells of the adult human heart. *Nature* **588**, 466–472 (2020).
https://doi.org/10.1038/s41586-020-2797-4

42   Feng, W. *et al.* Single-cell transcriptomic analysis identifies murine heart molecular features at
embryonic and neonatal stages. *Nature Communications* **13**, 7960 (2022).
https://doi.org/10.1038/s41467-022-35691-7

43   Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *Nature* **608**, 766–777
(2022). https://doi.org/10.1038/s41586-022-05060-x

44   Ko, T. *et al.* Cardiac fibroblasts regulate the development of heart failure via Htra3-TGF-β-
IGFBP7 axis. *Nature Communications* **13**, 3275 (2022). https://doi.org/10.1038/s41467-022-
30630-y

45   Walls, G. M. *et al.* Spatial Gene Expression Changes in the Mouse Heart After Base-Targeted
Irradiation. *International Journal of Radiation Oncology*Biology*Physics* **115**, 453–463 (2023).
https://doi.org/10.1016/j.ijrobp.2022.08.031

46   Farah, E. N. *et al.* Spatially organized cellular communities form the developing human heart.
Nature **627**, 854–864 (2024). https://doi.org/10.1038/s41586-024-07171-z

47   Lake, B. B. *et al.* An atlas of healthy and injured cell states and niches in the human kidney.
Nature **619**, 585–594 (2023). https://doi.org/10.1038/s41586-023-05769-3

1310	48	Abedini, A. *et al.* Spatially resolved human kidney multi-omics single cell atlas highlights the key
1311		role of the fibrotic microenvironment in kidney disease progression. *bioRxiv*,
1312		2022.2010.2024.513598 (2024). https://doi.org/10.1101/2022.10.24.513598

49 Raghubar, A. M. *et al.* Spatially Resolved Transcriptomes of Mammalian Kidneys Illustrate the
Molecular Complexity and Interactions of Functional Nephron Segments. *Frontiers in Medicine*
9 (2022). https://doi.org/10.3389/fmed.2022.873923

50 Ferreira, R. M. *et al.* Integration of spatial and single-cell transcriptomics localizes epithelial cell–
immune cross-talk in kidney injury. *JCI Insight* **6** (2021).
https://doi.org/10.1172/jci.insight.147703

51 Hansen, J. *et al.* A reference tissue atlas for the human kidney. *Science Advances* **8**, eabn4965
(2022). https://doi.org/10.1126/sciadv.abn4965

52 Hildebrandt, F. *et al.* Spatial Transcriptomics to define transcriptional patterns of zonation and
structural components in the mouse liver. *Nature Communications* **12**, 7046 (2021).
https://doi.org/10.1038/s41467-021-27354-w

53 Zhao, R. *et al.* Single-cell and spatiotemporal transcriptomic analyses reveal the effects of
microorganisms on immunity and metabolism in the mouse liver. *Computational and Structural*
Biotechnology Journal **21**, 3466–3477 (2023). https://doi.org/10.1016/j.csbj.2023.06.020

54 Ding, X. *et al.* Spatial Transcriptomics Sequencing of Mouse Liver at 2µm Resolution Using a
Novel Spatial DNA Chip. *bioRxiv*, 2024.2001.2008.574734 (2024).
https://doi.org/10.1101/2024.01.08.574734

55 Kasmani, M. Y. *et al.* A spatial sequencing atlas of age-induced changes in the lung during
influenza infection. *Nature Communications* **14**, 6597 (2023). https://doi.org/10.1038/s41467-
023-42021-y

56 Rendeiro, A. F. *et al.* The spatial landscape of lung pathology during COVID-19 progression.
Nature **593**, 564–569 (2021). https://doi.org/10.1038/s41586-021-03475-6

57 Franzén, L. *et al.* Mapping spatially resolved transcriptomes in human and mouse pulmonary
fibrosis. *Nature Genetics* **56**, 1725–1736 (2024). https://doi.org/10.1038/s41588-024-01819-2

58 Jiang, Y. *et al.* Spatial Transcriptome Uncovers the Mouse Lung Architectures and Functions.
Frontiers in Genetics **13** (2022). https://doi.org/10.3389/fgene.2022.858808

59 Madissoon, E. *et al.* A spatial multi-omics atlas of the human lung reveals a novel immune cell
survival niche. *bioRxiv*, 2021.2011.2026.470108 (2021).
https://doi.org/10.1101/2021.11.26.470108

60 Cardenas-Diaz, F. L. *et al.* Temporal and spatial staging of lung alveolar regeneration is
determined by the grainyhead transcription factor Tfcp2l1. *Cell Reports* **42**, 112451 (2023).
https://doi.org/10.1016/j.celrep.2023.112451

61 Xu, Z. *et al.* Integrative analysis of spatial transcriptome with single-cell transcriptome and
single-cell epigenome in mouse lungs after immunization. *iScience* **25**, 104900 (2022).
https://doi.org/10.1016/j.isci.2022.104900

62 Cerapio, J. P. *et al.* Integrated spatial and multimodal single-cell transcriptomics reveal patient-
dependent cell heterogeneity in splenic marginal zone lymphoma. *The Journal of Pathology*
263, 442–453 (2024). https://doi.org/10.1002/path.6296

63 Topchyan, P. *et al.* Spatial transcriptomics demonstrates the role of CD4 T cells in effector CD8
T cell differentiation during chronic viral infection. *Cell Reports* **41**, 111736 (2022).
https://doi.org/10.1016/j.celrep.2022.111736

64 Zhang, Y. *et al.* Microbiota-mediated shaping of mouse spleen structure and immune function
characterized by scRNA-seq and Stereo-seq. *Journal of Genetics and Genomics* **50**, 688–701
(2023). https://doi.org/10.1016/j.jgg.2023.04.012

65 Williams, C. G. *et al.* Plasmodium infection induces phenotypic, clonal, and spatial diversity
among differentiating CD4+ T cells. *Cell Reports* **43**, 114317 (2024).
https://doi.org/10.1016/j.celrep.2024.114317

66 Schäbitz, A. *et al.* Spatial transcriptomics landscape of lesions from non-communicable
inflammatory skin diseases. *Nature Communications* **13**, 7729 (2022).
https://doi.org/10.1038/s41467-022-35319-w

67 Lähnemann, D. *et al.* Eleven grand challenges in single-cell data science. *Genome Biology* **21**,
31 (2020). https://doi.org/10.1186/s13059-020-1926-6

68 Sziraki, A. *et al.* A global view of aging and Alzheimer's pathogenesis-associated cell population
dynamics and molecular signatures in human and mouse brains. *Nature Genetics* **55**, 2104–
2116 (2023). https://doi.org/10.1038/s41588-023-01572-y

69 Lee, H.-G., Lee, J.-H., Flausino, L. E. & Quintana, F. J. Neuroinflammation: An astrocyte
perspective. *Science Translational Medicine* **15**, eadi7828 (2023).
https://doi.org/10.1126/scitranslmed.adi7828

70 Barth, E. *et al.* Age-dependent expression changes of circadian system-related genes reveal a
potentially conserved link to aging. *Aging* **13**, 25694–25716
https://doi.org/10.18632/aging.203788

71 Fu, S. *et al.* Centenarian longevity is positively correlated with IgE levels but negatively
correlated with C3/C4 levels, abdominal obesity and metabolic syndrome. *Cellular & Molecular*
Immunology **17**, 1196–1197 (2020). https://doi.org/10.1038/s41423-020-0386-y

72 Rai, M., Curley, M., Coleman, Z. & Demontis, F. Contribution of proteases to the hallmarks of
aging and to age-related neurodegeneration. *Aging Cell* **21**, e13603 (2022).
https://doi.org/10.1111/acel.13603

73 Dunkelberger, J. R. & Song, W.-C. Complement and its role in innate and adaptive immune
responses. *Cell Research* **20**, 34–50 (2010). https://doi.org/10.1038/cr.2009.139

74 Aulak, K. S., Davis III, A. E., Donaldson, V. H. & Harrison, R. A. Chymotrypsin inhibitory activity
of normal C1-inhibitor and a P1 Arg to His mutant: Evidence for the presence of overlapping
reactive centers. *Protein Science* **2**, 727–732 (1993). https://doi.org/10.1002/pro.5560020504

75 Duranton, J., Boudier, C., Belorgey, D., Mellet, P. & Bieth, J. G. DNA Strongly Impairs the
Inhibition of Cathepsin G by α1-Antichymotrypsin and α1-Proteinase Inhibitor*. *Journal of*
Biological Chemistry **275**, 3787–3792 (2000). https://doi.org/10.1074/jbc.275.6.3787

76 Santamaria, M. *et al.* Nuclear α1-Antichymotrypsin Promotes Chromatin Condensation and
Inhibits Proliferation of Human Hepatocellular Carcinoma Cells. *Gastroenterology* **144**, 818–
828.e814 (2013). https://doi.org/10.1053/j.gastro.2012.12.029

77 Krishnarajah, S. *et al.* Single-cell profiling of immune system alterations in lymphoid, barrier and
solid tissues in aged mice. *Nature Aging* **2**, 74–89 (2022). https://doi.org/10.1038/s43587-021-
00148-x

78 Vicuña, L. *et al.* The serine protease inhibitor SerpinA3N attenuates neuropathic pain by
inhibiting T cell–derived leukocyte elastase. *Nature Medicine* **21**, 518–523 (2015).
https://doi.org/10.1038/nm.3852

79 Yang, A. C. *et al.* A human brain vascular atlas reveals diverse mediators of Alzheimer's risk.
Nature **603**, 885–892 (2022). https://doi.org/10.1038/s41586-021-04369-3

80 West, E. E. & Kemper, C. Complosome — the intracellular complement system. *Nature Reviews*
Nephrology **19**, 426–439 (2023). https://doi.org/10.1038/s41581-023-00704-1

81 Hadjilaou, A., Brandi, J., Riehn, M., Friese, M. A. & Jacobs, T. Pathogenetic mechanisms and
treatment targets in cerebral malaria. *Nature Reviews Neurology* **19**, 688–709 (2023).
https://doi.org/10.1038/s41582-023-00881-4

82 Grant, R. A. *et al.* Circuits between infected macrophages and T cells in SARS-CoV-2
pneumonia. *Nature* **590**, 635–641 (2021). https://doi.org/10.1038/s41586-020-03148-w

83 Wendisch, D. *et al.* SARS-CoV-2 infection triggers profibrotic macrophage responses and lung
fibrosis. *Cell* **184**, 6243–6261.e6227 (2021). https://doi.org/10.1016/j.cell.2021.11.033

84 Sounart, H. *et al.* Dual spatially resolved transcriptomics for human host–pathogen
colocalization studies in FFPE tissue sections. *Genome Biology* **24**, 237 (2023).
https://doi.org/10.1186/s13059-023-03080-y

85 Real, E. *et al.* A single-cell atlas of Plasmodium falciparum transmission through the mosquito.
Nature Communications **12**, 3196 (2021). https://doi.org/10.1038/s41467-021-23434-z

86 Afriat, A. *et al.* A spatiotemporally resolved single-cell atlas of the Plasmodium liver stage.
Nature **611**, 563–569 (2022). https://doi.org/10.1038/s41586-022-05406-5

87 Ruberto, A. A. *et al.* Single-cell RNA sequencing reveals developmental heterogeneity among
Plasmodium berghei sporozoites. *Scientific Reports* **11**, 4127 (2021).
https://doi.org/10.1038/s41598-021-82914-w

88 Bando, H. *et al.* CXCR4 regulates Plasmodium development in mouse and human hepatocytes.
Journal of Experimental Medicine **216**, 1733–1748 (2019).
https://doi.org/10.1084/jem.20182227

89 Hentzschel, F. *et al.* Host cell maturation modulates parasite invasion and sexual differentiation
in Plasmodium berghei. *Science Advances* **8**, eabm7348 (2022).
https://doi.org/10.1126/sciadv.abm7348

90 Howick, V. M. *et al.* The Malaria Cell Atlas: Single parasite transcriptomes across the complete
Plasmodium life cycle. *Science* **365**, eaaw2619 (2019).
https://doi.org/10.1126/science.aaw2619

91 Ngara, M. *et al.* Exploring parasite heterogeneity using single-cell RNA-seq reveals a gene
signature among sexual stage Plasmodium falciparum parasites. *Experimental Cell Research*
371, 130–138 (2018). https://doi.org/10.1016/j.yexcr.2018.08.003

92 Rood, J. E., Maartens, A., Hupalowska, A., Teichmann, S. A. & Regev, A. Impact of the Human
Cell Atlas on medicine. *Nature Medicine* **28**, 2486–2496 (2022). https://doi.org/10.1038/s41591-
022-02104-7

93 Flotho, M. *et al.* ZEBRA: a hierarchically integrated gene expression atlas of the murine and
human brain at single-cell resolution. *Nucleic Acids Research* **52**, D1089–D1096 (2024).
https://doi.org/10.1093/nar/gkad990
94 Han, L. *et al.* Cell transcriptomic atlas of the non-human primate Macaca fascicularis. *Nature*
604, 723–731 (2022). https://doi.org/10.1038/s41586-022-04587-3
95 Jacobs, T., Plate, T., Gaworski, I. & Fleischer, B. CTLA-4-dependent mechanisms prevent T cell
induced-liver pathology during the erythrocyte stage of Plasmodium berghei malaria. *European*
Journal of Immunology **34**, 972–980 (2004). https://doi.org/10.1002/eji.200324477
96 Chun, G. *et al.* SAW: an efficient and accurate data analysis workflow for Stereo-seq spatial
transcriptomics. *Gigabyte* **2024**, 0 (2024). https://doi.org/10.46471/gigabyte.111
97 Fang, S. *et al.* Stereopy: modeling comparative and spatiotemporal cellular heterogeneity via
multi-sample spatial transcriptomics. *bioRxiv*, 2023.2012.2004.569485 (2023).
https://doi.org/10.1101/2023.12.04.569485
98 Finak, G. *et al.* MAST: a flexible statistical framework for assessing transcriptional changes and
characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biology* **16**, 278
(2015). https://doi.org/10.1186/s13059-015-0844-5
99 Sun, S., Zhu, J. & Zhou, X. Statistical analysis of spatial expression patterns for spatially
resolved transcriptomic studies. *Nature Methods* **17**, 193–200 (2020).
https://doi.org/10.1038/s41592-019-0701-7
100 Zhu, J., Sun, S. & Zhou, X. SPARK-X: non-parametric modeling enables scalable and robust
detection of spatial expression patterns for large spatial transcriptomic studies. *Genome Biology*
22, 184 (2021). https://doi.org/10.1186/s13059-021-02404-0
101 Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nature Methods* **19**,
171–178 (2022). https://doi.org/10.1038/s41592-021-01358-2
102 Gerstner, N. *et al.* GeneTrail 3: advanced high-throughput enrichment analysis. *Nucleic Acids*
Research **48**, W515–W520 (2020). https://doi.org/10.1093/nar/gkaa306
103 Fehlmann, T. *et al.* miRMaster 2.0: multi-species non-coding RNA sequencing analyses at scale.
Nucleic Acids Research **49**, W397–W408 (2021). https://doi.org/10.1093/nar/gkab268
104 Kozomara, A., Birgaoanu, M. & Griffiths-Jones, S. miRBase: from microRNA sequences to
function. *Nucleic Acids Research* **47**, D155–D162 (2018). https://doi.org/10.1093/nar/gky1141
105 Aparicio-Puerta, E. *et al.* miEAA 2023: updates, new functional microRNA sets and improved
enrichment visualizations. *Nucleic Acids Research* (2023). https://doi.org/10.1093/nar/gkad392
106 Schmid, R. *et al.* Integrative analysis of multimodal mass spectrometry data in MZmine 3. *Nature*
Biotechnology **41**, 447–449 (2023). https://doi.org/10.1038/s41587-023-01690-2
107 Pang, Z. *et al.* MetaboAnalyst 6.0: towards a unified platform for metabolomics data processing,
analysis and interpretation. *Nucleic Acids Research* **52**, W398–W406 (2024).
https://doi.org/10.1093/nar/gkae253