Title: A high-resolution analysis of arrestin2 interactions responsible for CCR5 endocytosis

Authors: Ivana Petrovic[1,*], Samit Desai[1], Polina Isaikina[1,†], Layara Akemi Abiko[1], Anne Spang[1,*] and Stephan Grzesiek[1,*]

Affiliations:
[1] Biozentrum, University of Basel, CH-4056 Basel, Switzerland

† present address: Center for Life Sciences, Paul Scherrer Institut, CH-5232 Villigen, Switzerland

*Corresponding authors. Email: stephan.grzesiek@unibas.ch, anne.spang@unibas.ch, ivana.petrovic@unibas.ch

Keywords: arrestin, G protein-coupled receptor (GPCR); CCR5; chemokines, clathrin-mediated endocytosis (CME), clathrin, Adaptor protein 2 (AP2) complex, NMR, microscopy

Abstract

Clathrin-mediated endocytosis (CME) is crucial for regulating G protein-coupled receptors (GPCRs) via phosphorylation-dependent arrestin interactions. The interplay between receptor phosphorylation and arrestin coupling to the CME machinery has only been explored for a few GPCRs. Here we have investigated the CME of the chemokine receptor CCR5 induced by native and engineered CCL5 ligands. Upon CCL5 agonist stimulation, arrestin2 translocates to CCR5 at the plasma membrane forming a long-lived CCR5-arrestin2 complex that is internalized but does not reach the lysosomes. Solution NMR revealed that arrestin2 interacts weakly with clathrin through a single binding site, independent of arrestin2 activation by phosphorylated CCR5 C-terminal tail peptides. In contrast, the arrestin2-AP2 interaction is stronger and requires arrestin2 activation depending quantitatively on the CCR5 C-terminal tail phosphorylation. The *in vitro* results are corroborated by cellular assays, which establish a crucial role of the arrestin2 interaction with AP2, but not with clathrin for CCR5 internalization. These findings give insights how receptor phosphorylation regulates arrestin-mediated GPCR internalization.

Introduction

G protein-coupled receptors (GPCRs) represent a large family of transmembrane cell surface receptors that mediate the cellular response to a wide variety of signals, including hormones, neurotransmitters, and sensory stimuli. Their crucial role in many human physiological processes makes them an important drug target with more than 30% of existing small-molecule drugs binding to GPCRs.[1] To ensure precise signal transduction the receptor signaling cascade events must be tightly regulated. This is achieved by coordinating receptor interactions with various intracellular proteins, including G proteins,[2] GPCR kinases (GRKs),[3] and arrestins,[4] with the latter eventually leading to receptor internalization in a process known as endocytosis. Endocytosis is a major mechanism of GPCR signal downregulation.[5] However, some receptors rapidly recycle back to the plasma membrane,[5] which may also serve to scavenge external ligands and degrade them within the lysosomes.[6,7]

Although GPCR endocytosis can occur through several different pathways, the vast majority of GPCRs are internalized by clathrin-mediated endocytosis (CME).[8] This process relies on the clathrin protein for the formation of vesicular structures from the plasma membrane that later mature into endosomes.[9,10] While clathrin exhibits a diverse interaction spectrum,[11,12] it cannot directly interact with the membrane receptors or plasma membrane lipids and requires adaptor proteins to select its internalization cargo.[8,13–15]

The adaptor complex 2 (AP2) plays a central role in the formation of the clathrin-coated vesicles[16] and the selection of the cargo molecules.[17] It consists of two large (α and β2) and two smaller (μ2 and σ2) subunits.[18] While each subunit plays a role in CME,[19] recognition of endocytic cargo motifs has been attributed to the σ2 subunit (LL-motif) and μ2 subunit (YxxΦ-motif).[20]

For some GPCRs, such as the protease-activated receptor 1 (PAR1), the direct interaction with AP2 is sufficient for receptor internalization.[21,22] However, the majority of GPCRs require arrestins as endocytic adaptors for internalization.[23] The non-visual arrestins, arrestin2 and

arrestin3, bind to non-visual GPCRs upon intracellular phosphorylation,[4] participate in signaling

events,[24] and can mediate receptor internalization through direct interaction with clathrin[25,26]

and the β2 subunit of the AP2 complex.[27]

Once internalized, receptors are either recycled back to the plasma membrane or sent for

lysosomal degradation with the notion that the stability of the arrestin complex potentiates

intracellular receptor sequestration.[28] According to the strength and persistence of arrestin

interactions, GPCRs have been categorized into class A and class B subtypes.[28,29] To distinguish

these classes from the common IUPHAR classification of GPCRs, we refer in the following to

the arrestin classes as arr-class A and B, respectively. Arr-class A receptors (e.g. adrenergic

receptors) form transient complexes with arrestins that quickly dissociate near the plasma

membrane and the receptor traffics alone into endosomes. In contrast, arr-class B receptors (e.g.

the vasopressin receptor 2, V2R) have a higher affinity towards arrestin and form long-lived

complexes that traffic together into endosomes.[28]

The human chemokine receptor 5 (CCR5) is a GPCR, which acts as the main HIV co-

receptor[30] and plays a central role in inflammation,[31] COVID-19 infection,[32] chronic diseases,[31]

as well as cancer.[33] We have recently investigated the phosphorylation and arrestin2 interactions

of CCR5.[29] This revealed a key pXpp GPCR phosphorylation motif in its C-terminal tail, which

is specifically recognized by well-defined interactions of individual phosphates with positively

charged arrestin2 residues. The pXpp motif is also observed in many other receptors[34] known

to form a stable complex with arrestin2.[35–37] Moreover, the strength of the arrestin interaction

could be correlated to the phosphosite density and its distribution on the receptor C-terminal tail

and intracellular loops,[29,38] thus assigning the receptors with high phosphosite density (including

CCR5) to the arr-class B subtype. While our findings demonstrated how multi-site

phosphorylation controls GPCR-arrestin interactions, the interplay between arrestin activation

by receptor phosphorylation and its interactions with the clathrin/AP2 complex responsible for

internalization remains poorly understood.

Here we have characterized the chemokine-induced internalization of CCR5 in cellular

assays and the interactions of arrestin2 with clathrin and AP2 as a function of arrestin2 activation

by the phosphorylated CCR5 C-terminal tail. The results give insights how variation in receptor

phosphorylation propensities leads to the observed differences in arrestin-mediated arr-class A

and B GPCR internalization.

Results

Ligand-dependent CCR5 internalization monitored in HeLa cells

Previous reports indicated that CCR5 undergoes CME[39] and that interactions with arrestin2

and arrestin3 are essential for this process.[40] We recently established that the phosphorylated

CCR5 C-terminus forms a high-affinity complex with arrestin2.[29] To better understand the

downstream interaction with the endocytic machinery, we monitored CCR5 internalization upon

chemokine stimulation in HeLa (CCL2, ATCC nomenclature) cells (Fig. 1). HeLa cells were

transiently transfected with plasmids containing C-terminally FLAG-tagged CCR5 and YFP-

tagged arrestin2 (arrestin2-YFP) genes. This receptor construct was used to minimize CCR5

modifications and monitor the arrestin2 interaction in conditions as close to native as possible.

Before the addition of the chemokine ligands (0 min), CCR5 was observed at the plasma

membrane (magenta), while arrestin2 (green) was distributed throughout the cell (Fig. 1a-c).

Cells stimulated with the antagonist chemokine [5P12]CCL5[41] retained this CCR5 and arrestin2

distribution even 30 min after ligand addition (Fig. 1a). In contrast, when cells were stimulated

with the native agonist chemokine CCL5[42] (Fig. 1b) or the super-agonist [6P4]CCL5[41,43] (Fig.

1c), recruitment of arrestin2 from the cytoplasm to the plasma membrane could be detected

within 10 min. Prolonged ligand incubation times (30 min) lead to CCR5 internalization in

complex with arrestin2. To eliminate the possibility that arrestin2 recruitment to the membrane

was due to interactions with endogenous proteins in HeLa cells, we monitored internalization in

cells transfected only with arrestin2. In this case, the cells did not show characteristic arrestin2

recruitment to the plasma membrane or robust puncta formation, indicative of endosomes, even

in the presence of the super-agonist (Supplementary Fig. 1a). This control experiment also

revealed a few internal arrestin puncta devoid of receptor, presumably due to the arrestin2

overexpression. This phenomenon was also observed previously.[28,44]

Stronger phosphorylation and arrestin2/3 recruitment potencies have been reported for

[6P4]CCL5 as compared to wild-type CCL5.[45] In agreement with this, [6P4]CCL5 appears more

potent than CCL5 in recruiting arrestin2 to the plasma membrane and inducing CCR5-arrestin2

puncta formation after 10 min (Fig 1b,c). However, 30 min after stimulation, no significant

difference in CCR5 and arrestin2 colocalization was observed as quantified by the Mander's

coefficient (Fig. 1d). The Manders's coefficient is about 0.7 in cells stimulated by either agonist,

but close to zero for non-stimulated or antagonist-incubated cells (Fig. 1d). Apparently the two

agonist ligands induce a differing initial rate of arrestin2 recruitment by CCR5. However, once

arrestin2 is recruited, the CCR5-arrestin2 complex accumulates and remains stable for at least

30 min inside the cells.

To further corroborate the arr-class B-like behavior of CCR5, we counted the puncta positive

for CCR5 or both CCR5 and arrestin2 in cells at this time point after stimulation by [6P4]CCL5

(Fig. 1e). No significant difference was observed between the two values. This indicates that the

number of puncta containing only CCR5, but not arrestin2, must be very small and that the

majority of CCR5 puncta also contains arrestin2, consistent with stable complex formation.

Furthermore, the CCR5 and arrestin2 colocalization occurs not only close to the plasma

membrane but also close to the nucleus (Fig. 1f), indicative of the arr-class B-like behavior of

CCR5. For arr-class A receptors, it would be expected that arrestin2 is excluded from receptor-

containing structures, which traffic into the perinuclear region.[28]

As both recycling[7,46] and degradation[7] has been observed during CCR5 internalization, the

cellular fate of the CCR5-arrestin2 complex was probed using the lysosomal-associated

membrane protein 1 (LAMP1) as a marker of the lysosomal membranes (Fig. 1g). No

recruitment of the complex to the lysosomes (LAMP1, yellow) was detected in the presence of

[6P4]CCL5 even 30 min after ligand stimulation. Similar results were obtained for both CCL5

and [6P4]CCL5 using the LysoTracker[TM] dye that accumulates in acidic environments, such as

the lysosome (Supplementary Fig. 1b,c). These results indicate that the CCR5 internalization

pathway, although mediated by arrestin, does not lead to rapid receptor degradation in

lysosomes.

Localizing and quantifying the arrestin2-clathrin interaction by NMR spectroscopy

As CCR5 relies on arrestin2 for internalization[40] and utilizes clathrin-mediated pathways,[39]

we investigated the consequences of the CCR5-arrestin2 coupling on the interaction with

clathrin and AP2 proteins. Arrestin2 consists of two consecutive (N- and C-terminal) β-

sandwich domains (Fig. 2a), followed by the disordered clathrin-binding loop (CBL, residues

353–386), strand β20, and a disordered C-terminal tail after residue 393. The CBL contains a

clathrin-binding motif ([L/I]ΦXΦ[D/E] where X denotes any amino acid and Φ is a bulky

hydrophobic residue[11,19] (Supplementary Fig. 2a). Strand β20 forms a characteristic β-sheet with

the N-terminal strand β1 in inactive arrestin. Upon activation, the phosphorylated receptor tail

replaces β20 in the β-sheet and releases it from the arrestin core.

A structure of a complex between arrestin2 and the clathrin heavy chain N-terminal domain,

(residues 1-363, CLTC NTD) has been solved by X-ray crystallography (PDB:3GD1).[47] The

structure (Supplementary Fig. 2b-d) suggests a 2:1 binding model between arrestin2 and the

CLTC NTD. The first interaction occurs between the clathrin-binding motif of the arrestin2

CBL and the edge of the first two β-sheet blades of the CLTC NTD. This CLTC region is also

known to interact with many other proteins.[11] The second interaction is found between the splice

loop of the arrestin2 C-terminal β-sandwich and the 4th and 5th blade of the CLTC NTD. The

3GD1 structure had been solved in the absence of an arrestin2-activating phosphopeptide.

To elucidate the clathrin-arrestin interactions under solution conditions and the influence of

arrestin activation we utilized NMR spectroscopy. Fig. 2b depicts part of a ^{1}H-^{15}N TROSY

spectrum (full spectrum in Supplementary Fig. 3a) of the truncated ^{15}N-labeled arrestin2

construct arrestin2$^{1-393}$ (residues 1-393), which comprises the C-terminal strand β20, but lacks

the disordered C-terminal tail. Due to intrinsic microsecond dynamics, the assignment of the

^{1}H-^{15}N resonances by triple resonance methods was so far possible only for 16 residues (residues

367-381, 385-386) within the mobile CBL. Whereas this region of arrestin is typically not

visible in either crystal or cryo-EM structures due to its high flexibility, it provides excellent

probes for solution NMR studies. The addition of an equimolar amount of CLTC NTD to

arrestin2$^{1-393}$ induced significant line broadening between residues T374 and T381 with I377

being the most strongly affected residue (Fig. 2c, Supplementary Fig. 3b). This region almost

exactly matches and fully comprises the canonical arrestin2 clathrin-binding motif ^{376}LIELD^{380}.

Fig. 2a shows these interacting residues mapped on an arrestin2 AlphaFold2 model.[48]

As ^{1}H-^{13}C methyl resonances have more favorable relaxation properties than backbone

^{1}H-^{15}N resonances,[49] we used Ile-δ1-^{13}CH$_3$-labeled arrestin2 to quantitatively monitor the

formation of the arrestin2-clathrin complex (arrestin2 assignments transferred from BMRB,

ID:51131[50]). HMQC spectra of the 15 Ile δ1-^{13}CH$_3$ methyl groups of arrestin2$^{1-393}$ (Fig. 2d)

revealed changes in peak intensity (Fig. 2e, top) and chemical shift perturbations (Fig. 2e,

bottom) only for I377 at an equimolar addition of unlabeled CLTC NTD to arrestin2$^{1-393}$. An

apparent dissociation constant $K_D = 104 \pm 9$ μM was determined from titrating CLTC NTD and

fitting the arrestin2 I377 ^{13}C chemical shifts with a 1:1 binding isotherm (Fig. 2f). Although the

formed complex (total molecular weight 85 kDa) has low affinity, such interactions are common

in endocytosis, which involves many multi-site interactions within a large network of proteins.[19]

183 The arrestin2-clathrin interaction is independent of arrestin2 activation

Due to significant line broadening of the arrestin2 resonances upon phosphopeptide addition,

the influence of arrestin activation on the clathrin interaction could not be detected on either

backbone or methyl resonances. However, the CLTC NTD, although similar in size to

arrestin2$^{1-393}$ (41 kDa vs. 44 kDa), provided considerably better ^{1}H-^{15}N TROSY spectra due to

the absence of significant micro- to millisecond dynamics. This allowed to observe the effect of

arrestin2 activation on its interaction with clathrin by titrating arrestin2[1-393] to ^{15}N-labeled CLTC

NTD. Fig. 3a (left) shows the intensity changes (full spectra in Supplementary Fig. 4a) of the

CLTC NTD ^{1}H-^{15}N TROSY resonances (assignments transferred from BMRB, ID:25403[51])

upon addition of a one-molar equivalent of arrestin2[1-393]. A significant intensity reduction due

to line broadening is detected for CLTC residues 39-40, 48-50, 62-72, 83-90, 101-106, and 108.

These residues form a clearly defined binding region at the edges of blade 1 and blade 2 of the

CLTC NTD (Fig. 3a, right), which corresponds to the first interaction site observed in the 3GD1

crystal structure. However, no significant signal attenuation was observed for residues of the

second crystal interaction site between blade 4 and blade 5 (in proximity of residues 190 and

230), indicating that there is only one arrestin2 binding site in CLTC NTD. The second binding

site in the crystal structure may thus result from arrestin crystal packing.

Next, we tested whether the arrestin2-CLTC NTD interaction is enhanced in the presence of

the fully phosphorylated CCR5 phosphopeptide (CCR5pp6, where 6 indicates the number of

phosphosites, Supplementary Table 1), which displaces arrestin2 strand β20 by forming an

antiparallel β-sheet with strand β1, thereby activating arrestin2.[29] However, no additional signal

attenuation of the CLTC NTD ^{1}H-^{15}N resonances or extension to further CLTC NTD regions

was observed (Fig. 3b, Supplementary Fig. 4b), when arrestin2[1-393] was added in the presence

of a 5-molar equivalent of CCR5pp6 (corresponding to 80% of arrestin activation). Thus,

activation of arrestin2 has apparently no effect on the CLTC NTD interaction. These results

suggest that even for receptors with low phosphosite density, such as adrenergic and dopamine

receptors[29,38]), the arrestin2-clathrin interaction may still take place allowing efficient receptor

internalization.

Previous studies have suggested that the last residues of the arrestin2 C-terminus have an

inhibitory effect on clathrin binding.[52] We therefore tested the interaction of ^{15}N-labeled CLTC

NTD with full-length arrestin2 (arrestin2[1-418]). However, the obtained residue intensity

attenuation was indistinguishable from the one observed for the arrestin2[1-393] construct (Fig. 3c,

Supplementary Fig. 4c). Therefore, the residues of the arrestin2 C-terminal tail do not interfere

with clathrin binding. In contrast, the interaction of clathrin with arrestin2 (Fig. 3d,

Supplementary Fig. 4d) was completely abolished when the arrestin2 clathrin-binding loop was

replaced by the loop of visual arrestin (arrestin2$^{\Delta CBL}$, construct details in Methods). The latter

loop lacks the clathrin-binding motif making visual arrestin incapable of interacting with

clathrin.[25]

Taken together, the solution NMR experiments reveal that arrestin2 interacts with the CLTC

NTD through a single binding site, independently of arrestin2 activation and the conformation

of the arrestin2 core.

Arrestin activation is essential for AP2 interaction

An equally important element of the endocytic pathway is the arrestin interaction with AP2.

The β2 subunit of the AP2 complex (AP2β2) interacts with the arrestin2/3 C-terminus,[27,53,54]

and a crystal structure of the complex between the AP2β2 C-terminal domain and a C-terminal

arrestin2 peptide (residues 383-402) comprising strand β20 has been solved (PDB: 2IV8).[17]

Interestingly, the C-terminal arrestin2 peptide forms a helical structure in the complex with AP2

(Fig. 4a) as opposed to its β-sheet coordination with strand β1 in inactive arrestin2. No

experimental data exist on the formation of the AP2-arrestin2 complex in the context of full-

length arrestin2 and its activation by phosphorylated receptor tails.

To test how different phosphorylation levels influence arrestin2 recruitment by the C-

terminal domain of AP2β2 (AP2β2$^{701\text{-}937}$, residues 701-937), we performed a SEC binding

assay. In the absence of phosphopeptide an equimolar mixture of 10 μM arrestin2$^{1\text{-}418}$ and 10

μM AP2β2 is separated into two peaks (Fig. 4b, orange), which elute at the positions

corresponding to the individual apo proteins (Fig. 4b, blue, green). Upon addition of 100 μM

CCR5pp6 (Fig. 4c, purple), an additional SEC peak appears at an elution volume of 2.6 ml,

which corresponds to the formed arrestin2-AP2β2 complex. Monotonously decreasing complex

yields were observed when reducing the amounts of CCR5pp6 added to the mixture (Fig. 4d),

and no complex formation occurred in the presence of 100 μM non-phosphorylated CCR5

phosphopeptide (CCR5pp0) (Fig. 4e). The dissociation constant of the CCR5pp6-arrestin2

complex was determined previously as 45 μM (Supplementary Table 1).[29] Not surprisingly, the

formation of the arrestin2-AP2β2 complex depends quantitatively on the affinity of the

phosphopeptide-arrestin2 complex. Correspondingly, the addition of the less phosphorylated

CCR5pp3 and CCR5pp4 with lower affinities towards arrestin2 (both $K_D \sim$ 200 μM,

Supplementary Table 1),[29] resulted in a considerable, similar reduction of the formed arrestin2-

AP2β2 complex (Fig. 4f).

Using the extinction coefficients of all involved polypeptides, the apparent affinities

between AP2β2 and arrestin2 could be determined from the SEC profiles (Fig. 4g, see Methods

for details). In the presence of 100 μM CCR5pp6, this apparent K_D amounts to 9 μM. An

identical affinity value for the affinity between AP2β2 and an isolated arrestin β20 peptide

(residues 383-402, D402C) has been observed by EPR spectroscopy.[55] In the presence (100 μM)

of the less phosphorylated CCR5 peptides, the apparent AP2β2-arrestin2 affinities are reduced

by ~4–5 times (CCR5pp4: 44 μM, CCR5pp3: 40 μM), in good agreement with their ~4–5 times

reduced affinity for arrestin2.

It is interesting to note that CCR5pp3 and CCR5pp4 have similar affinities towards arrestin2

despite their different phosphorylation levels (3-fold vs. 4-fold phosphorylated), but only

CCR5pp3 contains the pXpp barcode which increases the arrestin2 activation, as assayed by the

binding of the Fab30 antibody.[29] Apparently, this barcode does not play a significant role in the

formation of arrestin2-AP2β2 complex as the complex formation appears solely governed by

the affinity of the phosphopeptide towards arrestin2.

To observe the atomic details of arrestin2 activation on its interaction with AP2β2, we used

[15]N-labeled AP2β2[701-937] (assignment transferred from BMRB ID:52320[56]). Upon addition of

arrestin2$^{1-418}$, no changes in the AP2β2 ^{1}H-^{15}N spectra are detected (Fig. 4h, Supplementary Fig.

5a), apparently due to the unavailability of arrestin2 β20 strand and C-terminal tail. However,

when arrestin2$^{1-418}$ was added in the presence of a 5-molar equivalent of CCR5pp6

(corresponding to 80% of arrestin activation), a significant intensity reduction due to line

broadening is detected for many AP2β2 ^{1}H-^{15}N resonances (Fig. 4i, left, Supplementary Fig.

5b). The strongest attenuated resonances clearly define a binding region within the C-terminal

domain of AP2β2 (Fig. 4i, right), which is an extension of the binding surface observed in the

crystal structure (PDB: 2IV8) of AP2β2 in complex with a 21-residue arrestin2 C-terminal

peptide comprising strand β20.[17] In agreement with the AP2β2 NMR observations, no

interaction was observed in the arrestin2 methyl and backbone NMR spectra upon addition of

AP2β2 in the absence of phosphopeptide (Supplementary Fig. 5c,d). However, the significant

line broadening of the arrestin2 resonances upon phosphopeptide addition (Supplementary Fig.

5e, f) precluded a meaningful assessment of the effect of the AP2β2 addition on arrestin2 in the

presence of phosphopeptide.

We then tested whether the presence of CLTC NTD influences complex formation between

arrestin2 and AP2β2, and vice versa. Arrestin2$^{1-418}$ does not form a complex in the SEC assay

with CLTC NTD (Supplementary Fig. 6a), which agrees with the weak affinity observed in the

NMR experiments (Fig. 2f). Furthermore, a SEC analysis of a mixture of arrestin2$^{1-418}$, CLTC

NTD, AP2β2$^{701-937}$, and CCR5pp6 (Supplementary Fig. 6b) indicates formation of a complex at

2.6 ml elution volume. An SDS analysis of this complex fraction showed the presence of

arrestin2$^{1-418}$ and AP2β2$^{701-937}$, but not of CLTC NTD (Supplementary Fig. 6c). This indicates

that CLTC NTD does not interact strongly with the arrestin2-AP2-CCR5 tail complex. As such

clathrin may not engage with the initial receptor internalization complex, but may be recruited

at later stages of the CME.

To test the stability of the formed arrestin2-AP2 interaction and its quaternary coordination, we performed a trypsin proteolysis assay (Supplementary Fig. 7, uncropped gels in Supplementary Fig. 8). Trypsin recognizes arrestin2 residue R393,[57] which is buried in the arrestin2 apo state, but is involved in the interaction with AP2β2.[27] Without phosphopeptide, the arrestin2$^{1-418}$ digestion midpoint occurs at 20-30 min due to the inaccessibility of the cleavage site (Supplementary Fig. 7a-c). In the presence of a 5-molar equivalent of CCR5pp6 (Supplementary Fig. 7a), the top band corresponding to full-length arrestin2 has almost completely disappeared (relative abundance 19%) already 5 min after incubation as a consequence of the β20 strand release and the exposure of R393 to trypsin. However, the presence of AP2β2$^{701-937}$ protects arrestin$^{1-418}$ from proteolysis, as evident from significantly higher abundance at 5 min and a shift of the proteolysis midpoints to 10, 20, and 30 min at 1-, 2-, and 5-molar equivalents of AP2β2$^{701-937}$ respectively.

Analogous digestion experiments using CCR5pp3 (Supplementary Fig. 7b) or CCR5pp4 (Supplementary Fig. 7c) phosphopeptides revealed that they did not enhance the rate of arrestin2 proteolysis by trypsin at a 3-molar equivalent, as expected from their weaker affinity towards arrestin2. In agreement with this, however, upon addition of 25-molar equivalents of either peptide the rate of arrestin2 proteolysis was significantly increased, as full-length arrestin2 almost completely disappeared already 5 min after incubation. Thus as for CCR5pp6, the presence of AP2β2$^{701-937}$ also protects arrestin$^{1-418}$ activated by CCR5pp3 or CCR5pp4 from proteolysis.

In summary, these results indicate that arrestin2 activation by the binding of receptor phosphopeptides and the subsequent release of its strand β20 is required for the interaction with AP2. This interaction is modulated by the phosphorylation level of the receptor C-terminal peptide and its relative abundance. The formed AP2 complex with arrestin is stable and more resistant to proteolysis.

Arrestin interaction with AP2 is essential for CCR5 internalization

The interactions of arrestin with clathrin and AP2 have mostly been studied for the internalization of the β2-adrenergic (β2AR)[23,44,47,54,58] and V2R[44,58] receptors. Here, we investigated how both interactions affect the internalization of CCR5 in a HeLa KO cell line (HeLa[-arr2/3]) devoid of arrestin2 and arrestin3 as well as in HeLa (CCL2) cells (Fig. 5, Supplementary Fig. 9, respectively).

To characterize the arrestin2-clathrin interaction, we generated a YFP-tagged arrestin2 construct with the clathrin-binding motif (arrestin2[ΔLIELD]-YFP) removed. No significant difference in CCR5 internalization was detected in either HeLa cell line between wild-type arrestin2-YFP and arrestin2[ΔLIELD]-YFP, when the cells were stimulated with [6P4]CCL5 (HeLa[-arr2/3]: Fig. 5a-c, HeLa: Supplementary Fig. 9a). Apparently, the arrestin2-clathrin interaction does not play a significant role in the internalization of the arr-class B CCR5 receptor. This is in contrast to findings for the arr-class A receptor β2AR where removal of the LIELD motif impairs receptor internalization.[54]

To test the influence of the arrestin2-AP2 interaction, we introduced mutations within the arrestin2 adaptin-binding motif, which is conserved across all arrestins (Supplementary Fig. 2a). The mutation R395A within this motif has been found to inhibit β2AR internalization by arrestin3.[27] Introducing the equivalent mutation into a fluorescent arrestin2 construct (arrestin2[R395A]-YFP), did not lead to significant changes in CCR5 internalization relative to wild-type arrestin2-YFP (HeLa[-arr2/3]: Fig. 5a-c, HeLa: Supplementary Fig. 9b).

The structure of AP2β2 in complex with the C-terminal arrestin2 peptide[17] shows arrestin2 residues D385, F388, F391, and R395 interacting via salt bridges and base stacking with AP2β2 (Fig. 4a). Apparently, these interactions lead to a rather stable complex of full-length arrestin2 with AP2β2[701-937] in solution as evidenced by our SEC binding assay, NMR and the protection of activated arrestin2 against proteolysis (Fig. 4, Supplementary Fig. 7). We hypothesized that due to this extended arrestin2-AP2β2 interaction interface, the mutation of a single interacting

residue may not suffice to inhibit the arrestin2-AP2 interaction leading to CCR5 internalization.

Hence, we generated a further arrestin2 mutant (arrestin2$^{\Delta AP2}$-YFP) with all four direct

interacting residues replaced (D385A, F388A, F391A, R395A). Similar to wild-type arrestin2-

YFP, arrestin2$^{\Delta AP2}$-YFP is recruited to CCR5 at the plasma membrane (HeLa: Supplementary

Fig. 9c). However, the subsequent CCR5 internalization is potently inhibited (HeLa$^{-arr2/3}$: Fig.

5a-c, HeLa: Supplementary Fig. 9d) as indicated by a reduction of the Mander's CCR5/arrestin2

colocalization coefficient (from ~0.8 to ~0.2) the number of puncta positive for CCR5 and

arrestin2 (from ~20–13 to ~8–6 per 100 μm^2, similar values obtained for CCR5 puncta). In

agreement with this finding, the arrestin2$^{\Delta AP2}$ mutant is unable to form a complex with AP2 in

the presence of CCR5pp6 as assayed by SEC (Supplementary Fig. 9e). Taken together, these

results show that the stable arrestin-AP2 interaction is crucial for CCR5 internalization, whereas

the arrestin-clathrin interaction is not critical for sequestration from the plasma membrane.

Discussion

In the present study, we have elucidated the role of CCR5 phosphorylation in arrestin-

clathrin and arrestin-AP2 interactions which are essential for CCR5 sequestration into cells.

Cellular experiments demonstrated chemokine agonist-induced recruitment of arrestin2 towards

CCR5 at the plasma membrane, which is followed by the sequestration of the CCR5-arrestin2

complex into the cytoplasm including the perinuclear region where it remains stable for an

extended period. This provides experimental evidence that CCR5 belongs to the arr-class B of

GPCRs as surmised previously from the large number of phosphorylation sites in its C-terminus

and the presence of the pXpp motif.[29] The sequestered receptor remained stable inside the cells

for at least 30 min, and was not rapidly degraded in lysosomes, even in the presence of

[6P4]CCL5. A similar CCR5 lysosome-evading internalization pattern has been observed for

the potent anti-HIV chemokine analog PSC-RANTES.[59] As a recent CCR5 trafficking study

demonstrated that CCR5 scavenges both CCL5 and [6P4]CCL5 chemokines in an arrestin-

dependent manner,[7] it is possible that the long retention of the CCR5-arrestin complex and the

ability of the receptor to undergo recycling[7,46] is utilized to remove the extracellular ligands

from the cell surface. This may help in the regulation of chemokine ligand concentrations.[60]

NMR and chromatographic data on arrestin2 interactions with clathrin and AP2 proteins in

the presence of arrestin2-activating CCR5 C-terminal phosphopeptides revealed the molecular

details underlying CCR5 sequestration. Whereas the analysis of arrestin2 ^{1}H-^{15}N resonances

was difficult due to significant line broadening from intrinsic microsecond dynamics,[29] the

better quality of the ^{1}H-^{15}N spectra of CLTC NTD allowed a detailed characterization of the

arrestin2-clathrin interaction from the clathrin side. Arrestin2 binds to clathrin with about 100

μM affinity. This interaction is independent of arrestin2 activation by phosphopeptides, which

may enable the internalization of receptors with low phosphorylation levels. The interaction is

mediated through a single binding site in a 1:1 stoichiometry involving the arrestin2 clathrin-

binding loop and the edges of blade 1 and blade 2 of the CLTC NTD. This binding site agrees

with one of the binding sites observed in the crystal structure of the arrestin2-CLTC NTD

complex (PDB: 3GD1),[47] but disagrees with the second crystal interaction site involving the

arrestin2 splice loop. Presumably, the latter is a result of crystal packing. Our findings are

supported by a recent study,[61] which suggests that the arrestin C-domain edge comprising the

splice loop anchors arrestin to the membrane. The membrane anchoring seems mutually

exclusive with clathrin binding.[61]

In contrast to the arrestin2-clathrin interaction, the arrestin2-AP2 interaction requires

arrestin2 activation by GPCR C-terminal tail phosphopeptides and depends quantitively on their

phosphorylation levels. The formed arrestin2-phosphopeptide-AP2 complexes are stable and

more resistant to proteolysis than binary arrestin2-phosphopeptide complexes. The dissociation

constant of the arrestin2-AP2 complex is about 10 μM in the presence of the fully

phosphorylated CCR5pp6, which is a 10-fold higher affinity than that of the arrestin2-clathrin

complex. This may indicate a preference towards AP2 complex formation for highly

phosphorylated receptors. However, as the phosphorylation decreases the arrestin2-AP2

dissociation constant weakens to about 40 μM, making it more comparable to the arrestin2-

clathrin interaction.

Based on the present CCR5 data, we envision the following CME mechanism, which may

also apply to other arr-class B receptors (Fig. 6, left). The high receptor phosphorylation leads

to a strong recruitment of arrestin even in the absence of interactions with negatively charged

PIP_2 in the plasma membrane.[62] The consequent tight binding of the phosphorylated receptor

tail to arrestin strand β1 stably activates arrestin thereby releasing its strand β20 and C-terminal

tail. The freed β20 strand can then interact with the β2 subunit of AP2, which becomes released

from the AP2 core upon AP2 activation by PIP_2 molecules.[16] Apparently, the interaction of

arrestin2 strand β20 with the AP2 β2 subunit is rather strong, since it could only be inhibited in

cellular assays by a fourfold, but not by a single mutation in the arrestin2 C-terminal region. In

contrast, removal of the clathrin-binding LIELD motif from arrestin2 did not impair receptor

internalization, which is in agreement with the determined relatively low affinity of the

arrestin2-clathrin interaction. This suggests that the clathrin interaction is less important for

receptor internalization than the AP2 interaction.

The AP2 activation by PIP_2 also exposes the AP2 clathrin-binding LLNLD motif,

responsible for clathrin recruitment and formation of the clathrin-coated pits.[16] In the stable

ternary complex of a highly phosphorylated arr-class B receptor with arrestin and AP2, the

exposed AP2 LLNLD clathrin-binding motif may then subsequently couple to the clathrin

network thereby inducing rapid sequestration from the plasma membrane into endosomes. The

direct clathrin interaction of arrestin via its LIELD motif appears less important in this respect.

Although the generalization of this mechanism from CCR5 to other arr-class B receptors has

to be explored further, it is indirectly corroborated in the visual rhodopsin-arrestin1 system. The

arr-class B receptor rhodopsin[29] also undergoes CME[55] with arrestin1 harboring the conserved

AP2 binding motif, but missing the clathrin-binding motif (Supplementary Fig. 2a). As arr-class

B receptors are almost all peptide-binding GPCRs,[28,29] we speculate that the strong arrestin

interaction may also be utilized to remove the extracellular ligands from the plasma membrane,

independent of receptor degradation.

In comparison to arr-class B receptors, arr-class A receptors have a lower phosphosite

density in their C-terminal tails. Previous studies on the arr-class A β2AR[27,47,54] revealed that

both clathrin and AP2 interaction are crucial for internalization. It is expected that the lower

phosphosite density of arr-class A receptors results in a less stable interaction with arrestin (Fig.

6, right). Therefore additional interactions with PIP_2 lipids are required to anchor arrestin to the

membrane and play a significant role in receptor endocytosis.[62] As a consequence of the low

phosphorylation of the receptor and the weakened interaction with arrestin strand β1, the

equilibrium between inactive (strand β20 forms β-sheet with β1) and active (β20 released)

arrestin is shifted towards the inactive conformation, even when arrestin is bound to the receptor

via core contacts. Due to the low availability of strand β20, the arrestin interaction with AP2 is

considerably weakened and comparable in strength to the clathrin interaction. This is consistent

with the finding that AP2 and clathrin participate with similar strength in arr-class A receptor

internalization.[27,47,54] However, as both interactions are weak, the sequestration is reduced. In

addition, the weaker receptor phosphorylation leads to fast dissociation of arrestin from the

receptor at or near the plasma membrane and in consequence the receptor traffics without

arrestin into endosomes.[28,38]

A recent study[63] using single-molecule microscopy and molecular dynamics simulations

suggests that arrestin3 preassociates with the plasma membrane and that, surprisingly, the

transient interaction with an active β2-adrenergic receptor (arr-class A) activates arrestin3,

which then diffuses to CCPs (clathrin-coated pits) independently of the activating receptor. It is

conceivable, that also arrestin2 first gets activated by the phosphorylated CCR5 and then enters

the CCPs without receptor, having its β20 strand released and then interact with AP2 present in

the CCPs. However, our microscopic data rather indicate that CCR5 and arrestin2 colocalize

both at the membrane as well as in the cytosol at a later stage of the CME (Fig. 1b,c). It would

be interesting to obtain single-molecule microscopic data on the CCR5-arrestin2 system to allow

a stringent comparison with our high-resolution NMR data of the receptor tail, clathrin and AP2

interactions.

In conclusion, this biophysical and cellular analysis of the interactions between the CCR5,

arrestin2, and clathrin/AP2 endocytic machinery elucidates the interplay of forces driving GPCR

internalization. The strength of the receptor phosphorylation modulates the interaction with

arrestin2, the release of its strand β20, and subsequently its interaction with the β2 subunit of

AP2. In comparison, the arrestin2-clathrin interaction is weaker. The interaction between the

receptor phosphosites and arrestin appears as a distinctive factor between arr-class B and arr-

class A receptor sequestration, since phosphorylation of the latter is considerably weaker. These

findings expand on the previously suggested mechanism for the reduced sequestration of the

arr-class A receptor β2AR by the Kim & Benovic[54] and Laporte et al[27] where the weakened

AP2-arrestin and the phosphorylation-independent clathrin-arrestin interactions appear of

similar importance. The phosphorylation dependence of the AP2-arrestin interaction may also

play a role in modulating the strength of downstream arrestin-mediated signaling from

endosomes.[64,65] As such, the detailed understanding of these molecular interactions may provide

a framework for the development of novel therapeutics interfering with GPCR sequestration.

Methods

Constructs

The genes encoding full-length (arrestin2$^{1-418}$) and truncated (arrestin2$^{1-393}$) human arrestin2 constructs have been described before.[29] The arrestin2$^{\Delta CBL}$ construct in pET-28a (+) vector was obtained from the GenScript by replacing the loop residues of arrestin2 K357 to D385 with visual arrestin residues E365 to N376. The construct also contained an N-terminal hexahistidine tag for purification followed by a TEV cleavage site (ENLYFQG), resulting in the following total sequence (arrestin2$^{\Delta CBL}$ in bold, visual arrestin loop underlined):

```
MGSSHHHHHHSSGENLYFQGMGDKGTRVFKKASPNGKLTVYLGKRDFVDHIDLVDPVDGVVLVDPEYLKERRVYVT
LTCAFRYGREDLDVLGLTFRKDLFVANVQSFPPAPEDKKPLTRLQERLIKKLGEHAYPFTFEIPPNLPCSVTLQPG
PEDTGKACGVDYEVKAFLAENLEEKIHKRNSVRLVIRKVQYAPERPGPQPTAETTRQFLMSDKPLHLEASLDKEIY
YHGEPISVNVHVTNNTNKTVKKIKISVRQYADIVLFNTAQYKVPVAMEEADDTVAPSSTFSKVYTLTPFLANNREK
RGLALDGKLKHEDTNLASSTLLREGANREILGIIVSYKVKVKLVVSRGGLLGDLASSDVAVELPFTLMHPKPEDPA
KESYQDANIVFEDFARQRLKGMKDDKEEEEDGTGSPQLNNR
```

CLTC NTD (pETM33_CLTC_Clathrin propeller, Addgene plasmid #178471)[66] and AP2β2$^{701-937}$ (pETM33_AP2B1_adaptin, Addgene plasmid #178467)[66] were a gift from Ylva Ivarsson. Arrestin2-YFP was a gift from Robert Lefkowitz (Addgene plasmid # 36916).[67]

The arrestin2$^{\Delta LIELD}$-YFP construct was obtained from the arrestin2-YFP template by standard PCR reaction using KAPA HiFi polymerase (Forward primer: 5'-GAG ACT CCA GTA GAC ACC AAT ACC AAT GAT GAC GAC A-3'; Reverse primer: 5'-TGT CGT CAT CAT TGG TAT TGG TGT CTA CTG GAG TCT C-3').

Arrestin2^{R395A}-YFP was generated in the same manner using forward primer: 5'-GGA CTT TGC TCG TCA GGC GCT GAA AGG CAT GAA G-3' and reverse primer: 5'-CTT CAT GCC TTT CAG CGC CTG ACG AGC AAA GTC C-3'.

The same protocol was applied for generating the arrestin2$^{\Delta AP2}$-YFP clone with the forward primer of the sequence: 5'-GAC ACC AAT GAT GAC GCC ATT GTG GCT GAG GAC GCT GCT CGT CAG GCG CTG AAA GGC ATG-3' and reverse primer 5'-CAT GCC TTT CAG CGC CTG ACG AGC AGC GTC CTC AGC CAC AAT GGC GTC ATC ATT GGT GTC-3'.

The CCR5 construct in pcDNA3.1 used in cellular assays was obtained from GenScript. The

complete construct contained the full-length human CCR5 sequence (bold) followed by a GS-

linker and a C-terminal FLAG-tag (underlined):

MDYQVSSPIYDINYYTSEPCQKINVKQIAARLLPPLYSLVFIFGFVGNMLVILILINCKRLKSMTDIYLLNLAISD
LFFLLTVPFWAHYAAAQWDFGNTMCQLLTGLYFIGFFSGIFFIILLTIDRYLAVVHAVFALKARTVTFGVVTSVIT
WVVAVFASLPGIIFTRSQKEGLHYTCSSHFPYSQYQFWKNFQTLKIVILGLVLPLLVMVICYSGILKTLLRCRNEK
KRHRAVRLIFTIMIVYFLFWAPYNIVLLLNTFQEFFGLNNCSSSNRLDQAMQVTETLGMTHCCINPIIYAFVGEKF
RNYLLVFFQKHIAKRFCKCCSIFQQEAPERASSVYTRSTGEQEISVGLGSGGGGSGGSSSGGGGSGGSSSGGEF<u>DY</u>
<u>KDDDDK</u>

The genes encoding [5P12]CCL5,[68] CCL5[68] and [6P4]CCL5[69] for expression in *E. coli* have

been described previously.

Protein expression and purification
All protein constructs were expressed in *E. coli* BL21 (DE3) cultured in Lysogeny broth

medium (LB). For the preparation of the deuterated ^{15}N-labeled (and ^{13}C-labeled) NMR

samples, cells were grown in $D_2O/^{15}NH_4Cl$ (^{13}C-glucose) M9 minimal medium. For the

preparation of Ile-$\delta1$-$^{13}CH_3$-labeled arrestin2$^{1-393}$, the minimal medium was supplemented with

80 mg per liter of 2-ketobutyric acid-4-^{13}C,3,3-d_2 sodium salt hydrate following previously

reported procedures.[70] In all cases, cells were grown at 37˚C until the optical density at 600 nm

($OD_{600\ nm}$) reached 0.7–0.8. Thereafter protein expression was induced by the addition of 25 μM

isopropyl β-D-thiogalactopyranoside (IPTG) (arrestin2 constructs) or 1 mM IPTG (CLTC NTD

and AP2β2$^{701-937}$). The temperature was lowered to 18˚C (arrestin2 constructs) or 22˚C (CLTC

NTD and AP2β2$^{701-937}$) for an overnight expression, after which the cells were harvested by

centrifugation (30 min at 6000 rpm).

Arrestin2
The purification of arrestin2 has been described before.[29] In short, arrestin2$^{1-418}$, arrestin2^{1-}

^{393}, arrestin2$^{\Delta CBL}$ and arrestin2$^{\Delta AP2}$ were purified on a Ni-NTA HiTrap HP column (GE Life

Sciences) and the His-tag was removed by overnight cleavage with TEV protease (homemade).

The cleaved protein was further separated from impurities by a reverse IMAC step on a Ni-NTA

HiTrap HP column. An ion exchange chromatography step using a 5-ml HiTrap Q HP anion

exchange column was then carried out before the final gel filtration step on a HiLoad 16/600

Superdex 200 pg gel filtration column (GE Healthcare) equilibrated with 20 mM HEPES,

150 mM NaCl, pH 7.4 (SEC buffer I). Protein purity was confirmed by SDS-PAGE.

CLTC NTD

CLTC NTD was purified similarly to the arrestin2 protocol using a Ni-NTA HiTrap HP

column. The His-tag was removed by overnight cleavage with 3C protease (homemade),

followed by a reverse IMAC step. The protein was concentrated in a Vivaspin 20 (Sartorius)

concentrator (10-kDa MWCO) before a final gel filtration step on a HiLoad 16/600 Superdex

S75 pg gel filtration column (GE Healthcare) equilibrated with SEC buffer I. Protein purity was

confirmed by SDS-PAGE.

AP2β2

530 AP2β2 was purified similarly to CLTC NTD. Due to a change of the isoelectric point from

531 6.6 to 7.7 after 3C protease cleavage, the pH of the SEC buffer was changed to 7.0 (SEC buffer

532 II, 20 mM HEPES, 150 mM NaCl, pH 7.0).

Peptide synthesis

534 Phosphorylated peptides (Supplementary Table 1) corresponding to the last 22 residues of

535 the human CCR5 receptor (CCR5pp3, CCR5pp4, CCR5pp6) were obtained from the Tufts

536 University Core Facility for peptide synthesis.

Cell culture

HeLa (ATCC CCL2) cells and HeLa KO cell line (HeLa$^{-arr2/3}$)[71] devoid of arrestin2 and

arrestin3 were grown in high-glucose Dulbecco's modified Eagle's medium (Sigma-Aldrich)

with 10% fetal bovine serum (FBS, Biowest), 2 mM L-glutamine (#25030; Gibco),

100 U ml^{-1} penicillin G and 100 ng ml^{-1} streptomycin (P433; Sigma-Aldrich Chemie GmbH),

1 mM sodium pyruvate (#S8636; Sigma-Aldrich Chemie GmbH) at 37°C and 5% CO_2. For

transient cell transfections, cells were plated into six-well plates to reach 70% confluency the

following day and transfected with 0.3 μg of each arrestin2-YFP and CCR5 plasmid DNA

complexed with Helix-IN transfection reagent (OZ Biosciences).

Immunostaining and microscopy

In HeLa cell lines, arrestin2 and CCR5 were overexpressed by transiently transfecting

arrestin2-YFP and CCR5 respectively. CCR5 was stained with a primary FLAG antibody

(1:200, Thermo Fisher F3165) in 5% FBS in PBS. A secondary mouse (1:500) Alexa 633

(A21052, Invitrogen) antibody was used in 5% FBS in PBS. Fluoromount-G mounting medium

(Thermo Fischer) was used to mount the coverslips onto the imaging slide. Fluorescence and

DIC images were acquired with an ORCA-Flash 4.0 camera (Hamamatsu) mounted on an Axio

Imager.M2 fluorescence microscope with a 63x Plan-Apochromat objective (Carl Zeiss) and an

HXP 120C light source with ZEN 2.6 software. Image processing was performed using

OMERO.insight client,[72] and analyzed with Fiji software.[73]

To assess the recruitment of arrestin2 by CCR5, arrestin2-YFP- and CCR5-expressing cells

were incubated with 1 μM chemokine ligands. Individual coverslips were removed and fixed

with 4% PFA for 10 min at 0, 10, and 30 min post-ligand stimulation. Fixed cells were

permeabilized with 0.1% Triton X-100 in PBS for 5 min, rinsed with PBS twice and blocked

with 5% FBS in PBS. CCR5 was stained as mentioned previously and imaged. Colocalization

between arrestin2 and CCR5 was assessed by measuring the Mander's coefficient in 2-3 regions

of interest (ROI) of 100 μm^2 per cell using the Fiji JACOP plug-in.[74] The number of internalized

CCR5 puncta with and without arrestin2 was quantified manually by taking 3 ROIs of 100 μm^2

per cell using FiJi.

To image arrestin2/CCR5 trafficking to lysosome, arrestin2-YFP and CCR5 expressing cells

were treated with [6P4]CCL5. Individual coverslips were removed and fixed with 4% PFA for

10 min at 0, 10, and 30 min post stimulation. Fixed cells were permeabilized with 0.1% Triton

X-100 in PBS for 5 min, rinsed with PBS twice and blocked with 5% FBS in PBS. Lysosomes

were stained with LAMP1 antibody (1:200, Cell Signaling Technology, D2D11). CCR5 was

stained with FLAG antibody (1:200, Thermo Fisher F3165). Secondary rabbit (1:500) Alexa-

Fluor 568 (A110042 Invitrogen) and secondary mouse (1:500) Alexa-Fluor 633 (A21052,

Invitrogen) fluorescent antibodies were used respectively. Fixation, imaging and analysis was

performed as described above.

For assessing the effect of arrestin2 mutations, respective mutants were overexpressed with

CCR5 in HeLa$^{-arr2/3}$ and HeLa cells, followed by immunostaining and microscopy analysis as

described above (HeLa$^{-arr2/3}$: 60 min , HeLa: 30 min post [6P4]CCL5 stimulation).

Live-cell imaging

For live-cell imaging, Hela cells expressing arrestin2-YFP were seeded on an ibidi μ-slide

VI 0.4 channel slide (#80606, Ibidi), 24 hours prior to data acquisition. Live cell imaging was

performed in complete growth medium lacking phenol red at 37˚C with 5% CO_2 508 using an

inverted Axio Observer microscope (Zeiss) with a Plan Apochromat N 63x/1.40 oil DIC 509

M27 objective and a Photometrics Prime 95B camera. Filters with standard specifications for

GFP were used to image arrestin2-YFP. After selecting cells to be imaged, images were acquired

at 5 min intervals. Image processing was performed using OMERO.insight client,[72] and

analyzed with Fiji software.[73]

To image lysosome stained with lysotracker, Hela cells expressing arrestin2-YFP and CCR5

were seeded on an ibidi μ-slide VI 0.4 channel slide (#80606, Ibidi), 24 hours prior to data

acquisition. LysoTracker™ Deep Red (L12492) and peptide ligand was added in complete

growth medium lacking phenol red. Images were taken at 0 and 30 min timepoints.

Statistics

591   Statistical analysis was performed using GraphPad Prism 10.1.1 (GraphPad Software, Inc.,

592   San Diego, CA, USA). For statistical evaluation, the normality of the data was routinely tested

593   using the Shapiro-Wilk or Kolmogorov-Smirnov normality test. For the comparison of two

594   groups (Fig. 5a,b), a t-test was employed, whereas to compare three or more groups (Fig. 1d), a

595   one-way ANOVA was performed followed by a Kruskal-Wallis test.

NMR Experiments

NMR samples were prepared in SEC buffer I supplemented with 5% D_2O, 2 mM EDTA and 0.03% NaN_3 as 270-µl volumes in Shigemi tubes. The assignment sample contained 300 mM NaCl. The spectra were recorded on a Bruker AVANCE 14.1 T (600 MHz) spectrometer (arrestin2[1-393], AP2β2[701-937]) or a Bruker AVANCE 21.2 T (900 MHz) spectrometer (CLTC NTD) equipped with a TCI cryoprobe at 303 K. The spectra were processed with NMRPipe,[75] assigned using CcpNmr AnalysisAssign[76] and analyzed with NMRFAM-SPARKY.[77]

Arrestin2 assignment

For the assignment of arrestin2 backbone resonances, HNCACB-TROSY, HNCA-TROSY and HNCO-TROSY spectra were recorded on ^{15}N-, ^{13}C-, ^{2}H-labeled arrestin2[1-393] (100-150 µM) samples.

Arrestin2 titrations

For detecting the CLTC NTD-arrestin2 interaction on arrestin2 by NMR, a titration of 50 µM ^{15}N-, ^{2}H-labeled arrestin2[1-393] with unlabeled CLTC NTD up to an equimolar amount was monitored by ^{1}H-^{15}N HSQC-TROSY spectra recorded as 100 (^{15}N) x 512 (^{1}H) complex points and acquisition times of 50 ms (^{15}N) and 55 ms (^{1}H) for 4.5 h. Peak intensities I were determined by integration and the interaction was quantified relative to the intensity I_0 in the absence of CLTC NTD. An intensity attenuation larger than one standard deviation of I/I_0 calculated from all resonances was considered significant.

Similarly, a titration of 50 µM Ile-$\delta1$-$^{13}CH_3$, ^{2}H-labeled arrestin2[1-393] with unlabeled CLTC NTD up to a concentration of 600 µM was monitored by ^{1}H-^{13}C HMQC spectra recorded as 32 (^{13}C) x 512 (^{1}H) complex points and acquisition times of 20 ms (^{13}C) and 55 ms (^{1}H) for 1 h. The assignments of the methyl groups were transferred from BMRB ID:51131.[50] K_D values were obtained by nonlinear least-squares fitting using Matlab (Matlab_R2021b, MathWorks, Inc.) and the following equation:

$$\Delta\delta = \Delta\delta_{max} \frac{[L] + [P] + K_D - \sqrt{([L] + [P] + K_D)^2 - 4[P][L]}}{2[P]} \qquad (1)$$

621 where $\Delta\delta = \delta_{apo} - \delta_{bound}$ is the difference in the ^{13}C chemical shift, δ_{max} is the difference between

622 apo and the fully ligand-bound state, and [P] and [L] are the total protein (arrestin2) and ligand

623 (CLTC NTD) concentrations, respectively.

624 Chemical shift perturbations (CSP) were calculated from the relation:

$$CSP = \sqrt{\Delta\delta_H{}^2 + \left(\frac{\Delta\delta_{13C}}{3.6}\right)^2} \qquad (2)$$

where $\Delta\delta_H$ is the methyl proton chemical shift difference, and $\Delta\delta_{13C}$ is the ^{13}C-methyl chemical

shift difference between the apo and bound state. CSPs greater than one standard deviation from

the mean were considered significant.

CLTC NTD titrations
For detecting the CLTC NTD-arrestin2 interaction on CLTC NTD by NMR, a titration of

50 μM ^{15}N-, ^2H-labeled CLTC NTD with unlabeled arrestin2$^{1-393}$, arrestin2$^{1-418}$, arrestin2$^{\Delta CBL}$ or

arrestin2$^{1-393}$•CCR5pp6 up to an equimolar amount was monitored by ^1H-^{15}N HSQC-TROSY

spectra recorded as 100 (^{15}N) x 1024 (^1H) complex points and acquisition times of 35 ms (^{15}N)

and 80 ms (^1H) for 4.5 h. Assignments for CLTC NTD were transferred from BMRB

ID:25403.[51] Peak intensities were determined by quantifying peak volumes. Binding regions

were determined by intensity ratios as described above.

AP2β2 titrations
For detecting the AP2β2-arrestin2 interaction on AP2β2$^{701-937}$ by NMR, a titration of 50 μM

^{15}N-, ^2H-labeled AP2β2$^{701-937}$ with unlabeled arrestin2$^{1-418}$ or arrestin2$^{1-418}$•CCR5pp6 up to an

equimolar amount was monitored by ^1H-^{15}N HSQC-TROSY spectra recorded as 100 (^{15}N) x

512 (^1H) complex points and acquisition times of 50 ms (^{15}N) and 55 ms (^1H) for 1 h.

Assignments for AP2β2$^{701-937}$ were transferred from BMRB ID: 52320.[56] Peak intensities were

determined by quantifying peak volumes. Binding regions were determined by intensity ratios

as described above.

SEC binding assay

SEC binding assays were performed on an UltiMate 3000 HPLC system. Samples

containing single polypeptides or their mixtures as indicated in the text and figure legends

(arrestin2$^{1\text{-}418}$: 10 μM, AP2β2: 10 μM, CLTC NTD: 10 μM, CCR5pp0: 100 μM, CCR5pp3: 100

μM, CCR5pp4: 100 μM, CCR5pp6: 25-100 μM) were prepared as 270 μL volumes in SEC

buffer II. Samples were injected with an autosampler using a 250-μL sample-loop onto a self-

packed 4.2-ml S200 10/300 SEC column (length 25 mm, diameter 4.6 mm), which had been

pre-equilibrated with SEC buffer II. Complex formation was monitored by following protein

absorbance at 280 nm at 2.6 ml elution volume and the composition of the respective SEC

fractions of all prepared complexes was analyzed by SDS-PAGE (Supplementary Fig. 6c).

The apparent dissociation constant of the arrestin2$^{1\text{-}418}$-AP2β2$^{701\text{-}937}$ complex in the presence

of the CCR5 phosphopeptides could be determined by integrating the SEC absorption peaks of

the complex and of the apo proteins and using their respective extinction coefficients. The

increase of absorbance around 2.6 ml is a consequence of the additional AP2β2$^{701\text{-}937}$ in that

elution region. This region overlaps to some extent with the elution profile of apo arrestin2$^{1\text{-}418}$.

Taking into account the 1:1 stoichiometry, the amount of formed complex was estimated by

integrating the profile between 2.4 and 2.8 ml, subtracting the respective integral of the apo

arrestin2$^{1\text{-}418}$ profile, and dividing the difference by the AP2β2$^{701\text{-}937}$ extinction coefficient.

Since the phosphopeptides do not contain tryptophan, their absorbance contribution was

ignored.

Trypsin proteolysis

Trypsin proteolysis assays were carried out by pre-incubating 100 μL of 20 μM arrestin2$^{1\text{-}418}$

and 100 μM CCR5pp6 or 500 μM CCR5pp4/CCR5pp3 phosphopeptide with 1- (CCR5pp6), 2-

(CCR5pp6) and 5-molar equivalents (CCR5pp6, CCR5pp4, CCR5pp3) of AP2β2 protein at

room temperature for 5-10 min in the SEC buffer II. Following pre-incubation, 1 ng of Trypsin

Gold (Promega) was added, and the mixture further incubated at 35 ˚C under gentle shaking

(500 rpm, ThermoMixer). Samples for SDS-PAGE (10 µL mixed with 10 µL of 4x SDS loading

buffer) were taken at 0, 5, 10, 20, 30, 60, and 90 min. The reaction was quenched by boiling the

samples for 10 min at 95˚C. Samples were then loaded on 4-20% precast gradient gels and

visualized with Instant Blue protein stain (Abcam). Control reactions were run with apo

arrestin2$^{1-418}$ and arrestin2$^{1-418}$ incubated with 100 µM CCR5pp6 or 60-500 µM

CCR5pp4/CCR5pp3 phosphopeptide.

SDS-PAGE gels were analyzed using ImageJ software[78] by integrating the peak area of the

two bands corresponding to the full length arrestin2$^{1-418}$ (top) and cleaved protein at residue

R393 (bottom). The relative abundance (in %) of arrestin2$^{1-418}$ was calculated from the

respective integrals at 5 min upon trypsin addition.

References

1. Hauser, A. S., Attwood, M. M., Rask-Andersen, M., Schiöth, H. B. & Gloriam, D. E. Trends in GPCR drug discovery: new agents, targets and indications. *Nat Rev Drug Discov* **16**, 829–842 (2017).

2. Weis, W. I. & Kobilka, B. K. The Molecular Basis of G Protein–Coupled Receptor Activation. *Annual Review of Biochemistry* **87**, 897–919 (2018).

3. Komolov, K. E. & Benovic, J. L. G protein-coupled receptor kinases: Past, present and future. *Cellular Signalling* **41**, 17–24 (2018).

4. Gurevich, V. V. & Gurevich, E. V. GPCR Signaling Regulation: The Role of GRKs and Arrestins. *Frontiers in Pharmacology* **10**, (2019).

5. Hanyaloglu, A. C. & Zastrow, M. von. Regulation of GPCRs by Endocytic Membrane Trafficking and Its Potential Implications. *Annu. Rev. Pharmacol. Toxicol.* **48**, 537–568 (2008).

6. Otero, C., Groettrup, M. & Legler, D. F. Opposite Fate of Endocytosed CCR7 and Its Ligands: Recycling versus Degradation1. *The Journal of Immunology* **177**, 2314–2323 (2006).

7. Gu, S. *et al.* Ligand-Dependent Mechanisms of C-C Chemokine Receptor 5 (CCR5) Trafficking Revealed by APEX2 Proximity Labeling Proteomics. 2023.11.01.565224 Preprint at https://doi.org/10.1101/2023.11.01.565224 (2023).

8. Wolfe, B. L. & Trejo, J. Clathrin-Dependent Mechanisms of G Protein-coupled Receptor Endocytosis. *Traffic* **8**, 462–470 (2007).

9. Kaksonen, M. & Roux, A. Mechanisms of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **19**, 313–326 (2018).

10. Smith, S. M. & Smith, C. J. Capturing the mechanics of clathrin-mediated endocytosis. *Current Opinion in Structural Biology* **75**, 102427 (2022).

11. Dell'Angelica, E. C. Clathrin-binding proteins: Got a motif? Join the network! *Trends in Cell Biology* **11**, 315–318 (2001).

12. Lemmon, S. K. & Traub, L. M. Getting in Touch with the Clathrin Terminal Domain. *Traffic* **13**, 511–519 (2012).

13. Smith, C. J., Grigorieff, N. & Pearse, B. M. F. Clathrin coats at 21 Å resolution: a cellular assembly designed to recycle multiple membrane receptors. *The EMBO Journal* **17**, 4943–4953 (1998).

14. Popova, N. V., Deyev, I. E. & Petrenko, A. G. Clathrin-Mediated Endocytosis and Adaptor Proteins. *Acta Naturae* **5**, 62–73 (2013).

15. Kovtun, O., Dickson, V. K., Kelly, B. T., Owen, D. J. & Briggs, J. A. G. Architecture of the AP2/clathrin coat on the membranes of clathrin-coated vesicles. *Science Advances* **6**, eaba8381 (2020).

16. Kelly, B. T. *et al.* AP2 controls clathrin polymerization with a membrane-activated switch. *Science* **345**, 459–463 (2014).

17. Schmid, E. M. *et al.* Role of the AP2 β-Appendage Hub in Recruiting Partners for Clathrin-Coated Vesicle Assembly. *PLOS Biology* **4**, e262 (2006).

18. Collins, B. M., McCoy, A. J., Kent, H. M., Evans, P. R. & Owen, D. J. Molecular Architecture and Functional Model of the Endocytic AP2 Complex. *Cell* **109**, 523–535 (2002).

19. Smith, S. M., Baker, M., Halebian, M. & Smith, C. J. Weak Molecular Interactions in Clathrin-Mediated Endocytosis. *Frontiers in Molecular Biosciences* **4**, (2017).

20. Jackson, L. P. *et al.* A Large-Scale Conformational Change Couples Membrane Recruitment to Cargo Binding in the AP2 Clathrin Adaptor Complex. *Cell* **141**, 1220–1229 (2010).

21. Paing, M. M., Johnston, C. A., Siderovski, D. P. & Trejo, J. Clathrin Adaptor AP2 Regulates Thrombin Receptor Constitutive Internalization and Endothelial Cell Resensitization. *Molecular and Cellular Biology* **26**, 3231–3242 (2006).

22. Marchese, A., Paing, M. M., Temple, B. R. S. & Trejo, J. G Protein–Coupled Receptor Sorting to Endosomes and Lysosomes. *Annual Review of Pharmacology and Toxicology* **48**, 601–629 (2008).

23. Goodman, O. B. *et al.* β-Arrestin acts as a clathrin adaptor in endocytosis of the β2-adrenergic receptor. *Nature* **383**, 447–450 (1996).

24. Lefkowitz, R. J. & Shenoy, S. K. Transduction of Receptor Signals by ß-Arrestins. *Science* **308**, 512–517 (2005).

25. Goodman, O. B., Krupnick, J. G., Gurevich, V. V., Benovic, J. L. & Keen, J. H. Arrestin/Clathrin Interaction: LOCALIZATION OF THE ARRESTIN BINDING LOCUS TO THE CLATHRIN TERMINAL DOMAIN*. *Journal of Biological Chemistry* **272**, 15017–15022 (1997).

26. Krupnick, J. G., Goodman, O. B., Keen, J. H. & Benovic, J. L. Arrestin/Clathrin Interaction: LOCALIZATION OF THE CLATHRIN BINDING DOMAIN OF NONVISUAL ARRESTINS TO THE CARBOXYL TERMINUS*. *Journal of Biological Chemistry* **272**, 15011–15016 (1997).

27. Laporte, S. A., Oakley, R. H., Holt, J. A., Barak, L. S. & Caron, M. G. The Interaction of β-Arrestin with the AP-2 Adaptor Is Required for the Clustering of β2-Adrenergic Receptor into Clathrin-coated Pits*. *Journal of Biological Chemistry* **275**, 23120–23126 (2000).

28. Oakley, R. H., Laporte, S. A., Holt, J. A., Caron, M. G. & Barak, L. S. Differential Affinities of Visual Arrestin, βArrestin1, and βArrestin2 for G Protein-coupled Receptors Delineate Two Major Classes of Receptors*. *Journal of Biological Chemistry* **275**, 17201–17210 (2000).

29. Isaikina, P. *et al.* A key GPCR phosphorylation motif discovered in arrestin2·CCR5 phosphopeptide complexes. *Molecular Cell* **83**, 2108-2121.e7 (2023).

30. Alkhatib, G. The biology of CCR5 and CXCR4. *Current Opinion in HIV and AIDS* **4**, 96–103 (2009).

31. Zeng, Z., Lan, T., Wei, Y. & Wei, X. CCL5/CCR5 axis in human diseases and related treatments. *Genes & Diseases* **9**, 12–27 (2022).

32. Patterson, B. K. *et al.* CCR5 inhibition in critical COVID-19 patients decreases inflammatory cytokines, increases CD8 T-cells, and decreases SARS-CoV2 RNA in plasma by day 14. *International Journal of Infectious Diseases* **103**, 25–32 (2021).

33. Jiao, X. *et al.* Recent Advances Targeting CCR5 for Cancer and Its Role in Immuno-Oncology. *Cancer Res* **79**, 4801–4807 (2019).

34. Maharana, J. *et al.* Structural snapshots uncover a key phosphorylation motif in GPCRs driving β-arrestin activation. *Molecular Cell* **83**, 2091-2107.e7 (2023).

35. Huang, W. *et al.* Structure of the neurotensin receptor 1 in complex with β-arrestin 1. *Nature* **579**, 303–308 (2020).

36. Bous, J. *et al.* Structure of the vasopressin hormone-V2 receptor-β-arrestin1 ternary complex. *Sci Adv* **8**, eabo7761 (2022).

37. Kang, Y. *et al.* Crystal structure of rhodopsin bound to arrestin by femtosecond X-ray laser. *Nature* **523**, 561–567 (2015).

38. Oakley, R. H., Laporte, S. A., Holt, J. A., Barak, L. S. & Caron, M. G. Molecular Determinants Underlying the Formation of Stable Intracellular G Protein-coupled Receptor-β-Arrestin Complexes after Receptor Endocytosis*. *Journal of Biological Chemistry* **276**, 19452–19460 (2001).

39. Signoret, N. *et al.* Agonist-induced Endocytosis of CC Chemokine Receptor 5 Is Clathrin Dependent. *Mol Biol Cell* **16**, 902–917 (2005).

40. Fraile-Ramos, A., Kohout, T. A., Waldhoer, M. & Marsh, M. Endocytosis of the Viral Chemokine Receptor US28 Does Not Require Beta-Arrestins But Is Dependent on the Clathrin-Mediated Pathway. *Traffic* **4**, 243–253 (2003).

41. Gaertner, H. *et al.* Highly potent, fully recombinant anti-HIV chemokines: Reengineering a low-cost microbicide. *PNAS* **105**, 17706–17711 (2008).

42. Hughes, C. E. & Nibbs, R. J. B. A guide to chemokines and their receptors. *The FEBS Journal* **285**, 2944–2971 (2018).

43. Isaikina, P. *et al.* Structural basis of the activation of the CC chemokine receptor 5 by a chemokine agonist. *Science Advances* (2021) doi:10.1126/sciadv.abg8685.

44. Beautrait, A. *et al.* A new inhibitor of the β-arrestin/AP2 endocytic complex reveals interplay between GPCR internalization and signalling. *Nat Commun* **8**, 15054 (2017).

45. Martins, E. *et al.* Arrestin Recruitment to C-C Chemokine Receptor 5: Potent C-C Chemokine Ligand 5 Analogs Reveal Differences in Dependence on Receptor Phosphorylation and Isoform-Specific Recruitment Bias. *Mol Pharmacol* **98**, 599–611 (2020).

46. Mueller, A., Kelly, E. & Strange, P. G. Pathways for internalization and recycling of the chemokine receptor CCR5. *Blood* **99**, 785–791 (2002).

47. Kang, D. S. *et al.* Structure of an Arrestin2-Clathrin Complex Reveals a Novel Clathrin Binding Domain That Modulates Receptor Trafficking. *J Biol Chem* **284**, 29860–29872 (2009).

48. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

49. Tugarinov, V., Hwang, P. M., Ollerenshaw, J. E. & Kay, L. E. Cross-Correlated Relaxation Enhanced 1H−13C NMR Spectroscopy of Methyl Groups in Very High Molecular Weight Proteins and Protein Complexes. *J. Am. Chem. Soc.* **125**, 10420–10428 (2003).

50. Shiraishi, Y. *et al.* Biphasic activation of β-arrestin 1 upon interaction with a GPCR revealed by methyl-TROSY NMR. *Nat Commun* **12**, 7158 (2021).

51. Zhuo, Y. *et al.* Nuclear Magnetic Resonance Structural Mapping Reveals Promiscuous Interactions between Clathrin-Box Motif Sequences and the N-Terminal Domain of the Clathrin Heavy Chain. *Biochemistry* **54**, 2571–2580 (2015).

52. Kern, R. C., Kang, D. S. & Benovic, J. L. Arrestin2/Clathrin Interaction Is Regulated by Key N- and C-Terminal Regions in Arrestin2. *Biochemistry* **48**, 7190–7200 (2009).

53. Laporte, S. A., Miller, W. E., Kim, K.-M. & Caron, M. G. β-Arrestin/AP-2 Interaction in G Protein-coupled Receptor Internalization: IDENTIFICATION OF A β-ARRESTIN BINDING SITE IN β2-ADAPTIN*. *Journal of Biological Chemistry* **277**, 9247–9254 (2002).

54. Kim, Y.-M. & Benovic, J. L. Differential Roles of Arrestin-2 Interaction with Clathrin and Adaptor Protein 2 in G Protein-coupled Receptor Trafficking*. *Journal of Biological Chemistry* **277**, 30760–30768 (2002).

55. Moaven, H. *et al.* Visual arrestin interaction with clathrin adaptor AP-2 regulates photoreceptor survival in the vertebrate retina. *Proceedings of the National Academy of Sciences* **110**, 9463–9468 (2013).

56. Naudi-Fabra, S. *et al.* An extended interaction site determines binding between AP180 and AP2 in clathrin mediated endocytosis. *Nat Commun* **15**, 5884 (2024).

57. Xiao, K., Shenoy, S. K., Nobles, K. & Lefkowitz, R. J. Activation-dependent Conformational Changes in β-Arrestin 2*. *Journal of Biological Chemistry* **279**, 55744–55753 (2004).

58. Barsi-Rhyne, B., Manglik, A. & von Zastrow, M. Discrete GPCR-triggered endocytic modes enable β-arrestins to flexibly regulate cell signaling. *eLife* **11**, e81563 (2022).

59. Escola, J.-M., Kuenzi, G., Gaertner, H., Foti, M. & Hartley, O. CC Chemokine Receptor 5 (CCR5) Desensitization: CYCLING RECEPTORS ACCUMULATE IN THE TRANS-GOLGI NETWORK*. *Journal of Biological Chemistry* **285**, 41772–41780 (2010).

60. Ariel, A. *et al.* Apoptotic neutrophils and T cells sequester chemokines during immune response resolution through modulation of CCR5 expression. *Nat Immunol* **7**, 1209–1216 (2006).

61. Lally, C. C. M., Bauer, B., Selent, J. & Sommer, M. E. C-edge loops of arrestin function as a membrane anchor. *Nat Commun* **8**, 14258 (2017).

62. Janetzko, J. *et al.* Membrane phosphoinositides regulate GPCR-β-arrestin complex assembly and dynamics. *Cell* **185**, 4560-4573.e19 (2022).

63. Grimes, J. *et al.* Plasma membrane preassociation drives β-arrestin coupling to receptors and activation. *Cell* **186**, 2238-2255.e20 (2023).

64. Tohgo, A. *et al.* The Stability of the G Protein-coupled Receptor-β-Arrestin Interaction Determines the Mechanism and Functional Consequence of ERK Activation *. *Journal of Biological Chemistry* **278**, 6258–6267 (2003).

65. Jean-Charles, P.-Y., Kaur, S. & Shenoy, S. K. G Protein–Coupled Receptor Signaling Through β-Arrestin–Dependent Mechanisms. *Journal of Cardiovascular Pharmacology* **70**, 142 (2017).

66. Benz, C. *et al.* Proteome-scale mapping of binding sites in the unstructured regions of the human proteome. *Molecular Systems Biology* **18**, e10584 (2022).

67. Violin, J. D., Ren, X.-R. & Lefkowitz, R. J. G-protein-coupled receptor kinase specificity for beta-arrestin recruitment to the beta2-adrenergic receptor revealed by fluorescence resonance energy transfer. *J Biol Chem* **281**, 20577–20588 (2006).

68. Wiktor, M., Hartley, O. & Grzesiek, S. Characterization of Structure, Dynamics, and Detergent Interactions of the Anti-HIV Chemokine Variant 5P12-RANTES. *Biophysical Journal* **105**, 2586–2597 (2013).

69. Isaikina, P. *et al.* Chapter 6 - Preparation of a stable CCL5·CCR5·Gi signaling complex for Cryo-EM analysis. in *Methods in Cell Biology* (ed. Shukla, A. K.) vol. 169 115–141 (Academic Press, 2022).

70. Tugarinov, V., Kanelis, V. & Kay, L. E. Isotope labeling strategies for the study of high-molecular-weight proteins by solution NMR spectroscopy. *Nat Protoc* **1**, 749–754 (2006).

71. D'Agostino, G. *et al.* β-Arrestin1 and β-Arrestin2 Are Required to Support the Activity of the CXCL12/HMGB1 Heterocomplex on CXCR4. *Front Immunol* **11**, 550824 (2020).

72. Allan, C. *et al.* OMERO: flexible, model-driven data management for experimental biology. *Nat Methods* **9**, 245–253 (2012).

73. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**, 676–682 (2012).

74. Bolte, S. & Cordelières, F. P. A guided tour into subcellular colocalization analysis in light microscopy. *Journal of Microscopy* **224**, 213–232 (2006).

75. Delaglio, F. *et al.* NMRPipe: A multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* **6**, 277–293 (1995).

76. Skinner, S. P. *et al.* CcpNmr AnalysisAssign: a flexible platform for integrated NMR analysis. *J Biomol NMR* **66**, 111–124 (2016).

77. Lee, W., Tonelli, M. & Markley, J. L. NMRFAM-SPARKY: enhanced software for biomolecular NMR spectroscopy. *Bioinformatics* **31**, 1325–1327 (2015).

78. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**, 671–675 (2012).

Acknowledgments

We gratefully acknowledge I. Hertel and M. Rogowski for the expression and purification

of proteases and CCL5 ligands. We thank Prof. Mariagrazia Uguccioni and Dr. Valentina

Cecchinato (Institute for Research in Biomedicine, Bellinzona, Switzerland) for the HeLa$^{-arr2/3}$

cells. This work was supported by the Swiss National Science Foundation (grants 31-201270

and IZLIZ3-200298 to S.G., 31-197779 and 31-185127 to A.S.), and by a Fellowship for

Excellence by the Biozentrum Basel International PhD Program to I.P.

Author contributions

859   I.P., S.D., P.I., A.S., and S.G. conceived the study. I.P. and L.A.A generated all the

860 constructs and mutants for the cellular assays. S.D. performed the cellular assays and analyzed

861 the cellular data with guidance from A.S. I.P. expressed and purified proteins. I.P. recorded

862 NMR experiments and analyzed NMR data. I.P. designed and analyzed SEC and trypsin

863 proteolysis assay. I.P. and S.G. wrote the manuscript with inputs from all co-authors.

Competing interest

865   The authors declare that they have no competing interests.

Data and materials availability

867   All data needed to evaluate the conclusions in the paper are present in the main text and/or

868 Supplementary information.

 Figures

Fig. 1: CCR5 internalization in the presence of the chemokine ligands. CCR5 internalization

monitored in HeLa cells co-transfected with plasmids containing arrestin2-YFP (green) and

CCR5 genes (magenta) stimulated with **a**, [5P12]CCL5 (antagonist) **b**, CCL5 (natural agonist)

c, [6P4]CCL5 (super-agonist). Cells were stimulated for 0, 10, and 30 min with chemokine

ligands before fixation and preparation for fluorescence microscopy. Each image is

accompanied by a zoomed region of interest (white squares) showing deconvolved CCR5 and

arrestin2-YFP signals. **d**, Mander's colocalization coefficients of the CCR5 and arrestin2-YFP

signals in the absence and presence of chemokine ligands 30 min after stimulation. Individual

values, as well as mean and standard deviation are shown for N=3 biological replicates and n=45

ROIs from 20 cells; ****: $P < 0.0001$; ns: not significant ($P > 0.9999$). **e** Number of CCR5-

positive (CCR5$^+$) and CCR5/arrestin2-positive (CCR5$^+$/arrestin2$^+$) puncta after 30 min

[6P4]CCL5 ligand stimulation. Individual values, as well as mean and standard deviation are

shown for N=3 biological replicates and n=90 ROIs from 30 cells; ns: not significant ($P >$

0.9999). **f**, CCR5 and arrestin2 fluorescence signals along the trajectory from the plasma

membrane to the nucleus (blue line in panel **c**) 30 min after [6P4]CCL5 ligand stimulation. **g**,

Lysosomal trafficking of the CCR5 (magenta) and arrestin2 (cyan) complex in presence of the

[6P4]CCL5 monitored in the HeLa cells using LAMP1 antibodies (yellow). No recruitment to

the lysosome is observed.

Fig. 2: Arrestin2-CLTC NTD interactions detected by arrestin2 NMR spectra. a,

AlphaFold2 model of Arrestin2$^{1-418}$ showing the N- and C-domain (gray) followed by the

clathrin-binding loop (CBL, orange), C-terminal β20 strand (yellow) and the disordered C-

terminal tail (blue). The C-terminal strand β20 forms a parallel β-sheet with the N-terminal

strand β1 (pink) of inactive arrestin2 and is released from the arrestin2 upon activation by

phosphopeptides. Arrestin2 residues interacting with CLTC NTD are shown as green (red for

main interacting residue I377) spheres. The clathrin and AP2 binding motifs are indicated in the

arrestin2 sequence on the top. **b**, Part of ^{1}H-^{15}N TROSY spectrum of apo arrestin2$^{1-393}$ (black)

and upon addition of an equimolar amount of CLTC NTD (green). Assigned CBL arrestin2$^{1-393}$

residues are indicated. **c**, Intensity ratios of assigned complexed vs. apo arrestin2$^{1-393}$ CBL

resonances. The region between T374 and T381 undergoes significant intensity attenuation. **d**,

^{1}H-^{13}C HMQC spectra of apo Ile-δ1-^{13}CH$_3$, ^{2}H-labeled arrestin2$^{1-393}$ (black) and upon addition

of an equimolar amount of the CLTC NTD (green). **e**, Intensity attenuation (top) and chemical

shift perturbation (bottom) of arrestin2$^{1-393}$ isoleucine ^{1}H$_3$-^{13}C^{δ1} resonances upon CLTC NTD

addition. Out of 15 isoleucine residues, significant changes are observed only for I377. **f**, Left:

part of ^{1}H-^{13}C HMQC spectrum showing resonance shifts of I377 upon CLTC NTD binding.

Right: detected chemical shift changes as a function of CLTC NTD concentration. The solid

line depicts a global non-linear least-square fit to the data points with respective dissociation

constant.

Fig. 3: Arrestin2-CLTC NTD interactions detected by CLTC NTD NMR spectra. (Left):

intensity attenuation I/I_0 of CLTC NTD ^{1}H-^{15}N TROSY resonances upon addition of equimolar

amounts of various arrestin2 constructs: **a**, apo arrestin2$^{1-393}$ **b**, CCR5pp6-activated arrestin2$^{1-393}$

c, full-length arrestin2$^{1-418}$ **d**, arrestin2$^{\Delta CBL}$ with removed clathrin-binding motif. Regions that

undergo significant attenuation upon interaction with arrestin2 are indicated by arrows. Right:

residues undergoing significant intensity attenuation (one standard deviation below the average

signal attenuation) are marked on the structure of CLTC NTD (PDB: 3GD1) together with a

schematic representation of the respective arrestin2 construct.

Fig. 4: Arrestin2 interaction with the C-terminal domain of the AP2β2. a, Structure of

AP2β2$^{701-937}$ (orange) in complex with arrestin C-terminal peptide (blue) (PDB: 2IV8).

Important residues from both chains stabilizing the interaction are depicted in stick

representation. **b-f** SEC profiles of arrestin2$^{1-418}$, AP2β2$^{701-937}$, various phosphopeptides and

their mixtures. The annotated color coding below the SEC profile indicates the individual

sample composition. In panels **b, c, e**, 'mixture' refers to a SEC sample containing all of the

individual components indicated on the left. In panels **d, f**, the primary sample composition is

indicated above the SEC profile, and the annotated color coding below the SEC profile indicates

the added component to the primary sample composition. **g**, Apparent affinities of the

complexes derived by integrating the SEC peaks marked in Fig. 4f and scaling the integrals by

the respective extinction coefficients. 'nd' indicates 'not detected'. **h, i** (Left): intensity

attenuation I/I_0 of AP2β2$^{701-937}$ ^{1}H-^{15}N TROSY resonances upon addition of equimolar amounts

of apo arrestin2$^{1-418}$ (**h**) or CCR5pp6-activated arrestin2$^{1-418}$ (**i**). Regions that undergo significant

attenuation upon interaction with arrestin2 are indicated by arrows. Right: residues undergoing

significant intensity attenuation are marked on the structure of AP2β2 (PDB: 2IV8) together

with a schematic representation of the respective arrestin2 construct.

Fig. 5: Dependence of CCR5 internalization on arrestin2 interactions with clathrin or AP2

monitored in HeLa$^{-arr2/3}$ cells. a, CCR5 internalization induced by 60 min [6P4]CCL5 ligand

stimulation in HeLa cells co-transfected with plasmids containing CCR5 and arrestin2-YFP,

arrestin2-YFP$^{\Delta LIELD}$, arrestin2-YFP^{R395A}, arrestin2-YFP$^{\Delta AP2}$ or empty pcDNA3.1. **b, c,**

Internalization quantified by counting the number of puncta (**b**) positive for CCR5 and arrestin2

(CCR5$^+$/arrestin2$^+$) or (**c**) only CCR5$^+$ 60 min after ligand stimulation in the HeLa$^{-arr2/3}$ cells. No

significant difference in CCR5 internalization is detected for the arrestin2-YFP$^{\text{ALIELD}}$ and
arrestin2-YFP$^{\text{R395A}}$ constructs (ns, not significant: P > 0.9999) whereas the absence of the AP2
binding motif in the arrestin2$^{\Delta\text{AP2}}$-YFP construct causes a significant (****: P < 0.0001)
reduction of CCR5 internalization 60 min after ligand incubation. Mean and standard deviation
are shown for N=3 biological replicates and n=90 ROIs from 30 cells.

Fig. 6: Overall scheme of arr-class A and b GPCR internalization. Schematic difference of
arrestin2-mediated internalization of arr-class B (left) vs. arr-class A (right) GPCRs. Arr-class
B GPCRs bind stably to arrestin due to their high levels of phosphorylation. This results in a
robust release of the arrestin C-terminus, a stable interaction with AP2 and formation of a long-
lived GPCR•arrestin complex. Arr-class A GPCRs bind weakly to arrestin due to their poor
phosphorylation. They require stabilization of the arrestin complex by membrane-bound PIP$_2$
molecules. The arrestin C-terminus is not fully released and consequently the interaction with
AP2 is unstable.

Figure 1

Figure 2

Figure 4

Figure 5

arr-class B GPCR: strong arrestin interaction

arr-class A GPCR: transient arrestin interaction