1 **Non-gonadal PIWI protein, Aubergine, regulates regenerative stem cell**
2 **proliferation and tumourigenesis in the *Drosophila* adult intestine.**
3

Karen Bellec[1], Lynsey R Carroll[1], Kathryn AF Pennel[1], Yuanliangzi Tian[1], Yachuan
Yu[2], Alexander R Cameron[1], Fabiana Herédia[3,4], Alisson M Gontijo[3,4], Joanne
Edwards[1], Rippei Hayashi[5,6, ‡], Julia B Cordero[1,2, ‡]

[1]School of Cancer Sciences, Wolfson Wohl Cancer Research Centre, University of
Glasgow, Garscube Estate, Switchback Road, Glasgow G61 1QH, UK. [2]CRUK
Scotland Institute, Garscube Estate, Switchback Road, Glasgow G61 1BD, UK. [3]
iNOVA4Health, Nova Medical School, Faculdade de Ciências Médicas, NMS, FCM,
Nova University of Lisbon, Lisbon, Portugal. [4]cE3c - Centre for Ecology, Evolution
and Environmental Changes & CHANGE - Global Change and Sustainability
Institute, Department of Animal Biology, Faculty of Sciences, University of Lisbon,
Lisbon, Portugal. [5]John Curtin School of Medical Research, The Australian National
University, Acton, Australian Capital Territory, Australia. [6]The Shine-Dalgarno Centre
for RNA Innovation, The Australian National University, Acton, Australian Capital
Territory, Australia.

[‡]Authors   for   correspondence   (rippei.hayashi@anu.edu.au,
julia.cordero@glasgow.ac.uk).

Abstract

Somatic stem cells are executors of physiological and pathological proliferation of adult self-renewing tissues, such as the intestine. The actions of intestinal stem cells (ISCs) rely on the integration of cell intrinsic and niche-derived signals, which are necessary to achieve a balanced response to the multiple stimuli that constantly challenge tissue homeostasis and organismal health. Disruption of such balance is causative of age-associated tissue dysfunction and hyperproliferative conditions, including inflammation and cancer.

The highly conserved PIWI-interacting RNAs (piRNAs) biosynthesis pathway, also known as the PIWI pathway, has been classically studied in the *Drosophila* germline for its role in the repression of transposable elements (TEs) and the regulation of germline stem cell homeostasis. Recent reports have emerged on a role of Piwi, the founding member of the pathway, in the maintenance of ISC homeostasis in the adult *Drosophila* midgut. The implications of these findings regarding a general role of the PIWI pathway and piRNAs in the intestine remain to be addressed. Here, we characterise a cell autonomous role of the PIWI family protein Aubergine (Aub) in ISCs. We show that, while dispensable for homeostatic self-renewal of the midgut, inducible Aub is essential to regulate ISC proliferation following acute damage of the intestinal epithelium and in oncogenic settings. Our work suggests that the role of Aub in ISCs is independent of its piRNAs regulatory function. Instead, Aub drives ISC proliferation through the regulation of protein synthesis, including induction of regenerative stem cell factors Myc and Sox21a, and a parallel pathway involving the eukaryotic initiation factor 3 complex (eIF3). In summary, our results discover a role for Aub in damage-induced proliferation of the adult *Drosophila* intestine involving the regulation of major regenerative pathways and the protein translation machinery. Furthermore, we present genetically defined hyperplastic settings fostering distinct functions of Aub in intestinal tumourigenesis.

Introduction

Since their discovery as regulators of germ line stem cell self-renewal in the fruit fly *Drosophila melanogaster* (Cox et al., 1998, Lin and Spradling, 1997, Cox et al., 2000), the highly conserved proteins of the PIWI family have been extensively studied for their role in protecting and maintaining genome integrity in animal germline through transcriptional and post-transcriptional silencing of TEs (Siomi et al., 2011, Vagin et al., 2006, Brennecke et al., 2007, Li et al., 2009, Aravin et al., 2007b, Houwing et al., 2007, Aravin et al., 2007a, Saito et al., 2006, Aravin et al., 2006, Girard et al., 2006). Genome defence mechanisms by the PIWI pathway rely on the biogenesis and maturation of piRNAs, a specific type of 24-32 nucleotides long small non-coding RNAs, that bind the PIWI-clade Argonaute proteins hereafter called PIWI proteins. piRNAs are derived from transposon-dense genomic loci called piRNA clusters (Brennecke et al., 2007). After a precursor RNA is transcribed and exported from the nucleus, a set of dedicated RNA helicases and nucleases process it and load the mature piRNA onto a PIWI protein. The PIWI protein loaded with a piRNA exerts gene silencing by cleaving mRNAs or recruiting chromatin modification enzymes to install heterochromatin. In the *Drosophila* germline, PIWI proteins Piwi and Aub mainly bind piRNAs that target TEs in the nucleus and the cytoplasm, respectively (Pane et al., 2007, Nishimasu et al., 2012, Brennecke et al., 2007, Ipsaro et al., 2012), while Argonaute 3 (AGO3) predominantly binds TE sense piRNAs to aid the production of TE antisense piRNAs (Gunawardane et al., 2007, Brennecke et al., 2007, Han et al., 2015, Wang et al., 2015b, Malone et al., 2009, Li et al., 2009).

While in-depth knowledge regarding the composition and molecular function of the PIWI pathway derives primarily from its study in the germline and its role in TEs repression, numerous reports have documented the presence of PIWI proteins in adult somatic tissues and/or roles for these proteins expanding beyond piRNAs and TE regulation (Ross et al., 2014, Pal-Bhadra et al., 2004, Brower-Toland et al., 2007, Huang et al., 2013, Mochizuki et al., 2002, Jones et al., 2016, Sousa-Victor et al., 2017, Tang et al., 2023, Rajasethupathy et al., 2012, Sharma et al., 2001).

Pioneering work in *Drosophila* reported ectopic expression and tumour promoter functions of germline genes, including PIWI pathway components, in a malignant brain tumour model (Janic et al., 2010). Similarly, growing evidence in humans suggests misexpression of PIWI proteins and/or piRNAs in multiple cancers (Qiao et

al., 2002, Shi et al., 2020, Wang et al., 2021, Huang et al., 2021, Wang et al., 2016,

Li et al., 2019, Martinez et al., 2015, Lin et al., 2021, Krishnan et al., 2016, Xie et al.,

2018, Martinez et al., 2016). However, the link between ectopic PIWI gene

expression or piRNAs as well as the identity of piRNAs in cancer remain debatable

(Shi et al., 2020, Li et al., 2020, Genzor et al., 2019, Zhang et al., 2013, Liu et al.,

2021, Tosar et al., 2021). Similarly, in colorectal cancer (CRC), upregulation of

several piRNAs has been correlated with cancer progression, metastasis, and poor

overall survival (Yin et al., 2019, Weng et al., 2018). Yet, the functional significance

of PIWI proteins and piRNAs in intestinal health and disease remains largely

unknown.

Most similar to the mammalian small intestine, the adult *Drosophila* midgut

epithelium is maintained and repaired by dedicated stem cells (Micchelli and

Perrimon, 2006, Ohlstein and Spradling, 2006) **(Fig. 1A)**. In addition to intestinal

stem cells (ISCs), the fly midgut epithelium consists of two subtypes of

undifferentiated stem cell progeny, namely, enteroblasts (EBs) and pre-

enteroendocrine cells (EEPs), which are precursors of absorptive enterocytes (ECs)

and secretory enteroendocrine cells (EEs), respectively (Beehler-Evans and

Micchelli, 2015, Zeng and Hou, 2015, Chen et al., 2018) (**Fig.1A'**). Epithelium

intrinsic cells, including EBs and ECs, or cells from gut associated tissues, such as

the visceral muscle (VM), terminal tracheal cells (TTCs) and enteric neurons (ENs)

are components of the intestinal microenvironment, which non-autonomously

influence ISC proliferation through paracrine signalling (Perochon et al., 2021,

Tamamouna et al., 2021, Petsakou and Perrimon, 2023) (**Fig.1A'**).

Recent work in the adult *Drosophila* midgut has identified a functional role of

stem cell expressed Piwi in the maintenance of ISCs during tissue homeostasis

(Tang et al., 2023) or following pathogenic and age-associated stress (Sousa-Victor

et al., 2017). Whether these observations reflect on a general role of the PIWI

pathway and piRNAs in the adult intestine remain to be addressed. In our study, we

discovered that the PIWI protein Aub is upregulated within the stem/progenitor

compartment of the adult midgut in response to intestinal damage or oncogenic

stress and it is required to regulate regenerative and hyperplastic ISC proliferation.

Despite the presence of a piRNA signature in the midgut and in ISCs/EBs revealed

by small RNA sequencing, combined genetic and functional studies indicate that the

induction of ISC proliferation by Aub is independent of its canonical piRNAs

regulatory function. Mechanistically, Aub drives regenerative proliferation in the
intestine through the regulation of protein synthesis, including induction of stem cell
Myc and Sox21a, and activation of a parallel pathway involving the eukaryotic
initiation factor 3 complex (eIF3). Interestingly, we identified genetically defined
tumour settings revealing distinct roles of Aub in hyperplastic ISC proliferation.
Altogether, our results uncover a previously unrecognised role of Aub in
physiological and pathological proliferation of the adult *Drosophila* midgut and
provide defined platforms for further *in vivo* studies on the role of PIWI proteins and
piRNAs in intestinal health and disease.
128

Results

Aub is required for regenerative proliferation of ISCs following damage to the adult midgut epithelium.

To study the role of the PIWI pathway in the adult *Drosophila* intestine, we first assessed the expression of its core components *piwi*, *aub* and *ago3,* by RT-qPCR in midguts from mated females (**Fig.1A**, **B**). We observed low but detectable levels of *piwi*, *aub* and *ago3* mRNAs in homeostatic midguts with a slight enrichment of *piwi* and *ago3* in the posterior half (R4-R5) of the midgut (**Fig.1A**, **B**), a region that is characterised for its higher degree of ISC proliferation and predominantly studied in intestinal regeneration research (Buchon et al., 2013, Marianes and Spradling, 2013). We next assessed the role of the PIWI pathway in intestinal regeneration by feeding wild type, *piwi*, *aub* or *ago3* whole mutant flies with sucrose or with the pathogen *Pseudomonas entomophila* (*Pe*) to induce intestinal damage (Jiang et al., 2009). We quantified proliferating ISCs in posterior midguts of mated females by staining against the mitotic marker phosphorylated histone H3 (PH3) (**Fig.1C-E**). As expected, *Pe* feeding caused robust ISC proliferation in wild type flies when compared to their sucrose fed counterparts (**Fig.1C, D**). Loss of *piwi* mildly diminished intestinal regeneration in agreement with previously described roles of the protein in homeostatic and post-stress ISC maintenance (Sousa-Victor et al., 2017, Tang et al., 2023) (**Fig.1C, D**).

Interestingly, aub^{HN2}/aub^{QC42} mutant animals showed a significantly greater impairment in midgut regeneration (**Fig.1C, D**). aub^{HN2}/aub^{QC42} trans heterozygotes are the gold standard and best characterised *aub* loss of function mutants, used in most studies where Aub function has been analysed (Malone et al., 2009, Wang et al., 2015b, Rojas-Ríos et al., 2017). These alleles were generated by ethyl methanesulfonate mutagenesis (Schüpbach and Wieschaus, 1991). aub^{HN2} contains a nonsense mutation while aub^{QC42} contains a point mutation, both resulting in a non-functional protein. We also tested other and less characterized loss of function alleles of *aub*, aub^{g1}/aub^{g2} and aub^{clash}, which showed consistent, albeit milder impairment of intestinal regeneration (**Fig.1E**). aub^{g1}/aub^{g2} have been generated using CRISPR/Cas9 and contain a frameshift mutation resulting in an early translational stop. This trans heterozygote combination is considered a molecular null based on morphological phenotypes, protein levels and expression of small

163   subset of TEs in germline studies (Senti et al., 2015). *aub^clash* is a loss of function

164   allele characterised in germline studies based on its effect on germline morphology

165   and female/male fertility (Sahin et al., 2016).

166   Unexpectedly, whole body loss of *ago3*, an obligate partner of *aub* in the

Ping-Pong pathway, or ISCs and EBs (stem/progenitors) gene knockdown by RNAi

expressed under the control of the temperature-sensitive *escargot-Gal4, UAS-GFP;*

tubGal80 (*esg^ts*), had no impact on intestinal regeneration upon damage (**Fig.1C, D**

and **Fig.S1A**). Importantly, conditional knockdown of *aub* in ISCs/EBs was sufficient

to abolish regenerative ISC proliferation (**Fig.1F, G** and **Fig.S1A**). However, unlike

piwi (Tang et al., 2023), knocking down *aub* did not cause any detectable effects in

homeostatic ISC renewal (**Fig.S1B, C**). These data point to a specific role of

stem/progenitor cell Aub in regenerative ISC proliferation of the adult *Drosophila*

midgut. Consistently, immunostaining experiments confirmed the post-transcriptional

upregulation of Aub expression in ISCs/EBs upon midgut damage (**Fig.1H, J, K**),

which was markedly reduced by RNAi driven *aub* knockdown in ISCs/EBs (**Fig.1I, J**).

Next, to distinguish the cell type where *aub* functions to regulate ISC proliferation, we

induced gene knockdown in EBs or ISCs only using cell specific drivers *Su(H)GBE-*

Gal4,UAS-GFP; tub-Gal80^ts or *esg-Gal4,UAS-GFP; Su(H)GBE-Gal80, tub-Gal80^ts*,

respectively (hereafter referred to as *EB^ts* and *ISC^ts*, respectively; **Fig.1L, M**)

(ZengChauhan and Hou, 2010, Wang et al., 2014). While no impact on normal

intestinal regeneration was observed upon knocking down of *aub* in EBs only

(**Fig.1L**), gene knockdown of *aub* in ISCs was sufficient to recapitulate the

impairment in regeneration observed in midguts with dual ISCs/EBs *aub* RNAi

expression (**Fig.1M**). Collectively, these results demonstrate a cell autonomous role

of damage induced Aub in regenerative ISC proliferation in the adult *Drosophila*

midgut.

Midgut stem/progenitor cells express TE piRNAs independently of Aub.

191   Aub and the PIWI pathway have been extensively described for their role in

the repression of germline TEs through piRNAs (Brennecke et al., 2007,

Gunawardane et al., 2007, Malone et al., 2009, Wang et al., 2015b). However,

broader biological functions of Aub have been described and include the regulation

of localisation, stability and translation of specific mRNAs in the *Drosophila* germline

and in the embryo (Barckmann et al., 2015, Dufourt et al., 2017, Ramat et al., 2020,

Becalska et al., 2011, WilsonConnell and Macdonald, 1996, Cook et al., 2004, Ma et

al., 2017, Rojas-Ríos et al., 2017). Such roles of Aub rely on piRNA-dependent

(Barckmann et al., 2015, Dufourt et al., 2017, Ramat et al., 2020, Rojas-Ríos et al.,

2017, Rouget et al., 2010, Vourekas et al., 2016, Rojas-Ríos et al., 2023) and

piRNA-independent mechanisms (Ma et al., 2017). Whether there is involvement of

piRNAs in the role of Piwi in the adult *Drosophila* midgut, remains unresolved

(Sousa-Victor et al., 2017, Tang et al., 2023).

We took multiple complementary approaches to understand the mechanisms

of action of Aub in the adult *Drosophila* midgut. Firstly, we tested whether we could

recapitulate the effect of *aub* knockdown on regenerative ISC proliferation by whole

body loss and/or adult ISC/EB specific knockdown of additional components of the

primary pathway (Armitage (Armi); Rhino (Rhi)) or the Ping-Pong pathway (Vasa;

Spindle-E (SpnE)) (**Fig. S1D, E**). As in the case of Ago3 (**Fig. 1C, D** and **Fig. S1A**),

we detected no effect on intestinal regeneration upon loss *spnE* (**Fig.S1D).** However,

whole body loss or ISC/EB knockdown of the other genes assessed led to a

significant decrease in ISC proliferation upon intestinal damage (**Fig.S1D, E**). Given

the role of complementary PIWI pathway members regulating each other's stability in

the germline (Senti et al., 2015, Wang et al., 2015b), we tested whether *piwi* or *vasa*

downregulation within ISCs/EBs might impact Aub stability and therefore explain

their effect on ISC proliferation through indirect Aub regulation. However, in neither

case did we see a change in the upregulation of Aub in regenerating midguts

(**Fig.S1F-I**). These data suggest that *piwi* and *vasa* act independently from Aub in

the midgut to regulate intestinal regeneration.

Next, we performed small RNA sequencing of oxidized samples of adult

whole midguts from wild type and *aub* mutant flies to assess the overall presence of

piRNAs in this tissue (**Fig.S2A**). Importantly, we took advantage of the 2′-*O*-

methylation present at the 3′ ends of piRNAs that confers resistance to sodium

periodate oxidation (Kirino and Mourelatos, 2007, Ohara et al., 2007, Saito et al.,

2007, Houwing et al., 2007). Consequently, RNA samples were enriched with

piRNAs versus other, and potentially more abundantly present, small RNAs

populations. Although much less prominent than in oxidized libraries from ovaries

(**Fig.S3A**), oxidation of gut libraries revealed an enrichment of 23-28 nucleotide long

TE antisense small RNAs (**Fig.S2B**). Importantly, those RNAs showed nucleotide

biases characteristic of piRNAs: Uridines at the 1st base position of TE antisense

reads (AS), and to a lesser extent Adenines at the 10th base position of TE sense

reads (AS), and to a lesser extent Adenines at the 10^{th} base position of TE sense

reads (S). Notably, this enrichment was not visible in previously published

unoxidized libraries (Siudeja et al., 2021)(**Fig.S3B**), emphasising the importance of

library oxidation to observe small enrichments in piRNAs. Interestingly, the

abundance of AS piRNAs observed in whole midguts appeared overall diminished in

aub mutant tissues (**Fig.S2B, B' versus Fig.S2C, C'** and **Fig.S2D, E**). While these

data may suggest an involvement of Aub in the regulation of piRNAs in the adult

midgut, whole midgut samples contain very heterogeneous cell populations,

including gut-associated-trachea and visceral muscle, in addition to the multiple

cellular subtypes present within the midgut epithelium itself. This cell heterogeneity

often underestimates molecular information from cells underrepresented in the

midgut and it is insufficient to address cell specific gene function. To better

determine if our results in whole adult midguts are reflective of the presence of

piRNAs and/or a piRNA regulatory function of Aub in ISCs, we next conducted small

RNA sequencing of oxidized libraries from sorted control or *aub-RNAi* expressing

ISCs/EBs (**Fig.2A**). Compared to whole midgut samples, small RNA sequencing

from ISCs/EBs of control and regenerating midguts showed a stronger piRNA

signature (**Fig.2B** versus **Fig.S2B**). However, piRNA abundance in ISCs/EBs was

not reduced by cell specific knockdown of *aub* (**Fig.2B, B'** versus **Fig.2C, C'**). If

anything, we observed an unexplained trend towards higher piRNA counts in

ISCs/EBs from *aub-RNAi* midguts (**Fig.2D, E**)**.**

The tiles analysis of genome-unique piRNA mappers in ISCs/EBs versus

those in ovaries showed that piRNA germline-expressed clusters 38C and 42AB are

more abundant in ISCs/EBs than those from the somatic cluster flamenco (**Fig.2F**).

This is consistent across all conditions and genotypes analysed (**Fig.S3C-F** and

Fig.S3G for ovaries). In summary, our results suggest that the overall piRNA

signature seen in ISCs/EBs from adult midguts is distinct from that observed in the

ovaries. However, *aub* downregulation does not impair ISC/EB piRNAs.

Aub regulates intestinal regeneration independently of its piRNA regulatory

function.

To further address the role of Aub in the midgut in relation to its piRNA

regulatory function, we performed mRNA sequencing from control or Aub-depleted

ISCs/EBs to measure the abundance of TE mRNAs in homeostasis (Sucrose) and

upon regeneration (*Pe*) (**Fig.3A, B** and **Fig.S4A, B**). While several TEs, such as
copia, *roo* and *Doc*, were abundantly expressed in ISCs/EBs in both conditions, this
was independent of the presence of Aub (**Fig.3A, B** and **Fig.S4A, B**). However,
these TEs are neither abundantly expressed nor under the control of Aub in the
ovaries (**Fig.S4C**) (Senti et al., 2015). Interestingly, we observed that *flea*
expression, a TE regulated by Aub in the germline (Sienski et al., 2015) (**Fig.S4C**),
was also upregulated in ISCs/EBs (**Fig.3A, B** and **Fig.S4A, B**). However, *flea*
upregulation was largely associated with the presence of *Pe* infection rather than
with Aub expression (**Fig.3A, B** and **Fig.S4A, B**). These data are consistent with
recent work suggesting that stress-associated TE regulation in the midgut may be
carried out by mechanisms independent of the piRNA machinery (Siudeja et al.,
2021). Next, we used site-directed transgenesis to generate fly lines carrying
inducible GFP tagged wild type Aub (*UAS-aub^{WT}*) or mutant isoforms (*UAS-aub^{AA}*
and *UAS-aub^{ADH}*, **Fig.S4D**). *UAS-aub^{AA}* contains a double point mutation in the PAZ
domain, responsible for the loading of piRNAs (Barckmann et al., 2015) while *UAS-*
aub^{ADH} carries a single mutation in the PIWI domain, responsible for Aub's
endonuclease activity and essential for piRNA biogenesis (Wang et al., 2015b).
Mutations in *aub* have been previously correlated with female sterility, a reduced size
of the ovaries and embryonic lethality (Sahin et al., 2016, Schüpbach and
Wieschaus, 1991, Barckmann et al., 2015, Wang et al., 2015b). We first verified the
general functionality of our transgenes by assessing their ability to revert female
sterility in *aub* mutant flies (**Fig.S4E**). Consistent with previous reports (Wang et al.,
2015b) we observed that only *UAS-aub^{WT}* could partially restore fertility when
expressed in the germline (**Fig.S4E**). Next, we used *esg-Gal4* to drive expression of
the different transgenes in the midgut of wild type flies and used anti-GFP and anti-
Aub staining as proxies to measure ISC/EB transgene expression levels and
confirmed comparable transgene expression in the midgut (**Fig.3C-E**). Next, we
expressed *UAS-aub^{WT}*, *UAS-aub^{AA}* or *UAS-aub^{ADH}* in ISCs/EBs from trans-
heterozygote *aub^{HN2/QC42}* flies (Harris and Macdonald, 2001, Schüpbach and
Wieschaus, 1991, Schüpbach and Wieschaus, 1989) using *esg-Gal4* (**Fig.3F-H**).
Aub expression by the different constructs was significantly increased when
compared to *aub^{HN2/QC42}* mutant flies, in both sucrose and *Pe* conditions (**Fig.3F, G**).
Despite the slightly reduced Aub levels upon the expression of *UAS-aub^{ADH}*
compared to the other transgenes, we observed a significant rescue of regenerative

ISC proliferation in all cases when compared to $aub^{HN2/QC42}$ midguts (**Fig.3F-H**).
Altogether, our results suggest that Aub is not necessary to repress TEs or induce
the piRNA signature of ISCs/EBs and that the piRNA regulatory function of Aub is
dispensable for the activation of ISC proliferation in the adult *Drosophila* midgut.

304 Aub regulates ISC proliferation through the translation of core regenerative
305 factors in the adult midgut.

To further characterize the processes underpinning the role of Aub in the
midgut, we next assessed the effect of *aub* knockdown on the activity of signalling
pathways essential for the regenerative capacity of the intestinal epithelium. The
EGFR/MAPK, Wnt and JAK-STAT pathways require ISC activation of their
respective effectors, Sox21a, Myc and Socs36E, to drive intestinal regeneration
(Buchon et al., 2010, Biteau and Jasper, 2011, Xu et al., 2011, Jiang et al., 2011,
Jiang et al., 2009, LinXu and Xi, 2010, Zhou et al., 2013, LinXu and Xi, 2008,
Cordero et al., 2012b). We hypothesized that these transcription factors would
represent good candidates to mediate the exclusive cell autonomous function of Aub
within ISCs (**Fig.1L, M**). In the first instance, we performed RT-qPCR analysis of
homeostatic and regenerating control and *aub* knockdown midguts, which showed
no effect of Aub on the upregulation of *sox21*a (**Fig.S5A**), *myc* (**Fig.S5B**) or *socs*
36e mRNA (**Fig.S5C**) in *Pe* treated midguts.

Well documented biological functions of Aub in the germline and embryo
include control of the translation of mRNAs coding for specific proteins, including
transcription factors (WilsonConnell and Macdonald, 1996, Harris and Macdonald,
2001, Ma et al., 2017, Ramat et al., 2020, Rojas-Ríos et al., 2017). Protein synthesis
is necessary for the adaptive proliferative response of ISCs to changes in diet and
nutrient availability (Obata et al., 2018, Bonfini et al., 2021). Protein translation,
measured by puromycin incorporation (Bonfini et al., 2021, David et al., 2012,
DeliuGhosh and Grewal, 2017), was significantly increased in ISCs of regenerating
midguts (**Fig.4A, C**), and this was dependent on *aub* expression (**Fig.4B, C**). We
therefore tested a possible translational control of *sox21a* and *myc* in ISCs by Aub.
Indeed, damage induced Sox21a (**Fig.4D-F**) and Myc (**Fig.4G-I**) expression was
significantly impaired upon *aub* knockdown. Unfortunately, the lack of available
antibodies precluded us from taking a similar approach for Socs36E. Altogether,
these data suggest that Aub drives regenerative ISC proliferation, at least in part

through cell autonomous upregulation of *sox21a* and *myc* translation and in a
manner that is independent of its association with piRNAs.

The multiprotein complex eIF3 is involved in the early steps of protein
translation where it actively participates in the formation and stabilisation of the pre-
initiation complex but also in the recruitment of mRNAs (Asano et al., 2000, Valásek
et al., 2003, MajumdarBandyopadhyay and Maitra, 2003, Jivotovskaya et al., 2006).
Previously identified roles of Aub in protein translation involves its physical
interaction with components of the early translation machinery, including eIF3 and
eIF4 (Ma et al., 2017, Ramat et al., 2020). Consistently, wild type midguts showed
significant upregulation of a subunit of the eIF3 complex, eIF3C, in ISCs in response
to damage (**Fig.5A, B**). Knocking down different subunits of the eIF3 complex in
ISCs mimicked the effect of *aub* downregulation on puromycin incorporation
(**Fig.S5D, D'**) and on ISC proliferation during midgut regeneration (**Fig.5C**).
Additionally, loss of eIF3C impaired ISC proliferation caused by *aub* overexpression
(**Fig.5D**).

We next tested a potential epistatic relationship between eIF3C and the
expression of Sox21a and Myc in regenerating ISCs. Unexpectedly, *eIF3C*
knockdown did not impact the damage-induced upregulation of either Sox21a or Myc
expression in ISCs (**Fig.5E-J**). Similarly, the loss of Sox21a or Myc did not decrease
eIF3C expression (**Fig.5K** and **Fig.S5E-G**), suggesting that the regulation of Sox21a
and Myc by Aub in ISCs does not involve the eIF3 complex (**Fig.5L**). Further work
will be necessary to fully understand the mechanisms of cell signalling regulation by
Aub in ISCs and identify specific targets of eIF3C in the system.

The human homologue of Aub, *PIWIL1,* is upregulated in colorectal cancer.

Aberrant expression of human *PIWIL1* has been reported in colorectal (Sellitto
et al., 2019, Wang et al., 2015a, Sun et al., 2017) and gastric cancers (Araújo et al.,
2018, Shi et al., 2020) where *in vitro* studies in 2D cell cultures suggest its role in cell
proliferation, cell migration, invasiveness and tumourigenesis. Consistently, our
analysis of *PIWIL1* gene expression in colon tissue samples using the TNMplot tool
(Bartha and Győrffy, 2021) (**Fig.6A-C**) revealed a significantly higher *PIWIL1* gene
expression in colon adenocarcinoma versus normal non-paired tissues (**Fig.6A**) as
well as versus adjacent normal tissues (**Fig.6B**). Chip-seq data analysis with the
TNMplot tool also revealed strong overexpression of *PIWIL1* in metastatic tissues

(**Fig.6C**). We next took advantage of the publicly available transcriptomic data from
the Cancer Genome Atlas Program (TCGA, PanCancer Atlas) and analysed *PIWIL1*
expression across different CRC subtypes (CMS1-4) as classified by (Guinney et al.,
2015) (**Fig.6D**). Interestingly, *PIWIL1 gene* expression was significantly upregulated
in the CMS1 human CRC subtype (**Fig.6D**), which presents high microsatellite
instability correlated with hypermethylated profiles and a strong immune activation
(Guinney et al., 2015). Consistently, we observed significant *PIWIL1* upregulation in
patients with tumours characterised by high microsatellite instability (**Fig.6E**). Next,
we assessed levels of PIWIL1 in tissue microarrays from a local cohort of CRC
patients who had undergone elective and potentially curative resection of stage I-III
colorectal adenocarcinoma in a single surgical unit at the Glasgow Royal Infirmary,
Western Infirmary or Stobhill Hospitals (Glasgow, UK) between 1997 and 2007. First,
we confirmed PIWIL1 antibody specificity in testis from wild type and *Piwil1* mutant
mice (**Fig.S6A**). In line with previous reports, we observed strong expression of
PIWIL1 in multiple CRC patient samples (**Fig.6F**) and a significant correlation
between PIWIL1 levels and poor patient survival (**Fig.6G**). Combined, these results
show a strong correlation between high PIWIL expression levels and CRC with
potential selectivity towards tumour subtypes.

Aub regulates Wnt and MAPK signalling driven intestinal hyperplasia *in vivo*.

Current knowledge on the role of PIWI in intestinal cancer is largely based on
tissue culture work. Oncogenic intestinal hyperplasia can be recapitulated in the
adult *Drosophila* intestine *in vivo* through stem cell specific oncogene activation or
tumour suppressor knockdown (Lee et al., 2009, Bangi et al., 2016, Jiang et al.,
2011, Cordero et al., 2012a). Our previous results in the regenerating midgut
suggest that Aub is functionally linked to the Wnt pathway through Myc (**Fig.4G-I**).
We therefore used genetically defined fly models to address the *in vivo* role of PIWI
proteins in CRC, with a focus on Aub. We first looked at Aub expression in intestinal
hyperplasia driven by hyperactivation of Wnt signalling activity, a major driver of
early intestinal tumourigenesis, including hereditary familial adenomatous polyposis
(FAP) (Segditsas and Tomlinson, 2006). Immunofluorescence staining for Aub
showed a significant increase of protein expression in ISCs/EBs from adult midguts
where Wnt signalling is hyperactivated upon loss of *Adenomatous polyposis Coli*
(*Apc1)* (Lee et al., 2009, Cordero et al., 2012a) (**Fig.7A-C**). We next used the

MARCM system (Lee and Luo, 1999) to induce 10 day-old control (*LacZ*) or *Apc1*
mutant clones in the presence or absence of *aub* knockdown (*aub^{RNAi}*) and
quantified tumourigenic ISC proliferation by measuring the size of the clones
(**Fig.7D-E**). As shown before (**Fig.S1B-C**), knocking down *aub* was not sufficient to
impact homeostatic clonal growth (**Fig.7D-E**). In line with our previous report
(Cordero et al., 2012a), loss of functional *Apc1* induced a significant increase of
clonal size (**Fig.7D-E**). Combined loss of functional *Apc1* and *aub* knockdown led to
a complete suppression of *Apc1* clonal growth (**Fig.7D-E**) suggesting that Aub
mediates intestinal tumourigenesis by *Apc1*. Activation of Wnt signalling through the
upregulation of the Wnt ligand Wingless (Wg) is necessary and sufficient to drive ISC
proliferation in adult midguts (Cordero et al., 2012b, Cordero et al., 2012a, LinXu and
Xi, 2008). Consistently, knocking down *aub* suppressed ISC proliferation induced by
the overexpression of Wg in adult midgut ISCs/EBS (**Fig.7F**).

Our previous analysis pointed to a selective upregulation of *PIWIL1* in CMS1
CRC tumour subtypes (**Fig.6D**). In addition to high microsatellite instability and
strong immune activation, CMS1 tumours are characterized by the presence of
activating mutations in the EGFR/MAPK signalling component BRAF (Guinney et al.,
2015). The functional association between Aub and Sox21a is consistent with a link
between Aub and MAPK signalling in intestinal regeneration (Meng and Biteau,
2015)(**Fig.4D-F**). We therefore used MAPK signalling driven fly models of intestinal
hyperplasia to address the functional role of Aub in such genetic tumour settings.
Overexpression of a constitutively active form of human BRAF (*Raf^{GOF}*) in ISCs/EBs
drives robust ISC hyperproliferation in the adult *Drosophila* midgut (**Fig.7G, H, J**)
(Brand and Perrimon, 1994, Markstein et al., 2014) and high levels of Aub (**Fig.7H,**
I). However, concomitant short-term (2 days) knockdown of *aub* by RNAi in these
midguts was insufficient to affect hyperplasia caused by *Raf^{GOF}* (**Fig.7J**). This is likely
due to the rapid and very high ISC hyperproliferation induced by *Raf^{GOF}*, and the
diminished RNAi efficiency to knock down *aub,* as evidenced by the only partial
impact of similar gene knockdown timelines on regenerative ISC proliferation
(**Fig.S6B**). We then tried longer periods of transgene co-expression, which allowed
for more effective *aub-RNAi* function (**Fig.S6C**). Longer periods (3 days) of *Raf^{GOF}*
overexpression had detrimental effects on gut integrity, including decreased ISC
proliferation, likely due to high apoptosis (**Fig.7K, L and Fig.S6D**). Unexpectedly, co-
expression of *aub RNAi* in such context maintained high levels of ISC proliferation

and impaired apoptosis in Raf^{GOF} overexpressing midguts (**Fig.7K, L and Fig.S6D**).

We observed a similar phenomenon when knocking down *aub* in midguts with

sustained *Ras* hyperactivation (**Fig.S6E-G**). Of note, while we could induce *Ras*

hyperactivation in midgut ISCs/EBs for longer periods of time (7 days), this was not

possible in the case of Raf^{GOF} due to catastrophic tissue disruption.

Altogether, our data suggest a context-dependent role of Aub in

tumourigenesis, which may be dictated by the genetic characteristics of the tumour.

Importantly, we provide valuable paradigms for broader studies on the role of PIWI

proteins and piRNAs in intestinal physiology and tumourigenesis.

Discussion

Multiple lines of evidence have demonstrated roles of PIWI proteins beyond their

well-defined function as repressors of TEs expression, including post-transcriptional

control of gene expression in the germline and embryo (Barckmann et al., 2015,

Dufourt et al., 2017, Ramat et al., 2020, Becalska et al., 2011, WilsonConnell and

Macdonald, 1996, Cook et al., 2004, Ma et al., 2017, Rojas-Ríos et al., 2017, Harris

and Macdonald, 2001). Here, we discover a new role of Aub in the adult midgut,

which is uncoupled of its piRNA regulatory function and involves translational

regulation of conserved pro-regenerative transcription factors within intestinal stem

cells.

Aub acts as a regulator of protein translation in ISCs.

Our work underscores a central role of inducible protein synthesis in the

capacity of stem cells to trigger the regenerative response of the intestine following

damage **(Figs.4 and Fig.5)**. This includes increased Aub production itself **(Fig.1)**

and the upregulation of a core component of the translation initiation machinery eIF3

(Fig.5). Physical interaction with translation initiation factors mediates the role of Aub

in post-transcriptional gene regulation through binding to specific mRNAs in the

germline and ovaries (Ma et al., 2017, Ramat et al., 2020). Unexpectedly, direct

interaction between Aub and eIF3 seems unlikely in midgut ISCs, as suggested by

the distinct regulation of Sox21a and Myc in ISCs by Aub but not eIF3c (**Figs.4** and

5). Our results therefore reveal not only a role Aub in protein translation beyond the

mainstream pathway involving eIF3, but also the existence of, yet to be identified,

eIF3 target mRNAs within ISCs, which are essential for intestinal regeneration.

Differential requirement of translation initiation factors and target specific mRNA

translation by the eIF3 complex regulate developmental processes in *Drosophila* and

mice (Fujii et al., 2021, Rode et al., 2018). Further studies will be needed to

determine by which mechanisms Aub regulates Sox21 and Myc in ISCs. Regulation

of germline stem cell self-renewal and differentiation by Aub has also been linked to

physical interaction with subunits of the eIF4 translation initiation complex (Ma et al.,

2017). It is therefore possible that a similar functional relationship exists between

Aub and eIF4 in ISCs. However, the reduced abundance of midgut piRNAs relative

to the germline, and current lack of *in vitro* cultured systems to increase tissue mass,
complicates the implementation of classical molecular and biochemical approaches
used in the PIWI/piRNA field for characterization of protein/protein and protein/RNA
interactions in ISCs.

What drives the observed upregulation of Aub in ISCs **(Fig.1)** is unknown.
Intriguingly, analysis of protein/mRNA complexes in the germline revealed Aub
binding to its own mRNA (Ma et al., 2017). While we don't find significant global
changes in *aub* mRNA in the midgut **(Fig.1)**, it is worth considering the possible
existence of an auto-regulatory mechanism of Aub regulating its own production in
ISCs. Additionally, stress signals commonly involved in driving regenerative and
tumourigenic ISC proliferation could be playing a role in Aub induction.

Aubergine functions independently of piRNAs in ISCs of the adult *Drosophila*
midgut.

While multiple studies have demonstrated dependency on piRNA binding for
broad functions of Aub, including regulation of mRNA localisation, stability and
translation (Barckmann et al., 2015, Dufourt et al., 2017, Rojas-Ríos et al., 2017,
Rouget et al., 2010, Ramat et al., 2020, Vourekas et al., 2016, Rojas-Ríos et al.,
2023), piRNA-independent roles of PIWI proteins are only recently emerging (Ma et
al., 2017, Shi et al., 2020, Li et al., 2020, Tang et al., 2023). Our experiments
suggest that the role of Aub in adult *Drosophila* ISC proliferation does not involve its
piRNA regulatory function (**Fig.3**). Additionally, loss of function of *ago3*, an essential
ping-pong partner of Aub in germline piRNAs biogenesis, did not impact ISC
proliferation (**Figs.1** and **Fig.S1**). On the other hand, we showed that the ping-pong
pathway component Vasa is required to drive ISC proliferation during midgut
regeneration (**Fig.S1**). However, biological functions of Vasa extend beyond its role
in the piRNA pathway, including the regulation of cell cycle genes (Yajima and
Wessel, 2011), which could potentially explain its role in *Drosophila* ISC proliferation.
Alternatively, but less likely, Vasa may physically interact with Aub to regulate ISC
proliferation without the need of piRNA binding. Nonetheless, given that Aub is
considered essential for ping-pong piRNAs biogenesis in the germline, our results
suggesting its dispensability for the ping-pong piRNAs observed in ISCs was
unexpected **(Fig.2)**. This could be explained by either a sufficiency to fulfil this role
with residual levels of Aub expected from RNAi partial knockdown, or by a non-

canonical Piwi and Ago3 dependent regulation of ISC ping-pong piRNAs in the absence of Aub.

Our results do not exclude a general role of piRNAs in ISCs. Previous reports highlighted the functional role of Piwi in ISC maintenance and TE regulation in ageing intestines (Sousa-Victor et al., 2017). Here, we report the presence of piRNAs in ISCs/EBs and a decrease of regenerative ISC proliferation upon the knockdown of Piwi and other components of the primary piRNA pathway (**Figs.1 and Fig.S1**). Further studies will be needed to directly address the role of PIWI proteins and piRNA biogenesis within ISCs.

Unlike in sorted ISCs/EBs (**Fig.2**), small RNA sequencing of whole midguts shows an overall decrease in the global piRNA signature of the midgut from *aub* mutant animals (**Fig.S2**). Independent reports of *aub* expression in the midgut include enteroendocrine cell progenitors (Korzelius et al., 2019). These results suggest the interesting prospect of a piRNA-dependent regulatory role of Aub in the midgut epithelium or associated tissues.

Aubergine and PIWI proteins in colorectal cancer.

Our initial observations suggest that, while the upregulation of Aub is a common feature of regenerative and oncogenic ISC hyperproliferation, the functional role of PIWI proteins in tumourigenesis is complex and likely context dependent **(Fig.7)**.

Alterations of the protein translation machinery is a cancer hallmark (Song et al., 2021). Over 95% of human CRC cases bear loss of function mutations in the tumour suppressor *APC,* leading to Wnt pathway hyperactivation and the subsequent upregulation of its target gene *myc* (SchatoffLeach and Dow, 2017). Myc expression in CRC relies on the activity of the eukaryotic initiation factor eIF4E (Wiegering et al., 2015, Knight et al., 2021) and eIF4A1 (Waldron et al., 2023). It is therefore possible that the mechanisms mediating the pro-proliferative function of Aub in intestinal regeneration **(Figs.4 and 5)** may extrapolate to pro-oncogenic roles of PIWI proteins in tumours with hyperactivating mutations in Wnt signalling components **(Fig.7)**.

On the other hand, upregulation of Aub upon MAPK signalling hyperactivation revealed a pro-apoptotic/tumour suppressive role of Aub in such context **(Fig.7),** which may predict a similar functional outcome for upregulated *PIWIL1* in CMS1

tumour subtypes containing BRAF hyperactivation **(Fig.6)** (Guinney et al., 2015)**.**
Anti- and pro-apoptotic roles have been described for PIWIL1 in tumour stem cells,
including glioma and leukaemia (Wang et al., 2015d, Huang et al., 2021). Our results
may therefore reveal general mechanisms mediating dual and antagonistic functions
of PIWI proteins in cancer.

Additional features of CMS1 tumours include epigenetic modifications, such
as genome hypermethylation (Guinney et al., 2015). PIWI proteins induce
transposon silencing partly through DNA hypermethylation, a function that might be
repurposed in oncogenic settings through the hypermethylation of tumour gene
promoters (WajedLaird and DeMeester, 2001). Previous studies have linked PIWI
proteins to DNA hypermethylation in tumourigenesis and cell invasion in sarcoma
and melanoma, respectively (SiddiqiTerry and Matushansky, 2012, Wang et al.,
2015c). PIWIL1 expression also correlates with promoter methylation of the tumour
suppressor gene PTEN in endometrial cancer (Chen et al., 2015). Further *in vivo*
studies in genetically defined tumour models would provide invaluable insights into
the role of PIWI proteins and piRNAs in colorectal cancer and our work presents
excellent paradigms to achieve this.

Material and Methods

A complete list of fly stocks, full genotypes, and reagents used in this study
can be found in Supplementary Table #1.

***Drosophila melanogaster* stock rearing and maintenance.**
Fly stocks were kept in temperature and humidity-controlled incubators set up with a
12h-12h light/dark cycle. For experiments with *aub* mutant lines, crosses were
performed at 25°C and the desired F1 progeny was sorted 2/3 days after adult
eclosion and kept at 25°C for 5-7 days before dissection. For experiments using the
GAL4/*GAL80*^ts^ system, flies were crossed at 18°C. The desired F1 progeny was
sorted 2/3 days after adult eclosion and was kept at 18°C for 5 days before being
transferred to 29°C to allow transgene activation. The incubation time at 29°C varied
from 2 days to 10 days depending on the transgene. In all cases, experimental flies
were flipped every two days on fresh food. Only mated females were used for
experiments.

Human patient information and ethics statement.
The human cohort analysed contains 1030 colorectal cancer (CRC) patients who
had undergone elective and potentially curative resection of stage I-III colorectal
adenocarcinoma in a single surgical unit at the Glasgow Royal Infirmary, Western
Infirmary or Stobhill Hospitals (Glasgow, UK) between 1997 and 2007. Resection
was considered curative based on pre-operative computed tomography and intra-
operative findings. The data are stored within the Glasgow Safehaven
(GSH21ON009) and ethical approval was in place for the study (MREC/01/0/36).

Damage-induced intestinal regeneration.
To induce intestinal regeneration, mated female flies were fed on filters (Whatman)
soaked in 5% sucrose only (control condition) or with the pathogenic bacteria
Pseudomonas entomophila (*Pe*) at OD_{600}=25 (damage condition) for 16 hours prior
to dissection. The bacterial solution was obtained after overnight culture in LB
medium at 29°C under 220 rpm shaking overnight, after which bacteria were pelleted

(Beckman Coulter JS-4.2 rotor, 10 min, 4000 rpm) and diluted in a solution of 5%

sucrose to reach OD_{600}=25.

Immunohistochemistry.

Drosophila tissue: The dissection of guts was performed in PBS at room temperature

and 16 hours after feeding the mated female flies with sucrose or _Pe_. After

dissection, guts were immediately fixed in a solution of 4% paraformaldehyde

(Polysciences Inc) for 1 hour. Guts were then washed in PBST (PBS with Triton

0.2%) before incubation overnight at 4°C with the primary antibodies diluted in

blocking solution PBT (with 0.5% BSA). The day after, guts were washed 3x15

minutes with PBST at room temperature before incubation with the secondary

antibodies for 2 hours at room temperature. Guts were then washed 3x15 minutes

with PBST at room temperature before mounting in DAPI containing mounting

medium. Slides were kept at 4°C before imaging. To improve Aub, Dcp-1, eIF3C and

Myc staining, guts were placed in methanol for 10 minutes at room temperature after

the fixation step with 4% paraformaldehyde.

Human and mouse tissue: Immunohistochemistry (IHC) staining was performed on a

previously constructed human tissue microarray or on 4µm formalin fixed paraffin

embedded sections, which had previously been incubated in a 60□C oven for 2

hours. Tissue slides for PIWIL1 detection were stained on a Leica Bond Rx

Automated Stainer. Samples underwent on-board dewaxing and epitope retrieval

using ER2 solution (Leica) for 20 minutes at 95°C. Sections were rinsed with Leica

wash buffer before peroxidase blocking was performed using an Intense R kit (Leica)

for 5 minutes. Sections were rinsed with wash buffer (Leica) before anti-PIWIL1

application at an optimal dilution (1/200). The sections were then rinsed with wash

buffer (Agilent) for 30 minutes, before secondary antibody application. The sections

were rinsed with wash buffer, visualized using DAB and counterstained with

Haematoxylin in the Intense R kit. To complete the IHC staining sections were rinsed

in tap water, dehydrated through a series of graded alcohols, and placed in xylene.

The stained sections were cover slipped in xylene using DPX mounting medium.

Details of antibodies sources and concentrations used as well as reagent'

information can be found in Supplementary Table #1.

Generation of _aub_ transgenes.

The three pUASt-attB-GFP constructs of *aub* were generated for this study. A
complementary DNA (cDNA) of *aub* was amplified from *w^1118* flies and cloned into
the pUASt-attB plasmid, that was originally made by the Basler group (GenBank
accession EF362409, PUBMED: 17360644) and received as a gift from Julius
Brennecke. A GFP cDNA was inserted N-terminally to *aub* cDNA. Standard site-
directed mutagenesis by Phusion DNA Polymerase (Thermo Fisher) was used to
introduce amino acid substitutions to generate *aub^AA* (Y345A/Y346A, (Barckmann et
al., 2015)) and *aub^ADH* (D636A, (Wang et al., 2015b)). Plasmid construction and
sequences are available at https://github.com/RippeiHayashi/gut_aubergine.
641

Egg laying/eclosion assay.
Crosses to obtain experimental animals were kept at 18□C. Virgin females of
the desired genotype were selected and placed into fresh vials and allowed to mate
with *w^1118* males. A maximum of 5 females and 5 males were housed in each vial
and allowed to mate over a period of 24 hours at 18□C. The following day, males
were removed, and females were moved onto fresh food and incubated at 25□C.
Females were flipped into new vials every 24 hours. The number of eggs in each vial
was counted and recorded at the same time. This process was repeated 3 more
times. Importantly, the number of live females was recorded each time, at the time of
flipping. Any females that died during the night were included in the eggs/adult ratio
but were excluded from the subsequent recordings. This experiment was repeated
three times.
654

Protein translation assay.
For general protein translation assessment, flies were fed on filters (Whatman)
soaked in 5% sucrose only or with *Pe* OD_{600}=25 16 hours prior to gut dissection in
Schneider's insect medium (Sigma-Aldrich) containing puromycin (stock solution 25
mg/mL, Sigma-Aldrich) with a final concentration of 25µg (1µL of stock solution) of
puromycin in 5mL of Schneider's insect medium. During dissection, Malpighian
tubules and crop were kept attached to the gut and precaution was taken to keep the
intestines intact. Guts were then incubated at room temperature in puromycin
containing medium for 45 minutes. After incubation, we applied the
immunochemistry protocols we described in this manuscript.
665

Clonal analysis.

Crosses were performed at 25°C and parental lines were flipped twice a week. Clones were induced using the MARCM technique (Lee and Luo, 1999). The desired progeny was collected and aged for 3 to 5 days at 25°C before undergoing three 30 minutes heat shocks in one day, in a 37°C water bath. Heat shocked adults were then aged for 10 days at 25°C before dissection.

Fluorescence-activated single cell sorting (FACS).

Control flies and flies expressing RNAi against *aub* were collected between 2 to 3 days after hatching and kept at 18°C for 5 days before being transferred to 29°C. Flies were kept at 29°C for 7 days and were flipped into fresh food vials every two days. Fly guts were dissected in cold PBS 16 hours after animals were fed with 5% sucrose only or 5% sucrose containing *Pe*. The crop, the Malpighian tubules and the hindgut were carefully removed. Midguts were then collected and transferred to an Eppendorf containing 400 µL of PBS and kept on ice. Each Eppendorf contained roughly 100 midguts. Once all the tissue was dissected, 10 µL of elastase (Sigma-Aldrich, 10 µg/ µL) was added to each Eppendorf. Samples were kept at 27°C in a heat block for at least 1 hour until midguts were fully dissociated. Tissues were further disrupted by pipetting up and down every 5 minutes. Samples were then centrifuged at 300 rcf for 20 minutes at 4°C. After removing the supernatant, the pellet was resuspended in cold PBS. To eliminate enterocytes as much as possible, the cell suspension was filtered with a 30 µm filter (Sysmex). 500 µL of cold PBS were added to wash the filter. After filtration, samples were ready for sorting. Sorting was performed by the CRUK Scotland Institute flow cytometry facility on a FACSAria Fusion Flow Cytometer (BD Biosciences). GFP or Red stinger positive controls were used to set up the flow cytometer parameters that were used for all the conditions. In parallel to the sorting, a few microliters of samples were taken and stained with 20 µL of DAPI to visualize cell death. Between 250 and 500 flies were dissected per condition, resulting in 250,000 to 750,000 sorted cells. Three biological replicates were dissected for each condition. Sorted cells were pelleted by centrifugation at 300 rcf for 20 minutes at 4°C. After carefully removing the supernatant, 800 µL of TRIzol (Invitrogen) was added to the samples, followed by RNA extraction and sequencing.

Small RNA sequencing.

Small RNA sequencing was performed on either whole midguts, ovaries or sorted
ISCs/EBs. For small RNA preparation from whole midguts, flies were dissected in
cold PBS 16 hours after being fed with 5% sucrose only or 5% sucrose containing
Pe. The crop, the Malpighian tubules and the hindgut were carefully removed. Three
replicates of 40 midguts each were dissected for each condition. Guts were then
collected and transferred to an Eppendorf containing 400 µL of PBS and kept on ice.
Once all the tissue was dissected, 10 µL of elastase (Sigma-Aldrich, 10 µg/µL) was
added to each Eppendorf. Samples were kept at 27°C in a heat block for at least 1
hour until the guts were totally degraded. Tissues were disrupted by pipetting up and
down every 5 minutes. Samples were then centrifuged at 300 rcf for 20 minutes at
4°C. After removing the supernatant, the pellet was resuspended in 800 µL of TRIzol
(Invitrogen). The same protocol was applied for the ovaries. Approximately 10 pairs
of ovaries were dissected. We generated small RNA libraries from 1 ~ 5 µg of total
RNA using a modified protocol from the original method (Hafner et al., 2008). 19 to
35 nucleotides-long RNA was first selected by polyacrylamide gel electrophoresis
(PAGE) with 7M Ureal using radio-labelled 19mer (5''-
CGUACGCGGGUUUAAACGA) and 35mer spikes (5''-
CUCAUCUUGGUCGUACGCGGAAUAGUUUAAACUGU). The size-selected RNA
was precipitated, oxidised by sodium periodate (Sigma)(Akbergenov et al., 2006),
and size-selected for the second time by PAGE. The size-selected oxidised small
RNAs were then ligated to the 3' adapter from IDT
(5rApp/NNNNAGATCGGAAGAGCACACGTCT/3ddC where Ns are randomised)
using the truncated T4 RNA Ligase 2, truncated KQ (NEB), followed by a third PAGE
to remove non-ligated adapters. Subsequently, the RNA was ligated to the 5' adapter
from IDT (ACACUCUUUCCCUACACGACGCUCUUCCGAUCUNNNN where Ns are
randomised) using the T4 RNA Ligase 1 (NEB). Adapter-ligated RNA was reverse-
transcribed using SuperScript II (Thermo Fisher) and amplified by KAPA LongRange
DNA polymerase (Sigma, KK3502) using the universal forward primer, Solexa_PCR-
fw: (5'-
AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCG
ATCT) and the barcode-containing reverse primer TruSeq_IDX: (5'-
CAAGCAGAAGACGGCATACGAGATyyyyyyGTGACTGGAGTTCAGACGTGTGCTC
TTCCGATCT where yyyyyy is the reverse-complemented barcode sequence).

Amplified libraries were multiplexed and sequenced on a HiSeq platform in the
paired-end 150 bp mode by GENEWIZ/Azenta.

Small RNA sequencing analysis.

The R1 sequencing reads were trimmed of the Illumina-adapter sequence using the
FASTX-Toolkit from the Hannon Laboratory (CRUK Cambridge Institute). The 4
random nucleotides at either ends of the read were further removed. The trimmed
reads of 18 to 40 nt in size were first mapped to the infrastructural RNAs, including
ribosomal RNAs, small nucleolar RNAs (snRNAs), small nuclear RNAs (snoRNAs),
microRNAs, and transfer RNAs (tRNAs) using Bowtie 1.2.3 allowing up to one
mismatch. Sequences annotated in the dm6 r6.31 assembly of the *Drosophila*
melanogaster genome were used. The trimmed and unfiltered reads were then
mapped to the dm6 genome using Bowtie allowing up to one mismatch. Reads that
uniquely mapped to the genome were analysed for the tile analysis. Reads that
mapped to the 100 nt upstream and downstream genomic regions of tRNA, snRNA
and snoRNA insertions as well as those that mapped to the *aubergine* gene locus
were removed from the tile analysis. Bedtools 2.28.0 was used to count the
coverage of the mapped reads. Endogenous siRNA reads (annotated as hpRNA)
were used for normalisation. The trimmed and unfiltered reads were separately
mapped to the curated sequences of *Drosophila melanogaster* transposons (Senti et
al., 2015) using Bowtie allowing up to three mismatches with the option of --all --best
--strata. The size distribution of the transposon-mapping reads was made using
ggplot2 3.4.4 in R 4.0.0. Nucleotide frequencies around the first nucleotide position
of antisense and the tenth nucleotide position of the sense transposon-mapping
piRNA reads (longer than 22 nucleotides) were counted and visualised using
weblogo 3.7.8. Frequencies were measured in the window of 11 nucleotides and the
z scores were calculated as the deviation of the frequency value at the first
(antisense reads) and tenth (sense reads) positions from the mean frequency
divided by the standard deviation of the frequencies. Small RNA sequencing libraries
from Siudeja et al. (Siudeja et al., 2021) were analysed in the same way except for
using the adapter sequence TGGAATTCTCGGGTGCCAAG, and not trimming the 4
nucleotides at either end.

PolyA-selected RNA sequencing.

Polyadenylated RNA was purified from the DNase-treated total RNA using oligo
d(T)25 magnetic beads (NEB, S1419), and then used for the library preparation.
Libraries were cloned using the NEBNext Ultra Directional II RNA Library Prep Kit for
Illumina (NEB, E7760), following the manufacturer's instruction, and amplified by
KAPA polymerase using the same primers as for the small RNA sequencing.
772

PolyA-selected RNA analysis.
Both R1 and R2 reads from the polyA-selected RNA sequencing reads were trimmed
of the Illumina-adapter sequences using the FASTX-Toolkit. The trimmed reads were
subsequently filtered by the sequencing quality. Only the paired and unfiltered reads
were then mapped to the dm6 r6.31 transcriptome combined with curated sequences
of *Drosophila melanogaster* transposons using salmon/1.1.0 with the options of --
validateMappings --incompatPrior 0.0 --seqBias --gcBias. Length-normalised
transposon mRNA reads per one million transcripts including host mRNAs were
compared between different libraries. polyA-selected RNA sequencing libraries from
ovarian samples from Senti et al. (Senti et al., 2015) were also analysed in the same
way.
784

***Drosophila* RNA extraction and RT-qPCR.**
Total RNA was extracted from tissues dissected in cold PBS. Three or four replicates
of 20 to 40 guts were dissected for each condition. Depending on the experiments,
whole guts or only R4-R5 regions were dissected. Ovaries were also dissected in
parallel as a positive control. Tissues were homogenised in 100 µl of Trizol (Thermo
Fischer Scientific) using a mortar and a pestle first, followed by adding further 700 µl
of Trizol. 200 µl of Chloroform was added and mixed well with Trizol by vortexing
before centrifugation on a bench top centrifuge. The aqueous phase was saved, and
RNA was precipitated with iso-propanol. RNA was treated with DNase (Thermo
Fischer Scientific) and quantified using a NanoDrop Spectrophotometer. About 1 µg
of RNA was converted to cDNA using a high-capacity cDNA reverse transcription kit
(Thermo Fisher Scientific). RT-qPCR was performed using Perfecta SYBR green
fast mix (Quantabio) as per the manufacturer's instructions on an Applied
Biosystems QuantStudio 3 fast real-time PCR system and each reaction was carried
out in triplicate. A standard curve was created using 1:10 dilutions from pooled cDNA
samples. The amount of sample was extrapolated from this standard curve and

normalised using data from the housekeeping gene *rpl32*. Melting curves were
generated previously to ensure only one product was formed from each pair of
primers. Data were exported and analysed in Excel. The results are shown as the
ratio of average of mRNA levels of the candidate gene/*rpl32*.

TCGA analysis.
To study *PIWIL1* RNA expression across the different CMS of CRC or according to
the microsatellite status of the patients, we took advantage of the TCGA datasets
from the PanCancer Atlas. Analysis has been performed on open access data from
colonic adenocarcinoma patients only. Differential gene expression was then
assessed, and statistics have been performed using Graphpad.

Sample visualization and image acquisition.
PH3-positive cells counting was performed using a BX51 Olympus. Fluorescence
images of *Drosophila* intestines were acquired using a confocal Zeiss 710 or a Zeiss
LSM 780. Stained TMAs slides were scanned and analysed using the HALO system
(indica labs).

TMA Scoring.
Scoring of PIWIL1 expression in the cytoplasm was independently performed by two
examiners (K.B and L.R.C) blinded to clinical data. PIWIL1 score was individually
determined in the cytoplasm using the equation: 0 × (% cells not stained) + 1 × (%
cells weakly stained) + 2 × (% cells moderately stained) + 3 × (% cells strongly
stained). This method gave a range of scores from 0 to 300. Damaged samples and
samples with tumour tissue representing less than 10% of the total tissue were
excluded.

Quantification and statistics.
Quantification of ISC proliferation and apoptosis was done by manually counting the
number of PH3-positive cells per posterior midguts and by manually counting the
number of Dcp-1-positive cells in the lumen per image, respectively. Quantification of
eIF3C, Myc, puromycin and Sox21a staining was performed by measuring the mean
fluorescence intensity of the protein in all GFP positive cells on a SUM projection
realised via Fiji (ImageJ 2.14.0/1.54f). Prior to the measurement, a mask was

created via Fiji to delineate the GFP positive cells only, using the tool "Create a
selection". In parallel, mean fluorescence intensity of the background was measured
on a specific region of interest on each SUM projection and subtracted from the
signal measured in the cells.

Quantification of Aub staining was performed by measuring the mean fluorescence
intensity per cell on a SUM projection done by Fiji. This measure was blindly
performed in three different GFP positive cells per gut for Aub. In parallel, mean
fluorescence intensity of the background was measured on a specific region of
interest on each SUM projection and subtracted from the signal measured in the cell.
This quantification was repeated in 5 different guts per biological replicate. A
minimum of 3 biological replicates per condition were quantified.

In the rescue experiments (aub^{WT}, aub^{AA} and aub^{ADH}), GFP and Aub stainings were
quantified by measuring the mean fluorescence intensity of the protein in all GFP
positive cells on a MAX projection realised via Fiji. Prior to the measurement, a mask
was created on Fiji to delineate the GFP positive cells only via using the tool "Create
a selection". In parallel, mean fluorescence intensity of the background was
measured on a specific region of interest on each SUM projection and subtracted
from the signal measured in the cells.

Quantification of ISC proliferation in MARCM clones was performed by manually
counting the number of DAPI positive cells in each GFP-positive clone. The statistics
were done with GraphPad Prism 6. The tests used for each experiment are
described in the figure legends.

For TMAs analysis, the cut off value for high and low expression of PIWIL1 was
determined using the Survminer package in RStudio software version 1.4. Patients
were excluded from analysis if mortality occurred within 30 days post-surgery, upon
presence of metastatic stage disease, and/or if subject to administration of
neoadjuvant therapy. Kaplan Meier survival analysis and univariate cox regression
analysis were performed using SPSS version 8 (IBM) to determine the association
between expression of PIWIL1 and cancer-specific survival.

Acknowledgements

We would like to thank Allison Bardin; Benoit Bîteau; Julius Brennecke; Angela Giangrande; Ginés Morata; Yulii Shidlovskii and Phillip Zamore for generously sharing flies and reagents. We thank the Bloomington Drosophila Stock Center; the Vienna Drosophila Resource Center and the *Drosophila* Studies Hybridoma Bank for fly stocks and antibodies. We would like to thank the Core Services and Advanced Technologies at the CRUK Scotland Institute, which is core funded by Cancer Research UK (A31287). With particular thanks to Leo Carlin, Claire Mitchell and Peter Thomason from the Beatson Advanced Imaging Resource; Yi-Hsia Liu and Thomas Gilbey from the FACS facility and Colin Nixon and the Histology Laboratory for assistance with mouse and human tissue sample preparation and staining. We are grateful to Henri Jasper and Pedro Souza-Victor for their advice and help with cell sorting and mRNA sequencing of midgut ISCs/EBs. We thank all members of the Cordero laboratory for general advice on the project.

Human tumour sample analysis was performed on data generated by the TCGA Research Network: https://www.cancer.gov/tcga.

The manuscript was critically reviewed by Catherine Winchester (CRUK Scotland Institute).

Funding

Work in the Cordero laboratory is funded through a Wellcome Trust and Royal Society Sir Henry Dale Fellowship (104103/Z/14/Z), and Wellcome Trust Senior Research Fellowship (223091/Z/21/Z) to J.B.C., core Institutional funds from CRUK to the CRUK Scotland Institute (A31287) and a China Scholarship Council studentship to Y.T. (202006910022). Research in the Hayashi laboratory is supported by the Australian Research Council (DP210102385). The Edwards laboratory is funded by Chief Scientific Office grants (CSO EPD/22/13; K.P. and J.E.) (CSO TCS/22/02; J.E.) and CRUK funding to the CRUK Scotland Centre (CTRQQR-2021\100006). The Gontijo and Heredia laboratory was supported by the FCT (LISBOA-01-0145-FEDER-030753;

10.54499/CEECINST/00102/2018/CP1567/CT0031;

10.54499/DL57/2016/CP1457/CT0016); EXPL/BIA-BID/1524/2021; EXPL/BIA-

898 COM/1296/2021; 10.54499/2022.03859.PTDC, 10.54499/UIDB/04462/2020;
899 10.54499/UIDB/00329/2020, 10.54499/LA/P/0087/2020; 10.54499/LA/P/0121/2020;
900 LISBOA-01-0145-FEDER-022170; UID/Multi/04462/2019.
901

902 **Conflict of interest**
903 None declared.
904

905 **Data availability**
906 Sequencing data and processed files have been deposited to Gene Expression
907 Omnibus: GSE253621 (small RNA sequencing); GSE253624 (mRNA sequencing).
908 Codes for the computational analyses are available at
909 https://github.com/RippeiHayashi/gut_aubergine.
910

Author contributions
K.B., L.R.C., R.H and J.B.C design the experiments. K.B., L.R.C., Y.T., Y.Y., A.R.C.
and R.H. performed the experiments. K.B., L.R.C., R.H. and J.B.C. analysed the
data. K.A.P. and J.E. provided methodology for TMA analysis. F.H. and A.M.G.
generated, aub^{NH2} recombinant fly line. K.B. wrote the original manuscript draft. K.B.,
R.H. and J.B.C. wrote, reviewed, and edited the manuscript.
917

Figure legends

Figure 1: Aub is required for the proliferation of ISCs in response to intestinal
damage.

(A) Representative confocal image of a mated female adult *Drosophila*'s
gastrointestinal tract. ISCs and EBs are identified with GFP (green). Dashed yellow
lines delineate the posterior midgut. Scale bar is 200µm. **(A')** Representative
scheme of the *Drosophila* intestinal epithelium and associated tissues. ISCs,
intestinal stem cells; EBs, enteroblasts; EEPs, enteroendocrine cell precursors; ECs,
enterocytes; EEs, enteroendocrine cells; VM, visceral muscle; EN, enteric neurons;
TTC, terminal tracheal cell. **(B)** RT-qPCR for mRNA expression of the PIWI genes
aub, *piwi* and *ago3* in the R4-R5 midgut region (pink), the whole adult *Drosophila*
midgut (red), and adult *Drosophila* ovaries (green) relative to *rpl32* mRNA levels.
Number of biological replicates is 3 with each replicate containing 20 to 40
midguts/R4-R5 regions dissected. **(C)** Representative confocal images of PH3
staining (green) in the posterior midguts of w^{1118}, $piwi^1/piwi^2$, aub^{QC42}/aub^{HN2} and
ago^{t2}/ago^{t3} from whole mutant female flies fed with sucrose or *Pe*. Scale bar is
100µm. **(D)** Quantification of PH3-positive cells in posterior midguts as in (C).
n = number of midguts/flies quantified. **(E)** Quantification of PH3-positive cells in the
posterior midguts of w^{1118}, $aub^{G1/G2}$ and $aub^{clash/clash}$ flies fed with sucrose or *Pe*.
n = number of midguts/flies quantified. **(F)** Representative confocal images of PH3
staining (green) in the posterior midguts of flies expressing GFP in ISCs/EBs,
aub^{RNAi} alone or overexpressing aub^{RNAi} within ISCs/EBs and fed with sucrose or *Pe*.
Scale bar is 100µm. **(G)** Quantification of PH3-positive cells in the posterior midguts
as in (F). n = number of midguts/flies quantified. **(H, I)** Representative confocal
images of Aub staining (red) in the posterior midguts of flies expressing GFP only **(H)**
or with aub^{RNAi} **(I)** within ISCs/EBs (green) and fed with sucrose or *Pe*. Dashed
yellow squares delineate the magnified area shown in the right panels. Scale bar is
50µm. **(J)** Quantification of Aub staining within ISCs/EBs in the posterior midguts as
in (H, I). n = number of cells quantified. 3 cells per gut/fly have been quantified and
10 flies per condition have been analysed. **(K)** RT-qPCR for mRNA expression of
aub in whole adult *Drosophila* midguts relative to *rpl32* mRNA levels. Number of
biological replicates is 4 with 30 midguts dissected per replicate. A Shapiro-Wilk

normality test followed by a Mann-Whitney t test were used. **(L, M)** Quantification of
PH3-positive cells in the posterior midguts of flies fed with sucrose or *Pe* and
expressing GFP alone or with *aub^RNAi* in either EBs **(L)** or ISCs **(M)**.

Unless otherwise noted, two-way ANOVA followed by a Sidak's multiple
comparisons test were used for statistical analysis of all data sets. n are indicated
below each graph following the same order of the genotypes and treatments
presented in the graph. Data represent the mean ± s.d. ns, not significant
($P > 0.05$); *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$. a.u.,
arbitrary units. Nuclei are identified with DAPI.

Figure 2: Aubergine does not impact the presence of piRNAs in *Drosophila*
intestinal stem/progenitor cells.

(A) Schematic representation of the *Drosophila* intestine. The green dots represent
ISCs/EBs that were sorted *via* FACS for small RNA sequencing. **(B, B', C, C')** Size
distribution (top graphs) of all transposon-mapping reads and the Uridine- and
Adenine-frequencies of transposon-mapping reads of piRNA size (>22nt) (bottom
graphs) in oxidized libraries from sorted ISCs/EBs of control flies fed with sucrose or
Pe **(B, B')** or in oxidized libraries from sorted ISCs/EBs of flies expressing *aub^RNAi* in
ISCs/EBs and fed with sucrose or *Pe* **(C, C')**. siRNAs are typically sized at 21 or 22nt
while piRNAs are longer than 22nt (see dashed black rectangles for higher
magnification). Uridine (red) at the 1st base position of TE antisense piRNA reads
(>22nt) is a common signature of all piRNAs while Adenine (green) at the 10th base
position of TE sense piRNA reads is a signature of Ping-Pong piRNAs. **(D, E)**
Quantification of TE antisense **(D)** and sense **(E)** piRNA abundance as in (B, B', C,
C'). Two-way ANOVA followed by a Sidak's multiple comparisons test were applied.
n = 3 biological replicates. **(F)** Representation of piRNA cluster expression in sorted
ISCs/EBs from control flies or from flies expressing *aub^RNAi* in ISCs/EBs and fed with
sucrose or *Pe,* compared to ovaries. The results are presented relative to control
ovaries. Germline clusters 38C left and 42AB are represented in orange and blue,
respectively. The somatic cluster flamenco is represented in green.

Data represent the mean ± s.d. ns, not significant ($P > 0.05$); *$P < 0.05$,
**$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$. a.u., arbitrary units. TE,
transposable elements; AS, antisense; S, sense.

Figure 3: Aub regulates intestinal regeneration independently of its piRNA regulatory function.

(A, B) Abundance of TE mRNAs in sorted ISCs/EBs from control flies **(A)** or in flies expressing *aub^RNAi* within ISCs/EBs **(B)** and fed with sucrose or *Pe*. TE mRNA counts were normalised to one million transcript counts including host mRNAs (TPM), and Suc versus *Pe* fed conditions were compared. Red dashed lines indicate 4-fold changes in either direction. TE mRNAs shown in red, blue and green represent TE mRNAs highly expressed in the gut, in the ovaries or in both, respectively. **(C)** Representative confocal images of GFP (green) and Aub staining (red) in posterior midguts from control flies or flies expressing *aub^WT*, *aub^AA* or *aub^ADH* within ISCs/EBs. Dashed white lines delineate the posterior midguts. **(D, E)** Quantification of GFP **(D)** or Aub staining **(E)** within ISCs/EBs in the posterior midguts as in (C). A Shapiro-Wilk normality test followed by a Kruskal-Wallis one-way ANOVA and a Dunn's multiple comparisons tests were applied. n = number of midguts/flies quantified. **(F)** Representative confocal images of Aub staining (red) in the posterior midguts of *aub^HN2*/*aub^QC42* mutant flies expressing *aub^WT*, *aub^AA* or *aub^ADH* within ISCs/EBs and fed with sucrose or *Pe*. Dashed white lines delineate the posterior midguts. **(G)** Quantification of Aub staining within ISCs/EBs in posterior midguts as in (F). **(H)** Quantification of PH3-positive cells in the posterior midguts of control flies; *aub^HN2*/*aub^QC42* flies expressing GFP alone or with *aub^WT*, *aub^AA* or *aub^ADH* within ISCs/EBs and fed with sucrose or *Pe*. Two-way ANOVA followed by a Sidak's multiple comparisons test were applied in (G) and (H). In both cases, n = number of midguts/flies quantified, which are indicated below each graph following the same order of the genotypes and treatments presented in the graph.

Data represent the mean ± s.d. ns, not significant ($P > 0.05$); *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$. a.u., arbitrary units. Nuclei are identified with DAPI. Scale bars are 50µm. TE, transposable elements.

Figure 4: Aub regulates core regenerative stem cell factors in the adult midgut.

(A, B) Representative confocal images of puromycin staining (red) in the posterior midguts of flies expressing GFP alone **(A)** or with *aub^RNAi* **(B)** within ISCs (green) and fed with sucrose or *Pe*. **(C)** Quantification of puromycin staining within ISCs in the posterior midguts as in (A and B). **(D, E)** Representative confocal images of Sox21a staining (red) in the posterior midgut of flies expressing GFP alone **(D)** or

with aub^{RNAi} **(E)** within ISCs (green) and fed with sucrose or *Pe*. **(F)** Quantification of
Sox21a staining within ISCs in the posterior midguts as in (D, E). **(G, H)**
Representative confocal images of Myc staining (red) in the posterior midguts of flies
expressing GFP alone **(G)** or with aub^{RNAi} **(H)** within ISCs (green) and fed with
sucrose or *Pe*. **(I)** Quantification of Myc staining within ISCs in the posterior midguts
as in (G, H). Two-way ANOVA followed by Sidak's multiple comparisons test were
applied in (C, F and I). n = number of midguts/flies quantified and are indicated
below each graph in the same order of the genotypes and treatments presented in
the graph.
Data represent the mean ± s.d. ns, not significant ($P > 0.05$); *$P < 0.05$,
**$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$. a.u., arbitrary units. Nuclei are
identified with DAPI. Scale bars are 50µm.
1032

Figure 5: Aub regulates the multiprotein complex eIF3.
(A) Representative confocal images of eIF3C staining (red) in the posterior midguts
of flies expressing GFP within ISCs (green) and fed with sucrose or *Pe*. Dashed
white squares delineate the high magnification shown in the right panels. Yellow
arrows highlight ISCs. **(B)** Quantification of eIF3C staining within ISCs in the
posterior midguts as in (A). A Shapiro-Wilk normality test followed by an unpaired t
test were applied. n = number of midguts/flies quantified. **(C)** Quantification of PH3-
positive cells in the posterior midguts of control flies expressing GFP alone or with
RNAi against different subunits of the eIF3 complex within ISCs. Flies were fed with
sucrose or *Pe*. n = number of midguts/flies quantified and are indicated below each
graph in the same order of the genotypes and treatments presented in the graph. **(D)**
Quantification of PH3-positive cells in the posterior midguts of control flies
expressing GFP only; with *aub::GFP* or with *aub::GFP* and $eIF3C^{RNAi}$ within ISCs. A
Shapiro-Wilk normality test followed by a Kruskal-Wallis one-way ANOVA and a
Dunn's multiple comparisons tests were applied. n = number of midguts/flies
quantified and are indicated below each graph in the same order of the genotypes
and treatments presented in the graph. **(E, F)** Representative confocal images of
Sox21a staining (red) in the posterior midgut of flies expressing GFP alone **(E)** or
with $eIF3C^{RNAi}$ **(F)** within ISCs (green) and fed with sucrose or *Pe*. **(G)** Quantification
of Sox21a staining within ISCs in the posterior midguts as in (E, F). **(H, I)**
Representative confocal images of Myc staining (red) in the posterior midguts of flies

expressing GFP alone **(H)** or with *eIF3C^{RNAi}* **(I)** within ISCs (green) and fed with sucrose or *Pe.* **(J)** Quantification of Myc staining within ISCs in the posterior midguts as in (H, I). **(K)** Quantification of eIF3C staining within ISCs in the posterior midguts as in (Fig.S5E-G). Two-way ANOVA followed by Sidak's multiple comparisons test were applied for statistical analysis in (C, G, J and K). n = number of midguts/flies quantified and are indicated below each graph in the same order of the genotypes and treatments presented in the graph. **(L)** Current working model on the regulation of regenerative ISC by Aub. ISCs: intestinal stem cells; EBs: enteroblasts, EC: enterocyte; VM: Visceral Muscle; TC: tracheal cell.

Data represent the mean ± s.d. ns, not significant ($P > 0.05$); *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$. a.u., arbitrary units. Nuclei are identified with DAPI. Scale bars are 50µm.

Figure 6: The mammalian homolog of Aub, PIWIL1, is upregulated in CRC patients.

(A, B) *PIWIL1* expression obtained from RNA sequencing experiments performed in non-paired normal and tumour tissues (**A**) or in paired adjacent normal and tumour tissues (**B**) from human colon adenocarcinoma. Graphs and statistics were done using the TNMplot tool. Mann-Whitney t test was applied. n = number of patients quantified. **(C)** *PIWIL1* expression from ChIP sequencing experiments performed in non-paired normal, tumour and metastatic tissues from human colon adenocarcinoma. Graph and statistics were done using the TNMplot tool. A Kruskal-Wallis one-way ANOVA test was applied. n = number of patients quantified. **(D)** *PIWIL1* expression obtained across 4 CRC subtypes (Guinney et al., 2015) (TCGA, PanCancer Atlas). Shapiro-Wilk normality test followed by a Kruskal-Wallis one-way ANOVA and Dunn's multiple comparisons tests were applied. n = number of patients quantified. **(E)** *PIWIL1* expression according to the microsatellite status of cancer patients (TCGA, PanCancer Atlas). MSS: microsatellite stability; MSI-L: low microsatellite instability; MSI-H: high microsatellite instability. Shapiro-Wilk normality test followed by a Kruskal-Wallis one-way ANOVA and Dunn's multiple comparisons tests were applied. n = number of patients quantified. **(F)** Representative microscopic images of PIWIL1 expression in TMA from CRC patients. Scale bar is 100 µm. **(G)** Kaplan-Meier survival analysis of cytoplasmic PIWIL1 expression in the cohort of CRC patients.

1088   For box plots, the line is the median and the box represents the values included

1089   between the 25th and 75th percentiles. For other graphs, data represent the

1090   mean $\pm$ s.d. ns, not significant ($P > 0.05$); *$P < 0.05$, **$P < 0.01$,

1091   ***$P < 0.001$ and ****$P < 0.0001$. a.u., arbitrary units.

1092

Figure 7: Aub mediates CRC-like intestinal hyperplasia in *Drosophila*.

(A, B) Representative confocal images of Aub staining (red) in the posterior midguts

of control flies **(A)** or *Apc^{1Q8}* mutant flies **(B)** expressing GFP within ISCs/EBs

(green). Dashed yellow squares delineate the high magnification shown in the right

panels. **(C)** Quantification of Aub staining within ISCs/EBs in posterior midguts as in

(A and B). Shapiro-Wilk normality test followed by Mann-Whitney t test were applied.

n = number of cells quantified. 3 cells per gut/fly have been quantified and 10 flies

per condition have been analysed. **(D)** Representative confocal images of MARCM

clones (green) expressing a *LacZ* control transgene only (41 flies analysed) or in

combination with *aub^{RNAi}* (33 flies analysed), and *Apc^{1Q8}* mutant clones only (35 flies

analysed) or co-expressing *aub^{RNAi}* (38 flies analysed). Dashed white lines delineate

the clonal margins. **(E)** Quantification of the number of cells in each MARCM clone of

genotypes as in (D). Shapiro-Wilk normality test followed by Kruskal-Wallis one-way

ANOVA and a Dunn's multiple comparisons tests were applied. n = number of clones

quantified. **(F)** Quantification of PH3-positive cells in the posterior midguts of control

flies expressing GFP alone; with *Wg::HA* or *aub^{RNAi}* or co-expressing *Wg::HA* and

aub^{RNAi} within ISCs/EBs. A Shapiro-Wilk normality test followed by a Kruskal-Wallis

one-way ANOVA and a Dunn's multiple comparisons tests were applied. n = number

of midguts/flies quantified and are indicated below each graph in the same order of

the genotypes and treatments presented in the graph. **(G, H)** Representative

confocal images of Aub staining (red) in the posterior midguts of control flies

expressing GFP only **(G)** or with the activated form of human *BRAF* (*Raf^{GOF}*) **(H)**

within ISCs/EBs (green) for 2 days. Dashed yellow squares delineate the high

magnification shown in the right panels. **(I)** Quantification of Aub staining within

ISCs/EBs in posterior midguts as in (G and H). Shapiro-Wilk normality test followed

by an unpaired t test were applied. n = number of cells quantified. 3 cells per gut/fly

have been quantified and 10 flies per condition have been analysed. **(J, K)**

Quantification of PH3-positive cells in the posterior midguts of control flies

expressing GFP only; *Raf^{GOF}* or *Raf^{GOF}* and *aub^{RNAi}* within ISCs/EBs for 2 **(J)** or 3

days **(K)**, respectively. Shapiro-Wilk normality test followed by Kruskal-Wallis one-

way ANOVA and Dunn's multiple comparisons tests were applied. n = number of

midguts/flies quantified and are indicated below each graph in the same order of the

genotypes and treatments presented in the graph. **(L)** Quantification of Dcp-1-

positive cells in the lumen of midguts as in (Fig.S6D). Shapiro-Wilk normality test

followed by one-way ANOVA and Sidak's multiple comparisons tests were applied.

n = number of midguts/flies quantified and are indicated below each graph in the

same order of the genotypes and treatments presented in the graph.

Data represent the mean□±□s.d. ns, not significant (P□>□0.05); *P□<□0.05,

**P□<□0.01, ***P□<□0.001 and ****P□<□0.0001. a.u., arbitrary units. Nuclei are

identified with DAPI. Scale bars are 50μm.

1133

Figure S1: Aub regulates intestinal regeneration independently of piRNAs.

(A) Quantification of PH3-positive cells in the posterior midguts of flies fed with

sucrose or *Pe* and expressing GFP only, with *piwi*^{iRNAi} or *ago3*^{RNAi} within ISCs/EBs.

n = number of midguts/flies quantified. **(B)** Representative confocal images of 7 or 14

days old MARCM clones (green) expressing a *LacZ* transgene only (4 and 9 flies

analysed, respectively) or with *aub*^{RNAi} (5 and 8 flies analysed, respectively). Dashed

white lines delineate clonal margins. **(C)** Quantification of the size of MARCM clones

as in (B). n = number of clones quantified. **(D)** Quantification of PH3-positive cells in

the posterior midguts of w^{1118} flies, *spnE*^{Δ125}/*spnE*$^{100-37}$; *vasa*^{PH165}/*vasa*^{D5} or *rhi*$^{18-}$

^{5}/*rhi*$^{18-7}$ flies fed with sucrose or *Pe*. n = number of midguts/flies quantified. **(E)**

Quantification of PH3-positive cells in the posterior midguts of flies fed with sucrose

or *Pe* and expressing GFP alone or with *vasa*^{RNAi}, *armi*^{RNAi} or *rhino*^{RNAi} within

ISCs/EBs. n = number of midguts/flies quantified. **(F, G, H)** Representative confocal

images of Aub staining (red) in the posterior midguts of flies expressing GFP alone

(F), with *piwi*^{RNAi} **(G)** or with *vasa*^{RNAi} **(H)** within ISCs (green) and fed with sucrose or

Pe. Dashed yellow squares delineate the high magnification shown in the right

panels. **(I)** Quantification of Aub staining within ISCs in the posterior midguts as in (F,

G, H). n=number of cells quantified. 3 cells per gut/fly have been quantified and 10

flies per condition have been analysed.

In all cases, two-way ANOVA followed by Sidak's multiple comparisons test were

used for statistical analysis. N are indicated below each graph in the same order of

the genotypes and treatments presented in the graph.

Data represent the mean□±□s.d. ns, not significant (P□>□0.05); *P□<□0.05,
**P□<□0.01, ***P□<□0.001 and ****P□<□0.0001. a.u., arbitrary units. Nuclei are
identified with DAPI. Scale bars are 50μm.

1159

Figure S2: Presence of piRNAs in the *Drosophila* midgut.
(A) Schematic representation of the *Drosophila* intestine. The green area highlights
the portion of the intestine that was dissected for small RNA sequencing. **(B, B', C,**
C') Size distribution (top graphs) of all transposon-mapping reads and the Uridine-
and Adenine-frequencies of transposon-mapping reads of piRNA size (>22nt)
(bottom graphs) in oxidized libraries from the whole intestine of control flies fed with
sucrose or *Pe* **(B, B')** or in oxidized libraries from the whole intestine of *aub* mutant
flies fed with sucrose or *Pe* **(C, C')**. siRNAs are typically sized at 21 or 22nt while
piRNAs are longer than 22nt (see dashed black rectangles for higher magnification).
Uridine (red) at the 1st base position of TE antisense piRNA reads (>22nt) is a
common signature of all piRNAs while Adenine (green) at the 10th base position of
TE sense piRNA reads is a signature of Ping-Pong piRNAs. **(D, E)** Quantification of
TE antisense **(D)** and sense **(E)** piRNAs abundance as in (B, B', C, C'). Two-way
ANOVA followed by a Sidak's multiple comparisons test were applied. n = 3
biological replicates.
Data represent the mean□±□s.d. ns, not significant (P□>□0.05); *P□<□0.05,
**P□<□0.01, ***P□<□0.001 and ****P□<□0.0001. a.u., arbitrary units. TE,
transposable elements; AS, antisense; S, sense.

1178

Figure S3: Presence of piRNAs in the *Drosophila* intestine.
(A, B) Size distribution (top graphs) of all transposon-mapping reads and the
Uridine- and Adenine-frequencies of transposon-mapping reads of piRNA size
(>22nt) (bottom graphs) in oxidized libraries from control ovaries **(A)** or in unoxidized
libraries from the whole intestine of control flies previously published (Siudeja et al.,
2021) **(B)**. siRNAs are typically sized at 21 or 22nt while piRNAs are longer than
22nt (see dashed black rectangles for higher magnification). Uridine (red) at the 1st
base position of TE antisense piRNA reads (>22nt) is a common signature of all
piRNAs while Adenine (green) at the 10th base position of TE sense piRNA reads is
a signature of Ping-Pong piRNAs. **(C-G)** Representation of the 0.5 kb tile analysis of
piRNA abundance measurement in sorted ISCs/EBs from control flies fed with

sucrose or *Pe* **(C, D)** or from flies expressing *aub^RNAi* in ISCs/EBs and fed with

sucrose or *Pe* **(E, F)** compared to ovaries. The same analysis was performed to

compare piRNA abundance between ovaries from control flies and from flies

expressing *aub^RNAi* in ISCs/EBs **(G)**. Tiles from piRNA clusters are coloured: cluster

38C in orange, cluster 42AB in blue and cluster flamenco in green. All other tiles are

coloured in grey. a.u., arbitrary units. TE, transposable elements; AS, antisense; S,

sense.

1197

Figure S4: TE mRNA expression in ISCs/EBs and ovaries.

(A, B, C) Abundance of TE mRNAs in sorted control **(A)**; or *aub^RNAi* ISCs/EBs **(B)**

from sucrose and *Pe* fed animals or from control and *aub^RNAi* ovaries obtained from a

previously published study **(C)** (Senti et al., 2015). TE mRNA counts were

normalised to one million transcript counts including host mRNAs (TPM). Red

dashed lines indicate 4-fold changes in either direction. Red, blue and green

represent TE mRNAs highly expressed in the gut, in the ovaries or in both tissues,

respectively. **(D)** Schematic representation of the site-directed transgenesis induced

to generate fly lines carrying inducible GFP tagged wild type Aub (*UAS-aub^WT*) or

Aub mutant isoforms (*UAS-aub^AA* and *UAS-aub^ADH*). *UAS-aub^AA* contains a double

point mutation in the PAZ domain (yellow), responsible for the loading of piRNAs,

while *UAS-aub^ADH* carries a single mutation in the PIWI domain (red), responsible for

Aub's endonuclease activity and essential for piRNA biogenesis. **(E)** Female fertility

assays in *aub^HN2/aub^QC42* mutant flies without or with expression of *aub^WT*, *aub^AA* or

aub^ADH in the germline. a.u., arbitrary units. TE, transposable elements.

1213

Figure S5: Aub regulates protein translation of Sox21a and Myc.

(A, B, C) RT-qPCR for mRNA expression of *sox21a* **(A)**, *myc* **(B)** *and socs36e* **(C)** in

the whole adult *Drosophila* midgut relative to *rpl32* mRNA levels. n = 4 biological

replicates with 30 midguts dissected for each replicate. Two-way ANOVA followed by

a Sidak's multiple comparisons test were applied for statistical analysis. n = number

of biological replicates and are indicated below each graph in the same order of the

genotypes and treatments presented in the graph. **(D)** Representative confocal

images of puromycin staining (red) in the posterior midguts of flies expressing

eIF3C^RNAi within ISCs (green) and fed with sucrose or *Pe*. **(D')** Quantification of

puromycin staining within ISCs in posterior midguts as in (Fig.4A-C and S5D). Two-

way ANOVA followed by Sidak's multiple comparisons test were applied. n = number of midguts/flies quantified and are indicated below each graph in the same order of the genotypes and treatments presented in the graph. **(E, F, G)** Representative confocal images of eIF3C staining (red) in the posterior midgut of flies expressing GFP alone **(E)**, with *myc^RNAi* **(F)** or *sox21a^RNAi* **(G)** within ISCs/EBs (green) and fed with sucrose or *Pe*. Quantification is shown in Fig.5K.

Data represent the mean □±□s.d. ns, not significant ($P□>□0.05$); *$P□<□0.05$, **$P□<□0.01$, ***$P□<□0.001$ and ****$P□<□0.0001$. a.u., arbitrary units. Nuclei are identified with DAPI. Scale bars are 50µm.

Figure S6: Aub role in intestinal tumourigenesis by MAPK signalling.

(A) Piwil1 expression in wild type and *Piwil1* mutant mouse testis. Dashed black squares delineate the high magnification shown in the right panels. **(B, C)** Quantification of PH3-positive cells in the posterior midguts of control flies expressing GFP alone or with *aub^RNAi* within ISCs/EBs. Flies were fed with sucrose or *Pe*. The quantification was performed after 2 (**B**) or 3 days (**C**) of RNAi induction. Two-way ANOVA followed by Sidak's multiple comparisons test were applied. n = number of midguts/flies quantified and are indicated below each graph in the same order of the genotypes and treatments presented in the graph. **(D)** Apical, basal and orthogonal confocal images of Dcp-1 staining (grey) to detect apoptosis in midguts overexpressing for 3 days GFP alone or with *Raf^GOF* in the absence or presence of *aub^RNAi*. Flies were fed with sucrose. Quantification is shown Fig.7L. (**E**) Quantification of Aub staining within ISCs/EBs in posterior midguts. A Shapiro-Wilk normality test followed by a Mann-Whitney t test were applied. n = number of cells quantified. 3 cells per gut/fly have been quantified and 15 flies per condition have been analysed. **(F, G)** Quantification of PH3-positive cells in the posterior midguts of flies expressing GFP alone; with *Ras^V12* or with *Ras^V12* and *aub^RNAi* within ISCs/EBs. Quantification was performed after 2 days **(F)** of 7 days **(G)** of RNAi co-expression with *Ras^V12*. Shapiro-Wilk normality test followed by a Kruskal-Wallis one-way ANOVA and Dunn's multiple comparisons tests were applied. n = number of midguts/flies quantified and are indicated below each graph in the same order of the genotypes and treatments presented in the graph.

1256　Data represent the mean ± s.d. ns, not significant ($P > 0.05$); *$P < 0.05$,

1257　**$P < 0.01$, ***$P < 0.001$ and ****$P < 0.0001$. a.u., arbitrary units. Nuclei are

1258　identified with DAPI. Scale bars are 50μm.

1259

1260

References

References

AKBERGENOV, R., SI-AMMOUR, A., BLEVINS, T., AMIN, I., KUTTER, C., VANDERSCHUREN, H., ZHANG,
P., GRUISSEM, W., MEINS, F., JR., HOHN, T. & POOGGIN, M. M. 2006. Molecular
characterization of geminivirus-derived small RNAs in different plant species. *Nucleic Acids*
Res, 34**,** 462-71.
ARAÚJO, T., KHAYAT, A., QUINTANA, L., CALCAGNO, D., MOURÃO, R., MODESTO, A., PAIVA, J., LIMA,
A., MOREIRA, F., OLIVEIRA, E., SOUZA, M., OTHMAN, M., LIEHR, T., ABDELHAY, E., GOMES,
R., SANTOS, S. & ASSUMPÇÃO, P. 2018. Piwi like RNA-mediated gene silencing 1 gene as a
possible major player in gastric cancer. *World J Gastroenterol,* 24**,** 5338-5350.
ARAVIN, A., GAIDATZIS, D., PFEFFER, S., LAGOS-QUINTANA, M., LANDGRAF, P., IOVINO, N., MORRIS,
P., BROWNSTEIN, M. J., KURAMOCHI-MIYAGAWA, S., NAKANO, T., CHIEN, M., RUSSO, J. J.,
JU, J., SHERIDAN, R., SANDER, C., ZAVOLAN, M. & TUSCHL, T. 2006. A novel class of small
RNAs bind to MILI protein in mouse testes. *Nature,* 442**,** 203-207.
ARAVIN, A. A., HANNON, G. J. & BRENNECKE, J. 2007. The Piwi-piRNA Pathway Provides an Adaptive
Defense in the Transposon Arms Race. *Science,* 318**,** 761-764.
ARAVIN, A. A., SACHIDANANDAM, R., GIRARD, A., FEJES-TOTH, K. & HANNON, G. J. 2007.
Developmentally Regulated piRNA Clusters Implicate MILI in Transposon Control. *Science,*
316**,** 744-747.
ASANO, K., CLAYTON, J., SHALEV, A. & HINNEBUSCH, A. G. 2000. A multifactor complex of eukaryotic
initiation factors, eIF1, eIF2, eIF3, eIF5, and initiator tRNA(Met) is an important translation
initiation intermediate in vivo. *Genes Dev,* 14**,** 2534-46.
BANGI, E., MURGIA, C., TEAGUE, A. G., SANSOM, O. J. & CAGAN, R. L. 2016. Functional exploration of
colorectal cancer genomes using Drosophila. *Nat Commun,* 7**,** 13615.
BARCKMANN, B., PIERSON, S., DUFOURT, J., PAPIN, C., ARMENISE, C., PORT, F., GRENTZINGER, T.,
CHAMBEYRON, S., BARONIAN, G., DESVIGNES, J.-P., CURK, T. & SIMONELIG, M. 2015.
Aubergine iCLIP Reveals piRNA-Dependent Decay of mRNAs Involved in Germ Cell
Development in the Early Embryo. *Cell Reports,* 12**,** 1205-1216.
BARTHA, Á. & GYŐRFFY, B. 2021. TNMplot.com: A Web Tool for the Comparison of Gene Expression
in Normal, Tumor and Metastatic Tissues. *Int J Mol Sci,* 22.
BECALSKA, A. N., KIM, Y. R., BELLETIER, N. G., LERIT, D. A., SINSIMER, K. S. & GAVIS, E. R. 2011.
Aubergine is a component of a nanos mRNA localization complex. *Dev Biol,* 349**,** 46-52.
BEEHLER-EVANS, R. & MICCHELLI, C. A. 2015. Generation of enteroendocrine cell diversity in midgut
stem cell lineages. *Development,* 142**,** 654-64.
BITEAU, B. & JASPER, H. 2011. EGF signaling regulates the proliferation of intestinal stem cells in
Drosophila. *Development,* 138**,** 1045-55.
BONFINI, A., DOBSON, A. J., DUNEAU, D., REVAH, J., LIU, X., HOUTZ, P. & BUCHON, N. 2021.
Multiscale analysis reveals that diet-dependent midgut plasticity emerges from alterations in
both stem cell niche coupling and enterocyte size. *Elife,* 10.
BRAND, A. H. & PERRIMON, N. 1994. Raf acts downstream of the EGF receptor to determine
dorsoventral polarity during Drosophila oogenesis. *Genes Dev,* 8**,** 629-39.
BRENNECKE, J., ARAVIN, A. A., STARK, A., DUS, M., KELLIS, M., SACHIDANANDAM, R. & HANNON, G.
J. 2007. Discrete Small RNA-Generating Loci as Master Regulators of Transposon Activity in
Drosophila. *Cell,* 128**,** 1089-1103.
BROWER-TOLAND, B., FINDLEY, S. D., JIANG, L., LIU, L., YIN, H., DUS, M., ZHOU, P., ELGIN, S. C. & LIN,
H. 2007. Drosophila PIWI associates with chromatin and interacts directly with HP1a. *Genes*
Dev, 21**,** 2300-11.
BUCHON, N., BRODERICK, N. A., KURAISHI, T. & LEMAITRE, B. 2010. Drosophila EGFR pathway
coordinates stem cell proliferation and gut remodeling following infection. *BMC Biol,* 8**,** 152.

BUCHON, N., OSMAN, D., DAVID, F. P., FANG, H. Y., BOQUETE, J. P., DEPLANCKE, B. & LEMAITRE, B. 2013. Morphological and molecular characterization of adult midgut compartmentalization in Drosophila. *Cell Rep,* 3**,** 1725-38.

CHEN, J., XU, N., WANG, C., HUANG, P., HUANG, H., JIN, Z., YU, Z., CAI, T., JIAO, R. & XI, R. 2018. Transient Scute activation via a self-stimulatory loop directs enteroendocrine cell pair specification from self-renewing intestinal stem cells. *Nat Cell Biol,* 20**,** 152-161.

CHEN, Z., CHE, Q., JIANG, F. Z., WANG, H. H., WANG, F. Y., LIAO, Y. & WAN, X. P. 2015. Piwil1 causes epigenetic alteration of PTEN gene via upregulation of DNA methyltransferase in type I endometrial cancer. *Biochem Biophys Res Commun,* 463**,** 876-80.

COOK, H. A., KOPPETSCH, B. S., WU, J. & THEURKAUF, W. E. 2004. The Drosophila SDE3 Homolog armitage Is Required for oskar mRNA Silencing and Embryonic Axis Specification. *Cell,* 116**,** 817-829.

CORDERO, J. B., STEFANATOS, R. K., MYANT, K., VIDAL, M. & SANSOM, O. J. 2012a. Non-autonomous crosstalk between the Jak/Stat and Egfr pathways mediates Apc1-driven intestinal stem cell hyperplasia in the Drosophila adult midgut. *Development,* 139**,** 4524-35.

CORDERO, J. B., STEFANATOS, R. K., SCOPELLITI, A., VIDAL, M. & SANSOM, O. J. 2012b. Inducible progenitor-derived Wingless regulates adult midgut regeneration in Drosophila. *The EMBO Journal,* 31**,** 3901-3917.

COX, D. N., CHAO, A., BAKER, J., CHANG, L., QIAO, D. & LIN, H. 1998. A novel class of evolutionarily conserved genes defined by piwi are essential for stem cell self-renewal. *Genes Dev,* 12**,** 3715-27.

COX, D. N., CHAO, A. & LIN, H. 2000. piwi encodes a nucleoplasmic factor whose activity modulates the number and division rate of germline stem cells. *Development,* 127**,** 503-514.

DAVID, A., DOLAN, B. P., HICKMAN, H. D., KNOWLTON, J. J., CLAVARINO, G., PIERRE, P., BENNINK, J. R. & YEWDELL, J. W. 2012. Nuclear translation visualized by ribosome-bound nascent chain puromycylation. *J Cell Biol,* 197**,** 45-57.

DELIU, L. P., GHOSH, A. & GREWAL, S. S. 2017. Investigation of protein synthesis in Drosophila larvae using puromycin labelling. *Biol Open,* 6**,** 1229-1234.

DUFOURT, J., BONTONOU, G., CHARTIER, A., JAHAN, C., MEUNIER, A.-C., PIERSON, S., HARRISON, P. F., PAPIN, C., BEILHARZ, T. H. & SIMONELIG, M. 2017. piRNAs and Aubergine cooperate with Wispy poly(A) polymerase to stabilize mRNAs in the germ plasm. *Nature Communications,* 8**,** 1305.

FUJII, K., ZHULYN, O., BYEON, G. W., GENUTH, N. R., KERR, C. H., WALSH, E. M. & BARNA, M. 2021. Controlling tissue patterning by translational regulation of signaling transcripts through the core translation factor eIF3c. *Dev Cell,* 56**,** 2928-2937.e9.

GENZOR, P., CORDTS, S. C., BOKIL, N. V. & HAASE, A. D. 2019. Aberrant expression of select piRNA-pathway genes does not reactivate piRNA silencing in cancer cells. *Proceedings of the National Academy of Sciences,* 116**,** 11111-11112.

GIRARD, A., SACHIDANANDAM, R., HANNON, G. J. & CARMELL, M. A. 2006. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature,* 442**,** 199-202.

GUINNEY, J., DIENSTMANN, R., WANG, X., DE REYNIÈS, A., SCHLICKER, A., SONESON, C., MARISA, L., ROEPMAN, P., NYAMUNDANDA, G., ANGELINO, P., BOT, B. M., MORRIS, J. S., SIMON, I. M., GERSTER, S., FESSLER, E., DE SOUSA E MELO, F., MISSIAGLIA, E., RAMAY, H., BARRAS, D., HOMICSKO, K., MARU, D., MANYAM, G. C., BROOM, B., BOIGE, V., PEREZ-VILLAMIL, B., LADERAS, T., SALAZAR, R., GRAY, J. W., HANAHAN, D., TABERNERO, J., BERNARDS, R., FRIEND, S. H., LAURENT-PUIG, P., MEDEMA, J. P., SADANANDAM, A., WESSELS, L., DELORENZI, M., KOPETZ, S., VERMEULEN, L. & TEJPAR, S. 2015. The consensus molecular subtypes of colorectal cancer. *Nature Medicine,* 21**,** 1350-1356.

GUNAWARDANE, L. S., SAITO, K., NISHIDA, K. M., MIYOSHI, K., KAWAMURA, Y., NAGAMI, T., SIOMI, H. & SIOMI, M. C. 2007. A Slicer-Mediated Mechanism for Repeat-Associated siRNA 5' End Formation in Drosophila. *Science,* 315**,** 1587-1590.

HAFNER, M., LANDGRAF, P., LUDWIG, J., RICE, A., OJO, T., LIN, C., HOLOCH, D., LIM, C. & TUSCHL, T.
2008. Identification of microRNAs and other small regulatory RNAs using cDNA library
sequencing. *Methods,* 44**,** 3-12.
HAN, B. W., WANG, W., LI, C., WENG, Z. & ZAMORE, P. D. 2015. piRNA-guided transposon cleavage
initiates Zucchini-dependent, phased piRNA production. *Science,* 348**,** 817-821.
HARRIS, A. N. & MACDONALD, P. M. 2001. aubergine encodes a Drosophila polar granule component
required for pole cell formation and related to eIF2C. *Development,* 128**,** 2823-2832.
HOUWING, S., KAMMINGA, L. M., BEREZIKOV, E., CRONEMBOLD, D., GIRARD, A., VAN DEN ELST, H.,
FILIPPOV, D. V., BLASER, H., RAZ, E., MOENS, C. B., PLASTERK, R. H., HANNON, G. J., DRAPER,
B. W. & KETTING, R. F. 2007. A role for Piwi and piRNAs in germ cell maintenance and
transposon silencing in Zebrafish. *Cell,* 129**,** 69-82.
HUANG, H., YU, X., HAN, X., HAO, J., ZHAO, J., BEBEK, G., BAO, S., PRAYSON, R. A., KHALIL, A. M.,
JANKOWSKY, E. & YU, J. S. 2021. Piwil1 Regulates Glioma Stem Cell Maintenance and
Glioblastoma Progression. *Cell Reports,* 34**,** 108522.
HUANG, XIAO A., YIN, H., SWEENEY, S., RAHA, D., SNYDER, M. & LIN, H. 2013. A Major Epigenetic
Programming Mechanism Guided by piRNAs. *Developmental Cell,* 24**,** 502-516.
IPSARO, J. J., HAASE, A. D., KNOTT, S. R., JOSHUA-TOR, L. & HANNON, G. J. 2012. The structural
biochemistry of Zucchini implicates it as a nuclease in piRNA biogenesis. *Nature,* 491**,** 279-
83.
JANIC, A., MENDIZABAL, L., LLAMAZARES, S., ROSSELL, D. & GONZALEZ, C. 2010. Ectopic Expression
of Germline Genes Drives Malignant Brain Tumor Growth in *Drosophila. Science,* 330**,** 1824-
1827.
JIANG, H., GRENLEY, M. O., BRAVO, M. J., BLUMHAGEN, R. Z. & EDGAR, B. A. 2011. EGFR/Ras/MAPK
signaling mediates adult midgut epithelial homeostasis and regeneration in Drosophila. *Cell
Stem Cell,* 8**,** 84-95.
JIANG, H., PATEL, P. H., KOHLMAIER, A., GRENLEY, M. O., MCEWEN, D. G. & EDGAR, B. A. 2009.
Cytokine/Jak/Stat Signaling Mediates Regeneration and Homeostasis in the Drosophila
Midgut. *Cell,* 137**,** 1343-1355.
JIVOTOVSKAYA, A. V., VALÁSEK, L., HINNEBUSCH, A. G. & NIELSEN, K. H. 2006. Eukaryotic translation
initiation factor 3 (eIF3) and eIF2 can promote mRNA binding to 40S subunits independently
of eIF4G in yeast. *Mol Cell Biol,* 26**,** 1355-72.
JONES, B. C., WOOD, J. G., CHANG, C., TAM, A. D., FRANKLIN, M. J., SIEGEL, E. R. & HELFAND, S. L.
2016. A somatic piRNA pathway in the Drosophila fat body ensures metabolic homeostasis
and normal lifespan. *Nat Commun,* 7**,** 13856.
KIRINO, Y. & MOURELATOS, Z. 2007. Mouse Piwi-interacting RNAs are 2'-O-methylated at their 3'
termini. *Nat Struct Mol Biol,* 14**,** 347-8.
KNIGHT, J. R. P., ALEXANDROU, C., SKALKA, G. L., VLAHOV, N., PENNEL, K., OFFICER, L., TEODOSIO, A.,
KANELLOS, G., GAY, D. M., MAY-WILSON, S., SMITH, E. M., NAJUMUDEEN, A. K., GILROY, K.,
RIDGWAY, R. A., FLANAGAN, D. J., SMITH, R. C. L., MCDONALD, L., MACKAY, C., CHEASTY, A.,
MCARTHUR, K., STANWAY, E., LEACH, J. D., JACKSTADT, R., WALDRON, J. A., CAMPBELL, A.
D., VLACHOGIANNIS, G., VALERI, N., HAIGIS, K. M., SONENBERG, N., PROUD, C. G., JONES, N.
P., SWARBRICK, M. E., MCKINNON, H. J., FALLER, W. J., LE QUESNE, J., EDWARDS, J., WILLIS,
A. E., BUSHELL, M. & SANSOM, O. J. 2021. MNK Inhibition Sensitizes KRAS-Mutant Colorectal
Cancer to mTORC1 Inhibition by Reducing eIF4E Phosphorylation and c-MYC Expression.
Cancer Discov, 11**,** 1228-1247.
KORZELIUS, J., AZAMI, S., RONNEN-ORON, T., KOCH, P., BALDAUF, M., MEIER, E., RODRIGUEZ-
FERNANDEZ, I. A., GROTH, M., SOUSA-VICTOR, P. & JASPER, H. 2019. The WT1-like
transcription factor Klumpfuss maintains lineage commitment of enterocyte progenitors in
the Drosophila intestine. *Nature Communications,* 10**,** 4123.

KRISHNAN, P., GHOSH, S., GRAHAM, K., MACKEY, J. R., KOVALCHUK, O. & DAMARAJU, S. 2016. Piwi-
interacting RNAs and PIWI genes as novel prognostic markers for breast cancer. *Oncotarget,*
7.
LEE, T. & LUO, L. 1999. Mosaic analysis with a repressible cell marker for studies of gene function in
neuronal morphogenesis. *Neuron,* 22**,** 451-61.
LEE, W. C., BEEBE, K., SUDMEIER, L. & MICCHELLI, C. A. 2009. Adenomatous polyposis coli regulates
Drosophila intestinal stem cell proliferation. *Development,* 136**,** 2255-64.
LI, C., VAGIN, V. V., LEE, S., XU, J., MA, S., XI, H., SEITZ, H., HORWICH, M. D., SYRZYCKA, M., HONDA,
B. M., KITTLER, E. L. W., ZAPP, M. L., KLATTENHOFF, C., SCHULZ, N., THEURKAUF, W. E.,
WENG, Z. & ZAMORE, P. D. 2009. Collapse of Germline piRNAs in the Absence of Argonaute3
Reveals Somatic piRNAs in Flies. *Cell,* 137**,** 509-521.
LI, F., YUAN, P., RAO, M., JIN, C.-H., TANG, W., RONG, Y.-F., HU, Y.-P., ZHANG, F., WEI, T., YIN, Q.,
LIANG, T., WU, L., LI, J., LI, D., LIU, Y., LOU, W., ZHAO, S. & LIU, M.-F. 2020. piRNA-
independent function of PIWIL1 as a co-activator for anaphase promoting
complex/cyclosome to drive pancreatic cancer metastasis. *Nature Cell Biology,* 22**,** 425-438.
LI, W., MARTINEZ-USEROS, J., GARCIA-CARBONERO, N., FERNANDEZ-ACEÑERO, M. J., ORTEGA-
MEDINA, L., GARCIA-BOTELLA, S., PEREZ-AGUIRRE, E., DIEZ-VALLADARES, L. & GARCIA-
FONCILLAS, J. 2019. The Prognosis Value of PIWIL1 and PIWIL2 Expression in Pancreatic
Cancer. *J Clin Med,* 8.
LIN, G., XU, N. & XI, R. 2008. Paracrine Wingless signalling controls self-renewal of Drosophila
intestinal stem cells. *Nature,* 455**,** 1119-1123.
LIN, G., XU, N. & XI, R. 2010. Paracrine unpaired signaling through the JAK/STAT pathway controls
self-renewal and lineage differentiation of drosophila intestinal stem cells. *J Mol Cell Biol,* 2**,**
37-49.
LIN, H. & SPRADLING, A. C. 1997. A novel group of pumilio mutations affects the asymmetric division
of germline stem cells in the Drosophila ovary. *Development,* 124**,** 2463-76.
LIN, Y., ZHENG, J. & LIN, D. 2021. PIWI-interacting RNAs in human cancer. *Seminars in Cancer*
Biology, 75**,** 15-28.
LIU, T., WANG, J., SUN, L., LI, M., HE, X., JIANG, J. & ZHOU, Q. 2021. Piwi-interacting RNA-651
promotes cell proliferation and migration and inhibits apoptosis in breast cancer by
facilitating DNMT1-mediated PTEN promoter methylation. *Cell Cycle,* 20**,** 1603-1616.
MA, X., ZHU, X., HAN, Y., STORY, B., DO, T., SONG, X., WANG, S., ZHANG, Y., BLANCHETTE, M.,
GOGOL, M., HALL, K., PEAK, A., ANOJA, P. & XIE, T. 2017. Aubergine Controls Germline Stem
Cell Self-Renewal and Progeny Differentiation via Distinct Mechanisms. *Developmental Cell,*
41**,** 157-169.e5.
MAJUMDAR, R., BANDYOPADHYAY, A. & MAITRA, U. 2003. Mammalian Translation Initiation Factor
eIF1 Functions with eIF1A and eIF3 in the Formation of a Stable 40 S Preinitiation Complex*.
Journal of Biological Chemistry, 278**,** 6580-6587.
MALONE, C. D., BRENNECKE, J., DUS, M., STARK, A., MCCOMBIE, W. R., SACHIDANANDAM, R. &
HANNON, G. J. 2009. Specialized piRNA Pathways Act in Germline and Somatic Tissues of the
Drosophila Ovary. *Cell,* 137**,** 522-535.
MARIANES, A. & SPRADLING, A. C. 2013. Physiological and stem cell compartmentalization within the
Drosophila midgut. *Elife,* 2**,** e00886.
MARKSTEIN, M., DETTORRE, S., CHO, J., NEUMÜLLER, R. A., CRAIG-MÜLLER, S. & PERRIMON, N.
2014. Systematic screen of chemotherapeutics in Drosophila stem cell tumors. *Proc Natl*
Acad Sci U S A, 111**,** 4530-5.
MARTINEZ, V. D., ENFIELD, K. S. S., ROWBOTHAM, D. A. & LAM, W. L. 2016. An atlas of gastric PIWI-
interacting RNA transcriptomes and their utility for identifying signatures of gastric cancer
recurrence. *Gastric Cancer,* 19**,** 660-665.
MARTINEZ, V. D., VUCIC, E. A., THU, K. L., HUBAUX, R., ENFIELD, K. S. S., PIKOR, L. A., BECKER-
SANTOS, D. D., BROWN, C. J., LAM, S. & LAM, W. L. 2015. Unique somatic and malignant

expression patterns implicate PIWI-interacting RNAs in cancer-type specific biology.
1461   *Scientific Reports,* 5, 10423.
1462 MENG, FANJU W. & BITEAU, B. 2015. A Sox Transcription Factor Is a Critical Regulator of Adult Stem
1463   Cell Proliferation in the Drosophila Intestine. *Cell Reports,* 13, 906-914.
1464 MICCHELLI, C. A. & PERRIMON, N. 2006. Evidence that stem cells reside in the adult Drosophila
1465   midgut epithelium. *Nature,* 439, 475-9.
1466 MOCHIZUKI, K., FINE, N. A., FUJISAWA, T. & GOROVSKY, M. A. 2002. Analysis of a piwi-related gene
1467   implicates small RNAs in genome rearrangement in tetrahymena. *Cell,* 110, 689-99.
1468 NISHIMASU, H., ISHIZU, H., SAITO, K., FUKUHARA, S., KAMATANI, M. K., BONNEFOND, L.,
1469   MATSUMOTO, N., NISHIZAWA, T., NAKANAGA, K., AOKI, J., ISHITANI, R., SIOMI, H., SIOMI, M.
1470   C. & NUREKI, O. 2012. Structure and function of Zucchini endoribonuclease in piRNA
1471   biogenesis. *Nature,* 491, 284-7.
1472 OBATA, F., TSUDA-SAKURAI, K., YAMAZAKI, T., NISHIO, R., NISHIMURA, K., KIMURA, M., FUNAKOSHI,
1473   M. & MIURA, M. 2018. Nutritional Control of Stem Cell Division through S-
1474   Adenosylmethionine in Drosophila Intestine. *Developmental Cell,* 44, 741-751.e3.
1475 OHARA, T., SAKAGUCHI, Y., SUZUKI, T., UEDA, H., MIYAUCHI, K. & SUZUKI, T. 2007. The 3' termini of
1476   mouse Piwi-interacting RNAs are 2'-O-methylated. *Nat Struct Mol Biol,* 14, 349-50.
1477 OHLSTEIN, B. & SPRADLING, A. 2006. The adult Drosophila posterior midgut is maintained by
1478   pluripotent stem cells. *Nature,* 439, 470-4.
1479 PAL-BHADRA, M., LEIBOVITCH, B. A., GANDHI, S. G., RAO, M., BHADRA, U., BIRCHLER, J. A. & ELGIN,
1480   S. C. R. 2004. Heterochromatic Silencing and HP1 Localization in *Drosophila* Are Dependent
1481   on the RNAi Machinery. *Science,* 303, 669-672.
1482 PANE, A., WEHR, K. & SCHÜPBACH, T. 2007. zucchini and squash encode two putative nucleases
1483   required for rasiRNA production in the Drosophila germline. *Dev Cell,* 12, 851-62.
1484 PEROCHON, J., YU, Y., AUGHEY, G. N., MEDINA, A. B., SOUTHALL, T. D. & CORDERO, J. B. 2021.
1485   Dynamic adult tracheal plasticity drives stem cell adaptation to changes in intestinal
1486   homeostasis in Drosophila. *Nat Cell Biol,* 23, 485-496.
1487 PETSAKOU, A. & PERRIMON, N. 2023. Bioelectric regulation of intestinal stem cells. *Trends Cell Biol,*
1488   33, 555-567.
1489 QIAO, D., ZEEMAN, A. M., DENG, W., LOOIJENGA, L. H. & LIN, H. 2002. Molecular characterization of
1490   hiwi, a human member of the piwi gene family whose overexpression is correlated to
1491   seminomas. *Oncogene,* 21, 3988-99.
1492 RAJASETHUPATHY, P., ANTONOV, I., SHERIDAN, R., FREY, S., SANDER, C., TUSCHL, T. & KANDEL,
1493   ERIC R. 2012. A Role for Neuronal piRNAs in the Epigenetic Control of Memory-Related
1494   Synaptic Plasticity. *Cell,* 149, 693-707.
1495 RAMAT, A., GARCIA-SILVA, M.-R., JAHAN, C., NAÏT-SAÏDI, R., DUFOURT, J., GARRET, C., CHARTIER, A.,
1496   CREMASCHI, J., PATEL, V., DECOURCELLE, M., BASTIDE, A., JUGE, F. & SIMONELIG, M. 2020.
1497   The PIWI protein Aubergine recruits eIF3 to activate translation in the germ plasm. *Cell*
1498   *Research,* 30, 421-435.
1499 RODE, S., OHM, H., ANHÄUSER, L., WAGNER, M., ROSING, M., DENG, X., SIN, O., LEIDEL, S. A.,
1500   STORKEBAUM, E., RENTMEISTER, A., ZHU, S. & RUMPF, S. 2018. Differential Requirement for
1501   Translation Initiation Factor Pathways during Ecdysone-Dependent Neuronal Remodeling in
1502   Drosophila. *Cell Rep,* 24, 2287-2299.e4.
1503 ROJAS-RÍOS, P., CHARTIER, A., ENJOLRAS, C., CREMASCHI, J., GARRET, C., BOUGHLITA, A., RAMAT, A.
1504   & SIMONELIG, M. 2023. piRNAs are regulators of metabolic reprogramming in stem cells.
1505   *bioRxiv,* 2023.10.31.564965.
1506 ROJAS-RÍOS, P., CHARTIER, A., PIERSON, S. & SIMONELIG, M. 2017. Aubergine and piRNAs promote
1507   germline stem cell self-renewal by repressing the proto-oncogene Cbl. *The EMBO Journal,*
1508   36, 3194-3211.
1509 ROSS, R. J., WEINER, M. M. & LIN, H. 2014. PIWI proteins and PIWI-interacting RNAs in the soma.
1510   *Nature,* 505, 353-359.

ROUGET, C., PAPIN, C., BOUREUX, A., MEUNIER, A.-C., FRANCO, B., ROBINE, N., LAI, E. C., PELISSON, A. & SIMONELIG, M. 2010. Maternal mRNA deadenylation and decay by the piRNA pathway in the early Drosophila embryo. *Nature,* 467, 1128-1132.

SAHIN, H. B., KARATAS, O. F., SPECCHIA, V., TOMMASO, S. D., DIEBOLD, C., BOZZETTI, M. P. & GIANGRANDE, A. 2016. Novel mutants of the aubergine gene. *Fly (Austin),* 10, 81-90.

SAITO, K., NISHIDA, K. M., MORI, T., KAWAMURA, Y., MIYOSHI, K., NAGAMI, T., SIOMI, H. & SIOMI, M. C. 2006. Specific association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the Drosophila genome. *Genes Dev,* 20, 2214-22.

SAITO, K., SAKAGUCHI, Y., SUZUKI, T., SUZUKI, T., SIOMI, H. & SIOMI, M. C. 2007. Pimet, the Drosophila homolog of HEN1, mediates 2'-O-methylation of Piwi-interacting RNAs at their 3' ends. *Genes Dev,* 21, 1603-8.

SCHATOFF, E. M., LEACH, B. I. & DOW, L. E. 2017. WNT Signaling and Colorectal Cancer. *Current Colorectal Cancer Reports,* 13, 101-110.

SCHÜPBACH, T. & WIESCHAUS, E. 1989. Female sterile mutations on the second chromosome of Drosophila melanogaster. I. Maternal effect mutations. *Genetics,* 121, 101-17.

SCHÜPBACH, T. & WIESCHAUS, E. 1991. Female sterile mutations on the second chromosome of Drosophila melanogaster. II. Mutations blocking oogenesis or altering egg morphology. *Genetics,* 129, 1119-36.

SEGDITSAS, S. & TOMLINSON, I. 2006. Colorectal cancer and genetic alterations in the Wnt pathway. *Oncogene,* 25, 7531-7.

SELLITTO, A., GELES, K., D'AGOSTINO, Y., CONTE, M., ALEXANDROVA, E., ROCCO, D., NASSA, G., GIURATO, G., TARALLO, R., WEISZ, A. & RIZZO, F. 2019. Molecular and Functional Characterization of the Somatic PIWIL1/piRNA Pathway in Colorectal Cancer Cells. *Cells,* 8, 1390.

SENTI, K. A., JURCZAK, D., SACHIDANANDAM, R. & BRENNECKE, J. 2015. piRNA-guided slicing of transposon transcripts enforces their transcriptional silencing via specifying the nuclear piRNA repertoire. *Genes Dev,* 29, 1747-62.

SHARMA, A. K., NELSON, M. C., BRANDT, J. E., WESSMAN, M., MAHMUD, N., WELLER, K. P. & HOFFMAN, R. 2001. Human CD34+ stem cells express the hiwigene, a human homologue of the Drosophila genepiwi. *Blood,* 97, 426-434.

SHI, S., YANG, Z.-Z., LIU, S., YANG, F. & LIN, H. 2020. PIWIL1 promotes gastric cancer via a piRNA-independent mechanism. *Proceedings of the National Academy of Sciences,* 117, 22390-22401.

SIDDIQI, S., TERRY, M. & MATUSHANSKY, I. 2012. Hiwi mediated tumorigenesis is associated with DNA hypermethylation. *PLoS One,* 7, e33711.

SIENSKI, G., BATKI, J., SENTI, K. A., DÖNERTAS, D., TIRIAN, L., MEIXNER, K. & BRENNECKE, J. 2015. Silencio/CG9754 connects the Piwi-piRNA complex to the cellular heterochromatin machinery. *Genes Dev,* 29, 2258-71.

SIOMI, M. C., SATO, K., PEZIC, D. & ARAVIN, A. A. 2011. PIWI-interacting small RNAs: the vanguard of genome defence. *Nature Reviews Molecular Cell Biology,* 12, 246-258.

SIUDEJA, K., VAN DEN BEEK, M., RIDDIFORD, N., BOUMARD, B., WURMSER, A., STEFANUTTI, M., LAMEIRAS, S. & BARDIN, A. J. 2021. Unraveling the features of somatic transposition in the Drosophila intestine. *Embo j,* 40, e106388.

SONG, P., YANG, F., JIN, H. & WANG, X. 2021. The regulation of protein translation and its implications for cancer. *Signal Transduction and Targeted Therapy,* 6, 68.

SOUSA-VICTOR, P., AYYAZ, A., HAYASHI, R., QI, Y., MADDEN, D. T., LUNYAK, V. V. & JASPER, H. 2017. Piwi Is Required to Limit Exhaustion of Aging Somatic Stem Cells. *Cell reports,* 20, 2527-2537.

SUN, R., GAO, C.-L., LI, D.-H., LI, B.-J. & DING, Y.-H. 2017. Expression Status of PIWIL1 as a Prognostic Marker of Colorectal Cancer. *Disease Markers,* 2017, 1204937.

TAMAMOUNA, V., RAHMAN, M. M., PETERSSON, M., CHARALAMBOUS, I., KUX, K., MAINOR, H., BOLENDER, V., ISBILIR, B., EDGAR, B. A. & PITSOULI, C. 2021. Remodelling of oxygen-

transporting tracheoles drives intestinal regeneration and tumorigenesis in Drosophila. *Nat*
Cell Biol, 23**,** 497-510.
TANG, X., LIU, N., QI, H. & LIN, H. 2023. Piwi maintains homeostasis in the Drosophila adult intestine.
Stem Cell Reports, 18**,** 503-518.
TOSAR, J. P., GARCÍA-SILVA, M. R. & CAYOTA, A. 2021. Circulating SNORD57 rather than piR-54265 is
a promising biomarker for colorectal cancer: common pitfalls in the study of somatic piRNAs
in cancer. *Rna,* 27**,** 403-410.
VAGIN, V. V., SIGOVA, A., LI, C., SEITZ, H., GVOZDEV, V. & ZAMORE, P. D. 2006. A distinct small RNA
pathway silences selfish genetic elements in the germline. *Science,* 313**,** 320-4.
VALÁSEK, L., MATHEW, A. A., SHIN, B. S., NIELSEN, K. H., SZAMECZ, B. & HINNEBUSCH, A. G. 2003.
The yeast eIF3 subunits TIF32/a, NIP1/c, and eIF5 make critical connections with the 40S
ribosome in vivo. *Genes Dev,* 17**,** 786-99.
VOUREKAS, A., ALEXIOU, P., VRETTOS, N., MARAGKAKIS, M. & MOURELATOS, Z. 2016. Sequence-
dependent but not sequence-specific piRNA adhesion traps mRNAs to the germ plasm.
Nature, 531**,** 390-394.
WAJED, S. A., LAIRD, P. W. & DEMEESTER, T. R. 2001. DNA methylation: an alternative pathway to
cancer. *Ann Surg,* 234**,** 10-20.
WALDRON, J. A., KANELLOS, G., SMITH, R. C. L., KNIGHT, J. R. P., MUNRO, J., ALEXANDROU, C.,
VLAHOV, N., PARDO-FERNANDEZ, L., MOORE, M., GILLEN, S. L., STRATHDEE, D., STEVENSON,
D., WARRANDER, F. C., GILROY, K., NIXON, C., CADDEN, B., POWLEY, I., OFFICER-JONES, L.,
BALLANTYNE, F., HAY, J., PENNEL, K., EDWARDS, J., CAMPBELL, A. D., RIDGWAY, R. A.,
COFFELT, S. B., NORMAN, J., QUESNE, J. L., BUSHELL, M. & SANSOM, O. J. 2023. eIF4A1 is
essential for reprogramming the translational landscape of Wnt-driven colorectal cancers.
*bioRxiv***,** 2023.11.10.566546.
WANG, H.-L., CHEN, B.-B., CAO, X.-G., WANG, J., HU, X.-F., MU, X.-Q. & CHEN, X.-B. 2015a. The
clinical significances of the abnormal expressions of Piwil1 and Piwil2 in colonic adenoma
and adenocarcinoma. *OncoTargets and Therapy,* 8**,** 1259-1264.
WANG, L., ZENG, X., RYOO, H. D. & JASPER, H. 2014. Integration of UPRER and Oxidative Stress
Signaling in the Control of Intestinal Stem Cell Proliferation. *PLOS Genetics,* 10**,** e1004568.
WANG, N., TAN, H.-Y., LU, Y., CHAN, Y.-T., WANG, D., GUO, W., XU, Y., ZHANG, C., CHEN, F., TANG, G.
& FENG, Y. 2021. PIWIL1 governs the crosstalk of cancer cell metabolism and
immunosuppressive microenvironment in hepatocellular carcinoma. *Signal Transduction and*
Targeted Therapy, 6**,** 86.
WANG, W., HAN, B. W., TIPPING, C., GE, D. T., ZHANG, Z., WENG, Z. & ZAMORE, P. D. 2015b. Slicing
and Binding by Ago3 or Aub Trigger Piwi-Bound piRNA Production by Distinct Mechanisms.
Mol Cell, 59**,** 819-30.
WANG, X., JIANG, C., FU, B., ZHU, R., DIAO, F., XU, N., CHEN, Z., TAO, W. & LI, C. J. 2015c. MILI, a PIWI
family protein, inhibits melanoma cell migration through methylation of LINE1. *Biochem*
Biophys Res Commun, 457**,** 514-9.
WANG, Y., JIANG, Y., MA, N., SANG, B., HU, X., CONG, X. & LIU, Z. 2015d. Overexpression of Hiwi
Inhibits the Growth and Migration of Chronic Myeloid Leukemia Cells. *Cell Biochem Biophys,*
73**,** 117-24.
WANG, Z., LIU, N., SHI, S., LIU, S. & LIN, H. 2016. The Role of PIWIL4, an Argonaute Family Protein, in
Breast Cancer*. *Journal of Biological Chemistry,* 291**,** 10646-10658.
WENG, W., LIU, N., TOIYAMA, Y., KUSUNOKI, M., NAGASAKA, T., FUJIWARA, T., WEI, Q., QIN, H., LIN,
H., MA, Y. & GOEL, A. 2018. Novel evidence for a PIWI-interacting RNA (piRNA) as an
oncogenic mediator of disease progression, and a potential prognostic biomarker in
colorectal cancer. *Mol Cancer,* 17**,** 16.
WIEGERING, A., UTHE, F. W., JAMIESON, T., RUOSS, Y., HÜTTENRAUCH, M., KÜSPERT, M., PFANN, C.,
NIXON, C., HEROLD, S., WALZ, S., TARANETS, L., GERMER, C. T., ROSENWALD, A., SANSOM,

1612    O. J. & EILERS, M. 2015. Targeting Translation Initiation Bypasses Signaling Crosstalk
1613    Mechanisms That Maintain High MYC Levels in Colorectal Cancer. *Cancer Discov,* 5, 768-781.
1614 WILSON, J. E., CONNELL, J. E. & MACDONALD, P. M. 1996. aubergine enhances oskar translation in
1615    the Drosophila ovary. *Development,* 122, 1631-9.
1616 XIE, K., ZHANG, K., KONG, J., WANG, C., GU, Y., LIANG, C., JIANG, T., QIN, N., LIU, J., GUO, X., HUO, R.,
1617    LIU, M., MA, H., DAI, J. & HU, Z. 2018. Cancer-testis gene PIWIL1 promotes cell proliferation,
1618    migration, and invasion in lung adenocarcinoma. *Cancer Medicine,* 7, 157-166.
1619 XU, N., WANG, S. Q., TAN, D., GAO, Y., LIN, G. & XI, R. 2011. EGFR, Wingless and JAK/STAT signaling
1620    cooperatively maintain Drosophila intestinal stem cells. *Dev Biol,* 354, 31-43.
1621 YAJIMA, M. & WESSEL, G. M. 2011. The multiple hats of Vasa: its functions in the germline and in cell
1622    cycle progression. *Mol Reprod Dev,* 78, 861-7.
1623 YIN, J., QI, W., JI, C. G., ZHANG, D. X., XIE, X. L., DING, Q., JIANG, X. Y., HAN, J. & JIANG, H. Q. 2019.
1624    Small RNA sequencing revealed aberrant piRNA expression profiles in colorectal cancer.
1625    *Oncol Rep,* 42, 263-272.
1626 ZENG, X., CHAUHAN, C. & HOU, S. X. 2010. Characterization of midgut stem cell- and enteroblast-
1627    specific Gal4 lines in drosophila. *Genesis (New York, N.Y. : 2000),* 48, 607-611.
1628 ZENG, X. & HOU, S. X. 2015. Enteroendocrine cells are generated from stem cells through a distinct
1629    progenitor in the adult Drosophila posterior midgut. *Development,* 142, 644-53.
1630 ZHANG, H., REN, Y., XU, H., PANG, D., DUAN, C. & LIU, C. 2013. The expression of stem cell protein
1631    Piwil2 and piR-932 in breast cancer. *Surgical Oncology,* 22, 217-223.
1632 ZHOU, F., RASMUSSEN, A., LEE, S. & AGAISSE, H. 2013. The UPD3 cytokine couples environmental
1633    challenge and intestinal stem cell division through modulation of JAK/STAT signaling in the
1634    stem cell microenvironment. *Dev Biol,* 373, 383-93.
1635

Fig.1

Fig.2

B Oxidized library (rep #2)
esg^ts>GFP Sucrose

TE AS, 1st nucleotides

Reads / endo siRNAs (a.u.)

siRNAs
piRNAs

nucleotide
- A (green)
- C (blue)
- G (orange)
- U (red)

20   25   30   35 /nt

TE S, 10th nucleotides

siRNAs
piRNAs

20   25   30   35 /nt

Probability

0.77
z = 17.3
+1

0.44
z = 7.9
+10

B' Oxidized library (rep #3)
esg^ts>GFP Pe

TE AS, 1st nucleotides

Reads / endo siRNAs

siRNAs
piRNAs

20   25   30   35 /nt

TE S, 10th nucleotides

siRNAs
piRNAs

20   25   30   35 /nt

Probability

0.78
z = 16.8
+1

0.43
z = 7.7
+10

C Oxidized library (rep #1)
esg^ts>GFP + aub^RNAi Sucrose

TE AS, 1st nucleotides

Reads / endo siRNAs (a.u.)

siRNAs
piRNAs

20   25   30   35 /nt

TE S, 10th nucleotides

siRNAs
piRNAs

20   25   30   35 /nt

Probability

0.77
z = 17.6
+1

0.44
z = 7.5
+10

C' Oxidized library (rep #1)
esg^ts>GFP + aub^RNAi *Pe*

TE AS, 1st nucleotides

Reads / endo siRNAs (a.u.)

siRNAs
piRNAs

20   25   30   35 /nt

TE S, 10th nucleotides

siRNAs
piRNAs

20   25   30   35 /nt

Probability

0.77
z = 17.2
+1

0.43
z = 7.3
+10

D piRNAs Antisense

Reads/endo siRNAs (a.u.)

ns
ns
ns
ns

- Sucrose (grey)
- *P.e* (blue)

esg^ts>GFP   *esg^ts>GFP + aub*^RNAi

E piRNAs Sense

Reads/endo siRNAs (a.u.)

ns
ns
ns
ns

- Sucrose (grey)
- *P.e* (blue)

esg^ts>GFP   *esg^ts>GFP + aub*^RNAi

F

RPKM / *esg^ts>GFP* ovaries (a.u.)

Cluster 38C Left

3.0

2.0

1.0

Ovaries aub^RNAi
ISCs/EBs Sucrose
ISCs/EBs P.e
ISCs/EBs aub^RNAi *Sucrose*
ISCs/EBs aub^RNAi *P.e*

Cluster 42AB

3.0

2.0

1.0

Ovaries aub^RNAi
ISCs/EBs Sucrose
ISCs/EBs P.e
ISCs/EBs aub^RNAi *Sucrose*
ISCs/EBs aub^RNAi *P.e*

Flamenco

1.5

1.0

0.5

Ovaries aub^RNAi
ISCs/EBs Sucrose
ISCs/EBs P.e
ISCs/EBs aub^RNAi *Sucrose*
ISCs/EBs aub^RNAi *P.e*

Fig.3

Fig.4

Fig.5

Fig.6

Fig.7

Fig.S1

Fig.S2

A

B — **Oxidized library (rep #3)** *w¹¹¹⁸* Sucrose

TE AS, 1st nucleotides | TE S, 10th nucleotides

siRNAs / piRNAs

nucleotide: A (green), C (blue), G (orange), U (red)

23 - 28 nt only

Reads / endo siRNAs (a.u.)

25 /nt

Probability

0.64 z = 10.7 | +1

0.35 z = 1.8 | +10

B' — **Oxidized library (rep #2)** *w¹¹¹⁸* Pe

TE AS, 1st nucleotides | TE S, 10th nucleotides

siRNAs / piRNAs

23 - 28 nt only

Reads / endo siRNAs

25 /nt

Probability

0.65 z = 13.7 | +1

0.37 z = 5.8 | +10

C — **Oxidized library (rep #2)** *aub* Sucrose

TE AS, 1st nucleotides | TE S, 10th nucleotides

siRNAs / piRNAs

23 - 28 nt only

Reads / endo siRNAs (a.u.)

25 /nt

Probability

0.57 z = 12.4 | +1

0.32 z = 1.2 | +10

C' — **Oxidized library (rep #3)** *aub* Pe

TE AS, 1st nucleotides | TE S, 10th nucleotides

siRNAs / piRNAs

23 - 28 nt only

Reads / endo siRNAs

25 /nt

Probability

0.49 z = 8.6 | +1

0.31 z = 1.1 | +10

D — **piRNAs Antisense**

Reads/endo siRNAs (a.u.)

ns

Sucrose / *P.e*

w^{1118} $aub^{HN2/QC42}$

E — **piRNAs Sense**

Reads/endo siRNAs (a.u.)

ns

Sucrose / *P.e*

w^{1118} $aub^{HN2/QC42}$

Fig.S3

Fig.S4

Fig.S5

Fig.S6