# ASYMMETRIC GENE FLOW MAINTAINS AN ABUNDANT RANGE CENTER

Zachary B. Hancock[*,1], Nicole E. Adams[1], Perry L. Wood Jr.[1], Gideon S. Bradburd[1]

[1]Department of Ecology & Evolutionary Biology, University of Michigan

[*]Corresponding author:

Zachary B. Hancock

Department of Ecology & Evolutionary Biology

University of Michigan

Ann Arbor, MI 48103

Email: hancockz@umich.edu

**Running title**: Asymmetric gene flow maintains an abundant range center

**ABSTRACT**

The abundant center hypothesis postulates that population density is the highest in the

center of the range and declines towards the periphery. While many evolutionary

hypotheses have been offered to explain what generates this pattern, ecologically it's

assumed the pattern emerges due to increasing distance from a species ideal niche

with edge populations occupying a "marginal" niche of low suitability. However, the

ubiquity of the abundant center pattern has recently been questioned. Furthermore, it

has proven difficult to assess both the pattern of the abundant center and the potential

generative scenarios (e.g., gene swamping, migration limitation) due to the inherent

complexity of species ranges. In this study, we evaluate the suitability of the abundant

center hypothesis to explain patterns of genetic diversity and inbreeding in shovel bugs

– small, sand-burrowing coastal crustaceans. In addition, we evaluate potential biases

in migration direction due to the dominant current regimes in the North Atlantic using a

common method for querying the site frequency spectrum, the directionality index ($\psi$).

While patterns of genetic diversity are consistent with an abundant center, we find a

strong signature of recent central colonization from the range periphery. Using

simulations, we evaluate the performance of $\psi$ under different migration regimes,

finding that, in the presence of asymmetric migration, it strongly infers an edge-to-center

colonization dynamic. We conclude that asymmetric migration towards the center can

lead to signatures of an abundant center without necessarily implying a reduction in

niche quality towards the range periphery.

**KEYWORDS:** abundant center hypothesis, range evolution, directionality index, migration, coastal, shovel bug, isolation-by-distance

**1 INTRODUCTION**

All species are restricted to finite ranges ultimately due to a failure to locally adapt along the periphery (Haldane 1956; Mayr 1963). Why populations fail to adapt and continue to expand their range, what is the long-term stability of range size, and how edges form and persist across a continuous environment are fundamental questions in the ecology and evolution of ranges (Kirkpatrick & Barton 1997; Sexton et al. 2009; Gaston 2003). Answering these questions has major implications for predicting species' responses to climate change (Doak & William 2010) as well as the propensity of invasive species to rapidly expand their range (Alexander & Edwards 2010).

A prime driver molding species' ranges is heterogeneity in population abundance, which has the potential to generate disparate evolutionary responses across populations. Grinnell (1922) envisioned a species range as an amoeboid organism with the center representing the "species metropolis," an area of high abundance and adaptiveness, with decreasing density towards the periphery. Along the edges of the range are "accidentals", occasional migrants that diffuse from the center like gas particles to the unsuitable fringe (Grinnell 1922). This concept of species ranges has since been formalized into the "abundant center" or "central marginal" hypothesis, which states that the region of the highest abundance is in the center of the

range and decreases monotonically towards the edge (Hengeveld & Haeck 1982;

Brown 1984). While intuitive, the generality of the abundant center hypothesis has

generated considerable debate (e.g., Dallas et al. 2020; Santini et al. 2019).

Ecologically, range limits are often conceptualized as the relationship between

the species' fundamental and realized niche, where the former represents the full

potential breadth of a species range whereas the latter reflects biotic and abiotic

constraints preventing full occupation of the fundamental niche (Hutchinson 1957,

1965). While appeals to a species' niche are intuitive for present distributions, often we

are curious about how they evolved in the first place and what forces are maintaining its

current bounds. These forces are fundamentally *evolutionary* in nature; niches are the

result of past adaptation; therefore, their limits should represent limits to a species

adaptive capacity. Haldane (1956) was among the first to suggest that population

replacement fails along the periphery due to the inability of edge populations to adapt

on account of their reduced size relative to central populations; genetic drift overwhelms

selection on the periphery. Mayr (1963) suggested this failure to adapt was due to the

influx of migrants from the range center, which carried with them alleles adapted to the

center but maladapted on the edge. This asymmetric migration was hypothesized to

swamp local adaptation, preventing edge populations from establishing. Kirkpatrick &

Barton (1997) formalized this "gene swamping" hypothesis by developing a quantitative

genetic model across environmental clines of varying steepness, demonstrating that

swamping could be a powerful mechanism maintaining species range edges. Thus,

gene swamping is a special case of the "migration load", the reduction in fitness of a

population due to the influx of migrant alleles maladapted to local conditions (Garcia-

Ramos & Kirkpatrick 1997; Bolnick & Nosil 2007).

However, in a large review of the literature, Kottler et al. (2021), find little

empirical evidence for gene swamping. Alternatively, they suggest that migration

limitation causes edge populations to suffer from a greater drift load and hence fail to

adapt; as opposed to gene swamping, an increase in migration is expected to rescue

edge populations, not doom them (Halbritter et al. 2015; Evans et al. 2014; Silva-Arias

et al. 2017). In this model, the rate at which stochastic extinction occurs in each patch

exceeds the rate at which migrants can colonize the patch. Migration limitation requires

that rates of dispersal decrease toward the edge, which implies some external barriers

exist along the edges but not, presumably, among central populations.

In the ocean, particularly for pelagic organisms, dispersal limitation may be a

product of the direction and strength of the dominant current regimes (Siegal et al.

2003). The average dispersal distance of a marine pelagic larva, for example, is a

function of the settling time and current velocity (Pringle & Wares 2007; Wares & Pringle

2008). Thus, marine ranges can be determined by complex abiotic factors that can

cause mass movement of individuals in asymmetric ways that may not be due to biases

in population abundance or reduction in niche quality. For example, in the North

Atlantic, a northward current, the Gulf Stream, emerges from the Loop Current in the

Gulf of Mexico, bringing warm water from the Caribbean, and travels up the coastline to

Chesapeake Bay before diverting to the mid-Atlantic (**Fig. 1A**). A second regime, the

Labrador Current, travels southward along Nova Scotia towards New England, bringing

with it cold Arctic water, before also diverting eastward. Hence, along the North Atlantic

coastline, two currents converge along the central latitude from opposite directions. For

a marine organism distributed along the entire coastline and whose dispersal is largely

dictated by the currents, this could generate a migration bias towards the center of the

range. Such an asymmetry in dispersal could theoretically generate an abundant center

pattern in the absence of reduced niche quality along the periphery if the current was

strong enough that successive colonization events became rarer and rarer at the range

extremes due to most individuals migrating downstream.

Determining the limits of ranges (e.g., Dallas et al. 2017; Pironon et al. 2017;

Soberón et al. 2018) and the underlying processes generating those limits remain

challenging (Kottler et al. 2021; Fristoe et al. 2023), especially in organisms not

amenable to transplant experiments. From a population genetic perspective, declines in

abundance along the periphery of ranges are often investigated using patterns of

genetic diversity (e.g., Ntuli et al. 2020; Dixon et al. 2013; Jin et al. 2020). Regions of

high abundance are expected to harbor the greatest diversity, whereas edge

populations are expected to have increased inbreeding (Haldane 1956; Arnaud-Haond

et al. 2006; De Ryck et al. 2016) and differentiation (Wright 1943; Eckert et al. 2008).

Furthermore, several techniques seek to detect asymmetries in the direction of

migration (Peter & Slatkin 2013, Beerli et al. 2019), whereas others model regions of the

range with high resistance to gene flow (Petkova et al. 2016; Marcus et al. 2021).

An expectation of the abundant center hypothesis is that edge populations may

be experiencing stochastic extinction and recolonization (in Grinnell's words, the

"accidentals" are expanding and contracting along the periphery; Grinnell 1922).

Theoretically, this might lead to a signature in the site frequency spectrum (SFS) of

recurrent range expansion in edge populations. Peter & Slatkin (2013; 2015) introduced

a method they call the "directionality index", $\psi$, which detects shifts in the frequency of

alleles across populations due to founder events in the expanding front. The logic

behind the index is that in large, stable populations there should be an abundance of

low frequency alleles, but subsequent founding events can cause those alleles to reach

high frequency via drift. Hence, alleles that are still segregating in the entire population

should have increasingly higher frequency in the direction of a range expansion. In the

case of repeated expansion along the edge in an abundant center range, the

directionality index should therefore detect a signature of higher frequency alleles on the

periphery relative to the center.

Small amphipod crustaceans called shovel bugs (Haustoriidae), which inhabit the

exposed sandy beaches of the western North Atlantic, are an excellent system in which

to test questions about range evolution. Due to their life history (see *Methods*), shovel

bugs are expected to be relatively poor dispersers, and this has been largely confirmed

in past studies (Jones 2005; Hancock et al. 2019). In the western North Atlantic, one

species of shovel bug (*Haustorius canadensis*, Bousfield 1962) can be found on

beaches from central Florida to southern Maine with some relict, isolated populations

reported in the Gulf of Saint Lawrence (Bousfield 1962; 1965). A curious feature of their

southernmost distribution is the abundance of suitable sandy beaches farther south

than they are found, with no shovel bug recorded south of Melbourne, Florida. The

northern range edge, alternatively, might represent a hard environmental transition, as

most of coastal Maine is composed of rocky outcrops with few fine sand beaches.

Given their relatively constrained niche — the intertidal of wave-exposed sandy

beaches — ease of identifying range boundaries, and the one-dimensional nature of

their range, shovel bugs are ideal organisms for evaluating hypotheses concerning

range evolution and for testing the abundant center hypothesis.

In this study, we evaluate whether the shovel bug *H. canadensis* in the western

North Atlantic constitutes a single, continuous population or contains discrete structure.

Next, we investigate whether patterns of genetic diversity and inbreeding are consistent

with expectations of the abundant center. Finally, we determine whether patterns of

migration and colonization are consistent with either gene swamping or dispersal

limitation as processes maintaining range edges. We find evidence for directional

colonization from the periphery of the ranges toward the center, which we hypothesize

is due to current regimes in the North Atlantic and consistent with the migration

limitation hypothesis.

**2 METHODS**

*2.1 Study system*

Deemed "the most interesting group of amphipods" by J. Laurens Barnard, shovel bugs

are haustoriid amphipods that are highly modified from their hypothesized gammarid

ancestor which include expanded articles on the 6th and 7th pereopods, extensive setal

elaboration, and spiny, cup-like structures on their 3rd and 4th pereopods that they

utilize as shovels (hence their common name; **Fig. 1A**) (Barnard 1969; Hancock &

Wicksten 2018). Shovel bugs are entirely fossorial, living in the first 1–10 cm of sand

and occupying almost exclusively the narrow swash zone (Croker & Hatfield 1980).

They lack a pelagic larva; instead, the females brood the juveniles through the first few

molts in their marsupium. Adult shovel bugs range in body size from 2–18 mm,

sometimes within the same species, and past work has found strong correlations

between body size and latitude (Hancock et al. 2021), though there may also be an

effect of sand grain size (LeCroy et al. 2002; Hancock & Wicksten 2018).

*2.2 Field collecting*

Shovel bugs were collected from 18 sites from Cocoa Beach, Florida, USA to Crescent

Beach, Maine, USA in May of 2023 (**Fig. 1A**). Collections began in West Palm Beach,

Florida, USA to identify the southernmost established population; no specimens were

located until Cocoa Beach, Florida. Similarly, sampling continued beyond Crescent

Beach, Maine, but no samples were located further north (last attempted location was

Popham Beach, ~93 km north). Specimens were sifted from sand using a 435 $\mu$m sieve

plate, plucked from the plate with forceps, and preserved in 95% ethanol. Individuals

were identified to species in the lab using a dissecting microscope and key from

Hancock & Wicksten (2018), and then stored at -20°C prior to DNA extraction.

*2.3 Molecular techniques*

Whole genomic DNA was extracted from either the whole individual or from pereopods

4–7 using Qiagen DNeasy Blood & Tissue Kit following manufacturer's protocols.

Library prep was performed using Restriction site Associated DNA sequencing (2RAD)

following Bayona-Vásquez et al. (2019), with slight modifications. All 96 samples were

quantified using a Qubit$^{TM}$ 1X dsDNA High Sensitivity (HS) kit on a Qubit 2.0

fluorometer. Samples were then normalized to have 100ng per sample (5 $\mu$L) on a 96

well plate, and then digested in a 15 $\mu$L solution with 1.5 $\mu$L 10X CutSmart Buffer (New

England Biolabs, USA), with 10 units of EcoRI (0.5 $\mu$L), 10 units MseI (0.5 $\mu$L), 3.5 $\mu$L

Invitrogen™ UltraPure™ DNase/RNase-Free Distilled Water, 2.5 $\mu$L of EcoRI

compatible adapter (2 $\mu$L), and 0.25 $\mu$M MseI compatible adapter ($\mu$L) on a Eppendorf

Mastercycler PRO S 6325 Thermal Cycler thermocycler at 37°C for one hour with the

heated lid turned off. Immediately after digestion we added 5 $\mu$L of ligation mastermix

composed of 2.75 $\mu$L of nuclease free water, 0.5 $\mu$L 10X ligation buffer (NEB), 1.5 $\mu$L 10

mM rATP (Promega), and 0.25 $\mu$L T4 DNA Ligase 400,000 U/ml to each sample. The

ligation reaction was implemented on the same thermocycler as above for two cycles of

22°C for 20 min, 37°C for 10 min, followed by 80°C for 20 min to stop the enzyme

activity. We then pulled 5 $\mu$L of each sample and split these into two microcenterfuge

tubes and performed a 1.25X SpeedBead cleanup (Rohland and Reich, 2012).

To reduce PCR duplicates and add the iTru5–8N barcode (which has a segment

of eight random base pairs) we implemented a single PCR cycle with six tubes each

with a 50 $\mu$L reaction. Each reaction contained 10 $\mu$L nuclease free water, 25 $\mu$L KAPA

HiFi hot start ready mix (Roche, USA), 5 $\mu$L iTru5–8N barcode (5 $\mu$M), and 10 $\mu$L of

library and were added using a single cycle of 98˚C for 1 min, 60˚C for 30s, and 72˚C

for 6 min. All six PCR reactions were then pooled and a 1.5X SpeedBead cleanup was

performed and resuspended in 30 $\mu$L of nuclease free water. Next, we split the library

into three 55 $\mu$L reactions with three unique iTru7 adapters. Each reaction had 10 $\mu$L of

nuclease free water, 25 $\mu$L KAPA HiFi hot start ready mix (Roche, USA), 5 $\mu$L P5 primer

(5 $\mu$M) and 5 $\mu$L of each iTru7 (5 $\mu$M) added separately. The PCR was implemented on

the thermocycler mentioned above under the following conditions: 95°C for 3 mins, 8

cycles of 98°C 20s, 60° 15s, 72°C 30s, and a final 72°C for 5 mins. All PCR reactions

were pooled and were clean using a 1.5X SpeedBead cleanup and resuspended in 30

$\mu$L of nuclease free water.

We performed size selection on the pooled library using a Pippin Prep (Sage

Science, USA) with internal standards on a 1.5% dye free cassette, target 350–480 bp

range (mean= 415 bp). After size selection to enrich the library, we performed a PCR

under the same conditions for the iTru7 reaction above, except with only four 50 $\mu$L

reactions and 6 cycles, also both the P5 and P7 primers (5 $\mu$M) were used. Finally, we

performed a 1X SpeedBead clean up and quantified the library on the Qubit 2.0 to

ensure that our library had a high enough concentration for sequencing. The library was

then sequenced on 12.5% percent of a NovaSeq X lane at the University of Michigan

Advanced Genomics Core.

Raw sequence reads were assembled and filtered using the de novo pipeline in

STACKS v.2.67 (Catchen et al. 2013; Rochette et al. 2019). A custom *process_radtags*

function was provided to us by J. Catchen to handle the unique FASTQ files of 2RAD,

which include an additional adapter sequence. Reads were assembled using the de

novo pipeline (Catchen et al. 2011); briefly, stacks were built in *ustacks* with a maximum

nucleotide distance allowed of 2 and minimum number of reads needed to seed a stack

of 3. Stacks were merged allowing a maximum of 2 gaps and 6 mismatches. Next, we

used the *populations* function, filtering out alleles that did not occur in at least 10

populations and keeping those that were present in 75% of individuals per population.

Sequences were output as two VCF files, one that included only variant sites and a

second that contained both variant and monomorphic sites. VCF files were additionally

filtered using *vcftools* v.1.16 (Danecek et al. 2011); first, individuals with >80% missing

data were removed. Next, we filtered out loci with more than two alleles (--max-alleles

2) and with a minor allele frequency of less than 0.05 (--maf 0.05). We then filtered out

loci with a minimum mean depth of less than 20 (--min-meanDP 20) and removed all

indels (--remove-indels). After filtering, we evaluated the relationship between mean

depth, estimated heterozygosity, and distance from the range center via linear

regression to determine if mean depth might be a confounding factor impacting the

detection of an abundant center.

*2.4 Population genetic analysis*

To evaluate potential patterns in genetic diversity, we estimated individual-level

heterozygosity using *pixy* (Korunes & Samuk 2021) on the VCF that included invariant

sites. The inbreeding coefficient ($F$; Wright, 1931), estimated as the difference between

the expected and observed homozygosity per individual, was estimated using *vcftools* (-

-het).

To test for geographic structure, we first used *pixy* to estimate population-level

pairwise $F_{ST}$ (Weir & Cockerham 1984), which we then linearized and regressed against

geographic distance to test for isolation-by-distance (Wright 1943; Rousset 1997). Next,

we performed a principal component analysis (PCA) to identify signs of genetic

clustering using *VCF2PCACluster* (He et al. 2024). Lastly, we performed a cluster

analysis using *fastSTRUCTURE* (Raj et al. 2014) with $k$ = 2–5. To determine whether

identified clusters and admixed individuals were merely a product of gapped sampling

or represented discrete populations, we performed a triangle plot analysis using

trianguIaR (Wiens & Colella 2024a). In the presence of discrete structure, individuals

with a hybrid index approaching 0.5 should have concomitantly high interclass

heterozygosity. However, a continuous population characterized solely by isolation-by-

distance will display a high hybrid index but no increase in interclass heterozygosity

(Wiens & Colella 2024b).

We evaluated potential barriers to migration using the program *FEEMS* (Marcus

et al. 2021), which estimates an effective migration surface using edge-specific

parameter estimation in a pseudo-likelihood framework. Lastly, we tested for a signature

of range expansion along the periphery of the population using Peter & Slatkin's (2013)

directionality index, $\psi$, as implemented in *rangeExpansion*

(https://github.com/BenjaminPeter/rangeexpansion). The directionality index is

calculated by comparing the difference in allele frequencies between pairwise

populations, conditioned on the allele being present in all populations. If the allele is not

lost by drift, then it is expected to be at higher frequency in founder populations than in

the source, because if founders carry the allele at all it will necessarily be at greater

frequency. Thus, successively more recently founded populations in an expanding

range will have increasingly positive values of $\psi$, permitting the identification of both the

origin and the direction of the range expansion (Peter & Slatkin 2013; 2015). In

populations at an isolation-by-distance equilibrium, simulations show that $\psi$ should be

zero. Therefore, we predict that if $\psi$ is zero, then there is no marked departure from

migration-drift equilibrium and hence no signature of reduced fitness or recurrent

colonization in edge populations. However, if dispersal limitation is driving edge

dynamics, we expect $\psi$ should be positive in the direction of the edge. We estimated the

hypothesized origin of the expansion both on the entire dataset and by subsetting it into

populations north and south of Chesapeake Bay.

*2.5 Simulations*

The direction of population expansion, as inferred from $\psi$, may be influenced by spatial

clines in population density even under scenarios with isotropic dispersal. To determine

whether the results of our $\psi$ analyses are robust to spatial variation in population

density (see Results/Discussion), we conducted forward time simulations using SLiM

v.4.0.1 (Haller & Messer 2023). We established a 10-deme stepping-stone model with

four different range scenarios: 1) equal population size per deme with symmetrical

migration; 2) equal population size per deme with asymmetrical migration from the edge

populations towards the center; 3) an abundant center model in which the central

demes had five times more individuals than the edge with symmetric migration; and 4)

abundant center with asymmetric migration from the edges. We performed 10 replicates

for each scenario. In addition, we performed simulations for the abundant center with

equal migration to evaluate how differences in population size between the central and

edge demes might drive patterns in $\psi$. We evaluated differences in population size of

5:1, 25:1, and 50:1, with 10 replicates each. Simulations were run for 100,000

generations (>10*N*) to ensure demographic equilibrium, and tree sequences were

recorded for subsequent processing using *pyslim* v.1.0.1 (Kelleher et al. 2018) and *tskit*

v.0.5.3 (Kelleher et al. 2018). For tree sequences with more than one root, we used

recapitation to simulate a neutral coalescent history until only a single root remained,

and then mutations were added at a rate of $1 \times 10^{-7}$ to the trees using *msprime* v.1.2.0

(Kelleher et al. 2016). Next, we randomly sampled five individuals per deme and output

their sequences in VCF format. From here, we trimmed the VCFs to only include

biallelic SNPs, and converted them to BED files using *plink* v.1.9 (Purcell et al. 2007).

Next, to use the *rangeExpansion* package, we converted the individual coordinates from

SLiM, which were in discrete demes, into Earth coordinates as required by

*rangeExpansion*; thus, we chose 10 coordinates 5° apart in latitude. Because the units

of distance are arbitrary in SLiM, we needed only to ensure that deme 1 is closest to

deme 2, etc., to adequately capture the SLiM environment in an Earth coordinate

system. We then calculated $\psi$ for each simulation to see how the different range

scenarios might cause bias inferences made from $\psi$.

**3 RESULTS**

*3.1 ddRAD assembly*

A total of 2,605,149 loci passed filtering in STACKS with a mean effective per-sample

coverage of 23.0, a standard deviation of 13.6, with a minimum of 4.0 and maximum of

67.4. The mean number of sites per locus was 220.1. After filtering for biallelic SNPs

and missingness in *vcftools*, the final dataset included 7,405 SNPs. We find a significant

positive relationship between mean depth per individual and our estimates of

heterozygosity from *pixy* ($P < 5.3e–7$, $R^2 = 0.27$; *Table S1*); in addition, we find a

significant negative relationship between mean depth and the distance from the center

of the range (taken as 36° latitude; $P < 0.002$, $R^2 = 0.10$) (see *Table S1*). To account for

the potential confounding between measures of heterozygosity, mean depth, and their

relationship with the distance from the range center, all subsequent multivariate linear

regressions are performed with mean depth as a confounding variable. In addition, we

tested for a significant correlation after dropping all individuals with a mean depth

greater than 30 to determine if a few individuals with high depth were driving the

patterns. While a significant correlation between heterozygosity and mean depth

remained ($P < 2.7e–5$; $R^2 = 0.38$), it was no longer significantly correlated with distance

from the center ($P = 0.57$; $R^2 = –0.02$) (*Fig. S1*).

*3.2 Genetic structure*

Results of the *fastSTRUCTURE* analysis supported three population clusters ($k = 3$) as

the model that maximized the marginal likelihood – one from Florida to South Carolina,

a second from North Carolina to Virginia, and a third that included the remaining

populations from Maryland to Maine (*Fig. S2*). However, the PCA reveals relatively

continuous structure, with a possible break between populations north and south of the

Chesapeake (**Fig. 1B**). A discrete pattern of ancestry was also not supported by the

triangle plot analysis, which found that, despite increasing hybrid ancestry towards

Chesapeake Bay, interclass heterozygosity did not increase dramatically (*Fig. S3*).

Furthermore, we found no discrete breaks in patterns of isolation-by-distance when

regressing linearized $F_{ST}$ against geographic distance between comparisons within

southern and northern populations and between them (**Fig. 1C**). Subsequent analyses

therefore treat the shovel bug population as if it is one continuous population without

discrete breaks.

*3.3 Patterns of diversity*

The patterns of genetic diversity and inbreeding we inferred are strongly indicative of an abundant center. After incorporating sequencing depth as a possible confounding effect, measures of individual heterozygosity are significantly predicted by the distance from the center of the range ($P < 3.79e{-}16$; mean depth $P = 5.1e{-}5$). This relationship remained significant after dropping individuals with mean depth greater than 30 ($P < 0.0009$; $R^2 = 0.25$) (*Fig. S4*). Similarly, measures of inbreeding are the highest at the range edges, declining as a function of distance from the center (**Fig. 2B**). These patterns are consistent with field observations, in which the ease of collecting and observed highest abundance was in the central range locations, with shovel bugs at the edge sample locations being rarer and thus more difficult to collect. However, we note that while distance is a much stronger predictor of patterns of genetic diversity and inbreeding, mean sequencing depth was still a significant predictor in each model (*Table S1*).

The effective migration surface estimated from *FEEMS* suggests the center of the range has the highest ease of movement, with reduced effective migration towards the range edges (*Fig. S5*). However, in accordance with results from *fastSTRUCTURE* and PCA, the migration surface indicates a strong depression in effective migration across the Chesapeake, which might suggest it acts as a partial barrier to gene flow.

*3.4 Directionality index & simulations*

The inferred origin of the range expansion of the entire sample of individuals, as

indicated by the lowest value of $\psi$, was 28.3° latitude in northern Florida (**Fig. 3A**).

When the data were subset into two regions, south and north (based on the cluster

analysis), the inferred origins were at 28.3° and 43.5° latitude, respectively. The sum of

$\psi$ for each population resulted in a pattern of increasing $\psi$ towards the range center,

mirroring the increase in heterozygosity (**Fig. 3A**). This pattern of $\psi$ superficially

suggests a range expansion toward the center of the range from the edges following the

dominant current regimes of the Gulf Stream running south to north and the Labrador

Current running north to south, each converging in the center of the range (**Fig. 1A,**

**3A**). This pattern of expansion is in opposition to expectations from the gene swamping

hypothesis and suggests that asymmetric migration due to the currents might be

maintaining range edges, which would be consistent with the migration limitation

hypothesis.

Simulation results from SLiM support this interpretation. Across replicates, the

strongest divergences from $\psi = 0$, which is expected at migration-drift equilibrium, occur

in simulations with asymmetric migration from the edges toward the center, with the

most striking inflation in the scenario in which there is both an abundant center and

asymmetric migration (**Fig. 3B**). While there is some deviation from zero along the

edges in scenarios with symmetric migration in both the equal and center-abundant

scenarios, they do not systematically produce a positive $\psi$ in the center as in the case

of asymmetric migration (**Fig. 3B**). Furthermore, increasing the population size

discrepancy between the center and the edge does not consistently lead to an increase

in central estimates of $\psi$ (*Fig. S6*), indicating that asymmetric migration, not increased

drift along the edges, is the dominant force shaping the observed dynamics.

**4 DISCUSSION**

The persistence of range edges despite the potential for local adaptation is an enduring

evolutionary mystery. Several hypotheses have been proposed to explain the finite

nature of species ranges, principle among these being gene swamping (Mayr 1963;

Kirkpatrick & Barton 1996) and dispersal limitation (Sexton et al. 2009; Hargraves et al.

2014). Each of these are expected to generate a pattern of an abundant range center in

which fitness, population density, migration, or some combination of the three, decline

as a function of distance from the center.

In this study, we investigated the range evolution of shovel bugs (*Haustorius*

*canadensis*; Bousfield 1962) distributed along the Eastern Seaboard of North America.

Past work has suggested that shovel bugs have the potential for large amounts of

cryptic diversity and are sensitive to environmental barriers (Jones 2005; Hancock et al.

2019), so our first aim was to establish that shovel bugs in the North Atlantic represent a

single, continuous population. Given that patterns of linearized $F_{ST}$ were consistent with

a continuous population and the triangle plot analyses did not reveal a strong

relationship with interclass heterozygosity and the hybrid index, we tentatively conclude

that shovel bugs in the North Atlantic represent a single, continuous population. We

identify reduced effective migration on the range periphery consistent with the dispersal

limitation hypothesis (*Fig. S5*); however, we also find evidence for resistance to

migration across the Chesapeake Bay. Shovel bugs of the genus *Haustorius* specialize

on open, wave-exposed beaches in which the sand is highly oxygenated (Bousfield

1970), and rarely occur in low salinity, brackish bays. While the Chesapeake Bay was

formed by an impact crater during the Eocene (Ivany et al. 2000), and this event has

been postulated to have driven haustoriid diversification in North America (Hancock et

al. 2022), its recent form as a large bay is young, filling in after the end of the Last

Glacial Maximum around 10,000 years ago (Colman & Mixon 1988). The bay has the

potential to be acting as a partial barrier to gene flow, and there is some support for this

from the PCA and *fastSTRUCTURE* analyses. However, the triangle plot and isolation-

by-distance regressions did not support discrete breaks (**Fig. 1C**; *Fig. S3*). This might

indicate that the barrier is weak or has occurred recently enough that it has not yet

influenced genome-wide patterns of allele frequency divergence, which can lag behind

other population summary statistics (Barton et al. 2013; Ringbauer et al. 2018).

Once the genetic structure of shovel bugs was characterized, our next aim was

to identify patterns of genetic diversity and inbreeding to determine if they were

consistent with the abundant center hypothesis. The abundant center hypothesis has

mostly been tested in terrestrial organisms; in the few instances in which it has been

tested in marine and coastal taxa, it has had mixed support. Rivadeneira et al. (2010)

found two of the five porcelain crab (Porcellanidae) species analyzed along the Chilean

coastline showed patterns of an abundant center, while the other three either showed

no pattern or greater diversity along an edge. Samis & Eckhert (2007), evaluating the

abundance and production of two species of dune flower, found no evidence for an

abundant center in one (pink sand verbena) while the other (beach evening primrose)

had the highest seed production in the center of the range. In this study, patterns of

genetic diversity and inbreeding strongly support the hypothesis of an abundant range

center (**Fig. 2**). We also observed higher densities of shovel bugs in the Carolinas and

southern New Jersey, and ease of collection decreased towards the range periphery.

Future work would benefit from quantitative estimates of census abundance across this

shovel bug's range. In the northern part of the range, the decline in density

corresponded with increasingly rocky beaches and, eventually, the replacement of

beaches entirely with sheer cliffs sculpted by the retreating Laurentide Ice Sheet. In

their southernmost range edge, however, suitable sandy beach habitat exists far

beyond their occurrence (LeCroy 2002; Hancock & Wicksten 2018). Hargraves et al.

(2013) propose that when the range limit is less than the niche limit, it is due either to

the recency of niche expansion relative to dispersal potential, the difference in the rate

of local extinction to migration, or fluctuations in suitability that exceed the rate of

recolonization. Each of these explanations rely on reductions in dispersal on the range

edge relative to the center, which our *FEEMS* analysis supports (*Fig. S5*).

If rates of recolonization of peripheral populations are less than that of stochastic

extinction due to environmental or demographic fluctuations, edge populations might

harbor signatures of repeated range expansions. Given that we found evidence of both

an abundant center and reduced migration along the periphery, we sought to test for

skews in the SFS consistent with an expansion (Peter & Slatkin 2013; 2015). However,

skews in the SFS were in opposite directions of what we expected; we detected a linear

increase in $\psi$ towards the range center from each edge, which implies a range

expansion from the periphery towards the center. This is contradictory to patterns of

inbreeding and genetic diversity, which clearly indicate that abundance is the highest in

the range center (**Fig. 2**). A possible explanation to these conflicting patterns is that the

$\psi$ index is biased in the presence of asymmetric dispersal.

As discussed in the Introduction, the dominant current regimes in the North

Atlantic flow in opposite directions, converging at a central latitude in the Chesapeake

Bay region (**Fig. 1A**). If dispersal is mostly passive in shovel bugs, we might expect a

net direction of movement towards the range center from the edges. Thus, asymmetric

dispersal may be responsible for generating an abundant center as it has the potential

to be acting as a population sink. In addition, because migrants are being pulled

towards the center, this could affect the shape of the SFS such that the signal is

consistent with recurrent colonization in the direction of the Chesapeake.

To evaluate whether our intuition about the impact of asymmetric migration on

the $\psi$ index was correct, we conducted simulations intended to tease apart the impacts

of differences in population size and migration (**Fig. 3B**; *Fig. S6*). Consistent with our

explanation above, we find that $\psi$ is highly sensitive to asymmetric migration and will

infer a population expansion in the direction of the range center even in equilibrium

scenarios. In addition, this bias is strongest when under the combined influence of

asymmetric migration and abundant center dynamics (**Fig. 3B**), but, importantly, does

not occur in the presence of differences in population size alone (*Fig. S6*). This

suggests that asymmetric migration may be playing the dominant role in shaping

patterns in the SFS that are being identified by the $\psi$ index.

The $\psi$ index is a popular statistic for testing for range expansions and has been

applied to a diverse array of different organisms (Manuel et al. 2016; Zhan et al. 2014;

Fifer et al. 2022). In this study, we uncover a potential confounding influence on $\psi$ that

is likely to impact any organism with high asymmetric migration. While asymmetric gene

flow has been identified in terrestrial organisms (Fedorka et al. 2012; Oswald et al.

2017), the taxa most likely to display directional dispersal are aquatic organisms with

passive dispersal via river flow or oceanic currents (Siegal et al. 2003; Pringle & Wares

2007; Morrissey & de Kerckhove 2009). Past work has shown that directional dispersal

in the ocean can mislead many population genetic analyses, including the inference of

random dispersal via patterns of isolation-by-distance and the probability of

coalescence (Nagylaki 1977; Lenormand 2002; Paz-Vinas et al. 2013). Using a

simulation approach, we uncover that asymmetric migration can also bias the detection

of a range expansion, potentially generating a false positive signature when, in fact, the

population is in demographic equilibrium.

In addition to the impact asymmetric migration has on the inference of a range

expansion, certain biased dispersal patterns may themselves be capable of generating

an abundant center. Specifically, asymmetric migration from the edges towards the

range center can maintain high abundance in the center, causing it to act like a

population sink, while colonization towards increasingly distant patches becomes less

likely as the number of available migrants dispersing away from the center becomes

increasingly small. This is a kind of migration limitation (Sexton et al. 2009; Hargraves et

al. 2013), in which the rate of extinction in peripheral patches exceeds that of

colonization, and the cause of this decreased migration is asymmetries in the probability

of migration. This hypothesis is in opposition to gene swamping, in which increased

rates of migration to the edge is detrimental (Kirkpatrick & Barton 1996).

The generality of the abundant center hypothesis remains hotly contested (Dallas

et al. 2020; Santini et al. 2019; Sagarin et al. 2006), as do the evolutionary processes

that give rise to them (Kottler et al. 2023). We uncover strong evidence for an abundant

center in shovel bugs in the western North Atlantic and strong support for the

hypothesis that this pattern is maintained by a combination of passive dispersal and

directional currents promoting gene flow towards the center of the range. We suggest

this mechanism for generating an abundant center might be widespread in coastal

marine taxa, especially those with passive dispersal, as oceanic currents regularly

converge within species ranges. Furthermore, focusing on asymmetric migration via

directional water or wind currents may help clarify what maintains range edges in cases

where the fundamental niche appears to exceed the range limit.

**DATA AVAILABILITY**

Bioinformatic pipelines, including filtered VCFs, are available at

https://github.com/zachbhancock/asym_gene_flow_proj. Raw sequences were

deposited to GENBANK.

**AUTHOR CONTRIBUTIONS**

ZBH and GSB conceptualized and designed the study. ZBH carried out the field

collections. ZBH and PLW performed the laboratory work. ZBH and NEA carried out the

bioinformatics and analyzed the data. All authors wrote and corrected the manuscript.

**ACKNOWLEDGEMENTS**

Thanks to members of the Bradburd lab who provided helpful comments on this

manuscript. Thanks also to Kristen Wacker and Teresa Pegan for thoughtful discussion

and feedback. We are all indebted to Leonard Jones who coined the common name

"shovel bug" for haustoriid amphipods, and proudly use it in print for the first time here.

May it catch on. Lastly, special thanks to Daniel Stern Cardinale and Robert Joel Duff

for their hospitality and conversations while ZBH was collecting.

**REFERENCES**

Alexander, J. M., & Edwards, P. J. (2010). Limits to the niche and range margins of alien
species. *Oikos*, *119*(9), 1377-1386.
554
555    ARNAUD-HAOND, S., Teixeira, S., Massa, S. I., Billot, C., Saenger, P., Coupland, G., Duarte, C.M., &
556    Serrao, E. A. (2006). Genetic structure at range edge: low diversity and high inbreeding in Southeast
557    Asian mangrove (Avicennia marina) populations. *Molecular Ecology*, *15*(12), 3515-3525.
558
559    Barnard, J.L. (1969) *The Families and Genera of Marine Gammaridean Amphipoda*. Smithsonian
560    Institution Press, Washington, D.C., 535 pp.
561
562    Barton, N. H., Etheridge, A. M., Kelleher, J., & Véber, A. (2013). Genetic hitchhiking in spatially extended
563    populations. *Theoretical population biology*, *87*, 75-89.
564
565    Bayona-Vásquez, N.J., Glenn, T.C., Kieran, T.J., Pierson, T.W., Hoffberg, S.L., Scott, P.A., Bentley, K.E.,
566    Finger, J.W., Louha, S., Troendle, N., Diaz-Jaimes, P., Mauricio, R., Faircloth, B.C. (2019). Adapterama
567    III: Quadruple-indexed, double/triple-enzyme RADseq libraries (2RAD/3RAD). *PeerJ*, 7:e7724
568    http://doi.org/10.7717/peerj.7724.
569
570    Beerli, P., Mashayekhi, S., Sadeghi, M., Khodaei, M., & Shaw, K. (2019). Population genetic inference
571    with MIGRATE. *Current protocols in bioinformatics*, *68*(1), e87.
572
573    Bolnick, D. I., & Nosil, P. (2007). Natural selection in populations subject to a migration
574    load. *Evolution*, *61*(9), 2229-2243.
575
576    Bousfield, E. L. (1962). *New Haustoriid Amphipods from the Canadian Atlantic Region*. Canada
577    Department of Northern Affairs and National Resources.
578
579    Bousfield, E. L. (1965). Haustoriidae of New England (Crustacea: Amphipoda). *Proceedings of the United*
580    *States National Museum*.
581
Bousfield, E. L. (1970). Adaptive radiation in sand-burrowing amphipod crustaceans. *Chesapeake*
*Science*, *11*(3), 143-154.

Brown, J. H. (1984). On the relationship between abundance and distribution of species. *The American*
*Naturalist*, *124*(2), 255-279.

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks: building and
genotyping loci de novo from short-read sequences. *G3: Genes, Genomes, Genetics*, *1*(3), 171-182.

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool
set for population genomics. *Molecular Ecology*, *22*(11), 3124-3140.

Colman, S. M., & Mixon, R. B. (1988). The record of major Quaternary sea-level changes in a large
coastal plain estuary, Chesapeake Bay, eastern United States. *Palaeogeography, Palaeoclimatology,*
*Palaeoecology*, *68*(2-4), 99-116.

Croker, R. A., & Hatfield, E. B. (1980). Space partitioning and interactions in an intertidal sand-burrowing
amphipod guild. *Marine Biology*, *61*, 79-88.

Dallas, T., Decker, R. R., & Hastings, A. (2017). Species are not most abundant in the centre of their
geographic range or climatic niche. *Ecology Letters*, *20*(12), 1526-1533.

Dallas, T. A., Santini, L., Decker, R., & Hastings, A. (2020). Weighing the evidence for the abundant-
centre hypothesis. *Biodiversity Informatics*, *15*(3), 81-91.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.E., Lunter,
G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., & 1000 Genomes Project Analysis Group. (2011).
The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156-2158.

De Ryck, D. J., Koedam, N., Van der Stocken, T., van der Ven, R. M., Adams, J., & Triest, L. (2016).
Dispersal limitation of the mangrove Avicennia marina at its South African range limit in strong contrast to
connectivity in its core East African region. *Marine Ecology Progress Series*, *545*, 123-134.

Dixon, A. L., Herlihy, C. R., & Busch, J. W. (2013). Demographic and population-genetic tests provide
mixed support for the abundant centre hypothesis in the endemic plant Leavenworthia stylosa. *Molecular*
*ecology*, *22*(7), 1777-1791.

Doak, D. F., & Morris, W. F. (2010). Demographic compensation and tipping points in climate-induced
range shifts. *Nature*, *467*(7318), 959-962.

Eckert, C. G., Samis, K. E., & Lougheed, S. C. (2008). Genetic variation across species' geographical
ranges: the central–marginal hypothesis and beyond. *Molecular Ecology*, *17*(5), 1170-1188.

Evans, S. M., Sinclair, E. A., Poore, A. G., Steinberg, P. D., Kendrick, G. A., & Vergés, A. (2014). Genetic
diversity in threatened Posidonia australis seagrass meadows. *Conservation Genetics*, *15*, 717-728.

Fedorka, K. M., Winterhalter, W. E., Shaw, K. L., Brogan, W. R., & Mousseau, T. A. (2012). The role of
gene flow asymmetry along an environmental gradient in constraining local adaptation and range
expansion. *Journal of Evolutionary Biology*, *25*(8), 1676-1685.

Fifer, J. E., Yasuda, N., Yamakita, T., Bove, C. B., & Davies, S. W. (2022). Genetic divergence and range
expansion in a western North Pacific coral. *Science of the Total Environment*, *813*, 152423.
634

635    Fristoe, T. S., Vilela, B., Brown, J. H., & Botero, C. A. (2023). Abundant-core thinking clarifies exceptions
636    to the abundant-center distribution pattern. *Ecography*, *2023*(2), e06365.
637

638    García-Ramos, G., & Kirkpatrick, M. (1997). Genetic models of adaptation and gene flow in peripheral
639    populations. *Evolution*, *51*(1), 21-28.
640

641    Gaston, K. J. (2009). Geographic range limits of species. *Proceedings of the Royal Society B: Biological*
642    *Sciences*, *276*(1661), 1391-1393.
643

644    Grinnell, J. (1922). The role of the" accidental". *The Auk*, *39*(3), 373-380.
645

646    Halbritter, A. H., Billeter, R., Edwards, P. J., & Alexander, J. M. (2015). Local adaptation at range edges:
647    comparing elevation and latitudinal gradients. *Journal of Evolutionary Biology*, *28*(10), 1849-1860.
648

649    Haldane, J. B. S. (1956). The relation between density regulation and natural selection. *Proceedings of*
650    *the Royal Society of London. Series B-Biological Sciences*, *145*(920), 306-308.
651

652    Haller, B. C., & Messer, P. W. (2023). SLiM 4: multispecies eco-evolutionary modeling. *The American*
653    *Naturalist*, *201*(5), E127-E139.
654

655    Hancock, Z. B., & Wicksten, M. K. (2018). Two new species of sand-burrowing amphipods of the genus
656    Haustorius Müller, 1775 (Amphipoda: Haustoriidae) from the northwestern Gulf of
657    Mexico. *Zootaxa*, *4459*(1), 101-127.
658

659    Hancock, Z. B., Hardin, F. O., & Light, J. E. (2019). Phylogeography of sand-burrowing amphipods
660    (Haustoriidae) supports an ancient suture zone in the Gulf of Mexico. *Journal of Biogeography*, *46*(11),
661    2532-2547.
662

663    Hancock, Z. B., Hardin, F. O., Murthy, A., Hillhouse, A., & Johnston, J. S. (2021). Rapid genomic
664    expansion and purging associated with habitat transitions in a clade of beach crustaceans (Amphipoda:
665    Haustoriidae). *Journal of Crustacean Biology*, *41*(3), ruab042.
666

667    Hancock, Z. B., Ogawa, H., Light, J. E., & Wicksten, M. K. (2022). Origin and evolution of the
668    Haustoriidae (Amphipoda): a eulogy for the Haustoriidira. *Zoological Journal of the Linnean*
669    *Society*, *194*(4), 1252-1267.
670

671    He, W., Xu, L., Wang, J., Yue, Z., Jing, Y., Tai, S., Yang,J., & Fang, X. (2024). VCF2PCACluster: a
672    simple, fast and memory-efficient tool for principal component analysis of tens of millions of SNPs. *BMC*
673    *Bioinformatics*, *25*(1), 173.
674

675    Hengeveld, R., & Haeck, J. (1982). The distribution of abundance. I. Measurements. *Journal of*
676    *Biogeography*, 303-316.
677

678    Hutchinson, G.E. (1957). Concluding remarks. Cold Spring Harbor Symposia on Quantitative Biology. 22
679    (2): 415–427.
680

681    Hutchinson, G.E. (1965). *The Ecological Theater and Evolutionary Play*. New Haven: Yale University
682    Press.
683
684    Ivany, L. C., Patterson, W. P., & Lohmann, K. C. (2000). Cooler winters as a possible cause of mass
685    extinctions at the Eocene/Oligocene boundary. *Nature*, *407*(6806), 887-890.
686
687    Jin, P. Y., Sun, J. T., Chen, L., Xue, X. F., & Hong, X. Y. (2020). Geography alone cannot explain
688    *Tetranychus truncatus* (Acari: Tetranychidae) population abundance and genetic diversity in the context
689    of the center–periphery hypothesis. *Heredity*, *124*(2), 383-396.
690
691    Jones, M.E. (2005) *Comparative Phylogeography, Population Structure, and Extended Maternal Care in
692    Three Intertidal Haustoriid Amphipods from the Western North Atlantic*. Unpublished Master's thesis,
693    College of Charleston.
694
695    Kelleher, J., Etheridge, A. M., & McVean, G. (2016). Efficient coalescent simulation and genealogical
696    analysis for large sample sizes. *PLoS Computational Biology*, *12*(5), e1004842.
697
698    Kelleher, J., Thornton, K. R., Ashander, J., & Ralph, P. L. (2018). Efficient pedigree recording for fast
699    population genetics simulation. *PLoS Computational Biology*, *14*(11), e1006581.
700
701    Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a species' range. *The American Naturalist*, *150*(1), 1-
702    23.
703
704    Kottler, E. J., Dickman, E. E., Sexton, J. P., Emery, N. C., & Franks, S. J. (2021). Draining the swamping
705    hypothesis: little evidence that gene flow reduces fitness at range edges. *Trends in Ecology &
706    Evolution*, *36*(6), 533-544.
707
708    Korunes, K. L., & Samuk, K. (2021). pixy: Unbiased estimation of nucleotide diversity and divergence in
709    the presence of missing data. *Molecular Ecology Resources*, *21*(4), 1359-1368.
710
711    LeCroy, S. E., Richardson, J. S., & Cobb, D. (2000). *An illustrated identification guide to the nearshore
712    marine and estuarine gammaridean Amphipoda of Florida* (Vol. 1, p. 2000). Florida Department of
713    Environmental Protection, Division of Resource Assessment and Management, Bureau of Laboratories.
714
715    Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, *17*(4),
716    183-189.
717
718    De Manuel, M., Kuhlwilm, M., Frandsen, P., Sousa, V. C., Desai, T., Prado-Martinez, J., et al. (2016).
719    Chimpanzee genomic diversity reveals ancient admixture with bonobos. *Science*, *354*(6311), 477-481.
720
721    Marcus, J., Ha, W., Barber, R. F., & Novembre, J. (2021). Fast and flexible estimation of effective
722    migration surfaces. *Elife*, *10*, e61927.
723
724    Mayr, E. (1963). *Animal species and evolution*. Harvard University Press.
725
726    Morrissey, M. B., & de Kerckhove, D. T. (2009). The maintenance of genetic variation due to asymmetric
727    gene flow in dendritic metapopulations. *The American Naturalist*, *174*(6), 875-889.
728

729    Nagylaki, T. (1977). Decay of genetic variability in geographically structured populations. *Proceedings of*
730    *the National Academy of Sciences*, *74*(6), 2523-2525.
731
732    Ntuli, N. N., Nicastro, K. R., Zardi, G. I., Assis, J., McQuaid, C. D., & Teske, P. R. (2020). Rejection of the
733    genetic implications of the "Abundant Centre Hypothesis" in marine mussels. *Scientific Reports*, *10*(1),
734    604.
735
736    Oswald, J. A., Overcast, I., Mauck III, W. M., Andersen, M. J., & Smith, B. T. (2017). Isolation with
737    asymmetric gene flow during the nonsynchronous divergence of dry forest birds. *Molecular*
738    *Ecology*, *26*(5), 1386-1400.
739
740    Paz-Vinas, I., Quéméré, E., Chikhi, L., Loot, G., & Blanchet, S. (2013). The demographic history of
741    populations experiencing asymmetric gene flow: combining simulated and empirical data. *Molecular*
742    *Ecology*, *22*(12), 3279-3291.
743
744    Peter, B. M., & Slatkin, M. (2013). Detecting range expansions from genetic data. *Evolution*, *67*(11),
745    3274-3289.
746
747    Peter, B. M., & Slatkin, M. (2015). The effective founder effect in a spatially expanding
748    population. *Evolution*, *69*(3), 721-734.
749
750    Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population structure with estimated
751    effective migration surfaces. *Nature Genetics*, *48*(1), 94-100.
752
753    Pironon, S., Papuga, G., Villellas, J., Angert, A. L., García, M. B., & Thompson, J. D. (2017). Geographic
754    variation in genetic and demographic performance: new insights from an old biogeographical
755    paradigm. *Biological Reviews*, *92*(4), 1877-1909.
756
757    Pringle, J. M., & Wares, J. P. (2007). Going against the flow: maintenance of alongshore variation in allele
758    frequency in a coastal ocean. *Marine Ecology Progress Series*, *335*, 69-84.
759
760    Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., et al. (2007). PLINK: a
761    tool set for whole-genome association and population-based linkage analyses. *The American Journal of*
762    *Human Genetics*, *81*(3), 559-575.
763
764    Raj, A., Stephens, M., & Pritchard, J. K. (2014). fastSTRUCTURE: variational inference of population
765    structure in large SNP data sets. *Genetics*, *197*(2), 573-589.
766
767    Ringbauer, H., Kolesnikov, A., Field, D. L., & Barton, N. H. (2018). Estimating barriers to gene flow from
768    distorted isolation-by-distance patterns. *Genetics*, *208*(3), 1231-1245.
769
770    Rivadeneira, M. M., Hernáez, P., Antonio Baeza, J., Boltana, S., Cifuentes, M., Correa, C., Cuevas, A.,
771    del Valle, E., Hinojosa, I., Ulrich, N., Valdivia, N., Vásquez, N., Zander, A., & Thiel, M. (2010). Testing the
772    abundant-centre hypothesis using intertidal porcelain crabs along the Chilean coast: linking abundance
773    and life-history variation. *Journal of Biogeography*, *37*(3), 486-498.
774
775    Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for paired-
776    end sequencing improve RADseq-based population genomics. *Molecular ecology*, *28*(21), 4737-4754.
777

778 Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under isolation by
779 distance. *Genetics*, *145*(4), 1219-1228.
780
781 Samis, K. E., & Eckert, C. G. (2007). Testing the abundant center model using range-wide demographic
782 surveys of two coastal dune plants. *Ecology*, *88*(7), 1747-1758.
783
784 Santini, L., Pironon, S., Maiorano, L., & Thuiller, W. (2019). Addressing common pitfalls does not provide
785 more support to geographical and ecological abundant-centre hypotheses. *Ecography*, *42*(4), 696-705.
786
787 Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and ecology of species range
788 limits. *Annual Review of Ecology, Evolution, & Systematics*, *40*, 415-436.
789
790 Siegel, D. A., Kinlan, B. P., Gaylord, B., & Gaines, S. D. (2003). Lagrangian descriptions of marine larval
791 dispersion. *Marine Ecology Progress Series*, *260*, 83-96.
792
793 Silva-Arias, G. A., Reck-Kortmann, M., Carstens, B. C., Hasenack, H., Bonatto, S. L., & Freitas, L. B.
794 (2017). From inland to the coast: spatial and environmental signatures on the genetic diversity in the
795 colonization of the South Atlantic Coastal Plain. *Perspectives in Plant Ecology, Evolution and*
796 *Systematics*, *28*, 47-57.
797
798 Soberón, J., Peterson, T.A., Osorio-Olvera, L. (2018). A comment on "Species are not most abundant in
799 the centre of their geographic range or climatic niche". *Rethinking Ecology*, 3: 13-18.
800
801 Wares, J. P., & Pringle, J. M. (2008). Drift by drift: effective population size is limited by advection. *BMC*
802 *Evolutionary Biology*, *8*(1), 1-7.
803
804 Wiens, B. J., & Colella, J. P. (2024a). triangulaR: an R package for identifying AIMs and building triangle
805 plots using SNP data from hybrid zones. *bioRxiv*, 2024-03.
806
807 Wiens, B. J., & Colella, J. P. (2024b). That's Not a Hybrid: How to Distinguish Patterns of Admixture and
808 Isolation By Distance. *Molecular Ecology Resources*, e14039.
809
810 Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population
811 structure. *Evolution*, 1358-1370.
812
813 Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*(2), 97.
814
815 Wright, S. (1943). Isolation by distance. *Genetics*, *28*(2), 114.
816
817 Zhan, S., Zhang, W., Niitepold, K., Hsu, J., Haeger, J. F., Zalucki, M. P., Altizer, S., de Roode, J.C.,
818 Reppert, S.M., & Kronforst, M. R. (2014). The genetics of monarch butterfly migration and warning
819 colouration. *Nature*, *514*(7522), 317-321.
820

 **FIGURE LEGENDS**

**Figure 1**. *Sample locations and characterization of population genetic structure*. A)

Points are sampled localities, colored by latitude. Northbound arrows are the Gulf

Stream, southbound are the Labrador Current, size differences represent different

current intensities. Inset is a photograph of a shovel bug collected from New Jersey. B)

PCA results; points are individuals, colored by the latitude of their sample site. C)

Isolation-by-distance; points are sample localities plotted by their pairwise linearized $F_{ST}$

against geographic distance in miles. Colors are comparisons within groups (north or

south of the Chesapeake) or between them.

**Figure 2**. *Abundant center results*. A) Individual genetic diversity (π) as a function of

latitude. B) Inbreeding coefficient as a function of the distance from the center of the

range, taken as 36° latitude, in miles. Points are colored by the latitude of their sample

location as in **Fig. 1A**.

**Figure 3**. *Directionality index (ψ) results*. A) Empirical estimates of ψ, points are the

summed values of ψ for all comparisons of population *i* to populations *j*. B) Simulation

results for each stepping-stone model: "center_asym" is abundant center with

asymmetric migration; "center_sym" is abundant center with symmetric migration;

"equal_asym" is equal deme size with asymmetric migration; and "equal_sym" is equal

deme size with symmetric migration. Each point per population is a different replicate.

Points are colored by estimated genetic diversity (π) of the sampled locality. Dashed

line is when ψ = 0, which is expected under equilibrium isolation-by-distance.

**FIGURES**

**Figure 1.** *Sample locations and characterization of population genetic structure.* A)

Points are sampled localities, colored by latitude. Northbound arrows are the Gulf

Stream, southbound are the Labrador Current, size differences represent different

current intensities. Inset is a photograph of a shovel bug collected from New Jersey. B)

PCA results; points are individuals, colored by the latitude of their sample site. C)

Isolation-by-distance; points are sample localities plotted by their pairwise linearized $F_{ST}$

against geographic distance in miles. Colors are comparisons within groups (north or

south of the Chesapeake) or between them.

**Figure 2**. *Abundant center results*. A) Individual genetic diversity (π) as a function of

latitude. B) Inbreeding coefficient as a function of the distance from the center of the

range, taken as 36° latitude, in miles. Points are colored by the latitude of their sample

location as in **Fig. 1A**.

**Figure 3**. *Directionality index (ψ) results*. A) Empirical estimates of $\psi$, points are the

summed values of $\psi$ for all comparisons of population *i* to populations *j*. B) Simulation

results for each stepping-stone model: "center_asym" is abundant center with

asymmetric migration; "center_sym" is abundant center with symmetric migration;

"equal_asym" is equal deme size with asymmetric migration; and "equal_sym" is equal

deme size with symmetric migration. Each point per population is a different replicate.

Points are colored by estimated genetic diversity (π) of the sampled locality. Dashed

line is when $\psi$ = 0, which is expected under equilibrium isolation-by-distance.