ASYMMETRIC GENE FLOW MAINTAINS RANGE EDGES IN A MARINE

INVERTEBRATE

Zachary B. Hancock[*,1], Nicole E. Adams[1], Perry L. Wood Jr.[1], Gideon S. Bradburd[1]

[1]Department of Ecology & Evolutionary Biology, University of Michigan

[*]Corresponding author:

Zachary B. Hancock

Department of Ecology & Evolutionary Biology

University of Michigan

Ann Arbor, MI 48103

Email: hancockz@umich.edu

Running title: Asymmetric gene flow maintains range edges in a marine invertebrate

ABSTRACT

Species are restricted to finite ranges, and when they are in demographic equilibrium

with the environment this implies a failure to adapt at the range periphery. There are

several competing evolutionary hypotheses for the maintenance and persistence of

range edges. One hypothesis is gene swamping of edge populations, in which migrants

from the center carrying alleles adapted to the range center but maladaptive along the

edge hamper local adaptation. A competing explanation is dispersal limitation, in which

reduced migration to the periphery keeps edge populations small, causing drift to

dominate over selection. It has proven difficult to assess these potential generative

scenarios due to the inherent complexity of species ranges. In this study, we leverage

range-wide genomic sequencing data to investigate the evolution of range edges in an

empirical system - *Haustorius canadensis*, the Canadian shovel bug. We find increased

genetic drift and inbreeding on the range edges and detect signatures of asymmetric

migration towards the range center. The inferred directionality of migration is consistent

with the direction of the dominant current regimes in the North Atlantic. We conclude

that anisotropic dispersal towards the range center leaves progressively fewer migrants

to replenish peripheral populations, consistent with the migration limitation hypothesis,

and generates the appearance of an abundant center.

KEYWORDS: gene swamping, range evolution, directionality index, migration, coastal,

shovel bug, isolation-by-distance

1 INTRODUCTION

All species are restricted to finite ranges. How range edges form and persist, why

populations fail to adapt and continue to expand their range, and what the long-term

stability of range size is, are all fundamental questions in the ecology and evolution of

ranges (Kirkpatrick & Barton 1997; Sexton et al. 2009; Gaston 2003; Louthan et al.

2015; Angert et al. 2020). Answering these questions has major implications for urgent

areas of research, such as predicting species' responses to climate change (Doak &

William 2010) or the propensity of invasive species to rapidly expand their range

(Alexander & Edwards 2010).

The simplest explanation for the persistence of a species' range limits is that they

reflect a species' niche limits (Holt 2003; Hargraves et al. 2014); i.e., the species'

distribution is determined by the environment (Angert et al. 2020). When range and

niche limits coincide, it suggests a failure of peripheral populations to adapt to local

conditions and expand both niche and range. There are several competing evolutionary

hypotheses that seek to explain the persistence of range edges due to a failure to

locally adapt. Haldane (1956) and Mayr (1963) were among the first to suggest that

asymmetric gene flow from central populations to the edge populations might limit the

latter's ability to adapt due to "gene swamping." Gene swamping is expected to occur if

migrants from central populations carry with them alleles adapted to "central" conditions

but, due to antagonistic pleiotropy, maladaptive at the range edge and conditions

beyond. If migration rates from the center populations are high enough, any adaptive

alleles segregating in edge populations may become "swamped" by the influx of

maladaptive alleles from migrants. Kirkpatrick & Barton (1997) formalized the gene

swamping hypothesis by developing a quantitative genetic model across environmental

clines of varying steepness, demonstrating that swamping could be a powerful

mechanism maintaining species range edges. Gene swamping is a special case of the

"migration load," the reduction in fitness of a population due to the influx of migrant

alleles maladapted to local conditions (Garcia-Ramos & Kirkpatrick 1997; Bolnick &

Nosil 2007).

However, empirical evidence for the prevalence of gene swamping has been

mixed. In a large review of the literature, Kottler et al. (2021) find little empirical

evidence for gene swamping. Indeed, multiple studies indicate that increased gene flow

to edge populations might have a positive effect (e.g., Sexton et al. 2011; Bontrager &

Angert 2019). Range edges often have lower genetic diversity (Eckert et al. 2008;

Pironon et al. 2016) and increased inbreeding depression (Arnaud-Haond et al. 2006;

De Ryck et al. 2016) compared to core populations, both conditions that could be

alleviated by increased migration. Therefore, Kottler et al. (2021) suggest that empirical

evidence supports an alternative mechanism of range edge evolution: dispersal

limitation (Sexton et al. 2009; Hargreaves et al. 2014). Under a dispersal limitation

scenario, the lack of migrants to edge populations may be depriving them of the needed

genetic variation to respond to strong selective pressures at the niche limit. As opposed

to gene swamping, which predicts that increased migration from the center will hinder

adaptation, the dispersal limitation hypothesis posits increased gene flow has the

potential to rescue edge populations (Halbritter et al. 2015; Evans et al. 2014; Silva-

Arias et al. 2017). However, due to the inherent complexity of species ranges,

identifying range limits (e.g., Dallas et al. 2017; Pironon et al. 2017; Soberón et al.

2018) and the underlying processes generating those limits remains challenging (Kottler

et al. 2021; Fristoe et al. 2023), especially in organisms not amenable to transplant

experiments. Thus, an ideal system for investigating questions of range evolution would

have a relatively simple range with well-defined edges.

In the ocean, particularly for pelagic organisms, an important cause of dispersal

limitation may be the direction and strength of the dominant current regimes (Siegal et

al. 2003). The average dispersal distance of a marine pelagic larva, for example, is a

function of its settling time and current velocity (Pringle & Wares 2007; Wares & Pringle

2008). White et al. (2010) found that incorporating larval duration and directional

oceanic currents dramatically improved model predictions of population differentiation in

the Kellet's whelk (*Kelletia kelletii*) over models relying on geographic distance alone.

Similarly, Dambach et al. (2016), investigating a deep-sea shrimp (*Nematocarcinus*

lanceopes), identified patterns of relatedness between populations consistent with

current connectivity in the Southern Ocean. Thus, marine ranges can be determined by

complex abiotic and biotic factors that can cause mass movement of individuals in

asymmetric ways. For example, in the North Atlantic, a northward current, the Gulf

Stream, emerges from the Loop Current in the Gulf of Mexico, bringing warm water from

the Caribbean, and travels up the coastline to Chesapeake Bay before diverting to the

mid-Atlantic (**Fig. 1A**). A second regime, the Labrador Current, travels southward along

Nova Scotia towards New England, bringing with it cold Arctic water, before also

diverting eastward. Hence, along the North Atlantic coastline, two currents converge

along the central latitude from opposite directions. For a marine organism distributed

along the entire coastline and whose dispersal is largely dictated by the currents, this

could generate a migration bias towards the center of the range. If the asymmetry in

dispersal is strong enough, it might even suggest that range limits are not reflective of

niche limits, but instead the lack of available migrants for successive colonization and

further range expansion into otherwise habitable patches (Hargraves et al. 2014;

Oldfather et al. 2019).

Small amphipod crustaceans in the family Haustoriidae, which inhabit the

exposed sandy beaches of the western North Atlantic, are an excellent system in which

to test questions about range evolution. Due to their life history (see *Methods*),

haustoriid amphipods are expected to be relatively poor dispersers, and this has been

largely confirmed in previous studies (Jones 2005; Hancock et al. 2019). In the western

North Atlantic, *Haustorius canadensis* (Bousfield 1962) can be found on beaches from

central Florida to southern Maine with some relict, isolated populations reported in the

Gulf of Saint Lawrence (Bousfield 1962; 1965). Haustoriid amphipods have no prior

common name; here we introduce the term "shovel bugs" (with reference to their

modified shovel limbs) to describe haustoriids, and "Canadian shovel bug" to refer to *H.*

canadensis. A curious feature of their southernmost distribution is the abundance of

suitable sandy beaches farther south than they are found, with no Canadian shovel bug

recorded south of Melbourne, Florida. The northern range edge, alternatively, might

represent a hard environmental transition, as most of coastal Maine is composed of

rocky outcrops with few fine sand beaches.

Given their relatively constrained niche (the intertidal of wave-exposed sandy

beaches), the ease of identifying range boundaries, and the one-dimensional nature of

their range, the Canadian shovel bug is an ideal organism for evaluating hypotheses of

the evolutionary mechanisms determining range limits. If gene swamping is maintaining

range edges in this species, we should detect a signature of asymmetric gene flow from

the range center towards the edge. Furthermore, we expect edge populations to harbor

the genetic signature of population sinks – lower genetic diversity relative to more

central populations, and the genetic diversity they do possess should be a subset of that

harbored in the center. Alternatively, range edges might be maintained by a failure to

colonize habitable patches due to asymmetries in the dispersal medium. Because

shovel bugs passively disperse via the currents, successively more distant patches in

the opposite direction of the current flow may have fewer available migrants to colonize

them, ultimately determining range limits. If dispersal limitation due to asymmetries in

current regimes are driving the persistence of range edges, we expect to find patterns of

asymmetric migration towards the range center. In addition, we would expect to identify

population-clustering associated with the two major currents, and a breakdown of this

clustering pattern at central latitudes where the currents converge.

In this study, we first characterize the population genetic structure of the

Canadian shovel bug across its entire range in the North Atlantic. Next, we estimate

genetic diversity and inbreeding across the range to test for patterns of reduced

abundance along the range edges. Finally, we determine whether patterns of migration

and colonization are consistent with either gene swamping or dispersal limitation as

processes maintaining range edges. We find evidence for directional colonization from

the periphery of the range toward the center, which we hypothesize is due to current

regimes in the North Atlantic and consistent with the migration limitation hypothesis.

2 METHODS

2.1 Natural history of shovel bugs

Deemed "the most interesting group of amphipods," shovel bugs are haustoriid

amphipods that have highly derived morphological adaptations, including expanded

articles on the 6th and 7th pereopods, extensive setal elaboration, and spiny, cup-like

structures on their 3rd and 4th pereopods that they utilize as shovels (**Fig. 1A**) (Barnard

1969; Hancock & Wicksten 2018).

Shovel bugs are blind and entirely fossorial, living in the first 1–10 cm of sand

and almost exclusively occupying the narrow swash zone (Croker & Hatfield 1980).

They lack a pelagic larva; instead, females brood juveniles through the first few molts in

their marsupium. Adult shovel bugs range in body size from 2–18 mm, sometimes within

the same species, and past work has found strong correlations between body size and

latitude (Hancock et al. 2021), though there may also be an effect of sand grain size

(LeCroy et al. 2002; Hancock & Wicksten 2018). Several species may occupy the same

beach, but there is evidence that they partition the beach by sex and by species to limit

competition for space (Croker & Hatfield 1980).

The relationship of shovel bugs to the broader Amphipoda has been shrouded in

mystery, with some hypothesizing a close kinship with other fossorial amphipods (Lowry

& Myers 2017). However, recent molecular phylogenies have cast doubt on the

taxonomic affinities between the fossorial amphipods, with multiple studies indicating

shovel bugs are gammarids and that modifications to the fossorial lifestyle found across

several families arose convergently (Verheye et al. 2016; Copilas-Ciocianu et al. 2020;

Hancock et al. 2022).

2.2 Field collecting

Shovel bugs were collected from 18 sites from Cocoa Beach, Florida, USA to Crescent

Beach, Maine, USA in May of 2023 (**Fig. 1A**; *Table S1*). Collections began in West

Palm Beach, Florida, USA to identify the southernmost established population; traveling

north from West Palm Beach, no specimens were located until Cocoa Beach, Florida.

Similarly, sampling continued beyond Crescent Beach, Maine, but no samples were

located further north (last attempted location was Popham Beach, ~93 km north of

Crescent Beach). Specimens were sifted from sand using a 435 μm sieve plate, plucked

from the plate with forceps, and preserved in 95% ethanol. Individuals were identified to

species in the lab using a dissecting microscope and key from Hancock & Wicksten

(2018), and then stored at -20°C prior to DNA extraction.

2.3 Molecular techniques

Whole genomic DNA was extracted from either the whole individual or from pereopods

4–7 using a Qiagen DNeasy Blood & Tissue Kit following manufacturer's protocols.

Library prep was performed using Restriction site Associated DNA sequencing (2RAD)

following Bayona-Vásquez et al. (2019), with slight modifications. All 96 samples were

quantified using a Qubit™ 1X dsDNA High Sensitivity (HS) kit on a Qubit 2.0

fluorometer. Samples were then normalized to have 100ng per sample (5 μL) on a 96

well plate, and then digested in a 15 μL solution with 1.5 μL 10X CutSmart Buffer (New

England Biolabs, USA), with 10 units of EcoRI (0.5 μL), 10 units MseI (0.5 μL), 3.5 μL

Invitrogen™ UltraPure™ DNase/RNase-Free Distilled Water, 2.5 μL of EcoRI

compatible adapter (2 μL), and 0.25 μM MseI compatible adapter (μL) on a Eppendorf

Mastercycler PRO S 6325 Thermal Cycler thermocycler at 37°C for one hour with the

heated lid turned off. Immediately after digestion we added 5 μL of ligation mastermix

composed of 2.75 μL of nuclease free water, 0.5 μL 10X ligation buffer (NEB), 1.5 μL 10

mM rATP (Promega), and 0.25 μL T4 DNA Ligase 400,000 U/ml to each sample. The

ligation reaction was implemented on the same thermocycler as above for two cycles of

22°C for 20 min, 37°C for 10 min, followed by 80°C for 20 min to stop the enzyme

activity. We then pulled 5 μL of each sample and split these into two microcenterfuge

tubes and performed a 1.25X SpeedBead cleanup (Rohland and Reich, 2012).

To reduce PCR duplicates and add the iTru5–8N barcode (which has a segment

of eight random base pairs) we implemented a single PCR cycle with six tubes each

with a 50 μL reaction. Each reaction contained 10 μL nuclease free water, 25 μL KAPA

HiFi hot start ready mix (Roche, USA), 5 μL iTru5–8N barcode (5 μM), and 10 μL of

library and were added using a single cycle of 98˚C for 1 min, 60˚C for 30s, and 72˚C

for 6 min. All six PCR reactions were then pooled and a 1.5X SpeedBead cleanup was

performed and resuspended in 30 μL of nuclease free water. Next, we split the library

into three 55 μL reactions with three unique iTru7 adapters. Each reaction had 10 μL of

nuclease free water, 25 μL KAPA HiFi hot start ready mix (Roche, USA), 5 μL P5 primer

(5 μM) and 5 μL of each iTru7 (5 μM) added separately. The PCR was implemented on

the thermocycler mentioned above under the following conditions: 95˚C for 3 mins, 8

cycles of 98°C 20s, 60° 15s, 72°C 30s, and a final 72°C for 5 mins. All PCR reactions

were pooled and then cleaned using a 1.5X SpeedBead cleanup and resuspended in 30

μL of nuclease free water.

We performed size selection on the pooled library using a Pippin Prep (Sage

Science, USA) with internal standards on a 1.5% dye free cassette, target 350–480 bp

range (mean= 415 bp). After size selection to enrich the library, we performed a PCR

under the same conditions for the iTru7 reaction above, except with only four 50 μL

reactions and 6 cycles, using both the P5 and P7 primers (5 μM). Finally, we performed

a 1X SpeedBead clean up and quantified the library on the Qubit 2.0 to ensure that our

library had a high enough concentration for sequencing. The library was then

sequenced on 12.5% percent of a NovaSeq X lane at the University of Michigan

Advanced Genomics Core.

Raw sequence reads were assembled and filtered using the de novo pipeline in

STACKS v.2.67 (Catchen et al. 2013; Rochette et al. 2019). A custom *process_radtags*

function was provided to us by J. Catchen to handle the unique FASTQ files of 2RAD,

which include an additional adapter sequence. Reads were assembled using the de

novo pipeline (Catchen et al. 2011); briefly, stacks were built in *ustacks* with a maximum

nucleotide distance allowed of 2 and minimum number of reads needed to seed a stack

of 3. Stacks were merged allowing a maximum of 2 gaps and 6 mismatches. Next, we

used the *populations* function, filtering out alleles that did not occur in at least 10

populations and keeping those that were present in 75% of individuals per population.

Lastly, we randomly selected one SNP per RAD locus (--write-random-snp).

Sequences were output as two VCF files, one that included only variant sites and a

second that contained both variant and monomorphic sites. VCF files were additionally

filtered using *vcftools* v.1.16 (Danecek et al. 2011); first, individuals with >80% missing

data were removed. Next, we filtered out loci with more than two alleles (--max-alleles

2) and with a minor allele frequency of less than 0.05 (--maf 0.05). We then filtered out

loci with a minimum mean depth of less than 20 (--min-meanDP 20) and removed all

indels (--remove-indels). After filtering, we evaluated the relationship between mean

depth, estimated heterozygosity, and distance from the range center via linear

regression to determine if mean depth might be a confounding factor.

2.4 Population genetic analysis

To evaluate potential patterns in genetic diversity, we estimated individual-level

heterozygosity using *pixy* (Korunes & Samuk 2021) on the VCF that included invariant

sites. The inbreeding coefficient (F; Wright, 1931), estimated as the difference between

the expected and observed homozygosity per individual, was estimated using *vcftools* (-

-het).

To test for geographic structure, we first used *pixy* to estimate population-level

pairwise F_{ST} (Weir & Cockerham 1984), which we then linearized and regressed against

geographic distance to test for isolation-by-distance (Wright 1943; Rousset 1997). Next,

we performed a principal component analysis (PCA) to identify signs of genetic

clustering using *VCF2PCACluster* (He et al. 2024). We then performed two cluster

analyses. First, we used *fastSTRUCTURE* (Raj et al. 2014) with k (the number of

discrete, ancestral clusters) from 2–5. Next, to account for continuous spatial

differentiation, we ran *conStruct* (Bradburd et al. 2018), varying k from 1–4. For each

value of *k*, we performed 5 independent MCMC chains each with 2000 iterations. We

then reported results from the chain with the highest mean posterior probability. As an

additional test of whether identified clusters and admixed individuals were the result of

increasing gene flow towards the range center or continuous isolation-by-distance, we

also performed a triangle plot analysis using *triangulaR* (Wiens & Colella 2024a). In the

presence of directional gene flow, individuals with a hybrid index approaching 0.5

should have concomitantly high interclass heterozygosity. However, a continuous

population characterized solely by isolation-by-distance will display a high hybrid index

but no increase in interclass heterozygosity (Wiens & Colella 2024b).

We evaluated potential barriers to migration using the program *FEEMS* (Marcus

et al. 2021), which estimates an effective migration surface using edge-specific

parameter estimation in a pseudo-likelihood framework. Lastly, we tested for a signature

of range expansion along the periphery of the population using Peter & Slatkin's (2013)

directionality index, ψ, as implemented in *rangeExpansion*

(https://github.com/BenjaminPeter/rangeexpansion). The directionality index is

calculated by comparing the difference in allele frequencies between pairs of

populations, conditioned on the allele being present in both populations. Large source

populations typically contain many rare alleles. When a new population is founded from

this larger source population, these same alleles, if they have not been lost due to drift,

are likely to drift to higher frequencies in the new population. Thus, successively more

recently founded populations in an expanding range will have increasingly positive

values of ψ, permitting the identification of both the origin and the direction of the range

expansion (Peter & Slatkin 2013; 2015). Following Peter & Slatkin (2015), the

directionality of range expansion is inferred by summing across the columns of the

matrix of pairwise estimates of ψ for each population (by row), where the value of ψ is

the strength of the signal and the sign is the direction. In populations at migration-drift

equilibrium in continuous space, simulations show that ψ should be zero (Peter &

Slatkin 2013). Therefore, if ψ is zero, there is no marked departure from migration-drift

equilibrium and hence no signature of reduced fitness or recurrent colonization in edge

populations. However, if gene swamping is driving edge dynamics, we expect ψ should

be positive in the direction of the edge. Alternatively, if dispersal limitation, driven by

directional currents, is the primary cause of range limits, then ψ should be negative at

the edges and increasingly positive toward the range center.

2.5 Simulations

Because the expansion statistic ψ is influenced by relative rates of drift, the inferred

direction of population expansion, as given by ψ, may be influenced by spatial clines in

population density even under scenarios with isotropic dispersal. To determine whether

the results of our ψ analyses are robust to spatial variation in population density (see

Results/Discussion), we conducted forward-time simulations using SLiM v.4.0.1 (Haller

& Messer 2023). We established a 10-deme stepping-stone model with four different

range scenarios: 1) equal population size per deme with symmetrical migration; 2) equal

population size per deme with asymmetrical migration from the edge populations

towards the center; 3) an abundant center model in which the central demes had a

fivefold higher census population size than the edge, with symmetric migration; and 4)

abundant center with asymmetric migration from the edges inward. For the asymmetric

migration scenarios, migration rates from the edge towards the center were twice that of

center to edge, and for scenarios with an abundant population center, population size

decayed linearly towards the edge. In addition, we performed simulations with

increasing differences in population sizes between the central and edge populations,

with equal migration, to evaluate how these relative differences in genetic drift might

drive patterns in ψ. We simulated ratios of population size between the center and edge

of 5:1, 25:1, and 50:1 and performed 10 replicates for each scenario. For each ratio,

population size decreased linearly, but more precipitously at higher ratios to balance

maintaining constant deme number and computational efficiency. Individuals were

diploid and hermaphroditic, and we simulated a 10 Mb chromosome with a

recombination rate of 1×10^{-8} per base pair. Simulations were run for 100,000 ticks

($>10N$) to reach demographic equilibrium, and tree sequences were recorded for

subsequent processing using *pyslim* v.1.0.1 (Kelleher et al. 2018) and *tskit* v.0.5.3

(Kelleher et al. 2018). For tree sequences with more than one root, we used recapitation

to simulate a neutral coalescent history until only a single root remained; mutations

were added at a rate of 1×10^{-7} per base-pair to the trees using *msprime* v.1.2.0

(Kelleher et al. 2016). Next, we randomly sampled five individuals per deme and output

their sequences in VCF format. From here, we trimmed the VCFs to only include

biallelic SNPs, and converted them to BED files using *plink* v.1.9 (Purcell et al. 2007).

To use the *rangeExpansion* package, we converted the individual coordinates from

SLiM, which were in discrete demes, into Earth coordinates as required by

rangeExpansion; thus, we chose 10 coordinates 5° apart in latitude. Because the units

of distance are arbitrary in SLiM, we needed only to ensure that deme 1 is closest to

deme 2, etc., to adequately capture the SLiM environment in an Earth coordinate

system. We then calculated ψ for each simulation to see how the different range

scenarios might impact inferences made from ψ.

3 RESULTS

3.1 ddRAD assembly

A total of 2,605,149 loci passed filtering in STACKS; the mean effective per-sample

coverage was 23.0 with a standard deviation of 13.6, a minimum of 4.0, and a maximum

of 67.4. The mean number of sites (bp) per locus was 220.1. After filtering for biallelic

SNPs and missingness in *vcftools*, the final dataset included 7,405 SNPs. We found a

significant positive relationship between mean depth per individual and our estimates of

heterozygosity from *pixy* (P < 5.3e–7, R^2 = 0.27, β = 8.77e–6; *Table S2*); in addition, we

found a significant negative relationship between mean depth and the distance from the

center of the range (taken as 36° latitude; P < 0.002, R^2 = 0.10, β = –0.051) (see *Table*

S2). To account for the potential confounding between measures of heterozygosity,

mean depth, and their relationship with the distance from the range center, all

subsequent multivariate linear regressions were performed with mean depth as an

additional predictor. To determine if a few individuals with high mean read depth were

driving the observed patterns, we re-tested for a significant correlation between depth

and heterozygosity after dropping all individuals with a mean depth greater than 30.

While a significant correlation between heterozygosity and mean depth remained (P <

2.7e–5; R^2 = 0.38, β = 3.1e–5), it was no longer significantly correlated with distance

from the center (P = 0.57; R^2 = –0.02, β = –2.03) (*Fig. S1*).

3.2 Genetic structure

The PCA revealed relatively continuous structure, with a possible break between

populations north and south of the Chesapeake (**Fig. 1B**), but there were no obvious

discrete breaks identified in patterns of isolation-by-distance when regressing linearized

F_{ST} against geographic distance (**Fig. 1C**). Results of the *fastSTRUCTURE* analysis

supported three geographically restricted population clusters (k = 3) as the model that

maximized the marginal likelihood – one from Florida to South Carolina, a second from

North Carolina to Virginia, and a third that included the remaining sample sites from

northern Virginia to Maine (*Fig. S2*). However, *conStruct*, which allows for continuous

structure to be accommodated within discrete populations, supported k = 2. Identified

clusters corresponded to the dominant current regimes in the Atlantic with increasing

admixture towards the center of the range (**Fig. 2**). This result was echoed by the

triangle plot analysis of the *conStruct* results, which found increasingly elevated

interclass heterozygosity at central latitudes (*Fig. S3*).

3.3 Patterns of diversity

After incorporating sequencing depth as a possible confounding effect, measures of

individual heterozygosity were significantly predicted by the distance from the center of

the range (P < 3.79e–16, β = –1.5e–6; mean depth P = 5.1e–5, β = 5.38e–6). This

relationship remained significant after dropping individuals with mean depth greater than

30 ($P < 0.0009$; $R^2 = 0.25$, $\beta = -1.2e-6$) (*Fig. S4*). Similarly, measures of inbreeding

were the highest at the range edges, declining as a function of distance from the center

(**Fig. 3B**). These patterns are consistent with field observations: the ease of collecting

and observed highest abundance was in the central range locations and shovel bugs at

the edge sample locations were rarer and thus more difficult to collect. However, we

note that while distance is a much stronger predictor of patterns of genetic diversity and

inbreeding, mean sequencing depth was still a significant predictor in each model

(*Table S2*).

The effective migration surface estimated from *FEEMS* suggested elevated

levels of dispersal from northern Florida to Virginia along the path of the Gulf Stream

and reduced effective migration towards the edges (**Fig. 2B**). In accordance with results

from the PCA, the migration surface indicated a strong depression in effective migration

across the Chesapeake, which suggests it might act as a partial barrier to gene flow.

3.4 Directionality index & simulations

The inferred origin of the range expansion of the entire sample of individuals, as

indicated by the lowest value of ψ, was 29.9°N in northern Florida (**Fig. 4A**). When the

data were subset into two regions, South and North (based on the *conStruct* analysis),

the inferred origins were at 29.9° and 43.6° latitude, respectively. The sum of ψ for each

population resulted in a pattern of increasing ψ towards the range center, mirroring the

increase in heterozygosity (**Fig. 4A**). This pattern of ψ superficially suggests a range

expansion from the edges toward the center of the range that follows the dominant

current regimes of the Gulf Stream running south to north and the Labrador Current

running north to south, both converging in the center of the range (**Fig. 1A, 4A**). This

pattern of expansion is in opposition to expectations from the gene swamping

hypothesis and suggests that asymmetric migration due to the currents might be

maintaining range edges, consistent with the migration limitation hypothesis.

Simulation results from SLiM supported this interpretation. Across replicates, the

strongest divergences from $\psi = 0$, which is expected at migration-drift equilibrium,

occurred in simulations with asymmetric migration from the edges toward the center,

with the most striking inflation in the scenario in which there is both an abundant center

and asymmetric migration (**Fig. 4B**). While there was some deviation from zero along

the edges in scenarios with symmetric migration in both the equal and center-abundant

scenarios, they did not systematically produce a positive ψ in the center as in the case

of asymmetric migration (**Fig. 4B**). Furthermore, increasing the population size

discrepancy between the center and the edge did not consistently lead to an increase in

central estimates of ψ (*Fig. S5*), indicating that asymmetric migration, not increased drift

along the edges, is the dominant force shaping the observed dynamics.

4 DISCUSSION

The persistence of range edges despite the potential for local adaptation to marginal

conditions and resulting continued expansion is an enduring evolutionary mystery.

Several hypotheses have been proposed to explain the finite nature of species ranges,

principle among these gene swamping (Mayr 1963; Kirkpatrick & Barton 1996) and

dispersal limitation (Sexton et al. 2009; Hargraves et al. 2014). In this study, we

investigated the suitability of these hypotheses to explain range evolution of the

Canadian shovel bug (*Haustorius canadensis*; Bousfield 1962), which is distributed

along the Eastern Seaboard of North America.

Past work suggests that shovel bugs have the potential for large amounts of

cryptic diversity and are sensitive to environmental barriers (Jones 2005; Hancock et al.

2019). Thus, our first aim was to characterize the population genetic structure of the

Canadian shovel bug. While we found that patterns of linearized F_{ST} were consistent

with a single continuous population, the PCA and *conStruct* results suggested the

presence of two populations associated with each current regime. In addition, we found

evidence for resistance to migration across the Chesapeake Bay. Shovel bugs of the

genus *Haustorius* specialize on open, wave-exposed beaches in which the sand is

highly oxygenated (Bousfield 1970), and rarely occur in low salinity, brackish bays.

While the Chesapeake Bay was formed by an impact crater during the Eocene (Ivany et

al. 2000), and this event has been postulated to have contributed to haustoriid

diversification in North America (Hancock et al. 2022), its recent form as a large bay is

more recent (around the end of the Last Glacial Maximum ~10,000 years ago (Colman

& Mixon 1988)). Thus, the bay has the potential to be acting as a partial barrier to

dispersal and gene flow. However, because the isolation-by-distance regression did not

support a discrete break (**Fig. 1C**; *Fig. S3*), the barrier may be weak or has occurred

recently enough that it has not yet influenced genome-wide patterns of allele frequency

divergence, which can lag behind other population summary statistics (Barton et al.

2013; Ringbauer et al. 2018).

Once the genetic structure of shovel bugs was characterized, our next aim was

to determine if range edges have reduced genetic diversity and increased inbreeding

relative to central populations. Past theoretical (Polechová & Barton 2015) and empirical

(Eckert et al. 2008; Pironon et al. 2016) work suggest that genetic diversity generally

decreases as a function of distance from the range center when the environment

changes gradually across space. Consistent with this past work, we found genetic

diversity and inbreeding strongly correlated with distance from the range center (**Fig. 3**);

edge populations harbor the lowest genetic diversity and highest rates of inbreeding.

This pattern is broadly consistent with the "abundant center hypothesis" (Grinnell 1922;

Hengeveld & Haeck 1982; Brown 1984), which predicts that population abundance is

highest in the core of the range and decreases monotonically towards the edge.

However, the generality of the abundant center hypothesis remains hotly contested, as

there is little empirical evidence that population abundance consistently declines from

the core to the edge (Dallas et al. 2020; Santini et al. 2019; Sagarin et al. 2006). While

we observed higher densities of shovel bugs in the central sample sites, anecdotally

consistent with the abundant center hypothesis, future work would benefit from

quantitative estimates of census abundance across the Canadian shovel bug's range.

Reduced genetic diversity and elevated rates of inbreeding in peripheral

populations might indicate a reduced adaptive capacity (Angert et al. 2020; Kottler et al.

2023). During a range expansion, this can be driven by allele "surfing," which describes

the process by which genetic drift is increased along an expanding front due to reduced

population densities, allowing alleles to reach high frequencies (Excoffier & Ray 2008).

When a range size is stable, reduced adaptive capacity might result from gene

swamping or dispersal limitation (Angert et al. 2020); both are expected to reduce

population density on the range periphery. If range limits are being determined by gene

swamping, we expected to infer asymmetric migration from the center toward the edge,

with edge populations harboring a subset of the genetic diversity of central populations.

Furthermore, we predicted that the shared alleles with central populations would be at

higher frequency on the edge, consistent with repeated founder events at increasingly

peripheral populations. Conversely, if range limits are being determined by dispersal

limitation, we expected to infer a net direction of movement towards the range center

from the edges, which might be reducing dispersal to peripheral populations. This is due

to the dominant current regimes in the North Atlantic flowing in opposite directions,

converging at a central latitude in the Chesapeake Bay region (**Fig. 1A**). In addition, if

dispersal limitation is driving the persistence of range edges, we expected increasing

admixture proportions towards the center of the range where the Gulf Stream and

Labrador Current converge.

Our results strongly support the finding that dispersal limitation is the dominant

mechanism shaping the persistence of range edges in shovel bugs. Our spatial

clustering analysis indicated there are two populations of the Canadian shovel bug

associated with each of the current regimes in the North Atlantic, with admixture

proportions increasing toward the range center (**Fig. 2**). We also identified reduced

effective migration on the range periphery (**Fig. 2B**). Interestingly, we observed that

effective migration is generally higher south of the Chesapeake than north, which is

consistent with the greater velocity of the Gulf Stream relative to the much weaker

Labrador Current (**Fig. 1**). Furthermore, we detected a linear increase in ψ towards the

range center from each edge. The standard interpretation of this index would imply a

range expansion from the periphery towards the center. However, this is contradictory

to patterns of inbreeding and genetic diversity, which clearly indicated that drift is likely

more dominant on the range edge (**Fig. 3**). A possible explanation for these conflicting

patterns is that the ψ index is influenced by asymmetric dispersal, and, if so, the

observed directionality would support the dispersal limitation hypothesis.

To evaluate whether inferences made on the basis of the ψ index could be

affected by heterogeneity in density as well as migration, we conducted forward-time,

individual-based simulations intended to tease apart the impacts of differences in

population size and migration (**Fig. 3B**; *Fig. 5*). Consistent with our explanation above,

we found that ψ was highly sensitive to asymmetric migration and would infer a

population expansion in the direction of the range center even in equilibrium (i.e., non-

expansion) scenarios. In addition, this bias was strongest when under the combined

influence of asymmetric migration and abundant center dynamics (**Fig. 3B**), but,

importantly, did not occur in the presence of differences in population size alone, at

least not at the ratios we tested (*Fig. S5*). This suggests that asymmetric migration may

be playing the dominant role in shaping patterns in the SFS that are being identified by

the ψ index.

The ψ index is a popular statistic for testing for range expansions and has been

applied to a diverse array of different organisms (Manuel et al. 2016; Zhan et al. 2014;

Fifer et al. 2022). In this study, we uncovered a potential confounding influence on ψ

that is likely to impact any organism with high asymmetric migration. While asymmetric

gene flow has been identified in terrestrial organisms (Fedorka et al. 2012; Oswald et al.

2017), the taxa most likely to display directional dispersal are aquatic organisms with

passive dispersal via river flow or oceanic currents (Siegal et al. 2003; Pringle & Wares

2007; Morrissey & de Kerckhove 2009). Past work has shown that directional dispersal

in the ocean can mislead many population genetic analyses, including the inference of

random dispersal via patterns of isolation-by-distance and the probability of

coalescence (Nagylaki 1977; Lenormand 2002; Paz-Vinas et al. 2013). Using a

simulation approach, we uncovered that asymmetric migration can also bias the

detection of a range expansion, potentially generating a false positive signature when,

in fact, the population is in demographic equilibrium.

In addition to the impact asymmetric migration has on the inference of a range

expansion, certain biased dispersal patterns may themselves be capable of generating

an abundant center pattern. Specifically, asymmetric migration from the edges towards

the range center can maintain high abundance in the center, causing it to act like a

population sink, while colonization towards increasingly distant patches becomes less

likely as the number of available migrants dispersing away from the center becomes

increasingly small. Our findings in this system suggest that this may be an

underappreciated form of migration limitation (Sexton et al. 2009; Hargraves et al.

2013), in which the rate of extinction in peripheral patches exceeds that of colonization;

the cause of this decreased migration is asymmetries in the probability of migration due

to the dispersal medium itself.

The evolutionary processes that maintain range edges, especially in systems in

which the environmental change is clinal, are still subjects of active debate (Kottler et al.

2023). We uncovered strong support for the hypothesis that range edges in this system

are maintained by dispersal limitation, itself driven by a combination of passive dispersal

and directional currents reducing the number of available migrants to colonize

peripheral patches. We suggest this mechanism for generating an apparent abundant

center might be widespread in coastal marine taxa, especially those with passive

dispersal, as oceanic currents regularly converge within species ranges. Furthermore,

focusing on asymmetric migration via directional water or wind currents may help clarify

what maintains range edges in cases where the fundamental niche appears to exceed

the range limit.

DATA AVAILABILITY

Bioinformatic and analysis pipelines, including filtered VCF data objects, are available at

https://github.com/zachbhancock/asym_gene_flow_proj. Raw sequences were

deposited to GENBANK (ascension PRJNA1183394).

AUTHOR CONTRIBUTIONS

ZBH and GSB conceptualized and designed the study. ZBH carried out the field

collections. ZBH and PLW performed the laboratory work. ZBH and NEA carried out the

bioinformatics and analyzed the data. All authors wrote and edited the manuscript.

ACKNOWLEDGEMENTS

Thanks to members of the Bradburd lab who provided helpful comments on this

manuscript. Thanks also to Kristen Wacker, Teresa Pegan, and Benjamin Peter for

thoughtful discussion and feedback. We are indebted to Leonard Jones who coined the

common name "shovel bug" for haustoriid amphipods, and proudly use it in print for the

first time here. May it catch on. Lastly, special thanks to Daniel Stern Cardinale and

Robert Joel Duff for their hospitality and conversations while ZBH was collecting.

Research reported was supported by the National Institute of General Medical Sciences

of the National Institutes of Health (NIH) under award R35GM137919 (awarded to

GSB). The content is solely the responsibility of the authors and does not necessarily

represent the official views of the NIH.

REFERENCES

Alexander, J. M., & Edwards, P. J. (2010). Limits to the niche and range margins of alien
species. *Oikos*, *119*(9), 1377-1386.
580
581   Angert, A. L., Bontrager, M. G., & Ågren, J. (2020). What do we really know about adaptation at range
582   edges?. *Annual Review of Ecology, Evolution, and Systematics*, *51*(1), 341-361.
583
584   Arnaud-Haond, S., Teixeira, S., Massa, S. I., Billot, C., Saenger, P., Coupland, G., Duarte, C.M., &
585   Serrao, E. A. (2006). Genetic structure at range edge: low diversity and high inbreeding in Southeast
586   Asian mangrove (Avicennia marina) populations. *Molecular Ecology*, *15*(12), 3515-3525.
587
588   Barnard, J.L. (1969) *The Families and Genera of Marine Gammaridean Amphipoda*. Smithsonian
589   Institution Press, Washington, D.C., 535 pp.
590
591   Barton, N. H., Etheridge, A. M., Kelleher, J., & Véber, A. (2013). Genetic hitchhiking in spatially extended
592   populations. *Theoretical population biology*, *87*, 75-89.
593
594   Bayona-Vásquez, N.J., Glenn, T.C., Kieran, T.J., Pierson, T.W., Hoffberg, S.L., Scott, P.A., Bentley, K.E.,
595   Finger, J.W., Louha, S., Troendle, N., Diaz-Jaimes, P., Mauricio, R., Faircloth, B.C. (2019). Adapterama
596   III: Quadruple-indexed, double/triple-enzyme RADseq libraries (2RAD/3RAD). *PeerJ*, 7:e7724
597   http://doi.org/10.7717/peerj.7724.
598
599   Beerli, P., Mashayekhi, S., Sadeghi, M., Khodaei, M., & Shaw, K. (2019). Population genetic inference
600   with MIGRATE. *Current protocols in bioinformatics*, *68*(1), e87.
601
602   Bolnick, D. I., & Nosil, P. (2007). Natural selection in populations subject to a migration
603   load. *Evolution*, *61*(9), 2229-2243.
604
605   Bousfield, E. L. (1962). *New Haustoriid Amphipods from the Canadian Atlantic Region*. Canada
606   Department of Northern Affairs and National Resources.
607
608   Bousfield, E. L. (1965). Haustoriidae of New England (Crustacea: Amphipoda). *Proceedings of the United*
609   *States National Museum*.
610

Bousfield, E. L. (1970). Adaptive radiation in sand-burrowing amphipod crustaceans. *Chesapeake*
Science, *11*(3), 143-154.

Bradburd, G. S., Coop, G. M., & Ralph, P. L. (2018). Inferring continuous and discrete population genetic
structure across space. *Genetics*, *210*(1), 33-52.

Brown, J. H. (1984). On the relationship between abundance and distribution of species. *The American*
Naturalist, *124*(2), 255-279.

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks: building and
genotyping loci de novo from short-read sequences. *G3: Genes, Genomes, Genetics*, *1*(3), 171-182.

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool
set for population genomics. *Molecular Ecology*, *22*(11), 3124-3140.

Colman, S. M., & Mixon, R. B. (1988). The record of major Quaternary sea-level changes in a large
coastal plain estuary, Chesapeake Bay, eastern United States. *Palaeogeography, Palaeoclimatology,*
Palaeoecology, *68*(2-4), 99-116.

Copilaş-Ciocianu, D., Borko, Š., & Fišer, C. (2020). The late blooming amphipods: Global change
promoted post-Jurassic ecological radiation despite Palaeozoic origin. *Molecular Phylogenetics and*
Evolution, *143*, 106664.

Croker, R. A., & Hatfield, E. B. (1980). Space partitioning and interactions in an intertidal sand-burrowing
amphipod guild. *Marine Biology*, *61*, 79-88.

Dallas, T., Decker, R. R., & Hastings, A. (2017). Species are not most abundant in the centre of their
geographic range or climatic niche. *Ecology Letters*, *20*(12), 1526-1533.

Dallas, T. A., Santini, L., Decker, R., & Hastings, A. (2020). Weighing the evidence for the abundant-
centre hypothesis. *Biodiversity Informatics*, *15*(3), 81-91.

Dambach, J., Raupach, M. J., Leese, F., Schwarzer, J., & Engler, J. O. (2016). Ocean currents determine
functional connectivity in an Antarctic deep-sea shrimp. *Marine Ecology*, *37*(6), 1336-1344.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R.E., Lunter,
G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., & 1000 Genomes Project Analysis Group. (2011).
The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156-2158.

De Ryck, D. J., Koedam, N., Van der Stocken, T., van der Ven, R. M., Adams, J., & Triest, L. (2016).
Dispersal limitation of the mangrove Avicennia marina at its South African range limit in strong contrast to
connectivity in its core East African region. *Marine Ecology Progress Series*, *545*, 123-134.

Dixon, A. L., Herlihy, C. R., & Busch, J. W. (2013). Demographic and population-genetic tests provide
mixed support for the abundant centre hypothesis in the endemic plant Leavenworthia stylosa. *Molecular*
ecology, *22*(7), 1777-1791.

Doak, D. F., & Morris, W. F. (2010). Demographic compensation and tipping points in climate-induced
range shifts. *Nature*, *467*(7318), 959-962.

Eckert, C. G., Samis, K. E., & Lougheed, S. C. (2008). Genetic variation across species' geographical
ranges: the central–marginal hypothesis and beyond. *Molecular Ecology*, *17*(5), 1170-1188.

Evans, S. M., Sinclair, E. A., Poore, A. G., Steinberg, P. D., Kendrick, G. A., & Vergés, A. (2014). Genetic
diversity in threatened Posidonia australis seagrass meadows. *Conservation Genetics*, *15*, 717-728.

Excoffier, L., & Ray, N. (2008). Surfing during population expansions promotes genetic revolutions and
structuration. *Trends in Ecology & Evolution*, *23*(7), 347-351.

Fedorka, K. M., Winterhalter, W. E., Shaw, K. L., Brogan, W. R., & Mousseau, T. A. (2012). The role of
gene flow asymmetry along an environmental gradient in constraining local adaptation and range
expansion. *Journal of Evolutionary Biology*, *25*(8), 1676-1685.

Fifer, J. E., Yasuda, N., Yamakita, T., Bove, C. B., & Davies, S. W. (2022). Genetic divergence and range
expansion in a western North Pacific coral. *Science of the Total Environment*, *813*, 152423.

Fristoe, T. S., Vilela, B., Brown, J. H., & Botero, C. A. (2023). Abundant-core thinking clarifies exceptions
to the abundant-center distribution pattern. *Ecography*, *2023*(2), e06365.

García-Ramos, G., & Kirkpatrick, M. (1997). Genetic models of adaptation and gene flow in peripheral
populations. *Evolution*, *51*(1), 21-28.

Gaston, K. J. (2009). Geographic range limits of species. *Proceedings of the Royal Society B: Biological
Sciences*, *276*(1661), 1391-1393.

Grinnell, J. (1922). The role of the" accidental". *The Auk*, *39*(3), 373-380.

Halbritter, A. H., Billeter, R., Edwards, P. J., & Alexander, J. M. (2015). Local adaptation at range edges:
comparing elevation and latitudinal gradients. *Journal of Evolutionary Biology*, *28*(10), 1849-1860.

Haldane, J. B. S. (1956). The relation between density regulation and natural selection. *Proceedings of
the Royal Society of London. Series B-Biological Sciences*, *145*(920), 306-308.

Haller, B. C., & Messer, P. W. (2023). SLiM 4: multispecies eco-evolutionary modeling. *The American
Naturalist*, *201*(5), E127-E139.

Hancock, Z. B., & Wicksten, M. K. (2018). Two new species of sand-burrowing amphipods of the genus
Haustorius Müller, 1775 (Amphipoda: Haustoriidae) from the northwestern Gulf of
Mexico. *Zootaxa*, *4459*(1), 101-127.

Hancock, Z. B., Hardin, F. O., & Light, J. E. (2019). Phylogeography of sand-burrowing amphipods
(Haustoriidae) supports an ancient suture zone in the Gulf of Mexico. *Journal of Biogeography*, *46*(11),
2532-2547.

Hancock, Z. B., Hardin, F. O., Murthy, A., Hillhouse, A., & Johnston, J. S. (2021). Rapid genomic
expansion and purging associated with habitat transitions in a clade of beach crustaceans (Amphipoda:
Haustoriidae). *Journal of Crustacean Biology*, *41*(3), ruab042.

Hancock, Z. B., Ogawa, H., Light, J. E., & Wicksten, M. K. (2022). Origin and evolution of the
Haustoriidae (Amphipoda): a eulogy for the Haustoriidira. *Zoological Journal of the Linnean*
Society, *194*(4), 1252-1267.
712
713　He, W., Xu, L., Wang, J., Yue, Z., Jing, Y., Tai, S., Yang,J., & Fang, X. (2024). VCF2PCACluster: a
714　simple, fast and memory-efficient tool for principal component analysis of tens of millions of SNPs. *BMC*
715　*Bioinformatics*, *25*(1), 173.
716
717　Hengeveld, R., & Haeck, J. (1982). The distribution of abundance. I. Measurements. *Journal of*
718　*Biogeography*, 303-316.
719
720　Hutchinson, G.E. (1957). Concluding remarks. Cold Spring Harbor Symposia on Quantitative Biology. 22
721　(2): 415–427.
722
723　Hutchinson, G.E. (1965). *The Ecological Theater and Evolutionary Play*. New Haven: Yale University
724　Press.
725
726　Ivany, L. C., Patterson, W. P., & Lohmann, K. C. (2000). Cooler winters as a possible cause of mass
727　extinctions at the Eocene/Oligocene boundary. *Nature*, *407*(6806), 887-890.
728
729　Jin, P. Y., Sun, J. T., Chen, L., Xue, X. F., & Hong, X. Y. (2020). Geography alone cannot explain
730　*Tetranychus truncatus* (Acari: Tetranychidae) population abundance and genetic diversity in the context
731　of the center–periphery hypothesis. *Heredity*, *124*(2), 383-396.
732
733　Jones, M.E. (2005) *Comparative Phylogeography, Population Structure, and Extended Maternal Care in*
734　*Three Intertidal Haustoriid Amphipods from the Western North Atlantic*. Unpublished Master's thesis,
735　College of Charleston.
736
737　Kelleher, J., Etheridge, A. M., & McVean, G. (2016). Efficient coalescent simulation and genealogical
738　analysis for large sample sizes. *PLoS Computational Biology*, *12*(5), e1004842.
739
740　Kelleher, J., Thornton, K. R., Ashander, J., & Ralph, P. L. (2018). Efficient pedigree recording for fast
741　population genetics simulation. *PLoS Computational Biology*, *14*(11), e1006581.
742
743　Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a species' range. *The American Naturalist*, *150*(1), 1-
744　23.
745
746　Kottler, E. J., Dickman, E. E., Sexton, J. P., Emery, N. C., & Franks, S. J. (2021). Draining the swamping
747　hypothesis: little evidence that gene flow reduces fitness at range edges. *Trends in Ecology &*
748　*Evolution*, *36*(6), 533-544.
749
750　Korunes, K. L., & Samuk, K. (2021). pixy: Unbiased estimation of nucleotide diversity and divergence in
751　the presence of missing data. *Molecular Ecology Resources*, *21*(4), 1359-1368.
752
753　LeCroy, S. E., Richardson, J. S., & Cobb, D. (2000). *An illustrated identification guide to the nearshore*
754　*marine and estuarine gammaridean Amphipoda of Florida* (Vol. 1, p. 2000). Florida Department of
755　Environmental Protection, Division of Resource Assessment and Management, Bureau of Laboratories.
756

757 Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, *17*(4),
758 183-189.
759
760 Louthan, A. M., Doak, D. F., & Angert, A. L. (2015). Where and when do species interactions set range
761 limits?. *Trends in Ecology & Evolution*, *30*(12), 780-792.
762
763 Lowry, J. K., & Myers, A. A. (2017). A phylogeny and classification of the Amphipoda with the
764 establishment of the new order Ingolfiellida (Crustacea: Peracarida). *Zootaxa*, *4265*(1), 1-89.
765
766 De Manuel, M., Kuhlwilm, M., Frandsen, P., Sousa, V. C., Desai, T., Prado-Martinez, J., et al. (2016).
767 Chimpanzee genomic diversity reveals ancient admixture with bonobos. *Science*, *354*(6311), 477-481.
768
769 Marcus, J., Ha, W., Barber, R. F., & Novembre, J. (2021). Fast and flexible estimation of effective
770 migration surfaces. *Elife*, *10*, e61927.
771
772 Mayr, E. (1963). *Animal species and evolution*. Harvard University Press.
773
774 Morrissey, M. B., & de Kerckhove, D. T. (2009). The maintenance of genetic variation due to asymmetric
775 gene flow in dendritic metapopulations. *The American Naturalist*, *174*(6), 875-889.
776
777 Nagylaki, T. (1977). Decay of genetic variability in geographically structured populations. *Proceedings of
778 the National Academy of Sciences*, *74*(6), 2523-2525.
779
780 Ntuli, N. N., Nicastro, K. R., Zardi, G. I., Assis, J., McQuaid, C. D., & Teske, P. R. (2020). Rejection of the
781 genetic implications of the "Abundant Centre Hypothesis" in marine mussels. *Scientific Reports*, *10*(1),
782 604.
783
784 Oldfather, M. F., Kling, M. M., Sheth, S. N., Emery, N. C., & Ackerly, D. D. (2020). Range edges in
785 heterogeneous landscapes: Integrating geographic scale and climate complexity into range
786 dynamics. *Global Change Biology*, *26*(3), 1055-1067.
787
788 Oswald, J. A., Overcast, I., Mauck III, W. M., Andersen, M. J., & Smith, B. T. (2017). Isolation with
789 asymmetric gene flow during the nonsynchronous divergence of dry forest birds. *Molecular
790 Ecology*, *26*(5), 1386-1400.
791
792 Paz-Vinas, I., Quéméré, E., Chikhi, L., Loot, G., & Blanchet, S. (2013). The demographic history of
793 populations experiencing asymmetric gene flow: combining simulated and empirical data. *Molecular
794 Ecology*, *22*(12), 3279-3291.
795
796 Peter, B. M., & Slatkin, M. (2013). Detecting range expansions from genetic data. *Evolution*, *67*(11),
797 3274-3289.
798
799 Peter, B. M., & Slatkin, M. (2015). The effective founder effect in a spatially expanding
800 population. *Evolution*, *69*(3), 721-734.
801
802 Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population structure with estimated
803 effective migration surfaces. *Nature Genetics*, *48*(1), 94-100.
804

Pironon, S., Papuga, G., Villellas, J., Angert, A. L., García, M. B., & Thompson, J. D. (2017). Geographic
variation in genetic and demographic performance: new insights from an old biogeographical
paradigm. *Biological Reviews*, *92*(4), 1877-1909.

Polechová, J., & Barton, N. H. (2015). Limits to adaptation along environmental gradients. *Proceedings of*
the National Academy of Sciences, *112*(20), 6401-6406.

Pringle, J. M., & Wares, J. P. (2007). Going against the flow: maintenance of alongshore variation in allele
frequency in a coastal ocean. *Marine Ecology Progress Series*, *335*, 69-84.

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., et al. (2007). PLINK: a
tool set for whole-genome association and population-based linkage analyses. *The American Journal of*
Human Genetics, *81*(3), 559-575.

Raj, A., Stephens, M., & Pritchard, J. K. (2014). fastSTRUCTURE: variational inference of population
structure in large SNP data sets. *Genetics*, *197*(2), 573-589.

Ringbauer, H., Kolesnikov, A., Field, D. L., & Barton, N. H. (2018). Estimating barriers to gene flow from
distorted isolation-by-distance patterns. *Genetics*, *208*(3), 1231-1245.

Rivadeneira, M. M., Hernáez, P., Antonio Baeza, J., Boltana, S., Cifuentes, M., Correa, C., Cuevas, A.,
del Valle, E., Hinojosa, I., Ulrich, N., Valdivia, N., Vásquez, N., Zander, A., & Thiel, M. (2010). Testing the
abundant-centre hypothesis using intertidal porcelain crabs along the Chilean coast: linking abundance
and life-history variation. *Journal of Biogeography*, *37*(3), 486-498.

Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for paired-
end sequencing improve RADseq-based population genomics. *Molecular ecology*, *28*(21), 4737-4754.

Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under isolation by
distance. *Genetics*, *145*(4), 1219-1228.

Samis, K. E., & Eckert, C. G. (2007). Testing the abundant center model using range-wide demographic
surveys of two coastal dune plants. *Ecology*, *88*(7), 1747-1758.

Santini, L., Pironon, S., Maiorano, L., & Thuiller, W. (2019). Addressing common pitfalls does not provide
more support to geographical and ecological abundant-centre hypotheses. *Ecography*, *42*(4), 696-705.

Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and ecology of species range
limits. *Annual Review of Ecology, Evolution, & Systematics*, *40*, 415-436.

Siegel, D. A., Kinlan, B. P., Gaylord, B., & Gaines, S. D. (2003). Lagrangian descriptions of marine larval
dispersion. *Marine Ecology Progress Series*, *260*, 83-96.

Silva-Arias, G. A., Reck-Kortmann, M., Carstens, B. C., Hasenack, H., Bonatto, S. L., & Freitas, L. B.
(2017). From inland to the coast: spatial and environmental signatures on the genetic diversity in the
colonization of the South Atlantic Coastal Plain. *Perspectives in Plant Ecology, Evolution and*
Systematics, *28*, 47-57.
852

853 Soberón, J., Peterson, T.A., Osorio-Olvera, L. (2018). A comment on "Species are not most abundant in
854 the centre of their geographic range or climatic niche". *Rethinking Ecology*, 3: 13-18.
855
856 Verheye, M. L., Backeljau, T., & d'Udekem d'Acoz, C. (2016). Looking beneath the tip of the iceberg:
857 diversification of the genus Epimeria on the Antarctic shelf (Crustacea, Amphipoda). *Polar Biology*, *39*,
858 925-945.
859
860 Wares, J. P., & Pringle, J. M. (2008). Drift by drift: effective population size is limited by advection. *BMC*
861 *Evolutionary Biology*, *8*(1), 1-7.
862
863 White, C., Selkoe, K. A., Watson, J., Siegel, D. A., Zacherl, D. C., & Toonen, R. J. (2010). Ocean currents
864 help explain population genetic structure. *Proceedings of the Royal Society B: Biological*
865 *Sciences*, *277*(1688), 1685-1694.
866
867 Wiens, B. J., & Colella, J. P. (2024a). triangulaR: an R package for identifying AIMs and building triangle
868 plots using SNP data from hybrid zones. *bioRxiv*, 2024-03.
869
870 Wiens, B. J., & Colella, J. P. (2024b). That's Not a Hybrid: How to Distinguish Patterns of Admixture and
871 Isolation By Distance. *Molecular Ecology Resources*, e14039.
872
873 Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population
874 structure. *Evolution*, 1358-1370.
875
876 Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*(2), 97.
877
878 Wright, S. (1943). Isolation by distance. *Genetics*, *28*(2), 114.
879
880 Zhan, S., Zhang, W., Niitepold, K., Hsu, J., Haeger, J. F., Zalucki, M. P., Altizer, S., de Roode, J.C.,
881 Reppert, S.M., & Kronforst, M. R. (2014). The genetics of monarch butterfly migration and warning
882 colouration. *Nature*, *514*(7522), 317-321.
883

 FIGURE LEGENDS

Figure 1. *Sample locations and characterization of population genetic structure*. A)

Points are sampled localities, colored by latitude. Northbound arrows are the Gulf

Stream, southbound are the Labrador Current, size differences represent different

current intensities. Inset is a photograph of a shovel bug collected from New Jersey. B)

PCA results; points are individuals, colored by the latitude of their sample site. C)

Isolation-by-distance; points are sample localities plotted by their pairwise linearized F_{ST}

against geographic distance in miles. Colors are comparisons within groups (north or

south of the Chesapeake) or between them.

Figure 2. *Admixture and effective migration surface*. A) Clustering (*conStruct*) analysis

results using $k = 2$. B) Effective migration surface from *FEEMS*. Cooler colors represent

regions of higher migration, warmer indicate potential barriers. Pies represent the

admixture proportions inferred from *conStruct* averaged across all individuals sampled

at a locality. Stars represent inferred origins of the population expansion from the ψ

analysis. Population abbreviations are defined in *Table S1*.

Figure 3. *Patterns of genetic diversity*. A) Individual genetic diversity (π) as a function of

latitude. B) Inbreeding coefficient as a function of the distance from the center of the

range, taken as 36° latitude, in miles. Points are colored by the latitude of their sample

location as in **Fig. 1A**.

Figure 4. *Directionality index (ψ) results*. A) Empirical estimates of ψ, points are the

summed values of ψ for all comparisons of population *i* to populations *j*. B) Simulation

results for each stepping-stone model: "center_asym" is abundant center with

asymmetric migration; "center_sym" is abundant center with symmetric migration;

"equal_asym" is equal deme size with asymmetric migration; and "equal_sym" is equal

deme size with symmetric migration. Each point per population is a different replicate.

Points are colored by estimated genetic diversity (π) of the sampled locality. Dashed

line is when ψ = 0, which is expected under equilibrium isolation-by-distance.

FIGURES

Figure 1. *Sample locations and visualization of population genetic variation*. A) Points are sampled localities, colored by latitude. Northbound arrows are the Gulf Stream, southbound are the Labrador Current, size differences represent different current intensities. Inset is a photograph of a shovel bug collected from New Jersey. B) PCA results; points are individuals, colored by the latitude of their sample site. C) Isolation-by-distance; points are sample localities plotted by their pairwise linearized F_{ST} against geographic distance in miles. Colors are comparisons within groups (north or south of the Chesapeake) or between them.

Figure 2. *Admixture and effective migration surface*. A) Clustering (*conStruct*) analysis

results using *k* = 2. B) Effective migration surface from *FEEMS*. Cooler colors represent

regions of higher migration, warmer indicate potential barriers. Pies represent the

admixture proportions inferred from *conStruct* averaged across all individuals sampled

at a locality. Stars represent inferred origins of the population expansion from the ψ

analysis. Population abbreviations are defined in *Table S1*.

933
934

Figure 3. *Patterns of genetic diversity*. A) Individual genetic diversity (π) as a function of

latitude. B) Inbreeding coefficient as a function of the distance from the center of the

range, taken as 36° latitude, in miles. Points are colored by the latitude of their sample

location as in **Fig. 1A**.

Figure 4. *Directionality index (ψ) results*. A) Empirical estimates of ψ, points are the

summed values of ψ for all comparisons of population *i* to populations *j*. B) Simulation

results for each stepping-stone model: "center_asym" is abundant center with

asymmetric migration; "center_sym" is abundant center with symmetric migration;

"equal_asym" is equal deme size with asymmetric migration; and "equal_sym" is equal

deme size with symmetric migration. Each point per population is a different replicate.

Points are colored by estimated genetic diversity (π) of the sampled locality. Dashed

line is when $\psi = 0$, which is expected under equilibrium isolation-by-distance.