1 **A common garden of *Halichondria* sponges: taxonomic revision of Northeast**
2 **Pacific Halichondriidae reveals many cryptic introduced species**
3

4 Thomas L. Turner[1], Christine Morrow[2], Bernard Picton[3], Claire Goodwin[4,5,6], Robert W. Thacker[7]
5
6 [1]Ecology, Evolution, and Marine Biology Department, University of California, Santa Barbara,
7 California, USA, tlturner@ucsb.edu
8 https://orcid.org/0000-0002-1380-1099
9
10 [2]School of Biological Sciences, Queen's University, Belfast, Northern Ireland, UK,
11 c.morrow@qub.ac.uk
12 https://orcid.org/0000-0001-5686-8747
13
14 [3]National Museums Northern Ireland, Northern Ireland, UK, bernard.picton@nmni.com
15 https://orcid.org/0000-0002-1500-2215
16
17 [4]Huntsman Marine Science Center, St. Andrew, New Brunswick, Canada
18 [5]University of New Brunswick, 100 Tucker Park Road Saint John, NB E2L 4L5, Canada.
19 [6]New Brunswick Museum, 277 Douglas Ave, Saint John NB E2K 1E5, Canada,
20 claire.goodwin@huntsmanmarine.ca
21 https://orcid.org/0000-0002-2428-1340
22
23 [7]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, USA,
24 and Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Republic of
25 Panama, robert.thacker@stonybrook.edu
26 https://orcid.org/0000-0002-9654-0073
27
28 Correspondence: tlturner@ucsb.edu

29
30 *Note about species names: this pre-print is not intended to be a publication of the associated*
31 *species names for the purposes of zoological nomenclature*
32

Abstract

Sponges (phylum Porifera) possess biochemical, cellular, and physiological traits with valuable biotechnical applications. However, our ability to harness these natural innovations is limited by a classification system that does not fully reflect their evolutionary history. In this study, we uncover numerous cryptic species within the genus *Halichondria* that are morphologically indistinguishable from the well-known *H. panicea*. Many of these species have habitat preferences and geographic distributions that strongly suggest they have been dispersed by human activity. Most of these species are broadly sympatric with their closest relatives, and these overlapping distributions allow us to use patterns of DNA variation to infer reproductive isolation between clades in nature. With reproductively isolated species thus delineated, we can use DNA states as taxonomic characters to formally describe them. Though much remains to be learned about these newly discovered species, the natural "common gardens" of introduced sponges in California, New York, and other locations provide opportunities to test hypotheses about their diversification in future work.

Introduction

The common ancestor of extant sponges is estimated to have lived over 700 million years ago (Dohrmann and Wörheide, 2017; Pankey et al., 2022; Rossi et al., 2024). This ancient origin suggests that some extant sponge lineages have been independently evolving for longer than the most distantly related animals within Bilateria. Sponges have undoubtedly diversified in biochemical, cellular and physiological dimensions during this prolonged time, developing numerous biological innovations with potential applications (Dunn et al., 2015; Mehbub et al., 2024). These innovations remain largely undiscovered, and it seems likely that a big impediment to advancing sponge biology is the lack of a stable classification system that reflects the evolutionary history of the phylum. The difficulty in assigning sponge specimens to species and determining where those species fall in the tree of life creates uncertainty that deters investigation across multiple disciplines.

Creating an accurate taxonomy of sponges based on morphological characters has proven challenging due to a paucity of informative character states. Sponge taxonomy is based mainly on skeletal characters: spicule morphology and the arrangement of spicules and/or spongin fibers into a three-dimensional skeleton (Hooper and Van Soest, 2002). While these features have been useful for diagnosing many species and higher taxonomic groups (Hooper and Van Soest, 2002; Morrow and Cárdenas, 2015), molecular phylogenies have revealed that skeletal traits alone are inadequate due to both insufficient differences between closely related species and convergently evolving traits in distantly related species (de Paula et al., 2012; Erwin and Thacker, 2007; Gazave et al., 2010; Morrow et al., 2013; Turner and Pankey, 2023; Xavier et al., 2010). These limitations are frequently encountered in sponge families with simple spicules and a loosely organized skeleton, such as the Halichondriidae. This family is of significant interest due to its pharmaceutical potential (Hirata and Uemura, 1986; Rodriguez Jimenez et al., 2021; Zhong et al., 2024) and as a model system, as this family includes one of the most studied sponges in the world, *Halichondria panicea* (Pallas, 1766). This species is found near many marine labs, is amenable to laboratory culture, and as a result, has been used to ask questions about sponge physiology, symbiosis, ecology, immunology, and more (Barthel, 1986; Barthel and Detmer, 1990; Barthel and Wolfrath, 1989; Carrier et al., 2023; Forester, 1979; Knobloch et

al., 2019; Schmittmann et al., 2021; Thomassen and Riisgård, 1995; Witte et al., 1994).
However, this research is built atop a shaky foundation, because the taxonomy and systematics
of *H. panicea* and its relatives is very poorly delineated. The World Register of Marine Species
lists over 70 previously named species that are now considered synonyms of *H. panicea* (de
Voogd, et al., 2024). Other species that remain putatively valid, like *H. bowerbanki* (Burton,
1930), are often difficult to differentiate from *H. panicea* and their validity remains in dispute.
Cryptic species within populations ascribed to *H. panicea* have remained a possibility, as this
species has supposedly been found in most of the world's oceans and in a wide variety of
climates. Under what can be called a "cryptic endemic species in different regions" hypothesis,
different regions of the world could contain morphologically indistinguishable species that are
endemic to each region (Erpenbeck, 2004). An alternative hypothesis, which can be called the
"single globally-distributed invasive species" hypothesis, is that *H. panicea* has been
unintentionally introduced by human activity to many regions of the world, as was recently
shown for the co-familial species *Hymeniacidon perlevis* (Samaai et al., 2022; Turner, 2020).
Below, we combine extensive field collections, genomic analysis, and traditional
morphological characters to reject both the "single globally-distributed invasive species" and
"cryptic endemic species in different regions" hypotheses. We reject the single-species
hypothesis by using multi-locus genomic data to infer reproductive isolation between sympatric
populations. When molecular phylogenies are built separately for mitochondrial genomes and
the nuclear-encoded ribosomal locus, we find numerous clades with reciprocal monophyly in
both compartments. Because the mitochondrial and nuclear genomes are unlinked, reciprocal
monophyly across these compartments is strong evidence of reproductive isolation when
species are sympatric (Jennings, 1917; Rannala and Yang, 2020). We are also able to reject the
"cryptic endemic species in different regions" hypothesis for many (but not all) of the delineated
species because we find a lack of correspondence between geographic region and species
range. Instead, we find numerous species that have likely been dispersed by human activity.
This newly characterized diversity presents an opportunity to use the sympatric, closely
related species of *Halichondria* as a natural "common garden" experiment to study sponge
ecology and evolution. This work will be greatly facilitated by formally describing the newly
discovered species because unique identifiers (such as Linnean names) are crucial for making
species visible to the research community and linking data across databases like GenBank, the
World Register of Marine Species, the Global Biodiversity Information Facility, and museum
inventories (Delić et al., 2017). Formally describing new species requires finding unique
characters that differentiate them from previously described species, which we failed to do
using traditional spicule and skeletal traits for most of the newly discovered species. We
therefore use DNA states as characters to define and diagnose species (Delić et al., 2017; Eitel et
al., 2018; Jörger and Schrödl, 2013; Lawley et al., 2021; Tessler et al., 2022). Using DNA
characters as taxonomic states has drawbacks, because identification of unknown samples will
be limited by preservation methods, sample age, and the expertise of investigators. However,
these problems would likely be even greater for alternative phenotypic traits such as
proteomics or histology, which would require less ubiquitous expertise and less common
preservation methods. We hope that this approach will lay the groundwork for future research
into the phenotypic differences among these newly described species.
120

Methods

Collections

Sponges were collected from the Atlantic Coast of Canada (Claire Goodwin), Northern Ireland (Christine Morrow and Bernard Picton), Sweden (Raquel Pereira), Portugal (Raquel Pereira), and three locations in the USA: New York (Robert Thacker), Washington (Thomas Turner, Robert Thacker, and Brooke Weigel), and California (Thomas Turner). Collections were most extensive in California, where 95 subtidal sites, 14 intertidal sites, and 23 marinas were surveyed. Subtidal sites were surveyed by divers, with no new collections from deep water (maximum depth 32 m). Previously vouchered material was acquired on loan from the Royal British Columbia Museum (voucher numbers with RBC), the Florida Museum (voucher numbers with UF or BULA), the Smithsonian National Museum (voucher numbers with USNM), the Naturalis Biodiversity Center (voucher numbers with ZMA), and the California Academy of Sciences (voucher numbers with CASIZ). Freshly collected material was deposited in these collections and also the Santa Barbara Museum of Natural history (voucher numbers with SBMNH), the Cheadle Center at the University of California, Santa Barbara (voucher numbers with UCSB), the Atlantic Reference Centre Museum (voucher numbers with ARC), the Ulster Museum (voucher numbers with BELEM), and the National History Museum of Los Angeles (voucher numbers with NHMLA). A table listing all examined samples with voucher numbers, GenBank accessions, collection dates and locations, and other metadata is available as supplementary data via Data Dryad.

Morphology

Spicules were examined after digesting sponge subsamples in bleach. For some samples, a single piece of sponge, including both the ectosome and choanosome, was used for digestion. In other cases, these domains were carefully separated with a razor and digested individually. Spicule measurements were taken from images using ImageJ (Schneider et al., 2012), with pixel-to-mm conversion determined using a calibration slide. The length of each spicule was measured as the longest straight line from tip to tip, even if the spicules were curved or bent. Width was measured at the widest point, excluding any adornments like swollen tyles. All spicule measurements are available as supplementary data via Data Dryad.

To analyze the choanosomal skeletons, sponge fragments were first frozen into ice cubes and then hand-cut into tissue sections. These sections were cleared using one of two methods. A few samples were transferred from a 95% ethanol bath to a 100% ethanol bath, and then cleared with histoclear (National Diagnostics). Most samples were digested in a mixture of 97% Nuclei Lysis Solution (Promega; from the Wizard Genomic DNA Purification Kit) and 3% 20 mg/ml Proteinase K (Promega). This method removed cellular material while preserving spicules and the spongin network (if present). For the ectosomal skeletons, the sponge surface was removed with a razor. In many cases, these skeletons could be imaged directly without further processing; if opaque, they were processed similarly to choanosomal sections.

To measure ectosomal tract thickness, we averaged the measurements of the three thickest visible tracts, excluding large matted areas. This process was difficult to standardize and the measurements are likely less reproducible than spicule size measurements. We also categorized ectosomal skeletons based on their morphology, noting that there were potentially distinct types. The five categories used were: 1. Multilayered and gothic-crypt-like; 2. Sieve-like

165 (a dense mat with approximately round open spaces); 3. Lace-like (a fairly isotropic mesh); 4.
166 Intermediate forms between categories 3 and 5; 5. A network of spicule tracts (resembling
167 either straight lines or meandering lines, similar to dry or wet spaghetti). In some cases,
168 ectosomal skeletons were found as dense mats of spicules without open spaces, but in these
169 cases, other areas of the same sponge exhibited one of the above types.
170
Illumina sequencing
DNA was extracted with several different kits including the Wizard Genomic DNA Purification kit
(Promega) and the Qiagen Blood & Tissue kit. Prior to Illumina sequencing, samples were
treated with RNase A and repurified with the DNA Clean and Concentrator (Zymo). Illumina
sequencing and library preparation were performed at the DNA Technologies Core Facility at the
University of California, Davis. Libraries were made with the Super-High-Throughput (SHT)
technique, which involved dual indexing each sample and pooling 96 samples before
sequencing. Paired-end 150 bp reads were generated using the Illumina NovaSeq instrument,
resulting in a median of 2.1 million reads per sample. The number of reads varied significantly
across samples, ranging from a maximum of 17 million reads to a minimum of 208 reads. The
reasons for this variability were not explored in detail, but DNA extractions appeared to be of
variable quality even before library construction and pooling, perhaps because of sample
preservation differences or the secondary chemistry of samples.
Reads were trimmed with fastP (Chen et al., 2018) and then de novo assembled into
contigs with Megahit (Li et al., 2015). The number of contigs per sample ranged from 0 to 1,080;
the median number of contigs for samples with at least one contig was 22. Blast searches
revealed that many contigs were microbial (or occasionally from other taxa), but the sponge's
nuclear ribosomal locus was usually the highest coverage contig. Indeed, in samples where only
one or two contigs assembled, the ribosomal locus was usually one of them. In samples where
the full ribosomal locus assembled de novo, this contig was then used as a reference sequence,
and reads from that sample were aligned to it with bwa-mem (Li and Durbin, 2009). SAMtools
(Danecek et al., 2021) was then used to convert SAM files from this alignment into a consensus
sequence, which served as the final sequence for that sample. For samples where no ribosomal
contig assembled at the megahit step, the most closely related sequence available (either from
another sample in our analysis or from GenBank) was used as reference sequence, and bwa-
mem was used to align that sample's reads to this reference. In some cases, ribosomal contigs
assembled with megahit were incomplete or fragmented, so these partial contigs were spliced
into the nearest available reference sequence to create a hybrid reference for bwa-mem
alignment. This approach successfully yielded nearly complete sequences of the 18S, 28S, and
intergenic transcribed spacer regions for most samples, though some had a significant
proportion of uncalled bases ("N"). Many of these samples had been previously sequenced at
the 28S locus with Sanger sequencing, and these sequences were combined with Illumina
sequencing when available. Combining Sanger and Illumina data often resulted in resolving
some uncalled bases but in no cases were Illumina consensus sequences found to have called
bases that conflicted with Sanger data, suggesting error rates of called bases are low. Ribosomal
sequences were assembled for 110 samples, with a median number of called bases (excluding
Ns) of 6,429 bp (ranging from 2,227 to 6,589 bp). The number of called bases for each sample is
listed in the supplementary table available from Data Dryad.

Mitochondrial contigs were present in some of the Megahit assemblies but were
generally less complete than ribosomal loci. To generate more complete mitochondrial
genomes, we aligned reads to existing reference sequences using bwa-mem. The references
used included *Halichondria okadai* (MG267395), *Halichondria galea* sp. nov. (KX244759),
Halichondria dokdoensis (MK105765), *Halichondria akesa* sp. nov. (MH756604), *Halichondria*
panicea (MH756603), and *Hymeniacidon perlevis* (KF192342).
For species lacking closely related reference sequences (e.g., the *Bowerbanki* group and
Hymeniacidon pierrei sp. nov.), we created hybrid reference sequences. This procedure involved
replacing parts of the closest available reference genome with de novo contigs from the
megahit assembly. Although this method allowed us to assemble substantial portions of the
mitochondrial genome for samples without good reference sequences, these assemblies were
less complete, often containing more uncalled bases ("Ns") due to regions being too divergent
for accurate short-read alignment. For instance, the average number of called bases in
Halichondria panicea mitogenomes was 19,228, while for *Halichondria bowerbanki*, it was
16,100. In total, we assembled mitochondrial genomes for 75 samples, with a median of 16,888
called bases. As seen with ribosomal loci, combining these sequences with Sanger data at cox1
did not find errors in called bases. The number of called bases for each sample is listed in the
supplementary table available from Data Dryad.
A sequence of the cox1 locus from the *H. panicea* neotype was generated using
sequence capture techniques (Agne et al., 2022) and was generously shared with us by Dirk
Erpenbeck.
230
Sanger sequencing
The cox1 region commonly sequenced in sponges is the Folmer region, a 709 bp segment
amplified using the primers LCO1490 and HCO2198 (Folmer et al., 1994). We used these
primers for a few samples when other primers failed to amplify, though they are less preferred
due to their shorter length and their tendency to co-amplify microbial sequences. We also
attempted to amplify DNA using the primers LCO1490 and cox1-R1, and were successful with
some samples (Rot et al., 2006). These primers target a larger 1264 bp region that includes both
the Folmer region and the "co1-ext" region, which has been useful for distinguishing sponge
species (Erpenbeck et al., 2006). However, we found that *Halichondria loma* and the
Bowerbanki group species had mismatches with the LCO1490 primer, and amplifications usually
failed with both primer sets. To improve amplification success, we designed new primers,
halcox1-L and halcox1-R, which were effective for all tested *Halichondria* species. These primers
amplify an 886 bp region that encompasses most of the Folmer region and part of the co1-ext
region. We used these new primers for all *Halichondria* samples not already processed with
Illumina or other primer sets. Primer sequences are provided in Table S2.
We sequenced two spans of the 28S nuclear locus. The D1-D2 region, which spans 940
bp and is highly variable, was amplified in all samples using primers 15F and 878R (Morrow et
al., 2012). We also amplified the adjacent D3-D5 region in some samples using primers 830F
and 1520R. This primer pair amplifies a 710 bp fragment that overlaps the D1-D2 fragment by
20 bp (Morrow et al., 2012).
The ND1 locus was amplified in selected taxa to verify that fixed differences detected in
Illumina sequencing were fixed in a larger sample. Primers were newly designed, but universal

primers for all target species were not found. For the *Panicea*, *Hygeia*, and *Dokdoensis* species
groups, except for *H. columbiana*, we used primers nd1-pan-R and nd1-pan-L. For *H.*
columbiana, nd1-pan-R was used with nd1-hygc-L. For the *Bowerbanki* species group, nd1-bow-
R and nd1-bow-L were used. For *H. loma*, nd1-loma-R was used with nd1-bow-L.
PCR was performed using a Biorad thermocycler (T100) using the following conditions:
95°C for 3 min, followed by 35 cycles of 94°C for 30 sec, [annealing temperature] for 30 sec,
72°C for [extension time], followed by 72°C for 5 minutes. The annealing temperatures and
extension times for each primer pair were as follows: halcox1 (54°C annealing, 60 second
extension), LCO & R1 primers (50°C annealing, 75 second extension), LCO & HCO primers (52°C
annealing, 60 second extension), 28S region (53°C annealing, 60 second extension for both
primer sets), ND1 locus (54°C annealing, 60 second extension, all primer sets). PCR was
performed in 50 µl reactions using the following recipe: 24 µl nuclease-free water, 10 µl 5x PCR
buffer (Gotaq flexi, Promega), 8 µl MgCl, 1 µl 10 mM dNTPs (Promega), 2.5 µl of each primer at
10 µM, 0.75 bovine serum albumin (10 mg/ml, final concentration 0.15 mg/ml), 0.25 µl Taq
(Gotaq flexi, Promega), 1 µl template. Post-PCR, samples were cleaned using ExoSAP-IT (Applied
Biosystems) and then sequenced by Functional Biosciences with Big Dye v3.1 on ABI 3730xl
instruments. We verified that the resulting sequences were of sponge origin using BLAST. All
sequences have been deposited in GenBank under the accession numbers listed in the
supplementary table available from Data Dryad..
272
Phylogenies
Sequences were aligned using MAFFT v.7 (Katoh et al., 2017), and the alignment files are
available as supplementary data. Phylogenies were estimated with maximum likelihood in IQ-
Tree (Nguyen et al., 2015; Trifinopoulos et al., 2016). The ultrafast bootstrap and Shimodaira-
Hasegawa approximate likelihood ratio test (Sh-aLrT) were used to measure node confidence
(Hoang et al., 2018). The optimal model for each tree was selected using ModelFinder
(Kalyaanamoorthy et al., 2017) based on the Bayesian Information Criterion. The GTR+F+I+G4
model was selected for all ribosomal trees/partitions, while mitochondrial models differed:
GTR+F+I+G4 for all concatenated mitochondrial genes, TN+F+I+G4 for cox1 only, and
TIM2+F+G4 for nd1 only. Figures were produced by exporting IQ-Tree files to the Interactive
Tree of Life webserver (Letunic and Bork, 2019).
Single-locus phylogenies of cox1, nd1, and 28S were made with a combination of Sanger
sequenced loci and Illumina sequences that were trimmed to a single locus. We used the NCBI
taxonomy browser and blast to assemble all available sequences related to the Halichondriidae,
along with appropriate outgroups. For cox1, sequences were excluded if they did not include
most of the Folmer barcoding region. For 28S, sequences in the D1 phylogeny were excluded if
they did not include the highly variable C2-D2 barcoding region.
Alignment files are available as supplementary files in Data Dryad.

Results

Molecular phylogenies

Our phylogeny for the 75 samples with the most comprehensive data includes both the nuclear ribosomal locus (averaging 6,445 bp) and the coding regions of the mitochondrial genome (averaging 17,153 bp; Figure 1). Previously, comparable data was available for only one species, *H. okadai*, which is included in the analysis. We successfully assembled nearly complete nuclear ribosomal loci for an additional 28 samples, including several additional species. However, we did not have sufficient reads to assemble the mitochondrial genome for these samples. Mitochondrial and ribosomal phylogenies are shown separately for the *Bowerbanki* group (Figure 2). For the remaining specimens, we constructed a ribosomal-only phylogeny (Figure 3).

Additionally, we constructed phylogenies using Sanger sequencing data from two mitochondrial loci (cox1 and nd1) and one nuclear locus (28S). These phylogenies were instrumental in validating genetic differences identified in the Illumina data and allowed us to include previously sequenced vouchers. In total, we sequenced at least one locus from 232 samples, and these phylogenies are included in the supplementary materials as (Figures S1-S3).

We consistently found highly differentiated clades across all phylogenies. Our first task was to determine which clade corresponded to the type species *H. panicea.* Due to the ambiguity of the original description, a neotype for the species was previously erected (Erpenbeck and van Soest, 2002). The neotype was collected from Burnham-on-Crouch, Essex, England, near the mouth of the Thames River, which is postulated to be near the original type location. Using sequence capture, Erpenbeck and colleagues generated 207 bp of sequence data from the hypervariable C2 region of the 28S barcoding locus from the neotype, and generously shared this previously unpublished sequence with us. Our 28S phylogeny confidently places this sequence within the clade designated *H. panicea* (Figure 1). It is an exact match to all three of the other samples from the United Kingdom, as well as samples from both the Atlantic and Pacific coasts of North America. These perfectly-matched samples were all previously identified as *H. panicea* based on their morphology, range, and habitat, and all fall into the *H. panicea* clade (Figure 1), supporting the genetic determination that this clade corresponds to the name *H. panicea*.

Similarly, we used a previously sequenced sample (Ulster Museum voucher BELUM:Mc4003; GenBank vouchers HQ379241, HQ379316, HQ379382, and KC902247) to determine which clade corresponds to the name *H. bowerbanki*. This name replaces *H. coalita* of Bowerbank, which was preoccupied by *H. coalita* (Müller, 1776). The type material consists only of microscope slides of isolated spicules. Like *H. panicea*, the type location is near the mouth of the Thames. However, two of the slides Bowerbank originally examined were from sponges collected at Strangford Lough, Northern Ireland. This location matches that of the Ulster Museum sample, which also aligns with the growth form described by Johnson (1842) and Bowerbank (1866). We are therefore confident that this clade corresponds to the name *H. bowerbanki*.

341
342 Figure 1. Maximum likelihood phylogeny for samples with assembled mitogenomes and nearly complete
343 ribosomal loci. Asterisks at nodes indicate a UFbootstrap ≥ 99%. Scale bar indicates substitutions per
344 site.
345
346
347
348

Inference of reproductive isolation: Bowerbanki group

Our next task was to determine which clades in our phylogeny correspond to species other than

H. panicea and *H. bowerbanki*. To infer reproductive isolation among species with overlapping

ranges, we compared the phylogenies of multiple loci that assort independently during meiosis.

For example, the mitochondrial and nuclear ribosomal phylogenies for the clade we named the

Bowerbanki group display reciprocal monophyly (Figure 2). Reciprocal monophyly in both

genomic compartments allows us to infer a minimum of four species in this group: *H.*

bowerbanki and three new species — *H. urca* sp. nov., *H. zabra* sp. nov., and *H. pinaza* sp. nov.

(new species descriptions are provided in the systematics section below). Among these species,

H. zabra and *H. pinaza* are the closest relatives. They are well-separated at the ribosomal locus,

but Sanger sequencing of the *cox1* mitochondrial locus did not reveal fixed genetic differences

between them (Figure S1). However, in the concatenated alignment of all mitochondrial coding

regions, we identified 24 fixed differences between these clades, including one fixed difference

in the cox1 gene outside the region bounded by our Sanger sequencing primers. With genetic

divergence (D_{xy}) of only 0.25% across the mitochondrial genes, the branch separating these

clades is short and difficult to visualize (Figure 2), but monophyly of each clade is well supported

(99% and 100% bootstrap support for *H. pinaza* and *H. zabra*, respectively).

368
Figure 2. Maximum likelihood phylogenies for the mitogenome (left) and ribosomal locus (right) for the

Bowerbanki group. Asterisks at nodes indicate a UFbootstrap ≥ 99%. Scale bar indicates substitutions per

site.

To confirm that these genetic differences are consistent in a larger sample, we

sequenced a region of the nd1 mitochondrial gene containing three apparent fixed differences.

In a larger sample of 16 *H. zabra* and 7 *H. pinaza* samples, these differences remained fixed.
Similarly, we sequenced the most variable region of the 28S nuclear ribosomal locus in 23 *H.*
zabra and 7 *H. pinaza* samples, confirming numerous fixed differences (divergence = 1.32%).
This co-assortment of genetic variation in the mitochondrial and nuclear genomes is strong
evidence of reproductive isolation due to the co-occurrence of these species in California. They
are sympatric even at a micro-geographic scale, with samples collected side-by-side on the
same floating dock in Tomales Bay in Northern California, and on floating docks in Santa Barbara
Harbor, over 500 km to the south. Likewise, *H. urca* and *H. bowerbanki* were sympatric at a
micro-geographic scale. All four species in this group were collected simultaneously from
floating docks in Santa Barbara Harbor, with varying degrees of sympatry across other harbors
and marinas throughout the state. There were numerous alleles fixed within the *H. urca* and *H.*
bowerbanki clades in both genetic compartments, with monophyly highly supported in both
cases (Figure 2).
Our inference of four species within the *Bowerbanki* group is likely conservative, as
phylogenetic analyses suggest at least one additional species. Within the clade we designate as
H. urca, one sample from Ireland is genetically distinct. Unlike the *H. pinaza* samples, where
Irish and California specimens are nearly identical at the 28S locus (divergence = 0.2%), the Irish
H. urca shows a 4.6% sequence divergence at this locus. A sample of *H. urca* from New York is
more similar to the California samples than to the Irish one, indicating that there might be a
single population spanning both ocean basins, with the Irish sample potentially representing a
different species or a highly divergent population. Additional *H. urca* data from the Atlantic
includes three sponges from Virginia (SERC/MarineGeo 2018 Lower Chesapeake BioBlitz,
available in GenBank) and one from New York (Pankey et al., 2022). These samples, previously
identified as *H. bowerbanki*, are recognized as *H. urca* in our cox1 phylogeny (Figure S1).
Combining these specimens with our samples results in a total of 5 Atlantic and 16 Pacific *H.*
urca at this locus, with a genetic divergence of 0.38% between them. Although this divergence
is slight, it is eight times greater than the divergence observed between *H. zabra* and *H. pinaza*,
which we can confidently say are reproductively isolated in sympatry. The differentiation seen in
cox1 for Atlantic versus Pacific *H. urca* is higher than for any other species found in both basins
(Fst = 0.64 for *H. urca*, compared to 0.00–0.58 for the six other species with distributions across
both oceans). Some or all Atlantic populations of *H. urca* may therefore be from an additional
species, but we conservatively assign all Atlantic samples to *H. urca* until additional data are
collected.
We also observed considerable variation in the mitochondrial genome within the clade
we assign to *H. bowerbanki* —far more variation than seen within any of the other species in
this study (Figure 2). In contrast to the clades we designate as species, however, the
mitochondrial clades within *H. bowerbanki* were not mirrored by divergence in the nuclear
locus. This lack of divergence could occur because there are multiple species within this clade,
but they are too recently diverged to have accumulated fixed differences in the ribosome.
Alternatively, this species may have a larger effective population size, higher mitochondrial
mutation rate, or balancing selection that has maintained mitochondrial variation within a
single species. We therefore assign all members of this clade to *H. bowerbanki*.
Finally, we note that there is disagreement between the mitochondrial and nuclear
compartments regarding the branching order of the species in this group. Mitochondrial genes

place *H. bowerbanki* as sister to the clade containing *H. zabra* and *H. pinaza*, while the nuclear
data position *H. bowerbanki* as sister to *H. urca*. This discrepancy could result from
mitochondrial introgression, incomplete lineage sorting, or homoplasy. Although this
discordance affects the inference of historical relationships, it does not impact conclusions
about current reproductive isolation and is the only instance of differing branching orders
between mitochondrial and nuclear genes in our Illumina-scale dataset.
425
Inference of reproductive isolation: Panicea group
Halichondria panicea was described from the North Atlantic, and there has been debate about
whether North Pacific populations belong to the same species (Erpenbeck, 2004; Erpenbeck and
van Soest, 2002). Our analysis reveals that Pacific and Atlantic populations of *H. panicea* are
sister clades distinguishable at the DNA level (Figures 1 & 3). Fixed genetic differences between
these ocean basins are evident in both nuclear and mitochondrial genomes, including two fixed
differences in each of the three Sanger-sequenced loci (cox1, nd1, and 28S). The genetic
divergence between Atlantic and Pacific populations of *H. panicea* at cox1 is similar to that
observed between Atlantic and Pacific populations of *H. urca* (0.40% divergence for both; Fst =
0.58 for *H. panicea*, Fst = 0.64 for *H. urca*). Unlike cases where populations are sympatric, this
divergence does not confirm that these populations are reproductively isolated. Indeed, *H.*
panicea have been reported from the Arctic. No genetic data are available to confirm the
identity of Arctic *Halichondria*, but if these populations are indeed *H. panicea*, it is possible that
populations from the Atlantic and Pacific meet and mix in these Northern waters. We therefore
leave the North Pacific population of *H. panicea* within this species, pending further data. We
note that a previous paper that attempted to compare *H. panicea* in the Atlantic and Pacific only
used two samples from the Atlantic, and alignment of those samples to our data reveal that the
Atlantic samples were, in fact, *H. bowerbanki* (Erpenbeck, 2004).
As noted in the morphological section, there are multiple cryptic species that are
morphologically indistinguishable from *H. panicea*. It is therefore surprising that, of the three
species most closely related to *H. panicea*, two are morphologically distinct. The most closely
related species is *H. sitiens* (Schmidt, 1870), which, like *H. panicea*, is found in both the North
Pacific and North Atlantic. *H. sitiens* is clearly differentiated from *H. panicea* by spicule size and
the presence of surface papillae. These papillae have historically been used to define subgenera
within *Halichondria*, with *H. sitiens* as the type species of subgenus *Eumastia* and *H. panicea* as
the type species of subgenus *Halichondria* (Erpenbeck and van Soest, 2002). The close
relationship between these two species clearly invalidates these subgenera. Although only one
H. sitiens was sequenced using Illumina technology, two Pacific and four Atlantic samples were
sequenced at each of the three Sanger loci. In contrast to *H. panicea*, all *H. sitiens* samples are
nearly identical, showing minimal genetic diversity both within and between populations
(divergence between ocean basins <0.1% at each locus, with no fixed differences).
Outside of the *H. panicea* + *H. sitiens* clade were two other species; short branches did
not allow us to resolve their branching order. One of these is represented by a single sample
that is morphologically indistinguishable from *H. panicea.* This sample (ESS07) was collected at
the same time and place as three other samples that were found to be *H. panicea* (Brig's Shoal,
in the Eastern Shore Islands of Nova Scotia). Though this clade is represented by only a single
sample in our collections, the mitochondrial genes are nearly identical (99.9%) to a previously

published mitochondrial genome from Iceland (GenBank accession MH756604). Previous work ascribed this genotype to *H. panicea* (Knobloch et al., 2019), but the closer relationship of *H. sitiens* to true *H. panicea* clearly indicates that a new name for this taxon is needed. We describe it here as *H. akesa* sp. nov.. This species differs from the *H. panicea* neotype DNA fragment at 3 sites (of 207); divergence vs. all *H. panicea samples* across the Sanger regions of cox1 and 28S is 0.50% and 0.80%, respectively.

The final species in the Panicea group serves to emphasize the inadequacy of spicule-based taxonomy in this family. *Topsentia fernaldi* (Sim and Bakus, 1986) has a very dense skeleton of long oxeas that make it quite distinct from *H. panicea*. The genus *Topsentia* is among the most polyphyletic of sponge genera, with previous molecular phylogenies placing some *Topsentia* species very far from other members of the Halichondriidae, in a sister clade to the order Axinellida (Pankey et al., 2022; Redmond et al., 2013).

Inference of reproductive isolation: Hygeia group
The most widely distributed species in the clade we dub the *Hygeia* group is the newly described *H. hygeia* sp. nov., which was found throughout the Northeast Pacific and in the Northwest Atlantic. In the Pacific, its range overlaps broadly with another new species, *H. californiana* sp. nov. These two species are distinctly separated at the ribosomal locus, showing 14 fixed differences (1.5% divergence) in the Sanger-sequenced region of 28S. Although they could not be differentiated in the Sanger-sequenced region of the cox1 mitochondrial locus, Illumina sequencing revealed numerous other fixed differences in the mitochondrial genome. Sanger sequencing of the nd1 mitochondrial gene confirmed 4 fixed differences in a larger sample (0.8% divergence, Fst = 0.73, N = 12 *H. californiana* vs. 17 *H. hygeia*). The fixed differences observed in both genomic partitions, coupled with their broad sympatry, strongly supports reproductive isolation in nature.

Two additional species are found in this group, both restricted to the Pacific Northwest of North America. Despite the limited number of samples (N=2 for each species), both are well differentiated from other species in the group based on mitochondrial and nuclear loci. Two *H. cascadia* sp. nov. samples were sequenced using Illumina technology, and these samples are strongly supported as a sister clade to all other species in the group (Figure 1). Given that *H. hygeia* and *H. californiana* are well-supported as isolated species, and *H. cascadia* is sympatric with *H. hygeia*, it is likely that *H. cascadia* is also an isolated lineage deserving of species status. We also have two samples of *H. columbiana* sp. nov., both from British Columbia, though only one was successfully sequenced with Illumina technology. This sample was quite unique, with many alleles differentiating it from the sympatric species *H. hygeia*. This sample formed a clade with the second sample across all three Sanger-sequenced loci, providing further evidence that this distinct clade is likely a separate species.

Figure 3 (below). Maximum likelihood phylogeny for samples with nearly complete ribosomal loci. Asterisks at nodes indicate a UFbootstrap ≥ 99%. Scale bar indicates substitutions per site.

Suberites lambei TLT296 California
Pseudosuberites latke TLT538 California
Pseudosuberites latke TLT1131 California
Halichondria sp. P19x63 Pacific Panama
Halichondria sp. P19x40 Pacific Panama
Halichondria galea sp. nov. TLT743 California
Halichondria galea sp. nov. TLT495 California
Halichondria okadai AB511881
Halichondria melanadocia P19x163 Caribbean Panama
Halichondria magniconulosa P19x171 Caribbean Panama

Okadai group

Hymeniacidon perlevis NIB0611 Ireland
Hymeniacidon perlevis NIB0616 Ireland
Hymeniacidon perlevis NIB0615 Ireland
Hymeniacidon perlevis TLT651 California
Hymeniacidon perlevis TLT1265 California
Hymeniacidon perlevis TLT736 California
Hymeniacidon perlevis TLT247 California
Hymeniacidon perlevis TLT658 California
Hymeniacidon perlevis TLT1074 California
Hymeniacidon pierrei sp. nov. TLT599 California
Hymeniacidon kitchingi TLT820 California
Semisuberites cribosa BHAK-09371 British Columbia
Hymeniacidon fusiformis TLT1194 California
Hymeniacidon sp. Quoddy115 Nova Scotia
Halichondria dokdoensis TLT386 California
Hymeniacidon actites TLT847 California
Hymeniacidon actites TLT227 California
Hymeniacidon actites TLT856 California
Hymeniacidon actites TLT454 California
Hymeniacidon actites TLT1158 California

Halichondria akesa sp. nov. ESS07 Nova Scotia
Topsentia fernaldi TLT1134 California
Halichondria sitiens ESS44 Nova Scotia
Halichondria sitiens ESS130 Nova Scotia
Halichondria panicea NIB0608 Ireland
Halichondria panicea 190424 Sweeden
Halichondria panicea 180923 Sweeden
Halichondria panicea 160504 Sweeden
Halichondria panicea Quoddy124 New Brunswick
Halichondria panicea 191026 Sweeden
Halichondria panicea Quoddy113 New Brunswick
Halichondria panicea ESS11 Nova Scotia
Halichondria panicea FH12x81 Washington
Halichondria panicea BHAK-00536 British Columbia
Halichondria panicea BHAK-03453 British Columbia
Halichondria panicea TLT858 California
Halichondria panicea TLT619 California
Halichondria panicea FH12x84 Washington
Halichondria panicea FH12x38 Washington
Halichondria panicea FH12x39 Washington

Panicea group

Halichondria cascadia sp. nov. FH12x93 Washington
Halichondria cascadia sp. nov. FH12x92 Washington
Halichondria californiana sp. nov. TLT417 California
Halichondria californiana sp. nov. TLT1093 California
Halichondria californiana sp. nov. TLT937 California
Halichondria californiana sp. nov. TLT860 California
Halichondria californiana sp. nov. TLT160 California
Halichondria columbiana sp. nov. BHAK-09466 British Columbia
Halichondria hygeia sp. nov. TLT742 California
Halichondria hygeia sp. nov. TLT645 California
Halichondria hygeia sp. nov. TLT1076 California
Halichondria hygeia sp. nov. TLT940 California
Halichondria hygeia sp. nov. TLT504 California
Halichondria hygeia sp. nov. F12x108 Washington
Halichondria hygeia sp. nov. BHAK-03597 British Columbia
Halichondria hygeia sp. nov. TLT724 California
Halichondria hygeia sp. nov. WMS21x10 New York
Halichondria hygeia sp. nov. WMS21x9 New York
Halichondria hygeia sp. nov. WMS19x5 New York

Hygeia group

Bowerbanki species group

Tree Scale: 0.05

Taxa are not diagnosable with traditional morphological characters
Our multi-locus molecular data, combined with broad sympatry, allows us to form strong
inferences regarding which clades are species. We can use these well-delineated species to
determine if known morphological characters are sufficient for their diagnosis. We first
considered only the previously named species *H. panicea* and *H. bowerbanki*. Previous attempts
to separate these species cite several differences including 1) longer spicules in *H. bowerbanki*,
2) a difference in spicule length in the choanosome vs. the ectosome in *H. panicea* but not *H.*
bowerbanki, and 3) differences in the arrangement of spicules in the ectosome between species
(Ackers et al., 2007; Hartman, 1958). These previous works did not use DNA to diagnose
samples, so it is likely that samples from the previously unknown species discovered and
described here were included in past work. Nonetheless, as detailed further below, we find that
these traits do differ, on average, between these species. However, we also find variability
within species that prevents them from being used to reliably diagnose all samples.
Spicules in *H. bowerbanki* were, on average, nearly 40% longer than those in *H. panicea*
(Wilcoxon p<0.0001, Figure 4). However, substantial overlap of these distributions prevents this
character from being used as a species diagnostic (Figure 4). Mean spicule length ranged from
202 to 357 μm for *H. panicea* and from 307 to 423 μm for *H. bowerbanki*. Both species had high
variability in mean spicule length within the Atlantic and Pacific basins. We also find that
spicules are, on average, 15% shorter in the ectosome than the choanosome for *H. panicea* but
not for *H. bowerbanki* (Figure S4). Again, this trait proved insufficient for diagnosing all samples
due to intraspecific variability. Some *H. panicea* specimens had spicules 30% longer in the
choanosome, and two samples (one from Nova Scotia and one from Sweden) had spicules of
equal length in both compartments. To verify that this variability within species was not entirely
due to technical variability and sampling error, we reisolated spicules from resampled tissues.
One *H. panicea* sample with no initial difference in spicule lengths by compartment and another
with a 32% difference were re-examined. Replicate data confirmed these results: the sponge
with no initial difference had only a 1% difference in the replicate (non-significant), while the
sponge with a 32% difference initially showed a 26% difference in the replicate (p<0.001). In
contrast, *H. bowerbanki* samples exhibited less variability, with ectosomal spicules varying from
about 10% longer to about 10% shorter than choanosomal spicules. Attempts to replicate
samples of *H. bowerbanki* with the greatest observed differences (11% longer choanosomal
spicules vs. 9% longer ectosomal spicules) showed both samples regressing towards a mean
difference of zero, consistent with the possibility that variation in *H. bowerbanki* is largely due
to tissue sampling variation and statistical noise.
When average spicule length and length differential between compartments are plotted
together, the average differences between *H. panicea* and *H. bowerbanki* are clear, and it
appears likely that a combination of these two traits would be sufficient to differentiate most,
but probably not all, samples (Figure 5). However, when we added the other species described
here to this plot, we observed substantial overlap among them.
The most comprehensive publication on the morphological distinction between *H.*
panicea and *H. bowerbanki* states that the arrangement of spicules in the ectosome is the most
reliable diagnostic character (Hartman, 1958). Typically, *H. bowerbanki* exhibits a network of
long, thin spicule tracts that delineate larger open spaces, while *H. panicea* has thicker spicule
tracts forming a dense mesh surrounding smaller openings. *H. panicea* sometimes also features

"gothic crypt-like supports" connecting the ectosomal and choanosomal skeletons (Hartman,
1958). Although these differences were observed in our samples, they proved insufficient for
diagnosing all samples reliably. This trait is highly variable within a single sponge and among
samples within species. While mean tract thickness differed significantly between species (69
μm for *H. panicea* vs. 38 μm for *H. bowerbanki*, Wilcoxon p<0.0001; Figure S5), there is
considerable overlap in the distributions (e.g., *H. panicea* 39–97 μm, *H. bowerbanki* 29–56 μm;
figure S5). We also attempted to categorize ectosomal skeletons into one of five structural types
(Figure S6); this categorization again highlighted average differences but also considerable
overlap. Principal components analysis combining ectosomal tract thickness, average spicule
length, and length differential between compartments did not enhance the ability of these traits
to diagnose species (Figure S7). Even when considering a small number of samples of only *H.*
panicea and *H. bowerbanki*, separation was incomplete.
561

562
563
564   Figure 4. Spicule length distributions for samples of *Halichondia panicea* (Hpan) and *H. bowerbanki*
565   (Hbow). Sample are sorted by mean within species, and names indicate if the sample was from the
566   Atlantic (Atl) or Pacific (Pac). Hpan samples generally have shorter average lengths and bimodal
567   distributions, due to length differences between the ectosome and choanosome, but some Hpan
568   samples are longer than the shortest Hbow samples, and bimodality is not always present.
569

Other morphological differences between (apparent) *H. panicea* and *H. bowerbanki* have
been proposed, such as smell (subjective and only available in fresh material), rigidity of
branches (branches often not present), color, and general form (Ackers et al., 2007; Hartman,
1958). It is likely that these traits and others will be partially informative, but our efforts to
differentiate samples based on these traits has proved inconsistent and frustrating. Moreover,
many morphological traits likely respond plastically to environmental conditions such as light
intensity, wave energy, and current strength (Palumbi, 1986). Some species are found primarily
in sheltered harbors and marinas, while others are found in wave-swept intertidal habitats, so
average morphological differences between them may be due more to habitat than genotype.
For instance, a sample of *H. urca* from an intertidal zone exhibited an ectosomal skeleton typical
of *H. panicea*, which is common in high-energy environments, whereas most *H. urca* specimens
from sheltered marinas displayed a sparsely spiculated ectosome (see *H. urca* description
below). Therefore, we conclude that the previously named species *H. panicea* and *H.*
bowerbanki cannot be reliably distinguished based on known morphological traits alone, and
the newly described *Halichondria* species should be considered cryptic species without clear
morphological differences.
586

587
588 Figure 5. Spicule measurements across *Halichondria* species. Only samples where spicules were
589 measured separately for the ectosome and the choanosome are shown. Mean spicule length is the mean
590 for both domains combined, and proportional length difference is the choanosomal mean minus the
591 ectosomal mean over the sample mean.
592
593

Conclusions

Here, we have documented a remarkable diversity of sympatric sponges in the family Halichondriidae. The discovery of cryptic species in this group isn't entirely surprising, given the few known morphological traits that distinguish them. It's also not surprising that some species have been spread through accidental human introductions, particularly since they are common on floating docks and other areas where introduced species have been observed, and other species in the family are known to be introduced (Carlton and Eldredge, 2009; Turner, 2020). However, we did not except to find so many cryptic introduced species, sympatricaly distributed at even small scales. *Halichondria bowerbanki*, *H. zabra*, *H. urca*, *H. pinaza*, and *H. hygeia* were all found growing on the same floating docks in a small harbor in Santa Barbara, California, and we document at least 4 of these species in Eastern North America and/or Europe. We hope that these unintentional "common gardens" of related sponges will facilitate future studies of their evolution and ecology.

We were unable to find morphological traits to diagnose many of these species. This presents challenges for ecological monitoring projects that lack genotyping capabilities. However, accurately delineating reproductively isolated groups is essential for monitoring both native diversity and potentially invasive species. Additionally, biologists studying physiology, biochemistry, or development need to know which species they are examining so that others can build on their findings. Despite their morphological similarities, we suspect that many of these species will prove be biologically distinct. For example, *Halichondria panicea* and *H. bowerbanki* are estimated to share a common ancestor in the mid-Cretaceous period, nearly 100 million years ago (Pankey et al., 2022). Even more dramatically, *Halichondria panicea* and *H. galea* have been independently evolving since the Jurassic (Pankey et al., 2022), splitting around the same time as the common ancestor of all Eutherian mammals (Archibald, 2003; Luo et al., 2011). While the Eutherian clade was diversifying into bats, whales, and primates, what were these sponges up to? We hope that our results will motivate future work on this question.

Systematics

The aims of this section are two-fold. First, we formally describe the *Halichondria* species that were found to be morphologically indistinguishable from *H. panicea*. Second, we aim to comprehensively revise the taxonomy of the family Halichondriidae for the Northeast Pacific, from British Columbia to California. This revision encompasses all newly discovered and previously described species within this region. The only species included here that has not been found in this area is *H. akesa* sp. nov., which is known solely from the North Atlantic.

Diagnoses include DNA character states with the format "number:letter", where the number indicates a position in one of the DNA alignments provided (where the first base is number 1), and the letter denotes the base state at that position, which is unique compared to all other species in the genus for which sequence data are available. Although many species of *Halichondria* lack DNA data, these previously described species were all morphologically distinguished from *H. panicea* and are therefore morphologically distinct from the newly described species as well.

In practice, informal diagnosis of all species is feasible by sequencing the D1-D3 region of the 28S locus and constructing a phylogeny with the provided sequences. However, clade membership alone is not sufficient for formal description under the Code of Zoological

638   Nomenclature, and not all species exhibit unique derived states at this locus. Consequently,
639   formal diagnosis requires sequencing different loci for some species. Morphological traits
640   should be employed to diagnose species when comparing them to previously described
641   *Halichondria* that have not been sequenced at these loci.
642
643   **List of species discussed in systematics section**
644   Genus Halichondria
645   Halichondria panicea (Pallas, 1766)
646   Halichondria akesa sp. nov.
647   Halichondria sitiens (Schmidt, 1870)
648   Halichondria hygeia sp. nov.
649   Halichondria californiana sp. nov.
650   Halichondria cascadia sp. nov.
651   Halichondria columbiana sp. nov.
652   Halichondria dokdoensis Kang, Kim & Kim 2022
653   Halichondria galea sp. nov.
654   Halichondria bowerbanki Burton, 1930
655   Halichondria urca sp. nov.
656   Halichondria zabra sp. nov.
657   Halichondria pinaza sp. nov.
658   Halichondria loma Turner & Lonhart 2023
659
660   Genus Hymeniacidon
661   Hymeniacidon perlevis (Montagu, 1814)
662   Hymeniacidon actites (Ristau, 1978)
663   Hymeniacidon ungodon de Laubenfels 1932
664   Hymeniacidon kitchingi (Burton 1935)
665   Hymeniacidon pierrei sp. nov.
666   Hymeniacidon fusiformis Turner & Lonhart 2023
667   Hymeniacidon globularis Ott, McDaniel & Humphrey 2024
668
669   Genus Semisuberites
670   Semisuberites cribrosa (Miklucho-Maclay, 1870)
671
672   Genus Topsentia
673   Topsentia fernaldi (Sim & Bakus, 1986)
674   Topsentia disparilis (Lambe 1894)
675
676   Genus Axinyssa
677   Axinyssa piloerecta sp. nov.
678   Axinyssa tuscara (Ristau 1978)
679
680   **Genus *Halichondria***
681

682 *Halichondria panicea* **(Pallas, 1766)**
683 Figures 6 & 7
684
685 **Synonyms**
686 Spongia panicea Pallas, 1766
687 The World Porifera Database lists over 70 additional synonyms (de Voogd, et al., 2024)
688
Material examined
160504, Spettan, Gullmarsfjorden, Sweden, (58.25770, 58.25770), 18 m, 2015-05-04; 160505,
Spettan, Gullmarsfjorden, Sweden, (58.25770, 58.25770), 18 m, 2015-05-04; 180911, Svartejan,
Idefjord, Sweden, (59.11020, 11.32190), 30 m, 2018-09-11; 180923, Lunneviken, Idefjord,
Sweden, (59.05460, 11.16900), 27 m, 2018-09-23; 181204, Svartejan, Idefjord, Sweden,
(59.11020, 11.32190), depth not recorded, 2018-12-04; 190424, Flatholmen, Gullmarsfjorden,
Sweden, (58.26080, 11.39820), 14 m, 2019-04-24; 190624, Trindeknubben, Idefjord, Sweden,
(58.78290, 10.99620), 12 m, 2019-06-24; 191026, Gamla Svinesundsbron, Idefjord, Sweden,
(59.09770, 11.26860), 19 m, 2019-10-26; 191116, Yttre Vattenholmen, Idefjord, Sweden,
(58.87540, 11.10560), 27 m, 2019-11-16; BHAK-00536, Fifth Beach, Shatner Point, Calvert
Island, British Columbia, (51.63870, -128.15700), intertidal, 2017-07-25; BHAK-03453, Crazy
Town, North Beach, Calvert Island, British Columbia, (51.66680, -128.13400), intertidal, 2017-
08-08; ESS11, Brig's Shoal, Eastern Shore Islands, Nova Scotia, (44.62781, -62.93603), 25 m ,
2021-07-19; ESS17, Brig's Shoal, Eastern Shore Islands, Nova Scotia, (44.62781, -62.93603), 25
m, 2021-07-19; ESS18, Brig's Shoal, Eastern Shore Islands, Nova Scotia, (44.62781, -62.93603),
25 m , 2021-07-19; ESS74, South of Brig's Shoal, Nova Scotia, (44.60336, -62.92724), 29 m,
2021-07-23; ESS80, South of Brig's Shoal, Nova Scotia, (44.60336, -62.92724), 29 m , 2021-07-
23; FH12x36, Cattle Point, San Juan Island, Washington, (48.45030, -122.96237), intertidal
(rocks), 2012-05-19; FH12x38, Cattle Point, San Juan Island, Washington, (48.45030, -
122.96237), intertidal (rocks), 2012-05-19; FH12x39, Cattle Point, San Juan Island, Washington,
(48.45030, -122.96237), intertidal (rocks), 2012-05-19; FH12x81, Eagle Cove, San Juan Island,
Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22; FH12x84, Eagle Cove, San
Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22; FH12x85, Eagle
Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22;
FH12x86, Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks),
2012-05-22; NIB0608, Carrick-a-Rede, Northern Ireland, (55.24238, -6.33396), 10m (growing on
kelp stipe), 2021-10-17; Quoddy113, Bald Head, Campobello, New Brunswick, (44.91220, -
66.95215), 11.6 m , 2022-10-31; Quoddy124, Marble Island, Off Indian Island, Western Isles,
New Brunswick, (44.91921, -66.97054), 10.6 m, 2022-11-01; TAT026, Strawberry Draw
Overhang, Tatoosh Island, Washington, (48.39190, -124.73800), intertidal, 2018-06-16; TAT031,
Toad Point, Tatoosh Island, Washington, (48.38970, -124.73500), intertidal, 2018-06-17; TAT032,
Toad Point, Tatoosh Island, Washington, (48.38970, -124.73500), intertidal, 2018-06-17;
TLT1293, Cape Alava, Washington, (48.17134, -124.75227), 4-12 m, 2023-07-29; TLT1370, Cape
Johnson, Washington, (47.97327, -124.68220), 4-7 m, 2023-07-27; TLT1371, Cape Johnson,
Washington, (47.97327, -124.68220), 4-7 m, 2023-07-27; TLT1430, Neah Bay, Washington,
(48.38676, -124.63223), 6-14 m, 2023-08-01; TLT619, Scott's Creek Beach, California, (37.04213,
-122.23366), beach wrack, 2020-11-14; TLT858, Cave Landings, California, (35.17535, -

120.72240), intertidal, 2021-02-06; ZMA6155, Jenny's Cove, Lundy Island, England, 24 m,
1985-01-08.

729
730 Figure 6. *Halichondria panicea*. Northwest Atlantic: A (ESS80), B (ESS17), D (ESS18), F (Quoddy124), H
731 (ESS11). Northeast Pacific: C (TLT1430), E (TLT1293), G (TLT1371). All are photographed in situ; C shows
732 scar where sample was taken to display internal color.

Diagnosis

Yellow or green *Halichondria* with a tangential ectosomal skeleton developed as a dense sieve-like mat of spicules or a lace-like network of spicule tracts. Choanosomal skeleton is confused with vague spicule tracts, sometimes accompanied by well-developed subectosomal spicule tracts descending from the ectosomal skeleton. Spicules oxeas, ranging from 100–500 μm in length, with averages per sample of 200–400 μm in length and 5–10 μm in width; usually slightly shorter in the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. 754:A, 952:A. N=55–69 depending on position. These characters differentiate *H. panicea* from all other sequenced *Halichondria* except *H. sitiens,* which is diagnosed relative to *H. panicea* based on morphological characters such as spicule length and surface papillae.

747 28S. No unique derived states shared by all *H. panicea*, but the Atlantic Ocean population has
748 two unique states, 605:T and 1549:C. N=13–21 depending on position
749
750 ND1. No unique derived states shared by all *H. panicea*, but the Pacific Ocean population has a
751 unique state, 451:T. N=10).
752
753 **Morphology**
754 Thinly or thickly encrusting, sometimes with digitate tendrils (Figure 6). Oscula usually
755 prominent, but vary from raised rims nearly flush with surface to tall oscular chimneys. Color
756 alive varies from pale translucent yellow to golden, often variegated with green, or with the
757 exterior entirely green or gray-green and the interior yellow. White or beige in ethanol.
758
Skeleton
Tangential ectosomal skeleton can be sieve-like, with a dense mat of spicules surrounding open
areas that are mostly circular or oval shaped (Figure 7A, 7F), or more lace-like, with a web of
spicule tracts surrounding open areas that tend towards triangular (Figure 7C, 7D). In some
samples, this tangential ectosomal skeleton is supported by prominent spicule tracts that rise
from the choanosome to fan out tangentially in the ectosome (Figure 7B, 7G). The choanosomal
skeleton in these samples becomes more confused towards the interior (Figure 7H). Other
samples have a confused jumble of multispicular tracts, spicule bundles, and individual spicules
both near the ectosome and interior (Figure 7I). No apparent spongin.
768
Spicules
Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have a typical halichondroid shape: evenly curved or with a slight central bend, thickest
in the center and tapering gently to sharp points at both ends. Spicules have a large range in
lengths in both the ectosome and the choanosome, but average 15.4% longer in the
choanosome. Several samples (ESS11, 190424) had no difference in length by compartment
(Figure S4). Dimensions are given for individual samples in table 1. When all samples are
combined, spicules measure 114–272–499 x 2–8–16 µm, n=1502. Average values varied
considerably across samples, with mean lengths from 202 to 358 µm, and mean widths from 5
to 10 µm.
779
Distribution and habitat
This species has been reported to occur from Iceland to the Mediterranean in the Northeast
Atlantic, from Northern Canada to New England in the Northwest Atlantic, and possibly the
Arctic and North Pacific (Erpenbeck and van Soest, 2002). DNA data is able to confirm only the
Northern portion of this range in the Northeast Atlantic, including Iceland, Sweden, the United
Kingdom, and the Netherlands. Genetic support in the Northwest Atlantic ranges from New
Brunswick to New Hampshire. Sequenced samples from farther South in the Atlantic (Portugal,
Connecticut, New York, Virginia) have thus far proved to be in the *Hygeia* or *Bowerbanki* species
groups, but additional sampling could certainly extend this range. In the North Pacific, DNA-
confirmed range is from the Bering Strait to Central California, with no DNA confirmation
available from the Northwest Pacific. The species is rare or absent from shallow waters in

Southern California, as the genotyping of over 100 *Halichondria* samples from this region found
none to be *H. panicea*.
Previously, this species was described as common on rocks and brown algae in the
intertidal and shallow subtidal zones, preferring exposed areas over sheltered bays, and
extending down to a depth of 500 meters (Erpenbeck and van Soest, 2002). DNA-confirmed
samples align with this description, although we cannot confirm a depth range beyond 29
meters. The species is common in exposed intertidal areas and the shallow subtidal zone but is
rare or absent on human-made structures like floating docks in marinas, where *H. hygeia* and
members of the *Bowerbanki* species group are prevalent.
800
801 Table 1. Spicule lengths of *Halichondria panicea*. Values given as min–mean–max in μm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
BHAK-00536	114–185–285 n=40	130–217–322 n=46
BHAK-03453	153–204–256 n=40	170–241–322 n=40
TLT619	130–270–451 n=40	199–334–442 n=40
TLT1370	172–286–451 n=45	146–364–499 n=50
TLT1430	147–233–353 n=96	162–299–367 n=98
FH12x38	155–247–335 n=49	169–272–377 n=50
FH12x39	134–247–357 n=42	169–291–367 n=47
FH12x81	145–238–351 n=55	166–262–354 n=54
Atlantic Ocean		
NIB0608	176–257–322 n=49	187–281–365 n=51
ZMA-6155	149–283–408 n=40	241–354–458 n=40
190424	178–273–347 n=40	196–267–342 n=40
180911	173–247–349 n=49	221–296–357 n=41
Quoddy124	212–320–421 n=48	298–397–463 n=44
ESS11	185–254–318 n=86	167–254–331 n=86

802
Remarks
As discussed in the genetic section, *H. panicea* from the Atlantic and Pacific basins form sister
clades that are distinguishable at the genetic level and could potentially be considered separate
species. However, if *H. panicea* occurs throughout the Arctic, it may link the populations across
these basins. We conservatively retain both populations under the same species concept
pending further data.
We lack genetic confirmation of *H. panicea* in the Northwest Pacific, but its presence
there seems likely, as we have confirmed its distribution in the Aleutian Islands and the Bering
Strait. This species has also been reported from various other global regions, including South
America, New Zealand, and the tropical South Pacific (gbif.org). Several lines of evidence suggest
that these records may actually pertain to other species, such as *H. hygeia*, members of the
Bowerbanki species group, or additional cryptic species that remain to be discovered. First, *H.*
panicea seems less likely than those species to be experiencing a human-assisted range
expansion for two reasons: 1) it does not seem to occur in human-modified habitats like

marinas, and 2) genetic differentiation between Atlantic and Pacific populations make it likely
that these populations do not share a common ancestor in the modern era. Second, the species
does not seem to tolerate warm temperate waters in Southern California, and is therefore
unlikely to occur in shallow tropical waters.
821

Figure 7. *Halichondria panicea* skeleton and spicules. Ectosomal skeletons: A (ESS80), B (TLT1430), C
(FH12x38), D (Quoddy124), E (NIB608), F (TLT619). B is imaged from below to show the twisting columns
of spicules connecting the ectosome to the choanosome in this sample. G: cross section at sponge
surface, showing spicule columns and subectosomal spaces (TLT1430), H: cross section of same sample
(TLT1430) farther towards the inferior, showing confused skeleton, I: cross section at sponge surface
without large subectosomal spaces (TLT1370), J: oxeas (TLT1293).
829

830 *Halichondria akesa* **sp. nov.**
831 Figure 8
832

833 **Material examined**
834 Holotype: ESS07, Brig's Shoal, Eastern Shore Islands, (44.62781, -62.93603), 25 m, 2021-07-19.
835

836 **Etymology**
837 Name inspired by the Greek goddess Akeso, sister of Panakeia.

Diagnosis

Yellow and green *Halichondria* with a tangential ectosomal skeleton developed as a thick mat of spicules and/or a lace-like network of spicules and spicule tracts. Choanosomal skeleton consists of prominent spicule tracts near the ectosome, transitioning to paucispicular tracts and spicules in confusion deeper in the choanosome. Spicules oxeas, ranging from 90–460 µm in length and 2–10 µm in width; slightly shorter in the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 621:T. N=1.

ND1. 39:T, 55:T. N=2.

Morphology

The single specimen examined was thickly encrusting with prominent oscula. Yellow alive, white in ethanol.

Figure 8. *Halichondria akesa*. A: field photos, B: tangential section showing ectosomal skeleton and subectosomal spicule tracts, C: oxeas. All photos from the holotype.

Skeleton

The single specimen examined has a tangential ectosomal skeleton which was a dense mat of spicules in some areas, and a network of tracts and single spicules surrounding oval-shaped open areas in other places. The ectosomal skeleton is supported by prominent spicule tracts that rise from the choanosome to fan out tangentially in the ectosome (Figure 8B, 8G). The choanosomal skeleton becomes more confused towards the interior, with some spicule tracts but many spicules in confusion. No apparent spongin.

Spicules

Nearly exclusively oxeas, though rare styles can be found. Oxeas have the typical halichondroid shape: gently curved or with a slight central bend, thickest in center and tapering gently to sharp points at both ends. Spicules have a large range in lengths in both the ectosome and the choanosome, but averaged 9% longer in the choanosome. Ectosomal spicules 110–256–438 x 4–6–10 µm, n=44; choanosomal spicules 98–281–459 x 2–6–9 µm, n=44; all spicules together

875 98–268–459 x 2–6–10 μm, n=88. Spicules likely to be indistinguishable from *H. panicea*, though
876 the minimum length was very slightly smaller than seen in any examined *H. panicea*.
877
878 **Distribution and habitat**
879 Currently known from three samples, all from the far North Atlantic / Arctic: Eastern Shore
880 Islands, Nova Scotia, Canada; Skútan, Northern Iceland; and Svalbard, Norway. Habitat
881 preferences are likely similar to *H. panicea* – though all three samples were collected from the
882 subtidal by divers, so it is possible this species does not occur in the intertidal.
883
Remarks
Though we were only able to examine one sample of this new species, the published
mitochondrial genome (MH756604) of another sample from Iceland is nearly identical. When
the entire mitochondrial genome is aligned (coding and non-coding regions), these two samples
differ at only 11 of 12,851 aligned sites (0.09%). In contrast, the average pairwise differences
(Dxy) between the two *H. akesa* mitochondria and 9 sequenced *H. panicea* mitochondria is
0.8%, leading to an Fst of 0.745 across the mitochondrial genome between species.
Furthermore, the ribosomal phylogeny and the phylogeny of concatenated mitochondrial genes
confidently place true *H. panicea* sequences closer to *H. sitiens* than to *H. akesa sp. nov.*. We
therefore feel confident in describing these samples as a new species that is sympatric with *H.*
panicea across the far North Atlantic. The sample from Nova Scotia was collected at the same
time and place as three other samples that were found to be *H. panicea*, so these species are
sympatric even at the scale of individual sites.
Assignment of the Svalbard sample to this species is based on only 610 bp of the cox1
gene (boldsystems.org sample ROPOR090-18). This sequence is, however, identical to the other
H. akesa samples, and has several SNPs that differentiate it from *H. panicea* and other similar
species.
901
902 *Halichondria sitiens* **(Schmidt, 1870)**
903 Figure 9
904
Material examined
ESS130, Faulker's Shoal, Nova Scotia, (44.60733, -62.96168), 30 m , 2021-08-12; ESS132,
Faulker's Shoal, Nova Scotia, (44.60733, -62.96168), 30 m, 2021-08-12; ESS153, N. of Bull Rock,
Nova Scotia, (44.65375, -62.90491), 30 m , 2021-08-14; ESS44, Bull Shoal, Nova Scotia,
(44.63589, -62.89906), 28 m , 2021-07-21; RBC11, Knight Inlet, British Columbia, (50.68577, -
125.99457), 9 m, 2009-11-08; RBC13, Kuldekduma Island; Weynton Passage, British Columbia,
depth not known, 2019-04-12; RBC8, Knight Inlet, British Columbia, (50.68500, -126.03667), 15-
20 m, 2011-03-29.
913
914 **Diagnosis**
915 Yellow, white, green, or slightly blueish *Halichondria* with numerous papillae decorating the
916 surface. Scattered oscula usually visible, but often not at the ends of papillae. Tangential
917 ectosomal skeleton developed as a web of meandering spicule tracts and individual spicules.
918 Choanosomal skeleton consists of multispicular tracts and many spicules in confusion. Spicules

oxeas, 130–1000 μm long, with averages per sample of 440–630 μm in length and 9–10 μm in width; usually shorter in the ectosome than the choanosome. Additional DNA characters that may be useful in identification are as follows:

COX1. 754:A, 952:A. N=6. These characters differentiate *H. sitiens* from all other sequenced *Halichondria* except *H. panicea,* which is diagnosed relative to *H. sitiens* based on morphological characters such as spicule length and surface papillae.

28S. 2301:G. N=2.

ND1. 97:C. N=5.

Figure 9. *Halichondria sitiens*. A (ESS44), B (ESS132), C (ESS153), D (RBC13, photo by Neil McDonald). E: cross-section showing choanosomal skeleton (ESS153), F: oxeas (RBC8). Tangential ectosomal skeletons: G (ESS130), H (ESS44).

Morphology
Thinly to thickly encrusting or growing as a semi-globular cushion. Surface is often replete with cone-shaped papillae that sometimes branch, but these are less apparent in small sponges. A few scattered oscula are usually visible, often large and prominent and sometimes atop oscular

940   chimneys, but not found at the ends of the papillae. Color generally a pale, translucent yellow
941   but surface can appear nearly white, and field photos of sponges that are likely this species have
942   also documented pale green and slightly bluish individuals. White in ethanol.
943
Skeleton
Tangential ectosomal skeleton is well-developed, but less dense and regularly-patterned than
observed in *H. panicea*. Instead, the ectosomal skeleton is comprised of a mesh of meandering
multispicular tracts and individual spicules, with spaces between them variable in size and
shape (similar to species in the *Bowerbanki* group). Choanosomal skeleton typically
halichondroid, with ill-defined spicule tracts and many spicules in confusion. No apparent
spongin.
951
Spicules
Exclusively oxeas with typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering gently to sharp points at both ends. Spicules have a large
range in lengths in both the ectosome and the choanosome, but average 14% longer in the
choanosome. The difference between ectosome and choanosome is quite variable, with up to
30% longer or 5% shorter spicules in the choanosome. Dimensions are given for individual
samples in table 2. When all samples are combined, spicules measure 138–531–1000 x 4–12–25
µm, n=616. Average values varied considerably across samples, with notably longer spicules
seen in Pacific Canada (means 552–628 µm) than Atlantic Canada (means 448–531 µm).
961
962   Table 2. Spicule lengths of *Halichondria sitiens*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
RBC13	318–533–868 n=47	297–724–1000 n=47
RBC11	335–582–892 n=40	332–648–964 n=40
RBC8	308–505–844 n=41	295–597–893 n=43
Atlantic Ocean		
ESS130	312–498–623 n=53	325–455–619 n=59
ESS153	270–466–625 n=40	302–518–698 n=40
ESS44	138–398–680 n=43	332–497–659 n=43
ESS132	258–468–728 n=40	370–563–723 n=40

963
964
Distribution and habitat
Known from the Arctic, North Atlantic, and North Pacific, including Greenland, Svalbard, the
White Sea, Alaska, and both coasts of Canada. Samples examined here were from British
Columbia in the Pacific and Nova Scotia in the Atlantic, and these are the only samples with
DNA data available. Previously described as occurring to 100 m depth (Erpenbeck and van
Soest, 2002), but the samples examined here were all collected in the shallow subtidal by
divers.
972

Remarks

It is remarkable that *H. sitiens* is the closest relative of *H. panicea*, of all the species with DNA data available. Many of the species described here are morphologically indistinguishable from *H. panicea*, while *H. sitiens* has longer spicules and surface papillae that distinguish it. The presence of surface papillae has been used as a genus level trait in the past, later downgraded to the subgenus *Eumastia.* Given the close relationship between *H. panicea* (type species of subgenus *Halichondria*) and *H. sitiens* (type species of subgenus *Eumastia*), we recommend eliminating the subgenera.

In contrast to *H. panicea*, where fixed genetic differences were seen between Atlantic and Pacific populations, *H sitiens* has nearly no genetic variation within or between populations at any of the sequenced loci.

Halichondria hygeia **sp. nov.**

Figure 10

Material examined

Holotype: TLT645, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16. Paratypes: BHAK-03597, Pruth Bay dock, Calvert Island, British Columbia, (51.65500, -128.12900), floating dock, 2017-08-08; FH12x108, Roche Harbor, San Juan Island, Washington, (48.60970, -123.15436), floating dock, 2012-05-23; FH12x114, Roche Harbor, San Juan Island, Washington, (48.60970, -123.15436), floating dock, 2012-05-23; TLT1076, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1417, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2023-09-01; TLT308, Monterey Marina, California, (36.60797, -121.89288), floating dock, 2019-11-22; TLT504, Point Pinos, California, (36.63447, -121.93923), intertidal, 2019-11-24; TLT706, Oceanside Harbor, California, (33.20623, -117.39208), floating dock, 2021-01-09; TLT724, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2020-12-14; TLT742, Dana Point Harbor, California, (33.46090, -117.69156), floating dock, 2021-01-09; TLT940, Lazy Days, San Diego, California, (32.69415, -117.27110), 12-25 m, 2021-05-15; WMS19x5, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2019-05-19; WMS19x6, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2019-05-19; WMS19x7, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2019-05-19; WMS21x10, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x19, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x6, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x7, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x8, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x9, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS23x10, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2023-06-07; WMS23x9, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2023-06-07.

Etymology

1017   Name inspired by the Greek goddess Hygeia, sister of Panakeia.
1018

1019
1020   Figure 10. *Halichondria hygeia*. A & B (TLT504), C (TLT645), D (TLT706). E: cross-section showing
1021   choanosomal skeleton at sponge surface (TLT504), F: spicules (FH12x108). Tangential ectosomal
1022   skeletons: G (TLT742), H (TLT645).
1023

Diagnosis
Yellow *Halichondria* with well-developed tangential ectosomal skeleton manifested as a dense
mat of spicules or a mesh of meandering multispicular tracts and individual spicules.
Choanosomal skeleton confused, sometimes accompanied by multispicular tracts. Spicules
oxeas, 170–430 µm in length, with averages per sample of 250–350 µm in length and 5–10 µm
in width. On average, no difference in length is seen between the ectosome than the
choanosome. Diagnosis relative to similar species requires DNA characters, as follows:
1031

1032   COX1. No unique derived states.
1033

1034   28S. 659:G, 670-671:GG, 675:A. N=24.
1035

1036 ND1. 877-880:AACC. State 877:A is unique compared to all close relatives, but homoplastic with
1037 *Halichondria loma.* The full sequence 877-880:AACC is not shared with any other sequenced
1038 *Halichondria*. N=14.
1039
1040 **Morphology**
1041 Thinly to thickly encrusting, sometimes with digitate tendrils. Oscula vary from not apparent, to
1042 slightly volcano-shaped with raised rims, to atop oscular chimneys. Color varies from pale
1043 yellow to golden; not seen in green. See Figure 10 for examples.
1044
Skeleton
Tangential ectosomal skeleton can be sieve-like, with a dense mat of spicules surrounding oval-
shaped open areas (Figure 10G), or more lace-like, with a web of spicule tracts and individual
spicules delineating more irregularly shaped open areas (Figure 10H). This ectosomal skeleton is
supported by poorly developed spicule tracts that delineate large subectosomal spaces, with
the choanosomal skeleton becoming more confused towards the interior. No apparent spongin.
1051
Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering to sharp points at both ends. Spicules have a large range in
lengths, though slightly less range than seen in *H. panicea*, and spicules in the ectosome and the
choanosome average the same length. Dimensions for five samples where ectosome and
choanosome were measured separately are given in table 3. When spicules from all seven
measured samples are combined, spicules measure 175–276–424 x 3–7–13 μm, n=561 for
length, n=405 for width. Average values varied considerably across samples, with mean lengths
254–346 μm, and mean widths 5–10 μm.
1061
1062 Table 3. Spicule lengths of *Halichondria hygeia*. Values given as min–mean–max in μm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
FH12x108	284–343–386 n=40	236–349–424 n=40
TLT645	194–259–313 n=40	216–269–312 n=40
TLT742	190–267–301 n=40	193–259–319 n=43
TLT706	188–284–343 n=42	232–279–328 n=43
Atlantic Ocean		
WMS21x9	246–283–336 n=40	204–256–302 n=44

1063
Distribution and habitat
In the Northeast Pacific, this species was found from Calvert Island, British Columbia to
Oceanside, Southern California. In contrast to *H. panicea*, *H. hygeia* was found primarily on
floating docks in sheltered marinas, but it was not limited to this habitat. Of 12 samples
collected in the Northeast Pacific, 10 were on floating docks but one was in found in a wave-
swept tide pool at Point Pinos (Monterey Bay, Central California), and one was found in subtidal
rocky reef off Point Loma (extreme Southern California, collected between 12 and 25 m depth).

The species is also found in the North Atlantic, as it was common on intertidal cobbles at West Meadow Beach, Stony Brook, Long Island, New York. Publicly available sequences of cox1 may indicate that the species is also found in Connecticut and Portugal (Genbank accessions MN448237, MZ487266, figure S1). These sequences are consistent with either *H. hygeia* or *H. californiana*; as *H. californiana* is thus far known only from California, these sequences are likely to be *H. hygeia*, but confirmation would require sequencing 28S or ND1.

Remarks

Two pieces of evidence suggest that *H. hygeia* has experienced a recent human-assisted range expansion. First, artificial substrates like floating docks — where this species is primarily found — are known to harbor many introduced species (Bulleri and Chapman, 2010; Glasby et al., 2007; Ruiz et al., 2000; Simkanin et al., 2012). Second, samples from New York and California are nearly identical across the mitochondrial genes and the ribosomal DNA (Figure 1). Other halichondriids have been shown to have limited larval dispersal (Maldonado, 2006; Xue et al., 2009), so together, these observations make it likely that the species has been moved around by human activity.

Halichondria hygeia and *H. panicea* are likely to be indistinguishable with respect to spicule and skeletal traits, but some differences between then are apparent. These include habitat (*H. hygeia* is common on floating docks in marinas, while *H. panicea* is common in the wave-exposed intertidal), southern range limit (*H. hygeia* was found in Southern California, while *H. panicea* was not found in Southern California), and color (*H. hygeia* is yellow while *H. panicea* can be yellow or green).

Halichondria californiana **sp. nov.**
Figure 11

Material examined

Holotype: TLT937, Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15. Paratypes: TLT1010, Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-11 m, 2021-07-22; TLT1093, Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-11 m, 2021-07-22; TLT160, Isla Vista Reef, California, (34.40278, -119.85755), 9-12 m, 2019-08-01; TLT256, Isla Vista Reef, California, (34.40278, -119.85755), 9-12 m, 2019-08-01; TLT417, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2019-10-25; TLT428, Elwood Reef, California, (34.41775, -119.90150), 9-15 m, 2019-10-23; TLT812, Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15; TLT860, Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06; TLT921, Point Fermin, California, (33.70664, -118.28595), intertidal, 2021-03-10; TLT928, Point Fermin, California, (33.70664, -118.28595), intertidal, 2021-03-10; TLT934, Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15.

Etymology

Named for the state of California.

Diagnosis

Yellow *Halichondria* with tangential ectosomal skeleton developed as a sieve-like mat of spicules or a lace-like network of spicule tracts. Choanosomal skeleton consists of poorly delineated multispicular tracts and spicules in confusion. Spicules oxeas, ranging from 90–290 μm in length, with averages per sample of 130–230 μm long and 3–7 μm wide. On average, no difference in length is seen between the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 1011:A, 1037:T, 1319:C. N=9–11 depending on position.

ND1. 214-217:AAAT. State 217:T is unique compared to all close relatives, but homoplastic with members of the Okadai group (fig). The full sequence 214–217:AAAT is not shared with any other sequenced *Halichondria*. N=10.

Figure 11. *Halichondria californiana*. Field photos: A (TLT937), B (TLT880), C (TLT1093). Ectosomal skeletons: D (TLT160), E (TLT937), F (TLT1010), G (TLT860). H: cross-section showing choanosomal skeleton at sponge surface (TLT160), I: spicules (TLT860).

Morphology

1136 Encrusting yellow sponges; oscula often small and flush with sponge surface, but can also occur
1137 on oscular chimneys. Color varies from pale, translucent yellow to golden, sometimes with red
1138 blotches; not seen in green. Beige in ethanol. See Figure 11 for examples.
1139

1140 **Skeleton**
1141 Tangential ectosomal skeleton can be sieve-like, with a dense mat of spicules, or more lace-like,
1142 with a web of spicule tracts and single spicules. Open areas are generally oval or polygonal and
1143 relatively regular in size. Choanosomal skeleton with poorly developed spicules tracts and many
1144 spicules in confusion. No apparent spongin.
1145

Spicules
Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have a typical halichondroid shape: evenly curved or with a slight central bend, thickest
in the center and tapering gently to sharp points at both ends. Spicules have a large range in
lengths, though slightly less range than seen in *H. panicea*, and spicules in the ectosome and the
choanosome average the same length. Dimensions for the six samples where ectosome and
choanosome were measured separately are given in table 4. When spicules from 10 measured
samples are combined, spicules measure 99–194–286 x 2–5–9 µm, n=813 for length, n=733 for
width. Average values varied considerably across samples, with means of 134–226 µm long and
3–7 µm wide.
The maximum length of 286 µm is notably shorter than some similar species, such as *H.
hygeia* (424 µm), *H. panicea* (499 µm), and *H. cascadia* (444 µm) — though *H. columbiana* is
similar (266 µm). Spicule size is insufficient to differentially diagnose all samples, however. For
example, some individual *H. californiana* samples had average and maximum lengths exceeding
some *H. hygiea* samples, and some *H. californiana* had average lengths exceeding some *H.
panicea*.
1162

1163 Table 4. Spicule lengths of *Halichondria californiana*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT937	149–207–239 n=40	177–215–259 n=40
TLT1010	113–136–172 n=51	99–132–161 n=51
TLT921	121–177–199 n=40	145–180–208 n=40
TLT256	107–146–164 n=40	122–155–190 n=41
TLT860	187–235–274 n=40	190–236–286 n=59
TLT428	167–213–258 n=40	166–218–266 n=40

1164

Distribution and habitat
Known only from California, from San Luis Obispo Bay in Central California to Point Loma in
Southern California. Nearly allopatric with *H. panicea*: the two species were found growing side-
by-side in a tidal pool near Avila Beach, California, but this was the Southernmost location that
H. panicea was found and the Northernmost location for *H. californiana.* Like *H. panicea,* this
species was found growing on rock or algae, in natural subtidal reefs and exposed intertidal

areas, and not found on human structures in sheltered marinas. The deepest samples were
found between 9 and 18 m.
1173
Remarks
In contrast to its closest known relative *H. hygeia*, this species was found only in natural
habitats, and not on human structures. It is also different from *H. hygeia* in that it was only
found in California and thus could be endemic to the Northeast Pacific. However, these two
species cannot be differentiated at the commonly sequenced regions of cox1, so publicly
available sequences from Connecticut and Portugal could be this species (Genbank accessions
MN448237, MZ487266). The nudibranch *Doris montereyensis* was seen feeding on this species
in multiple locations.
1182
1183 ***Halichondria cascadia* sp. nov.**
1184 Figure 12
1185

1186
1187 Figure 12. *Halichondria cascadia*. A: sample post-preservation (FH12x92), B: tangential ectosomal
1188 skeleton (FH12x93), C: choanosomal skeleton (FH12x92).
1189
1190 **Material examined**
1191 Holotype: FH12x92, Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal,
1192 2012-05-22. Paratype: FH12x93, Eagle Cove, San Juan Island, Washington, (48.46128, -
1193 123.03155), intertidal, 2012-05-22.
1194
1195 **Etymology**
1196 Name inspired by the term "Cascadia region" which is sometimes used to refer to an area
1197 including Oregon, Washington, and British Columbia.
1198
Diagnosis
Halichondria with a tangential ectosomal skeleton developed as a mesh of meandering
multispicular tracts and individual spicules. Choanosomal skeleton consists of spicules and
spicule bundles in confusion. Spicules oxeas, 160–450 µm long, with averages per sample of
270–320 µm long and 8-9 µm wide. Infrequent styles and rare strongyles can be observed.
Some samples average shorter spicules in the ectosome than the choanosome. Diagnosis based
primarily on DNA characters, as follows:
1206
1207 COX1. 949:A. N=3.

28S. 2061:T. N=2.

ND1. 847:A. N=2.

Morphology
Known only from two small samples, so little is known about the morphology of this species. Preserved samples are white; color alive is yellow. Both samples have oscular chimneys approximately 1 cm in height.

Skeleton
Tangential ectosomal skeleton a web of spicule tracts and individual spicules, with open areas irregular in size and shape (Figure 12B). Choanosomal skeleton of spicules and spicule bundles in confusion. No apparent spongin.

Spicules
Oxeas with a typical halichondroid shape: evenly curved or with a slight central bend, thickest in the center and tapering to sharp points at both ends. Infrequent styles and rare strongyles can be observed. Spicules have a large range in lengths, with one sample having significantly shorter spicules in the ectosome while the other sample has no difference in length between ectosome and choanosome. When spicules from both samples are combined, spicules measure 162–294–444 x 3–8–12 µm, n=169. Table 5 shows dimensions for samples and domains separately.

Table 5. Spicule lengths of *Halichondria cascadia*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
FH12x92	178–267–353 n=41	162–273–373 n=46
FH12x93	199–287–280 n=42	260–353–444 n=40

Distribution and habitat
Samples examined here were both collected from rocky intertidal habitats at Eagle Cove, San Juan Island, Washington. Publicly available sequence data suggests a third sample was found at Carl Moses Harbor, Captains Bay, Unalaska Island (Genbank accession MZ580608). Though only 658 bp of cox1 is available for this sample, it is identical to sample FH12x93.

Remarks
The two samples examined here are small fragments. Though they were collected at the same location, they are genetically different from one another: they differ at 1.17% of bases in the nuclear ribosomal locus and at 0.6% of bases in the protein-coding mitochondrial genome. They form a well-supported clade in the phylogeny, with 98.5% bootstrap support, and are the sister group to the other three species in the *Hygeia* group (*H. hygeia, H. californiana, and H. columbiana*). They were collected at the same time and place as *H. panicea*, and *H. hygiea* was collected at a marina on the same island the following day, so this species is sympatric with several closely related *Halichondria.* Despite the limited sampling of this species, the strong

1248 support for *H. hygiea* and *H. californiana* as different species compels us to formally describe
1249 these samples in the hope that more can be learned about this enigmatic group.
1250
1251 ***Halichondria columbiana* sp. nov.**
1252 Figure 13
1253
1254 **Material examined**
1255 Holotype: BHAK-09466, Port Reef, Hakai Pass, British Columbia, (51.697, -128.106), 28 m, 2019-
1256 05-28. Paratype: BHAK-09479, Whidbey Point, Fitz Hugh Sound, British Columbia, (51.71700, -
1257 127.89200), 19 m, 2019-05-29.
1258
1259 **Etymology**
1260 Named for the British Columbia region of Canada.
1261
1262 **Diagnosis**
1263 *Halichondria* with tangential ectosomal skeleton developed as a chaotic mesh of meandering
1264 multispicular tracts and individual spicules. Choanosomal skeleton confused. Spicules oxeas,
1265 150–270 µm long, with means per sample of 220–240 µm long and 8–9 µm wide. Diagnosis
1266 based primarily on DNA characters, as follows:
1267
1268 COX1. 973:C, 1012:C, 1102:G, 1207:A. N=2.
1269
1270 28S. No unique derived states.
1271
1272 ND1. 79:C, 88:C, 172:T, 340:G, 352:C, 733:G, 811:G. N=2.
1273

1274
1275 Figure 13. *Halichondria columbiana*. Photos of sponges after collection, before preservation: A (BHAK-
1276 9466), B (BHAK-9479). C: tangential ectosomal skeleton (BHAK-9466).
1277
1278 **Morphology**
1279 Known only from two small samples, little is known about the morphology of this species, but
1280 they appear similar to encrusting *H. panicea*. Samples were yellow alive and are white in
1281 ethanol.
1282
1283 **Skeleton**

1284 Tangential ectosomal skeleton is a web of spicule tracts and individual spicules (Figure 13C).
1285 Choanosomal skeleton difficult to characterize in such small samples, but appears to be
1286 confused, as seen in all closely related species. No apparent spongin.
1287

Spicules
1289 Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend,
1290 thickest in the center and tapering to sharp points at both ends. When spicules from both
1291 samples are combined, spicules measure 149–228–266 x 4–8–11 μm, n=151. Spicules from the
1292 ectosome and choanosome were measured separately for one sample and the sizes did not
1293 differ by compartment (the ectosome could not be located in the other small sample).
1294

Distribution and habitat
1296 The two known samples were collected 15 km apart in British Columbia. Both were subtidal,
1297 between 19 and 28 m depth.
1298

Remarks
The two samples examined here are very small fragments. The samples are identical at the
ribosomal locus and only slightly different at mitochondrial loci (sequence differences of 0.4%
and 0.2% at ND1 and cox1, respectively). Contrastingly, they are genetically distinct from the
other species in the *Hygeia* group: when compared to *H. hygeia,* average pairwise differences
are 1.0% (Fst=0.9) at the ribosomal locus and 1.2% (Fst=0.84) at cox1. Despite the limited
sampling of this species, the strong support for *H. hygiea* and *H. californiana* as different species
compels us to formally describe these samples in the hope that more can be learned about this
enigmatic group.
1308

1309 *Halichondria dokdoensis* **Kang, Kim & Kim, 2022**
1310 Figure 14
1311

1312
1313 Figure 14. *Halichondria dokdoensis*. A: field photo, B: tangential ectosomal skeleton, C: cross-section
1314 showing choanosomal skeleton at sponge surface, D: oxea. All photos from TLT386.
1315

Material examined
1317 TLT386, Oil Platform C, Santa Barbara, California, (34.33293, -119.63173), 21 m, 2019-08-05.
1318

Diagnosis

Yellow, reddish-purple, or orange *Halichondria* with tangential ectosomal skeleton developed as a lace-like network of spicule tracts surrounding oval-shaped open areas. Choanosomal skeleton with multispicular tracts supporting the ectosome and delineating large subectosomal spaces, with an increasingly confused skeleton towards the interior. Spicules oxeas, 130–310 μm long and 1–10 μm wide; slightly shorter in the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. 100:C, 247:G, 361:G, 844:C, 892:T. N=2-3 depending on position.

28S. 162:T, 195:C, 260:G, 584:C, 614:T, 671:T, 1660:G. N=1.

ND1. 286:C, 322:A, 346:G, 403:T, 769:G. N=2.

Morphology
Samples from Korea were previously described as reddish-purple to orange with prominent oscular chimneys (Kang et al., 2022). A sample from California was yellow, with oscula nearly flush with the sponge surface, so color and shape appear to be variable, as seen in many other *Halichondria* species.

Skeleton
Well-developed tangential ectosomal skeleton with a lace-like web of spicules and spicule tracts surrounding open spaces that are oval shaped and fairly consistently sized (Figure 14B). This ectosomal skeleton is supported by multispicular tracts that delineate large subectosomal spaces. The choanosomal skeleton becomes more confused towards the interior. No apparent spongin.

Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend, thickest in the center and tapering gently to sharp points at both ends (rare spicules occasionally found with a centrotylote swelling). The single sponge we examined had spicules 177–230–287 x 2–6–8 μm, n=146 for length, n=106 or width. When measured separately, spicules in the choanosome averaged 238 μm in length vs. 221 μm in the ectosome (p=0.001). Spicules from Korean samples previously reported as 135–310 x 1–8 μm (means and sample size not reported). Spicules were previously reported as having "size classes", with some being thin (1–2.5 μm) and some thick (5–8 μm). Of the 146 spicules we measured, only one would fall in the thin category, with 10 falling between thick and thin. The abundance of thin oxea is therefore variable and/or universally rare, perhaps because they are immature thick spicules.

Distribution and habitat
The holotype location for this species is the Dokdo Islands, South Korea, with additional samples from mainland South Korea and Jeju Island. The sample examined here was found on an oil platform in the Santa Barbara Channel in Southern California, and a publicly available sequence matching this species was obtained near Goes, Zeeland, the Netherlands (see remarks). The

1363 California sample was subtidal (21 m), and though depths are not reported for the other
1364 samples, the Korean samples were also collected by SCUBA divers.
1365
Remarks
Halichondria dokdoensis was described based on morphological characters, but it is our view
that it cannot be morphologically distinguished from *H. panicea*. Characters previously listed as
distinguishing the species were color (but we show here that *H. dokdoensis* can be found in
yellow) and larger spicules than *H. panicea* (but we show here that the spicules of *H. panicea*
span a large size range encompassing that of *H. dokdoensis*). However, a mitochondrial genome
sequence was previously published for *H. dokdoensis*, and we are able to use this sequence to
"rescue" this previously published name (Kim et al., 2019). This sequence is nearly identical to
the mitochondrial genome of our California sample: they differ by only 12 of 13,033 aligned
base pairs (<0.1%). These samples are more closely related to *Hymeniacidon actites* than to any
other sequenced *Halichondria*, and together these two species form a clade that is the sister to
the combined *Panicea* and *Hygeia* species groups.
This species is likely to have experienced a recent human-mediated range expansion.
Finding samples in both California and Korea could plausibly be due to a natural range across
the North Pacific, though this is made less likely by the observation that it was only found (thus
far) on an artificial structure in California. However, a publicly available sequence of cox1
indicates that the species is also likely found in the Netherlands (NLMAR324-20,
boldsystems.org). Though this sequence is only 618 bp of the cox1 gene, it is identical to other
H. dokdoensis and does not match any other sequenced species. This species is therefore likely
to be introduced to multiple regions of the Northern Hemisphere, similar to other species
described here.
1387
1388 ***Halichondria galea* sp. nov.**
1389 Figure 15
1390
1391 **Material examined**
1392 Holotype: TLT495, Point La Jolla, California, (32.85227, -117.27239), intertidal, 2020-02-08.
1393 Paratypes: TLT496, Point La Jolla, California, (32.85227, -117.27239), intertidal, 2020-02-08;
1394 TLT743, Dana Point Harbor, California, (33.46090, -117.69156), floating dock, 2021-01-09.
1395
1396 **Etymology**
1397 Name inspired by the word "galleon", one of the ships of the Spanish armada in the age of sail.
1398
Diagnosis
Encrusting green, yellow, or brownish *Halichondria*, sometimes with a rugose surface.
Tangential ectosomal skeleton is a sieve-like mat of spicules surrounding open areas that are
generally oval-shaped and consistent in size. Choanosomal skeleton of multispicular tracts and
abundant spicules in confusion. Spicules oxeas, 100–500 µm long, with averages per sample of
250–290 µm long and 5–6 µm wide. Spicules 10–20% longer in choanosome than ectosome, but
the full range of spicules lengths is seen in both compartments. Diagnosis relative to similar
species requires DNA characters, as follows:

1407

1408 COX1. 1069:A. N=8.

1409

1410 28S. 516:T, 520:G, 547:G, 1943:T, 2111:T. N=2–3 depending on position.

1411

1412 ND1. No unique derived states.

1413

1414

1415
1416 Figure 15. *Halichondria galea*. A (TLT495), B & C (TLT743, C shows underside that was attached to
1417 floating dock). D: tangential ectosomal skeleton (TLT496), E: cross-section showing choanosomal
1418 skeleton at sponge surface (TLT495), F: oxeas (TLT495).

1419

1420 **Morphology**
1421 Thinly to thickly encrusting with uneven thickness, leading to a lumpy, undulating, or rugose
1422 appearance. Oscula, when apparent, nearly flush with surface. Surface is a mix of yellow and
1423 green regions, or yellow, green, and brownish regions, with the interior yellow. See Figure 15 for
1424 examples.

1425

1426 **Skeleton**
1427 Tangential ectosomal skeleton a sieve-like mat of spicules surrounding oval-shaped open areas
1428 with fairly consistent size. Choanosomal skeleton with meandering multispicular tracts and
1429 many spicules in confusion. No apparent spongin.

1430

Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering to sharp points at both ends. Like *H. panicea*, spicules have a
very large range in lengths. Though spicules in the ectosome are 10–20% shorter than the
choanosome, the full range of spicule lengths is seen in both compartments, with size

distributions continuous but bimodal within each compartment. Dimensions for three samples
where ectosome and choanosome were measured separately are given in table 6. When
spicules from all samples are combined, spicules measure 105–265–483 x 2–5–10 µm, n=308.
Average dimensions per sample were 249–270 µm long and 5–6 µm wide.
1440
1441 Table 6. Spicule lengths of *Halichondria galea*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT495	114–256–440 n=50	137–284–445 n=50
TLT496	126–269–472 n=40	125–300–483 n=40
TLT743	105–225–401 n=65	106–273–412 n=63

1442

Distribution and habitat
Known from China, Korea, and California. In the Eastern Pacific, this species is known from
Orange and San Diego Counties in Southern California. We found it growing on a floating dock in
Dana Point Harbor and in natural tidepools at Point La Jolla; it was also previously reported from
Mission Bay, San Diego (Park et al., 2007). In the Western Pacific it has been collected from the
Gulei Peninsula in China (Wang et al., 2016) and from rocky intertidal areas around the Korean
Peninsula and islands of the Korean Strait (Park et al., 2007).
1450

Remarks
The first mitochondrial genome published for a *Halichondria* species was for a "*Halichondria*
sp." sponge sampled from Fujian Province, China (Wang et al., 2016). Sequencing of
mitochondrial genes later documented this same genotype from the intertidal zone of mainland
South Korea and nearby islands, and from Mission Bay, San Diego, California (Park et al., 2007).
This sponge has remained undescribed, likely due to it being morphologically indistinguishable
from *H. panicea*.
Our Southern California samples have mitochondrial genomes that are nearly identical
with the published sequence from China (99.95% identical), confirming that we have recollected
this same species from California. Consistent with previously published data, it is only distantly
related to *H. panicea*, despite their morphological similarity. The clade we designate here as the
Okadai species group contains *H. okadai* and *H. galea*, which are sympatric in the Korean Strait,
as well as the Caribbean species *H. magniconulosa* and *H. melanodocia,* and samples we
collected from Pacific Panama that are likely from an undescribed species (Figure 3).
It is probable that *H. galea* has experienced a recent human-mediated range expansion,
though the evidence is less compelling than for the other species reported here. It has only
been found in the North Pacific thus far, and this could indicate a natural range connected
through the Bering Strait. It has been found in artificial habitats like floating docks, but most
samples reported to date have been collected from rocky intertidal areas that host fewer
introduced species. However, finding nearly identical mitochondrial genomes in China and
California makes it less likely that those populations are found at extreme ends of a wide
distribution, and more likely that they share a recent common ancestor. When combined with
the observation that many other *Halichondria* species have been moved around by human
activity, we feel this species should be treated as a possible introduction across some part of its
range (though which part remains to be determined).

1476

1477 *Halichondria bowerbanki* Burton 1930

1478 Figure 16

1479

Material examined

190615, Praia to Lumiar, Viana do Castelo, Portugal, (41.73540, -8.87578), 7.1 m, 2019-06-15;
190616, Canto Marinho, Viana do Castelo, Portugal, (41.72910, -8.88071), 12.5 m, 2019-06-16;
NIB0309, The Dorn, Northern Ireland, (54.43459, -5.54536), intertial (boulders), 2021-04-28;
NIB0612, Carnalea , Northern Ireland, (54.66844, -5.70801), intertidal (rocky shore), 2021-10-
27; NIB0614, Mount Stewart, Northern Ireland, (54.54482, -5.61017), intertidal (low shore
boulders), 2021-04-29; TLT1052, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1072, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1087b, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1357, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT1358, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT1418, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT234, Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-
11 m, 2019-07-29; TLT337, Saddleback Ridge, Santa Cruz Island, California, (34.03817, -
119.52470), 5-12 m, 2019-10-18; TLT342, Coal Oil Point, California, (34.40450, -119.87890), 3-8
m, 2019-08-30; TLT415, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2019-10-25;
TLT490, Coal Oil Point, California, (34.40450, -119.87890), 5 m, 2020-07-17; TLT738, Marina del
Rey, California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT980, Santa Barbara
Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT981, Santa Barbara
Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01.

1500

Diagnosis

Yellow, green, or blue-green Halichondria with a tangential ectosomal skeleton developed as a
web of spicule tracts and individual spicules, with open spaces generally triangular or polygonal
rather than oval. Choanosomal skeleton confused, with a slight tendency to form multispicular
tracts. Spicules oxeas, 180–550 μm long, with averages per sample of 300–430 μm long and 7–
12 μm wide; usually the same length in the ectosome and the choanosome. Diagnosis relative
to similar species requires DNA characters, as follows:

1508

1509 COX1. 301:C, 484:A . N=12.

1510

1511 28S. 8:G, 848:T. 853:G, 865:C, 1060:C, 1937:C, 2032:A, 2327:A. N=13–20 depending on position.

1512

1513 ND1. 264:G, 684:A, 691:C, 739:T. N=13–14 depending on position.

1514

1515 **Morphology**

1516 Previously described as variable in shape, from encrusting to cushion-shaped, with oscular
1517 chimneys and/or thread-like tendrils, and sometimes highly ramose with anastomosing
1518 branches (Ackers et al., 2007; Burton, 1930; Evcen et al., 2023). Samples examined here, with
1519 identities confirmed by DNA sequencing, do not include any forms with branches, threads, or

oscular chimneys, though the previously sequenced sample from Ireland (BELUM:Mc4003) had these features.

Sponges from sheltered marinas were thinly encrusting, usually green or blue-green but one sample was yellow. Samples from subtidal rocky reefs were usually large cushions, pale yellow, but one sample was encrusting and green. Oscula are not always apparent, but when they are, they are nearly flush with the sponge surface or with raised rims, and often accompanied by a translucent channel running up one side (Figure 16B). White or beige in ethanol.

Figure 16. *Halichondria bowerbanki*. A & B (TLT490), C (TLT337), D (TLT738), E (TLT1052), F (TLT234). G: spicules (TLT234). Tangential ectosomal skeletons: H (TLT337), I (TLT234). J: cross-section showing choanosomal skeleton at sponge surface (TLT342).

Skeleton

Tangential ectosomal skeleton is a web of spicule tracts and individual spicules, with open spaces generally triangular or polygonal rather than oval. The choanosomal skeleton is confused, with many free spicules, but meandering multispicular tracts are usually evident. No apparent spongin.

Spicules

Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have the typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering gently to sharp points at both ends. Spicules have a large
range in lengths in both the ectosome and the choanosome, with no average difference
between compartments (though individual samples varied from 5% shorter to 7% longer in the
choanosome). Dimensions for 10 samples where ectosome and choanosome were measured
separately are given in table 7. When spicules from all 16 samples with data are combined,
spicules measure 185–379–547 x 2–8–17 µm, n=1312 for length, n=1092 for width. Average
values varied considerably across samples, 307–423 µm long, and 7–12 µm wide.
1550
1551 Table 7. Spicule lengths of *Halichondria bowerbanki*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
TLT1052	273–356–454 n=45	203–359–410 n=40
TLT342	350–433–490 n=40	291–411–528 n=41
TLT490	340–412–477 n=40	265–430–506 n=40
TLT980	231–333–384 n=40	249–323–371 n=46
TLT981	251–322–389 n=40	213–323–379 n=40
TLT234	228–424–539 n=97	196–422–547 n=96
Atlantic Ocean		
NIB0512	185–307–365 n=40	206–306–399 n=40
NIB0614	292–389–467 n=45	320–416–491 n=45
NIB0309	263–362–488 n=96	210–370–472 n=96
190615	334–408–469 n=40	249–391–507 n=40

1552

Distribution and habitat
This species was previously described as occurring throughout Europe, both coasts of North
America, and the Korean Peninsula (de Voogd *et al.* 2024) However, many of these samples are
likely to be members of the cryptic species newly described here, so the confirmed range of this
species is limited to samples with genetic data. European samples with DNA confirmation are
from Ireland, the Netherlands, Portugal, and the Black Sea. Confirmed Pacific samples were
found in a small number of Southern California locations. Samples previously sequenced from
Argentina expand the known range of this species into the Southern Hemisphere (Figure S1).
This distribution is likely a result of accidental human transport, and the species seems likely to
be confirmed from additional locations in the future.
Previous publications have stated that this species is less tolerant of tidal exposure than
H. panicea, but more tolerant of silted and brackish conditions (Ackers et al., 2007; Hartman,
1958; Vethaak et al., 1982)*. DNA-confirmed samples are consistent with this description, as the
species was found in sheltered marinas and subtidal reefs, but intertidal samples were verified
only in Ireland. Samples were collected on floating docks in several marinas, but *H. urca, H.
zabra,* and *H. hygeia* were much more common and widespread on floating docks than *H.
bowerbanki.* Subtidal samples were collected in natural rocky reef habitat, up to 12 m deep,
usually growing on holdfasts of *Macrocystis pyrifera*. Though not quantitative, our observations

of this species in the California kelp forest demonstrate substantial differences in abundance
through time. The species was among the most abundant sponges at two Southern California
locations near Santa Barbara (Coal Oil Point and Arroyo Quemado Reef) in 2019 and 2020, but
could not be relocated in those areas in 2023 or 2024.

1575

Remarks
Though *H. panicea* and *H. bowerbanki* cannot always be distinguished without genetic data,
they do have average differences in morphological characters and habitat. However, *H.*
bowerbanki is much harder to distinguish from other *Halichondria* such as *H. zabra, H. urca,* and
H. pinaza, and many previous reports of *H. bowerbanki* are likely to be conflated with these
species. In California, for example, ramose *Halichondria* from harbors and marinas have long
been attributed to *H. bowerbanki*, but we found *H. bowerbanki* to be uncommon in California
marinas, and found it to be thinly encrusting in those habitats when present.
It is likely that *H. bowerbanki* is introduced in some parts of its range. The presence of
this species in the North Atlantic and North Pacific could potentially be explained by a natural
range that includes the Arctic, though this seems less likely than for *H. panicea* because 1)
genetic variation does not seem to be associated with geographic distance, 2) the species is
found in human-modified habitats where introduced species are common, and 3) samples
confirmed thus far suggest a less cold-tolerant range than for *H. panicea* (in the Pacific, where
our sampling is most extensive, it has been found only in Southern California so far). Moreover,
several *H. bowerbanki* were collected from Punta Verde, a beach in San Antonio Bay, Argentina,
by Marianela Gastaldi (*pers. comm.*). Sequencing of the cox1 locus confirms these are *H.*
bowerbanki (MZ487239, MZ487236, MZ487221). It is highly unlikely that this population was
founded by natural means.

1595

1596 *Halichondria urca* **sp. nov.**
1597 Figures 17 & 18

1598

Material examined
Holotype: TLT1087, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock,
2021-06-01. Paratypes: 15492, Cabrillo Beach, California, (33.70800, -118.28500), intertidal,
2019-08-20; 15925, Port of Long Beach, California, (33.72865, -118.23638), 10 m, 2019-08-20;
MSH23x2, Mount Sinai Harbor, New York, (40.96345, -73.03646), 1 m, 2023-08-16; NIB0610,
Carrickfergus Marina , Northern Ireland, (54.71028, -5.81237), floating dock, 2021-10-24;
TLT1002, Farnsworth Bank, California, (33.34380, -118.51650), 21-27 m, 2021-07-10; TLT1050,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1051,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1053,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1058,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1060,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1062,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1063,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1065,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1066,
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1068,

Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1069, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1070, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1071, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1073, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1075, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1078, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1081, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1083, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1085, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT534, Point La Jolla, California, (32.85227, -117.27239), intertidal, 2020-02-08; TLT590, Mission Bay: Quiviera Basin, California, (32.76421, -117.23829), floating dock, 2020-02-08; TLT591, Mission Bay: Quiviera Basin, California, (32.76421, -117.23829), floating dock, 2020-02-08; TLT593, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16; TLT594, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16; TLT661, Stearn's, California, (34.41023, -119.68563), 3-5 m, 2020-08-25; TLT715, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2020-12-14; TLT720, Mission Bay: South Shores Boat Launch, California, (32.76406, -117.21750), floating dock, 2021-01-10; TLT733, Marina del Rey, California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT740, Marina del Rey, California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT747, San Diego Bay: Shelter Island Marinas, California, (32.71094, -117.23423), floating dock, 2021-01-10; TLT764, Newport Bay: Balboa Peninsula, California, (33.60375, -117.90052), intertidal (on piling), 2021-01-09; TLT766, Newport Bay: Balboa Peninsula, California, (33.60375, -117.90052), floating dock, 2021-01-09; TLT926, Mission Bay: Quiviera Basin, California, (32.76421, -117.23829), floating dock, 2021-05-15; TLT951, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-05-24; TLT951B, Nudi Wall, California, (32.69872, -117.27580), 18-28 m, 2021-05-16; TLT960, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24.

Etymology
Named for the urca, one of the ships of the Spanish armada in the age of sail.

Diagnosis
Yellow *Halichondria* with a tangential ectosomal skeleton developed as a web of spicule tracts and individual spicules, or as a sieve-like mat of spicules surrounding oval-shaped open spaces. Spicules oxeas, 100–440 µm long, with averages per sample of 230–360 µm long and 5–10 µm wide; usually very slightly longer in the choanosome than the ectosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. 72:T, 93:A, 128:G, 139:C, 247:C, 310:C, 313:T, 364:T. 397:G, 412:T, 475:T, 538:G, 559:G, 634:T, 709:C, 823:G, 880:C, 979:C, 1040:T, 1041:G, 1102:T, 1120:T, 1150:A, 1321:T, 1432:G. N=9–21 depending on position.

28S. 693:A, 704:G, 829:C, 846:A, 884:T, 1359:G, 1373:C, 1656:G, 2331:G, 2102:C, 2328:T. N=9–40 depending on position.

1659

1660   ND1. 10:G, 121:C, 204:C, 228:C, 364:T, 448:G, 543:A, 658:T, 838:T. N=9–15 depending on
1661   position.

1662

1663
1664   Figure 17. *Halichondria urca*. A (TLT 661), B (TLT740), C (TLT747), D (TLT1063), E (TLT534), F (TLT1069,
1665   sponge torn to show brooded larvae inside). A (subtidal) & E (intertidal) are in situ, others are pictured
1666   after removal from floating docks, before preservation.

1667

Morphology
Samples discovered to date were bright yellow to yellowish brown, and did not contain any
traces of green. Shape is variable from thin encrustations to semi-globular cushions. Oscula can
be flush with surface, with raised rims, on volcano-shaped protrusions, or atop oscular
chimneys (Figure 17). Sometimes has tendril-like projections. Translucent areas usually present,
including channels leading up the sides of oscula, as often seen in *H. bowerbanki*. When
removed from the substrate or torn, brooded bright yellow larvae may be present internally and
where the sponge contacted the substrate (Figure 17F). In Santa Barbara Harbor, Southern
California, larvae were seen in one of two sponges collected in February, a single sponge
collected in May, and 5 of 20 sponges collected in June.

1678

Skeleton
The tangential ectosomal skeleton is usually developed as a web of long meandering spicule
tracts and single spicules, with open spaces between them irregular in size and shape. However,
some samples have more regular open spaces that are triangular and polygonal, and two
samples had a mat of spicules that surrounded oval-shaped openings (Figure 18C).
Choanosomal skeleton typically halichondroiid, with meandering spicules tracts and spicules in
confusion, without apparent spongin.

1686

Spicules

Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have a typical halichondroid shape: evenly curved or with a slight central bend, thickest
in the center and tapering to sharp points at both ends. Spicules have a large range in lengths in
both the ectosome and the choanosome, with a very slight tendency to be longer in the
choanosome (average of 3% longer across samples). Table 8 shows dimensions for 6 samples
where ectosome and choanosome were measured separately. When all 19 samples with
measurements are combined, spicules are 107–308–435 x 1–7–14 µm, n=1047 for length,
n=863 for width. Average values varied across samples, with means of 238–359 µm long, 5–10
µm wide.

1697

1698 Table 8. Spicule lengths of *Halichondria urca*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
TLT926	264–315–368 n=50	273–327–362 n=50
TLT951	241–296–335 n=40	242–320–364 n=40
TLT1051	277–308–361 n=40	241–310–359 n=42
TLT661	183–254–315 n=50	181–239–296 n=50
TLT715	211–315–365 n=42	288–332–391 n=42
Atlantic Ocean		
NIB0610	259–346–416 n=40	316–372–425 n=40

1699

Distribution and habitat
Known from the intertidal to at least 28 m depth, but this species is much more common on
floating docks in sheltered marinas than in other habitats investigated. Exceptions include
encrusting samples found exposed at low tide at Point La Jolla and on a rocky reef in the Point
Loma kelp forests, both near San Diego, California. Other samples found outside of marinas
were on man-made debris: entangled trash 3 m below the Santa Barbara Wharf and an
entangled rope at 21 m depth in the Farnsworth Offshore State Marine Conservation Area at
Catalina Island, California.
Pacific samples are all from Southern California. Atlantic samples are from New York,
Virginia, and Ireland. It was the most abundant species of *Halichondria* found in marinas in
Southern California, where 48% of *Halichondria* (34 of 71 samples) were this species (sampling
strategies were aimed at maximizing diversity rather than estimating relative abundance, so this
number is likely an underestimate). *Halichondria urca* was not found at higher latitudes in the
Northeast Pacific: only *H. zabra*, *H. pinaza*, and *H. hygeia* were collected in marinas from Central
California to British Columbia, suggesting that *H. urca* may be less cold-tolerant than those
species (though sampling in these areas was more limited than in Southern California).

1716

Remarks
This species is often morphologically indistinguishable from other members of the *Bowerbanki*
group, but there were some features that made it more likely a sample would prove to be *H.*
urca when genotyped. These include large size (semiglobular *Halichondria* ~10 cm across, from

floating docks, have consistently proved to be this species) and color (*H. urca* has not yet been
found with any traces of green color).
It is possible that *H. urca* is introduced in some parts of its range. Its strong association
with artificial substrate is consistent with an introduced species, as is finding samples in both
the North Atlantic and North Pacific. The genetic differences seen between the Atlantic and
Pacific are intriguing, however, and it is possible that additional sampling will reveal some of all
Atlantic samples to be from an additional undescribed species.
1728

1729
1730 Figure 18. *Halichondria urca* skeleton and spicules. Tangential ectosomal skeletons: A (TLT926), B
1731 (TLT661), C (TLT534), E (TLT591), F (TLT594). D: spicules (TLT733). G: cross-section showing choanosomal
1732 skeleton at sponge surface (TLT594).
1733

1734 ***Halichondria zabra* sp. nov.**
1735 Figures 19 & 20

1736

Material examined
Holotype: TLT1064, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock,
2021-06-01. Paratypes: TLT1077, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1082, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1086, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1317, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT612, Jack London Square Marina, California, (37.79371, -
122.27757), floating dock, 2020-11-15; TLT623, Jack London Square Marina, California,
(37.79371, -122.27757), floating dock, 2020-11-15; TLT624, Jack London Square Marina,
California, (37.79371, -122.27757), floating dock, 2020-11-15; TLT712, Ventura Harbor,
California, (34.24801, -119.26550), floating dock, 2020-12-14; TLT722, Ventura Harbor,
California, (34.24801, -119.26550), floating dock, 2020-12-14; TLT737, Marina del Rey,
California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT751, San Diego Bay: Shelter

Island Marinas, California, (32.71094, -117.23423), floating dock, 2021-01-10; TLT756, Mission Bay: South Shores Boat Launch, California, (32.76406, -117.21750), floating dock, 2021-01-10; TLT757, Rainbow Harbor, California, (33.76184, -118.19111), floating dock, 2021-01-09; TLT759, Rainbow Harbor, California, (33.76184, -118.19111), floating dock, 2021-01-09; TLT760, Rainbow Harbor, California, (33.76184, -118.19111), floating dock, 2021-01-09; TLT761, Rainbow Harbor, California, (33.76184, -118.19111), floating dock, 2021-01-09; TLT780, San Diego Bay: Shelter Island Marinas, California, (32.71094, -117.23423), floating dock, 2021-01-10; TLT815, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24; TLT874, Tomales Bay Resort & Marina, California, (38.10792, -122.86237), floating dock, 2021-02-08; TLT967, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24; TLT978, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24; TLT979, Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24.

Figure 19. *Halichondria zabra*. A (TLT1317), B (TLT1077), C (TLT978), D (TLT759), E (TLT624), F (TLT761), G (TLT1086, sponge torn to show brooded larvae inside). A is in situ, others are pictured after removal from floating docks, before preservation.

Etymology
Named for the zabra, one of the ships of the Spanish armada in the age of sail.

Diagnosis
Yellow or green *Halichondria* with a tangential ectosomal skeleton developed as a web of spicule tracts and individual spicules, with open spaces generally triangular or polygonal rather than oval. Choanosomal skeleton with spicule tracts and spicules in confusion. Spicules oxeas, 180–570 μm long, with averages per sample of 290–410 μm long and 6–12 μm wide; little-to-no difference in length in the ectosome and the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

1779   COX1. No unique derived states.

1780

1781   28S. 4–5:GG. State 4:G is unique compared to all close relatives, but homoplastic with members

1782   of the *Okadai* group. The full sequence 4–5:GG is not shared with any other sequenced

1783   *Halichondria*. 831–832:GG. State 831:G is unique compared to all close relatives, but

1784   homoplastic with *H. dokdoensis.* The full sequence 831–832:GG is not shared with any other

1785   sequenced *Halichondria*. N=6.

1786

1787   ND1. 238:A. N=14.

1788

Morphology

Samples discovered to date were yellow, green, or a mix of the two; some yellow samples were

very pale and nearly white (Figure 19F). Except in thinly encrusting samples, green pigment was

more pronounced on the external surfaces of the sponge. Found as thin or thick encrustations,

all with oscula elevated to various degrees: as volcano-shape protrusions, or atop oscular

chimneys. Tendril-like projections may also be present, and these and oscular branches are

sometimes anastomosing (Figure 19C). Translucent channels sometimes present, including

those leading up one side of oscular protrusions, as often seen in other members of the

Bowerbanki group. In Santa Barbara Harbor, Southern California, brooded bright yellow larvae

were seen in one of three sponges collected in June (Figure 19G).

1799

1800
1801   Figure 20. *Halichondria zabra* skeleton and spicules. Tangential ectosomal skeletons: A (TLT751), B

1802   (TLT624), E (TLT760). C: cross-section showing choanosomal skeleton at sponge surface (TLT623), D:

1803   spicules (TLT712).

1804

1805   **Skeleton**

1806   Tangential ectosomal skeleton with a web of spicule tracts and single spicules, with open spaces

1807   fairly regular in size and tending towards triangular or polygonal shapes. Choanosomal skeleton

1808 typically halichondroid, with vague spicules tracts and spicules in confusion, without apparent
1809 spongin.
1810

Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering to sharp points at both ends. Spicules have a large range in
lengths in both the ectosome and the choanosome, with little average difference in length
between compartments. Table 9 shows dimensions for 5 samples where ectosome and
choanosome were measured separately. When all 17 samples with measurements are
combined, spicules measure 186–347–566 x 3–9–15 µm, n=1080 for length, n=912 for width.
Average values varied across samples, with means of 294 to 407 µm long and 6 to 12 µm wide.
1819

1820 Table 9. Spicule lengths of *Halichondria zabra*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT712	231–365–458 n=40	328–405–457 n=40
TLT780	234–309–364 n=40	188–294–344 n=41
TLT978	305–397–474 n=40	312–416–500 n=40
TLT1077	225–319–380 n=40	203–327–413 n=40
TLT1064	244–361–411 n=40	304–352–433 n=40

1821

Distribution and habitat
This species is known only from floating docks in California marinas, where it was found from
Tomales Bay (the most Northen site investigated) to San Diego Bay (the most Southern site
investigated). It was the second most common *Halichondria* on docks in Southern California,
where 19 of 71 *Halichondria* (27%) were this species (sampling strategies were aimed at
maximizing diversity rather than estimating relative abundance, so this number is very
approximate). Only *H. zabra* and *H. pinaza* have thus far been found on floating docks in
Northern California, where they were at similar abundance (with much less sampling than
Southern California).
1831

Remarks
Though this species has only been found in California thus far, we predict it will be found in
additional regions as more samples are sequenced. Its strong association with floating dock
habitat, where many introduced species are found, make it likely that it has been
unintentionally introduced. Previously published cox1 sequences from New Hampshire, New
York, Virginia, and Germany are consistent with a possible Atlantic distribution (see
supplementary data), but could also be from *H. pinaza* (these two species cannot be
differentiated at the most commonly sequenced region of this gene).
1840

1841 ***Halichondria pinaza* sp. nov.**
1842 Figure 21
1843

1844 **Material examined**

Holotype: TLT869, Sausalito Yacht Harbor, California, (37.85930, -122.48044), floating dock, 2021-02-08. Paratypes: NIB0603, Ballycastle Marina, Northern Ireland, (55.20687, -6.23962), floating dock, 2021-10-16; NIB0604, Ballycastle Marina, Northern Ireland, (55.20687, -6.23962), floating dock, 2021-10-16; TLT1056, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1080, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT873, Tomales Bay Resort & Marina, California, (38.10792, -122.86237), floating dock, 2021-02-08; TLT892, Home Bay, Drake's Estero, California, (38.06934, -122.91711), intertidal, 2021-02-08; WMS21x22, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal (rocks), 2021-05-26; WMS23x1, West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal (rocks), 2023-06-07.

Figure 21. *Halichondria pinaza*. A (TLT892), B (TLT1080), C (TLT869), D & E (TLT1056, E shows underside where sponge attached to dock, with brooded larvae visible). Tangential ectosomal skeletons: F (NIB0603), G (NIB0604). H: cross-section showing choanosomal skeleton at sponge surface (TLT1080), I: spicules (TLT869).

Etymology
Named for the pinaza, one of the ships of the Spanish armada in the age of sail.

Diagnosis

Yellow or green *Halichondria* with a tangential ectosomal skeleton developed as a web of multispicular tracts and individual spicules, with open spaces tending towards triangular or polygonal shapes rather than ovals. Choanosomal skeleton with spicule tracts and spicules in confusion. Spicules oxeas, ranging from 90–370 μm in length, with averages per sample of 264–308 μm in length and 6–10 μm in width; little-to-no difference in length in the ectosome and the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 4:T, 628:A, 669:T, 786:A, 803:C, 998:T. N=7–9 depending on position.

ND1. No unique derived states.

Morphology
All samples found to date were encrustations with protruding oscular chimneys (Figure 21). Occurs in yellow, green, or both. Tendril-like projections not seen. Translucent channels sometimes present, including those leading up one side of oscular protrusions, as often seen in other members of the *Bowerbanki* group. In Santa Barbara Harbor, California, brooded bright yellow larvae were seen in two sponges collected in June (Figure 21E).

Skeleton
Tangential ectosomal skeleton with a web of spicule tracts and single spicules, with open spaces tending towards triangular or polygonal shapes (Figure 21F and 21G). Choanosomal skeleton typically halichondroid, with vague spicule tracts and spicules in confusion, without apparent spongin.

Spicules
Nearly exclusively oxeas, though one sample (TLT892) had some aberrant spicules including styles, centrotylote oxeas, and oxeas with x-shaped cross pieces at one end. Oxeas have the typical halichondroid shape: evenly curved or with a slight central bend, thickest in the center and tapering to sharp points at both ends. Spicules have a large range in lengths in both the ectosome and the choanosome, with a very slight tendency to be longer in the choanosome (average of 3% longer across samples). Table 10 shows dimensions for 3 samples where ectosome and choanosome were measured separately. When all 6 samples with measurements are combined, spicules are 92–288–365 x 1–7–12 μm, n=432. Average values varied across samples, with means of 264 to 308 μm long, 6 to 10 μm wide.

Distribution and habitat
This species is thus-far known from California, New York, and Ireland. All samples were found in sheltered bays and marinas. All were on floating docks except one sample on the underside of an intertidal boulder. It may be less tolerant of warm water than congeners such as *H. bowerbanki* and *H. zabra*, as California samples were found from Drake's Estero in Northern California to Santa Barbara in Southern California, but not in sites farther South. Of all the *Halichondria* collected on floating docks in Southern California, only 2 of 71 (3%) were this

1910 species; in contrast, *H. zabra* and *H. pinaza* had similar abundance in Northern California
1911 (though sampling effort was more limited there).
1912
1913 Table 10. Spicule lengths of *Halichondria pinaza*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
TLT1056	158–272–311 n=55	236–281–330 n=57
Atlantic Ocean		
NIB0603	268–296–343 n=40	265–310–351 n=40
NIB0604	261–305–357 n=41	179–302–346 n=40

1914
1915 **Remarks**
1916 This species is likely introduced in some portion of its range. First, it was found primarily on
1917 floating docks, which are known to harbor many introduced species (Bulleri and Chapman,
1918 2010; Glasby et al., 2007; Ruiz et al., 2000; Simkanin et al., 2012). Second, samples with very
1919 similar DNA sequences were found in California, New York, and Ireland, despite this species
1920 likely having limited larval dispersal (Maldonado, 2006; Xue et al., 2009). Sequences from the
1921 cox1 locus from samples collected by others from New Hampshire, New York, Virginia, and
1922 Germany may indicate this species is also present in those areas, but these could also be *H.*
1923 *zabra* (these two species cannot be differentiated at the most commonly sequenced region of
1924 this gene).
1925
1926 *Halichondria loma* Turner & Lonhart 2023
1927
1928 **Material examined**
1929 Holotype: TLT1225, North Monastery Beach, California, (36.52647, -121.92730), 12-29 m, 2021-
1930 09-22. Paratypes: TLT1012, Acropolis Street (Pt. Pinos), California, (36.64183, -121.93060), 9-18
1931 m, 2021-08-09; TLT1024, Butterfly House, California, (36.53908, -121.93520), 9-20 m, 2021-08-
1932 10; TLT1130, Fire Rock Pescadero Point, California, (36.55898, -121.95110), 10-22 m, 2021-08-
1933 10; TLT514, Goalpost, San Diego, California, (32.69438, -117.26860), 12-15 m, 2020-02-08;
1934 TLT936, Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15; TLT976,
1935 Wreck of the Ruby E, California, (32.76680, -117.27620), 18-25 m, 2021-05-16; TLT989, Inner
1936 Pinnacle, California, (36.55910, -121.96630), 10-18 m, 2021-08-10.
1937
1938 **Diagnosis**
1939 Yellow *Halichondria* with a tangential ectosomal skeleton developed as a web of mulitspicular
1940 tracts and individual spicules, with open spaces tending towards triangular or polygonal shapes,
1941 or as a lace-like mat of spicules surrounding oval-shaped open areas. Choanosomal skeleton
1942 with spicule tracts and spicules in confusion. Spicules oxeas, 290–650 µm long, with averages
1943 per sample of 460–550 µm long and 5–16 µm wide. This species is diagnosed relative to *H.*
1944 *sitiens* by a lack of papillae, and from all other *Halichondria* examined here by having longer
1945 spicules However, this difference is slight (the longest *H. bowerbanki* sample averaged only 39
1946 µm shorter), so diagnosis is facilitated by DNA characters, as follows:

COX1. No unique derived states.

28S. 5:A, 13:G, 46:G, 621:G, 674:T, 894:C, 2091:T, 2198:G. N=6–8 depending on position.

ND1. 199:C, 205:G, 209:G, 220:A, 231:G, 248:T, 250:T, 700:G, 880:A. N=7.

Remarks

This species was recently described by Turner and Lonhart (2023), and no additional samples have been discovered or examined since. Our contribution to the understanding of this species comes from DNA data: we used Illumina sequencing to increase the genetic data available for 7 of the 8 previously known samples, and used that data to locate DNA characters that aid in diagnosis.

This species is currently the closest known relative of the *Bowerbanki* group (Figure 1). In contrast to the species in that group, *H. loma* has not been found in sheltered bays or on human structures, and there is no indication that it may be introduced in California.

Genus *Hymeniacidon* Bowerbank, 1858

***Hymeniacidon perlevis* (Montagu, 1814)**
Figure 22

Synonyms
Spongia perlevis Montagu 1814
Hymeniacidon sinapium de Laubenfels 1930
The World Porifera Database lists 27 additional synonyms (de Voogd, et al., 2024)

Material examined

CASIZ 180246, Newport Bay (H. sinapium type location), California, depth not known, 2006-09-14; NIB0611, Carnalea, Bangor, Northern Ireland, (54.66844, -5.70801), intertidal (rocky shore crevices), 2021-10-27; NIB0615, Mount Stewart, Strangford Lough, Northern Ireland, (54.54482, -5.61017), intertidal (boulders), 2021-04-29; NIB0616, Portaferry, Strangford Lough, Northern Ireland, (54.38226, -5.55393), intertidal (boulders), 2021-10-10; TLT1054, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1057, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1059, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1067, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1074, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1079, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1265, Zebra Cove, California, (34.01000, -119.44000), 6-13 m, 2022-06-17; TLT129, Carpinteria Reef, California, (34.39163, -119.54169), 3-6 m, 2019-07-31; TLT247, Carpinteria Reef, California, (34.39163, -119.54169), 3-6 m, 2019-07-31; TLT595, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-07-17; TLT643, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16; TLT648, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16;

1991 TLT651, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-07-17; TLT658, Lechuza
1992 Point, California, (34.03437, -118.86146), intertidal, 2020-12-14; TLT730, Lechuza Point,
1993 California, (34.03437, -118.86146), intertidal, 2020-12-14; TLT736, Marina del Rey, California,
1994 (33.97228, -118.44653), floating dock, 2021-01-09; TLT762, Rainbow Harbor, California,
1995 (33.76184, -118.19111), floating dock, 2021-01-09; TLT79, Tajigus, California, (34.46279, -
1996 120.10185), 3-8 m, 2019-07-01; TLT823, Elwood Beach, California, (34.42395, -119.90604),
1997 intertidal, 2020-12-29.
1998

2000 Figure 22. *Hymeniacidon perlevis*. A (TLT1265), B (TLT550), C (TLT730), D (TLT1067). A-C are in situ, with A
2001 and B subtidal and C exposed at low tide. D is removed from a floating dock. E: tangential ectosomal
2002 skeleton (TLT129). F: cross-section showing choanosomal skeleton at sponge surface (TLT129). G: typical
2003 style (TLT1057).
2004

2005 **Diagnosis**
2006 Yellow, green, orange, or red-orange *Hymeniacidon,* thinly or thickly encrusting, frequently with
2007 oscular chimneys, tendrils, or other fistulae. Ectosomal skeleton developed as a disorganized
2008 network of tangential spicule bundles, with some paratangential and protruding from the
2009 sponge surface. Choanosomal skeleton with multispicular tracts and spicules in confusion.
2010 Spicules styles; in the Northeast Pacific, they range from 110–420 μm in length, with averages
2011 per sample of 170–290 μm in length and 4–6 μm in width. Diagnosed relative to all other
2012 *Hymeniacidon* in the Northeast Pacific, except *H. fusiformis*, by having a large range (>100 μm)
2013 in lengths within any one sample, with some spicules in excess of 300 μm. Diagnosis relative to
2014 *H. fusiformis* based on thinner, non-fusiform spicules. It is not well established if this species can
2015 be diagnosed morphologically in all other regions where it occurs, so DNA confirmation of clade
2016 membership is recommended.
2017

2018 **Morphology**

Yellow, green, orange, or red-orange encrusting sponges, frequently with erect projections such as oscular chimneys and digitate tendrils. Dark red and pinkish forms reported from Europe and southern Africa (Ackers et al., 2007; Samaai et al., 2022). In silted habitats, it is often partially buried, with only vertical projections protruding from sediment. Translucent surface channels sometimes present, but it generally has a more opaque surface than *Halichondria* in the *Panicea* and *Bowerbanki* groups. Sometimes containing brooded yellow larvae.

Skeleton
Ectosomal skeleton is more disorganized than *H. panicea*, but similar in that it is developed as a tangential mesh of spicule bundles (Figure 22E). Bundles of spicules also protrude from this mesh to create a microscopically hispid surface. Choanosomal skeleton with wispy spicule tracts and spicules in confusion (Figure 22F). No apparent spongin.

Spicules
Exclusively styles, usually with uniform thickness but a significant minority have faint swelling at or near the head (Figure 22G). When data from 18 samples from California are combined, spicules measure 116–242–415 x 2–5–9 µm, n=660 for length, n=645 for width. Mean lengths per sponge varied from 177 to 289 µm, and mean widths from 4 to 6 µm. Sponges from other regions are reported to be similar in length but sometimes thicker (Harbo et al., 2021).

Distribution and habitat
In the Northeastern Pacific this species is known from Ladysmith Harbor, British Columbia to Southern California. It is present in many other regions as well, including China, Korea, Europe, Brazil, Argentina, and South Africa (Samaai et al., 2022; Turner, 2020). Future DNA sequencing will likely confirm its presence in other regions such as Peru (Cóndor-Luján et al., 2023) and possibly New Zealand (Bergquist, 1970).
        This species has a broad environmental niche, thriving in the intertidal zone, the shallow subtidal, areas with high levels of siltation, and brackish bays and estuaries. It is abundant on human structures in marinas but can also be found in relatively undisturbed natural habitats, such as those found at the Channel Islands in Southern California.

Remarks
Turner (2020) compiled genetic and morphological data from many previous studies to argue that this species has a worldwide distribution. Here, we provide additional genetic data consistent with that conclusion. A complete mitochondrial sequence was previously published from Korea (Jun et al., 2015), and we find it to be nearly identical to mitochondrial genomes from California and Ireland.
        Eleven samples were collected in Santa Barbara Harbor from April to July 2021, and checked for brooded larvae. None were seen in the 10 samples collected between April 21st and June 1st, but they were present in the final sample collected July 1st.

Hymeniacidon actites (Ristau, 1978)
Figure 23

Figure 23. *Hymeniacidon actites*. A (TLT227), B (TLT1158), C (TLT454). Spicules: D (USNM 24526). E: tangential ectosomal skeleton (TLT227). F & G: cross-sections showing choanosomal skeleton (TLT227; ectosomal skeleton is visible at the top of G).

Synonyms

Leucophloeus actites Ristau 1978

Material examined

Holotype: USNM 24526, Horeshoe Cove, Bodega Head, California, intertidal, 1977-01-06. Other samples: RBC1, Execution Rock Cave, Bamfield, British Columbia, (48.83170, -125.17830), not recorded, 2010-08-10; TLT1158, CRABS, California, (36.55377, -121.93840), 10-17 m, 2021-09-21; TLT227, Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-11 m, 2019-07-29; TLT454, Middle Reef, Whaler's Cove, Point Lobos, California, (36.52172, -121.93894), 6-15 m, 2019-11-23; TLT847, Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06; TLT856, Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06.

Diagnosis

Encrusting yellow *Hymeniacidon* with partially translucent ectosome and no tassels or tendril-like projections. Spicules exclusively styles 150–200 µm long, with less than 75 µm difference between shortest and longest spicules in any one sample. Well-developed sieve-like tangential ectosomal skeleton reminiscent of *H. panicea*. Much more genetically similar to *H. panicea* than other species of *Hymeniacidon.*

Morphology

Encrusting sponges, bright yellow to pale yellow-tan alive, with partially translucent ectosome. Beige or white preserved. Oscula usually apparent but small and flush with sponge surface. General morphology indistinguishable from sympatric *Halichondria* such as *H. californiana*. See Figure 23 for examples.

Skeleton

Tangential ectosomal skeleton generally sieve-like, with a dense mat of spicules surrounding open areas that are mostly circular or oval shaped (Figure 23E). Choanosomal skeleton with many spicules in confusion, but wavy, meandering tracts can be found in some samples. No apparent spongin.

Spicules

Exclusively styles, generally straight or evenly curved. Some samples contain spicules with very slight swelling at spicule heads or with multiple bends that make spicules somewhat sinuous. When all samples are combined, spicules measure 139–179–237 x 2–5–8 µm, n=250. Mean values per sample 156 to 199 µm long, 4 to 6 µm wide. Spicules from holotype: 168–196–237 x 4–6–8 µm, n=43.

Distribution and habitat

Occurs in wave-exposed rocky intertidal habitats and shallow rocky reefs; not known from bays, harbors, or artificial structures. Known range previously limited to Northern and Central California (Lee et al., 2007), but is now extended North to British Columbia and South to Southern California. One sample was found on a *Loxorhynchus* decorator crab.

Remarks

This species can be differentiated from *H. kitchingi* and *H. pierrei* due to the absence of tendrils and tufts, and from *H. perlevis* by color, surface patterning, smaller spicules, and a narrower range in spicule lengths. It is harder to distinguish from *H. ungodon* de Laubenfels 1932. As described below, the spicules and choanosomal skeletons in the two holotypes are likely indistinguishable. Indeed, the British Columbian sample examined here (RBC 018-00195-002) was previously identified as *H. ungodon* by William Austin. We assign this sample, and the others we collected, to *H. actites* based on two apparent differences between the species: ectosomal skeleton and color (Ristau, 1978).

Regular sieve-like ectosomal patterning was reported in the original *H. actites* description and was seen in samples we examined (including the British Columbian sample). The *H. actites* holotype fragment examined by us was too small to investigate the skeleton, but the holotype was previously examined by Willard D. Hartman, and online images of his sections show this patterning as well (Yale Peabody Museum, sample IZ.074262). In contrast, the original description of *H. ungodon* states it has a rugose surface with a dense mat of ectosomal spicules in confusion (de Laubenfels, 1932). Absence of sieve-like ectosomal patterning in the *H. ungodon* holotype is not conclusive evidence of these species being distinct, as other samples we examined that show a sieve-like pattern in some portions of the ectosome will show a disordered mat in other places. *Hymeniacidon ungodon* is also described as having contrasting "mahogany-brown ectosome over yellowish-drab endosome", and this was called-out in the original description as being "characteristic and unique in the genus" (de Laubenfels, 1932). Color is known to be highly variable, so it is unclear if this trait is a legitimate reason to separate these species either. Genetic data from samples matching the original description of *H. ungodon*

are therefore needed to resolve whether these two species are truly different, but for now, we
assign these samples to *H. actites*.

Hymeniacidon ungodon de Laubenfels 1932

Material examined
Holotype: USNM 22061, Point Lobos, Carmel, California, intertidal, 1930-07-12.

Diagnosis
Encrusting sponges with mahogany-brown ectosome over yellowish-drab endosome and no
tassels or tendril-like projections. Ectosome densely packed with spicules in confusion. Spicules
exclusively styles 160–230 µm in length.

Morphology
We were unable to locate any new samples that we could definitively assign to this species, and
examined only a small fragment of the holotype. The original description states that they are
encrusting sponges 1 cm thick, with a coarsely rugose surface; mahogany-brown ectosome over
yellowish-drab endosome (de Laubenfels, 1932).

Skeleton
The original description states that the ectosome is "densely packed with spicules in confusion".
Only a small piece of ectosome could be located on the holotype fragment we examined, and it
was a mat of tangential spicules in confusion, matching the previous description. The
choanosomal skeleton was previously described as confused with vague ascending tracts, like
other species of *Hymeniacidon*.

Spicules
Exclusively styles, generally curved and of consistent thickness. 161–206–227 x 4–7–8 µm, n=50.

Distribution and habitat
The holotype is from the intertidal zone in Central California, and was the only sample
vouchered by de Laubenfels; the original description states that he has "seen this species
several times in collections made by students" but the locations are not stated (de Laubenfels,
1932). Samples identified by others as this species have ranged from British Columbia to
Southern California, from the intertidal to the subtidal (Lee et al., 2007). We examined many of
these samples; the only one from Southern California is likely *H. perlevis*, and the others were
either assigned to *H. actites* or were of uncertain affinity.

Remarks
When Ristau (1978) described *H. actites,* he stated that it differed from *H. ungodon* in
"coloration, growth form, and disposition of the ectosomal skeleton". We have been unable to
locate any fresh samples matching the coloration described for *H. ungodon,* and all samples we
examined had ectosomal skeletons matching *H. actites*. DNA was successfully amplified from
only one sample previously identified as *H. ungodon*, from British Columbia, and it matched

sequences from *H. actites*. Genetic data from samples matching the original description of *H.*
ungodon are needed to resolve whether these two species are truly different.
2181
2182 ***Hymeniacidon kichingi* (Burton, 1935)**
2183 Figure 24
2184

2185
2186 Figure 24. *Hymeniacidon kitchingi*. A (TLT597), B (TLT154), C (TLT412). D: cross-section showing
2187 choanosomal skeleton at sponge surface (TLT154). Spicules: E (TLT154), F (TLT491).
2188
Material examined
TLT154, Elwood Reef, California, (34.41775, -119.90150), 9-15 m, 2019-05-15; TLT378, Coal Oil
Point, California, (34.40450, -119.87890), 3-8 m, 2019-10-25; TLT412, Elwood Reef, California,
(34.41775, -119.90150), 9-15 m, 2019-10-23; TLT424, Elwood Reef, California, (34.41775, -
119.90150), 9-15 m, 2019-10-23; TLT491, Coal Oil Point, California, (34.40450, -119.87890), 7 m,
2020-07-17; TLT597, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-07-17;
TLT820, Platform Holly, California, (34.38993, -119.90656), 3-21 m, 2021-05-12; TLT941, Lazy
Days, San Diego, California, (32.69415, -117.27110), 12-25 m, 2021-05-15; TLT977, Wreck of the
Ruby E, California, (32.76680, -117.27620), 18-25 m, 2021-05-16.
2198
Diagnosis
Encrusting or cushion-shaped sponges, pale yellow, bright yellow, or beige shaded with pink or
purple. Surface bearing many thin tassel-like extensions: in Californian samples these have a
distinctive "frayed rope" appearance that differentiates this species from all others known from
the region except *H. pierrei*. Styles are the only spicule, 110–260 µm long and 1–8 µm wide.
Choanosomal skeleton is typically halichondroid, with meandering spicule tracts and spicules in
confusion; differentiated ectosomal skeleton lacking. Diagnosis relative to *H. pierrei* requires
DNA characters, as follows:
2207

COX1. 718:T, 742:A, 769:G785:C, 786:T, 792:C, 796:C, 820:G, 937:C, 949:A, 1011:C, 1030:C, 1099:C, 1108:A, 1186:C, 1369:A, 1372:T, 1405:A, 1423:A, 1444:G, 1447:G, 1455:G, 1480:G, 1489:A, 1492:C. N=5–8 depending on position.

28S. 190:T, 206:G, 286:G, 471:G, 525:T, 529:T, 536:A, 540:C, 552:C, 558:A, 579:A, 603:T, 637:A, 641:T, 649:C, 660:G, 682:T, 697:T, 720:T, 789:C, 793:T, 797:C, 820:G, 826:A, 852:C, 859:T. N=5–9.

ND1. Not assessed.

Morphology
Form varies from thin encrustations to semiglobular cushions. California samples were replete with tassels: digitate projections that appear frayed due to profusely branching tips. All but the smallest samples in the British Isles are also covered in thin tassels, but available images do not clearly show the "frayed ends" appearance (Ackers et al., 2007). California samples pale to bright yellow; British Isles samples greyish-beige, with a tinge of pink or purple. Translucent surface channels, pores, and small scattered oscula visible on some living samples.

Skeleton
Choanosomal skeleton typically halichondroid, with meandering multispicular tracts and spicules in confusion. No special ectosomal skeleton is apparent, though some spicules and spicule tracts are tangential in the surface of the sponge. Spicule tracts pierce the sponge surface at tassel tips. No apparent spongin.

Spicules
Thin styles, straight or slightly bent. Most are of consistent thickness until tapering to a point, but a significant minority have very slight swelling at the head, with at least one sample (TLT491) containing some tylostyles. Some are slightly thinner at the head, and therefore weakly fusiform. The original description notes that tips are telescope-like, in that they taper in a stepwise fashion. This feature was subtly present in Californian samples, but only in some spicules; a more recent publication found it to be present on only about 10% of spicules in samples from the British Isles as well (Ackers et al., 2007). When data from 7 California samples is pooled, spicules measure 114–216–260 x 1–4–8 μm, n=192, very similar to reports on European samples (Ackers et al., 2007). Mean values per sponge were 206 to 223 μm long, 4 to 6 μm wide.

Distribution and habitat
Previously known from just below the intertidal zone in the British Isles and Brittany. Samples examined here were found in 5 locations in Southern California, all subtidal, from 3 to 25 m depth. Two of these locations were artificial substrates (an oil platform and a scuttled ship), and three were natural reefs where the sponge was attached to rock.

Remarks
This species was not previously known outside of the British Isles and Brittany, so it was surprising to find samples matching its description in Southern California. California samples are

genetically identical to a 2085 bp sequence of the 28S ribosomal locus that was previously published from Ireland (Morrow et al., 2012; Thacker et al., 2013). This sequence includes the hypervariable "C2" barcoding region, making it highly likely that these samples are the same species. It also seems probable that the species has experienced a human-assisted range expansion, as a natural circumboreal distribution would likely result in some genetic divergence between these distant locations. This species can therefore be added to the growing list of Halichondriidae potentially introduced by human activity.

This species is not closely related to the type species *H. perlevis*, and instead forms a distinct clade with *H. pierrei*, *H. fusiformis*, an unidentified *Hymeniacidon* from the North Atlantic, and *Semisuberites cribrosa* (Figure 1). Published sequences of cox1 indicate additional samples in this clade, identified as belonging to a variety of genera, with a Hawaiian sample identified as *H. gracilis* being the only additional sample identified as a *Hymeniacidon* (Figure S1).

Hymeniacidon pierrei sp. nov.
Figure 25

Material examined
Holotype: TLT599, Coal Oil Point, California, (34.40450, -119.87890), 6.7 m, 2020-07-17.
Paratype: TLT531, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-06-30.

Etymology
Named for Christoph Pierre, marine naturalist and Director of Marine Operations at the University of California, Santa Barbara.

Diagnosis
Encrusting *Hymeniacidon* bearing many thin tassel-like extensions that differentiate this species from all others known from the region except *H. kitchingi*. Styles are the only spicule, 100–230 µm long and 1–7 µm wide. Choanosomal skeleton is typically halichondroid, with meandering spicule tracts and spicules in confusion; differentiated ectosomal skeleton lacking. Diagnosis relative to *H. kitchingi* requires DNA characters, as follows:

COX1. 486:C, 502:C, 598:C, 727:G, 792:A, 886:C, 1034:C. N=1.

28S. 170:T, 181:T, 182:T, 203:A, 262:T, 277:C, 278:A, 377:A, 463:G, 500:A, 512:C, 523:T, 530:T, 535:A, 551:T, 552:A, 559:T, 562:T, 603:G, 606:T, 608:C, 609:A, 611:G, 615:A, 623:A, 625:A, 642:C, 654:G, 672:T, 682:G, 704:C, 720:C, 769:A, 771:A, 778:A, 789:T, 792:A, 795:G, 796:A, 818:A, 819:C, 827:T, 841:T, 850:T, 884:G, 894:C. N=2.

ND1. Not assessed.

Morphology

2294 Thinly encrusting sponges replete with tassels: digitate projections that appear frayed due to
2295 profusely branching tips (Figure 25A & 25B). Translucent surface channels, pores, and small
2296 scattered oscula visible on living samples. Pale yellow alive, white when preserved.
2297

2298
2299 Figure 25. *Hymeniacidon pierrei*. A (TLT599), B (TLT531). C: cross-section showing choanosomal skeleton
2300 in tuft (TLT531). D: tangential spicules and tracts in ectosome (TLT599). Spicules: E (TLT531).
2301

Skeleton

2302
2303 Choanosomal skeleton typically halichondroid, with meandering multispicular tracts and
2304 spicules in confusion. Branching and somewhat plumose spicule tracts are well developed in
2305 tassels, where they pierce the sponge surface at tassel tips. No special ectosomal skeleton is
2306 apparent, though some spicules and spicule tracts are tangential in the ectosome. No apparent
2307 spongin.
2308

Spicules

2309
2310 Thin styles, straight or slightly bent. Most are of consistent thickness until tapering to a point,
2311 but a minority have very slight swelling at the head. Like *H. kitchingi*, the tips of some spicules
2312 taper in a subtly stepwise fashion. When data from both samples is pooled, spicules measure
2313 109–187–229 x 1–4–7 µm, n=121. Means per sponge were 183 and 199 µm in length and 4 and
2314 5 µm in width.
2315

Distribution and habitat

2316
2317 This species is only known from the shallow (3–8 m) subtidal at Coal Oil Point, Santa Barbara,
2318 California. It was found on the decaying holdfasts of dead *Macrocyctis pyrifera*.
2319

Remarks

2320
Hymeniacidon kitchingi was discovered in California in 2019 at Coal Oil Point and nearby Elwood
Reef. An additional tasseled *Hymeniacidon* was collected on a return trip to Coal Oil Point in

2020, but when it was sequenced, it was very genetically distinct from *H. kitchingi*. An
additional trip later in 2020 yielded both genotypes, collected side-by-side. With 13% sequence
divergence (Fst = 0.97) at the 28S locus and 10% sequence divergence (Fst = 0.32) at the cox1
locus, these sequences are clearly from a different species. Indeed, at cox1, *H. pierrei* are more
closely related to samples collected in Hawaii (Genbank accession KY565326, identified as *H.*
gracilis) and Australia (Genbank accession KJ620398, unidentified) than to *H. kitchingi.*
It is unclear whether *H. pierrei* is native to California or introduced. It was found on a
natural reef that is now part of a marine protected area, but this site was previously highly
disturbed, as it housed a pier and oil derricks as part of the Elwood Oil Field. This area is one of
two places that the (possibly introduced) *H. kitchingi* has been found, so it seems possible that
H. pierrei is introduced as well.
2334
2335 *Hymeniacidon fusiformis* **Turner & Lonhart 2023**
2336
2337 **Material examined**
2338 Holotype: CASIZ236657/IZC00048452, Inner Carmel Pinnacle, Carmel, California, (36.55852, -
2339 121.96820), 10–24 m, 2021-09-22. Additional sample: FH12x28, Bell Island (West Face),
2340 Washington, (48.59667, -122.98150), 15 m, 2012-05-18.
2341
Diagnosis
Thickly encrusting white or yellow sponges, sometimes with irregularly projecting fingers and
lobes, but not thin tentacle-like extensions or tassels. Styles are the only spicule, slightly
fusiform, 150–410 μm in length and 4–19 μm in width. Maximum style length and large range in
spicule lengths differentiate this species from sympatric *Hymeniacidon* except *H. perlevis*, while
the wide range in spicule widths is unique in this geographic region. Choanosomal skeleton is
typically halichondroid, with spicule tracts and spicules in confusion. The ctosomal skeleton is
developed as a sieve-like mesh surrounding triangular open spaces or as a dense unstructured
mat of tangential spicules.
2351
Spicules
Exclusively styles; most are very slightly fusiform, so that they are thicker in the center than
near the head of style. Length and width are both highly variable; distributions are continuous
but strongly bimodal, despite not showing any differences in size in the ecotosome vs. the
choanosome. Spicules in the holotype measure 186–281–409 x 4–10–18 μm (n=345), with
length modes 210 and 350 μm and width modes 9 and 14 μm. The new sample from
Washington is similar, 153–270–396 x 6–11–19 μm (n=80), with length modes 260 and 330 μm
and width modes 8 and 13 μm.
2360
2361 **Distribution and habitat**
2362 Known from two samples, the first from a natural subtidal pinnacle in Carmel Bay, Central
2363 California, 10–24 m depth, the second from a submerged rock wall, Bell Island, Washington, 15
2364 m depth.
2365
Remarks

This species was recently described by Turner & Lonhart (2023) from a single sample. Here we
report additional genetic data from this sample. We used Illumina sequencing to completely
sequence the nuclear ribosomal locus; coverage was insufficient to assemble the mitochondrial
genome, but we were able to extract 777 bp of the cox1 locus. We also report the discovery of
one additional sample. This new sample conforms well to the morphological description of the
holotype, and is only 0.4% divergent at the 28S locus.
The closest relative to this species in our molecular phylogenies is a *Hymeniacidon* sp.
we collected and sequenced from the Bay of Fundy, in the Canadian North Atlantic (sample
Quoddy115 in Figures 1 & 3). This samples is 1.2% divergent at the cox1 locus and 0.4%
divergent at the 28S locus when compared to Pacific samples, which could indicate a species-
level difference or not. However, it is also has much shorter and thinner spicules that are not
fusiform, and these combined morphological and genetic differences indicate it is likely a
different, possibly undescribed, species. These species fall in a larger clade including *H. kitchingi*,
H. pierrei, *S. cribrosa* and samples from Genbank identified as belonging to a variety of other
genera (Figure S1).
2382
2383 ***Hymeniacidon globularis* Ott, McDaniel & Humphrey, 2024**
2384
2385 **Material examined**
2386 None
2387
2388 **Diagnosis**
2389 Light yellow globular sponges with a prominent central oscule; subtylostyles with elliptical
2390 heads are the only spicule. Distinguished from all other *Hymeniacidon* in the region by
2391 "puffball" shape, large styles (300–450 µm long and 5–13 µm wide), and having elliptical
2392 subtylostyles as the primary spicule shape.
2393
2394 **Distribution and habitat**
2395 Known only from the subtidal, 17-24 m, in British Columbia.
2396
2397 **Remarks**
2398 This species was recently described by Ott, McDaniel & Humphrey (Montagu 1814), and is
2399 included here for completeness. See their description for additional details.
2400
2401 **Genus *Semisuberites* Carter, 1877**
2402
2403 ***Semisuberites cribrosa* (Miklucho-Maclay, 1870)**
2404 Figure 26
2405
2406
2407 **Synonyms**
2408 Veluspa polymorpha var. cribrosa Miklucho-Maclay, 1870
2409 Semisuberites arctica Carter, 1877
2410 Stylissa stipitata de Laubenfels, 1961

2411 The World Porifera Database lists 22 additional synonyms (de Voogd, et al., 2024)

2412

2413

2414 Figure 26. *Semisuberites cribrosa*. A (BHAK-08809), B (BHAK-09371). C: cross-section showing
2415 choanosomal skeleton near sponge rim, with external surface on the left and internal surface on the
2416 right; round structures in center are cross-sections of spicule tracts (BHAK-08809). Spicules: E (BHAK-
2417 10379).

2418

2419 **Material examined**
2420 BHAK-08809, Nalau Pass, British Columbia, (51.784, -128.101), 30 m, 2019-05-25; BHAK-09371,
2421 Wedgeborough Point, British Columbia, (51.647, -127.956), 20 m, 2019-05-27; BHAK-10379,
2422 Wedgeborough Point, British Columbia, (51.647, -127.956), 25 m, 2019-05-27.

2423

2424 **Diagnosis**
2425 Soft, fragile, funnel-shaped sponges arising from a stalk. Styles are the only spicule, slightly
2426 fusiform, 110–450 µm long 1–12 µm wide (styles reportedly up to 600 µm long and 16 µm wide
2427 in other regions). Prominent spicule tracts run parallel to the surface of the sponge, branching
2428 off to intersect the sponge surface, where they form spicule brushes in the ectosome.

2429

2430 **Morphology**
2431 Funnel or trumpet-shaped sponges arising from stalks that attach to the substrate with root-like
2432 structures (Figure 26B). Stalk may branch and give rise to multiple funnels (Figure 26A). Sponges
2433 soft and fragile, with distal lip often frayed (Figure 26B). Velvety surface. Tan alive, white in
2434 ethanol.

2435

2436 **Skeleton**
2437 Branching spicule tracts run mostly parallel to the sponge surface, but give rise to tracts that
2438 branch off to intersect the sponge surface. These flare to form bouquets that pierce the surface
2439 of the sponge. Some spicules also confused throughout the choanosome and tangentially
2440 arranged within the ectosome (Figure 26C).

2441

Spicules
Nearly exclusively styles (Figure 26BD), straight or evenly curved, some subtylote but many
simple; rare strongyles occasionally seen. Distributions of length were bimodal or trimodal for
each sponge, but continuous with unclear size classes. When spicules from three samples are
combined, they measure 118–285–446 x 1–8–12 µm, n=260 for length, n=253 for width. The

combined length distribution is trimodal, with modes at approximately 210, 310, and 390 µm.
Means per sponge were 259 to 301 µm in length and 7 to 8 µm in width.
2449
2450 **Distribution and habitat**
2451 Known from the Arctic, North Atlantic, and North Pacific, from 20–289 m depth (Dinn et al.,
2452 2020). The Southernmost Atlantic specimens are known from the Gulf of St. Lawrence (Dinn et
2453 al., 2020), while the Southernmost Pacific specimens are known from the San Juan Islands,
2454 Washington (de Laubenfels, 1961).
2455
Remarks
The choanosomal skeletons of halichondriid sponges are often described as "confused", but this
term can also be applied to the taxonomy of many species in this group. This species is no
exception. Populations of this species have been described as members of various families and
genera; in the Northeast Pacific, these sponges were first described as *Stylissa stipitata* (de
Laubenfels, 1961); it now seems likely that this name is a junior synonym of *Semisuberites*
cribrosa. The samples we examined here agree with the recent morphological description of
samples from the North Atlantic, though Atlantic samples had slightly larger spicules (Dinn et
al., 2020).
The genus *Semisuberites* is monotypic and has been difficult to place. It was assigned to
the Suberitidae by Topsent (1928); Van Soest and Hajdu (2002) noted similarities with
Hymeniacidon but tentatively placed the genus in the Esperiopsidae (Order Poecilosclerida).
Van Soest (2016) later proposed that this genus is more likely to be in the order Suberitida. Here
we present the first genetic data for this species, which clearly places it within the Suberitida.
Based on morphological data and genetic similarity to some *Hymeniacidon* species, we propose
placement in the Halichondriidae. The upright brushes of styles in the ectosome are unusual for
this family, but there do appear to be some tangential spicules in the ectosome as well.
2473
2474 ***Topsentia fernaldi* (Sim and Bakus, 1986)**
2475 Figure 27
2476
2477 **Synonyms**
2478 Oxeostilon fernaldi Sim & Bakus, 1986
2479
2480 **Material examined**
2481 Holotype: USNM 33629, Bird Rock, Santa Catalina Island, California, (33.45000, -118.41667), 46
2482 m, 1968-11-20. Other samples: TLT1134, Honeymooners, California, (36.50390, -121.94100), 10-
2483 20 m, 2021-09-22; TLT1261, Honeymooners, California, (36.50390, -121.94100, 10-20 m, 2021-
2484 09-22.
2485
Diagnosis
Thickly encrusting, irregular masses, yellow or chocolate brown exteriorly with yellow interiors.
Sponges are hard and incompressible due to dense spiculation. The ectosomal skeleton is a
crust of tangential spicules in a sieve-like pattern or a disordered mat. The choanosome is very
dense with disordered spicules, with many oriented vertically near the surface, their tips

piercing the ectosome. Spicules are oxeas in a large size range, 160–1170 μm long and 4–35 μm
wide. Distribution of length is strongly bimodal, with modes near 250 and 850 μm. Spicules
average 50% shorter in the ectosome, where most spicules are less than 600 μm long, but the
full size-range is present in the choanosome.
2495

2496
2497 Figure 27. *Topsentia fernaldi*. A (TLT1134), B (TLT1261). C & D: tangential ectosomal skeleton (TLT1261).
2498 Cross-sections show choanosomal skeleton at sponge surface (E) and farther towards the interior (F),
2499 TLT1261. G: long and short oxeas from holotype, USNM 33629.
2500

Morphology

The holotype was previously described as a thickly encrusting yellow mass, firm but friable, with
a crusty surface texture and sparse oscula. Newly collected samples were thick, irregular
encrustations, chocolate brown on the surface and pale yellow interiorly. Their interiors were
full of bryozoan skeletons and worm tubes which are presumed to have been overgrown by the
sponge, and the exterior was heavily fouled with hydroids, bryozoans, and other organisms.
Oscula were few, small, scattered, and flush with surface. Samples are white in ethanol.
2508

Skeleton

The original description states only that spicules are "distributed in an irregular fashion". We
examined only a small portion of the holotype, and did not locate ectosomal regions. The
choanosomal skeleton was confused, with a higher density of spicules than *H. panicea*, but a
lower density than the recently collected samples from Central California.
Freshly collected samples from Central California had tangential ectosomal skeletons
developed in some places as a sieve-like pattern around open spaces that were oval or
polygonal, and in other places as a dense mat of discorded spicules. The choanosome was
densely populated with spicules in confusion. Spicules became somewhat less confused towards
the ectosome, where the majority are oriented vertically, with the final ones piercing the
surface to create microscopic hispidity. No apparent spongin.

Spicules

Nearly exclusively curved oxeas, with rare styles or strongyles sometimes present. Distributions of length were strongly bimodal, but continuous distributions made it difficult to define size classes. Spicules from the holotype measure 224–568–1129 x 7–15–31 µm, n=115; length modes at about 250 µm and 850 µm. When data from the holotype are combined with the two newly collected samples, spicules measure 168–511–1161 x 4–15–35 µm, n=499. See table 11 for data from each sample individually. When spicules are isolated from the ectosome and choanosome separately, the full range of lengths is seen in the choanosome, but ectosomal spicules are mostly from the shorter half of the length distribution (ectosome mean length 330 µm vs. choanosome mean length 614 µm).

The holotype was originally described as having small oxeas, large oxeas, styles, and strongyles, with "styles and especially strongyles uncommon". Our spicule preparation from the holotype found only 2 styles, one tylostyle, and no strongyles, out of several hundred spicules, so we believe these to be rare aberrations (one style was also seen in another sample). We also do not consider oxeas to be present in size classes, because the length distribution is continuous, but the modes of the length distribution are very similar to the means of the size classes previously reported (239 and 813 µm).

Table 11. Spicule lengths of *Topsentia fernaldi*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT1134	177–406–880 n=45	218–648–1021 n=46
TLT1261	168–282–1161 n=70	194–592–974 n=70
Topsentia aff. fibrosa:		
RBC10	288–463–977 n=95	300–667–1020 n=72

Distribution and habitat

Described from a single sample collected at Catalina Island, in Southern California, at 46 m. Two additional samples were recently collected in Carmel Bay in Central California, between 10-20 m. Both locations were natural rocky reefs.

Remarks

The two sponges collected from Central California fit well into the current concept of *Topsentia*: they have a well-developed tangential ectosomal spicule crust that is difficult to remove, a densely confused choanosomal skeleton, and oxeas in a large size range, with smaller spicules at the sponge surface (Erpenbeck and van Soest, 2002). However, we have reservations in assigning them to *T. fernaldi.* The original description for this species is sparse, but it was described as "moderately dense, firm but friable", rather than stony and incompressible, as the newly collected samples are. The small fragment of the holotype we examined did appear to be less densely spiculated, consistent with this difference. However, the spicules from the new samples are very similar to the *T. fernaldi* holotype, and DNA extractions from the holotype were not successful. We therefore assign the new samples to this species, but would not be surprised that future collections yield new material that would result in splitting them.

We also examined a sponge from British Columbia, previously identified by William
Austin as *Topsentia* aff. *fibrosa*. This sponge was morphologically indistinguishable from the
Central California samples (table 11), but DNA extractions were again unsuccessful. It is
therefore possible that *T. fernaldi* ranges farther North, but additional sampling of this genus in
the Northeast Pacific is badly needed. *Topsentia fibrosa* is an Arctic species that may range into
British Columbia, but has not been confirmed there. The original description of *T. fibrosa* is
limited, but does state that samples are dark brown after preservation in spirit (Fristedt, 1887),
while all the samples examined here were white after preservation in ethanol.
The presence of *T. fernaldi* within the *Panicea* species group is very surprising (Figure 1).
While much more distantly related species are morphologically indistinguishable from *H.*
panicea, the spicule dimensions and density make this species quite distinct. A few other
Topsentia species have genetic data available, and some appear to fall within entirely different
orders of demosponges (Pankey et al., 2022; Redmond et al., 2013). These results serve to
emphasize how evolutionary liable spicule size, shape, and density are in this clade.
2572
2573 ***Topsentia disparilis* (Lambe, 1893)**
2574
2575 **Synonyms**
2576 Halichondria disparilis Lambe, 1893
2577
2578 **Material examined**
2579 None.
2580
Description
First described from a single massive, flat sponge, 35 x 28 mm across and 11 mm thick. Spicules
are exclusively or nearly exclusively oxeas, but in a very wide range of lengths, with two size-
classes (or perhaps modes in a continuous distribution). Large oxeas originally described as
438–1287 μm long, averaging 13 μm thick (Lambe, 1893); a later report states they are 900 x 20
μm (de Laubenfels, 1953). Short oxeas reported as 130 x 8 μm (de Laubenfels, 1953) or 91 x 4
(Lambe, 1893). Choanosomal skeleton densely confused; ectosomal skeleton variously reported
as a tangential reticulation of oxeas (Lambe, 1893) or bristling with erect spicules (de
Laubenfels, 1953). The latter report of this species also notes it has scattered areas covered in
low, small tubercles less than 1 mm high (de Laubenfels, 1953).
2591
2592 **Distribution and habitat**
2593 The holotype was collected at 73 m near Comox, Vancouver Island, British Columbia, and
2594 additional samples were reported from 38–45 m near Point Barrow, AK (de Laubenfels, 1953).
2595
Remarks
We did not examine any material from this species, but it is included here for completeness.
Published descriptions do not clearly differentiate this species from *T. fibrosa* (Fristedt, 1887),
which is also known from the Arctic, and may be a senior synonym. As stated in the *T. fernaldi*
section above, further revision of *Topsentia* diversity in the Northeast Pacific is badly needed,
but likely requires freshly collected material and additional DNA data.

2602

2603   **Genus *Axinyssa* Lendenfeld, 1897**

2604

2605   ***Axinyssa piloerecta* sp. nov.**

2606   Figure 28

2607

2608
2609   Figure 28. *Axinyssa piloerecta*. Sample shown in situ (A) and cross-sectioned after collection, before
2610   preservation (B). C: sample TLT1132 after collection, before preservation. Cross-sections show
2611   choanosomal skeleton at sponge surface (D) and farther towards the interior (E). F: example oxeas, G:
2612   example style. All photos from TLT1007 except C.

2613

2614   **Material examined**
2615   TLT1007 (holotype) and TLT1132 (paratype), both collected at Yellowbanks, Santa Cruz Island
2616   (33.99432, -119.51910), 24-30 m, 2021-09-15.

2617

2618   **Etymology**
2619   From the word piloerection, the process that causes skin to form "goosebumps" and hair to
2620   stand up, creating a pattern similar to the surface of this sponge.

2621

Diagnosis
Yellow cushion-shaped sponges with widely-spaced surface conules. The combination of a
smooth, slick-looking surface and conules with protruding spicule bundles is distinctive in the
region. This contrasts with the sympatric *Obruta collector* Turner & Lonhart 2024, which has
similar spicules but lacks raised bumps and is hispid over the entire surface. The large oxeas
(530–1480 x 5–33 µm) and the significant fraction of spicules that are styles (18%) further
distinguish this species from others in the region.

2629

2630   **Morphology**

Cushion-shaped irregular mounds, pale yellow when alive and beige in ethanol. The surface is covered in raised projections, 1–3 mm apart, 1–3 mm in height, with spicules projecting an additional 0.5 mm. When alive, the holotype was approximately 60 x 35 mm across, 24 mm in maximum thickness; sampled portion approximately 35 x 35 x 24 mm. A distinct surface membrane covers the sponge, creating a smooth and slick-looking surface except where spicules protrude from raised conules.

Skeleton
A dense, confused mass of spicules internally, with no apparent organization. Spicules become roughly parallel towards the periphery, pointing outwards, and form vague plumose tracts. Some of these tracts end in tufts of spicules that pierce the surface of the sponge. No ectosomal skeleton. Spongin not apparent, but body of sponge is collagenous and dissolves very slowly in bleach, in contrast to all other Halichondridae described here, which dissolved very rapidly.

Spicules Styles and oxeas, with some intermediates; one strongyle also present. Oxeas are curved or bent, usually near center, sometimes with weak centrotylote swelling (Figure 28F). Dimensions highly variable, but without clear size categories; 533–957–1476 x 5–24–33 µm (n=78). Styles are less common than oxeas (18% of spicules), but with similar dimensions. 606–1178–2166 x 5–17–27 µm (n=17).

Distribution and habitat. Found at only a single location: a rocky reef near Yellowbanks, Santa Cruz Island. The site is a rocky outcrop, typical of other rocky reefs in the region, but most other sites searched in the Northern Channel Islands have been in less than 20 m of water, while this site was 24-30 m in depth. The species appeared to be common at the site at the time of collection.

Remarks
The morphology of this species places it firmly within the *Axinyssa* (Erpenbeck and van Soest, 2002). Like the genus *Topsentia*, the genus *Axinyssa* is highly polyphyletic, with some species, including the type species, falling among the order Bubarida and other species falling in the order Biemnida (Morrow and Cárdenas, 2015; Morrow et al., 2013; Pankey et al., 2022; Thacker et al., 2013). Genetic data from this species was previously reported as placing it clearly in the order Axinellida (Turner and Pankey, 2023). We describe it here because in the current taxonomy, this genus remains within the Halichondriidae, but we expect future taxonomic revision to move this species to the Axinellida.

Axinyssa tuscara **(Ristau, 1978)**

Synonyms
Axinomimus tuscarus Ristau, 1978

Material examined
Holotype: USNM 24527, Carmel River Beach Point, Carmel, California, 7 m, 1973-04-10.

Description

Thickly encrusting dark brown sponges that remain brown after preservation in ethanol. Soft and compressible. Choanosomal skeleton of vertical spicule tracts that pierce sponge and project up to 1.5 mm beyond sponge surface, connected by a confused reticulation of individual spicules. No special ectosomal skeleton. Spicules curved oxeas, longer and considerably thicker than sympatric *Halichondria*, 355–476–552 x 10–18–22 µm, n=47.

Distribution and habitat

Collected in situ only from the type location: Carmel River Beach Point, Central California, on a rock reef at 7 m depth. It was described as rare at this location (Ristau, 1978). Several samples were also collected in beach wrack in Northern California at Bodega Head.

Remarks

All known samples of this species were collected from 1973 to 1977 and were included in the original description (Ristau, 1978). We examined a small piece of the holotype in order to attempt DNA extraction, which failed. We also performed a spicule preparation in order to publish more detailed information on the distribution of spicule sizes, as described above. It was difficult to characterize the skeleton in the small piece we examined, and could only determine that the chromosomal skeleton was confused.

Acknowledgements

We are grateful for the help and support of many people in UCSB's Marine Science Institute and Diving & Boating Program, especially Robert Miller, Clint Nelson, Christoph Pierre, and Christian Orsini. Numerous NOAA personal provided logistics and planning support for collections in the Monterey Bay and Olympic Coast National Marine Sanctuaries. Raquel Pereira graciously shared samples from Sweden and Portugal, and Brooke Weigel shared samples from Tatoosh, Washington. Gustav Paulay, Hugh MacIntosh, Abigail Reft and Christina Piotrowski provided access to their collections at the Florida Museum of Natural History, Royal British Columbia Museum, Smithsonian National Museum of Natural History, California Academy of Sciences, respectively. We also thank Matt Whalen, Margot Hessing-Lewis, and the staff, students and volunteers of the Hakai Institute for support during the 2017 Hakai-MarineGeo bioblitz, and the Makah Tribal Nation for supporting the study of sponges from Tatoosh Island. We thank DFO colleagues Andrew Cooper, Peter Lawton, Laura Teed and Torben Brydges from Saint Andrews Biological Station, who partnered on the Eastern Shores habitat survey project and participated in specimen collection.

Funding

Financial support was provided by UCSB and by the National Aeronautics and Space Administration Biodiversity and Ecological Forecasting Program (Grant NNx14Ar62A); the Bureau of Ocean Energy Management Environmental Studies Program (BOEM Agreement MC15AC00006); the National Oceanic and Atmospheric Administration (NOAA) in support of the Santa Barbara Channel Marine Biodiversity observation Network; and the U.S. National Science Foundation (NSF) in support of the Santa Barbara Coastal Long Term Ecological research program under Awards OCE-9982105, OCE-0620276, OCE-1232779, OCE-1831937, and the Tula

Foundation. Additional support was provided by the NSF under award EF-2025121 to RWT and funding from the Department of Agriculture Environment and Rural Affairs Environment Fund to CM. Funding for collection of specimens from the Eastern Shore Islands, Nova Scotia was covered as part of a habitat and sponge biodiversity survey funded by Fisheries and Oceans Canada. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Supporting Information

Data available from the Dryad Digital Repository: link to be generated upon manuscript submission.

References

Ackers, R.G., Moss, D., Picton, B.E., Stone, S.M.K., Morrow, C.C., 2007. Sponges of the British Isles: A color guide and working document. Marine Conservation Society, Ross-on-Wye.

Agne, S., Ekins, M., Galitz, A., Hofreiter, M., Preick, M., Straube, N., Wörheide, G., Erpenbeck, D., 2022. Keratose sponge MuseOMICS: setting reference points in dictyoceratid demosponge phylogeny. Zootaxa 5195, 296–300. https://doi.org/10.11646/zootaxa.5195.3.9

Archibald, D.J., 2003. Timing and biogeography of the eutherian radiation: fossils and molecules compared. Mol. Phylogenet. Evol. 28, 350–359. https://doi.org/10.1016/S1055-7903(03)00034-4

Barthel, D., 1986. On the ecophysiology of the sponge Halichondria panicea in Kiel Bight. I. Substrate specificity, growth and reproduction. Mar. Ecol. Prog. Ser. 32, 291–298. https://doi.org/10.3354/meps032291

Barthel, D., Detmer, A., 1990. The spermatogenesis of Halichondria panicea (Porifera, Demospongiae). Zoomorphology 110, 9–15. https://doi.org/10.1007/BF01632807

Barthel, D., Wolfrath, B., 1989. Tissue sloughing in the sponge Halichondria panicea: a fouling organism prevents being fouled. Oecologia 78, 357–360. https://doi.org/10.1007/BF00379109

Bergquist, P.R., 1970. The marine fauna of New Zealand: Porifera, Demospongiae, part 2 (Axinellida and Halichondrida). N. Z. Oceanogr. Inst. Mem. 51, 9–85.

Bowerbank, J.S., 1866. A Monograph of the British Spongiadae. Volume 2. Ray Society, London.

Bulleri, F., Chapman, M.G., 2010. The introduction of coastal infrastructure as a driver of change in marine environments. J. Appl. Ecol. 47, 26–35. https://doi.org/10.1111/j.1365-2664.2009.01751.x

Burton, M., 1935. Notes on British sponges, with a description of a new genus and species. Ann. Mag. Nat. Hist. 10, 651–653.

Burton, M., 1930. Additions to the sponge fauna at Plymouth. J. Mar. Biol. Assoc. U. K. 16, 489–507.

Carlton, J.T., Eldredge, L.G., 2009. Marine bioinvasions of Hawaiʻi: the introduced and cryptogenic marine and estuarine animals and plants of the Hawaiian Archipelago, Bishop Museum bulletin in cultural and environmental studies. Bernice P. Bishop Museum/Bishop Museum Press, Honolulu, Hawaiʻi.

Carrier, T.J., Schmittmann, L., Jung, S., Pita, L., Hentschel, U., 2023. Maternal provisioning of an
obligate symbiont in a sponge. Ecol. Evol. 13, e10012.
https://doi.org/10.1002/ece3.10012
Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34, i884–i890. https://doi.org/10.1093/bioinformatics/bty560
Cóndor-Luján, B., Arteaga, A., Polo, C., Arroyo, Y., Willenz, P., Hajdu, E., 2023. Shallow
Suberitida (Porifera, Demospongiae) from Peru. Zootaxa 5264, 451–489.
https://doi.org/10.11646/zootaxa.5264.4.1
Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane,
T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and BCFtools.
GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008
de Laubenfels, M.W., 1961. Porifera of Friday Harbor and Vicinity. Pac. Sci. 15, 192–202.
de Laubenfels, M.W., 1953. Sponges of the Alaska Arctic. Smithson. Misc. Collect. 121, 1–21.
de Laubenfels, M.W., 1932. The marine and fresh-water sponges of California. Proc. U. S. Natl.
Mus. 81, 1–140.
de Laubenfels, M.W., 1930. The Sponges of California. Stanf. Univ. Bull., Abstracts of
dissertations for the degree of doctor of philosophy 5, 24–29.
de Paula, T.S., Zilberberg, C., Hajdu, E., Lôbo-Hajdu, G., 2012. Morphology and molecules on
opposite sides of the diversity gradient: Four cryptic species of the Cliona celata
(Porifera, Demospongiae) complex in South America revealed by mitochondrial and
nuclear markers. Mol. Phylogenet. Evol. 62, 529–541.
https://doi.org/10.1016/j.ympev.2011.11.001
de Voogd, N.J., Alvarez, B., Boury-Esnault, N., Cárdenas, P., Díaz, C., Dohrmann, M., Downey, R.,
Goodwin, C., Hajdu, E., Hooper, J.N.A., Kelly, M., Klautau, M., Lim, S.C., Monconi, R.,
Morrow, C.C., Pinheiro, U., Pisera, A.B., Ríos, P., Rützler, K., Schönberg, C., Turner, T.L.,
Vacelet, J., van Soest, R.W.M., Xavier, J., 2024. World Porifera Database [WWW
Document]. URL https://www.marinespecies.org/porifera on 2023-01-03.
doi:10.14284/359 (accessed 1.1.24).
Delić, T., Trontelj, P., Rendoš, M., Fišer, C., 2017. The importance of naming cryptic species and
the conservation of endemic subterranean amphipods. Sci. Rep. 7.
Dinn, C., Leys, S.P., Roussel, M., Méthé, D., 2020. Geographic range extensions of stalked,
flabelliform sponges (Porifera) from eastern Canada with a new combination of a
species of Plicatellopsis in the North Atlantic. Zootaxa 4755, 301–321.
https://doi.org/10.11646/zootaxa.4755.2.6
Dohrmann, M., Wörheide, G., 2017. Dating early animal evolution using phylogenomic data. Sci.
Rep. 7, 3599. https://doi.org/10.1038/s41598-017-03791-w
Dunn, C.W., Leys, S.P., Haddock, S.H.D., 2015. The hidden biology of sponges and ctenophores.
Trends Ecol. Evol. 30, 282–291. https://doi.org/10.1016/j.tree.2015.03.003
Eitel, M., Francis, W.R., Varoqueaux, F., Daraspe, J., Osigus, H.-J., Krebs, S., Vargas, S., Blum, H.,
Williams, G.A., Schierwater, B., Wörheide, G., 2018. Comparative genomics and the
nature of placozoan species. PLOS Biol. 16, e2005359.
https://doi.org/10.1371/journal.pbio.2005359

Erpenbeck, D., 2004. A molecular comparison of Alaskan and North East Atlantic Halichondria panicea (Pallas 1766) (Porifera:Demospongiae) populations. BMIB - Boll. Dei Musei E Degli Ist. Biol. 68.

Erpenbeck, D., Hooper, J.N.A., Wörheide, G., 2006. CO1 phylogenies in diploblasts and the 'Barcoding of Life'— are we sequencing a suboptimal partition? Mol. Ecol. Notes 6, 550–553. https://doi.org/10.1111/j.1471-8286.2005.01259.x

Erpenbeck, D., van Soest, R.W.M., 2002. Family Halichondriidae Gray, 1867, in: Hooper, J.N.A., van Soest, R.W.M. (Eds.), Systema Porifera. A Guide to the Classification of Sponges. Kluwer Academic/Plenum Publ., New York, pp. 787–816.

Erwin, P.M., Thacker, R.W., 2007. Phylogenetic analyses of marine sponges within the order Verongida: a comparison of morphological and molecular data. Invertebr. Biol. 126, 220–234. https://doi.org/10.1111/j.1744-7410.2007.00092.x

Evcen, A., Gözcelioğlu, B., Çinar, M.E., 2023. New sponge records (Porifera) from the Black Sea. Reg. Stud. Mar. Sci. 65, 103043. https://doi.org/10.1016/j.rsma.2023.103043

Folmer, O., Black, M., Wr, H., Vrijenhoek, R., 1994. DNA primers for amplification of mitochondrial Cytochrome C oxidase subunit I from diverse metazoan invertebrates. Mol. Mar. Biol. Biotechnol. 3, 294–299.

Forester, A.J., 1979. The association between the sponge Halichondria panicea (Pallas) and scallop Chlamys varia (L.): A commensal-protective mutualism. J. Exp. Mar. Biol. Ecol. 36, 1–10. https://doi.org/10.1016/0022-0981(79)90096-0

Fristedt, K., 1887. Sponges from the Atlantic and Arctic Oceans and the Behring Sea. Vega-Exped. Vetensk. Iakttag. Nord. 4, 410–471.

Gazave, E., Carteron, S., Chenuil, A., Richelle-Maurer, E., Boury-Esnault, N., Borchiellini, C., 2010. Polyphyly of the genus Axinella and of the family Axinellidae (Porifera: Demospongiae). Mol. Phylogenet. Evol. 57, 35–47. https://doi.org/10.1016/j.ympev.2010.05.028

Glasby, T.M., Connell, S.D., Holloway, M.G., Hewitt, C.L., 2007. Nonindigenous biota on artificial structures: could habitat creation facilitate biological invasions? Mar. Biol. 151, 887–895. https://doi.org/10.1007/s00227-006-0552-5

Harbo, R., Ott, B., Reiswig, H., McDaniel, N., 2021. First Canadian record (Ladysmith Harbour, British Columbia) of the non-native European sponge Hymeniacidon perlevis (Montagu, 1814) (Porifera, Demospongiae). BioInvasions Rec. 10, 277–286. https://doi.org/10.3391/bir.2021.10.2.05

Hartman, W., 1958. Natural history of the marine sponges of southern New England. Bull. Peabody Mus. Nat. Hist. 1–155.

Hirata, Y., Uemura, D., 1986. Halichondrins - antitumor polyether macrolides from a marine sponge. Pure Appl. Chem. 58, 701–710. https://doi.org/10.1351/pac198658050701

Hoang, D., Chernomor, O., von Haeseler, A., Minh, B., Vinh, L., 2018. UFBoot2: Improving the ultrafast bootstrap approximation. Mol. Biol. Evol. 35, 518–522. https://doi.org/10.1093/molbev/msx281

Hooper, J.N.A., Van Soest, R.W.M., 2002. Systema Porifera. A Guide to the Classification of Sponges, in: Hooper, J.N.A., Van Soest, R.W.M., Willenz, P. (Eds.), Systema Porifera: A Guide to the Classification of Sponges. Springer US, Boston, MA, pp. 1–7. https://doi.org/10.1007/978-1-4615-0747-5_1

Jennings, H.S., 1917. The numerical results of diverse systems of breeding, with respect to two
pairs of characters, linked or independent, with special relation to the effects of linkage.
Genetics 2, 97–154.
Johnson, G., 1842. A history of British sponges and lithophytes. W.H. Lizars, Edinburgh.
Jörger, K.M., Schrödl, M., 2013. How to describe a cryptic species? Practical challenges of
molecular taxonomy. Front. Zool. 10, 59. https://doi.org/10.1186/1742-9994-10-59
Jun, J., Yu, J.-N., Choi, E.H., 2015. Complete mitochondrial genome of Hymeniacidon sinapium
(Demospongiae, Halichondriidae). Mitochondrial DNA 26, 261–262.
https://doi.org/10.3109/19401736.2013.823189
Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods
14, 587–589. https://doi.org/10.1038/nmeth.4285
Kang, D.W., Kim, H., Kim, H.J., 2022. New species of Halichondria (Suberitida: Halichondriidae)
from Korea. J. Asia-Pac. Biodivers. 15, 98–101.
https://doi.org/10.1016/j.japb.2021.10.004
Katoh, K., Rozewicki, J., Yamada, K.D., 2017. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief. Bioinform. 20, 1160–
1166. https://doi.org/10.1093/bib/bbx108
Kim, H., Kang, D.W., Yu, C., Jung, Y.H., Yoon, M., Kim, H.J., 2019. The complete mitochondrial
genome of sponge Halichondria sp. (Demospongiae, Suberitida, Halichondriidae) from
Dok-do. Mitochondrial DNA Part B 4, 1729–1730.
https://doi.org/10.1080/23802359.2019.1610099
Knobloch, S., Jóhannsson, R., Marteinsson, V., 2019. Bacterial diversity in the marine sponge
Halichondria panicea from Icelandic waters and host-specificity of its dominant
symbiont "Candidatus Halichondribacter symbioticus." FEMS Microbiol. Ecol. 95, fiy220.
https://doi.org/10.1093/femsec/fiy220
Lambe, L.M., 1893. Sponges from the Pacific coast of Canada. Trans. R. Soc. Can. 11, 25–43.
Lawley, J.W., Gamero-Mora, E., Maronna, M.M., Chiaverano, L.M., Stampar, S.N., Hopcroft,
R.R., Collins, A.G., Morandini, A.C., 2021. The importance of molecular characters when
morphological variability hinders diagnosability: systematics of the moon jellyfish genus
Aurelia (Cnidaria: Scyphozoa). PeerJ 9, e11954. https://doi.org/10.7717/peerj.11954
Lee, W., Elvin, D., Reiswig, H., 2007. The sponges of California: A guide and key to the marine
sponges of California. Monterey Bay Sanctuary Foundation, Monterey, CA.
Letunic, I., Bork, P., 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Res. 47, W256–W259. https://doi.org/10.1093/nar/gkz239
Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2015. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics 31, 1674–1676. https://doi.org/10.1093/bioinformatics/btv033
Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinforma. Oxf. Engl. 25, 1754–1760.
https://doi.org/10.1093/bioinformatics/btp324
Luo, Z.-X., Yuan, C.-X., Meng, Q.-J., Ji, Q., 2011. A Jurassic eutherian mammal and divergence of
marsupials and placentals. Nature 476, 442–445. https://doi.org/10.1038/nature10291

Maldonado, M., 2006. The ecology of the sponge larva. Can. J. Zool. 84, 175–194.
https://doi.org/10.1139/z05-177
Mehbub, M.F., Yang, Q., Cheng, Y., Franco, C.M.M., Zhang, W., 2024. Marine sponge-derived
natural products: trends and opportunities for the decade of 2011-2020. Front. Mar. Sci.
11. https://doi.org/10.3389/fmars.2024.1462825
Miklucho-Maclay, N.N., 1870. Uber einige Schwämme des nördlichen Stillen Oceans und des
Eismeeres, welche im Zoologischen Museum der Kaiserlichen Akademie der
Wissenschaften in St. Petersburg aufgestellt sind. Ein Beitrag zur Morphologie und
Verbreitung der Spongien. Mém. Académie Impériale Sci. St Petersbourg 15, 1–24.
Montagu, G., 1814. An Essay on sponges, with descriptions of all the species that have been
discovered on the coast of Great Britain. Mem. Wernerian Nat. Hist. Soc. 2, 67–122.
Morrow, C., Cárdenas, P., 2015. Proposal for a revised classification of the Demospongiae
(Porifera). Front. Zool. 12, 1–27. https://doi.org/10.1186/s12983-015-0099-8
Morrow, C.C., Picton, B.E., Erpenbeck, D., Boury-Esnault, N., Maggs, C.A., Allcock, A.L., 2012.
Congruence between nuclear and mitochondrial genes in Demospongiae: A new
hypothesis for relationships within the G4 clade (Porifera: Demospongiae). Mol.
Phylogenet. Evol. 62, 174–190. https://doi.org/10.1016/j.ympev.2011.09.016
Morrow, C.C., Redmond, N.E., Picton, B.E., Thacker, R.W., Collins, A.G., Maggs, C.A., Sigwart,
J.D., Allcock, A.L., 2013. Molecular phylogenies support homoplasy of multiple
morphological characters used in the taxonomy of Heteroscleromorpha (Porifera:
Demospongiae). Integr. Comp. Biol. 53, 428–446. https://doi.org/10.1093/icb/ict065
Müller, O.F., 1776. Zoologiae Danicae prodromus, seu Animalium Daniae et Norvegiae
indigenarum: characteres, nomina, et synonyma imprimis popularium. Typis Hallagerii,
Havni, Copenhagen.
Nguyen, L.-T., Schmidt, H., von Haeseler, A., Minh, B., 2015. IQ-TREE: A fast and effective
stochastic algorithm for estimating maximum likelihood phylogenies. Mol. Biol. Evol. 32,
268–274. https://doi.org/10.1093/molbev/msu300
Ott, B., Mcdaniel, N., Humphrey, E., 2024. Fourteen new species of demosponges (Porifera)
from three coastal fjords in southern British Columbia, Canada. Zootaxa 5463, 151–200.
https://doi.org/10.11646/zootaxa.5463.2.1
Pallas, P.S., 1766. Elenchus zoophytorum sistens generum adumbrationes generaliores et
specierum cognitarum succintas descriptiones, cum selectis auctorum synonymis.
Fransiscum Varrentrapp, Hagae.
Palumbi, S.R., 1986. How body plans limit acclimation: Responses of a Demosponge to wave
force. Ecology 67, 208–214. https://doi.org/10.2307/1938520
Pankey, M.S., Plachetzki, D.C., Macartney, K.J., Gastaldi, M., Slattery, M., Gochfeld, D.J., Lesser,
M.P., 2022. Cophylogeny and convergence shape holobiont evolution in sponge-microbe
symbioses. Nat. Ecol. Evol. 6, 750–762. https://doi.org/10.1038/s41559-022-01712-3
Park, M.-H., Sim, C.-J., Baek, J., Min, G.-S., 2007. Identification of genes suitable for DNA
barcoding of morphologically indistinguishable Korean Halichondriidae sponges. Mol.
Cells 23, 220–227. https://doi.org/10.1016/S1016-8478(23)07377-6
Rannala, B., Yang, Z., 2020. Chapter 5.5 Species Delimitation, in: Scornavacca, C., Delsuc, F.,
Galtier, N. (Eds.), Phylogenetics in the Genomic Era. No commercial publisher, pp. 1–
568.

Redmond, N.E., Morrow, C.C., Thacker, R.W., Diaz, M.C., Boury-Esnault, N., Cárdenas, P., Hajdu,
E., Lôbo-Hajdu, G., Picton, B.E., Pomponi, S.A., Kayal, E., Collins, A.G., 2013. Phylogeny
and systematics of Demospongiae in light of new small-subunit ribosomal DNA (18S)
sequences. Integr. Comp. Biol. 53, 388–415. https://doi.org/10.1093/icb/ict078
Ristau, D.A., 1978. Six new species of shallow-water marine demosponges from California. Proc.
Biol. Soc. Wash. 91, 569–589.
Rodriguez Jimenez, A., Dechamps, E., Giaux, A., Goetghebuer, L., Bauwens, M., Willenz, P.,
Flahaut, S., Laport, M.S., George, I.F., 2021. The sponges Hymeniacidon perlevis and
Halichondria panicea are reservoirs of antibiotic-producing bacteria against multi-drug
resistant Staphylococcus aureus. J. Appl. Microbiol. 131, 706–718.
https://doi.org/10.1111/jam.14999
Rossi, M.E., Kenny, N.J., Giacomelli, M., Keating, J.N., Álvarez-Carretero, S., Schuster, A.,
Cárdenas, P., Taboada, S., Koutsouveli, V., Donoghue, P., Riesgo, A., Pisani, D., 2024.
Independent origins of spicules reconcile the evolutionary history of sponges.
https://doi.org/10.1101/2024.06.24.600355
Rot, C., Goldfarb, I., Ilan, M., Huchon, D., 2006. Putative cross-kingdom horizontal gene transfer
in sponge (Porifera) mitochondria. BMC Evol. Biol. 6, 1–11.
https://doi.org/10.1186/1471-2148-6-71
Ruiz, G.M., Fofonoff, P.W., Carlton, J.T., Wonham, M.J., Hines, A.H., 2000. Invasion of coastal
marine communities in North America: apparent patterns, processes, and biases. Annu.
Rev. Ecol. Evol. Syst. 31, 481–531. https://doi.org/10.1146/annurev.ecolsys.31.1.481
Samaai, T., Turner, T.L., Kara, J., Yemane, D., Ngwakum, B.B., Payne, R.P., Kerwath, S., 2022.
Confirmation of the southern African distribution of the marine sponge Hymeniacidon
perlevis (Montagu, 1814) in the context of its global dispersal. PeerJ 10, e14388.
https://doi.org/10.7717/peerj.14388
Schmidt, O., 1870. Grundzüge einer Spongien-Fauna des atlantischen Gebietes. Wilhelm
Engelmann, Leipzig.
Schmittmann, L., Franzenburg, S., Pita, L., 2021. Individuality in the immune repertoire and
induced response of the sponge Halichondria panicea. Front. Immunol. 12.
https://doi.org/10.3389/fimmu.2021.689051
Schneider, C., Rasband, W., Eliceiri, K., 2012. NIH Image to ImageJ: 25 years of image analysis.
Nat. Methods 9, 671–675. https://doi.org/10.1038/nmeth.2089
Sim, C.J., Bakus, G.J., 1986. Marine sponges of Santa Catalina Island, California. Occas. Pap.
Allan Hancock Found. 5, 1–23.
Simkanin, C., Davidson, I.C., Dower, J.F., Jamieson, G., Therriault, T.W., 2012. Anthropogenic
structures and the infiltration of natural benthos by invasive ascidians. Mar. Ecol. 33,
499–511. https://doi.org/10.1111/j.1439-0485.2012.00516.x
Tessler, M., Neumann, J.S., Kamm, K., Osigus, H.-J., Eshel, G., Narechania, A., Burns, J.A.,
DeSalle, R., Schierwater, B., 2022. Phylogenomics and the first higher taxonomy of
Placozoa, an ancient and enigmatic animal phylum. Front. Ecol. Evol. 10.
https://doi.org/10.3389/fevo.2022.1016357
Thacker, R., Hill, A., Hill, M., Redmond, N., Collins, A., Morrow, C., Spicer, L., Carmack, C., Zappe,
M., Pohlmann, D., Hall, C., Diaz, M., Bangalore, P., 2013. Nearly complete 28S rRNA gene

sequences confirm new hypotheses of sponge evolution. Integr. Comp. Biol. 53, 373–
387. https://doi.org/10.1093/icb/ict071
Thomassen, S., Riisgård, H.U., 1995. Growth and energetics of the sponge Halichondria panicea.
Mar. Ecol. Prog. Ser. 128, 239–246. https://doi.org/10.3354/meps128239
Topsent, E., 1928. Spongiaires de l'Atlantique et de la Mediterranee provenant des croisieres du
Prince Albert ler de Monaco. Result. Camp. Sci. Accompl. Par Ie Prince Albert Monaco
74, 1–376.
Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., Minh, B., 2016. W-IQ-TREE: a fast online
phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res. 44, W232–W235.
https://doi.org/10.1093/nar/gkw256
Turner, T.L., 2020. The marine sponge Hymeniacidon perlevis is a globally-distributed invasive
species. Bioarxve doi: 10.1101/2020.02.13.948000.
Turner, T.L., Lonhart, S.I., 2023. The sponges of the Carmel Pinnacles marine protected area.
Zootaxa 5318, 151–194. https://doi.org/10.11646/ZOOTAXA.5318.2.1
Turner, T.L., Pankey, M.S., 2023. The order Axinellida (Porifera: Demospongiae) in California.
Zootaxa 5230, 501–539. https://doi.org/10.11646/zootaxa.5230.5.1
van Soest, R.W.M., 2016. Sponge-collecting from a drifting ice floe: the Porifera obtained in the
Kara Sea by the Dutch Polar Expedition 1882-83. Contrib. Zool. 85, 311–336.
van Soest, R.W.M., Hajdu, E., 2002. Family Esperiopsidae Hentschel, 1923, in: Hooper, J.N.A.,
van Soest, R.W.M. (Eds.), Systema Porifera: A Guide to the Classification of Sponges.
Springer, Boston, pp. 656–664.
Vethaak, A.D., Cronie, J.A., van Soest, R.W.M., 1982. Ecology and distribution of two sympatric,
closely related sponge species, Halichondria panicea (Pallas, 1766) and H. bowerbanki
Burton, 1930 (Porifera, Demospongiae), with remarks on their speciation. Bijdr. Tot
Dierkd. 52.
Wang, D., Zhang, Y., Huang, D., 2016. The complete mitochondrial genome of sponge
Halichondria (Halichondria) sp. (Demospongiae, Suberitida, Halichondriidae).
Mitochondrial DNA Part B 1, 512–514. https://doi.org/10.1080/23802359.2016.1192516
Witte, U., Barthel, D., Tendal, O., 1994. The reproductive cycle of the sponge Halichondria
panicea Pallas (1766) and its relationship to temperature and salinity. J. Exp. Mar. Biol.
Ecol. 183, 41–52. https://doi.org/10.1016/0022-0981(94)90155-4
Xavier, J.R., Rachello-Dolmen, P.G., Parra-Velandia, F., Schönberg, C.H.L., Breeuwer, J. a. J., van
Soest, R.W.M., 2010. Molecular evidence of cryptic speciation in the "cosmopolitan"
excavating sponge Cliona celata (Porifera, Clionaidae). Mol. Phylogenet. Evol. 56, 13–20.
https://doi.org/10.1016/j.ympev.2010.03.030
Xue, L., Zhang, X., Zhang, W., 2009. Larval release and settlement of the marine sponge
Hymeniacidon perlevis (Porifera, Demospongiae) under controlled laboratory
conditions. Aquaculture 290, 132–139.
https://doi.org/10.1016/j.aquaculture.2009.01.037
Zhong, W., Olugbami, J.O., Rathakrishnan, P., Mohanty, I., Moore, S.G., Garg, N., Oyelere, A.K.,
Turner, T.L., McShan, A.C., Agarwal, V., 2024. Discovery and folding dynamics of a fused
bicyclic cysteine knot undecapeptide from the marine sponge Halichondria bowerbanki.
J. Org. Chem. 89, 12748–12752. https://doi.org/10.1021/acs.joc.4c01104
3022