A common garden of *Halichondria* sponges: taxonomic revision of Northeast Pacific Halichondriidae reveals many cryptic introduced species

Thomas L. Turner[1], Christine Morrow[2], Bernard Picton[3], Claire Goodwin[4,5,6], Robert W. Thacker[7]

[1]Ecology, Evolution, and Marine Biology Department, University of California, Santa Barbara, California, USA, tlturner@ucsb.edu
https://orcid.org/0000-0002-1380-1099

[2]School of Biological Sciences, Queen's University, Belfast, Northern Ireland, UK, c.morrow@qub.ac.uk
https://orcid.org/0000-0001-5686-8747

[3]National Museums Northern Ireland, Northern Ireland, UK, bernard.picton@nmni.com
https://orcid.org/0000-0002-1500-2215

[4]Huntsman Marine Science Center, St. Andrew, New Brunswick, Canada
[5]University of New Brunswick, 100 Tucker Park Road Saint John, NB E2L 4L5, Canada.
[6]New Brunswick Museum, 277 Douglas Ave, Saint John NB E2K 1E5, Canada, claire.goodwin@huntsmanmarine.ca
https://orcid.org/0000-0002-2428-1340

[7]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, USA, and Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Republic of Panama, robert.thacker@stonybrook.edu
https://orcid.org/0000-0002-9654-0073

Correspondence: tlturner@ucsb.edu

Zoobank: urn:lsid:zoobank.org:pub:866481F8-9313-45AE-8BC5-0CF458DCBF6E

Note about species names: this pre-print is not intended to be a publication of the associated species names for the purposes of zoological nomenclature

Abstract
Sponges (phylum Porifera) possess biochemical, cellular, and physiological traits with valuable
biotechnical applications. However, our ability to harness these natural innovations is limited by
a classification system that does not fully reflect their evolutionary history. In this study, we
uncover numerous cryptic species within the genus *Halichondria* that are morphologically
indistinguishable from the well-known *Ha. panicea*. Many of these species have habitat
preferences and geographic distributions that strongly suggest they have been dispersed by
human activity. Most of these species are broadly sympatric with their closest relatives, and
these overlapping distributions allow us to use patterns of DNA variation to infer reproductive
isolation between clades in nature. With reproductively isolated species thus delineated, we
can use DNA states as taxonomic characters to formally describe them. Though much remains
to be learned about these newly discovered species, the natural "common gardens" of these
sponges in California, New York, and other locations provide opportunities to test hypotheses
about their diversification in future work.
50

Introduction
The common ancestor of extant sponges is estimated to have lived over 700 million years ago
(Dohrmann and Wörheide, 2017; Pankey et al., 2022; Rossi et al., 2024). This ancient origin
suggests that some sponge lineages have been independently evolving for longer than the most
distantly related animals within Bilateria. Sponges have undoubtedly diversified in biochemical,
cellular and physiological dimensions during this prolonged time, developing numerous
innovations with potential applications (Dunn et al., 2015; Mehbub et al., 2024). These
innovations remain largely undiscovered, and it seems likely that a major impediment to
advancing sponge biology is the lack of a stable classification system that reflects the
evolutionary history of the phylum. The difficulty in assigning sponge specimens to species and
determining where those species fall in the tree of life creates uncertainty that deters
investigation across multiple disciplines.
Creating an accurate taxonomy of sponges based on morphological characters has
proven challenging due to a paucity of informative character states. Sponge taxonomy is based
mainly on skeletal characters: spicule morphology and the arrangement of spicules and/or
spongin fibers into a three-dimensional skeleton (Hooper and Van Soest, 2002). While these
features have been useful for diagnosing many species and higher taxonomic groups (Hooper
and Van Soest, 2002; Morrow and Cárdenas, 2015), molecular phylogenies have revealed that
skeletal traits alone are inadequate due to both insufficient differences between closely related
species and convergently evolving traits in distantly related species (de Paula et al., 2012; Erwin
and Thacker, 2007; Gazave et al., 2010; Morrow et al., 2013; Turner and Pankey, 2023; Xavier et
al., 2010). These limitations are frequently encountered in sponge families with simple spicules
and a loosely organized skeleton, such as the Halichondriidae. This family is of significant
interest due to its pharmaceutical potential (Hirata and Uemura, 1986; Rodriguez Jimenez et al.,
2021; Zhong et al., 2024) and as a model system, as this family includes one of the most studied
sponges in the world, *Halichondria panicea* (Pallas, 1766). This species is found near many
marine labs, is amenable to laboratory culture, and as a result, has been used to ask questions
about sponge physiology, symbiosis, ecology, immunology, and more (Barthel, 1986; Barthel
and Detmer, 1990; Barthel and Wolfrath, 1989; Carrier et al., 2023; Forester, 1979; Knobloch et

al., 2019; Schmittmann et al., 2021; Thomassen and Riisgård, 1995; Witte et al., 1994).
However, this research is built atop a shaky foundation, because the taxonomy and systematics
of *Ha. panicea* and its relatives is very poorly delineated. The World Register of Marine Species
lists over 70 previously named species that are now considered synonyms of *Ha. panicea* (de
Voogd, et al., 2024). Other species that remain putatively valid, like *Ha. bowerbanki* (Burton,
1930), are often difficult to differentiate from *Ha. panicea* and their validity remains in dispute.
Cryptic species within populations ascribed to *Ha. panicea* have remained a possibility, as this
species has supposedly been found in most of the world's oceans and in a wide variety of
climates. Under what can be called a "cryptic endemic species in different regions" hypothesis,
different regions of the world could contain morphologically indistinguishable species that are
endemic to each region (Erpenbeck, 2004). An alternative hypothesis, which can be called the
"single globally-distributed invasive species" hypothesis, is that *Ha. panicea* has been
unintentionally introduced by human activity to many regions of the world, as was recently
shown for the co-familial species *Hymeniacidon perlevis* (Samaai et al., 2022; Turner, 2020).
Below, we test these hypotheses by integrating extensive field collections, genomic
analyses, and traditional morphological characters. For the genomic analyses, we use a
combination of Illumina and Sanger sequencing to assemble mitochondrial genomes and
nuclear ribosomal loci. Because the mitochondrial and nuclear genomes are unlinked,
reciprocal monophyly across these compartments is strong evidence of reproductive isolation
when species are sympatric (Jennings, 1917; Rannala and Yang, 2020). In cases where we find
strong evidence of reproductive isolation, we formally describe these new species. Formally
describing newly discovered species is important because unique identifiers (such as Linnean
names) are crucial for making species visible to the research community and linking data across
databases like GenBank, the World Register of Marine Species, the Global Biodiversity
Information Facility, and museum inventories (Delić et al., 2017). Formally describing new
species requires finding unique characters that differentiate them from previously described
species. When morphological characters cannot be found, DNA states can provide characters to
define and diagnose species (Delić et al., 2017; Eitel et al., 2018; Jörger and Schrödl, 2013;
Lawley et al., 2021; Tessler et al., 2022). The use of DNA sequences to formally describe new
species has a complicated history, with some recent works claiming that the publication of a
consensus DNA "barcode" (a few hundred basepairs of the *cox1* locus) is sufficient to describe a
new species by itself (Sharkey et al., 2021). This latter approach has been heavily criticized, but
these criticisms often explicitly note that the use of DNA characters to describe morphologically
cryptic species is valid when done properly (Ahrens et al., 2021; Meier et al., 2022; Zamani et
al., 2022). For the species described herein, we were therefore careful to avoid the pitfalls of
these earlier works, and instead follow the example of authors (e.g., Eitel et al., 2018; Lawley et
al., 2021) who used DNA-based descriptions that were consistent with the International Code of
Zoological Nomenclature (ICZN, 1999). It remains true that using DNA characters as taxonomic
states has drawbacks: for example, the identification of unknown samples will be limited by
preservation methods, sample age, and the expertise of investigators. However, these
problems would likely be even greater for alternative phenotypic traits such as proteomics or
histology, which would require less ubiquitous expertise and less common preservation
methods. We hope that our approach will lay the groundwork for future research into the
phenotypic differences among these newly described species.

124
Methods
126

Collections

Sponges were collected from the Atlantic Coast of Canada (13 samples, Claire Goodwin), Northern Ireland (10 samples, Christine Morrow and Bernard Picton), Sweden (9 samples, Raquel Pereira), Portugal (2 samples, Raquel Pereira), Panama (6 samples, Robert Thacker) and three locations in the USA: New York (14 samples, Robert Thacker), Washington (19 samples, Thomas Turner, Robert Thacker, and Brooke Weigel), and California (165 samples, Thomas Turner). Collections were most extensive in California, where 95 subtidal sites, 14 intertidal sites, and 23 marinas were surveyed. Subtidal sites were surveyed by divers, with no new collections from deep water (maximum depth 32 m). Previously vouchered material was acquired on loan from the Royal British Columbia Museum (voucher numbers with RBC), the Florida Museum (voucher numbers with UF or BULA), the Smithsonian Institution's National Museum of Natural History (voucher numbers with USNM), the Naturalis Biodiversity Center (voucher numbers with ZMA), and the California Academy of Sciences (voucher numbers with CASIZ). Freshly collected material was deposited in these collections and also the Santa Barbara Museum of Natural history (voucher numbers with SBMNH), the Cheadle Center at the University of California, Santa Barbara (voucher numbers with UCSB), the Atlantic Reference Centre Museum (voucher numbers with ARC), the Ulster Museum (voucher numbers with BELEM), the Yale Peabody Museum (voucher numbers with YPM), and the National History Museum of Los Angeles (voucher numbers with NHMLA). A table listing all examined samples with voucher numbers, GenBank accessions, collection dates and locations, and other metadata is available as supplementary data via Data Dryad (https://doi.org/10.5061/dryad.bvq83bkk6).

Morphology

Spicules were examined after digesting sponge subsamples in bleach. For some samples, a single piece of sponge, including both the ectosome and choanosome, was used for digestion. When sizes of spicules were compared between ectosome and choanosome, these domains were carefully separated with a razor and digested individually. Spicule measurements were taken from images using ImageJ (Schneider et al., 2012), with pixel-to-mm conversion determined using a calibration slide. The length of each spicule was measured as the longest straight line from tip to tip, even if the spicules were curved or bent. Width was measured at the widest point, excluding any adornments like swollen tyles. All spicule measurements are available as supplementary data via Data Dryad.

To analyze the choanosomal skeletons, sponge fragments were first frozen into ice cubes and then hand-cut into tissue sections. These sections were cleared using one of two methods. A few samples were transferred from a 95% ethanol bath to a 100% ethanol bath, and then cleared with histoclear (National Diagnostics). Most samples were digested in a mixture of 97% Nuclei Lysis Solution (Promega; from the Wizard Genomic DNA Purification Kit) and 3% 20 mg/ml Proteinase K (Promega). This method removed cellular material while preserving spicules and the spongin network (if present). For the ectosomal skeletons, the sponge surface was removed with a razor. In many cases, these skeletons could be imaged

directly without further processing; if opaque, they were processed similarly to choanosomal
sections.
To measure ectosomal tract thickness, we averaged the measurements of the three
thickest visible tracts, excluding large matted areas. This process was difficult to standardize
and the measurements are likely less reproducible than spicule size measurements. We also
categorized ectosomal skeletons based on their morphology, noting that there were potentially
distinct types. The five categories used were: 1. Multilayered and gothic-crypt-like; 2. Sieve-like
(a dense mat with approximately round open spaces); 3. Lace-like (a fairly isotropic mesh); 4.
Intermediate forms between categories 3 and 5; 5. A network of spicule tracts (resembling
either straight lines or meandering lines, similar to dry or wet spaghetti). In some cases,
ectosomal skeletons were found as dense mats of spicules without open spaces, but in these
cases, other areas of the same sponge exhibited one of the above types.
179
Illumina sequencing
DNA was extracted with several different kits including the Wizard Genomic DNA Purification kit
(Promega) and the Qiagen Blood & Tissue kit. Prior to Illumina sequencing, samples were
treated with RNase A and repurified with the DNA Clean and Concentrator (Zymo). Illumina
sequencing and library preparation were performed at the DNA Technologies Core Facility at
the University of California, Davis. Libraries were made with the Super-High-Throughput (SHT)
technique, which involved dual indexing each sample and pooling 96 samples before
sequencing. Paired-end 150 bp reads were generated using the Illumina NovaSeq instrument,
resulting in a median of 2.1 million reads per sample. The number of reads varied significantly
across samples, ranging from a maximum of 17 million reads to a minimum of 208 reads. The
reasons for this variability were not explored in detail, but DNA extractions appeared to be of
variable quality even before library construction and pooling, perhaps because of sample
preservation differences or the secondary chemistry of samples.
Reads were trimmed with fastP (Chen et al., 2018) and then de novo assembled into
contigs with Megahit (Li et al., 2015). The number of contigs per sample ranged from 0 to
1,080; the median number of contigs for samples with at least one contig was 22. Blast
searches revealed that many contigs were microbial (or occasionally from other taxa), but the
sponge's nuclear ribosomal locus was usually the highest coverage contig. Indeed, in samples
where only one or two contigs assembled, the ribosomal locus was usually one of them. In
samples where the full ribosomal locus assembled de novo, this contig was then used as a
reference sequence, and reads from that sample were aligned to it with bwa-mem (Li and
Durbin, 2009). SAMtools (Danecek et al., 2021) was then used to convert SAM files from this
alignment into a consensus sequence, which served as the final sequence for that sample. For
samples where no ribosomal contig assembled at the megahit step, the most closely related
sequence available (either from another sample in our analysis or from GenBank) was used as
reference sequence, and bwa-mem was used to align that sample's reads to this reference. In
some cases, ribosomal contigs assembled with megahit were incomplete or fragmented, so
these partial contigs were spliced into the nearest available reference sequence to create a
hybrid reference for bwa-mem alignment. This approach successfully yielded nearly complete
sequences of the 18S, 28S, and intergenic transcribed spacer regions for most samples, though
some had a significant proportion of uncalled bases ("N"). Many of these samples had been

previously sequenced at the 28S locus with Sanger sequencing, and these sequences were
combined with Illumina sequencing when available. Combining Sanger and Illumina data often
resulted in resolving some uncalled bases but in no cases were Illumina consensus sequences
found to have called bases that conflicted with Sanger data, suggesting error rates of called
bases are low. Ribosomal sequences were assembled for 110 samples, with a median number
of called bases (excluding Ns) of 6,429 bp (ranging from 2,227 to 6,589 bp). The number of
called bases for each sample is listed in the supplementary table available from Data Dryad.
Mitochondrial contigs were present in some of the Megahit assemblies but were
generally less complete than ribosomal loci. To generate more complete mitochondrial
genomes, we aligned reads to existing reference sequences using bwa-mem. The references
used included *Halichondria okadai* (MG267395), *Halichondria galea* sp. nov. (KX244759),
Halichondria dokdoensis (MK105765), *Halichondria akesa* sp. nov. (MH756604), *Halichondria*
panicea (MH756603), and *Hymeniacidon perlevis* (KF192342).
For species lacking closely related reference sequences (e.g., the Bowerbanki species
group, as defined in figure 1, and *Hymeniacidon pierrei* sp. nov.), we created hybrid reference
sequences. This procedure involved replacing parts of the closest available reference genome
with de novo contigs from the megahit assembly. Although this method allowed us to assemble
substantial portions of the mitochondrial genome for samples without good reference
sequences, these assemblies were less complete, often containing more uncalled bases ("Ns")
due to regions being too divergent for accurate short-read alignment. For instance, the average
number of called bases in *Halichondria panicea* mitogenomes was 19,228, while for
Halichondria bowerbanki, it was 16,100. In total, we assembled mitochondrial genomes for 75
samples, with a median of 16,888 called bases. As seen with ribosomal loci, combining these
sequences with Sanger data at cox1 did not find errors in called bases. The number of called
bases for each sample is listed in the supplementary table available from Data Dryad.
A sequence of the cox1 locus from the *Ha. panicea* neotype was generated using
sequence capture techniques (Agne et al., 2022) and was generously shared with us by Dirk
Erpenbeck.
239
Sanger sequencing
The cox1 region commonly sequenced in sponges is the Folmer region, a 709 bp segment
amplified using the primers LCO1490 and HCO2198 (Folmer et al., 1994). We used these
primers for a few samples when other primers failed to amplify, though they are less preferred
due to their shorter length and their tendency to co-amplify microbial sequences. We also
attempted to amplify DNA using the primers LCO1490 and cox1-R1, and were successful with
some samples (Rot et al., 2006). These primers target a larger 1264 bp region that includes both
the Folmer region and the "co1-ext" region, which has been useful for distinguishing sponge
species (Erpenbeck et al., 2006). However, we found that *Halichondria loma* and the
Bowerbanki species group (Figure 1) had mismatches with the LCO1490 primer, and
amplifications usually failed with both primer sets. To improve amplification success, we
designed new primers, halcox1-L and halcox1-R, which were effective for all tested *Halichondria*
species. These primers amplify an 886 bp region that encompasses most of the Folmer region
and part of the co1-ext region. We used these new primers for all *Halichondria* samples not

already processed with Illumina or other primer sets. Primer sequences are provided in Table
S2.
We sequenced two spans of the 28S nuclear locus. The D1-D2 region, which spans 940
bp and is highly variable, was amplified in all samples using primers 15F and 878R (Morrow et
al., 2012). We also amplified the adjacent D3-D5 region in some samples using primers 830F
and 1520R. This primer pair amplifies a 710 bp fragment that overlaps the D1-D2 fragment by
20 bp (Morrow et al., 2012).
The ND1 locus was amplified in selected taxa to verify that fixed differences detected in
Illumina sequencing were fixed in a larger sample. Primers were newly designed, but universal
primers for all target species were not found. For the *Ha. dokdoensis*, *Ha. actities*, the Panicea
species group (Figure 1), and Hygeia species group except for *Ha. columbiana* sp. nov., we used
primers nd1-pan-R and nd1-pan-L. For *Ha. columbiana* sp. nov., nd1-pan-R was used with nd1-
hygc-L. For the Bowerbanki species group, nd1-bow-R and nd1-bow-L were used. For *Ha. loma*,
nd1-loma-R was used with nd1-bow-L.
PCR was performed using a Biorad thermocycler (T100) using the following conditions:
95°C for 3 min, followed by 35 cycles of 94°C for 30 sec, [annealing temperature] for 30 sec,
72°C for [extension time], followed by 72°C for 5 minutes. The annealing temperatures and
extension times for each primer pair were as follows: halcox1 (54°C annealing, 60 second
extension), LCO & R1 primers (50°C annealing, 75 second extension), LCO & HCO primers (52°C
annealing, 60 second extension), 28S region (53°C annealing, 60 second extension for both
primer sets), ND1 locus (54°C annealing, 60 second extension, all primer sets). PCR was
performed in 50 µl reactions using the following recipe: 24 µl nuclease-free water, 10 µl 5x PCR
buffer (Gotaq flexi, Promega), 8 µl MgCl, 1 µl 10 mM dNTPs (Promega), 2.5 µl of each primer at
10 µM, 0.75 bovine serum albumin (10 mg/ml, final concentration 0.15 mg/ml), 0.25 µl Taq
(Gotaq flexi, Promega), 1 µl template. Post-PCR, samples were cleaned using ExoSAP-IT (Applied
Biosystems) and then sequenced by Functional Biosciences with Big Dye v3.1 on ABI 3730xl
instruments. We verified that the resulting sequences were of sponge origin using BLAST. All
sequences have been deposited in GenBank under the accession numbers listed in the
supplementary table available from Data Dryad.
283
Phylogenies
Sequences were aligned using MAFFT v.7 (Katoh et al., 2017), and the alignment files are
available as supplementary data. Phylogenies were estimated with maximum likelihood in IQ-
Tree (Nguyen et al., 2015; Trifinopoulos et al., 2016). The ultrafast bootstrap and Shimodaira-
Hasegawa approximate likelihood ratio test (Sh-aLrT) were used to measure node confidence
(Hoang et al., 2018). The optimal model for each tree was selected using ModelFinder
(Kalyaanamoorthy et al., 2017) based on the Bayesian Information Criterion. Mitochondrial
phylogenies were estimated from the concatenation of all known mitochondrial genes; when
they were combined with ribosomal loci (Figure 1), they were analyzed as a separate partition.
The GTR+F+I+G4 model was selected for all ribosomal trees/partitions, while mitochondrial
models differed: GTR+F+I+G4 for all concatenated mitochondrial genes, TN+F+I+G4 for cox1
only, and TIM2+F+G4 for nd1 only. Figures were produced by exporting IQ-Tree files to the
Interactive Tree of Life webserver (Letunic and Bork, 2019).

Single-locus phylogenies of cox1, nd1, and 28S were made with a combination of Sanger
sequenced loci and Illumina sequences that were trimmed to a single locus. We used the NCBI
taxonomy browser and blast to assemble all available sequences related to the Halichondriidae,
along with appropriate outgroups. For cox1, sequences were excluded if they did not include
most of the Folmer barcoding region. For 28S, sequences in the D1 phylogeny were excluded if
they did not include the highly variable C2-D2 barcoding region. We named four clades of the
phylogeny, to facilitate discussions of component taxa, as shown on figures 1 and 3. Alignment
files are available as supplementary files in Data Dryad.
305
306   **Results and Discussion**
307
308   *Molecular phylogenies*
309
Our phylogeny for the 75 samples with the most comprehensive data includes both the nuclear
ribosomal locus (averaging 6,445 bp) and the coding regions of the mitochondrial genome
(averaging 17,153 bp; Figure 1). Previously, comparable data was available for only one species,
Ha. okadai, which is included in the analysis. We successfully assembled nearly complete
nuclear ribosomal loci for an additional 28 samples, including several additional species.
However, we did not have sufficient reads to assemble the mitochondrial genome for these
samples. Mitochondrial and ribosomal phylogenies are shown separately for the Bowerbanki
group (Figure 2). For the remaining specimens, we constructed a ribosomal-only phylogeny
(Figure 3).
Additionally, we constructed phylogenies using Sanger sequencing data from two
mitochondrial loci (cox1 and nd1) and one nuclear locus (28S). These phylogenies were
instrumental in validating genetic differences identified in the Illumina data and allowed us to
include previously sequenced vouchers. In total, we sequenced at least one locus from 232
samples, and these phylogenies are included in the supplementary materials as (Figures S1-S3).
We consistently found highly differentiated clades across all phylogenies. Our first task
was to determine which clade corresponded to the type species *Ha. panicea.* Due to the
ambiguity of the original description, a neotype for the species was previously erected
(Erpenbeck and van Soest, 2002). The neotype was collected from Burnham-on-Crouch, Essex,
England, near the mouth of the Thames River, which is postulated to be near the original type
location. Using sequence capture, Erpenbeck and colleagues generated 207 bp of sequence
data from the hypervariable C2 region of the 28S barcoding locus from the neotype, and
generously shared this previously unpublished sequence with us. Our 28S phylogeny
confidently places this sequence within the clade designated *Ha. panicea* (Figure 1). It is an
exact match to all three of the other samples from the United Kingdom, as well as samples from
both the Atlantic and Pacific coasts of North America. These perfectly-matched samples were
all previously identified as *Ha. panicea* based on their morphology, range, and habitat, and all
fall into the *Ha. panicea* clade (Figure 1), supporting the genetic determination that this clade
corresponds to the name *Ha. panicea*.
338
339
340

341
342 Figure 1. Maximum likelihood phylogeny for samples with assembled mitogenomes and nearly complete
343 ribosomal loci. Asterisks at nodes indicate a UFbootstrap ≥ 99%. Scale bar indicates substitutions per
344 site.
345
Similarly, we used a previously sequenced sample (Ulster Museum voucher
BELUM:Mc4003; GenBank vouchers HQ379241, HQ379316, HQ379382, and KC902247) to
determine which clade corresponds to the name *Ha. bowerbanki*. This name replaces *Ha.*

coalita of Bowerbank, which was preoccupied by *Ha. coalita* (Müller, 1776). The type material
consists only of microscope slides of isolated spicules. Like *Ha. panicea*, the type location is near
the mouth of the Thames. However, two of the slides Bowerbank originally examined were
from sponges collected at Strangford Lough, Northern Ireland. This location matches that of the
Ulster Museum sample, which also aligns with the growth form described by Johnson (1842)
and Bowerbank (1866). We are therefore confident that this clade corresponds to the name *Ha.*
bowerbanki.
356
357 *Inference of reproductive isolation: Bowerbanki group*
358
Our next task was to determine which clades in our phylogeny correspond to species other than
Ha. panicea and *Ha. bowerbanki*. To infer reproductive isolation among species with
overlapping geographic ranges, we compared the phylogenies of multiple loci that assort
independently during meiosis. For example, the mitochondrial and nuclear ribosomal
phylogenies for the clade we named the Bowerbanki group display reciprocal monophyly
(Figure 2). Reciprocal monophyly in both genomic compartments allows us to infer a minimum
of four species in this group: *Ha. bowerbanki* and three new species — *Ha. urca* sp. nov., *Ha.*
zabra sp. nov., and *Ha. pinaza* sp. nov. (new species descriptions are provided in the
systematics section below). Among these species, *Ha. zabra* sp. nov. and *Ha. pinaza* sp. nov. are
the closest relatives. They are well-separated at the ribosomal locus, but Sanger sequencing of
the *cox1* mitochondrial locus did not reveal fixed genetic differences between them (Figure S1).
However, in the concatenated alignment of all mitochondrial coding regions, we identified 24
fixed differences between these clades, including one fixed difference in the cox1 gene outside
the region bounded by our Sanger sequencing primers. With genetic divergence (D_{xy}) of only
0.25% across the mitochondrial genes, the branch separating these clades is short and difficult
to visualize (Figure 2), but monophyly of each clade is well supported (99% and 100% bootstrap
support for *Ha. pinaza* sp. nov. and *Ha. zabra* sp. nov., respectively).
To confirm that these genetic differences are consistent in a larger sample, we
sequenced a region of the nd1 mitochondrial gene containing three apparent fixed differences.
In a larger sample of 16 *Ha. zabra* sp. nov. and 7 *Ha. pinaza* sp. nov. samples, these differences
remained fixed. Similarly, we sequenced the most variable region of the 28S nuclear ribosomal
locus in 23 *Ha. zabra* sp. nov. and 7 *Ha. pinaza* sp. nov. samples, confirming numerous fixed
differences (divergence = 1.32%). This co-assortment of genetic variation in the mitochondrial
and nuclear genomes is strong evidence of reproductive isolation due to the co-occurrence of
these species in California. They are sympatric even at a micro-geographic scale, with samples
collected side-by-side on the same floating dock in Tomales Bay in Northern California, and on
floating docks in Santa Barbara Harbor, over 500 km to the south. Likewise, *Ha. urca* sp. nov.
and *Ha. bowerbanki* were sympatric at a micro-geographic scale. All four species in this species
group were collected simultaneously from floating docks in Santa Barbara Harbor, with varying
degrees of sympatry across other harbors and marinas throughout the state. There were
numerous alleles fixed within the *Ha. urca* sp. nov. and *Ha. bowerbanki* clades in both genetic
compartments, with monophyly highly supported in both cases (Figure 2).
391

392
393 Figure 2. Maximum likelihood phylogenies for the mitogenome (left) and ribosomal locus (right) for the
394 Bowerbanki group. Asterisks at nodes indicate a UFbootstrap ≥ 99%. Scale bar indicates substitutions
395 per site.
396

Our inference of four species within the Bowerbanki group is likely conservative, as
phylogenetic analyses suggest at least one additional species. Within the clade we designate as
Ha. urca sp. nov., one sample from Ireland is genetically distinct. Unlike the *Ha. pinaza* sp. nov.
samples, where Irish and California specimens are nearly identical at the 28S locus (divergence
= 0.2%), the Irish *Ha. urca* sp. nov. shows a 4.6% sequence divergence at this locus. A sample of
Ha. urca sp. nov. from New York is more similar to the California samples than to the Irish one,
indicating that there might be a single population spanning both ocean basins, with the Irish
sample potentially representing a different species or a highly divergent population. Additional
Ha. urca sp. nov. data from the Atlantic includes three sponges from Virginia (SERC/MarineGeo
2018 Lower Chesapeake BioBlitz, available in GenBank) and one from New York (Pankey et al.,
2022). These samples, previously identified as *Ha. bowerbanki*, are recognized as *H. urca* sp.
nov. in our cox1 phylogeny (Figure S1). Combining these specimens with our samples results in
a total of 5 Atlantic and 16 Pacific *Ha. urca* sp. nov. at this locus, with a genetic divergence of
0.38% between them. Although this divergence is slight, it is eight times greater than the
divergence observed between *Ha. zabra* sp. nov. and *Ha. pinaza* sp. nov., which we can
confidently say are reproductively isolated in sympatry. The differentiation seen in cox1 for
Atlantic versus Pacific *Ha. urca* sp. nov. is higher than for any other species found in both basins
(Fst = 0.64 for *Ha. urca* sp. nov., compared to 0.00–0.58 for the six other species with
distributions across both oceans). Some or all Atlantic populations of *Ha. urca* sp. nov. may
therefore be from an additional species, but we conservatively assign all Atlantic samples to *Ha.*
urca sp. nov. until additional data are collected.

We also observed considerable variation in the mitochondrial genome within the clade
we assign to *Ha. bowerbanki* —far more variation than seen within any of the other species in
this study (Figure 2). In contrast to the clades we designate as species, however, the
mitochondrial clades within *Ha. bowerbanki* were not mirrored by divergence in the nuclear
locus. This lack of divergence could occur because there are multiple species within this clade,
but they are too recently diverged to have accumulated fixed differences in the ribosome
(Rittmeyer and Austin, 2012). Alternatively, this species may have a larger effective population
size, higher mitochondrial mutation rate, or balancing selection that has maintained
mitochondrial variation within a single species (Ballard and Whitlock, 2004; Galtier et al., 2009).
We therefore assign all members of this clade to *Ha. bowerbanki*.
Finally, we note that there is disagreement between the mitochondrial and nuclear
compartments regarding the branching order of the species in this group. Mitochondrial genes
place *Ha. bowerbanki* as sister to the clade containing *Ha. zabra* sp. nov. and *Ha. pinaza* sp.
nov., while the nuclear data position *Ha. bowerbanki* as sister to *Ha. urca* sp. nov.. This
discrepancy could result from mitochondrial introgression, incomplete lineage sorting, or
homoplasy. Although this discordance affects the inference of historical relationships, it does
not impact conclusions about current reproductive isolation and is the only instance of differing
branching orders between mitochondrial and nuclear genes in our Illumina-scale dataset.
436
Inference of reproductive isolation: Panicea group
Halichondria panicea was described from the North Atlantic, and there has been debate about
whether North Pacific populations belong to the same species (Erpenbeck, 2004; Erpenbeck
and van Soest, 2002). Our analysis reveals that Pacific and Atlantic populations of *Ha. panicea*
are sister clades distinguishable at the DNA level (Figures 1 & 3). Fixed genetic differences
between these ocean basins are evident in both nuclear and mitochondrial genomes, including
two fixed differences in each of the three Sanger-sequenced loci (cox1, nd1, and 28S). The
genetic divergence between Atlantic and Pacific populations of *Ha. panicea* at cox1 is similar to
that observed between Atlantic and Pacific populations of *Ha. urca* sp. nov. (0.40% divergence
for both; Fst = 0.58 for *Ha. panicea*, Fst = 0.64 for *Ha. urca* sp. nov.). Unlike cases where
populations are sympatric, this divergence does not confirm that these populations are
reproductively isolated (Rannala and Yang, 2020). Indeed, *Ha. panicea* have been reported from
the Arctic. No genetic data are available to confirm the identity of Arctic *Halichondria*, but if
these populations are indeed *Ha. panicea*, it is possible that populations from the Atlantic and
Pacific meet and mix in these Northern waters. We therefore leave the North Pacific population
of *Ha. panicea* within this species, pending further data. We note that a previous paper that
attempted to compare *Ha. panicea* in the Atlantic and Pacific only used two samples from the
Atlantic, and alignment of those samples to our data reveal that the Atlantic samples were, in
fact, *Ha. bowerbanki* (Erpenbeck, 2004).
As noted in the morphological section, there are multiple cryptic species that are
morphologically indistinguishable from *Ha. panicea*. It is therefore surprising that, of the three
species most closely related to *Ha. panicea*, two are morphologically distinct. The most closely
related species is *Ha. sitiens* (Schmidt, 1870), which, like *Ha. panicea*, is found in both the North
Pacific and North Atlantic. *Ha. sitiens* is clearly differentiated from *Ha. panicea* by spicule size
and the presence of surface papillae. These papillae have historically been used to define

subgenera within *Halichondria*, with *Ha. sitiens* as the type species of subgenus *Eumastia* and
Ha. panicea as the type species of subgenus *Halichondria* (Erpenbeck and van Soest, 2002). The
close relationship between these two species clearly invalidates these subgenera. Although
only one *Ha. sitiens* was sequenced using Illumina technology, two Pacific and four Atlantic
samples were sequenced at each of the three Sanger loci. In contrast to *Ha. panicea*, all *Ha.*
sitiens samples are nearly identical, showing minimal genetic diversity both within and between
populations (divergence between ocean basins <0.1% at each locus, with no fixed differences).
Outside of the *Ha. panicea* + *Ha. sitiens* clade were two other species; short branches
did not allow us to resolve their branching order. One of these is represented by a single sample
that is morphologically indistinguishable from *Ha. panicea.* This sample (ESS07) was collected at
the same time and place as three other samples that were found to be *Ha. panicea* (Brig's
Shoal, in the Eastern Shore Islands of Nova Scotia). Though this clade is represented by only a
single sample in our collections, the mitochondrial genes are nearly identical (99.9%) to a
previously published mitochondrial genome from Iceland (GenBank accession MH756604).
Previous work ascribed this genotype to *Ha. panicea* (Knobloch et al., 2019), but the closer
relationship of *Ha. sitiens* to true *Ha. panicea* clearly indicates that a new name for this taxon is
needed. We describe it here as *Ha. akesa* sp. nov. This species differs from the *Ha. panicea*
neotype DNA fragment at 3 sites (of 207); divergence vs. all *Ha. panicea samples* across the
Sanger regions of cox1 and 28S is 0.50% and 0.80%, respectively.
The final species in the Panicea group serves to emphasize the inadequacy of spicule-
based taxonomy in this family. *Topsentia fernaldi* (Sim and Bakus, 1986) has a very dense
skeleton of long oxeas that make it quite distinct from *Ha. panicea*. The genus *Topsentia* is
among the most polyphyletic of sponge genera, with previous molecular phylogenies placing
some *Topsentia* species very far from other members of the Halichondriidae, in a sister clade to
the order Axinellida (Pankey et al., 2022; Redmond et al., 2013). Reconciling the taxonomy and
phylogeny of this genus (and the other polyphyletic genera included here) will require data
from additional species, especially type species, and is beyond the scope of this paper.
489
Inference of reproductive isolation: Hygeia group
The most widely distributed species in the clade we dub the Hygeia group is the newly
described *Ha. hygeia* sp. nov., which was found throughout the Northeast Pacific and in the
Northwest Atlantic. In the Pacific, its range overlaps broadly with another new species, *Ha.*
californiana sp. nov. These two species are distinctly separated at the ribosomal locus, showing
14 fixed differences (1.5% divergence) in the Sanger-sequenced region of 28S (Figure S2).
Although they could not be differentiated in the Sanger-sequenced region of the cox1
mitochondrial locus (Figure S1), Illumina sequencing revealed numerous other fixed differences
in the mitochondrial genome. Sanger sequencing of the nd1 mitochondrial gene (Figure S3)
confirmed 4 fixed differences in a larger sample (0.8% divergence, Fst = 0.73, N = 12 *Ha.*
californiana sp. nov. vs. 17 *Ha. hygeia* sp. nov.). The fixed differences observed in both genomic
partitions, coupled with their broad sympatry, strongly supports reproductive isolation in
nature.
Two additional species are found in this group, both restricted to the Pacific Northwest
of North America. Despite the limited number of samples (N=2 for each species), both are well
differentiated from other species in the group based on mitochondrial and nuclear loci. Two *Ha.*

cascadia sp. nov. samples were sequenced using Illumina technology, and these samples are
strongly supported as a sister clade to all other species in the group (Figure 1). Given that *Ha.*
hygeia sp. nov. and *Ha. californiana* sp. nov. are well-supported as isolated species, and *Ha.*
cascadia sp. nov. is sympatric with *Ha. hygeia* sp. nov., it is likely that *Ha. cascadia* sp. nov. is
also an isolated lineage deserving of species status. We also have two samples of *Ha.*
columbiana sp. nov., both from British Columbia, though only one was successfully sequenced
with Illumina technology. This sample was quite unique, with many alleles differentiating it
from the sympatric species *Ha. hygeia* sp. nov.. This sample formed a clade with the second
sample across all three Sanger-sequenced loci, providing further evidence that this distinct
clade is likely a separate species. Our inference of four species within the Hygeia species group
is likely conservative, as phylogenetic analyses suggest at least one additional species. Three
samples (TLT1076, TLT742, TLT645) formed a reciprocally monophyletic clade at the nuclear
ribosomal locus and the mitochondrial nd1 locus (Figures S2 and S3). Given the limited sample
sizes and the close relationship of these samples with other *Ha. hygeia* sp. nov., we
conservatively include these samples within the species concept for *Ha. hygeia* sp. nov.,
pending further data.
522
Additional Halichondria species found to span the temperate North Pacific
The first mitochondrial genome published for a *Halichondria* species was for a "*Halichondria*
sp." sponge sampled from Fujian Province, China (Wang et al., 2016). Sequencing of
mitochondrial genes later documented this same genotype from the intertidal zone of mainland
South Korea and nearby islands, and from Mission Bay, San Diego, California (Park et al., 2007).
This sponge has remained undescribed, likely due to it being morphologically indistinguishable
from *Ha. panicea*. Our Southern California samples have mitochondrial genomes that are nearly
identical with the published sequence from China (99.95% identical), confirming that we have
recollected this same species from California. Consistent with previously published data, it is
only distantly related to *Ha. panicea*, despite their morphological similarity. The clade we
designate here as the Okadai species group contains *Ha. okadai* and *Ha. galea* sp. nov., which
are sympatric in the Korean Strait; the Caribbean species *Ha. magniconulosa* and *Ha.*
melanodocia; and samples we collected from Pacific Panama that are likely from an
undescribed species (Figure 3).
Remarkably, an additional species of *Halichondria* known from Asia was also discovered
in California for the first time: *Ha. dokdoensis* Kang, Kim & Kim, 2022. A mitochondrial genome
sequence was previously published for this species (Kim et al., 2019), and this sequence is
nearly identical to the mitochondrial genome of our California sample: they differ by only 12 of
13,033 aligned base pairs (<0.1%). These samples are more closely related to *Hymeniacidon*
actites than to any other sequenced *Halichondria*, and together these two species form a clade
that is the sister to the combined Panicea and Hygeia species groups. We discovered a single
sample of this species in California, on an oil extraction platform off the coast of Santa Barbara.
As discussed in the systematics section, a publicly available sequence of the cox1 locus indicates
that the species is also likely found in the Netherlands. These discoveries serve to highlight that
there are likely additional species of widespread, introduced *Halichondria* that will be
discovered by additional genotyping.
549

Suberites lambei TLT296 California
Pseudosuberites latke TLT538 California
Pseudosuberites latke TLT1131 California

Okadai group
Halichondria sp. P19x63 Pacific Panama
Halichondria sp. P19x40 Pacific Panama
Halichondria galea sp. nov. TLT743 California
Halichondria galea sp. nov. TLT495 California
Halichondria okadai AB511881
Halichondria melanadocia P19x163 Caribbean Panama
Halichondria magniconulosa P19x171 Caribbean Panama

Hymeniacidon perlevis NIB0611 Ireland
Hymeniacidon perlevis NIB0616 Ireland
Hymeniacidon perlevis NIB0615 Ireland
Hymeniacidon perlevis TLT651 California
Hymeniacidon perlevis TLT1265 California
Hymeniacidon perlevis TLT736 California
Hymeniacidon perlevis TLT247 California
Hymeniacidon perlevis TLT658 California
Hymeniacidon perlevis TLT1074 California

Hymeniacidon pierrei sp. nov. TLT599 California
Hymeniacidon kitchingi TLT820 California
Semisuberites cribosa BHAK-09371 British Columbia
Hymeniacidon fusiformis TLT1194 California
Hymeniacidon sp. Quoddy115 Nova Scotia

Halichondria dokdoensis TLT386 California
Hymeniacidon actites TLT847 California
Hymeniacidon actites TLT227 California
Hymeniacidon actites TLT856 California
Hymeniacidon actites TLT454 California
Hymeniacidon actites TLT1158 California

Panicea group
Halichondria akesa sp. nov. ESS07 Nova Scotia
Topsentia fernaldi TLT1134 California
Halichondria sitiens ESS44 Nova Scotia
Halichondria sitiens ESS130 Nova Scotia
Halichondria panicea NIB0608 Ireland
Halichondria panicea 190424 Sweeden
Halichondria panicea 180923 Sweeden
Halichondria panicea 160504 Sweeden
Halichondria panicea Quoddy124 New Brunswick
Halichondria panicea 191026 Sweeden
Halichondria panicea Quoddy113 New Brunswick
Halichondria panicea ESS11 Nova Scotia
Halichondria panicea FH12x81 Washington
Halichondria panicea BHAK-00536 British Columbia
Halichondria panicea BHAK-03453 British Columbia
Halichondria panicea TLT858 California
Halichondria panicea TLT619 California
Halichondria panicea FH12x84 Washington
Halichondria panicea FH12x38 Washington
Halichondria panicea FH12x39 Washington

Hygeia group
Halichondria cascadia sp. nov. FH12x93 Washington
Halichondria cascadia sp. nov. FH12x92 Washington
Halichondria californiana sp. nov. TLT417 California
Halichondria californiana sp. nov. TLT1093 California
Halichondria californiana sp. nov. TLT937 California
Halichondria californiana sp. nov. TLT860 California
Halichondria californiana sp. nov. TLT160 California
Halichondria columbiana sp. nov. BHAK-09466 British Columbia
Halichondria hygeia sp. nov. TLT742 California
Halichondria hygeia sp. nov. TLT645 California
Halichondria hygeia sp. nov. TLT1076 California
Halichondria hygeia sp. nov. TLT940 California
Halichondria hygeia sp. nov. TLT504 California
Halichondria hygeia sp. nov. F12x108 Washington
Halichondria hygeia sp. nov. BHAK-03597 British Columbia
Halichondria hygeia sp. nov. TLT724 California
Halichondria hygeia sp. nov. WMS21x10 New York
Halichondria hygeia sp. nov. WMS21x9 New York
Halichondria hygeia sp. nov. WMS19x5 New York

Bowerbanki species group

Tree Scale: 0.05

Figure 3 (above). Maximum likelihood phylogeny for samples with nearly complete ribosomal loci. Asterisks at nodes indicate a UFbootstrap ≥ 99%. Scale bar indicates substitutions per site.

Taxa are not diagnosable with traditional morphological characters
Our multi-locus molecular data, combined with broad sympatry, allows us to form strong inferences regarding which clades are species. We can use these well-delineated species to determine if known morphological characters are sufficient for their diagnosis. We first considered only the previously named species *Ha. panicea* and *Ha. bowerbanki*. Previous attempts to separate these species cite several differences including 1) longer spicules in *Ha. bowerbanki*, 2) a difference in spicule length in the choanosome vs. the ectosome in *Ha. panicea* but not *Ha. bowerbanki*, and 3) differences in the arrangement of spicules in the ectosome between species (Ackers et al., 2007; Hartman, 1958). These previous works did not use DNA to diagnose samples, so it is likely that samples from the previously unknown species discovered and described here were included in past work. Nonetheless, as detailed further below, we find that these traits do differ, on average, between these species. However, we also find variability within species that prevents them from being used to reliably diagnose all samples.

   Spicules in *Ha. bowerbanki* were, on average, nearly 40% longer than those in *Ha. panicea* (Wilcoxon p<0.0001, Figure 4). However, substantial overlap of these distributions prevents this character from being used as a species diagnostic (Figure 4). Mean spicule length ranged from 202 to 357 µm for *Ha. panicea* and from 307 to 423 µm for *Ha. bowerbanki*. Both species had high variability in mean spicule length within the Atlantic and Pacific basins. We also find that spicules are, on average, 15% shorter in the ectosome than the choanosome for *Ha. panicea* but not for *Ha. bowerbanki* (Figure S4). Again, this trait proved insufficient for diagnosing all samples due to intraspecific variability. Some *Ha. panicea* specimens had spicules 30% longer in the choanosome, and two samples (one from Nova Scotia and one from Sweden) had spicules of equal length in both compartments. To verify that this variability within species was not entirely due to technical variability and sampling error, we reisolated spicules from resampled tissues. One *Ha. panicea* sample with no initial difference in spicule lengths by compartment and another with a 32% difference were re-examined. Replicate data confirmed these results: the sponge with no initial difference had only a 1% difference in the replicate (non-significant), while the sponge with a 32% difference initially showed a 26% difference in the replicate (p<0.001). In contrast, *Ha. bowerbanki* samples exhibited less variability, with ectosomal spicules varying from about 10% longer to about 10% shorter than choanosomal spicules. Attempts to replicate samples of *Ha. bowerbanki* with the greatest observed differences (11% longer choanosomal spicules vs. 9% longer ectosomal spicules) showed both samples regressing towards a mean difference of zero, consistent with the possibility that variation in *Ha. bowerbanki* is largely due to tissue sampling variation and statistical noise.

   When average spicule length and length differential between compartments are plotted together, the average differences between *Ha. panicea* and *Ha. bowerbanki* are clear, and it appears likely that a combination of these two traits would be sufficient to differentiate most, but probably not all, samples (Figure 5). However, when we added the other species described here to this plot, we observed substantial overlap among them.

595
596
597 Figure 4. Spicule length distributions for samples of *Halichondia panicea* (Hpan) and *Ha. bowerbanki*
598 (Hbow). Sample are sorted by mean within species, and names indicate if the sample was from the
599 Atlantic (Atl) or Pacific (Pac). Hpan samples generally have shorter average lengths and bimodal
600 distributions, due to length differences between the ectosome and choanosome, but some Hpan
601 samples are longer than the shortest Hbow samples, and bimodality is not always present.

602
The most comprehensive publication on the morphological distinction between *Ha.*
panicea and *Ha. bowerbanki* states that the arrangement of spicules in the ectosome is the
most reliable diagnostic character (Hartman, 1958). Typically, *Ha. bowerbanki* exhibits a
network of long, thin spicule tracts that delineate larger open spaces, while *Ha. panicea* has
thicker spicule tracts forming a dense mesh surrounding smaller openings. *Ha. panicea*
sometimes also features "gothic crypt-like supports" connecting the ectosomal and
choanosomal skeletons (Hartman, 1958). Although these differences were observed in our
samples, they proved insufficient for diagnosing all samples reliably. This trait is highly variable
within a single sponge and among samples within species. While mean tract thickness differed
significantly between species (69 µm for *Ha. panicea* vs. 38 µm for *Ha. bowerbanki*, Wilcoxon
p<0.0001; Figure S5), there is considerable overlap in the distributions (e.g., *Ha. panicea* 39–97
µm, *Ha. bowerbanki* 29–56 µm; figure S5). We also attempted to categorize ectosomal
skeletons into one of five structural types (Figure S6); this categorization again highlighted
average differences but also considerable overlap. Principal components analysis combining

ectosomal tract thickness, average spicule length, and length differential between
compartments did not enhance the ability of these traits to diagnose species (Figure S7). Even
when considering a small number of samples of only *Ha. panicea* and *Ha. bowerbanki*,
separation was incomplete.
621

622
623　Figure 5. Spicule measurements across *Halichondria* species. Only samples where spicules were
624　measured separately for the ectosome and the choanosome are shown. Mean spicule length is the
625　mean for both domains combined, and proportional length difference is the choanosomal mean minus
626　the ectosomal mean over the sample mean.
627
Other morphological differences between (apparent) *Ha. panicea* and *Ha. bowerbanki*
have been proposed, such as smell (subjective and only available in fresh material), rigidity of
branches (branches often not present), color, and general form (Ackers et al., 2007; Hartman,
1958). It is likely that these traits and others will be partially informative, but our efforts to
differentiate samples based on these traits has proved inconsistent and frustrating. Moreover,
many morphological traits likely respond plastically to environmental conditions such as light
intensity, wave energy, and current strength (Palumbi, 1986). Some species are found primarily
in sheltered harbors and marinas, while others are found in wave-swept intertidal habitats, so
average morphological differences between them may be due more to habitat than genotype.
For instance, a sample of *Ha. urca* sp. nov. from an intertidal zone exhibited an ectosomal
skeleton typical of *Ha. panicea*, which is common in high-energy environments, whereas most
Ha. urca sp. nov. specimens from sheltered marinas displayed a sparsely spiculated ectosome
(see *Ha. urca* sp. nov. description below). Therefore, we conclude that the previously named

species *Ha. panicea* and *Ha. bowerbanki* cannot be reliably distinguished based on known
morphological traits alone, and the newly described *Halichondria* species should be considered
cryptic species without clear morphological differences.
644
Systematics
The aims of this section are two-fold. First, we formally describe the *Halichondria* species that
were found to be morphologically indistinguishable from *Ha. panicea*. Second, we aim to
comprehensively revise the taxonomy of the family Halichondriidae for the Northeast Pacific,
from British Columbia to California. This revision encompasses all newly discovered and
previously described species within this region. The only species included here that has not
been found in this area is *Ha. akesa* sp. nov., which is known solely from the North Atlantic.
Diagnoses include DNA character states with the format "number:letter", where the
number indicates a position in one of the DNA alignments provided (where the first base is
number 1), and the letter denotes the base state at that position, which is unique compared to
all other species in the genus for which sequence data are available. The notation "number-
number" indicates a sequence of DNA starting at the first number and proceeding to the last
number. The sample sizes (N) for each character are indicated; these can be variable as some
samples were sequenced at more loci than others, or because of the presence of missing data
in the sequences. Although many species of *Halichondria* lack DNA data, these previously
described species were all morphologically distinguished from *Ha. panicea* and are therefore
morphologically distinct from the newly described species as well.
In practice, informal diagnosis of all species is feasible by sequencing the D1-D3 region
of the 28S locus and constructing a phylogeny with the provided sequences. However, clade
membership alone is not sufficient for formal description under the Code of Zoological
Nomenclature, and not all species exhibit unique derived states at this locus. Consequently,
formal diagnosis requires sequencing different loci for some species. Morphological traits
should be employed to diagnose species when comparing them to previously described
Halichondria that have not been sequenced at these loci.
669
670 **List of species discussed in systematics section**
671
672 **Order Suberitida, Family Halichondriidae**
673
Genus Halichondria
Halichondria panicea (Pallas, 1766)
Halichondria akesa sp. nov.
Halichondria sitiens (Schmidt, 1870)
Halichondria hygeia sp. nov.
Halichondria californiana sp. nov.
Halichondria cascadia sp. nov.
Halichondria columbiana sp. nov.
Halichondria dokdoensis Kang, Kim & Kim 2022
Halichondria galea sp. nov.
Halichondria bowerbanki Burton, 1930

Halichondria urca sp. nov.
Halichondria zabra sp. nov.
Halichondria pinaza sp. nov.
Halichondria loma Turner & Lonhart 2023
689
Genus Hymeniacidon
Hymeniacidon perlevis (Montagu, 1814)
Hymeniacidon actites (Ristau, 1978)
Hymeniacidon ungodon de Laubenfels 1932
Hymeniacidon kitchingi (Burton 1935)
Hymeniacidon pierrei sp. nov.
Hymeniacidon fusiformis Turner & Lonhart 2023
Hymeniacidon globularis Ott, McDaniel & Humphrey 2024
698
699   Genus Semisuberities
700   *Semisuberites cribrosa* (Miklucho-Maclay, 1870)
701
702   Genus Topsentia
703   *Topsentia fernaldi* (Sim & Bakus, 1986)
704   *Topsentia disparilis* (Lambe 1894)
705
706   Genus Axinyssa
707   *Axinyssa piloerecta* sp. nov.
708   *Axinyssa tuscara* (Ristau 1978)
709

Family Halichondriidae
The Systema Porifera defines the Halichondriidae as members of order "Halichondrida with a
confused arrangement of smooth oxeas and/or styles in the choanosome and usually an
organized special ectosomal skeleton consisting of tangentially arranged or densely confusedly
arranged crust of oxeas and/or styles of sizes similar to or smaller than those of the
choanosome" (Erpenbeck and Van Soest, 2002). Order Halichondrida was abandoned based on
a lack of synapomorphic characters and its families were distributed into several orders. Family
Halichondriidae joined families Suberitidae and Stylocordylidae in order Suberitida (Morrow
and Cárdenas, 2015).
719

720   **Genus *Halichondria***
721   Halichondriidae with a tangential ectosomal skeleton carried by subectosomal spicule tracts or
722   brushes separated by subdermal spaces. Megascleres exclusively oxeas or derivates in a wide
723   size range (paraphrased from Erpenback and Van Soest, 2002). This broadly defined genus
724   includes over 95 species (de Voogd et al., 2024).
725

726   ***Halichondria panicea* (Pallas, 1766)**
727   Figures 6 & 7
728

Synonyms
Spongia panicea Pallas, 1766
The World Porifera Database lists over 70 additional synonyms (de Voogd, et al., 2024)

Material examined
160504, Spettan, Gullmarsfjorden, Sweden, (58.25770, 58.25770), 18 m, 2015-05-04; 160505, Spettan, Gullmarsfjorden, Sweden, (58.25770, 58.25770), 18 m, 2015-05-04; 180911, Svartejan, Idefjord, Sweden, (59.11020, 11.32190), 30 m, 2018-09-11; 180923, Lunneviken, Idefjord, Sweden, (59.05460, 11.16900), 27 m, 2018-09-23; 181204, Svartejan, Idefjord, Sweden, (59.11020, 11.32190), depth not recorded, 2018-12-04; 190424, Flatholmen, Gullmarsfjorden, Sweden, (58.26080, 11.39820), 14 m, 2019-04-24; 190624, Trindeknubben, Idefjord, Sweden, (58.78290, 10.99620), 12 m, 2019-06-24; 191026, Gamla Svinesundsbron, Idefjord, Sweden, (59.09770, 11.26860), 19 m, 2019-10-26; 191116, Yttre Vattenholmen, Idefjord, Sweden, (58.87540, 11.10560), 27 m, 2019-11-16; BHAK-00536, Fifth Beach, Shatner Point, Calvert Island, British Columbia, (51.63870, -128.15700), intertidal, 2017-07-25; BHAK-03453, Crazy Town, North Beach, Calvert Island, British Columbia, (51.66680, -128.13400), intertidal, 2017-08-08; ESS11, Brig's Shoal, Eastern Shore Islands, Nova Scotia, (44.62781, -62.93603), 25 m , 2021-07-19; ESS17, Brig's Shoal, Eastern Shore Islands, Nova Scotia, (44.62781, -62.93603), 25 m, 2021-07-19; ESS18, Brig's Shoal, Eastern Shore Islands, Nova Scotia, (44.62781, -62.93603), 25 m , 2021-07-19; ESS74, South of Brig's Shoal, Nova Scotia, (44.60336, -62.92724), 29 m, 2021-07-23; ESS80, South of Brig's Shoal, Nova Scotia, (44.60336, -62.92724), 29 m , 2021-07-23; FH12x36 (YPM 111889), Cattle Point, San Juan Island, Washington, (48.45030, -122.96237), intertidal (rocks), 2012-05-19; FH12x38 (YPM 111890), Cattle Point, San Juan Island, Washington, (48.45030, -122.96237), intertidal (rocks), 2012-05-19; FH12x39 (YPM 111891), Cattle Point, San Juan Island, Washington, (48.45030, -122.96237), intertidal (rocks), 2012-05-19; FH12x81 (YPM 111892), Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22; FH12x84 (YPM 111893), Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22; FH12x85 (YPM 111894), Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22; FH12x86 (YPM 111895), Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal (rocks), 2012-05-22; NIB0608, Carrick-a-Rede, Northern Ireland, (55.24238, -6.33396), 10m (growing on kelp stipe), 2021-10-17; Quoddy113, Bald Head, Campobello, New Brunswick, (44.91220, -66.95215), 11.6 m , 2022-10-31; Quoddy124, Marble Island, Off Indian Island, Western Isles, New Brunswick, (44.91921, -66.97054), 10.6 m, 2022-11-01; TAT026, Strawberry Draw Overhang, Tatoosh Island, Washington, (48.39190, -124.73800), intertidal, 2018-06-16; TAT031, Toad Point, Tatoosh Island, Washington, (48.38970, -124.73500), intertidal, 2018-06-17; TAT032, Toad Point, Tatoosh Island, Washington, (48.38970, -124.73500), intertidal, 2018-06-17; TLT1293, Cape Alava, Washington, (48.17134, -124.75227), 4-12 m, 2023-07-29; TLT1370, Cape Johnson, Washington, (47.97327, -124.68220), 4-7 m, 2023-07-27; TLT1371, Cape Johnson, Washington, (47.97327, -124.68220), 4-7 m, 2023-07-27; TLT1430, Neah Bay, Washington, (48.38676, -124.63223), 6-14 m, 2023-08-01; TLT619, Scott's Creek Beach, California, (37.04213, -122.23366), beach wrack, 2020-11-14; TLT858, Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06; ZMA6155, Jenny's Cove, Lundy Island, England, 24 m, 1985-01-08.

774
Figure 6. *Halichondria panicea*. Northwest Atlantic: A (ESS80), B (ESS17), D (ESS18), F (Quoddy124), H
(ESS11). Northeast Pacific: C (TLT1430), E (TLT1293), G (TLT1371). All are photographed in situ; C shows
wound where sample was taken to display internal color.

Diagnosis
Yellow or green *Halichondria* with a tangential ectosomal skeleton developed as a dense sieve-
like mat of spicules or a lace-like network of spicule tracts. Choanosomal skeleton is confused
with vague spicule tracts, sometimes accompanied by well-developed subectosomal spicule
tracts descending from the ectosomal skeleton. Spicules oxeas, ranging from 100–500 μm in
length, with averages per sample of 200–400 μm in length and 5–10 μm in width; usually
slightly shorter in the ectosome than the choanosome. Diagnosis relative to similar species
requires DNA characters, as follows:

COX1. 754:A, 952:A. N=55–69 depending on position. These characters differentiate *Ha.*
panicea from all other sequenced *Halichondria* except *Ha. sitiens,* which is diagnosed relative to
Ha. panicea based on morphological characters such as spicule length and surface papillae.

28S. No unique derived states shared by all *Ha. panicea*, but the Atlantic Ocean population has
two unique states, 605:T and 1549:C. N=13–21 depending on position

ND1. No unique derived states shared by all *Ha. panicea*, but the Pacific Ocean population has a
unique state, 451:T. N=10).

Morphology
Thinly or thickly encrusting, sometimes with digitate tendrils (Figure 6). Oscula usually
prominent, but vary from raised rims nearly flush with surface to tall oscular chimneys. Color
alive varies from pale translucent yellow to golden, often variegated with green, or with the
exterior entirely green or gray-green and the interior yellow. White or beige in ethanol.

Skeleton
Tangential ectosomal skeleton can be sieve-like, with a dense mat of spicules surrounding open
areas that are mostly circular or oval shaped (Figure 7A, 7F), or more lace-like, with a web of
spicule tracts surrounding open areas that tend towards triangular (Figure 7C, 7D). In some
samples, this tangential ectosomal skeleton is supported by prominent spicule tracts that rise
from the choanosome to fan out tangentially in the ectosome (Figure 7B, 7G). The choanosomal
skeleton in these samples becomes more confused towards the interior (Figure 7H). Other
samples have a confused jumble of multispicular tracts, spicule bundles, and individual spicules
both near the ectosome and interior (Figure 7I). No apparent spongin.

Spicules
Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have a typical halichondroid shape: evenly curved or with a slight central bend, thickest
in the center and tapering gently to sharp points at both ends. Spicules have a large range in
lengths in both the ectosome and the choanosome, but average 15.4% longer in the
choanosome. Several samples (ESS11, 190424) had no difference in length by compartment
(Figure S4). Dimensions are given for individual samples in table 1. When all samples are
combined, spicules measure 114–272–499 x 2–8–16 µm, n=1502. Average values varied
considerably across samples, with mean lengths from 202 to 358 µm, and mean widths from 5
to 10 µm.

Distribution and habitat
This species has been reported to occur from Iceland to the Mediterranean in the Northeast
Atlantic, from Northern Canada to New England in the Northwest Atlantic, and possibly the
Arctic and North Pacific (Erpenbeck and van Soest, 2002). DNA data is able to confirm only the
Northern portion of this range in the Northeast Atlantic, including Iceland, Sweden, the United
Kingdom, and the Netherlands. Genetic support in the Northwest Atlantic ranges from New
Brunswick to New Hampshire. Sequenced samples from farther South in the Atlantic (Portugal,
Connecticut, New York, Virginia) have thus far proved to be in the Hygeia or Bowerbanki species
groups, but additional sampling could certainly extend this range. In the North Pacific, DNA-
confirmed range is from the Bering Strait to Central California, with no DNA confirmation
available from the Northwest Pacific. The species is rare or absent from shallow waters in
Southern California, as the genotyping of over 100 *Halichondria* samples from this region found
none to be *Ha. panicea*.

Previously, this species was described as common on rocks and brown algae in the
intertidal and shallow subtidal zones, preferring exposed areas over sheltered bays, and
extending down to a depth of 500 meters (Erpenbeck and van Soest, 2002). DNA-confirmed
samples align with this description, although we cannot confirm a depth range beyond 29
meters. The species is common in exposed intertidal areas and the shallow subtidal zone but is
rare or absent on human-made structures like floating docks in marinas, where *Ha. hygeia* sp.
nov. and members of the Bowerbanki species group are prevalent.
845
846 Table 1. Spicule lengths of *Halichondria panicea*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
BHAK-00536	114–185–285 n=40	130–217–322 n=46
BHAK-03453	153–204–256 n=40	170–241–322 n=40
TLT619	130–270–451 n=40	199–334–442 n=40
TLT1370	172–286–451 n=45	146–364–499 n=50
TLT1430	147–233–353 n=96	162–299–367 n=98
FH12x38	155–247–335 n=49	169–272–377 n=50
FH12x39	134–247–357 n=42	169–291–367 n=47
FH12x81	145–238–351 n=55	166–262–354 n=54
Atlantic Ocean		
NIB0608	176–257–322 n=49	187–281–365 n=51
ZMA-6155	149–283–408 n=40	241–354–458 n=40
190424	178–273–347 n=40	196–267–342 n=40
180911	173–247–349 n=49	221–296–357 n=41
Quoddy124	212–320–421 n=48	298–397–463 n=44
ESS11	185–254–318 n=86	167–254–331 n=86

847
Remarks
As discussed in the genetic section, *Ha. panicea* from the Atlantic and Pacific basins form sister
clades that are distinguishable at the genetic level and could potentially be considered separate
species. However, if *Ha. panicea* occurs throughout the Arctic, it may link the populations
across these basins. We conservatively retain both populations under the same species concept
pending further data.
We lack genetic confirmation of *Ha. panicea* in the Northwest Pacific, but its presence
there seems likely, as we have confirmed its distribution in the Aleutian Islands and the Bering
Strait. This species has also been reported from various other global regions, including South
America, New Zealand, and the tropical South Pacific (gbif.org). Several lines of evidence
suggest that these records may actually pertain to other species, such as *Ha. hygeia* sp. nov.,
members of the Bowerbanki species group, or additional cryptic species that remain to be
discovered. First, *Ha. panicea* seems less likely than those species to be experiencing a human-
assisted range expansion for two reasons: 1) it does not seem to occur in human-modified
habitats like marinas, and 2) genetic differentiation between Atlantic and Pacific populations
make it likely that these populations do not share a common ancestor in the modern era.

Second, the species does not seem to tolerate warm temperate waters in Southern California,
and is therefore unlikely to occur in shallow tropical waters.
866

867
Figure 7. *Halichondria panicea* skeleton and spicules. Ectosomal skeletons: A (ESS80), B (TLT1430), C
(FH12x38), D (Quoddy124), E (NIB608), F (TLT619). B is imaged from below to show the twisting columns
of spicules connecting the ectosome to the choanosome in this sample. G: cross section at sponge
surface, showing spicule columns and subectosomal spaces (TLT1430), H: cross section of same sample
(TLT1430) farther towards the inferior, showing confused skeleton, I: cross section at sponge surface
without large subectosomal spaces (TLT1370), J: oxeas (TLT1293).

874
875 ***Halichondria akesa* sp. nov.**
876 Figure 8

877
878 **Material examined**
879 Holotype: ESS07 (ARC 82019, CASIZ 245200), Brig's Shoal, Eastern Shore Islands, (44.62781, -
880 62.93603), 25 m, 2021-07-19.

881
882 **Etymology**
883 Name inspired by the Greek goddess Akeso, sister of Panakeia.

Diagnosis

Yellow and green *Halichondria* with a tangential ectosomal skeleton developed as a thick mat of spicules and/or a lace-like network of spicules and spicule tracts. Choanosomal skeleton consists of prominent spicule tracts near the ectosome, transitioning to paucispicular tracts and spicules in confusion deeper in the choanosome. Spicules oxeas, ranging from 90–460 µm in length and 2–10 µm in width; slightly shorter in the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 621:T. N=1.

ND1. 39:T, 55:T. N=2.

Morphology

The single specimen examined was thickly encrusting with prominent oscula. Yellow alive, white in ethanol.

Figure 8. *Halichondria akesa*. A: field photos, B: tangential section showing ectosomal skeleton and subectosomal spicule tracts, C: oxeas. All photos from the holotype.

Skeleton

The single specimen examined has a tangential ectosomal skeleton which was a dense mat of spicules in some areas, and a network of tracts and single spicules surrounding oval-shaped open areas in other places. The ectosomal skeleton is supported by prominent spicule tracts that rise from the choanosome to fan out tangentially in the ectosome (Figure 8B, 8G). The choanosomal skeleton becomes more confused towards the interior, with some spicule tracts but many spicules in confusion. No apparent spongin.

Spicules

Nearly exclusively oxeas, though rare styles can be found. Oxeas have the typical halichondroid shape: gently curved or with a slight central bend, thickest in center and tapering gently to sharp points at both ends. Spicules have a large range in lengths in both the ectosome and the choanosome, but averaged 9% longer in the choanosome. Ectosomal spicules 110–256–438 x 4–6–10 µm, n=44; choanosomal spicules 98–281–459 x 2–6–9 µm, n=44; all spicules together

921 98–268–459 x 2–6–10 µm, n=88. Spicules likely to be indistinguishable from *Ha. panicea*,
922 though the minimum length was very slightly smaller than seen in any examined *Ha. panicea*.
923
924 **Distribution and habitat**
925 Currently known from three samples, all from the far North Atlantic / Arctic: Eastern Shore
926 Islands, Nova Scotia, Canada; Skútan, Northern Iceland; and Svalbard, Norway. Habitat
927 preferences are likely similar to *H. panicea* – though all three samples were collected from the
928 subtidal by divers, so it is possible this species does not occur in the intertidal*.*
929
Remarks
Though we were only able to examine one sample of this new species, the published
mitochondrial genome (MH756604) of another sample from Iceland is nearly identical. When
the entire mitochondrial genome is aligned (coding and non-coding regions), these two samples
differ at only 11 of 12,851 aligned sites (0.09%). In contrast, the average pairwise differences
(Dxy) between the two *Ha. akesa* sp. nov. mitochondria and 9 sequenced *Ha. panicea*
mitochondria is 0.8%, leading to an Fst of 0.745 across the mitochondrial genome between
species. Furthermore, the ribosomal phylogeny and the phylogeny of concatenated
mitochondrial genes confidently place true *Ha. panicea* sequences closer to *Ha. sitiens* than to
Ha. akesa sp. nov. We therefore feel confident in describing these samples as a new species
that is sympatric with *Ha. panicea* across the far North Atlantic. The sample from Nova Scotia
was collected at the same time and place as three other samples that were found to be *Ha.*
panicea, so these species are sympatric even at the scale of individual sites.
Assignment of the Svalbard sample to this species is based on only 610 bp of the cox1
gene (boldsystems.org sample ROPOR090-18). This sequence is, however, identical to the other
Ha. akesa sp. nov. samples, and has several SNPs that differentiate it from *Ha. panicea* and
other similar species.
947
948 *Halichondria sitiens* **(Schmidt, 1870)**
949 Figure 9
950
Material examined
ESS130, Faulker's Shoal, Nova Scotia, (44.60733, -62.96168), 30 m , 2021-08-12; ESS132,
Faulker's Shoal, Nova Scotia, (44.60733, -62.96168), 30 m, 2021-08-12; ESS153, N. of Bull Rock,
Nova Scotia, (44.65375, -62.90491), 30 m , 2021-08-14; ESS44, Bull Shoal, Nova Scotia,
(44.63589, -62.89906), 28 m , 2021-07-21; RBC11, Knight Inlet, British Columbia, (50.68577, -
125.99457), 9 m, 2009-11-08; RBC13, Kuldekduma Island; Weynton Passage, British Columbia,
depth not known, 2019-04-12; RBC8, Knight Inlet, British Columbia, (50.68500, -126.03667), 15-
20 m, 2011-03-29.
959
960 **Diagnosis**
961 Yellow, white, green, or slightly blueish *Halichondria* with numerous papillae decorating the
962 surface. Scattered oscula usually visible, but often not at the ends of papillae. Tangential
963 ectosomal skeleton developed as a web of meandering spicule tracts and individual spicules.
964 Choanosomal skeleton consists of multispicular tracts and many spicules in confusion. Spicules

oxeas, 130–1000 μm long, with averages per sample of 440–630 μm in length and 9–10 μm in width; usually shorter in the ectosome than the choanosome. Additional DNA characters that may be useful in identification are as follows:

COX1. 754:A, 952:A. N=6. These characters differentiate *Ha. sitiens* from all other sequenced *Halichondria* except *Ha. panicea,* which is diagnosed relative to *Ha. sitiens* based on morphological characters such as spicule length and surface papillae.

28S. 2301:G. N=2.

ND1. 97:C. N=5.

Figure 9. *Halichondria sitiens*. A (ESS44), B (ESS132), C (ESS153), D (RBC13, photo by Neil McDonald). E: cross-section showing choanosomal skeleton (ESS153), F: oxeas (RBC8). Tangential ectosomal skeletons: G (ESS130), H (ESS44).

Morphology
Thinly to thickly encrusting or growing as a semi-globular cushion. Surface is often replete with cone-shaped papillae that sometimes branch, but these are less apparent in small sponges. A few scattered oscula are usually visible, often large and prominent and sometimes atop oscular

986   chimneys, but not found at the ends of the papillae. Color generally a pale, translucent yellow
987   but surface can appear nearly white, and field photos of sponges that are likely this species
988   have also documented pale green and slightly bluish individuals. White in ethanol.
989
Skeleton
Tangential ectosomal skeleton is well-developed, but less dense and regularly-patterned than
observed in *Ha. panicea*. Instead, the ectosomal skeleton is comprised of a mesh of meandering
multispicular tracts and individual spicules, with spaces between them variable in size and
shape (similar to species in the Bowerbanki group). Choanosomal skeleton typically
halichondroid, with ill-defined spicule tracts and many spicules in confusion. No apparent
spongin.
997
Spicules
Exclusively oxeas with typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering gently to sharp points at both ends. Spicules have a large
range in lengths in both the ectosome and the choanosome, but average 14% longer in the
choanosome. The difference between ectosome and choanosome is quite variable, with up to
30% longer or 5% shorter spicules in the choanosome. Dimensions are given for individual
samples in table 2. When all samples are combined, spicules measure 138–531–1000 x 4–12–25
µm, n=616. Average values varied considerably across samples, with notably longer spicules
seen in Pacific Canada (means 552–628 µm) than Atlantic Canada (means 448–531 µm).
1007
1008 Table 2. Spicule lengths of *Halichondria sitiens*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
RBC13	318–533–868 n=47	297–724–1000 n=47
RBC11	335–582–892 n=40	332–648–964 n=40
RBC8	308–505–844 n=41	295–597–893 n=43
Atlantic Ocean		
ESS130	312–498–623 n=53	325–455–619 n=59
ESS153	270–466–625 n=40	302–518–698 n=40
ESS44	138–398–680 n=43	332–497–659 n=43
ESS132	258–468–728 n=40	370–563–723 n=40

1009
1010
Distribution and habitat
Known from the Arctic, North Atlantic, and North Pacific, including Greenland, Svalbard, the
White Sea, Alaska, and both coasts of Canada. Samples examined here were from British
Columbia in the Pacific and Nova Scotia in the Atlantic, and these are the only samples with
DNA data available. Previously described as occurring to 100 m depth (Erpenbeck and van
Soest, 2002), but the samples examined here were all collected in the shallow subtidal by
divers.
1018

Remarks

It is remarkable that *Ha. sitiens* is the closest relative of *Ha. panicea*, of all the species with DNA data available. Many of the species described here are morphologically indistinguishable from *Ha. panicea*, while *Ha. sitiens* has longer spicules and surface papillae that distinguish it. The presence of surface papillae has been used as a genus level trait in the past, later downgraded to the subgenus *Eumastia.* Given the close relationship between *Ha. panicea* (type species of subgenus *Halichondria*) and *Ha. sitiens* (type species of subgenus *Eumastia*), we recommend eliminating the subgenera.

In contrast to *Ha. panicea*, where fixed genetic differences were seen between Atlantic and Pacific populations, *Ha. sitiens* has nearly no genetic variation within or between populations at any of the sequenced loci.

Halichondria hygeia **sp. nov.**

Figure 10

Material examined

Holotype: TLT645 (CASIZ 245213), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16. Other samples: BHAK-03597 (UF 4157), Pruth Bay dock, Calvert Island, British Columbia, (51.65500, -128.12900), floating dock, 2017-08-08; FH12x108 (YPM 111876), Roche Harbor, San Juan Island, Washington, (48.60970, -123.15436), floating dock, 2012-05-23; FH12x114 (YPM 111877), Roche Harbor, San Juan Island, Washington, (48.60970, -123.15436), floating dock, 2012-05-23; TLT1076 (SBMNH 718624), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1417 (UCSB-IZC00069011), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2023-09-01; TLT308 (UCSB-IZC00069012), Monterey Marina, California, (36.60797, -121.89288), floating dock, 2019-11-22; TLT504 (CASIZ 245212), Point Pinos, California, (36.63447, -121.93923), intertidal, 2019-11-24; TLT706 (SBMNH 718625), Oceanside Harbor, California, (33.20623, -117.39208), floating dock, 2021-01-09; TLT724 (CASIZ 245214), Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2020-12-14; TLT742 (SBMNH 718626), Dana Point Harbor, California, (33.46090, -117.69156), floating dock, 2021-01-09; TLT940 (UCSB-IZC00069013), Lazy Days, San Diego, California, (32.69415, -117.27110), 12-25 m, 2021-05-15; WMS19x5 (YPM 111878), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2019-05-19; WMS19x6 (YPM 111879), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2019-05-19; WMS19x7 (YPM 111880), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2019-05-19; WMS21x10 (YPM 111881), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x19 (YPM 111882), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x6 (YPM 111883), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x7 (YPM 111884), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x8 (YPM 111885), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS21x9 (YPM 111886), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2021-05-26; WMS23x10 (YPM 111887), West Meadow Beach, Stony Brook, New

1062 York, (40.92673, -73.14807), intertidal, 2023-06-07; WMS23x9 (YPM 111888), West Meadow
1063 Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal, 2023-06-07.
1064
1065 **Etymology**
1066 Name inspired by the Greek goddess Hygeia, sister of Panakeia.
1067

1068
1069 Figure 10. *Halichondria hygeia*. A & B (TLT504), C (TLT645), D (TLT706). E: cross-section showing
1070 choanosomal skeleton at sponge surface (TLT504), F: spicules (FH12x108). Tangential ectosomal
1071 skeletons: G (TLT742), H (TLT645).
1072

Diagnosis
Yellow *Halichondria* with well-developed tangential ectosomal skeleton manifested as a dense
mat of spicules or a mesh of meandering multispicular tracts and individual spicules.
Choanosomal skeleton confused, sometimes accompanied by multispicular tracts. Spicules
oxeas, 170–430 µm in length, with averages per sample of 250–350 µm in length and 5–10 µm
in width. On average, no difference in length is seen between the ectosome than the
choanosome. Diagnosis relative to similar species requires DNA characters, as follows:
1080
1081 COX1. No unique derived states.
1082

1083   28S. 659:G, 670-671:GG, 675:A. N=24.

1084

1085   ND1. 877-880:AACC. State 877:A is unique compared to all close relatives, but homoplastic with
1086   *Halichondria loma.* The full sequence 877-880:AACC is not shared with any other sequenced
1087   *Halichondria*. N=14.

1088

1089   **Morphology**
1090   Thinly to thickly encrusting, sometimes with digitate tendrils. Oscula vary from not apparent, to
1091   slightly volcano-shaped with raised rims, to atop oscular chimneys. Color varies from pale
1092   yellow to golden; not seen in green. See Figure 10 for examples.

1093

Skeleton
Tangential ectosomal skeleton can be sieve-like, with a dense mat of spicules surrounding oval-
shaped open areas (Figure 10G), or more lace-like, with a web of spicule tracts and individual
spicules delineating more irregularly shaped open areas (Figure 10H). This ectosomal skeleton is
supported by poorly developed spicule tracts that delineate large subectosomal spaces, with
the choanosomal skeleton becoming more confused towards the interior. No apparent spongin.

1100

Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central
bend, thickest in the center and tapering to sharp points at both ends. Spicules have a large
range in lengths, though slightly less range than seen in *Ha. panicea*, and spicules in the
ectosome and the choanosome average the same length. Dimensions for five samples where
ectosome and choanosome were measured separately are given in table 3. When spicules from
all seven measured samples are combined, spicules measure 175–276–424 x 3–7–13 µm, n=561
for length, n=405 for width. Average values varied considerably across samples, with mean
lengths 254–346 µm, and mean widths 5–10 µm.

1110

1111   Table 3. Spicule lengths of *Halichondria hygeia*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
FH12x108	284–343–386 n=40	236–349–424 n=40
TLT645	194–259–313 n=40	216–269–312 n=40
TLT742	190–267–301 n=40	193–259–319 n=43
TLT706	188–284–343 n=42	232–279–328 n=43
Atlantic Ocean		
WMS21x9	246–283–336 n=40	204–256–302 n=44

1112

1113   **Distribution and habitat**
1114   In the Northeast Pacific, this species was found from Calvert Island, British Columbia to
1115   Oceanside, Southern California. In contrast to *Ha. panicea*, *Ha. hygeia* sp. nov. was found
1116   primarily on floating docks in sheltered marinas, but it was not limited to this habitat. Of 12
1117   samples collected in the Northeast Pacific, 10 were on floating docks but one was in found in a

wave-swept tide pool at Point Pinos (Monterey Bay, Central California), and one was found in subtidal rocky reef off Point Loma (extreme Southern California, collected between 12 and 25 m depth). The species is also found in the North Atlantic, as it was common on intertidal cobbles at West Meadow Beach, Stony Brook, Long Island, New York. Publicly available sequences of cox1 may indicate that the species is also found in Connecticut and Portugal (Genbank accessions MN448237, MZ487266, figure S1). These sequences are consistent with either *Ha. hygeia* sp. nov. or *Ha. californiana* sp. nov.; as *Ha. californiana* sp. nov. is thus far known only from California, these sequences are likely to be *Ha. hygeia* sp. nov., but confirmation would require sequencing 28S or ND1.

Remarks

Two pieces of evidence suggest that *Ha. hygeia* sp. nov. has experienced a recent human-assisted range expansion. First, artificial substrates like floating docks — where this species is primarily found — are known to harbor many introduced species (Bulleri and Chapman, 2010; Glasby et al., 2007; Ruiz et al., 2000; Simkanin et al., 2012). Second, samples from New York and California are nearly identical across the mitochondrial genes and the ribosomal DNA (Figure 1). Other halichondriids have been shown to have limited larval dispersal (Maldonado, 2006; Xue et al., 2009), so together, these observations make it likely that the species has been moved around by human activity.

     Halichondria hygeia sp. nov. and *Ha. panicea* are likely to be indistinguishable with respect to spicule and skeletal traits, but some differences between them are apparent. These include habitat (*Ha. hygeia* sp. nov. is common on floating docks in marinas, while *Ha. panicea* is common in the wave-exposed intertidal), southern range limit (*Ha. hygeia* sp. nov. was found in Southern California, while *Ha. panicea* was not found in Southern California), and color (*Ha. hygeia* sp. nov. is yellow while *Ha. panicea* can be yellow or green).

Halichondria californiana **sp. nov.**
Figure 11

Material examined

Holotype: TLT937 (CASIZ 245207), Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15. Other samples: TLT1010 (UCSB-IZC00069005), Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-11 m, 2021-07-22; TLT1093 (UCSB-IZC00069006), Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-11 m, 2021-07-22; TLT160 (CASIZ 245204), Isla Vista Reef, California, (34.40278, -119.85755), 9-12 m, 2019-08-01; TLT256 (UCSB-IZC00069007), Isla Vista Reef, California, (34.40278, -119.85755), 9-12 m, 2019-08-01; TLT417 (CASIZ 245205), Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2019-10-25; TLT428 (CASIZ 245206), Elwood Reef, California, (34.41775, -119.90150), 9-15 m, 2019-10-23; TLT812 (SBMNH 718620), Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15; TLT860 (SBMNH 718621), Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06; TLT921 (SBMNH 718622), Point Fermin, California, (33.70664, -118.28595), intertidal, 2021-03-10; TLT928 (UCSB-IZC00069008), Point Fermin, California, (33.70664, -118.28595), intertidal, 2021-03-10; TLT934 (UCSB-IZC00069009), Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15.

1162

Etymology

Named for the state of California.

Diagnosis

Yellow *Halichondria* with tangential ectosomal skeleton developed as a sieve-like mat of spicules or a lace-like network of spicule tracts. Choanosomal skeleton consists of poorly delineated multispicular tracts and spicules in confusion. Spicules oxeas, ranging from 90–290 μm in length, with averages per sample of 130–230 μm long and 3–7 μm wide. On average, no difference in length is seen between the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 1011:A, 1037:T, 1319:C. N=9–11 depending on position.

ND1. 214-217:AAAT. State 217:T is unique compared to all close relatives, but homoplastic with members of the Okadai group (Figure 1). The full sequence 214–217:AAAT is not shared with any other sequenced *Halichondria*. N=10.

1183 Figure 11. *Halichondria californiana*. Field photos: A (TLT937), B (TLT880), C (TLT1093). Ectosomal
1184 skeletons: D (TLT160), E (TLT937), F (TLT1010), G (TLT860). H: cross-section showing choanosomal
1185 skeleton at sponge surface (TLT160), I: spicules (TLT860).
1186

1187 **Morphology**
1188 Encrusting yellow sponges; oscula often small and flush with sponge surface, but can also occur
1189 on oscular chimneys. Color varies from pale, translucent yellow to golden, sometimes with red
1190 blotches; not seen in green. Beige in ethanol. See Figure 11 for examples.
1191

1192 **Skeleton**
1193 Tangential ectosomal skeleton can be sieve-like, with a dense mat of spicules, or more lace-like,
1194 with a web of spicule tracts and single spicules. Open areas are generally oval or polygonal and
1195 relatively regular in size. Choanosomal skeleton with poorly developed spicules tracts and many
1196 spicules in confusion. No apparent spongin.
1197

Spicules
Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have a typical halichondroid shape: evenly curved or with a slight central bend, thickest
in the center and tapering gently to sharp points at both ends. Spicules have a large range in
lengths, though slightly less range than seen in *Ha. panicea*, and spicules in the ectosome and
the choanosome average the same length. Dimensions for the six samples where ectosome and
choanosome were measured separately are given in table 4. When spicules from 10 measured
samples are combined, spicules measure 99–194–286 x 2–5–9 µm, n=813 for length, n=733 for
width. Average values varied considerably across samples, with means of 134–226 µm long and
3–7 µm wide.
The maximum length of 286 µm is notably shorter than some similar species, such as
Ha. hygeia sp. nov. (424 µm), *Ha. panicea* (499 µm), and *Ha. cascadia* sp. nov. (444 µm) —
though *Ha. columbiana* sp. nov. is similar (266 µm). Spicule size is insufficient to differentially
diagnose all samples, however. For example, some individual *Ha. californiana* sp. nov. samples
had average and maximum lengths exceeding some *Ha. hygiea* sp. nov. samples, and some *Ha.*
californiana sp. nov. had average lengths exceeding some *Ha. panicea*.
1214

1215 Table 4. Spicule lengths of *Halichondria californiana*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT937	149–207–239 n=40	177–215–259 n=40
TLT1010	113–136–172 n=51	99–132–161 n=51
TLT921	121–177–199 n=40	145–180–208 n=40
TLT256	107–146–164 n=40	122–155–190 n=41
TLT860	187–235–274 n=40	190–236–286 n=59
TLT428	167–213–258 n=40	166–218–266 n=40

1216

Distribution and habitat
Known only from California, from San Luis Obispo Bay in Central California to Point Loma in
Southern California. Nearly allopatric with *Ha. panicea*: the two species were found growing

side-by-side in a tidal pool near Avila Beach, California, but this was the southernmost location
where *Ha. panicea* was found and the northernmost location for *Ha. californiana* sp. nov.. Like
Ha. panicea, this species was found growing on rock or algae, in natural subtidal reefs and
exposed intertidal areas, and not found on human structures in sheltered marinas. The deepest
samples were found between 9 and 18 m.
1225
Remarks
In contrast to its closest known relative *Ha. hygeia* sp. nov., this species was found only in
natural habitats, and not on human structures. It is also different from *Ha. hygeia* sp. nov. in
that it was only found in California and thus could be endemic to the Northeast Pacific.
However, these two species cannot be differentiated at the commonly sequenced regions of
cox1, so publicly available sequences from Connecticut and Portugal could be this species
(Genbank accessions MN448237, MZ487266). The nudibranch *Doris montereyensis* was seen
feeding on this species in multiple locations.
1234
1235 ***Halichondria cascadia* sp. nov.**
1236 Figure 12
1237

1238
1239 Figure 12. *Halichondria cascadia*. A: sample post-preservation (FH12x92), B: tangential ectosomal
1240 skeleton (FH12x93), C: choanosomal skeleton (FH12x92).
1241
1242 **Material examined**
1243 Holotype: FH12x92 (CASIZ 245208, YPM 111874), Eagle Cove, San Juan Island, Washington,
1244 (48.46128, -123.03155), intertidal, 2012-05-22. Additional sample: FH12x93 (YPM 111875),
1245 Eagle Cove, San Juan Island, Washington, (48.46128, -123.03155), intertidal, 2012-05-22.
1246
1247 **Etymology**
1248 Name inspired by the term "Cascadia region" which is sometimes used to refer to an area
1249 including Oregon, Washington, and British Columbia.
1250
Diagnosis
Halichondria with a tangential ectosomal skeleton developed as a mesh of meandering
multispicular tracts and individual spicules. Choanosomal skeleton consists of spicules and
spicule bundles in confusion. Spicules oxeas, 160–450 μm long, with averages per sample of
270–320 μm long and 8-9 μm wide. Infrequent styles and rare strongyles can be observed.

Some samples average shorter spicules in the ectosome than the choanosome. Diagnosis based
primarily on DNA characters, as follows:

1258

1259 COX1. 949:A. N=3.

1260

1261 28S. 2061:T. N=2.

1262

1263 ND1. 847:A. N=2.

1264

1265 **Morphology**
1266 Known only from two small samples, so little is known about the morphology of this species.
1267 Preserved samples are white; color alive is yellow. Both samples have oscular chimneys
1268 approximately 1 cm in height.

1269

1270 **Skeleton**
1271 Tangential ectosomal skeleton a web of spicule tracts and individual spicules, with open areas
1272 irregular in size and shape (Figure 12B). Choanosomal skeleton of spicules and spicule bundles
1273 in confusion. No apparent spongin.

1274

Spicules
Oxeas with a typical halichondroid shape: evenly curved or with a slight central bend, thickest in
the center and tapering to sharp points at both ends. Infrequent styles and rare strongyles can
be observed. Spicules have a large range in lengths, with one sample having significantly
shorter spicules in the ectosome while the other sample has no difference in length between
ectosome and choanosome. When spicules from both samples are combined, spicules measure
162–294–444 x 3–8–12 μm, n=169. Table 5 shows dimensions for samples and domains
separately.

1283

1284 Table 5. Spicule lengths of *Halichondria cascadia*. Values given as min–mean–max in μm.

Specimen	Ectosome	Choanosome
FH12x92	178–267–353 n=41	162–273–373 n=46
FH12x93	199–287–280 n=42	260–353–444 n=40

1285

1286 **Distribution and habitat**
1287 Samples examined here were both collected from rocky intertidal habitats at Eagle Cove, San
1288 Juan Island, Washington. Publicly available sequence data suggests a third sample was found at
1289 Carl Moses Harbor, Captains Bay, Unalaska Island (Genbank accession MZ580608). Though only
1290 658 bp of cox1 is available for this sample, it is identical to sample FH12x93.

1291

Remarks
The two samples examined here are small fragments. Though they were collected at the same
location, they are genetically different from one another: they differ at 1.17% of bases in the
nuclear ribosomal locus and at 0.6% of bases in the protein-coding mitochondrial genome. They
form a well-supported clade in the phylogeny, with 98.5% bootstrap support, and are the sister

group to the other three species in the Hygeia group (*Ha. hygeia* sp. nov.*, Ha. californiana* sp.
nov.*, and Ha. columbiana* sp. nov.). They were collected at the same time and place as *Ha.*
panicea, and *Ha. hygiea* sp. nov. was collected at a marina on the same island the following
day, so this species is sympatric with several closely related *Halichondria.* Despite the limited
sampling of this species, the strong support for *Ha. hygiea* sp. nov. and *Ha. californiana* sp. nov.
as different species compels us to formally describe these samples in the hope that more can
be learned about this enigmatic group.
1304
1305   **Halichondria columbiana sp. nov.**
1306   Figure 13
1307
1308   **Material examined**
1309   Holotype: BHAK-09466 (UF 4408), Port Reef, Hakai Pass, British Columbia, (51.697, -128.106),
1310   28 m, 2019-05-28. Additional sample: BHAK-09479, Whidbey Point, Fitz Hugh Sound, British
1311   Columbia, (51.71700, -127.89200), 19 m, 2019-05-29.
1312
1313   **Etymology**
1314   Named for the British Columbia region of Canada.
1315
1316   **Diagnosis**
1317   *Halichondria* with tangential ectosomal skeleton developed as a chaotic mesh of meandering
1318   multispicular tracts and individual spicules. Choanosomal skeleton confused. Spicules oxeas,
1319   150–270 µm long, with means per sample of 220–240 µm long and 8–9 µm wide. Diagnosis
1320   based primarily on DNA characters, as follows:
1321
1322   COX1. 973:C, 1012:C, 1102:G, 1207:A. N=2.
1323
1324   28S. No unique derived states.
1325
1326   ND1. 79:C, 88:C, 172:T, 340:G, 352:C, 733:G, 811:G. N=2.
1327

1328
1329   Figure 13. *Halichondria columbiana*. Photos of sponges after collection, before preservation: A (BHAK-
1330   9466), B (BHAK-9479). C: tangential ectosomal skeleton (BHAK-9466).
1331
1332   **Morphology**

1333 Known only from two small samples, little is known about the morphology of this species, but
1334 they appear similar to encrusting *Ha. panicea*. Samples were yellow alive and are white in
1335 ethanol.
1336
1337 **Skeleton**
1338 Tangential ectosomal skeleton is a web of spicule tracts and individual spicules (Figure 13C).
1339 Choanosomal skeleton difficult to characterize in such small samples, but appears to be
1340 confused, as seen in all closely related species. No apparent spongin.
1341
Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central
bend, thickest in the center and tapering to sharp points at both ends. When spicules from both
samples are combined, spicules measure 149–228–266 x 4–8–11 µm, n=151. Spicules from the
ectosome and choanosome were measured separately for one sample and the sizes did not
differ by compartment (the ectosome could not be located in the other small sample).
1348
1349 **Distribution and habitat**
1350 The two known samples were collected 15 km apart in British Columbia. Both were subtidal,
1351 between 19 and 28 m depth.
1352
Remarks
The two samples examined here are very small fragments. The samples are identical at the
ribosomal locus and only slightly different at mitochondrial loci (sequence differences of 0.4%
and 0.2% at ND1 and cox1, respectively). Contrastingly, they are genetically distinct from the
other species in the Hygeia group: when compared to *Ha. hygeia* sp. nov., average pairwise
differences are 1.0% (Fst=0.9) at the ribosomal locus and 1.2% (Fst=0.84) at cox1. Despite the
limited sampling of this species, the strong support for *Ha. hygiea* sp. nov. and *Ha. californiana*
sp. nov. as different species compels us to formally describe these samples in the hope that
more can be learned about this enigmatic group.
1362
1363 *Halichondria dokdoensis* **Kang, Kim & Kim, 2022**
1364 Figure 14
1365

1366
1367 Figure 14. *Halichondria dokdoensis*. A: field photo, B: tangential ectosomal skeleton, C: cross-section
1368 showing choanosomal skeleton at sponge surface, D: oxea. All photos from TLT386.
1369

Material examined

TLT386, Oil Platform C, Santa Barbara, California, (34.33293, -119.63173), 21 m, 2019-08-05.

Diagnosis

Yellow, reddish-purple, or orange *Halichondria* with tangential ectosomal skeleton developed as a lace-like network of spicule tracts surrounding oval-shaped open areas. Choanosomal skeleton with multispicular tracts supporting the ectosome and delineating large subectosomal spaces, with an increasingly confused skeleton towards the interior. Spicules oxeas, 130–310 µm long and 1–10 µm wide; slightly shorter in the ectosome than the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. 100:C, 247:G, 361:G, 844:C, 892:T. N=2-3 depending on position.

28S. 162:T, 195:C, 260:G, 584:C, 614:T, 671:T, 1660:G. N=1.

ND1. 286:C, 322:A, 346:G, 403:T, 769:G. N=2.

Morphology

Samples from Korea were previously described as reddish-purple to orange with prominent oscular chimneys (Kang et al., 2022). A sample from California was yellow, with oscula nearly flush with the sponge surface, so color and shape appear to be variable, as seen in many other *Halichondria* species.

Skeleton

Well-developed tangential ectosomal skeleton with a lace-like web of spicules and spicule tracts surrounding open spaces that are oval shaped and fairly consistently sized (Figure 14B). This ectosomal skeleton is supported by multispicular tracts that delineate large subectosomal spaces. The choanosomal skeleton becomes more confused towards the interior. No apparent spongin.

Spicules

Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend, thickest in the center and tapering gently to sharp points at both ends (rare spicules occasionally found with a centrotylote swelling). The single sponge we examined had spicules 177–230–287 x 2–6–8 µm, n=146 for length, n=106 or width. When measured separately, spicules in the choanosome averaged 238 µm in length vs. 221 µm in the ectosome (p=0.001). Spicules from Korean samples previously reported as 135–310 x 1–8 µm (means and sample size not reported). Spicules were previously reported as having "size classes", with some being thin (1–2.5 µm) and some thick (5–8 µm). Of the 146 spicules we measured, only one would fall in the thin category, with 10 falling between thick and thin. The abundance of thin oxea is therefore variable and/or universally rare, perhaps because they are immature thick spicules.

Distribution and habitat

The holotype location for this species is the Dokdo Islands, South Korea, with additional samples from mainland South Korea and Jeju Island. The sample examined here was found on an oil platform in the Santa Barbara Channel in Southern California, and a publicly available sequence matching this species was obtained near Goes, Zeeland, the Netherlands (see remarks). The California sample was subtidal (21 m), and though depths are not reported for the other samples, the Korean samples were also collected by SCUBA divers.

Remarks

Halichondria dokdoensis was described based on morphological characters, but it is our view that it cannot be morphologically distinguished from *Ha. panicea*. Characters previously listed as distinguishing the species were color (but we show here that *Ha. dokdoensis* can be found in yellow) and larger spicules than *H. panicea* (but we show here that the spicules of *Ha. panicea* span a large size range encompassing that of *Ha. dokdoensis*). However, a mitochondrial genome sequence was previously published for *Ha. dokdoensis*, and we are able to use this sequence to "rescue" this previously published name (Kim et al., 2019). This sequence is nearly identical to the mitochondrial genome of our California sample: they differ by only 12 of 13,033 aligned base pairs (<0.1%). These samples are more closely related to *Hymeniacidon actites* than to any other sequenced *Halichondria*, and together these two species form a clade that is the sister to the combined Panicea and Hygeia species groups.

This species is likely to have experienced a recent human-mediated range expansion. Finding samples in both California and Korea could plausibly be due to a natural range across the North Pacific, though this is made less likely by the observation that it was only found (thus far) on an artificial structure in California. However, a publicly available sequence of cox1 indicates that the species is also likely found in the Netherlands (NLMAR324-20, boldsystems.org). Though this sequence is only 618 bp of the cox1 gene, it is identical to other *Ha. dokdoensis* and does not match any other sequenced species. This species is therefore likely to be introduced to multiple regions of the Northern Hemisphere, similar to other species described here.

Halichondria galea **sp. nov.**

Figure 15

Material examined

Holotype: TLT495 (CASIZ 245211), Point La Jolla, California, (32.85227, -117.27239), intertidal, 2020-02-08. Other samples: TLT496 (SBMNH 718623), Point La Jolla, California, (32.85227, -117.27239), intertidal, 2020-02-08; TLT743 (UCSB-IZC00069010), Dana Point Harbor, California, (33.46090, -117.69156), floating dock, 2021-01-09.

Etymology

Name inspired by the word "galleon", one of the ships of the Spanish armada in the age of sail.

Diagnosis

Encrusting green, yellow, or brownish *Halichondria*, sometimes with a rugose surface. Tangential ectosomal skeleton is a sieve-like mat of spicules surrounding open areas that are

1457   generally oval-shaped and consistent in size. Choanosomal skeleton of multispicular tracts and
1458   abundant spicules in confusion. Spicules oxeas, 100–500 μm long, with averages per sample of
1459   250–290 μm long and 5–6 μm wide. Spicules 10–20% longer in choanosome than ectosome,
1460   but the full range of spicules lengths is seen in both compartments. Diagnosis relative to similar
1461   species requires DNA characters, as follows:

1462

1463   COX1. 1069:A. N=8.

1464

1465   28S. 516:T, 520:G, 547:G, 1943:T, 2111:T. N=2–3 depending on position.

1466

1467   ND1. No unique derived states.

1468
1469

1470
1471   Figure 15. *Halichondria galea*. A (TLT495), B & C (TLT743, C shows underside that was attached to
1472   floating dock). D: tangential ectosomal skeleton (TLT496), E: cross-section showing choanosomal
1473   skeleton at sponge surface (TLT495), F: oxeas (TLT495).

1474
1475 **Morphology**
1476   Thinly to thickly encrusting with uneven thickness, leading to a lumpy, undulating, or rugose
1477   appearance. Oscula, when apparent, nearly flush with surface. Surface is a mix of yellow and
1478   green regions, or yellow, green, and brownish regions, with the interior yellow. See Figure 15
1479   for examples.

1480
1481 **Skeleton**
1482   Tangential ectosomal skeleton a sieve-like mat of spicules surrounding oval-shaped open areas
1483   with fairly consistent size. Choanosomal skeleton with meandering multispicular tracts and
1484   many spicules in confusion. No apparent spongin.
1485

Spicules

Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central bend, thickest in the center and tapering to sharp points at both ends. Like *Ha. panicea*, spicules have a very large range in lengths. Though spicules in the ectosome are 10–20% shorter than the choanosome, the full range of spicule lengths is seen in both compartments, with size distributions continuous but bimodal within each compartment. Dimensions for three samples where ectosome and choanosome were measured separately are given in table 6. When spicules from all samples are combined, spicules measure 105–265–483 x 2–5–10 µm, n=308. Average dimensions per sample were 249–270 µm long and 5–6 µm wide.

Table 6. Spicule lengths of *Halichondria galea*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT495	114–256–440 n=50	137–284–445 n=50
TLT496	126–269–472 n=40	125–300–483 n=40
TLT743	105–225–401 n=65	106–273–412 n=63

Distribution and habitat

Known from China, Korea, and California. In the Eastern Pacific, this species is known from Orange and San Diego Counties in Southern California. We found it growing on a floating dock in Dana Point Harbor and in natural tidepools at Point La Jolla; it was also previously reported from Mission Bay, San Diego (Park et al., 2007). In the Western Pacific it has been collected from the Gulei Peninsula in China (Wang et al., 2016) and from rocky intertidal areas around the Korean Peninsula and islands of the Korean Strait (Park et al., 2007).

Remarks

The first mitochondrial genome published for a *Halichondria* species was for a "*Halichondria sp.*" sponge sampled from Fujian Province, China (Wang et al., 2016). Sequencing of mitochondrial genes later documented this same genotype from the intertidal zone of mainland South Korea and nearby islands, and from Mission Bay, San Diego, California (Park et al., 2007). This sponge has remained undescribed, likely due to it being morphologically indistinguishable from *Ha. panicea*.

Our Southern California samples have mitochondrial genomes that are nearly identical with the published sequence from China (99.95% identical), confirming that we have recollected this same species from California. Consistent with previously published data, it is only distantly related to *Ha. panicea*, despite their morphological similarity. The clade we designate here as the Okadai species group contains *Ha. okadai* and *Ha. galea* sp. nov., which are sympatric in the Korean Strait, as well as the Caribbean species *Ha. magniconulosa* and *Ha. melanadocia,* and samples we collected from Pacific Panama that are likely from an undescribed species (Figure 3).

It is probable that *Ha. galea* sp. nov. has experienced a recent human-mediated range expansion, though the evidence is less compelling than for the other species reported here. It has only been found in the North Pacific thus far, and this could indicate a natural range connected through the Bering Strait. It has been found in artificial habitats like floating docks, but most samples reported to date have been collected from rocky intertidal areas that host

fewer introduced species. However, finding nearly identical mitochondrial genomes in China
and California makes it less likely that those populations are found at extreme ends of a wide
distribution, and more likely that they share a recent common ancestor. When combined with
the observation that many other *Halichondria* species have been moved around by human
activity, we feel this species should be treated as a possible introduction across some part of its
range (though which part remains to be determined).
1532
1533 ***Halichondria bowerbanki* Burton 1930**
1534 Figure 16
1535
Material examined
190615, Praia to Lumiar, Viana do Castelo, Portugal, (41.73540, -8.87578), 7.1 m, 2019-06-15;
190616, Canto Marinho, Viana do Castelo, Portugal, (41.72910, -8.88071), 12.5 m, 2019-06-16;
NIB0309, The Dorn, Northern Ireland, (54.43459, -5.54536), intertial (boulders), 2021-04-28;
NIB0612, Carnalea , Northern Ireland, (54.66844, -5.70801), intertidal (rocky shore), 2021-10-
27; NIB0614, Mount Stewart, Northern Ireland, (54.54482, -5.61017), intertidal (low shore
boulders), 2021-04-29; TLT1052, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1072, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1087b, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01; TLT1357, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT1358, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT1418, Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2023-09-01; TLT234, Arroyo Quemado Reef, California, (34.46775, -120.11905),
7-11 m, 2019-07-29; TLT337, Saddleback Ridge, Santa Cruz Island, California, (34.03817, -
119.52470), 5-12 m, 2019-10-18; TLT342, Coal Oil Point, California, (34.40450, -119.87890), 3-8
m, 2019-08-30; TLT415, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2019-10-25;
TLT490, Coal Oil Point, California, (34.40450, -119.87890), 5 m, 2020-07-17; TLT738, Marina del
Rey, California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT980, Santa Barbara
Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT981, Santa Barbara
Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01.
1556
Diagnosis
Yellow, green, or blue-green Halichondria with a tangential ectosomal skeleton developed as a
web of spicule tracts and individual spicules, with open spaces generally triangular or polygonal
rather than oval. Choanosomal skeleton confused, with a slight tendency to form multispicular
tracts. Spicules oxeas, 180–550 μm long, with averages per sample of 300–430 μm long and 7–
12 μm wide; usually the same length in the ectosome and the choanosome. Diagnosis relative
to similar species requires DNA characters, as follows:
1564
1565 COX1. 301:C, 484:A . N=12.
1566
1567 28S. 8:G, 848:T. 853:G, 865:C, 1060:C, 1937:C, 2032:A, 2327:A. N=13–20 depending on position.
1568
1569 ND1. 264:G, 684:A, 691:C, 739:T. N=13–14 depending on position.

1570

Morphology

Previously described as variable in shape, from encrusting to cushion-shaped, with oscular chimneys and/or thread-like tendrils, and sometimes highly ramose with anastomosing branches (Ackers et al., 2007; Burton, 1930; Evcen et al., 2023). Samples examined here, with identities confirmed by DNA sequencing, do not include any forms with branches, threads, or oscular chimneys, though the previously sequenced sample from Ireland (BELUM:Mc4003) had these features.

Sponges from sheltered marinas were thinly encrusting, usually green or blue-green but one sample was yellow. Samples from subtidal rocky reefs were usually large cushions, pale yellow, but one sample was encrusting and green. Oscula are not always apparent, but when they are, they are nearly flush with the sponge surface or with raised rims, and often accompanied by a translucent channel running up one side (Figure 16B). White or beige in ethanol.

Figure 16. *Halichondria bowerbanki*. A & B (TLT490), C (TLT337), D (TLT738), E (TLT1052), F (TLT234). G: spicules (TLT234). Tangential ectosomal skeletons: H (TLT337), I (TLT234). J: cross-section showing choanosomal skeleton at sponge surface (TLT342).

Skeleton

1591 Tangential ectosomal skeleton is a web of spicule tracts and individual spicules, with open
1592 spaces generally triangular or polygonal rather than oval. The choanosomal skeleton is
1593 confused, with many free spicules, but meandering multispicular tracts are usually evident. No
1594 apparent spongin.
1595
Spicules
Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have the typical halichondroid shape: evenly curved or with a slight central bend,
thickest in the center and tapering gently to sharp points at both ends. Spicules have a large
range in lengths in both the ectosome and the choanosome, with no average difference
between compartments (though individual samples varied from 5% shorter to 7% longer in the
choanosome). Dimensions for 10 samples where ectosome and choanosome were measured
separately are given in table 7. When spicules from all 16 samples with data are combined,
spicules measure 185–379–547 x 2–8–17 µm, n=1312 for length, n=1092 for width. Average
values varied considerably across samples, 307–423 µm long, and 7–12 µm wide.
1606
1607 Table 7. Spicule lengths of *Halichondria bowerbanki*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
TLT1052	273–356–454 n=45	203–359–410 n=40
TLT342	350–433–490 n=40	291–411–528 n=41
TLT490	340–412–477 n=40	265–430–506 n=40
TLT980	231–333–384 n=40	249–323–371 n=46
TLT981	251–322–389 n=40	213–323–379 n=40
TLT234	228–424–539 n=97	196–422–547 n=96
Atlantic Ocean		
NIB0512	185–307–365 n=40	206–306–399 n=40
NIB0614	292–389–467 n=45	320–416–491 n=45
NIB0309	263–362–488 n=96	210–370–472 n=96
190615	334–408–469 n=40	249–391–507 n=40

1608
Distribution and habitat
This species was previously described as occurring throughout Europe, both coasts of North
America, and the Korean Peninsula (de Voogd *et al.* 2024) However, many of these samples are
likely to be members of the cryptic species newly described here, so the confirmed range of this
species is limited to samples with genetic data. European samples with DNA confirmation are
from Ireland, the Netherlands, Portugal, and the Black Sea. Confirmed Pacific samples were
found in a small number of Southern California locations. Samples previously sequenced from
Argentina expand the known range of this species into the Southern Hemisphere (Figure S1).
This distribution is likely a result of accidental human transport, and the species seems likely to
be confirmed from additional locations in the future.
Previous publications have stated that this species is less tolerant of tidal exposure than
Ha. panicea, but more tolerant of silted and brackish conditions (Ackers et al., 2007; Hartman,

1958; Vethaak et al., 1982). DNA-confirmed samples are consistent with this description, as the
species was found in sheltered marinas and subtidal reefs, but intertidal samples were verified
only in Ireland. Samples were collected on floating docks in several marinas, but *Ha. urca* sp.
nov.*, Ha. zabra* sp. nov.*,* and *Ha. hygeia* sp. nov. were much more common and widespread on
floating docks than *Ha. bowerbanki.* Subtidal samples were collected in natural rocky reef
habitat, up to 12 m deep, usually growing on holdfasts of *Macrocystis pyrifera*. Though not
quantitative, our observations of this species in the California kelp forest demonstrate
substantial differences in abundance through time. The species was among the most abundant
sponges at two Southern California locations near Santa Barbara (Coal Oil Point and Arroyo
Quemado Reef) in 2019 and 2020, but could not be relocated in those areas in 2023 or 2024.
1631
Remarks
Though *Ha. panicea* and *Ha. bowerbanki* cannot always be distinguished without genetic data,
they do have average differences in morphological characters and habitat. However, *Ha.*
bowerbanki is much harder to distinguish from other *Halichondria* such as *Ha. zabra* sp. nov.*,*
Ha. urca sp. nov.*,* and *Ha. pinaza* sp. nov.*,* and many previous reports of *H. bowerbanki* are
likely to be conflated with these species. In California, for example, ramose *Halichondria* from
harbors and marinas have long been attributed to *Ha. bowerbanki*, but we found *Ha.*
bowerbanki to be uncommon in California marinas, and found it to be thinly encrusting in those
habitats when present.
It is likely that *Ha. bowerbanki* is introduced in some parts of its range. The presence of
this species in the North Atlantic and North Pacific could potentially be explained by a natural
range that includes the Arctic, though this seems less likely than for *Ha. panicea* because 1)
genetic variation does not seem to be associated with geographic distance, 2) the species is
found in human-modified habitats where introduced species are common, and 3) samples
confirmed thus far suggest a less cold-tolerant range than for *Ha. panicea* (in the Pacific, where
our sampling is most extensive, it has been found only in Southern California so far). Moreover,
several *Ha. bowerbanki* were collected from Punta Verde, a beach in San Antonio Bay,
Argentina, by Marianela Gastaldi (*pers. comm.*). Sequencing of the cox1 locus confirms these
specimens are *Ha. bowerbanki* (MZ487239, MZ487236, MZ487221). It is highly unlikely that this
population was founded by natural means.
1652
1653 *Halichondria urca* **sp. nov.**
1654 Figures 17 & 18
1655
Material examined
Holotype: TLT1087 (CASIZ 245218), Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01. Other samples: 15492 (BULA-0171), Cabrillo Beach, California,
(33.70800, -118.28500), intertidal, 2019-08-20; 15925 (BULA-0236), Port of Long Beach,
California, (33.72865, -118.23638), 10 m, 2019-08-20; MSH23x2 (YPM 111898), Mount Sinai
Harbor, New York, (40.96345, -73.03646), 1 m, 2023-08-16; NIB0610 (BELUM.Mc2024.7),
Carrickfergus Marina , Northern Ireland, (54.71028, -5.81237), floating dock, 2021-10-24;
TLT1002 (SBMNH 718631), Farnsworth Bank, California, (33.34380, -118.51650), 21-27 m, 2021-
07-10; TLT1050 (UCSB-IZC00069017), Santa Barbara Harbor, California, (34.40559, -119.68964),

floating dock, 2021-06-01; TLT1051 (SBMNH 718632), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1053 (UCSB-IZC00069018), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1058 (UCSB-IZC00069019), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1060 (UCSB-IZC00069020), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1062 (UCSB-IZC00069021), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1063 (SBMNH 718633), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1065 (UCSB-IZC00069022), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1066 (UCSB-IZC00069023), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1068 (SBMNH 718634), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1069 (UCSB-IZC00069024), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1070 (UCSB-IZC00069025), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1071 (UCSB-IZC00069026), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1073 (SBMNH 718635), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1075 (UCSB-IZC00069027), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1078 (UCSB-IZC00069028), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1081 (SBMNH 718636), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1083 (SBMNH 718637), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1085 (SBMNH 718638), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT534 (UCSB-IZC00069029), Point La Jolla, California, (32.85227, -117.27239), intertidal, 2020-02-08; TLT590 (UCSB-IZC00069030), Mission Bay: Quiviera Basin, California, (32.76421, -117.23829), floating dock, 2020-02-08; TLT591 (SBMNH 718639), Mission Bay: Quiviera Basin, California, (32.76421, -117.23829), floating dock, 2020-02-08; TLT593 (CASIZ 245219), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16; TLT594 (UCSB-IZC00069031), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16; TLT661 (UCSB-IZC00069032), Stearn's, California, (34.41023, -119.68563), 3-5 m, 2020-08-25; TLT715 (UCSB-IZC00069033), Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2020-12-14; TLT720 (SBMNH 718640), Mission Bay: South Shores Boat Launch, California, (32.76406, -117.21750), floating dock, 2021-01-10; TLT733 (CASIZ 245220), Marina del Rey, California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT740 (CASIZ 245221), Marina del Rey, California, (33.97228, -118.44653), floating dock, 2021-01-09; TLT747 (SBMNH 718641), San Diego Bay: Shelter Island Marinas, California, (32.71094, -117.23423), floating dock, 2021-01-10; TLT764 (SBMNH 718642), Newport Bay: Balboa Peninsula, California, (33.60375, -117.90052), intertidal (on piling), 2021-01-09; TLT766 (SBMNH 718643), Newport Bay: Balboa Peninsula, California, (33.60375, -117.90052), floating dock, 2021-01-09; TLT926 (UCSB-IZC00069034), Mission Bay: Quiviera Basin, California, (32.76421, -117.23829), floating dock, 2021-05-15; TLT951, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-05-24; TLT951B (UCSB-IZC00069035), Nudi Wall, California, (32.69872, -117.27580), 18-28 m, 2021-05-16; TLT960 (UCSB-IZC00069036), Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24.

1709

Etymology

1710

1711 Named for the urca, one of the ships of the Spanish armada in the age of sail.

1712

Diagnosis

Yellow *Halichondria* with a tangential ectosomal skeleton developed as a web of spicule tracts
and individual spicules, or as a sieve-like mat of spicules surrounding oval-shaped open spaces.
Spicules oxeas, 100–440 µm long, with averages per sample of 230–360 µm long and 5–10 µm
wide; usually very slightly longer in the choanosome than the ectosome. Diagnosis relative to
similar species requires DNA characters, as follows:

1719

1720 COX1. 72:T, 93:A, 128:G, 139:C, 247:C, 310:C, 313:T, 364:T. 397:G, 412:T, 475:T, 538:G, 559:G,
1721 634:T, 709:C, 823:G, 880:C, 979:C, 1040:T, 1041:G, 1102:T, 1120:T, 1150:A, 1321:T, 1432:G.
1722 N=9–21 depending on position.

1723

1724 28S. 693:A, 704:G, 829:C, 846:A, 884:T, 1359:G, 1373:C, 1656:G, 2331:G, 2102:C, 2328:T. N=9–
1725 40 depending on position.

1726

1727 ND1. 10:G, 121:C, 204:C, 228:C, 364:T, 448:G, 543:A, 658:T, 838:T. N=9–15 depending on
1728 position.

1729

1730
1731 Figure 17. *Halichondria urca*. A (TLT 661), B (TLT740), C (TLT747), D (TLT1063), E (TLT534), F (TLT1069,
1732 sponge torn to show brooded larvae inside). A (subtidal) & E (intertidal) are in situ, others are pictured
1733 after removal from floating docks, before preservation.

1734

Morphology

Samples discovered to date were bright yellow to yellowish brown, and did not contain any
traces of green. Shape is variable from thin encrustations to semi-globular cushions. Oscula can

be flush with surface, with raised rims, on volcano-shaped protrusions, or atop oscular
chimneys (Figure 17). Sometimes has tendril-like projections. Translucent areas usually present,
including channels leading up the sides of oscula, as often seen in *Ha. bowerbanki*. When
removed from the substrate or torn, brooded bright yellow larvae may be present internally
and where the sponge contacted the substrate (Figure 17F). In Santa Barbara Harbor, Southern
California, larvae were seen in one of two sponges collected in February, a single sponge
collected in May, and 5 of 20 sponges collected in June.
1745

Skeleton
The tangential ectosomal skeleton is usually developed as a web of long meandering spicule
tracts and single spicules, with open spaces between them irregular in size and shape.
However, some samples have more regular open spaces that are triangular and polygonal, and
two samples had a mat of spicules that surrounded oval-shaped openings (Figure 18C).
Choanosomal skeleton typically halichondroiid, with meandering spicules tracts and spicules in
confusion, without apparent spongin.
1753

Spicules
Nearly exclusively oxeas, though rarely a spicule can be found that is modified into a style.
Spicules have a typical halichondroid shape: evenly curved or with a slight central bend, thickest
in the center and tapering to sharp points at both ends. Spicules have a large range in lengths in
both the ectosome and the choanosome, with a very slight tendency to be longer in the
choanosome (average of 3% longer across samples). Table 8 shows dimensions for 6 samples
where ectosome and choanosome were measured separately. When all 19 samples with
measurements are combined, spicules are 107–308–435 x 1–7–14 µm, n=1047 for length,
n=863 for width. Average values varied across samples, with means of 238–359 µm long, 5–10
µm wide.
1764

1765 Table 8. Spicule lengths of *Halichondria urca*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
TLT926	264–315–368 n=50	273–327–362 n=50
TLT951	241–296–335 n=40	242–320–364 n=40
TLT1051	277–308–361 n=40	241–310–359 n=42
TLT661	183–254–315 n=50	181–239–296 n=50
TLT715	211–315–365 n=42	288–332–391 n=42
Atlantic Ocean		
NIB0610	259–346–416 n=40	316–372–425 n=40

1766

Distribution and habitat
Known from the intertidal to at least 28 m depth, but this species is much more common on
floating docks in sheltered marinas than in other habitats investigated. Exceptions include
encrusting samples found exposed at low tide at Point La Jolla and on a rocky reef in the Point
Loma kelp forests, both near San Diego, California. Other samples found outside of marinas

were on man-made debris: entangled trash 3 m below the Santa Barbara Wharf and an
entangled rope at 21 m depth in the Farnsworth Offshore State Marine Conservation Area at
Catalina Island, California.
Pacific samples are all from Southern California. Atlantic samples are from New York,
Virginia, and Ireland. It was the most abundant species of *Halichondria* found in marinas in
Southern California, where 48% of *Halichondria* (34 of 71 samples) were this species (sampling
strategies were aimed at maximizing diversity rather than estimating relative abundance, so
this number is likely an underestimate). *Halichondria urca* sp. nov. was not found at higher
latitudes in the Northeast Pacific: only *Ha. zabra* sp. nov., *Ha. pinaza* sp. nov., and *Ha. hygeia*
sp. nov. were collected in marinas from Central California to British Columbia, suggesting that
Ha. urca sp. nov. may be less cold-tolerant than those species (though sampling in these areas
was more limited than in Southern California).
1784
Remarks
This species is often morphologically indistinguishable from other members of the Bowerbanki
group, but there were some features that made it more likely a sample would prove to be *Ha.*
urca sp. nov. when genotyped. These include large size (semiglobular *Halichondria* ~10 cm
across, from floating docks, have consistently proved to be this species) and color (*Ha. urca* sp.
nov. has not yet been found with any traces of green color).
It is possible that *Ha. urca* sp. nov. is introduced in some parts of its range. Its strong
association with artificial substrate is consistent with an introduced species, as is finding
samples in both the North Atlantic and North Pacific. The genetic differences seen between the
Atlantic and Pacific are intriguing, however, and it is possible that additional sampling will
reveal some of all Atlantic samples to be from an additional undescribed species.
1796

1797
1798 Figure 18. *Halichondria urca* skeleton and spicules. Tangential ectosomal skeletons: A (TLT926), B
1799 (TLT661), C (TLT534), E (TLT591), F (TLT594). D: spicules (TLT733). G: cross-section showing choanosomal
1800 skeleton at sponge surface (TLT594).

1801

1802 *Halichondria zabra* sp. nov.

1803 Figures 19 & 20

1804

Material examined

Holotype: TLT1064 (CASIZ 245222), Santa Barbara Harbor, California, (34.40559, -119.68964),
floating dock, 2021-06-01. Other samples: TLT1077 (SBMNH 718644), Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1082 (UCSB-IZC00069037),
Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1086
(UCSB-IZC00069038), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock,
2021-06-01; TLT1317, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock,
2023-09-01; TLT612 (CASIZ 245223), Jack London Square Marina, California, (37.79371, -
122.27757), floating dock, 2020-11-15; TLT623 (UCSB-IZC00069039), Jack London Square
Marina, California, (37.79371, -122.27757), floating dock, 2020-11-15; TLT624 (SBMNH
718645), Jack London Square Marina, California, (37.79371, -122.27757), floating dock, 2020-
11-15; TLT712 (UCSB-IZC00069040), Ventura Harbor, California, (34.24801, -119.26550),
floating dock, 2020-12-14; TLT722 (SBMNH 718646), Ventura Harbor, California, (34.24801, -
119.26550), floating dock, 2020-12-14; TLT737 (CASIZ 245224), Marina del Rey, California,
(33.97228, -118.44653), floating dock, 2021-01-09; TLT751 (UCSB-IZC00069041), San Diego Bay:
Shelter Island Marinas, California, (32.71094, -117.23423), floating dock, 2021-01-10; TLT756
(CASIZ 245225), Mission Bay: South Shores Boat Launch, California, (32.76406, -117.21750),
floating dock, 2021-01-10; TLT757 (SBMNH 718647), Rainbow Harbor, California, (33.76184, -
118.19111), floating dock, 2021-01-09; TLT759 (UCSB-IZC00069042), Rainbow Harbor,
California, (33.76184, -118.19111), floating dock, 2021-01-09; TLT760 (SBMNH 718648),
Rainbow Harbor, California, (33.76184, -118.19111), floating dock, 2021-01-09; TLT761 (UCSB-
IZC00069043), Rainbow Harbor, California, (33.76184, -118.19111), floating dock, 2021-01-09;
TLT780 (SBMNH 718649), San Diego Bay: Shelter Island Marinas, California, (32.71094, -
117.23423), floating dock, 2021-01-10; TLT815 (SBMNH 718650), Ventura Harbor, California,
(34.24801, -119.26550), floating dock, 2021-05-24; TLT874 (UCSB-IZC00069044), Tomales Bay
Resort & Marina, California, (38.10792, -122.86237), floating dock, 2021-02-08; TLT967 (UCSB-
IZC00069045), Ventura Harbor, California, (34.24801, -119.26550), floating dock, 2021-05-24;
TLT978 (UCSB-IZC00069046), Ventura Harbor, California, (34.24801, -119.26550), floating dock,
2021-05-24; TLT979 (SBMNH 718651), Ventura Harbor, California, (34.24801, -119.26550),
floating dock, 2021-05-24.

1835

Figure 19. *Halichondria zabra*. A (TLT1317), B (TLT1077), C (TLT978), D (TLT759), E (TLT624), F (TLT761), G (TLT1086, sponge torn to show brooded larvae inside). A is in situ, others are pictured after removal from floating docks, before preservation.

Etymology

Named for the zabra, one of the ships of the Spanish armada in the age of sail.

Diagnosis

Yellow or green *Halichondria* with a tangential ectosomal skeleton developed as a web of spicule tracts and individual spicules, with open spaces generally triangular or polygonal rather than oval. Choanosomal skeleton with spicule tracts and spicules in confusion. Spicules oxeas, 180–570 µm long, with averages per sample of 290–410 µm long and 6–12 µm wide; little-to-no difference in length in the ectosome and the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 4–5:GG. State 4:G is unique compared to all close relatives, but homoplastic with members of the Okadai group. The full sequence 4–5:GG is not shared with any other sequenced *Halichondria*. 831–832:GG. State 831:G is unique compared to all close relatives, but homoplastic with *H. dokdoensis.* The full sequence 831–832:GG is not shared with any other sequenced *Halichondria*. N=6.

ND1. 238:A. N=14.

Morphology

Samples discovered to date were yellow, green, or a mix of the two; some yellow samples were very pale and nearly white (Figure 19F). Except in thinly encrusting samples, green pigment was

more pronounced on the external surfaces of the sponge. Found as thin or thick encrustations,
all with oscula elevated to various degrees: as volcano-shape protrusions, or atop oscular
chimneys. Tendril-like projections may also be present, and these and oscular branches are
sometimes anastomosing (Figure 19C). Translucent channels sometimes present, including
those leading up one side of oscular protrusions, as often seen in other members of the
Bowerbanki group. In Santa Barbara Harbor, Southern California, brooded bright yellow larvae
were seen in one of three sponges collected in June (Figure 19G).
1872

1873
1874 Figure 20. *Halichondria zabra* skeleton and spicules. Tangential ectosomal skeletons: A (TLT751), B
1875 (TLT624), E (TLT760). C: cross-section showing choanosomal skeleton at sponge surface (TLT623), D:
1876 spicules (TLT712).
1877

1878 **Skeleton**
1879 Tangential ectosomal skeleton with a web of spicule tracts and single spicules, with open spaces
1880 fairly regular in size and tending towards triangular or polygonal shapes. Choanosomal skeleton
1881 typically halichondroid, with vague spicules tracts and spicules in confusion, without apparent
1882 spongin.
1883

Spicules
Exclusively oxeas with a typical halichondroid shape: evenly curved or with a slight central
bend, thickest in the center and tapering to sharp points at both ends. Spicules have a large
range in lengths in both the ectosome and the choanosome, with little average difference in
length between compartments. Table 9 shows dimensions for 5 samples where ectosome and
choanosome were measured separately. When all 17 samples with measurements are
combined, spicules measure 186–347–566 x 3–9–15 µm, n=1080 for length, n=912 for width.
Average values varied across samples, with means of 294 to 407 µm long and 6 to 12 µm wide.
1892

1893 Table 9. Spicule lengths of *Halichondria zabra*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT712	231–365–458 n=40	328–405–457 n=40
TLT780	234–309–364 n=40	188–294–344 n=41
TLT978	305–397–474 n=40	312–416–500 n=40
TLT1077	225–319–380 n=40	203–327–413 n=40
TLT1064	244–361–411 n=40	304–352–433 n=40

Distribution and habitat

This species is known only from floating docks in California marinas, where it was found from Tomales Bay (the most northern site investigated) to San Diego Bay (the most Southern site investigated). It was the second most common *Halichondria* on docks in Southern California, where 19 of 71 *Halichondria* (27%) were this species (sampling strategies were aimed at maximizing diversity rather than estimating relative abundance, so this number is very approximate). Only *Ha. zabra* sp. nov. and *Ha. pinaza* sp. nov. have thus far been found on floating docks in Northern California, where they were at similar abundance (with much less sampling than Southern California).

Remarks

Though this species has only been found in California thus far, we predict it will be found in additional regions as more samples are sequenced. Its strong association with floating dock habitat, where many introduced species are found, make it likely that it has been unintentionally introduced. Previously published cox1 sequences from New Hampshire, New York, Virginia, and Germany are consistent with a possible Atlantic distribution (see supplementary data), but could also be from *Ha. pinaza* (these two species cannot be differentiated at the most commonly sequenced region of this gene).

Halichondria pinaza **sp. nov.**

Figure 21

Material examined

Holotype: TLT869 (CASIZ 245217), Sausalito Yacht Harbor, California, (37.85930, -122.48044), floating dock, 2021-02-08. Other samples: NIB0603 (BELUM.Mc2024.5), Ballycastle Marina, Northern Ireland, (55.20687, -6.23962), floating dock, 2021-10-16; NIB0604 (BELUM.Mc2024.6), Ballycastle Marina, Northern Ireland, (55.20687, -6.23962), floating dock, 2021-10-16; TLT1056 (CASIZ 245216), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1080 (UCSB-IZC00069016), Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT873 (SBMNH 718629), Tomales Bay Resort & Marina, California, (38.10792, -122.86237), floating dock, 2021-02-08; TLT892 (SBMNH 718630), Home Bay, Drake's Estero, California, (38.06934, -122.91711), intertidal, 2021-02-08; WMS21x22 (YPM 111896), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal (rocks), 2021-05-26; WMS23x1 (YPM 111897), West Meadow Beach, Stony Brook, New York, (40.92673, -73.14807), intertidal (rocks), 2023-06-07.

Figure 21. *Halichondria pinaza*. A (TLT892), B (TLT1080), C (TLT869), D & E (TLT1056, E shows underside where sponge attached to dock, with brooded larvae visible). Tangential ectosomal skeletons: F (NIB0603), G (NIB0604). H: cross-section showing choanosomal skeleton at sponge surface (TLT1080), I: spicules (TLT869).

Etymology

Named for the pinaza, one of the ships of the Spanish armada in the age of sail.

Diagnosis

Yellow or green *Halichondria* with a tangential ectosomal skeleton developed as a web of multispicular tracts and individual spicules, with open spaces tending towards triangular or polygonal shapes rather than ovals. Choanosomal skeleton with spicule tracts and spicules in confusion. Spicules oxeas, ranging from 90–370 µm in length, with averages per sample of 264–308 µm in length and 6–10 µm in width; little-to-no difference in length in the ectosome and the choanosome. Diagnosis relative to similar species requires DNA characters, as follows:

COX1. No unique derived states.

28S. 4:T, 628:A, 669:T, 786:A, 803:C, 998:T. N=7–9 depending on position.

1952 ND1. No unique derived states.
1953
1954 **Morphology**
1955 All samples found to date were encrustations with protruding oscular chimneys (Figure 21).
1956 Occurs in yellow, green, or both. Tendril-like projections not seen. Translucent channels
1957 sometimes present, including those leading up one side of oscular protrusions, as often seen in
1958 other members of the Bowerbanki group. In Santa Barbara Harbor, California, brooded bright
1959 yellow larvae were seen in two sponges collected in June (Figure 21E).
1960
1961 **Skeleton**
1962 Tangential ectosomal skeleton with a web of spicule tracts and single spicules, with open spaces
1963 tending towards triangular or polygonal shapes (Figure 21F and 21G). Choanosomal skeleton
1964 typically halichondroid, with vague spicule tracts and spicules in confusion, without apparent
1965 spongin.
1966
1967 **Spicules**
1968 Nearly exclusively oxeas, though one sample (TLT892) had some aberrant spicules including
1969 styles, centrotylote oxeas, and oxeas with x-shaped cross pieces at one end. Oxeas have the
1970 typical halichondroid shape: evenly curved or with a slight central bend, thickest in the center
1971 and tapering to sharp points at both ends. Spicules have a large range in lengths in both the
1972 ectosome and the choanosome, with a very slight tendency to be longer in the choanosome
1973 (average of 3% longer across samples). Table 10 shows dimensions for 3 samples where
1974 ectosome and choanosome were measured separately. When all 6 samples with measurements
1975 are combined, spicules are 92–288–365 x 1–7–12 µm, n=432. Average values varied across
1976 samples, with means of 264 to 308 µm long, 6 to 10 µm wide.
1977
1978 **Distribution and habitat**
1979 This species is thus-far known from California, New York, and Ireland. All samples were found in
1980 sheltered bays and marinas. All were on floating docks except one sample on the underside of
1981 an intertidal boulder. It may be less tolerant of warm water than congeners such as *Ha.*
1982 *bowerbanki* and *Ha. zabra* sp. nov., as California samples were found from Drake's Estero in
1983 Northern California to Santa Barbara in Southern California, but not in sites farther South. Of all
1984 the *Halichondria* collected on floating docks in Southern California, only 2 of 71 (3%) were this
1985 species; in contrast, *Ha. zabra* sp. nov. and *Ha. pinaza* sp. nov. had similar abundance in
1986 Northern California (though sampling effort was more limited there).
1987
1988 Table 10. Spicule lengths of *Halichondria pinaza*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
Pacific Ocean		
TLT1056	158–272–311 n=55	236–281–330 n=57
Atlantic Ocean		
NIB0603	268–296–343 n=40	265–310–351 n=40
NIB0604	261–305–357 n=41	179–302–346 n=40

Remarks

This species is likely introduced in some portion of its range. First, it was found primarily on floating docks, which are known to harbor many introduced species (Bulleri and Chapman, 2010; Glasby et al., 2007; Ruiz et al., 2000; Simkanin et al., 2012). Second, samples with very similar DNA sequences were found in California, New York, and Ireland, despite this species likely having limited larval dispersal (Maldonado, 2006; Xue et al., 2009). Sequences from the cox1 locus from samples collected by others from New Hampshire, New York, Virginia, and Germany may indicate this species is also present in those areas, but these could also be *Ha. zabra* sp. nov. (these two species cannot be differentiated at the most commonly sequenced region of this gene).

Halichondria loma Turner & Lonhart 2023

Material examined

Holotype: TLT1225 (CASIZ 236654 & UCSB-IZC00048445), North Monastery Beach, California, (36.52647, -121.92730), 12-29 m, 2021-09-22. Paratypes: TLT1012, Acropolis Street (Pt. Pinos), California, (36.64183, -121.93060), 9-18 m, 2021-08-09; TLT1024, Butterfly House, California, (36.53908, -121.93520), 9-20 m, 2021-08-10; TLT1130, Fire Rock Pescadero Point, California, (36.55898, -121.95110), 10-22 m, 2021-08-10; TLT514, Goalpost, San Diego, California, (32.69438, -117.26860), 12-15 m, 2020-02-08; TLT936, Six Fathoms, San Diego, California, (32.71000, -117.26860), 9-18 m, 2021-05-15; TLT976, Wreck of the Ruby E, California, (32.76680, -117.27620), 18-25 m, 2021-05-16; TLT989, Inner Pinnacle, California, (36.55910, -121.96630), 10-18 m, 2021-08-10.

Diagnosis

Yellow *Halichondria* with a tangential ectosomal skeleton developed as a web of mulitspicular tracts and individual spicules, with open spaces tending towards triangular or polygonal shapes, or as a lace-like mat of spicules surrounding oval-shaped open areas. Choanosomal skeleton with spicule tracts and spicules in confusion. Spicules oxeas, 290–650 μm long, with averages per sample of 460–550 μm long and 5–16 μm wide. This species is diagnosed relative to *H. sitiens* by a lack of papillae, and from all other *Halichondria* examined here by having longer spicules However, this difference is slight (the longest *Ha. bowerbanki* sample averaged only 39 μm shorter), so diagnosis is facilitated by DNA characters, as follows:

COX1. No unique derived states.

28S. 5:A, 13:G, 46:G, 621:G, 674:T, 894:C, 2091:T, 2198:G. N=6–8 depending on position.

ND1. 199:C, 205:G, 209:G, 220:A, 231:G, 248:T, 250:T, 700:G, 880:A. N=7.

Remarks

This species was recently described by Turner and Lonhart (2023), and no additional samples have been discovered or examined since. Our contribution to the understanding of this species

2033 comes from DNA data: we used Illumina sequencing to increase the genetic data available for 7
2034 of the 8 previously known samples, and used that data to locate DNA characters that aid in
2035 diagnosis.
2036   This species is currently the closest known relative of the Bowerbanki group (Figure 1).
2037 In contrast to the species in that group, *Ha. loma* has not been found in sheltered bays or on
2038 human structures, and there is no indication that it may be introduced in California.
2039
2040 **Genus *Hymeniacidon* Bowerbank, 1858**
2041 Encrusting or lobate Halichondriidae with small styles (<500 μm) for megascleres. Ectosomal
2042 tangential skeleton of intercrossing bundles end single megascleres. Choanosomal skeleton
2043 with ascending vague bundles and many loose, confusedly arranged megascleres (paraphrased
2044 from Erpenback and Van Soest, 2002). Approximately 50 species are currently considered valid
2045 (de Voogd et al., 2024).
2046
2047 ***Hymeniacidon perlevis* (Montagu, 1814)**
2048 Figure 22
2049
2050 **Synonyms**
2051 *Spongia perlevis* Montagu 1814
2052 *Hymeniacidon sinapium* de Laubenfels 1930
2053 The World Porifera Database lists 27 additional synonyms (de Voogd, et al., 2024)
2054
Material examined
CASIZ 180246, Newport Bay (*Hy. sinapium* type location), California, depth not known, 2006-09-
14; NIB0611, Carnalea, Bangor, Northern Ireland, (54.66844, -5.70801), intertidal (rocky shore
crevices), 2021-10-27; NIB0615, Mount Stewart, Strangford Lough, Northern Ireland, (54.54482,
-5.61017), intertidal (boulders), 2021-04-29; NIB0616, Portaferry, Strangford Lough, Northern
Ireland, (54.38226, -5.55393), intertidal (boulders), 2021-10-10; TLT1054, Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1057, Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1059, Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1067, Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1074, Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1079, Santa Barbara Harbor,
California, (34.40559, -119.68964), floating dock, 2021-06-01; TLT1265, Zebra Cove, California,
(34.01000, -119.44000), 6-13 m, 2022-06-17; TLT129, Carpinteria Reef, California, (34.39163, -
119.54169), 3-6 m, 2019-07-31; TLT247, Carpinteria Reef, California, (34.39163, -119.54169), 3-
6 m, 2019-07-31; TLT595, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-07-17;
TLT643, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16;
TLT648, Santa Barbara Harbor, California, (34.40559, -119.68964), floating dock, 2020-02-16;
TLT651, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-07-17; TLT658, Lechuza
Point, California, (34.03437, -118.86146), intertidal, 2020-12-14; TLT730, Lechuza Point,
California, (34.03437, -118.86146), intertidal, 2020-12-14; TLT736, Marina del Rey, California,
(33.97228, -118.44653), floating dock, 2021-01-09; TLT762, Rainbow Harbor, California,
(33.76184, -118.19111), floating dock, 2021-01-09; TLT79, Tajigus, California, (34.46279, -

120.10185), 3-8 m, 2019-07-01; TLT823, Elwood Beach, California, (34.42395, -119.90604), intertidal, 2020-12-29.

Figure 22. *Hymeniacidon perlevis*. A (TLT1265), B (TLT550), C (TLT730), D (TLT1067). A-C are in situ, with A and B subtidal and C exposed at low tide. D is removed from a floating dock. E: tangential ectosomal skeleton (TLT129). F: cross-section showing choanosomal skeleton at sponge surface (TLT129). G: typical style (TLT1057).

Diagnosis

Yellow, green, orange, or red-orange *Hymeniacidon,* thinly or thickly encrusting, frequently with oscular chimneys, tendrils, or other fistulae. Ectosomal skeleton developed as a disorganized network of tangential spicule bundles, with some paratangential and protruding from the sponge surface. Choanosomal skeleton with multispicular tracts and spicules in confusion. Spicules styles; in the Northeast Pacific, they range from 110–420 µm in length, with averages per sample of 170–290 µm in length and 4–6 µm in width. Diagnosed relative to all other *Hymeniacidon* in the Northeast Pacific, except *Hy. fusiformis*, by having a large range (>100 µm) in lengths within any one sample, with some spicules in excess of 300 µm. Diagnosis relative to *Hy. fusiformis* based on thinner, non-fusiform spicules. It is not well established if this species can be diagnosed morphologically in all other regions where it occurs, so DNA confirmation of clade membership is recommended.

Morphology

Yellow, green, orange, or red-orange encrusting sponges, frequently with erect projections such as oscular chimneys and digitate tendrils. Dark red and pinkish forms reported from Europe and southern Africa (Ackers et al., 2007; Samaai et al., 2022). In silted habitats, it is often partially buried, with only vertical projections protruding from sediment. Translucent surface channels sometimes present, but it generally has a more opaque surface than *Halichondria* in the Panicea and Bowerbanki groups. Sometimes containing brooded yellow larvae.

2106

Skeleton

Ectosomal skeleton is more disorganized than *Ha. panicea*, but similar in that it is developed as a tangential mesh of spicule bundles (Figure 22E). Bundles of spicules also protrude from this mesh to create a microscopically hispid surface. Choanosomal skeleton with wispy spicule tracts and spicules in confusion (Figure 22F). No apparent spongin.

Spicules

Exclusively styles, usually with uniform thickness but a significant minority have faint swelling at or near the head (Figure 22G). When data from 18 samples from California are combined, spicules measure 116–242–415 x 2–5–9 µm, n=660 for length, n=645 for width. Mean lengths per sponge varied from 177 to 289 µm, and mean widths from 4 to 6 µm. Sponges from other regions are reported to be similar in length but sometimes thicker (Harbo et al., 2021).

Distribution and habitat

In the Northeastern Pacific this species is known from Ladysmith Harbor, British Columbia to Southern California. It is present in many other regions as well, including China, Korea, Europe, Brazil, Argentina, and South Africa (Samaai et al., 2022; Turner, 2020). Future DNA sequencing will likely confirm its presence in other regions such as Peru (Cóndor-Luján et al., 2023) and possibly New Zealand (Bergquist, 1970).

This species has a broad environmental niche, thriving in the intertidal zone, the shallow subtidal, areas with high levels of siltation, and brackish bays and estuaries. It is abundant on human structures in marinas but can also be found in relatively undisturbed natural habitats, such as those found at the Channel Islands in Southern California.

Remarks

Turner (2020) compiled genetic and morphological data from many previous studies to argue that this species has a worldwide distribution. Here, we provide additional genetic data consistent with that conclusion. A complete mitochondrial sequence was previously published from Korea (Jun et al., 2015), and we find it to be nearly identical to mitochondrial genomes from California and Ireland.

Eleven samples were collected in Santa Barbara Harbor from April to July 2021, and checked for brooded larvae. None were seen in the 10 samples collected between April 21st and June 1st, but they were present in the final sample collected July 1st.

Hymeniacidon actites **(Ristau, 1978)**

Figure 23

Synonyms

Leucophloeus actites Ristau 1978

Figure 23. *Hymeniacidon actites*. A (TLT227), B (TLT1158), C (TLT454). Spicules: D (USNM 24526). E: tangential ectosomal skeleton (TLT227). F & G: cross-sections showing choanosomal skeleton (TLT227; ectosomal skeleton is visible at the top of G).

Material examined

Holotype: USNM 24526, Horeshoe Cove, Bodega Head, California, intertidal, 1977-01-06. Other samples: RBC1, Execution Rock Cave, Bamfield, British Columbia, (48.83170, -125.17830), not recorded, 2010-08-10; TLT1158, CRABS, California, (36.55377, -121.93840), 10-17 m, 2021-09-21; TLT227, Arroyo Quemado Reef, California, (34.46775, -120.11905), 7-11 m, 2019-07-29; TLT454, Middle Reef, Whaler's Cove, Point Lobos, California, (36.52172, -121.93894), 6-15 m, 2019-11-23; TLT847, Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06; TLT856, Cave Landings, California, (35.17535, -120.72240), intertidal, 2021-02-06.

Diagnosis

Encrusting yellow *Hymeniacidon* with partially translucent ectosome and no tassels or tendril-like projections. Spicules exclusively styles 150–200 µm long, with less than 75 µm difference between shortest and longest spicules in any one sample. Well-developed sieve-like tangential ectosomal skeleton reminiscent of *Ha. panicea*. Much more genetically similar to *Ha. panicea* than other species of *Hymeniacidon.*

Morphology

Encrusting sponges, bright yellow to pale yellow-tan alive, with partially translucent ectosome. Beige or white preserved. Oscula usually apparent but small and flush with sponge surface. General morphology indistinguishable from sympatric *Halichondria* such as *Ha. californiana* sp. nov. See Figure 23 for examples.

Skeleton

2175 Tangential ectosomal skeleton generally sieve-like, with a dense mat of spicules surrounding
2176 open areas that are mostly circular or oval shaped (Figure 23E). Choanosomal skeleton with
2177 many spicules in confusion, but wavy, meandering tracts can be found in some samples. No
2178 apparent spongin.
2179

Spicules
Exclusively styles, generally straight or evenly curved. Some samples contain spicules with very
slight swelling at spicule heads or with multiple bends that make spicules somewhat sinuous.
When all samples are combined, spicules measure 139–179–237 x 2–5–8 μm, n=250. Mean
values per sample 156 to 199 μm long, 4 to 6 μm wide. Spicules from holotype: 168–196–237 x
4–6–8 μm, n=43.
2186

2187 **Distribution and habitat**
2188 Occurs in wave-exposed rocky intertidal habitats and shallow rocky reefs; not known from bays,
2189 harbors, or artificial structures. Known range previously limited to Northern and Central
2190 California (Lee et al., 2007), but is now extended North to British Columbia and South to
2191 Southern California. One sample was found on a *Loxorhynchus* decorator crab.
2192

Remarks
This species can be differentiated from *Hy. kitchingi* and *Hy. pierrei* sp. nov. due to the absence
of tendrils and tufts, and from *Hy. perlevis* by color, surface patterning, smaller spicules, and a
narrower range in spicule lengths. It is harder to distinguish from *Hy. ungodon* de Laubenfels
1932. As described below, the spicules and choanosomal skeletons in the two holotypes are
likely indistinguishable. Indeed, the British Columbian sample examined here (RBC 018-00195-
002) was previously identified as *Hy. ungodon* by William Austin. We assign this sample, and the
others we collected, to *Hy. actites* based on two apparent differences between the species:
ectosomal skeleton and color (Ristau, 1978).
Regular sieve-like ectosomal patterning was reported in the original *Hy. actites*
description and was seen in samples we examined (including the British Columbian sample).
The *Hy. actites* holotype fragment examined by us was too small to investigate the skeleton,
but the holotype was previously examined by Willard D. Hartman, and online images of his
sections show this patterning as well (Yale Peabody Museum, sample IZ.074262). In contrast,
the original description of *Hy. ungodon* states it has a rugose surface with a dense mat of
ectosomal spicules in confusion (de Laubenfels, 1932). Absence of sieve-like ectosomal
patterning in the *Hy. ungodon* holotype is not conclusive evidence of these species being
distinct, as other samples we examined that show a sieve-like pattern in some portions of the
ectosome will show a disordered mat in other places. *Hymeniacidon ungodon* is also described
as having contrasting "mahogany-brown ectosome over yellowish-drab endosome", and this
was called-out in the original description as being "characteristic and unique in the genus" (de
Laubenfels, 1932). Color is known to be highly variable, so it is unclear if this trait is a legitimate
reason to separate these species either. Genetic data from samples matching the original
description of *Hy. ungodon* are therefore needed to resolve whether these two species are truly
different, but for now, we assign these samples to *Hy. actites*.
2218

Hymeniacidon ungodon de Laubenfels 1932

Material examined

Holotype: USNM 22061, Point Lobos, Carmel, California, intertidal, 1930-07-12.

Diagnosis

Encrusting sponges with mahogany-brown ectosome over yellowish-drab endosome and no tassels or tendril-like projections. Ectosome densely packed with spicules in confusion. Spicules exclusively styles 160–230 μm in length.

Morphology

We were unable to locate any new samples that we could definitively assign to this species, and examined only a small fragment of the holotype. The original description states that they are encrusting sponges 1 cm thick, with a coarsely rugose surface; mahogany-brown ectosome over yellowish-drab endosome (de Laubenfels, 1932).

Skeleton

The original description states that the ectosome is "densely packed with spicules in confusion". Only a small piece of ectosome could be located on the holotype fragment we examined, and it was a mat of tangential spicules in confusion, matching the previous description. The choanosomal skeleton was previously described as confused with vague ascending tracts, like other species of *Hymeniacidon*.

Spicules

Exclusively styles, generally curved and of consistent thickness. 161–206–227 x 4–7–8 μm, n=50.

Distribution and habitat

The holotype is from the intertidal zone in Central California, and was the only sample vouchered by de Laubenfels; the original description states that he has "seen this species several times in collections made by students" but the locations are not stated (de Laubenfels, 1932). Samples identified by others as this species have ranged from British Columbia to Southern California, from the intertidal to the subtidal (Lee et al., 2007). We examined many of these samples; the only one from Southern California is likely *Hy. perlevis*, and the others were either assigned to *Hy. actites* or were of uncertain affinity.

Remarks

When Ristau (1978) described *Hy. actites,* he stated that it differed from *Hy. ungodon* in "coloration, growth form, and disposition of the ectosomal skeleton". We have been unable to locate any fresh samples matching the coloration described for *Hy. ungodon,* and all samples we examined had ectosomal skeletons matching *Hy. actites*. DNA was successfully amplified from only one sample previously identified as *Hy. ungodon*, from British Columbia, and it matched sequences from *Hy. actites*. Genetic data from samples matching the original description of *H. ungodon* are needed to resolve whether these two species are truly different.

2263

Hymeniacidon kichingi (Burton, 1935)

Figure 24

Figure 24. *Hymeniacidon kitchingi*. A (TLT597), B (TLT154), C (TLT412). D: cross-section showing choanosomal skeleton at sponge surface (TLT154). Spicules: E (TLT154), F (TLT491).

Material examined

TLT154, Elwood Reef, California, (34.41775, -119.90150), 9-15 m, 2019-05-15; TLT378, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2019-10-25; TLT412, Elwood Reef, California, (34.41775, -119.90150), 9-15 m, 2019-10-23; TLT424, Elwood Reef, California, (34.41775, -119.90150), 9-15 m, 2019-10-23; TLT491, Coal Oil Point, California, (34.40450, -119.87890), 7 m, 2020-07-17; TLT597, Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-07-17; TLT820, Platform Holly, California, (34.38993, -119.90656), 3-21 m, 2021-05-12; TLT941, Lazy Days, San Diego, California, (32.69415, -117.27110), 12-25 m, 2021-05-15; TLT977, Wreck of the Ruby E, California, (32.76680, -117.27620), 18-25 m, 2021-05-16.

Diagnosis

Encrusting or cushion-shaped sponges, pale yellow, bright yellow, or beige shaded with pink or purple. Surface bearing many thin tassel-like extensions: in Californian samples these have a distinctive "frayed rope" appearance that differentiates this species from all others known from the region except *Hy. pierrei* sp. nov.. Styles are the only spicule, 110–260 µm long and 1–8 µm wide. Choanosomal skeleton is typically halichondroid, with meandering spicule tracts and spicules in confusion; differentiated ectosomal skeleton lacking. Diagnosis relative to *Hy. pierrei* sp. nov. requires DNA characters, as follows:

COX1. 718:T, 742:A, 769:G785:C, 786:T, 792:C, 796:C, 820:G, 937:C, 949:A, 1011:C, 1030:C, 1099:C, 1108:A, 1186:C, 1369:A, 1372:T, 1405:A, 1423:A, 1444:G, 1447:G, 1455:G, 1480:G, 1489:A, 1492:C. N=5–8 depending on position.

28S. 190:T, 206:G, 286:G, 471:G, 525:T, 529:T, 536:A, 540:C, 552:C, 558:A, 579:A, 603:T, 637:A, 641:T, 649:C, 660:G, 682:T, 697:T, 720:T, 789:C, 793:T, 797:C, 820:G, 826:A, 852:C, 859:T. N=5–9.

ND1. Not assessed.

Morphology
Form varies from thin encrustations to semiglobular cushions. California samples were replete with tassels: digitate projections that appear frayed due to profusely branching tips. All but the smallest samples in the British Isles are also covered in thin tassels, but available images do not clearly show the "frayed ends" appearance (Ackers et al., 2007). California samples pale to bright yellow; British Isles samples greyish-beige, with a tinge of pink or purple. Translucent surface channels, pores, and small scattered oscula visible on some living samples.

Skeleton
Choanosomal skeleton typically halichondroid, with meandering multispicular tracts and spicules in confusion. No special ectosomal skeleton is apparent, though some spicules and spicule tracts are tangential in the surface of the sponge. Spicule tracts pierce the sponge surface at tassel tips. No apparent spongin.

Spicules
Thin styles, straight or slightly bent. Most are of consistent thickness until tapering to a point, but a significant minority have very slight swelling at the head, with at least one sample (TLT491) containing some tylostyles. Some are slightly thinner at the head, and therefore weakly fusiform. The original description notes that tips are telescope-like, in that they taper in a stepwise fashion. This feature was subtly present in Californian samples, but only in some spicules; a more recent publication found it to be present on only about 10% of spicules in samples from the British Isles as well (Ackers et al., 2007). When data from 7 California samples is pooled, spicules measure 114–216–260 x 1–4–8 µm, n=192, very similar to reports on European samples (Ackers et al., 2007). Mean values per sponge were 206 to 223 µm long, 4 to 6 µm wide.

Distribution and habitat
Previously known from just below the intertidal zone in the British Isles and Brittany. Samples examined here were found in 5 locations in Southern California, all subtidal, from 3 to 25 m depth. Two of these locations were artificial substrates (an oil platform and a scuttled ship), and three were natural reefs where the sponge was attached to rock.

Remarks

This species was not previously known outside of the British Isles and Brittany, so it was surprising to find samples matching its description in Southern California. California samples are genetically identical to a 2085 bp sequence of the 28S ribosomal locus that was previously published from Ireland (Morrow et al., 2012; Thacker et al., 2013). This sequence includes the hypervariable "C2" barcoding region, making it highly likely that these samples are the same species. It also seems probable that the species has experienced a human-assisted range expansion, as a natural circumboreal distribution would likely result in some genetic divergence between these distant locations. This species can therefore be added to the growing list of Halichondriidae potentially introduced by human activity.

This species is not closely related to the type species *Hy. perlevis*, and instead forms a distinct clade with *Hy. pierrei* sp. nov., *Hy. fusiformis*, an unidentified *Hymeniacidon* from the North Atlantic, and *Semisuberites cribrosa* (Figure 1). Published sequences of cox1 indicate additional samples in this clade, identified as belonging to a variety of genera, with a Hawaiian sample identified as *Hy. gracilis* being the only additional sample identified as a *Hymeniacidon* (Figure S1).

Hymeniacidon pierrei sp. nov.
Figure 25

Material examined
Holotype: TLT599 (CASIZ 245233), Coal Oil Point, California, (34.40450, -119.87890), 6.7 m, 2020-07-17. Other sample: TLT531 (SBMNH 718663), Coal Oil Point, California, (34.40450, -119.87890), 3-8 m, 2020-06-30.

Etymology
Named for Christoph Pierre, marine naturalist and Director of Marine Operations at the University of California, Santa Barbara.

Diagnosis
Encrusting *Hymeniacidon* bearing many thin tassel-like extensions. Styles are the only spicule, 100–230 µm long and 1–7 µm wide. Choanosomal skeleton is typically halichondroid, with meandering spicule tracts and spicules in confusion; differentiated ectosomal skeleton lacking. The combination of tassel-like extensions, no differentiated ectosomal skeleton, and very small, thin styles in a small size range differentiate this species for all others save *Hy. kitchingi*. Diagnosis relative to *Hy. kitchingi* requires DNA characters, as follows:

COX1. 486:C, 502:C, 598:C, 727:G, 792:A, 886:C, 1034:C. N=1.

28S. 170:T, 181:T, 182:T, 203:A, 262:T, 277:C, 278:A, 377:A, 463:G, 500:A, 512:C, 523:T, 530:T, 535:A, 551:T, 552:A, 559:T, 562:T, 603:G, 606:T, 608:C, 609:A, 611:G, 615:A, 623:A, 625:A, 642:C, 654:G, 672:T, 682:G, 704:C, 720:C, 769:A, 771:A, 778:A, 789:T, 792:A, 795:G, 796:A, 818:A, 819:C, 827:T, 841:T, 850:T, 884:G, 894:C. N=2.

ND1. Not assessed.

2377

Morphology

Thinly encrusting sponges replete with tassels: digitate projections that appear frayed due to profusely branching tips (Figure 25A & 25B). Translucent surface channels, pores, and small scattered oscula visible on living samples. Pale yellow alive, white when preserved.

Figure 25. *Hymeniacidon pierrei*. A (TLT599), B (TLT531). C: cross-section showing choanosomal skeleton in tuft (TLT531). D: tangential spicules and tracts in ectosome (TLT599). Spicules: E (TLT531).

Skeleton

Choanosomal skeleton typically halichondroid, with meandering multispicular tracts and spicules in confusion. Branching and somewhat plumose spicule tracts are well developed in tassels, where they pierce the sponge surface at tassel tips. No special ectosomal skeleton is apparent, though some spicules and spicule tracts are tangential in the ectosome. No apparent spongin.

Spicules

Thin styles, straight or slightly bent. Most are of consistent thickness until tapering to a point, but a minority have very slight swelling at the head. Like *H. kitchingi*, the tips of some spicules taper in a subtly stepwise fashion. When data from both samples is pooled, spicules measure 109–187–229 x 1–4–7 μm, n=121. Means per sponge were 183 and 199 μm in length and 4 and 5 μm in width.

Distribution and habitat

This species is only known from the shallow (3–8 m) subtidal at Coal Oil Point, Santa Barbara, California. It was found on the decaying holdfasts of dead *Macrocyctis pyrifera*.

Remarks

Hymeniacidon kitchingi was discovered in California in 2019 at Coal Oil Point and nearby Elwood Reef. An additional tasseled *Hymeniacidon* was collected on a return trip to Coal Oil Point in 2020, but when it was sequenced, it was very genetically distinct from *Hy. kitchingi*. An additional trip later in 2020 yielded both genotypes, collected side-by-side. With 13% sequence divergence (Fst = 0.97) at the 28S locus and 10% sequence divergence (Fst = 0.32) at the cox1 locus, these sequences are clearly from a different species. Indeed, at cox1, *Hy. pierrei* sp. nov. are more closely related to samples collected in Hawaii (Genbank accession KY565326, identified as *Hy. gracilis*) and Australia (Genbank accession KJ620398, unidentified) than to *Hy. kitchingi.*

It is unclear whether *H. pierrei* sp. nov. is native to California or introduced. It was found on a natural reef that is now part of a marine protected area, but this site was previously highly disturbed, as it housed a pier and oil derricks as part of the Elwood Oil Field. This area is one of two places that the (possibly introduced) *Hy. kitchingi* has been found, so it seems possible that *Hy. pierrei* sp. nov. is introduced as well.

Hymeniacidon fusiformis Turner & Lonhart 2023

Material examined
Holotype: TLT1194 (CASIZ236657 & UCSB-IZC00048452), Inner Carmel Pinnacle, Carmel, California, (36.55852, -121.96820), 10–24 m, 2021-09-22. Additional sample: FH12x28 (YPM 111899), Bell Island (West Face), Washington, (48.59667, -122.98150), 15 m, 2012-05-18.

Diagnosis
Thickly encrusting white or yellow sponges, sometimes with irregularly projecting fingers and lobes, but not thin tentacle-like extensions or tassels. Styles are the only spicule, slightly fusiform, 150–410 µm in length and 4–19 µm in width. Maximum style length and large range in spicule lengths differentiate this species from sympatric *Hymeniacidon* except *Hy. perlevis*, while the wide range in spicule widths is unique in this geographic region. Choanosomal skeleton is typically halichondroid, with spicule tracts and spicules in confusion. The ectosomal skeleton is developed as a sieve-like mesh surrounding triangular open spaces or as a dense unstructured mat of tangential spicules.

Spicules
Exclusively styles; most are very slightly fusiform, so that they are thicker in the center than near the head of style. Length and width are both highly variable; distributions are continuous but strongly bimodal, despite not showing any differences in size in the ecotosome vs. the choanosome. Spicules in the holotype measure 186–281–409 x 4–10–18 µm (n=345), with length modes 210 and 350 µm and width modes 9 and 14 µm. The new sample from Washington is similar, 153–270–396 x 6–11–19 µm (n=80), with length modes 260 and 330 µm and width modes 8 and 13 µm.

Distribution and habitat

2448 Known from two samples, the first from a natural subtidal pinnacle in Carmel Bay, Central
2449 California, 10–24 m depth, the second from a submerged rock wall, Bell Island, Washington, 15
2450 m depth.
2451

Remarks
This species was recently described by Turner & Lonhart (2023) from a single sample. Here we
report additional genetic data from this sample. We used Illumina sequencing to completely
sequence the nuclear ribosomal locus; coverage was insufficient to assemble the mitochondrial
genome, but we were able to extract 777 bp of the cox1 locus. We also report the discovery of
one additional sample. This new sample conforms well to the morphological description of the
holotype, and is only 0.4% divergent at the 28S locus.
The closest relative to this species in our molecular phylogenies is a *Hymeniacidon* sp.
we collected and sequenced from the Bay of Fundy, in the Canadian North Atlantic (sample
Quoddy115 in Figures 1 & 3). This sample is 1.2% divergent at the cox1 locus and 0.4%
divergent at the 28S locus when compared to Pacific samples, which could indicate a species-
level difference or not. However, it is also has much shorter and thinner spicules that are not
fusiform, and these combined morphological and genetic differences indicate it is likely a
different, possibly undescribed, species. These species fall in a larger clade including *Hy.*
kitchingi, *Hy. pierrei* sp. nov., *S. cribrosa* and samples from Genbank identified as belonging to a
variety of other genera (Figure S1).
2468

2469 ***Hymeniacidon globularis* Ott, McDaniel & Humphrey, 2024**
2470

2471 **Material examined**
2472 None
2473

2474 **Diagnosis**
2475 Light yellow globular sponges with a prominent central oscule; subtylostyles with elliptical
2476 heads are the only spicule. Distinguished from all other *Hymeniacidon* in the region by
2477 "puffball" shape, large styles (300–450 µm long and 5–13 µm wide), and having elliptical
2478 subtylostyles as the primary spicule shape.
2479

2480 **Distribution and habitat**
2481 Known only from the subtidal, 17-24 m, in British Columbia.
2482

2483 **Remarks**
2484 This species was recently described by Ott, McDaniel & Humphrey (Montagu 1814), and is
2485 included here for completeness. See their description for additional details.
2486

Genus *Semisuberites* Carter, 1877
Funnel-shaped sponges arising from a stalk. Choanosomal skeleton of branching spicule tracts
run mostly parallel to the sponge surface, but give rise to tracts that branch off to intersect the
sponge surface. These flare to form bouquets that pierce the surface of the sponge. Some

spicules are also confused throughout the choanosome and tangentially arranged within the
ectosome. Spicules styles and subtylostyles. Genus is monotypic.
2493
2494   ***Semisuberites cribrosa* (Miklucho-Maclay, 1870)**
2495   Figure 26
2496
2497   **Synonyms**
2498   *Veluspa polymorpha* var. *cribrosa* Miklucho-Maclay, 1870
2499   *Semisuberites arctica* Carter, 1877
2500   *Stylissa stipitata* de Laubenfels, 1961
2501   The World Porifera Database lists 22 additional synonyms (de Voogd, et al., 2024)
2502

2503
2504   Figure 26. *Semisuberites cribrosa*. A (BHAK-08809), B (BHAK-09371). C: cross-section showing
2505   choanosomal skeleton near sponge rim, with external surface on the left and internal surface on the
2506   right; round structures in center are cross-sections of spicule tracts (BHAK-08809). Spicules: E (BHAK-
2507   10379).
2508
2509   **Material examined**
2510   BHAK-08809, Nalau Pass, British Columbia, (51.784, -128.101), 30 m, 2019-05-25; BHAK-09371,
2511   Wedgeborough Point, British Columbia, (51.647, -127.956), 20 m, 2019-05-27; BHAK-10379,
2512   Wedgeborough Point, British Columbia, (51.647, -127.956), 25 m, 2019-05-27.
2513
2514   **Diagnosis**
2515   Soft, fragile, funnel-shaped sponges arising from a stalk. Styles are the only spicule, slightly
2516   fusiform, 110–450 µm long 1–12 µm wide (styles reportedly up to 600 µm long and 16 µm wide
2517   in other regions). Prominent spicule tracts run parallel to the surface of the sponge, branching
2518   off to intersect the sponge surface, where they form spicule brushes in the ectosome.
2519
2520   **Morphology**
2521   Funnel or trumpet-shaped sponges arising from stalks that attach to the substrate with root-
2522   like structures (Figure 26B). Stalk may branch and give rise to multiple funnels (Figure 26A).
2523   Sponges soft and fragile, with distal lip often frayed (Figure 26B). Velvety surface. Tan alive,
2524   white in ethanol.
2525
2526   **Skeleton**

2527   Branching spicule tracts run mostly parallel to the sponge surface, but give rise to tracts that
2528   branch off to intersect the sponge surface. These flare to form bouquets that pierce the surface
2529   of the sponge. Some spicules are also confused throughout the choanosome and tangentially
2530   arranged within the ectosome (Figure 26C).
2531
Spicules
Nearly exclusively styles (Figure 26BD), straight or evenly curved, some subtylote but many
simple; rare strongyles occasionally seen. Distributions of length were bimodal or trimodal for
each sponge, but continuous with unclear size classes. When spicules from three samples are
combined, they measure 118–285–446 x 1–8–12 µm, n=260 for length, n=253 for width. The
combined length distribution is trimodal, with modes at approximately 210, 310, and 390 µm.
Means per sponge were 259 to 301 µm in length and 7 to 8 µm in width.
2539
2540   **Distribution and habitat**
2541   Known from the Arctic, North Atlantic, and North Pacific, from 20–289 m depth (Dinn et al.,
2542   2020). The Southernmost Atlantic specimens are known from the Gulf of St. Lawrence (Dinn et
2543   al., 2020), while the Southernmost Pacific specimens are known from the San Juan Islands,
2544   Washington (de Laubenfels, 1961).
2545
Remarks
The choanosomal skeletons of halichondriid sponges are often described as "confused", but this
term can also be applied to the taxonomy of many species in this group. This species is no
exception. Populations of this species have been described as members of various families and
genera; in the Northeast Pacific, these sponges were first described as *Stylissa stipitata* (de
Laubenfels, 1961); it now seems likely that this name is a junior synonym of *Semisuberites*
cribrosa. The samples we examined here agree with the recent morphological description of
samples from the North Atlantic, though Atlantic samples had slightly larger spicules (Dinn et
al., 2020).
The genus *Semisuberites* is monotypic and has been difficult to place. It was assigned to
the Suberitidae by Topsent (1928); Van Soest and Hajdu (2002) noted similarities with
Hymeniacidon but tentatively placed the genus in the Esperiopsidae (Order Poecilosclerida).
Van Soest (2016) later proposed that this genus is more likely to be in the order Suberitida.
Here we present the first genetic data for this species, which clearly places it within the
Suberitida. Based on morphological data and genetic similarity to some *Hymeniacidon* species,
we propose placement in the Halichondriidae. The upright brushes of styles in the ectosome
are unusual for this family, but there do appear to be some tangential spicules in the ectosome
as well.
2564
Genus Topsentia Berg, 1899
Halichondriidae with a densely confused choanosomal skeleton and a crust-like tangential or
paratangential arrangement of spicules in the ectosome. The ectosome is difficult to detach
due to limited subdermal cavities. Spicules oxeas or modifications, in a large size range,
including smaller spicules concentrated at the surface (paraphrased from Erpenback and Van
Soest, 2002). Approximately 36 species are currently considered valid (de Voogd et al., 2024).

2571

2572 ***Topsentia fernaldi* (Sim and Bakus, 1986)**

2573 Figure 27

2574

2575 **Synonyms**

2576 *Oxeostilon fernaldi* Sim & Bakus, 1986

2577

2578 **Material examined**

2579 Holotype: USNM 33629, Bird Rock, Santa Catalina Island, California, (33.45000, -118.41667), 46

2580 m, 1968-11-20. Other samples: TLT1134, Honeymooners, California, (36.50390, -121.94100),

2581 10-20 m, 2021-09-22; TLT1261, Honeymooners, California, (36.50390, -121.94100, 10-20 m,

2582 2021-09-22.

2583

Diagnosis

Thickly encrusting, irregular masses, yellow or chocolate brown exteriorly with yellow interiors.

Sponges are hard and incompressible due to dense spiculation. The ectosomal skeleton is a

crust of tangential spicules in a sieve-like pattern or a disordered mat. The choanosome is very

dense with disordered spicules, with many oriented vertically near the surface, their tips

piercing the ectosome. Spicules are oxeas in a large size range, 160–1170 µm long and 4–35 µm

wide. Distribution of length is strongly bimodal, with modes near 250 and 850 µm. Spicules

average 50% shorter in the ectosome, where most spicules are less than 600 µm long, but the

full size-range is present in the choanosome.

2593

2594
2595 Figure 27. *Topsentia fernaldi*. A (TLT1134), B (TLT1261). C & D: tangential ectosomal skeleton (TLT1261).

2596 Cross-sections show choanosomal skeleton at sponge surface (E) and farther towards the interior (F),

2597 TLT1261. G: long and short oxeas from holotype, USNM 33629.

2598

2599 **Morphology**

The holotype was previously described as a thickly encrusting yellow mass, firm but friable, with a crusty surface texture and sparse oscula. Newly collected samples were thick, irregular encrustations, chocolate brown on the surface and pale yellow interiorly. Their interiors were full of bryozoan skeletons and worm tubes which are presumed to have been overgrown by the sponge, and the exterior was heavily fouled with hydroids, bryozoans, and other organisms. Oscula were few, small, scattered, and flush with surface. Samples are white in ethanol.

Skeleton

The original description states only that spicules are "distributed in an irregular fashion". We examined only a small portion of the holotype, and did not locate ectosomal regions. The choanosomal skeleton was confused, with a higher density of spicules than *Ha. panicea*, but a lower density than the recently collected samples from Central California.

Freshly collected samples from Central California had tangential ectosomal skeletons developed in some places as a sieve-like pattern around open spaces that were oval or polygonal, and in other places as a dense mat of discorded spicules. The choanosome was densely populated with spicules in confusion. Spicules became somewhat less confused towards the ectosome, where the majority are oriented vertically, with the final ones piercing the surface to create microscopic hispidity. No apparent spongin.

Spicules

Nearly exclusively curved oxeas, with rare styles or strongyles sometimes present. Distributions of length were strongly bimodal, but continuous distributions made it difficult to define size classes. Spicules from the holotype measure 224–568–1129 x 7–15–31 µm, n=115; length modes at about 250 µm and 850 µm. When data from the holotype are combined with the two newly collected samples, spicules measure 168–511–1161 x 4–15–35 µm, n=499. See table 11 for data from each sample individually. When spicules are isolated from the ectosome and choanosome separately, the full range of lengths is seen in the choanosome, but ectosomal spicules are mostly from the shorter half of the length distribution (ectosome mean length 330 µm vs. choanosome mean length 614 µm).

The holotype was originally described as having small oxeas, large oxeas, styles, and strongyles, with "styles and especially strongyles uncommon". Our spicule preparation from the holotype found only 2 styles, one tylostyle, and no strongyles, out of several hundred spicules, so we believe these to be rare aberrations (one style was also seen in another sample). We also do not consider oxeas to be present in size classes, because the length distribution is continuous, but the modes of the length distribution are very similar to the means of the size classes previously reported (239 and 813 µm).

Table 11. Spicule lengths of *Topsentia fernaldi*. Values given as min–mean–max in µm.

Specimen	Ectosome	Choanosome
TLT1134	177–406–880 n=45	218–648–1021 n=46
TLT1261	168–282–1161 n=70	194–592–974 n=70
Topsentia aff. *fibrosa*:		
RBC10	288–463–977 n=95	300–667–1020 n=72

Distribution and habitat

Described from a single sample collected at Catalina Island, in Southern California, at 46 m. Two additional samples were recently collected in Carmel Bay in Central California, between 10-20 m. Both locations were natural rocky reefs.

Remarks

The two sponges collected from Central California fit well into the current concept of *Topsentia*: they have a well-developed tangential ectosomal spicule crust that is difficult to remove, a densely confused choanosomal skeleton, and oxeas in a large size range, with smaller spicules at the sponge surface (Erpenbeck and van Soest, 2002). However, we have reservations in assigning them to *T. fernaldi.* The original description for this species is sparse, but it was described as "moderately dense, firm but friable", rather than stony and incompressible, as the newly collected samples are. The small fragment of the holotype we examined did appear to be less densely spiculated, consistent with this difference. However, the spicules from the new samples are very similar to the *T. fernaldi* holotype, and DNA extractions from the holotype were not successful. We therefore assign the new samples to this species, but would not be surprised that future collections yield new material that would result in splitting them.

We also examined a sponge from British Columbia, previously identified by William Austin as *Topsentia* aff. *fibrosa*. This sponge was morphologically indistinguishable from the Central California samples (table 11), but DNA extractions were again unsuccessful. It is therefore possible that *T. fernaldi* ranges farther North, but additional sampling of this genus in the Northeast Pacific is badly needed. *Topsentia fibrosa* is an Arctic species that may range into British Columbia, but has not been confirmed there. The original description of *T. fibrosa* is limited, but does state that samples are dark brown after preservation in spirit (Fristedt, 1887), while all the samples examined here were white after preservation in ethanol.

The presence of *T. fernaldi* within the Panicea species group is very surprising (Figure 1). While much more distantly related species are morphologically indistinguishable from *Ha. panicea*, the spicule dimensions and density make this species quite distinct. A few other *Topsentia* species have genetic data available, and some appear to fall within entirely different orders of demosponges (Pankey et al., 2022; Redmond et al., 2013). These results serve to emphasize how evolutionary liable spicule size, shape, and density are in this clade.

***Topsentia disparilis* (Lambe, 1893)**

Synonyms

Halichondria disparilis Lambe, 1893

Material examined

None.

Description

First described from a single massive, flat sponge, 35 x 28 mm across and 11 mm thick. Spicules are exclusively or nearly exclusively oxeas, but in a very wide range of lengths, with two size-

classes (or perhaps modes in a continuous distribution). Large oxeas originally described as
438–1287 µm long, averaging 13 µm thick (Lambe, 1893); a later report states they are 900 x 20
µm (de Laubenfels, 1953). Short oxeas reported as 130 x 8 µm (de Laubenfels, 1953) or 91 x 4
(Lambe, 1893). Choanosomal skeleton densely confused; ectosomal skeleton variously reported
as a tangential reticulation of oxeas (Lambe, 1893) or bristling with erect spicules (de
Laubenfels, 1953). The latter report of this species also notes it has scattered areas covered in
low, small tubercles less than 1 mm high (de Laubenfels, 1953).
2689
2690 **Distribution and habitat**
2691 The holotype was collected at 73 m near Comox, Vancouver Island, British Columbia, and
2692 additional samples were reported from 38–45 m near Point Barrow, AK (de Laubenfels, 1953).
2693
Remarks
We did not examine any material from this species, but it is included here for completeness.
Published descriptions do not clearly differentiate this species from *T. fibrosa* (Fristedt, 1887),
which is also known from the Arctic, and may be a senior synonym. As stated in the *T. fernaldi*
section above, further revision of *Topsentia* diversity in the Northeast Pacific is badly needed,
but likely requires freshly collected material and additional DNA data.
2700
2701 **Genus *Axinyssa* Lendenfeld, 1897**
2702 Halichondriidae lacking an ectosomal tangential skeleton. Choanosomal skeleton confused
2703 interiorly but with bundles near the periphery that protrude and form surface conules.
2704 (paraphrased from Erpenback and Van Soest, 2002). Approximately 28 species are currently
2705 considered valid (de Voogd et al., 2024).
2706
2707 *Axinyssa piloerecta* **sp. nov.**
2708 Figure 28
2709
2710 **Material examined**
2711 Holotype: TLT1007 (CASIZ 245235 & UCSB-IZC00069001) and additional sample TLT1132
2712 (SBMNH 718614), both collected at Yellowbanks, Santa Cruz Island (33.99432, -119.51910), 24-
2713 30 m, 2021-09-15.
2714
2715 **Etymology**
2716 From the word piloerection, the process that causes skin to form "goosebumps" and hair to
2717 stand up, creating a pattern similar to the surface of this sponge.
2718

Figure 28. *Axinyssa piloerecta*. Sample shown in situ (A) and cross-sectioned after collection, before preservation (B). C: sample TLT1132 after collection, before preservation. Cross-sections show choanosomal skeleton at sponge surface (D) and farther towards the interior (E). F: example oxeas, G: example style. All photos from TLT1007 except C.

Diagnosis

Yellow cushion-shaped sponges with widely-spaced surface conules. The combination of a smooth, slick-looking surface and conules with protruding spicule bundles is distinctive in the region. This contrasts with the sympatric *Obruta collector* Turner & Lonhart 2024, which has similar spicules but lacks raised bumps and is hispid over the entire surface. The large oxeas (530–1480 x 5–33 µm) and the significant fraction of spicules that are styles (18%) further distinguish this species from others in the region.

Morphology

Cushion-shaped irregular mounds, pale yellow when alive and beige in ethanol. The surface is covered in raised projections, 1–3 mm apart, 1–3 mm in height, with spicules projecting an additional 0.5 mm. When alive, the holotype was approximately 60 x 35 mm across, 24 mm in maximum thickness; sampled portion approximately 35 x 35 x 24 mm. A distinct surface membrane covers the sponge, creating a smooth and slick-looking surface except where spicules protrude from raised conules.

Skeleton

A dense, confused mass of spicules internally, with no apparent organization. Spicules become roughly parallel towards the periphery, pointing outwards, and form vague plumose tracts. Some of these tracts end in tufts of spicules that pierce the surface of the sponge. No ectosomal skeleton. Spongin not apparent, but body of sponge is collagenous and dissolves very slowly in bleach, in contrast to all other Halichondridae described here, which dissolved very rapidly.

Spicules

Styles and oxeas, with some intermediates; one strongyle also present. Oxeas are curved or bent, usually near center, sometimes with weak centrotylote swelling (Figure 28F). Dimensions highly variable, but without clear size categories; 533–957–1476 x 5–24–33 µm (n=78). Styles are less common than oxeas (18% of spicules), but with similar dimensions. 606–1178–2166 x 5–17–27 µm (n=17).

Distribution and habitat

Found at only a single location: a rocky reef near Yellowbanks, Santa Cruz Island. The site is a rocky outcrop, typical of other rocky reefs in the region, but most other sites searched in the Northern Channel Islands have been in less than 20 m of water, while this site was 24-30 m in depth. The species appeared to be common at the site at the time of collection.

Remarks

The morphology of this species places it firmly within the *Axinyssa* (Erpenbeck and van Soest, 2002). Like the genus *Topsentia*, the genus *Axinyssa* is highly polyphyletic, with some species, including the type species, falling among the order Bubarida and other species falling in the order Biemnida (Morrow and Cárdenas, 2015; Morrow et al., 2013; Pankey et al., 2022; Thacker et al., 2013). Genetic data from this species was previously reported as placing it clearly in the order Axinellida (Turner and Pankey, 2023). We describe it here because in the current taxonomy, this genus remains within the Halichondriidae, but we expect future taxonomic revision to move this species to the Axinellida.

Two species of *Axinyssa* are known from the North Pacific: *A. tuscara* (Ristau, 1978) and *A. isabela* (Carballo and Cruz-Barraza, 2008). The new species is distinguished from these species by its long spicules and high fraction of spicules that are styles. These same features distinguish it from species known from the temperate Atlantic.

Axinyssa tuscara **(Ristau, 1978)**

Synonyms

Axinomimus tuscarus Ristau, 1978

Material examined

Holotype: USNM 24527, Carmel River Beach Point, Carmel, California, 7 m, 1973-04-10.

Diagnosis

Thickly encrusting dark brown sponges that remain brown after preservation in ethanol. Soft and compressible. Choanosomal skeleton of vertical spicule tracts that pierce sponge and project up to 1.5 mm beyond sponge surface, connected by a confused reticulation of individual spicules. No special ectosomal skeleton. Spicules oxeas only, longer and considerably thicker than sympatric *Halichondria*, 355–552 x 10–22 µm.

Morphology

Previously described as dark brown exteriorly and dark chocolate brown interiorly; thickly encrusting and amorphous; soft and compressable. The small piece of the holotype we examined was consistent with this description, though we were unable to compare ectosome and choanosome.

Skeleton
Previously described as having vauge and somewhat plumose spicule tracts, with spicules and spicule bundles protruding through the ectosome. Lacking a special ectosomal skeleton. In the small piece we examined, and could only determine that the chromosomal skeleton was confused.

Spicules
Curved oxeas, 355–476–552 x 10–18–22 µm, n=47.

Distribution and habitat
Collected in situ only from the type location: Carmel River Beach Point, Central California, on a rock reef at 7 m depth. It was described as rare at this location (Ristau, 1978). Several samples were also collected in beach wrack in Northern California at Bodega Head.

Remarks
All known samples of this species were collected from 1973 to 1977 and were included in the original description (Ristau, 1978). We examined a small piece of the holotype in order to attempt DNA extraction, which failed. We also performed a spicule preparation in order to publish more detailed information on the distribution of spicule sizes, as described above. It was difficult to characterize the skeleton in the small piece we examined, and could only determine that the chromosomal skeleton was confused.

Conclusions
Here, we have documented a remarkable diversity of sympatric sponges in the family Halichondriidae. We combined extensive field collections, genomic analysis, and traditional morphological characters to reject both the "single globally-distributed invasive species" and "cryptic endemic species in different regions" hypotheses. We reject the single-species hypothesis by using multi-locus genomic data to infer reproductive isolation between sympatric populations. When molecular phylogenies are built separately for mitochondrial genomes and the nuclear-encoded ribosomal locus, we find numerous clades with reciprocal monophyly in both compartments. Because the mitochondrial and nuclear genomes are unlinked, reciprocal monophyly across these compartments is strong evidence of reproductive isolation when species are sympatric (Jennings, 1917; Rannala and Yang, 2020). We are also able to reject the "cryptic endemic species in different regions" hypothesis for many (but not all) of the delineated species because we find a lack of correspondence between geographic region and species range. Instead, we find numerous species that have likely been dispersed by human activity.

The discovery of cryptic species in this group is not entirely surprising, given the few known morphological traits that distinguish them. It is also not surprising that some species

have been spread through accidental human introductions, particularly since they are common
on floating docks and other areas where introduced species have been observed, and other
species in the family are known to be introduced (Carlton and Eldredge, 2009; Turner, 2020).
However, we did not expect to find so many cryptic introduced species, sympatrically
distributed at even small scales. *Halichondria bowerbanki*, *Ha. zabra* sp. nov., *Ha. urca* sp. nov.,
Ha. pinaza sp. nov., and *Ha. hygeia* sp. nov. were all found growing on the same floating docks
in a small harbor in Santa Barbara, California, and we document at least 4 of these species in
Eastern North America and/or Europe. We hope that these unintentional "common gardens" of
related sponges will facilitate future studies of their evolution and ecology.
We were unable to find morphological traits to diagnose many of these species. This
presents challenges for ecological monitoring projects that lack genotyping capabilities.
However, accurately delineating reproductively isolated groups is essential for monitoring both
native diversity and potentially invasive species. Additionally, biologists studying physiology,
biochemistry, or development need to know which species they are examining so that others
can build on their findings. Despite their morphological similarities, we suspect that many of
these species will prove to be biologically distinct. For example, *Halichondria panicea* and *Ha.*
bowerbanki are estimated to share a common ancestor in the mid-Cretaceous period, nearly
100 million years ago (Pankey et al., 2022). Even more dramatically, *Halichondria panicea* and
Ha. galea sp. nov. have been independently evolving since the Jurassic (Pankey et al., 2022),
splitting around the same time as the common ancestor of all Eutherian mammals (Archibald,
2003; Luo et al., 2011). While the Eutherian clade was diversifying into bats, whales, and
primates, what were these sponges up to? We hope that our results will motivate future work
on this question.
2859

Acknowledgements
We are grateful for the help and support of many people in UCSB's Marine Science Institute and
Diving & Boating Program, especially Robert Miller, Clint Nelson, Christoph Pierre, and Christian
Orsini. Numerous NOAA personnel provided logistics and planning support for collections in the
Monterey Bay and Olympic Coast National Marine Sanctuaries. Raquel Pereira graciously
shared samples from Sweden and Portugal, and Brooke Weigel shared samples from Tatoosh,
Washington. Gustav Paulay, Hugh MacIntosh, Abigail Reft and Christina Piotrowski provided
access to their collections at the Florida Museum of Natural History, Royal British Columbia
Museum, Smithsonian National Museum of Natural History, California Academy of Sciences,
respectively. We also thank Matt Whalen, Margot Hessing-Lewis, and the staff, students and
volunteers of the Hakai Institute for support during the 2017 Hakai-MarineGeo bioblitz, and the
Makah Tribal Nation for supporting the study of sponges from Tatoosh Island. We thank DFO
colleagues Andrew Cooper, Peter Lawton, Laura Teed and Torben Brydges from Saint Andrews
Biological Station, who partnered on the Eastern Shores habitat survey project and participated
in specimen collection.
2875
Funding
Financial support was provided by UCSB and by the National Aeronautics and Space
Administration Biodiversity and Ecological Forecasting Program (Grant NNx14Ar62A); the
Bureau of Ocean Energy Management Environmental Studies Program (BOEM Agreement

MC15AC00006); the National Oceanic and Atmospheric Administration (NOAA) in support of
the Santa Barbara Channel Marine Biodiversity observation Network; and the U.S. National
Science Foundation (NSF) in support of the Santa Barbara Coastal Long Term Ecological
research program under Awards OCE-9982105, OCE-0620276, OCE-1232779, OCE-1831937, and
the Tula Foundation. Additional support was provided by the NSF under award EF-2025121 to
RWT and funding from the Department of Agriculture Environment and Rural Affairs
Environment Fund to CM. Funding for collection of specimens from the Eastern Shore Islands,
Nova Scotia was covered as part of a habitat and sponge biodiversity survey funded by Fisheries
and Oceans Canada. The funders had no role in study design, data collection and analysis,
decision to publish, or preparation of the manuscript.

Supporting Information

Data available from the Dryad Digital Repository (https://doi.org/10.5061/dryad.bvq83bkk6).

References

Ackers, R.G., Moss, D., Picton, B.E., Stone, S.M.K., Morrow, C.C., 2007. Sponges of the British Isles: A color guide and working document. Marine Conservation Society, Ross-on-Wye.

Agne, S., Ekins, M., Galitz, A., Hofreiter, M., Preick, M., Straube, N., Wörheide, G., Erpenbeck, D., 2022. Keratose sponge MuseOMICS: setting reference points in dictyoceratid demosponge phylogeny. Zootaxa 5195, 296–300. https://doi.org/10.11646/zootaxa.5195.3.9

Ahrens, D., Ahyong, S.T., Ballerio, A., Barclay, M.V.L., Eberle, J., Espeland, M., Huber, B.A., Mengual, X., Pacheco, T.L., Peters, R.S., Rulik, B., Vaz-DE-Mello, F., Wesener, T., Krell, F.-T., 2021. Is it time to describe new species without diagnoses? A comment on Sharkey et al. (2021). Zootaxa 5027, 151–159. https://doi.org/10.11646/zootaxa.5027.2.1

Archibald, D.J., 2003. Timing and biogeography of the eutherian radiation: fossils and molecules compared. Mol. Phylogenet. Evol. 28, 350–359. https://doi.org/10.1016/S1055-7903(03)00034-4

Ballard, J.W.O., Whitlock, M.C., 2004. The incomplete natural history of mitochondria. Mol. Ecol. 13, 729–744. https://doi.org/10.1046/j.1365-294X.2003.02063.x

Barthel, D., 1986. On the ecophysiology of the sponge Halichondria panicea in Kiel Bight. I. Substrate specificity, growth and reproduction. Mar. Ecol. Prog. Ser. 32, 291–298. https://doi.org/10.3354/meps032291

Barthel, D., Detmer, A., 1990. The spermatogenesis of Halichondria panicea (Porifera, Demospongiae). Zoomorphology 110, 9–15. https://doi.org/10.1007/BF01632807

Barthel, D., Wolfrath, B., 1989. Tissue sloughing in the sponge Halichondria panicea: a fouling organism prevents being fouled. Oecologia 78, 357–360. https://doi.org/10.1007/BF00379109

Bergquist, P.R., 1970. The marine fauna of New Zealand: Porifera, Demospongiae, part 2 (Axinellida and Halichondrida). N. Z. Oceanogr. Inst. Mem. 51, 9–85.

Bowerbank, J.S., 1866. A Monograph of the British Spongiadae. Volume 2. Ray Society, London.

Bulleri, F., Chapman, M.G., 2010. The introduction of coastal infrastructure as a driver of
change in marine environments. J. Appl. Ecol. 47, 26–35. https://doi.org/10.1111/j.1365-
2664.2009.01751.x
Burton, M., 1935. Notes on British sponges, with a description of a new genus and species. Ann.
Mag. Nat. Hist. 10, 651–653.
Burton, M., 1930. Additions to the sponge fauna at Plymouth. J. Mar. Biol. Assoc. U. K. 16,
489–507.
Carballo, J.L., Cruz-Barraza, J.A., 2008. First record of Axinyssa Lendenfeld, 1897
(Demospongiae, Halichondrida) from the East Pacific Ocean, with the description of
Axinyssa isabela sp. nov. Zootaxa 1784, 58–68.
https://doi.org/10.11646/zootaxa.1784.1.4
Carlton, J.T., Eldredge, L.G., 2009. Marine bioinvasions of Hawai'i: the introduced and
cryptogenic marine and estuarine animals and plants of the Hawaiian Archipelago,
Bishop Museum bulletin in cultural and environmental studies. Bernice P. Bishop
Museum/Bishop Museum Press, Honolulu, Hawai'i.
Carrier, T.J., Schmittmann, L., Jung, S., Pita, L., Hentschel, U., 2023. Maternal provisioning of
an obligate symbiont in a sponge. Ecol. Evol. 13, e10012.
https://doi.org/10.1002/ece3.10012
Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34, i884–i890. https://doi.org/10.1093/bioinformatics/bty560
Cóndor-Luján, B., Arteaga, A., Polo, C., Arroyo, Y., Willenz, P., Hajdu, E., 2023. Shallow
Suberitida (Porifera, Demospongiae) from Peru. Zootaxa 5264, 451–489.
https://doi.org/10.11646/zootaxa.5264.4.1
Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and
BCFtools. GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008
de Laubenfels, M.W., 1961. Porifera of Friday Harbor and Vicinity. Pac. Sci. 15, 192–202.
de Laubenfels, M.W., 1953. Sponges of the Alaska Arctic. Smithson. Misc. Collect. 121, 1–21.
de Laubenfels, M.W., 1932. The marine and fresh-water sponges of California. Proc. U. S. Natl.
Mus. 81, 1–140.
de Laubenfels, M.W., 1930. The Sponges of California. Stanf. Univ. Bull., Abstracts of
dissertations for the degree of doctor of philosophy 5, 24–29.
de Paula, T.S., Zilberberg, C., Hajdu, E., Lôbo-Hajdu, G., 2012. Morphology and molecules on
opposite sides of the diversity gradient: Four cryptic species of the Cliona celata
(Porifera, Demospongiae) complex in South America revealed by mitochondrial and
nuclear markers. Mol. Phylogenet. Evol. 62, 529–541.
https://doi.org/10.1016/j.ympev.2011.11.001
de Voogd, N.J., Alvarez, B., Boury-Esnault, N., Cárdenas, P., Díaz, C., Dohrmann, M., Downey,
R., Goodwin, C., Hajdu, E., Hooper, J.N.A., Kelly, M., Klautau, M., Lim, S.C., Monconi,
R., Morrow, C.C., Pinheiro, U., Pisera, A.B., Ríos, P., Rützler, K., Schönberg, C., Turner,
T.L., Vacelet, J., van Soest, R.W.M., Xavier, J., 2024. World Porifera Database [WWW
Document]. URL https://www.marinespecies.org/porifera on 2023-01-03.
doi:10.14284/359 (accessed 1.1.24).
Delić, T., Trontelj, P., Rendoš, M., Fišer, C., 2017. The importance of naming cryptic species
and the conservation of endemic subterranean amphipods. Sci. Rep. 7.

Dinn, C., Leys, S.P., Roussel, M., Méthé, D., 2020. Geographic range extensions of stalked,
flabelliform sponges (Porifera) from eastern Canada with a new combination of a species
of Plicatellopsis in the North Atlantic. Zootaxa 4755, 301–321.
https://doi.org/10.11646/zootaxa.4755.2.6
Dohrmann, M., Wörheide, G., 2017. Dating early animal evolution using phylogenomic data.
Sci. Rep. 7, 3599. https://doi.org/10.1038/s41598-017-03791-w
Dunn, C.W., Leys, S.P., Haddock, S.H.D., 2015. The hidden biology of sponges and
ctenophores. Trends Ecol. Evol. 30, 282–291. https://doi.org/10.1016/j.tree.2015.03.003
Eitel, M., Francis, W.R., Varoqueaux, F., Daraspe, J., Osigus, H.-J., Krebs, S., Vargas, S., Blum,
H., Williams, G.A., Schierwater, B., Wörheide, G., 2018. Comparative genomics and the
nature of placozoan species. PLOS Biol. 16, e2005359.
https://doi.org/10.1371/journal.pbio.2005359
Erpenbeck, D., 2004. A molecular comparison of Alaskan and North East Atlantic Halichondria
panicea (Pallas 1766) (Porifera:Demospongiae) populations. BMIB - Boll. Dei Musei E
Degli Ist. Biol. 68.
Erpenbeck, D., Hooper, J.N.A., Wörheide, G., 2006. CO1 phylogenies in diploblasts and the
'Barcoding of Life'— are we sequencing a suboptimal partition? Mol. Ecol. Notes 6,
550–553. https://doi.org/10.1111/j.1471-8286.2005.01259.x
Erpenbeck, D., van Soest, R.W.M., 2002. Family Halichondriidae Gray, 1867, in: Hooper,
J.N.A., van Soest, R.W.M. (Eds.), Systema Porifera. A Guide to the Classification of
Sponges. Kluwer Academic/Plenum Publ., New York, pp. 787–816.
Erwin, P.M., Thacker, R.W., 2007. Phylogenetic analyses of marine sponges within the order
Verongida: a comparison of morphological and molecular data. Invertebr. Biol. 126,
220–234. https://doi.org/10.1111/j.1744-7410.2007.00092.x
Evcen, A., Gözcelioğlu, B., Çinar, M.E., 2023. New sponge records (Porifera) from the Black
Sea. Reg. Stud. Mar. Sci. 65, 103043. https://doi.org/10.1016/j.rsma.2023.103043
Folmer, O., Black, M., Wr, H., Vrijenhoek, R., 1994. DNA primers for amplification of
mitochondrial Cytochrome C oxidase subunit I from diverse metazoan invertebrates.
Mol. Mar. Biol. Biotechnol. 3, 294–299.
Forester, A.J., 1979. The association between the sponge Halichondria panicea (Pallas) and
scallop Chlamys varia (L.): A commensal-protective mutualism. J. Exp. Mar. Biol. Ecol.
36, 1–10. https://doi.org/10.1016/0022-0981(79)90096-0
Fristedt, K., 1887. Sponges from the Atlantic and Arctic Oceans and the Behring Sea. Vega-
Exped. Vetensk. Iakttag. Nord. 4, 410–471.
Galtier, N., Nabholz, B., Glémin, S., Hurst, G.D.D., 2009. Mitochondrial DNA as a marker of
molecular diversity: a reappraisal. Mol. Ecol. 18, 4541–4550.
https://doi.org/10.1111/j.1365-294X.2009.04380.x
Gazave, E., Carteron, S., Chenuil, A., Richelle-Maurer, E., Boury-Esnault, N., Borchiellini, C.,
2010. Polyphyly of the genus Axinella and of the family Axinellidae (Porifera:
Demospongiae). Mol. Phylogenet. Evol. 57, 35–47.
https://doi.org/10.1016/j.ympev.2010.05.028
Glasby, T.M., Connell, S.D., Holloway, M.G., Hewitt, C.L., 2007. Nonindigenous biota on
artificial structures: could habitat creation facilitate biological invasions? Mar. Biol. 151,
887–895. https://doi.org/10.1007/s00227-006-0552-5
Harbo, R., Ott, B., Reiswig, H., McDaniel, N., 2021. First Canadian record (Ladysmith Harbour,
British Columbia) of the non-native European sponge Hymeniacidon perlevis (Montagu,

1814) (Porifera, Demospongiae). BioInvasions Rec. 10, 277–286.
https://doi.org/10.3391/bir.2021.10.2.05
Hartman, W., 1958. Natural history of the marine sponges of southern New England. Bull.
Peabody Mus. Nat. Hist. 1–155.
Hirata, Y., Uemura, D., 1986. Halichondrins - antitumor polyether macrolides from a marine
sponge. Pure Appl. Chem. 58, 701–710. https://doi.org/10.1351/pac198658050701
Hoang, D., Chernomor, O., von Haeseler, A., Minh, B., Vinh, L., 2018. UFBoot2: Improving the
ultrafast bootstrap approximation. Mol. Biol. Evol. 35, 518–522.
https://doi.org/10.1093/molbev/msx281
Hooper, J.N.A., Van Soest, R.W.M., 2002. Systema Porifera. A Guide to the Classification of
Sponges, in: Hooper, J.N.A., Van Soest, R.W.M., Willenz, P. (Eds.), Systema Porifera: A
Guide to the Classification of Sponges. Springer US, Boston, MA, pp. 1–7.
https://doi.org/10.1007/978-1-4615-0747-5_1
ICZN, 1999. International Code of Zoological Nomenclature, 4th ed. International Trust for
Zoological Nomenclature, London.
Jennings, H.S., 1917. The numerical results of diverse systems of breeding, with respect to two
pairs of characters, linked or independent, with special relation to the effects of linkage.
Genetics 2, 97–154.
Johnson, G., 1842. A history of British sponges and lithophytes. W.H. Lizars, Edinburgh.
Jörger, K.M., Schrödl, M., 2013. How to describe a cryptic species? Practical challenges of
molecular taxonomy. Front. Zool. 10, 59. https://doi.org/10.1186/1742-9994-10-59
Jun, J., Yu, J.-N., Choi, E.H., 2015. Complete mitochondrial genome of Hymeniacidon sinapium
(Demospongiae, Halichondriidae). Mitochondrial DNA 26, 261–262.
https://doi.org/10.3109/19401736.2013.823189
Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermiin, L.S., 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods 14,
587–589. https://doi.org/10.1038/nmeth.4285
Kang, D.W., Kim, H., Kim, H.J., 2022. New species of Halichondria (Suberitida:
Halichondriidae) from Korea. J. Asia-Pac. Biodivers. 15, 98–101.
https://doi.org/10.1016/j.japb.2021.10.004
Katoh, K., Rozewicki, J., Yamada, K.D., 2017. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief. Bioinform. 20, 1160–
1166. https://doi.org/10.1093/bib/bbx108
Kim, H., Kang, D.W., Yu, C., Jung, Y.H., Yoon, M., Kim, H.J., 2019. The complete
mitochondrial genome of sponge Halichondria sp. (Demospongiae, Suberitida,
Halichondriidae) from Dok-do. Mitochondrial DNA Part B 4, 1729–1730.
https://doi.org/10.1080/23802359.2019.1610099
Knobloch, S., Jóhannsson, R., Marteinsson, V., 2019. Bacterial diversity in the marine sponge
Halichondria panicea from Icelandic waters and host-specificity of its dominant symbiont
"Candidatus Halichondribacter symbioticus." FEMS Microbiol. Ecol. 95, fiy220.
https://doi.org/10.1093/femsec/fiy220
Lambe, L.M., 1893. Sponges from the Pacific coast of Canada. Trans. R. Soc. Can. 11, 25–43.
Lawley, J.W., Gamero-Mora, E., Maronna, M.M., Chiaverano, L.M., Stampar, S.N., Hopcroft,
R.R., Collins, A.G., Morandini, A.C., 2021. The importance of molecular characters
when morphological variability hinders diagnosability: systematics of the moon jellyfish

genus Aurelia (Cnidaria: Scyphozoa). PeerJ 9, e11954.
https://doi.org/10.7717/peerj.11954

Lee, W., Elvin, D., Reiswig, H., 2007. The sponges of California: A guide and key to the marine
sponges of California. Monterey Bay Sanctuary Foundation, Monterey, CA.

Letunic, I., Bork, P., 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Res. 47, W256–W259. https://doi.org/10.1093/nar/gkz239

Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2015. MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct de Bruijn
graph. Bioinformatics 31, 1674–1676. https://doi.org/10.1093/bioinformatics/btv033

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinforma. Oxf. Engl. 25, 1754–1760.
https://doi.org/10.1093/bioinformatics/btp324

Luo, Z.-X., Yuan, C.-X., Meng, Q.-J., Ji, Q., 2011. A Jurassic eutherian mammal and divergence
of marsupials and placentals. Nature 476, 442–445. https://doi.org/10.1038/nature10291

Maldonado, M., 2006. The ecology of the sponge larva. Can. J. Zool. 84, 175–194.
https://doi.org/10.1139/z05-177

Mehbub, M.F., Yang, Q., Cheng, Y., Franco, C.M.M., Zhang, W., 2024. Marine sponge-derived
natural products: trends and opportunities for the decade of 2011-2020. Front. Mar. Sci.
11. https://doi.org/10.3389/fmars.2024.1462825

Meier, R., Blaimer, B.B., Buenaventura, E., Hartop, E., von Rintelen, T., Srivathsan, A., Yeo, D.,
2022. A re-analysis of the data in Sharkey et al.'s (2021) minimalist revision reveals that
BINs do not deserve names, but BOLD Systems needs a stronger commitment to open
science. Cladistics 38, 264–275. https://doi.org/10.1111/cla.12489

Miklucho-Maclay, N.N., 1870. Uber einige Schwämme des nördlichen Stillen Oceans und des
Eismeeres, welche im Zoologischen Museum der Kaiserlichen Akademie der
Wissenschaften in St. Petersburg aufgestellt sind. Ein Beitrag zur Morphologie und
Verbreitung der Spongien. Mém. Académie Impériale Sci. St Petersbourg 15, 1–24.

Montagu, G., 1814. An Essay on sponges, with descriptions of all the species that have been
discovered on the coast of Great Britain. Mem. Wernerian Nat. Hist. Soc. 2, 67–122.

Morrow, C., Cárdenas, P., 2015. Proposal for a revised classification of the Demospongiae
(Porifera). Front. Zool. 12, 1–27. https://doi.org/10.1186/s12983-015-0099-8

Morrow, C.C., Picton, B.E., Erpenbeck, D., Boury-Esnault, N., Maggs, C.A., Allcock, A.L.,
2012. Congruence between nuclear and mitochondrial genes in Demospongiae: A new
hypothesis for relationships within the G4 clade (Porifera: Demospongiae). Mol.
Phylogenet. Evol. 62, 174–190. https://doi.org/10.1016/j.ympev.2011.09.016

Morrow, C.C., Redmond, N.E., Picton, B.E., Thacker, R.W., Collins, A.G., Maggs, C.A.,
Sigwart, J.D., Allcock, A.L., 2013. Molecular phylogenies support homoplasy of multiple
morphological characters used in the taxonomy of Heteroscleromorpha (Porifera:
Demospongiae). Integr. Comp. Biol. 53, 428–446. https://doi.org/10.1093/icb/ict065

Müller, O.F., 1776. Zoologiae Danicae prodromus, seu Animalium Daniae et Norvegiae
indigenarum: characteres, nomina, et synonyma imprimis popularium. Typis Hallagerii,
Havni, Copenhagen.

Nguyen, L.-T., Schmidt, H., von Haeseler, A., Minh, B., 2015. IQ-TREE: A fast and effective
stochastic algorithm for estimating maximum likelihood phylogenies. Mol. Biol. Evol.
32, 268–274. https://doi.org/10.1093/molbev/msu300

Ott, B., Mcdaniel, N., Humphrey, E., 2024. Fourteen new species of demosponges (Porifera) from three coastal fjords in southern British Columbia, Canada. Zootaxa 5463, 151–200. https://doi.org/10.11646/zootaxa.5463.2.1

Pallas, P.S., 1766. Elenchus zoophytorum sistens generum adumbrationes generaliores et specierum cognitarum succintas descriptiones, cum selectis auctorum synonymis. Fransiscum Varrentrapp, Hagae.

Palumbi, S.R., 1986. How body plans limit acclimation: Responses of a Demosponge to wave force. Ecology 67, 208–214. https://doi.org/10.2307/1938520

Pankey, M.S., Plachetzki, D.C., Macartney, K.J., Gastaldi, M., Slattery, M., Gochfeld, D.J., Lesser, M.P., 2022. Cophylogeny and convergence shape holobiont evolution in sponge-microbe symbioses. Nat. Ecol. Evol. 6, 750–762. https://doi.org/10.1038/s41559-022-01712-3

Park, M.-H., Sim, C.-J., Baek, J., Min, G.-S., 2007. Identification of genes suitable for DNA barcoding of morphologically indistinguishable Korean Halichondriidae sponges. Mol. Cells 23, 220–227. https://doi.org/10.1016/S1016-8478(23)07377-6

Rannala, B., Yang, Z., 2020. Chapter 5.5 Species Delimitation, in: Scornavacca, C., Delsuc, F., Galtier, N. (Eds.), Phylogenetics in the Genomic Era. No commercial publisher, pp. 1–568.

Redmond, N.E., Morrow, C.C., Thacker, R.W., Diaz, M.C., Boury-Esnault, N., Cárdenas, P., Hajdu, E., Lôbo-Hajdu, G., Picton, B.E., Pomponi, S.A., Kayal, E., Collins, A.G., 2013. Phylogeny and systematics of Demospongiae in light of new small-subunit ribosomal DNA (18S) sequences. Integr. Comp. Biol. 53, 388–415. https://doi.org/10.1093/icb/ict078

Ristau, D.A., 1978. Six new species of shallow-water marine demosponges from California. Proc. Biol. Soc. Wash. 91, 569–589.

Rittmeyer, E.N., Austin, C.C., 2012. The effects of sampling on delimiting species from multi-locus sequence data. Mol. Phylogenet. Evol. 65, 451–463. https://doi.org/10.1016/j.ympev.2012.06.031

Rodriguez Jimenez, A., Dechamps, E., Giaux, A., Goetghebuer, L., Bauwens, M., Willenz, P., Flahaut, S., Laport, M.S., George, I.F., 2021. The sponges Hymeniacidon perlevis and Halichondria panicea are reservoirs of antibiotic-producing bacteria against multi-drug resistant Staphylococcus aureus. J. Appl. Microbiol. 131, 706–718. https://doi.org/10.1111/jam.14999

Rossi, M.E., Kenny, N.J., Giacomelli, M., Keating, J.N., Álvarez-Carretero, S., Schuster, A., Cárdenas, P., Taboada, S., Koutsouveli, V., Donoghue, P., Riesgo, A., Pisani, D., 2024. Independent origins of spicules reconcile the evolutionary history of sponges. https://doi.org/10.1101/2024.06.24.600355

Rot, C., Goldfarb, I., Ilan, M., Huchon, D., 2006. Putative cross-kingdom horizontal gene transfer in sponge (Porifera) mitochondria. BMC Evol. Biol. 6, 1–11. https://doi.org/10.1186/1471-2148-6-71

Ruiz, G.M., Fofonoff, P.W., Carlton, J.T., Wonham, M.J., Hines, A.H., 2000. Invasion of coastal marine communities in North America: apparent patterns, processes, and biases. Annu. Rev. Ecol. Evol. Syst. 31, 481–531. https://doi.org/10.1146/annurev.ecolsys.31.1.481

Samaai, T., Turner, T.L., Kara, J., Yemane, D., Ngwakum, B.B., Payne, R.P., Kerwath, S., 2022. Confirmation of the southern African distribution of the marine sponge Hymeniacidon

perlevis (Montagu, 1814) in the context of its global dispersal. PeerJ 10, e14388. https://doi.org/10.7717/peerj.14388

Schmidt, O., 1870. Grundzüge einer Spongien-Fauna des atlantischen Gebietes. Wilhelm Engelmann, Leipzig.

Schmittmann, L., Franzenburg, S., Pita, L., 2021. Individuality in the immune repertoire and induced response of the sponge Halichondria panicea. Front. Immunol. 12. https://doi.org/10.3389/fimmu.2021.689051

Schneider, C., Rasband, W., Eliceiri, K., 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9, 671–675. https://doi.org/10.1038/nmeth.2089

Sharkey, M.J., Janzen, D.H., Hallwachs, W., Chapman, E.G., Smith, M.A., Dapkey, T., Brown, A., Ratnasingham, S., Naik, S., Manjunath, R., Perez, K., Milton, M., Hebert, P., Shaw, S.R., Kittel, R.N., Solis, M.A., Metz, M.A., Goldstein, P.Z., Brown, J.W., Quicke, D.L.J., van Achterberg, C., Brown, B.V., Burns, J.M., 2021. Minimalist revision and description of 403 new species in 11 subfamilies of Costa Rican braconid parasitoid wasps, including host records for 219 species. ZooKeys 1013, 1–665. https://doi.org/10.3897/zookeys.1013.55600

Sim, C.J., Bakus, G.J., 1986. Marine sponges of Santa Catalina Island, California. Occas. Pap. Allan Hancock Found. 5, 1–23.

Simkanin, C., Davidson, I.C., Dower, J.F., Jamieson, G., Therriault, T.W., 2012. Anthropogenic structures and the infiltration of natural benthos by invasive ascidians. Mar. Ecol. 33, 499–511. https://doi.org/10.1111/j.1439-0485.2012.00516.x

Tessler, M., Neumann, J.S., Kamm, K., Osigus, H.-J., Eshel, G., Narechania, A., Burns, J.A., DeSalle, R., Schierwater, B., 2022. Phylogenomics and the first higher taxonomy of Placozoa, an ancient and enigmatic animal phylum. Front. Ecol. Evol. 10. https://doi.org/10.3389/fevo.2022.1016357

Thacker, R., Hill, A., Hill, M., Redmond, N., Collins, A., Morrow, C., Spicer, L., Carmack, C., Zappe, M., Pohlmann, D., Hall, C., Diaz, M., Bangalore, P., 2013. Nearly complete 28S rRNA gene sequences confirm new hypotheses of sponge evolution. Integr. Comp. Biol. 53, 373–387. https://doi.org/10.1093/icb/ict071

Thomassen, S., Riisgård, H.U., 1995. Growth and energetics of the sponge Halichondria panicea. Mar. Ecol. Prog. Ser. 128, 239–246. https://doi.org/10.3354/meps128239

Topsent, E., 1928. Spongiaires de l'Atlantique et de la Mediterranee provenant des croisieres du Prince Albert ler de Monaco. Result. Camp. Sci. Accompl. Par Ie Prince Albert Monaco 74, 1–376.

Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., Minh, B., 2016. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. Nucleic Acids Res. 44, W232–W235. https://doi.org/10.1093/nar/gkw256

Turner, T.L., 2020. The marine sponge Hymeniacidon perlevis is a globally-distributed invasive species. Bioarxve doi: 10.1101/2020.02.13.948000.

Turner, T.L., Lonhart, S.I., 2023. The sponges of the Carmel Pinnacles marine protected area. Zootaxa 5318, 151–194. https://doi.org/10.11646/ZOOTAXA.5318.2.1

Turner, T.L., Pankey, M.S., 2023. The order Axinellida (Porifera: Demospongiae) in California. Zootaxa 5230, 501–539. https://doi.org/10.11646/zootaxa.5230.5.1

van Soest, R.W.M., 2016. Sponge-collecting from a drifting ice floe: the Porifera obtained in the Kara Sea by the Dutch Polar Expedition 1882-83. Contrib. Zool. 85, 311–336.

van Soest, R.W.M., Hajdu, E., 2002. Family Esperiopsidae Hentschel, 1923, in: Hooper, J.N.A.,
van Soest, R.W.M. (Eds.), Systema Porifera: A Guide to the Classification of Sponges.
Springer, Boston, pp. 656–664.
Vethaak, A.D., Cronie, J.A., van Soest, R.W.M., 1982. Ecology and distribution of two
sympatric, closely related sponge species, Halichondria panicea (Pallas, 1766) and H.
bowerbanki Burton, 1930 (Porifera, Demospongiae), with remarks on their speciation.
Bijdr. Tot Dierkd. 52.
Wang, D., Zhang, Y., Huang, D., 2016. The complete mitochondrial genome of sponge
Halichondria (Halichondria) sp. (Demospongiae, Suberitida, Halichondriidae).
Mitochondrial DNA Part B 1, 512–514. https://doi.org/10.1080/23802359.2016.1192516
Witte, U., Barthel, D., Tendal, O., 1994. The reproductive cycle of the sponge Halichondria
panicea Pallas (1766) and its relationship to temperature and salinity. J. Exp. Mar. Biol.
Ecol. 183, 41–52. https://doi.org/10.1016/0022-0981(94)90155-4
Xavier, J.R., Rachello-Dolmen, P.G., Parra-Velandia, F., Schönberg, C.H.L., Breeuwer, J. a. J.,
van Soest, R.W.M., 2010. Molecular evidence of cryptic speciation in the "cosmopolitan"
excavating sponge Cliona celata (Porifera, Clionaidae). Mol. Phylogenet. Evol. 56, 13–
20. https://doi.org/10.1016/j.ympev.2010.03.030
Xue, L., Zhang, X., Zhang, W., 2009. Larval release and settlement of the marine sponge
Hymeniacidon perlevis (Porifera, Demospongiae) under controlled laboratory conditions.
Aquaculture 290, 132–139. https://doi.org/10.1016/j.aquaculture.2009.01.037
Zamani, A., Fric, Z.F., Gante, H.F., Hopkins, T., Orfinger, A.B., Scherz, M.D., Bartoňová, A.S.,
Pos, D.D., 2022. DNA barcodes on their own are not enough to describe a species. Syst.
Entomol. 47, 385–389. https://doi.org/10.1111/syen.12538
Zhong, W., Olugbami, J.O., Rathakrishnan, P., Mohanty, I., Moore, S.G., Garg, N., Oyelere,
A.K., Turner, T.L., McShan, A.C., Agarwal, V., 2024. Discovery and folding dynamics
of a fused bicyclic cysteine knot undecapeptide from the marine sponge Halichondria
bowerbanki. J. Org. Chem. 89, 12748–12752. https://doi.org/10.1021/acs.joc.4c01104
3221