Immunologic and Biophysical Features of the BNT162b2 JN.1- and KP.2-Adapted

COVID-19 Vaccines

Authors

Wei Chen[1*], Kristin R. Tompkins[1*], Ian W. Windsor[2*], Lyndsey T. Martinez[1], Minah Ramos[1],

Weiqiang Li[1], Shikha Shrivastava[1], Swati Rajput[1], Jeanne S. Chang[2], Parag Sahasrabudhe[2],

Kimberly F. Fennell[2], Thomas J. McLellan[2], Graham M. West[2], Kristianne P. Dizon[1], Aaron

Yam[1], Siddartha Mitra[1], Subrata Saha[1], Daiana Sharaf [1], Andrew P. McKeen[3], Carla I. Cadima[4],

Alexander Muik[4], Wesley Swanson[1], Raquel Munoz Moreno[1], Pilar Mendoza Daroca[1], Ugur

Sahin[4], Annaliesa S. Anderson[1], Huixian Wu[2], Kena A. Swanson[1▫] & Kayvon Modjarrad[1▫]

* Contributed equally

▫ Corresponding authors: kayvon.modjarrad@pfizer.com, kena.swanson@pfizer.com

Affiliations

[1]Pfizer Vaccine Research and Development; Pearl River, NY, USA.

[2]Pfizer Discovery Sciences; Groton, CT, USA.

[3]Pfizer Global Biometrics and Data Management; Pearl River, NY, USA.

[4]BioNTech; Mainz, Germany.

ABSTRACT Vaccines remain a vital public health tool to reduce the burden of COVID-19.

COVID-19 vaccines that are more closely matched to circulating SARS-CoV-2 lineages elicit

more potent and relevant immune responses that translate to improved real-world vaccine

effectiveness. The rise in prevalence of the Omicron JN.1 lineage, and subsequent derivative

sublineages such as KP.2 and KP.3, coincided with reduced neutralizing activity and

effectiveness of Omicron XBB.1.5-adapted vaccines. Here, we characterized the biophysical

and immunologic attributes of BNT162b2 JN.1- and KP.2-adapted mRNA vaccine-encoded

spike (S) protein immunogens. Biophysical interrogations of S revealed the structural

consequences of hallmark amino acid substitutions and a potential molecular mechanism of

immune escape employed by JN.1 and KP.2. The vaccine candidates were evaluated for their

immunogenicity when administered as fourth or fifth doses in BNT162b2-experienced mice or

as a primary series in naïve mice. In both vaccine-experienced and naïve settings, JN.1- and

KP.2-adapted vaccines conferred improved neutralizing responses over the BNT162b2 XBB.1.5

vaccine against a broad panel of emerging JN.1 sublineages, including the predominant

KP.3.1.1 and emerging XEC lineages. Antigenic mapping of neutralizing responses indicated

greater antigenic overlap of JN.1- and KP.2-adapted vaccine responses with currently

circulating sublineages compared to an XBB.1.5-adapted vaccine. $CD4^+$ and $CD8^+$ T cell

responses were generally conserved across all three vaccines. Together, the data support the

selection of JN.1- or KP.2-adapted vaccines for the 2024-25 COVID-19 vaccine formula.

KEYWORDS

COVID-19; SARS-CoV-2; BNT162b2; mRNA vaccine; preclinical; variants; immunogenicity;

spike protein; structure; cryogenic-electron microscopy

ONE-SENTENCE SUMMARY

The Omicron JN.1- and KP.2-adapted BNT162b2 mRNA vaccines encoding prefusion S proteins elicit similar preclinical neutralizing antibody responses against circulating JN.1 sublineage pseudoviruses that are more potent than those elicited by past iterations of BNT162b2 licensed vaccines, thus demonstrating the importance of annual strain changes to the COVID-19 vaccine.

INTRODUCTION

As coronavirus disease-2019 (COVID-19) transitions from a pandemic to an endemic state, questions remain around the evolutionary trajectory of its causative pathogen, Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), the periodicity of its incidence, and the necessary frequency of variant-adapted vaccine updates to ensure optimal protection against a range of clinical outcomes. It remains clear, however, that the annual burden of COVID-19 cases, hospitalizations and deaths still places it among the leading global causes of infectious disease morbidity and mortality, similar to or higher than pre-pandemic levels observed for influenza and other pneumonias [1-4].

Since the emergence of the first SARS-CoV-2 Omicron variant of concern (VOC) in late 2021, the virus has differentiated into more than 4300 distinct genetic lineages [5]. As these lineages have trended toward increased transmissibility and greater antigenic distance from the original ancestral strain, approved vaccines have been updated to include variant-specific mutations in the SARS-CoV-2 spike (S) [6-10], to maintain optimal protection against the most relevant lineages causing COVID-19. Although updated vaccines have been typically introduced to the general public during the early Autumn season of the Northern Hemisphere,

vaccine manufacturers must conduct at-risk development activities throughout the preceding

year in anticipation of the variant composition recommendations issued by regulators and

international normative bodies in the Spring of that year [9-11].

During the Northern Hemisphere Winter of 2023 to 2024, the variant epidemiology of

SARS-CoV-2 shifted from a dominance of Omicron XBB derivative lineages to BA.2.86

descendants, most notably the variant of interest, JN.1 [12] and its sublineages, achieving a peak

prevalence that exceeded all prior Omicron lineages since the emergence of BA.1 and BA.2. As

the global dominance of JN.1 plateaued and declined in prevalence, derivative sublineages

acquired additional mutations, translating to amino acid substitutions in the S protein that

conferred improved viral fitness or immune escape. Some of those sublineages (*e.g.*, KP.2)

gained epidemiologic relevance during the summer months of 2024, particularly those that

contained convergent F456L and R346T amino acid substitutions in the S receptor binding

domain (RBD). More recently, the KP.3 sublineage, which contains the F456L, but not the

R346T substitution, has emerged as globally dominant, likely owing to its acquisition of the

Q493E substitution, also located within the RBD. This sublineage and its derivatives,

particularly KP.3.1.1, now account for most SARS-CoV-2 infections [13].

The JN.1 cluster of sublineages occupies a unique antigenic space separate from all

other prior Omicron lineages. Real-world studies of the XBB.1.5-adapted COVID-19 vaccines

demonstrated robust immune response and effectiveness against XBB lineages that diminished

in response to the BA.2.86 and JN.1 lineages [14-19]. To better understand the structural and

functional consequences of the antigenic shift from this new lineage cluster, and subsequent

antigenic drift within it, we evaluated the biophysical and immunologic features of the

prefusion-stabilized full-length SARS-CoV-2 S-proteins of JN.1 and KP.2. To our knowledge,

we present the first in-depth structural analysis of JN.1 and KP.2 full-length, prefusion-

stabilized S immunogens, building prior reports detailing the impact of JN.1 sublineage-specific

S mutations on neutralization resistance. We also assessed whether an Omicron JN.1 or KP.2-

updated BNT162b2 vaccine could improve immune responses that are important for conferring

protection against COVID-19 [20], against the JN.1 lineage and a panel of contemporary JN.1-

derived sublineages, as compared to the BNT162b2 XBB.1.5 vaccine.

Both monovalent JN.1- and KP.2-adapted COVID-19 vaccines have been recommended

for the 2024-2025 formula. The two lineages differ in their S amino acid sequences by 3

residues [21]. In prior years, small genetic differences have not necessarily translated into

antigenic differences. Given that two variant-adapted vaccines have been introduced in the same

season, we sought to understand the impact of these few, but potentially consequential, genetic

differences on biophysical, structural and immunologic attributes of vaccine-encoded JN.1 and

KP.2 prefusion stabilized S proteins. The BNT162b2 JN.1 and KP.2 vaccines use the same

mRNA backbone as prior approved versions of BNT162b2 and are minimally modified to

contain lineage-specific sequence changes of the SARS-CoV-2 prefusion stabilized S protein

that has been evaluated in previous preclinical and clinical studies . We employed a vaccine-

experienced murine model, with varying schedules of prior vaccination regimens, together with

a vaccine-naïve murine model, to assess humoral and cellular immunogenicity, so as to

approximate the diversity of the immune-experience of the general population across all age

groups [22]. An evaluation of both structural and functional consequences of the most recent

evolution of SARS-CoV-2 lineages in varied models may then elucidate the potential impact on

vaccine performance.

METHODS

SARS-CoV-2 S(P2) and RBD Protein Expression and Purification

Protein sequences of the full-length prefusion stabilized S(P2) of the SARS-CoV-2 JN.1 lineage and KP.2 sublineage contained amino acid changes relative to the ancestral Wuhan-Hu-1 S (GenBank accession number: MN908947.3) as listed in Table S1**Error! Reference source not found.**. S(P2) contains two proline substitutions at residues 986 and 987 (K986P, V987P) and utilizes a C-terminal TwinStrep tag. The RBD constructs contain an N-terminal S protein leader peptide and coding regions from 321-527 for JN.1 and KP.2, and 327-528 for the ancestral strain, followed by a C-terminal affinity tag as indicated in Table S1. Details on S(P2) and RBD protein expression and purification can be found in the Supplementary Materials.

Stability of Wild Type (WT), Omicron JN.1 and KP.2 FL S(P2) by Thermal Shift Assay (TSA)

Stability of FL S(P2) proteins was measured by Tycho NT.6 (NanoTemper, firmware version: 1.10.3) as described previously [23]. The inflection temperature for each thermal melting curve reported by the Tycho NT.6 software was reported as the thermal melting temperature (T_m) of S(P2) proteins.

Binding Kinetics of Purified FL S(P2) Protein and RBD to Immobilized Human ACE-2-PD

FL S(P2) and RBD proteins of JN.1, KP.2 and WT strains were assessed by biolayer interferometry (BLI) binding to immobilized human ACE-2-PD, as described previously [23].

Cryogenic Electron Microscopy (Cryo-EM) of JN.1 and KP.2 S(P2)

SARS-CoV-2 FL S(P2) (K986P and V987P substituted) proteins of the JN.1 and KP.2 lineages were interrogated by electronic cryogenic electron microscopy (cryo-EM) analysis following membrane extraction and purification. Grids were prepared with JN.1 and KP.2 S(P2) proteins at 5.6 and 5.8 mg/ml by applying 4 µL to Quantifoil R1.2/1.3 200 mesh gold grids that were glow-discharged for 60 seconds at 15 mA in a Pelco easiGlow cleaning system and blotted for 3 seconds with a blot force of +3 before plunging into liquid nitrogen-cooled liquid ethane using a Mark IV Vitrobot with a sample chamber maintained at 4°C and 100% humidity. Data collection was performed with a Titan Krios G2 electron microscope operated at 300 kV, controlled by EPU, and equipped with a Falcon 4i direct electron detector and Selectris energy filter configured with a 10 eV slit (ThermoFisher Scientific). A total of 7k and 9k movies were collected for JN.1 and KP.2 S(P2) protein samples with a total dose of 40.0 $e^-/Å^2$ and a -0.4 µm to -2.4 µm defocus range. Details on cryo-EM data processing and model building can be found in the Supplementary Materials and Table S3.

Mass Spectrometry Characterization of JN.1 and KP.2 S(P2) N-linked Glycosylation

Mapping of N-linked glycosylation sites was conducted on recombinant purified JN.1 and KP.2 S(P2) as described previously [23]. For glycosylation complexity determination, S(P2) protein digests, before and after O18 water and PNGase F treatment, were analyzed using liquid chromatography-mass spectrometry (LC-MS). The data were processed using Protein Metrics PMI-Byos Byologic PTM workflow with glycan modifications enabled. Relative levels of the various glycan forms were determined by summing the extracted ion chromatogram peak areas

for all detected charge states and forms for a given isobaric glycosylation state. Relative

estimation of abundance was obtained by comparing frequencies of different glycans, with an

assumption that all glycopeptides have similar ionization.

Animal Ethics

All mouse studies were performed at Pfizer, Inc. (Pearl River, NY, USA), an Association for

Assessment and Accreditation of Laboratory Animal Care (AAALAC) accredited facility.

Animals were observed after all procedures and injection sites were monitored following each

vaccination. All procedures performed on animals were in accordance with regulations and

established guidelines and were reviewed and approved by an Institutional Animal Care and

Use Committee or through an ethical review process.

Immunogenicity Studies

4th or 5th Vaccination in BNT162b2-Experienced Mice

Monovalent XBB.1.5, JN.1, and KP.2-adapted BNT162b2 vaccines were evaluated as either a

4th or 5th dose in female BALB/c mice (10 per group; Jackson Laboratory). In the 5th dose study,

mice were vaccinated intramuscularly (i.m.) at 6-8 weeks of age with 2-doses (Day 0, 21) of

original BNT162b2 WT vaccine, followed by a 3rd dose booster (Day 49) of bivalent WT +

Omicron BA.4/5 vaccine, a 4th dose (Day 84) of the monovalent Omicron XBB.1.5 adapted

vaccine, and a 5th dose (Day 111) of either the monovalent XBB.1.5, JN.1, or KP.2-adapted

vaccine. All vaccinations used a 0.5 µg total dose in a 50 µL volume (bivalent formulations

contained equal quantities of each mRNA, 0.25 µg each). A control group of five mice received

saline injections (50 µL) according to the same schedule, in place of active vaccines. Sera were

collected for evaluation of pseudovirus neutralizing antibody responses prior to the 5th dose

(Day 111) and at the study end (Day 140). Spleens were collected from all mice at Day 140 to

evaluate cell-mediated immune responses (Fig. S4A).

In the 4th dose study, female BALB/c mice (10 per group; Jackson Laboratory) were first

vaccinated i.m. at 6-8 weeks of age and subsequent doses were administered according to the

same schedule as the 5th dose study, except that the 4th dose was administered on Day 64 as

either the monovalent XBB.1.5, JN.1, or KP.2-adapted vaccine and there was no 5th dose. As

above, sera were collected for evaluation of pseudovirus neutralization responses prior to the 4th

dose (Day 64) and two weeks later at study end (Day 78) (Fig. S4B).

2-Dose Primary Series in Naïve Mice

Female BALB/c mice (10 per group Jackson Laboratory) were vaccinated i.m. at 6-8 weeks of

age on Days 0 and 21 with either monovalent XBB.1.5, JN.1, or KP.2-adapted vaccine. As

described above, vaccine formulations contained a total dose level of 0.5 µg mRNA in a 50 µL

volume and a control group (10 mice) received saline injections adjacent to the vaccine groups.

Sera and spleens (5 mice/group) were collected 28 days after the second dose (day 49) for

evaluation of pseudovirus neutralizing antibody responses and cell-mediated immune responses,

respectively (Fig. S4C).

VSV-SARS-CoV-2 S Pseudovirus Neutralization Assay

The pseudovirus neutralization assay (PNA) was defined and performed as previously described

[23]. VSV-based pseudoviruses contained the S protein from the following SARS-CoV-2 lineages:

WT (Wuhan-Hu-1, ancestral strain), BA.4/5, XBB.1.5, JN.1, JN.1.6,1, JN.1.7, JN.1.13.1,

KW.1.1, KP.2, KP.2.2, KP.1.1, KS.1, JN.1.16.1, KZ.1.1.1, KP.2.3, KP.3, KP.3.1.1, LB.1 and

XEC. Amino acid sequence substitutions relative to JN.1 for all tested pseudoviruses are

provided in Fig. S5.

Omicron XBB.1.5, JN.1 and KP.2 Antigenic Cartography

Antigenic maps were constructed using the antigenic cartography toolkit Racmacs v1.2.9

(https://acorg.github.io/Racmacs/index.html). In brief, antigenic cartography is a method to

quantify and visualize neutralization data. An antigenic map takes titration data that measures

strength of reactivity of a group of antisera (sera where an individual mouse has been vaccinated

with a unique antigen of a particular SARS-CoV-2 lineage) against a group of antigens

(different SARS-CoV-2 lineages). Antigenic mapping uses multidimensional scaling to position

antigens (viruses) and sera in a map to represent their antigenic relationships. Antigenically

similar strains are spatially close to one another on the map, while antigenically distinct strains

are further apart. The spacing between grid lines is 1 unit of antigenic distance, corresponding

to a two-fold dilution of antiserum in the neutralization assay. The maps were constructed by

Racmacs package in R using 2000 optimizations, with the minimum column basis parameter set

to "none."

T-cell Response by Flow Cytometry Assay

Murine splenocytes were stimulated *ex vivo* as previously described (*1*) with DMSO only

(unstimulated) or specific amino acid (aa) peptide libraries (15aa, 11aa overlap, 1 to 2

μg/mL/peptide) representing SARS-CoV-2 S amino acid sequences. Five individual peptide

pools represented the full-length S sequence of the ancestral Wuhan (WT) (JPT), BA.4/5 (JPT),

XBB.1.5 (JPT), JN.1 (JPT) and KP.2 lineages (Mimotopes). Following stimulation, splenocytes

were stained for CD154 (CD40L), IFN-γ, TNF-α, and IL-2 positive CD4$^+$ and CD8$^+$ T cells, as

previously described [23]. Samples were acquired on a 5-Laser Aurora system (Cytek ®) using

SpectroFlo® software (version 3.1.2). The instrument was subject to daily quality control

procedures using SpectroFlo® QC Beads per manufacturer recommendations. Acquired data

files were analyzed using OMIQ® software. The T cell gating strategy is shown in Fig. S6.

Data Availability Statement

The final full-length S(P2) cryo-EM density maps and models for the JN.1 (3-down and 1-up)

and KP.2 (3-down, 1-up, and 2-up) conformations are deposited in the Electron Microscopy

Data Bank (EMDB) and the Protein Data Bank (PDB) under accession codes EMD-46637 and

PDB ID 9D8H, EMD-46638 and PDB ID 9D8H, EMD-46639 and PDB ID 9D8I, EMD-46640

and PDB ID 9D8J, and EMD-46641 and PDB ID 9D8K, respectively.

Statistical Analysis

Mouse immunogenicity data were analyzed using SAS version 9.4. All statistical analyses were

performed using ANOVA on log-transformed data. Comparisons were made on mouse sera

across pseudoviruses at the last post-vaccination timepoint of each study with Dunnett's test for

multiple comparisons. For intergroup comparisons, the XBB.1.5 vaccine group was the

reference; for intragroup comparisons (pseudoviruses within a vaccine group), the vaccine

target lineage (XBB.1.5, JN.1 or KP.2) served as the reference. All tests were two-tailed. A

p-value of less than 0.05 was considered statistically significant.

RESULTS

Enhanced ACE2 receptor binding and reduced thermostability of JN.1 and KP.2 S

proteins compared to WT SARS-CoV-2 S protein

Full-length prefusion-stabilized S(P2) proteins of Wuhan-Hu-1 (WT), Omicron JN.1, and KP.2

were expressed from DNA corresponding to the adapted BNT162b2 RNA coding sequences

using similar methods as previously reported[23]. All S(P2) proteins were expressed on the cell

surface following in vitro transfection of Expi293F cells and showed similar binding to human

ACE2 (Fig. S1A). Analysis of membrane-extracted and affinity-purified full-length S(P2)

showed a single peak by size exclusion chromatography (SEC), with S predominantly cleaved

into S1 and S2 subunits (Fig. 1A and Fig. S1B). Thermal shift assay (TSA) analyses showed the

melting temperature (T_m) of JN.1 (61.6 $\pm$ 0.24 ° C) and KP.2 (60.1 $\pm$ 0.19 ° C) were similar to

one another (Fig. 1B) and approximately 5-7 °C lower than WT S(P2) (67.1 $\pm$ 0.17 °C) [23].

Biolayer interferometry (BLI) analysis of purified RBDs only, which contain the receptor

binding motif (RBM) critical for ACE2 binding, showed significantly increased binding of

soluble JN.1 and KP.2 RBDs to fixed human ACE2 peptidase domain (ACE2-PD) (K_D 1.70 nM

and 2.07 nM, respectively) compared to the WT RBD (K_D 31.3 nM) [23] (Fig. 1C).

Acquisition of glycosylation sites within the JN.1 and KP.2 S proteins

The purified S(P2) proteins of JN.1 and KP.2 were analyzed by liquid chromatography mass

spectrometry (LCMS) to identify N-linked glycosylation sites. Glycosylation mapping

confirmed that JN.1 and KP.2 have conserved glycosylation sites on the S protein, despite

undergoing extensive mutations compared to WT S. This is consistent with previous studies of

earlier lineages, suggesting a correlation between viral fitness and glycosylation [24]. Compared to

XBB.1.5, we discovered two new glycosylation sites (Table 1) in the RBD and N-terminal

domain (NTD) of JN.1 and KP.2 S, respectively. The novel RBD glycosylation site is located

on N354 which is gained by the K356T substitution, whereas the novel NTD glycosylation site

is a result of the H245N substitution within the peptide ALN[245]RSYLT. Analysis of the

glycosylation heterogeneity confirmed that both sites are fully modified by glycans with a major

composition being HexNAc(2)Hex(5) (Table S2).

JN.1 and KP.2 S(P2) cryo-EM structure

Structures of JN.1 and KP.2 full-length, prefusion stabilized SARS-CoV-2 S proteins were

solved by cryo-EM to validate the conformation of the encoded antigen of both vaccines and to

evaluate the structural consequences of the lineages' hallmark mutations in the context of S(P2).

Both JN.1 and KP.2 S(P2) proteins exhibited prefusion conformations, as anticipated. The cryo-

EM processing employed herein sought to identify all RBD (up/down) configurations (Fig. S2)

within the S trimer. Unlike the WT S(P2) protein, which was predominantly (~80%) found in

the 3-down conformation[25], the prefusion-stabilized JN.1 S protein was observed in the 3-down

(65%) conformation, and at a lower frequency in a 1-up (35%) conformations, while KP.2 S

was more evenly observed in 3-down (41%) and 1-up (48%) conformations, with a small

fraction in a 2-up (11%) conformation, indicating KP.2 exhibits a greater tendency than earlier

variants for adopting the RBD-up conformations (Fig. 2A).

In both JN.1 and KP.2 S, the C3-symmetric, 3-down conformation of the RBD core is

resolved at ~4Å, whereas the receptor binding motif (RBM), comprising residues 450-490, were

absent at high resolution. In the 1-up conformations, the RBM was best resolved for the down-

RBD adjacent to the up-RBD (Fig. S3). The RBM of the down-RBD, adjacent to the up-RBD,

was poorly resolved, though it contained high-resolution features for the core. In contrast, very

little of the second up-RBD was seen in the 2-RBD-up KP.2 structure (Fig. S3).

Structural changes for both lineages were likely caused by two amino acid substitutions

shared by JN.1 and KP.2—P621S and P1143L—that differentiate them from XBB.1.5. The

P621 residue is in a region of the S subdomain 2 (SD2) called the 630 loop that is unstructured

in full-length, 2P-stabilized XBB.1.5 and Wuhan S protein cryo-EM densities. Despite being

situated in this unstructured region, the P621S substitution leads to a well-resolved alpha helix

(Fig. 2B). The structured region is found in every protomer of the five solved structures of the

JN.1 and KP.2 S proteins. In the context of the 1-up RBD structures of JN.1 and KP.2, a region

adjacent to the 630 helix—residues 829-848 or the fusion peptide proximal region (FPPR) of

the adjacent protomer—is also well-resolved (Fig. 2B, Fig. S3). Though lower in resolution,

there is a clear density, including a disulfide bond between C836 and C847. As a result, the

structured FPPR interacts with the 630-helix from the adjacent protomer, contributing to an

enhanced interprotomer stability which may reduce S1/S2 shedding in the context of the RBD-

up conformation. The P1143 residue is located near the N-terminus of the stem helix. The cryo-

EM density revealed two discrete conformations of the stem for both JN.1 and KP.2 S. The

alternative poses were best resolved in the JN.1 3-down, C3-symmetry reconstruction (Fig. 2C,

Fig. S3) and diverge near the mutated proline residue.

Two additional mutations, H245N and K356T, differentiate JN.1 and KP.2 from

XBB.1.5 and earlier lineages, and are shared with the parental BA.2.86 lineage. These

mutations have yielded novel glycosylation sites that were fully glycosylated in the resolved

structures (Fig. 2D, Tables 2 and S1). However, H245N within the NTD is located in a low-

resolution region of the EM densities of the two JN.1 and three KP.2 structures. As a result, the

possible glycosylation modification of H245N is absent from the cryo-EM map reconstruction.

The K356T mutation led to glycosylation at N354 in JN.1, KP.2 and previous variants

harboring the mutation [26,27]. This glycan is located near the interface of the RBD and the NTD

of an adjacent protomer (Fig. 2D).

Overall, JN.1 and KP.2 S proteins differ by three residues: R346T and F456L in the

RBD and V1104L in the S2 domain. R346T has been observed in several earlier lineages and

may impact antibodies recognizing the RBD class 3 epitope [28]. The V1104L mutant residue

exhibited density consistent with the additional methylene group and no discernable structural

changes. F456L is located in the RBM and in a relatively low-resolution region of the map. Of

note, the RBD is best resolved in the down conformation adjacent to the up conformation and

the mutant L456 is located at the protomer interface (Fig. 2D, Fig. S3).

BNT162b2 JN.1- and KP.2-adapted vaccines neutralizing responses to JN.1 sublineages in

a vaccine-experienced mouse model

BNT162b2 JN.1- and KP.2-adapted vaccines were evaluated in two murine studies that

examined booster immunogenicity in a BNT162b2-experienced immune setting. To

approximate the immune background of a vaccinated human population, a vaccine-experienced

animal model was generated by vaccinating mice with all licensed BNT162b2 vaccines

sequentially spanning 2021 to the present [Original (2-doses), Bivalent Original+BA.4/5 (1-

dose), XBB.1.5 (1-dose)]. BNT162b2 XBB.1.5, JN.1, or KP.2-adapted vaccines were

administered as a 5th dose to female BALB/c mice 27 days following the XBB.1.5 vaccination

(4th dose) (Fig. S4A). Sera were collected prior to and one month following administration of

the 5[th] dose for assessment of pseudovirus neutralization against contemporary JN.1 lineages

(JN.1, JN.1.16.1, KP.2, KP.2.3, KP.3, KP.3.1.1, LB.1, and XEC).

As a 5[th] dose, JN.1- and KP.2-adapted vaccines elicited much higher 50% geometric

mean neutralizing titers (GMTs) against JN.1 and all JN.1 sublineages tested, as compared to

the XBB.1.5 vaccine (Fig. 3A). The JN.1 and KP.2 vaccine responses were 3-to-4 times and 7-

to-10 times higher, respectively, compared to the XBB.1.5 vaccine group, including against the

globally prevalent KP.3.1.1 sublineage and the rapidly rising XEC sublineage (Fig. 3B). In

XBB.1.5 vaccinated animals, neutralizing responses against the antigenically distant JN.1

lineage were approximately 20-fold lower than against the vaccine-matched lineage, XBB.1.5

(Fig. 3A). In contrast, sera from JN.1- and KP.2-vaccinated mice neutralized all JN.1

sublineages with similar potency, indicating a broadly robust and cross-protective immune

response. From pre- to post-5[th] dose, the JN.1-adapted vaccine boosted neutralizing responses

(GMT fold rise (GMFR)) against JN.1 and KP.2 by 3.5- and 3.9-fold, respectively. The KP.2-

adapted vaccine increased neutralizing responses by 5.9-fold and 7.4-fold against the JN.1 and

KP.2 lineages, respectively, as compared to the XBB.1.5 vaccine (Fig. S7). The XBB.1.5-

adapted vaccine elicited the lowest GMFRs against JN.1 and KP.2 (0.9 and 1.8, respectively).

In a separate study, BNT162b2 XBB.1.5, JN.1, or KP.2 vaccines were administered as a

4[th] dose to female BALB/c mice two weeks after the 3[rd] dose (BNT162b2 bivalent WT +

BA.4/5) (Fig. S4B). As observed in the 5[th] dose study, sera from JN.1- and KP.2-vaccinated

mice neutralized a broad panel of JN.1 sublineages with similar potency (Fig. S8A). Here again,

the JN.1- and KP.2-adapted vaccines elicited significantly (p <0.05) higher neutralizing

responses against JN.1 sublineages, on the order of 2-to-4-fold and 3-to-7-fold, respectively,

than the XBB.1.5 vaccine. GMFRs from pre-to-post 4[th] dose were also similar between the JN.1

and KP.2 vaccines, as compared to the XBB.1.5-adapted vaccine and similar to the magnitude

of rise observed when the vaccines were given as a 5th dose (Fig. S8B). Despite the variations in

prior vaccination regimens, the JN.1 and KP.2-adapted vaccines elicited consistent

neutralization trends and significantly improved immunogenicity against contemporary JN.1

lineage pseudoviruses, including against the more recently dominant sublineages (*i.e.*, KP.3.1.1

and XEC), as compared to the XBB.1.5 vaccine.

BNT162b2 JN.1 and KP.2-adapted vaccines neutralizing responses to JN.1 sublineages in

naïve mice

The Omicron XBB.1.5, JN.1 and KP.2 vaccines were administered on Days 0 and 21 to naïve

female BALB/c mice as a primary series (Fig. S4C). Sera were collected one month after the 2nd

dose and tested against the pseudovirus panel used in the vaccine experienced studies described

above. Although the BNT162b2 XBB.1.5 vaccine induced robust neutralizing responses against

the XBB.1.5 sublineage, it failed to elicit similar neutralizing titers against the antigenically

distant JN.1 lineage and sublineages (Fig. 4A). Overall, the JN.1 and KP.2-adapted vaccines

elicited significantly higher neutralizing responses against JN.1 and other relevant JN.1

sublineages, compared to the XBB.1.5-adapted vaccine (Fig. 4), by an order of magnitude

greater than the differences observed in the vaccine-experienced models. Neutralizing GMTs in

the JN.1 and KP.2 vaccine groups were 9-to-14-fold and 15-to-29-fold higher against JN.1 and

JN.1 sublineages, respectively, as compared to the XBB.1.5 vaccine group (Fig. 4B). Although

KP.2-adapted vaccine-elicited GMTs trended 2-to-3-fold higher than JN.1 vaccine-elicited

GMTs, these differences were not statistically significant (Fig. 4B).

Mapping of neutralizing antibody responses reveals antigenic shifts and drifts of SARS-CoV-2 Omicron lineages

To investigate the relative antigenic differences among lineages, as reflected in variant-adapted vaccine humoral immunogenicity, serum neutralizing titers from XBB.1.5-, JN.1- and KP.2-adapted BNT162b2 vaccinated mice (vaccine-experienced and naïve) described above, were used to generate antigenic maps of contemporary SARS-CoV-2 lineages, relative to one another and to the ancestral Wuhan-Hu-1 (WT) strain (Fig. 5). In the map of vaccine-elicited sera from naïve mice, JN.1 and all JN.1-derived sublineages lie within 2 antigenic units (1 unit equals 2-fold change in neutralization titer) from each other, suggesting a high antigenic similarity and indicative of limited antigenic drift within the JN.1 cluster thus far (Fig. 5A). In contrast, JN.1 and JN.1-derived sublineages were found to lie more than 4 antigenic units away from the original WT strain, and previously dominant lineages BA.4/5 and Omicron XBB.1.5, indicating a major antigenic shift from earlier SARS-CoV-2 lineages. Sera from vaccine-experienced mice yielded similar spatial relationships (Fig. 5B); however, these data are likely confounded by the influence of cross-reactive antibodies recognizing conserved epitopes. Antigenic maps generated in a naïve background more accurately reflect the true antigenic differences between virus strains or species. The cartographies of both backgrounds, however, clearly demonstrate that evolution toward JN.1 lineages marks a major antigenic shift from prior dominant lineages, such as those belonging to the XBB cluster. The distance of JN.1 lineages from XBB.1.5 is even greater than the latter is from its epidemiologically dominant predecessor, BA.4/5. However, the JN.1 lineages, particularly the parental JN.1 and its derivative KP.2, occupy proximal or overlapping coordinates of the antigenic maps, which is consistent with the similarity in neutralizing activity elicited by both vaccines against the entire panel of JN.1 sublineages.

BNT162b2 JN.1- and KP.2-adapted vaccines induce comparable S-specific CD4$^+$ and

CD8$^+$ T cell responses in BNT162b2-experienced and naïve mice

In both vaccine-experienced and naïve mice, spleens collected one-month following final

vaccination were interrogated for S-specific T cell cytokine responses using a flow cytometry-

based intracellular cytokine staining (ICS) assay (Fig. S6). Peptide pools representing the full-

length S protein from the Wuhan (WT) virus, BA.4/5, XBB.1.5, JN.1 and KP.2 variants were

used to assess lineage-specific CD4$^+$ and CD8$^+$ T cell responses *ex vivo*.

In vaccine-experienced mice, the XBB.1.5, JN.1 and KP.2 vaccines induced overall high

frequencies of S-specific CD4$^+$ and CD8$^+$ T cells as compared to the saline control (Fig. 6, Fig.

S9). Mice administered the 5th dose (Fig. S4A) of KP.2 or JN.1-adapted vaccine elicited similar

frequencies of IFN-γ^+, TNF-α^+, and IL-2$^+$ CD4$^+$ T cells, as compared to the XBB.1.5 vaccine

(Fig. 6A-C). Within each vaccine group the frequencies of cytokine expressing CD4$^+$ T cells

were similar across all the lineages tested, indicating that the T cell responses were highly cross-

reactive. Similarly, for the cytokine expressing (IFN-γ^+ or TNF-α^+ or IL-2$^+$) CD8$^+$ T cells, the

KP.2 and JN.1-adapted vaccine elicited comparable frequencies as the XBB.1.5-adapted

vaccine (Fig. 6D-F). Within each vaccine group, individual lineage-specific T cell frequencies

varied slightly; however, these variations did not impact the overall magnitude of the T cell

response, and each vaccine was capable of inducing robust T cell cytokine responses against all

the lineages analyzed. The same was true for polyfunctional (IFN-γ^+ TNF-α^+ IL-2$^+$) CD4$^+$ (Fig.

S9A) and CD8$^+$ T cells (Fig. S9B).

In naïve mice, the trends for both CD4$^+$ (Fig. 7A-C, Fig. S10A) and CD8$^+$ (Fig. 7D-F,

Fig. S10B) T cell responses elicited by the JN.1 and KP.2 vaccines, with respect to cross-

reactivity and similarity to one another and to the XBB.1.5 vaccine, were generally consistent

with those observed in the vaccine-experienced study. In both models, minimal IL-4 secreting

$CD4^+$ T cells were detected (data not shown), particularly in relation to those secreting IFN-γ,

TNF-α and IL-2, indicating a strong bias toward a Th1 over a Th2 cytokine profile, consistent

with character of the T cell responses elicited by the original and earlier versions of BNT162b2

[23,25].

DISCUSSION

The dynamic epidemiologic landscape of SARS-CoV-2 presents a unique challenge for

the development of periodically updated COVID-19 vaccines. Since its emergence, SARS-

CoV-2 has evolved quickly and broadly, comparable to other respiratory viruses that account

for a large burden of disease, such as influenza [29]. The periodicity of disease incidence, though,

has not aligned with the more predictable seasonality of other respiratory viral infections [30,31].

These attributes pose challenges for vaccine manufacturers and regulatory authorities alike in

making decisions about what lineages to target in vaccine updates and when to introduce those

updated vaccines into the general population, to make the greatest impact in reducing the

burden of COVID-19 morbidity and mortality.

The FDA recently approved the 2024-2025 formula of BNT162b2 for individuals 12

years of age and older and granted emergency use authorization (EUA) for individuals 6 months

through 11 years of age in the US [32], following a recommendation to mRNA vaccine

manufacturers to target the KP.2 lineage in the vaccine update [11]. The genetic drift of lineages,

and questions around the relative antigenic proximity to future lineages that may emerge in the

Northern Hemisphere Winter of 2024/2025, has prompted slightly different recommendations

from regulatory authorities on the composition of a vaccine update [9-11]. As such, both JN.1- and

KP.2-adapted formulas of the BNT162b2 vaccine have been approved and authorized in

different countries. The nonclinical evaluation of both formulas in multiple immune

backgrounds against a broad panel of immunologically and epidemiologically relevant

contemporary lineages, summarized here, has enabled an informed decision-making process in

the development of these variant-adapted vaccines.

JN.1 differs from most of its descendant lineages by 1 to 5 amino acid residues in the S

protein, which is in the same range of difference that separated XBB.1.5 from other dominant

XBB sublineages [33]. As the XBB.1.5 vaccine conferred cross-protective immunity against other

sublineages within the XBB lineage cluster [15-19,34], we hypothesized that JN.1 and KP.2-adapted

vaccines would be similarly effective in generating broad immune responses against

contemporary JN.1 sublineages. Virus neutralizing antibody responses trend closely with

protection from COVID-19 disease [20,35,36], and therefore have been a useful surrogate for

COVID-19 vaccine performance against emerging lineages. The degree to which those lineages

escape vaccine elicited neutralization has generally served as a metric of how well updated

vaccines will protect against COVID-19.

To elucidate potential molecular mechanisms of immune escape, we sought to first

characterize the JN.1 and KP.2 full-length prefusion stabilized S(P2) proteins and RBDs. Our

data revealed the unique molecular features of JN.1 and KP.2 S(P2) gained from the extensive

mutations during viral evolution. The S(P2) proteins of both JN.1 and KP.2 have lower T_m as

compared to the WT and XBB.1.5 S(P2) proteins [23]. This finding is consistent with the idea that

SARS-CoV-2 may be evolving toward gradually reduced thermostability, which, along with the

increased population of S in RBD-up conformations, could contribute to increased infectivity

and transmissibility, and account for the epidemiologic growth advantage of these lineages

relative to their predecessors. When assessing ACE2 binding affinity of the new JN.1 lineages,

the RBDs of JN.1 and KP.2 have inherited several mutations also contained in XBB.1.5 that

enhance receptor binding, such as G446S, Q498R, N501Y and Y505H in the RBM. The

additional E484K substitution acquired by JN.1 and KP.2 may contribute to further

enhancement of receptor binding via a salt bridge with E35 from ACE2 (Fig. 1D). This

likelihood is supported by data that shows JN.1 and KP.2 RBD bind to ACE2 with a higher

potency, as compared to the WT RBD[23].

Few cryo-EM structures of SARS-CoV-2 S proteins have been resolved with more than

one RBD-up conformation and not bound to ACE2 or antibodies [37]. KP.2 exhibits a greater

propensity to occupy the RBD-up conformation than does JN.1, despite these lineages differing

at only three residues in the S protein. We reason that inter-RBD interactions conferred by

F456L substitution may stabilize the RBD up conformation. In the 1- and 2-up KP.2 structures,

L456 of the down-RBD is in the immediate vicinity of P384 of the up-RBD, and likely

participates in hydrophobic interactions (Fig. 2D, Fig. S3). The FPPR is only well-resolved in

the protomer bearing the up conformation in the 1-up structures. The improved resolution of

this feature in the KP.2 versus the JN.1 structure may be due in part to the stabilization imposed

by the inter-RBD contact mediated by L456, especially as RBD dynamics have been implicated

in interactions within this distal region [38,39].

Previous non-prefusion-stabilized structures of the Omicron BA.1 lineage revealed that

the 630-loop was well resolved in the down conformation [40], like the D614G and other earlier

lineages [38,39]. The P612S mutation may play a role similar to D614G by stabilizing the FPPR

via a structured 630-loop, thereby enhancing the stability of the cleaved trimer and bolstering

interactions between protomers, as well as between S1 and S2; however, doing so in the setting

of the up conformation, like the Gamma variant [38].

The P1143 residue is located at the N-terminus of the stem helix. The secondary amide

between the side chain and the main chain renders this residue unable to donate an upstream

hydrogen bond. Conversely, L1143, a new mutation acquired by both JN.1 and KP.2 can

capitalize on this hydrogen bonding interaction, where this additional secondary structure is

likely to enforce an alternative trajectory of the stem helix. The interior facing residues of the

stem are the target of weak, broadly neutralizing antibodies and the P1443L mutation may,

therefore, represent an initial step toward stabilizing a conformation of the stem helix that

occludes this epitope, yet is not detrimental to attaining the postfusion conformation.

Neutralizing antibodies that are overrepresented at the population level are key drivers

of antigenic drift, as most that have conserved gene usage and stereotyped epitope recognition

have been rendered ineffective by point mutations [41]. For example, P4J15 is a rare antibody that

recognizes the class 1 epitope and has maintained breadth from the original Wuhan strain

through XBB.1 [42]. However, the F456L mutation, present in KP.2 and located within the P4J15

epitope, likely impacts its binding (Fig. 2E). The N364 glycan is similarly located in a region

recognized by broadly neutralizing antibodies, such as G32R7 (Fig. 2E). This region is known

as the RBD class 3 or the RBD1 competition cluster, and represents the largest memory B cell

epitope cluster among convalescent patients infected at the start of the pandemic [28]. The

mutations found in KP.2 likely perturb the epitopes of the few remaining neutralizing antibodies

elicited by the Wuhan strain and underpin much of its fitness and immune escape among the

SARS-CoV-2-experienced population.

As nonclinical data have closely aligned with clinical responses in prior cycles of

variant-adapted vaccine updates [14], we assessed the immunogenicity of BNT162b2 JN.1 and

KP.2 vaccines against JN.1 and contemporary circulating sublineages of epidemiologic

relevance in BNT162b2-experienced and naïve mice. As most adults have developed hybrid

immunity from a mixture of prior infections and vaccinations, recapitulating the collective

history of antigenic exposures has become an increasingly complex endeavor, particularly with

the successive emergence of new lineage clusters each year . Nonetheless, models that account

for prior exposure history provide a more relevant model for evaluating vaccine responses in

immune experienced populations. To understand the impact of varying immune backgrounds on

variant-adapted vaccine immunogenicity, we assessed vaccine-elicited immune responses in

studies with differing compositions and schedules of prior vaccinations. The most parsimonious

model that maintains immunologic relevance is likely to streamline preparatory activities for

future vaccine updates.

Prior preclinical data that supported FDA approval of the BNT162b2 XBB.1.5 vaccine

in 2023 showed that vaccine-elicited neutralizing responses against the vaccine-matched

XBB.1.5 and related lineages (*e.g*. XBB.1.16, XBB.2.3) were 4-to-5 times higher than

responses elicited against the same lineage by the previously authorized vaccine (bivalent WT +

Omicron BA.4/5) [23]. Similarly, in the present studies, JN.1 and KP.2-adapted vaccines, elicited

neutralizing titers against JN.1 sublineages that were consistently improved over BNT162b2

XBB.1.5 vaccine responses. Overall, there was a 3-to-10-fold improved response over the

XBB.1.5 vaccine in BNT162b2-experienced mice across two studies, depending on the vaccine

formulation and the sublineage tested. JN.1 and KP.2 vaccines both elicited responses that

effectively neutralized all 16 JN.1 pseudoviruses evaluated, including those sublineages that

acquired advantageous R346T and F456L substitutions (*e.g.*, JN.1.16.1, KP.2), and a next

generation of sublineages that have an additional advantageous deletion at the S31 position (*e.g.*,

KP.2.3, LB.1) [43]. KP.3 and its most dominant KP.3.1.1 sublineage, as well as the rapidly

emerging recombinant XEC lineage [13], were also evaluated and neutralized by JN.1 and KP.2

vaccines in the studies reported here.

As multiple animal models are employed in the evaluation of COVID-19 vaccines,

many still frequently perform primary series studies in naïve animals [44-48], which does not

account for the current immune experience of most adults. Even when vaccine-experienced

animal studies are employed to better recapitulate the antigenic exposures of the general

population, there remains a lack of consensus on how those studies should be designed. To

inform investigations into the influence of varied immune backgrounds, we assessed JN.1 and

KP.2 vaccine immunogenicity against two different vaccine-experience regimens, varying the

prior vaccine composition and schedule. Across both studies, we observed a similar trend in

neutralizing antibody responses for both JN.1 and KP.2 vaccines, regardless of prior BNT162b2

XBB.1.5 vaccine exposure or timing between vaccine administrations. Mice boosted with JN.1

or KP.2-adapted vaccines as a 4th dose elicited greater immunity against JN.1 and a broader

panel of JN.1 sublineages, consistent with findings of the 5th dose study. These findings indicate

that the exact sequence and scheme of prior antigenic exposures may not affect trends in

responses elicited by variant-updated vaccines. It may be more important to reproduce

exposures with the ancestral strain and an Omicron lineage as a means to approximate the

immune experience of the general population.

Despite the importance of immune-experienced models, it is critical to also evaluate

vaccine immunogenicity in naïve models that are particularly relevant to younger age cohorts

that have not yet been exposed to SARS-CoV-2 through either infection or vaccination.

Children under the age of four experience a high burden of COVID-19 hospitalization. During

the 2023–2024 respiratory season in the US, the cumulative COVID-19 hospitalization rate for

children under four was 104.3 per 100,000, and was higher only among persons ≥50 years of

age [4]. Notably, 50% of infants, children, and adolescents ≤17 years hospitalized with COVID-

19 from July 2023–March 2024, and 40% of those admitted to the ICU, had no underlying

medical conditions [49]. According to 2023 data in the US, COVID-19 was the third leading cause

of death due to infectious disease in individuals 0–17 years of age [50]. Given this public health

relevance, we evaluated a two-dose primary series in naïve mice, wherein we found that both

JN.1 and KP.2-adapted vaccines elicited a similar breadth of neutralizing activity against all

JN.1 sublineages that were improved over the XBB.1.5 vaccine responses by an order of

magnitude (8-to-29-fold higher). Responses elicited by the XBB.1.5 vaccine were also higher

(10-fold) than the prior vaccine (bivalent WT + Omicron BA.4/5) in a previous study [23].

Although the comparative trends in neutralizing responses across vaccine groups were

consistent across vaccine-experienced and naïve mouse studies, the magnitude of those

differences were much greater in the naïve model, reflecting the considerable antigenic distance

between the XBB and JN.1 lineage clusters.

Overall, JN.1 and KP.2 BNT162b2 vaccines elicited improved neutralizing antibody

responses over the XBB.1.5 vaccine in mice against the contemporary JN.1 lineages.

Additionally, the maintenance of comparable T cell responses between vaccines across a broad

set of variant-specific peptides in mice after booster vaccination indicates that cellular immune

responses are not substantially impacted by the mutational differences among recent Omicron

lineages. Biophysical characterization indicated that JN.1 and KP.2 S proteins are relatively

similar to one another but differ in their structural features from prior Omicron lineages. As

XBB.1.5 vaccines have demonstrated diminished effectiveness against the JN.1 lineage cluster,

and preclinical data have trended closely with clinical responses and real-world effectiveness

outcomes, it is anticipated that the potent responses elicited by both the JN.1 and KP.2-adapted

BNT162b2 vaccines will confer robust protection against antigenically drifted JN.1 sublineages,

including the most globally dominant and recently rising strains, similar to how the XBB.1.5

vaccine performed against the XBB lineage cluster [16-18,34]. As reported here, and moving

forward into the current and future peak seasons of COVID-19 disease activity, a

multidimensional approach is needed to monitor the evolutionary dynamics and epidemiology

of novel SARS-CoV-2 lineages and the effectiveness of the JN.1- and KP.2-adapted vaccines in

protecting against a wide range of clinical outcomes.

Acknowledgements

We thank the Pfizer Comparative Medicine department and veterinary staff at Pearl River, NY

for their contributions to the *in vivo* studies; members of Pfizer Vaccine Research and

Development at Pearl River, NY, for their contributions to assay development and

implementation, virus variant monitoring, material production and characterization, and

material and technical support in molecular biology, virology and immunology; and members of

Pfizer Discovery Sciences in Groton, CT, for their material and technical support on

recombinant protein production and characterization. The authors also thank Teresa Hauguel for

review of the manuscript, Christina D'Arco for writing and graphics support and Charlotte Ford

for editorial assistance.

Funding

This study was supported by Pfizer.

Author Contributions

WC contributed to writing of the original draft, methodology, investigation, and data generation

and interpretation. KRT contributed to the writing of the original draft, investigation, data

generation and interpretation. IWW contributed to writing of the original draft, cryo-EM

structural determination and visualization, data analysis and interpretation. LTM, MR, WL, SS,

SR, KPD, AY, DS, SM, SS, WS, RMM and PMD contributed to data generation, analysis,

visualization and interpretation; LTM also contributed to serologic methodology. JSC

performed antigen purification and thermal shift experiments. PS performed BLI experiments.

KFF performed recombinant protein expression. TJM and GMW performed site-specific glycan

mapping by MS studies. APM contributed to data analysis. CIC and AM contributed to data

interpretation. US and ASA supervised the work. HW contributed to writing of the original draft,

supervision, investigation, visualization, data interpretation and analysis related to the structural

and biophysical work. KAS contributed to writing of the original draft, data interpretation,

investigation, and overall supervision and conceptualization of the work. KM contributed to

writing of the original draft, experimental designs, data analysis and interpretation, investigation

and overall supervision and conceptualization of the work. All authors contributed to the

development and critical review of the manuscript and have read and agreed to the published

version.

Competing Interests

All authors are current or former employees of Pfizer or BioNTech and may, therefore, be respective shareholders. Pfizer and BioNTech participated in the design, analysis and interpretation of the data as well as the writing of this report and the decision to publish. WL, AM, UŞ, KAS and KM are inventors on patents and patent applications related to the COVID-19 vaccine. AM and UŞ are inventors on patents and patent applications related to RNA technology.

References

HHS. *COVID-19 Reported Patient Impact and Hospital Capacity by State Timeseries (RAW)*, <https://healthdata.gov/Hospital/COVID-19-Reported-Patient-Impact-and-Hospital-Capa/g62h-syeh/about_data> (2024).
CDC. *Disease Burden of Flu*, <https://www.cdc.gov/flu/about/burden/> (2023).
CDC. *Preliminary Estimated Flu Disease Burden 2023–2024 Flu Season*, <https://www.cdc.gov/flu-burden/php/data-vis/2023-2024.html> (2024).
CDC. *COVID Data Tracker*, <https://covid.cdc.gov/covid-data-tracker/#datatracker-home> (2024).
corneliusroemer. *cov-lineages / pango-designation*, <https://github.com/cov-lineages/pango-designation/blob/master/lineage_notes.txt> (2024).
FDA. *Vaccines and Related Biological Products Advisory Committee June 5, 2024 Meeting Announcement*, <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-june-5-2024-meeting-announcement#:~:text=On%20June%205%2C%202024%2C%20the,and%2For%20video%20conferencing%20platform.> (2024).
FDA. *Vaccines and Related Biological Products Advisory Committee June 15, 2023 Meeting Announcement*, <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-june-15-2023-meeting-announcement> (2023).
FDA. *Vaccines and Related Biological Products Advisory Committee June 28, 2022 Meeting Announcement*, <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-june-28-2022-meeting-announcement> (2022).
W.H.O. *Statement on the antigen composition of COVID-19 vaccines*, <https://www.who.int/news/item/26-04-2024-statement-on-the-antigen-composition-of-covid-19-vaccines> (2024).
EMA. *ETF recommends updating COVID-19 vaccines to target new JN.1 variant*, <https://www.ema.europa.eu/en/news/etf-recommends-updating-covid-19-vaccines-target-new-jn1-variant> (2024).

11 FDA. *Updated COVID-19 Vaccines for Use in the United States Beginning in
Fall 2024*, <https://www.fda.gov/vaccines-blood-biologics/updated-covid-19-
vaccines-use-united-states-beginning-fall-2024> (2024).
12 W.H.O. *Initial risk evaluation of JN.1, 19 December 2023*,
<https://www.who.int/docs/default-
source/coronaviruse/18122023_jn.1_ire_clean.pdf> (2023).
13 W.H.O. *COVID-19 epidemiological update – 9 October 2024*,
<https://www.who.int/publications/m/item/covid-19-epidemiological-update-
edition-172> (2024).
14 Gayed, J. *et al.* Immunogenicity of the Monovalent Omicron XBB.1.5-Adapted
BNT162b2 COVID-19 Vaccine against XBB.1.5, BA.2.86, and JN.1
Sublineages: A Phase 2/3 Trial. *Vaccines (Basel)* **12** (2024).
https://doi.org:10.3390/vaccines12070734
15 Gayed, J. *et al.* Safety and Immunogenicity of the Monovalent Omicron
XBB.1.5-Adapted BNT162b2 COVID-19 Vaccine in Individuals >/=12 Years Old:
A Phase 2/3 Trial. *Vaccines (Basel)* **12** (2024).
https://doi.org:10.3390/vaccines12020118
16 Wang, Q. *et al.* XBB.1.5 monovalent mRNA vaccine booster elicits robust
neutralizing antibodies against XBB subvariants and JN.1. *Cell Host Microbe* **32**,
315-321 e313 (2024). https://doi.org:10.1016/j.chom.2024.01.014
17 Wang, Q. *et al.* Robust SARS-CoV-2-neutralizing antibodies sustained through
6 months post XBB.1.5 mRNA vaccine booster. *Cell Rep Med* **5**, 101701 (2024).
https://doi.org:10.1016/j.xcrm.2024.101701
18 Tartof, S. Y. *et al.* Effectiveness of BNT162b2 XBB Vaccine Against XBB and
JN.1 Sublineages. *Open Forum Infect Dis* **11**, ofae370 (2024).
https://doi.org:10.1093/ofid/ofae370
19 Ma, K. C. *et al.* Effectiveness of Updated 2023-2024 (Monovalent XBB.1.5)
COVID-19 Vaccination Against SARS-CoV-2 Omicron XBB and BA.2.86/JN.1
Lineage Hospitalization and a Comparison of Clinical Severity-IVY Network, 26
Hospitals, October 18, 2023-March 9, 2024. *Clin Infect Dis* (2024).
https://doi.org:10.1093/cid/ciae405
20 Khoury, D. S. *et al.* Neutralizing antibody levels are highly predictive of immune
protection from symptomatic SARS-CoV-2 infection. *Nat Med* **27**, 1205-1211
(2021). https://doi.org:10.1038/s41591-021-01377-8
21 Kaku, Y. *et al.* Virological characteristics of the SARS-CoV-2 KP.2 variant.
Lancet Infect Dis **24**, e416 (2024). https://doi.org:10.1016/S1473-
3099(24)00298-6
22 W.H.O. *Interim statement on hybrid immunity and increasing population
seroprevalence rates*, <https://www.who.int/news/item/01-06-2022-interim-
statement-on-hybrid-immunity-and-increasing-population-seroprevalence-rates>
(2022).
23 Modjarrad, K. *et al.* Preclinical Characterization of the Omicron XBB.1.5-
Adapted BNT162b2 COVID-19 Vaccine. *bioRxiv* **accepted in Npj Vaccines**,
2023.2011.2017.567633 (2023). https://doi.org:10.1101/2023.11.17.567633

24 Shajahan, A., Pepi, L. E., Kumar, B., Murray, N. B. & Azadi, P. Site specific N-
and O-glycosylation mapping of the spike proteins of SARS-CoV-2 variants of
concern. *Sci Rep* **13**, 10053 (2023). https://doi.org:10.1038/s41598-023-33088-0
25 Vogel, A. B. *et al.* BNT162b vaccines protect rhesus macaques from SARS-
CoV-2. *Nature* **592**, 283-289 (2021). https://doi.org:10.1038/s41586-021-03275-
y
26 Liu, P. *et al.* Spike N354 glycosylation augments SARS-CoV-2 fitness for human
adaptation through structural plasticity. *Natl Sci Rev* **11**, nwae206 (2024).
https://doi.org:10.1093/nsr/nwae206
27 Li, L. *et al.* Spike structures, receptor binding, and immune escape of recently
circulating SARS-CoV-2 Omicron BA.2.86, JN.1, EG.5, EG.5.1, and HV.1 sub-
variants. *Structure* **32**, 1055-1067 e1056 (2024).
https://doi.org:10.1016/j.str.2024.06.012
28 Tong, P. *et al.* Memory B cell repertoire for recognition of evolving SARS-CoV-2
spike. *Cell* **184**, 4969-4980 e4915 (2021).
https://doi.org:10.1016/j.cell.2021.07.025
29 Wang, X. *et al.* SARS-CoV-2 versus Influenza A Virus: Characteristics and Co-
Treatments. *Microorganisms* **11** (2023).
https://doi.org:10.3390/microorganisms11030580
30 Gong, Z. *et al.* Natural and socio-environmental factors in the transmission of
COVID-19: a comprehensive analysis of epidemiology and mechanisms. *BMC*
Public Health **24**, 2196 (2024). https://doi.org:10.1186/s12889-024-19749-3
31 Barosa, M., Ioannidis, J. P. A. & Prasad, V. Evidence base for yearly respiratory
virus vaccines: Current status and proposed improved strategies. *Eur J Clin*
Invest, e14286 (2024). https://doi.org:10.1111/eci.14286
32 FDA. *Pfizer-BioNTech COVID-19 Vaccine: Pfizer-BioNTech COVID-19 Vaccine*
(2024-2025 Formula) Authorized For Individuals 6 Months through 11 Years of
Age, <https://www.fda.gov/vaccines-blood-biologics/coronavirus-covid-19-cber-
regulated-biologics/pfizer-biontech-covid-19-vaccine> (2024).
33 Tamura, T. *et al.* Virological characteristics of the SARS-CoV-2 Omicron
XBB.1.5 variant. *Nat Commun* **15**, 1176 (2024). https://doi.org:10.1038/s41467-
024-45274-3
34 Chalkias, S. *et al.* Interim Report of the Reactogenicity and Immunogenicity of
Severe Acute Respiratory Syndrome Coronavirus 2 XBB-Containing Vaccines.
The Journal of infectious diseases **230**, e279-e286 (2024).
https://doi.org:10.1093/infdis/jiae067
35 Cromer, D. *et al.* Neutralising antibody titres as predictors of protection against
SARS-CoV-2 variants and the impact of boosting: a meta-analysis. *Lancet*
Microbe **3**, e52-e61 (2022). https://doi.org:10.1016/S2666-5247(21)00267-6
36 Earle, K. A. *et al.* Evidence for antibody as a protective correlate for COVID-19
vaccines. *Vaccine* **39**, 4423-4428 (2021).
https://doi.org:10.1016/j.vaccine.2021.05.063
37 Cai, Y. *et al.* Structural basis for enhanced infectivity and immune evasion of
SARS-CoV-2 variants. *Science* **373**, 642-648 (2021).
https://doi.org:10.1126/science.abi9745

38   Zhang, J. *et al.* Membrane fusion and immune evasion by the spike protein of
SARS-CoV-2 Delta variant. *Science* **374**, 1353-1360 (2021).
https://doi.org:10.1126/science.abl9463
39   Zhang, J. *et al.* Structural impact on SARS-CoV-2 spike protein by D614G
substitution. *Science* **372**, 525-530 (2021).
https://doi.org:10.1126/science.abf2303
40   Zhang, J. *et al.* Structural and functional impact by SARS-CoV-2 Omicron spike
mutations. *Cell Rep* **39**, 110729 (2022).
https://doi.org:10.1016/j.celrep.2022.110729
41   Cao, Y. *et al.* Omicron escapes the majority of existing SARS-CoV-2
neutralizing antibodies. *Nature* **602**, 657-663 (2022).
https://doi.org:10.1038/s41586-021-04385-3
42   Fenwick, C. *et al.* Broadly potent anti-SARS-CoV-2 antibody shares 93% of
epitope with ACE2 and provides full protection in monkeys. *J Infect* **87**, 524-537
(2023). https://doi.org:10.1016/j.jinf.2023.10.008
43   Li, P. *et al.* Neutralization and Stability of JN.1-derived LB.1, KP.2.3, KP.3 and
KP.3.1.1 Subvariants. *bioRxiv* (2024).
https://doi.org:10.1101/2024.09.04.611219
44   Slamanig, S. *et al.* Intranasal SARS-CoV-2 Omicron variant vaccines elicit
humoral and cellular mucosal immunity in female mice. *EBioMedicine* **105**,
105185 (2024). https://doi.org:10.1016/j.ebiom.2024.105185
45   Zhang, Y. *et al.* Three SARS-CoV-2 spike protein variants delivered intranasally
by measles and mumps vaccines are broadly protective. *Nat Commun* **15**, 5589
(2024). https://doi.org:10.1038/s41467-024-49443-2
46   Yang, Z. H. *et al.* Measles Virus-Based Vaccine Expressing Membrane-
Anchored Spike of SARS-CoV-2 Inducing Efficacious Systemic and Mucosal
Humoral Immunity in Hamsters. *Viruses* **16** (2024).
https://doi.org:10.3390/v16040559
47   Hoffmann, M. A. G. *et al.* ESCRT recruitment to SARS-CoV-2 spike induces
virus-like particles that improve mRNA vaccines. *Cell* **186**, 2380-2391 e2389
(2023). https://doi.org:10.1016/j.cell.2023.04.024
48   Singh, O. N. *et al.* Immunogenicity and protection efficacy of self-amplifying and
circular mRNA vaccines for SARS-CoV-2. *bioRxiv*, 2024.2008.2023.609366
(2024). https://doi.org:10.1101/2024.08.23.609366
49   Havers, F. P. *COVID-19–Associated Hospitalizations among Children and*
Adults — COVID-NET, <https://www.cdc.gov/acip/downloads/slides-2024-06-
26-28/02-COVID-Havers-508.pdf> (2024).
50   CDC. *About Provisional Mortality Statistics, 2018 through Last Week*,
<https://wonder.cdc.gov/mcd-icd10-provisional.html> (
808
809
810
811

814
Figure 1. Biophysical Characteristics of SARS-CoV-2 Omicron JN.1 and KP.2 Prefusion-

Stabilized Spike and Receptor Binding Domain. a SEC profiles of the S(P2) proteins of JN.1

and KP.2. The S(P2) proteins were purified in DDM (*n*-dodecyl-*β*-D-maltopyranoside, see

Supplementary Materials). SDS-PAGE fractions from 12 mL to 15 mL of each FL S construct

are shown in Fig. S1B. **b** Melting temperature (Tm) of purified S(P2) proteins determined by

the inflection point of the first derivatives of protein fluorescence signals collected at 330 nm

over that of 350 nm. Experiments were conducted by Tycho NT.6 and run in triplicates (n=3). **c**

Biolayer interferometry (BLI) sensorgrams of RBD binding to immobilized human angiotensin

converting enzyme-2 peptidase domain (ACE2-PD). The binding curves (black) were globally

fit to a 1:1 Langmuir binding model (light blue). Calculated apparent K_D, k_{on}, k_{off} values are

listed in the figure. **d** Crystal structure of ACE2-PD (PDB: 6M0J) is aligned to the KP.2 1-up

RBD. ACE2-PD is shown in maroon ribbon with surface representation. KP.2 RBD is shown in

blue cartoon with amino acid residue changes in the receptor binding motif (RBM) from WT to

KP.2 shown in yellow spheres. RBM residue changes (blue circle) that enhance receptor

binding via putative specific interactions (dotted lines) to ACE2 residues (maroon circle) are

shown in the interaction scheme on the right.

Figure 2. Structural and Functional Consequences of SARS-CoV-2 JN.1 and KP.2 S

Protein Mutations. a Local resolution-filtered maps of the final 3D reconstructions are shown

for each of the resolved RBD conformational states and colored by protomer. Local resolution

estimations for maps shown in panels A though D are found in Figure S3. The first up RBD is

colored green in both the JN.1 and KP.2 maps and the second up RBD in the KP.2 is colored

blue. Proportions of particles derived from reference-based classification are expressed as a

percentage below each map. **b** Magnified view of the 630-helix and fusion peptide proximal

region (FPPR) from the KP.2 1-RBD-up map overlaid with the model shows the quality of the

EM density in this region. **c** Magnified view of the stem from the JN.1 3-RBD-down map

overlaid with the model shows the two partially occupied conformations of this helix beginning

at the P1143L mutation. **d** Top view of the KP.2 1-RBD-up conformation structure shows the

location of the N354 glycan due to the K356T substitution and a putative interprotomer

hydrophobic interaction with P384 due to the F456L substitution. **e** Broad monoclonal

antibodies (mAbs) G32R7 (PDB: 7N65) and P4J15 (PDB: 8PQ2), which bind early omicron

lineages, are aligned to the KP.2 1-RBD-up. The proximity of these epitopes and the K356T and

F456L substitution-induced surface alterations implicate these changes with evasion of early

immunity.

Figure 3. Pseudovirus Neutralization Elicited by BNT162b2 XBB.1.5, JN.1 And KP.2-Adapted Vaccines Administered to Vaccine-Experienced Mice. Female mice were immunized i.m. according to Fig. S4A on Days 0 and 21 with the BNT162b2 Wuhan (WT), on Day 49 with the bivalent BNT162b2 (WT + BA.4/5), on Day 84 with BNT162b2 XBB.1.5, and on Day 111 with the BNT162b2 XBB.1.5, JN.1, or KP.2 vaccine. One month post-5th dose, sera were collected from the terminal bleed and neutralizing antibody responses were measured against a panel of 11 pseudoviruses that included the Wuhan (WT) reference strain and Omicron lineages BA.4/5, XBB.1.5, JN.1, JN.16.1, KP.2, KP.2.3, KP.3, KP.3.1.1, LB.1, and XEC. **a** The number above each bar indicates the 50% neutralizing geometric mean titer (GMT) with 95% CI of 10 mice per vaccine group. **b** The geometric mean ratio (GMR) is shown as the ratio of the BNT162b2 KP.2 or JN.1 vaccine GMT to the BNT162b2 XBB.1.5 vaccine GMT of the corresponding pseudovirus. The number above each bar indicates the GMR with 95% CI. The limit of detection (LOD) is the lowest serum dilution, 1:20. Asterisks indicate statistical significance of pseudovirus GMR relative to the corresponding pseudovirus in the monovalent XBB.1.5 vaccine group. ****p<0.0001, ***p<0.001, ** p<0.01, * p<0.05.

Figure 4. Pseudovirus Neutralization Elicited by BNT162b2 XBB.1.5, JN.1 and KP.2-

Adapted Vaccines Administered as a Primary Series to Naïve Mice. Female mice were

immunized i.m. according to Fig. S4A on Days 0 and 21 with the BNT162b2 XBB.1.5, JN.1, or

KP.2-adapted vaccine. Sera were collected from the terminal bleed (Day 49) and neutralizing

antibody responses were measured against a panel of 11 pseudoviruses that included the Wuhan

(WT) reference strain and Omicron lineages BA.4/5, XBB.1.5, JN.1, JN.16.1, KP.2, KP.2.3,

KP.3, KP.3.1.1, LB.1, and XEC. **a** The number above each bar indicates the 50% neutralizing

geometric mean titer (GMT) with 95% CI of 10 mice per vaccine group. **b** The geometric mean

ratio (GMR) is shown as the ratio of the BNT162b2 KP.2 or JN.1 vaccine GMT to the

BNT162b2 XBB.1.5 vaccine GMT of the analogous pseudovirus. The number above each bar

indicates the GMR with 95% CI. The limit of detection (LOD) is the lowest serum dilution,

1:20. Asterisks indicate statistical significance of pseudovirus GMR relative to the analogous

pseudovirus in the monovalent XBB.1.5 vaccine group. ****p<0.0001, ***p<0.001, ** p<0.01,

* p<0.05. N/A = Not available as both values to calculate GMR are at limit of detection.

Figure 5. Antigenic Map of SARS-CoV-2 Omicron JN.1 Lineages Relative to Lineages Contained in Prior BNT162b2 Lineage-Adapted Vaccines. a-b Antigenic map visualizes cross-reactivity among a panel of 11 SARS-CoV-2 lineages showing **(a)** three groups of post-vaccination sera from naïve mice and **(b)** three groups of post-vaccination sera from vaccine-experienced mice. SARS-CoV-2 lineages are shown as circles and sera are indicated as squares. Each square corresponds to sera of one individual mouse and is colored by the vaccine that mouse received (BNT162b2 XBB.1.5, JN.1, or KP.2). Antigenic distance is represented in both horizontal and vertical axes. Each square in the matrix represents 1 antigenic unit, which reflects a two-fold difference in neutralization titer. The points that are more closely together reflect higher cross-neutralization and are therefore antigenically more similar.

898
Figure 6. T Cell Mediated Immune Response Elicited by BNT162b2 XBB.1.5, JN.1 and

KP.2-Adapted Vaccines Administered to Vaccine-Experienced Mice. One-month after the

5th dose of BNT162b2 variant-adapted vaccine (XBB.1.5, JN.1, or KP.2), S-specific splenocytes

(n=5/group) were characterized by a flow cytometry-based intracellular cytokine staining (ICS)

assay. All samples were stimulated *ex vivo* with S peptide pools from the WT reference strain

and Omicron BA.4/5, XBB.1.5, JN.1, and KP.2 sublineages. **a-f** Graphs show the frequency of

CD4+ T cells expressing (**a**) IFN-γ , (**b**) TNF-α , and (**c**) IL-2 , and the frequency of CD8+ T

cells expressing (**d**) IFN-γ , (**e**) TNF-α , (**f**) IL-2 in response to stimulation with each peptide

pool across vaccine groups. Bars depict mean frequency + SEM.

Figure 7. **T Cell Mediated Immune Response Elicited by BNT162b2 XBB.1.5, JN.1 and**

KP.2-Adapted Vaccines Administered to Naïve Mice. At one-month post-second dose of

BNT162b2 variant-adapted vaccine (XBB.1.5, JN.1, or KP.2) (completion of primary series), S-

specific splenocytes (n=5/group) were measured by intracellular cytokine staining (ICS) assay.

All samples were stimulated *ex vivo* with S peptide pools from the WT reference strain and

Omicron BA.4/5, XBB.1.5, JN.1, and KP.2 sublineages. **a-f** Graphs show the frequency of

CD4$^+$ T cells expressing **(a)** IFN-γ, **(b)** TNF-α ,and **(c)** IL-2 , and the frequency of CD8$^+$ T cells

expressing **(d)** IFN-γ, **(e)** TNF-α , **(f)** IL-2 in response to stimulation with each peptide pool

across vaccine groups. Bars depict mean frequency + SEM.

Table 1. Emerging N-link glycosylation sites at JN.1 and KP.2 Spike proteins

Variant	Mutation	Peptide	% Glycosylation
JN.1	K356T	R.FASVYAW**N**R.**T**	99.8%
	H245N	R.FQTLLAL**N**R.**S**	99.6%
KP.2	K356T	W.**N**R**T**RISNCVADY.S	98.3%
	H245N	R.FQTLLAL**N**R.**S**	96.2%