# Nucleus accumbens D2-expressing neurons: Balancing reward and licking disruption through rhythmic optogenetic stimulation

Nikte Requejo-Mendoza[1,2], José-Antonio Arias-Montaño[1], Ranier Gutierrez[2,3]*

[1] Departamento de Fisiología, Biofísica y Neurociencias, Centro de Investigación y de Estudios Avanzados (Cinvestav), Ciudad de México, México

[2] Laboratory Neurobiology of Appetite; Departamento Farmacología, Centro de Investigación y de Estudios Avanzados (Cinvestav), Ciudad de México, México

[3] Laboratory Neurobiology of Appetite; Centro de Investigación sobre el Envejecimiento, Centro de Investigación y de Estudios Avanzados (Cinvestav Sede sur), Ciudad de México, México.

*Corresponding author: Email: ranier@cinvestav.mx

# Abstract

Dopamine D1 receptor-expressing neurons in the nucleus accumbens (NAc) are known to be critical for processing reward and regulating food intake. However, the role of D2-expressing neurons in this region remains less understood. This study employed optogenetic manipulations to investigate the role of NAc D2-expressing neurons in reward processing and sucrose consumption. Optogenetic activation of these neurons decreased sucrose preference (at 20 Hz), disrupted licking patterns (particularly at 8 and 20 Hz), and increased self-stimulation. Conversely, synchronizing stimulation with the animal's licking rhythm mitigated licking disruption and even increased sucrose intake, suggesting a rewarding effect. Furthermore, 20 Hz stimulation (but not 8 Hz) induced place preference in a real-time place preference (RTPP) test, establishing a long-lasting spatial reward memory. In contrast, inhibiting D2 neurons produced a negative hedonic state, although not reaching a complete aversion, influencing food choices in specific contexts. For instance, while the RTPP test per se was not sensitive enough to observe place aversion when mice could choose between consuming a high-fat diet (HFD) pellet in a context associated with or without inhibition of D2 neurons, they preferred to consume HFD on the non-inhibited side. This suggests that the palatability of HFD can unmask (but also overshadow) the negative hedonic state associated with D2 neuron inhibition. A negative reinforcement paradigm further confirmed the active avoidance behavior induced by D2 neuron inhibition. In conclusion, NAc D2 neuron inhibition induces a negative hedonic state, while activation has a dual effect—it is rewarding yet disrupts licking behavior—highlighting its complex role in reward and consummatory behavior. Importantly, self-paced stimulation entrained to licking rhythms offers a more "physiologically" relevant approach to brain activation.

35
36 **Keywords:** Reward; Sucrose; Licking; Aversion; Nucleus Accumbens Shell; Medium
37 Spiny Neurons.
38
39
40

# Introduction

The nucleus accumbens (NAc) is known to be involved in reward processing and food

intake.[1–3] While the role of dopamine D1 receptor-expressing neurons in the NAc has

been extensively studied, the role of D2 neurons remains unclear.[4,5] D1 medium spiny

neurons (MSNs) in the NAc are well-established as key players in processing rewards

and regulating food intake.[6,7] For instance, D1 neuron inhibition permits food

consumption, whereas activation suppresses licking behavior, a fundamental component

of the feeding-gating hypothesis of the NAc, particularly of the shell region.[8,9] In

contrast, feeding has been reported to have no modulatory effect on D2 receptor-

expressing neurons.[9,10] Accordingly, a recent study using rabies virus has traced that

D1 NAc neurons receive direct inputs from taste bud cells on the tongue and send direct

projections to the reticular formation in which the Central Pattern Generator (CPG) for

rhythmic licking is controlled. In contrast, the same study showed that D2 neurons are not

directly connected with the CPG controlling tongue movements.[11]

Beyond its involvement in food consumption, the NAc has been recognized as a crucial

player in reward processing since the groundbreaking work of James Olds (1956) on the

brain pleasure centers.[12] This role is also supported by evidence that mice will self-

stimulate to activate all glutamatergic inputs to the NAc shell, [13,14] indicating the

capability of glutamate to modulate reward and even suppress feeding behavior,

presumably by influencing the activity of either D1 or D2 receptor-expressing neurons.[14]

The classic view of D1- and D2-expressing neurons posited opposing roles in feeding[15]

and in reward, D1 neuron activation being rewarding and D2 activation aversive, with

inhibition leading to the opposite effects.[16] However, current research suggests a more

complex dynamic. While D1-expressing neurons predominantly drive positive

reinforcement and reward behaviors,[16–18] D2-expressing neurons appear to have

more complex, bidirectional effects, depending on the context and neural stimulation

patterns.[4]  Soares-Cunha work further challenged the classic view, demonstrating that

brief (1 s, 40 Hz) optogenetic stimulation of D1- or D2-expressing neurons is rewarding,

whereas prolonged stimulation (60 s) elicits aversion.[19] This highlights that the

rewarding or aversive nature of D1 and D2 neuron activity hinges on stimulation

parameters.[5] Thus, while the role of NAc D1-expressing neurons in reward is more

established,[20,21] the contribution of D2-expressing neurons yields variable results

across different behaviors and experimental setups.[4,5,11,17]

This study employed optogenetic manipulations to investigate the specific role of NAc D2-

expressing neurons in reward processing and consummatory behavior related to sucrose

and HFD intake. By exploring multiple frequency-dependent parameters for optogenetic

manipulation, we aimed to provide further insights into how D2 neurons fine-tune reward

processing and consummatory behaviors. Our data demonstrate the complex nature of

NAc D2 neuron activation, revealing its dual role in promoting reward while disrupting

certain aspects of consummatory behavior. In contrast, NAc D2 neuron inhibition induces

a negative hedonic value. Furthermore, our findings highlight the importance of

stimulation parameters in understanding NAc D2 neuron function

# Results

## Optogenetic activation of NAc D2-expressing neurons at 20 Hz reduces sucrose preference by disrupting the microstructure of licking and increasing self-stimulation

A two-bottle freely licking preference test was conducted to assess the role of NAc D2-expressing neurons in modulating sucrose preference. This experiment was conducted under three conditions, each lasting two days: 1) Baseline (No-laser), 2) Laser activated by the sucrose bottle, and 3) Laser activated by the water bottle. In the D2$^{tdTomato}$ (control) and D2$^{ArchT}$ (inhibition) groups, animals maintained a stable preference for sucrose over water throughout all conditions. However, in the D2$^{ChR2}$ (activation) group, optogenetic stimulation at 20 Hz during sucrose consumption significantly reduced sucrose preference (**Fig 1c** [* one-way ANOVA; main factor: groups (Laser sucrose condition); $F_{(2, 62)}$ = 9.20, p < 0.001]), a *post hoc* analysis further revealed that D2$^{ChR2}$ exhibited a significant decrease compared to D2$^{tdTomato}$ and D2$^{ArchT}$ (all *p's < 0.003). This reduction in sucrose preference was accompanied by a marked increase in water preference, as indicated by fewer total sucrose licks and more water licks in the D2$^{ChR2}$ group (**Fig 1d** [* one-way ANOVA; main factor: groups (within Laser sucrose condition); $F_{(1, 40)}$ = 13.81, p < 0.001], [# one-way ANOVA; main factor: conditions; for water $F_{(1, 44)}$ = 24.13, p < 0.001; for sucrose $F_{(1, 44)}$ = 13.30, p < 0.001]). As well as an increase in total licks (water+ sucrose) within the same condition for the D2$^{ArchT}$ group [@ one-way ANOVA; main factor: conditions; $F_{(1,39)}$ = 4.17, p = 0.04]. Additionally, when stimulation was paired with the

water bottle, the D2$^{ChR2}$ group exhibited a significant increase in sucrose licking compared

to the baseline condition [# one-way ANOVA; main factor: conditions; $F_{(1,36)}$ = 7.75, p =

0.008], the D2$^{tdTomato}$ group [* one-way ANOVA; main factor: groups; $F_{(1,26)}$ =27.91,

p<0.001] and generating a significant increase for total licks [@ one-way ANOVA; main

factor: groups; $F_{(1,26)}$ = 31.66, p<0.001], as well the D2$^{ArchT}$ group showed a significant

increase compared with D2$^{tdTomato}$ group [* one-way ANOVA; main factor: groups; $F_{(1,21)}$

= 12.40, p = 0.002], resulting in a significant increase for total licks [@ one-way ANOVA;

main factor: groups; $F_{(1,21)}$ = 11.09, p =  0.003]. This increase in licking behavior was

accompanied by a higher number of trials initiated (a measure of self-stimulations) (**Fig**

**1e** [*one-way ANOVA; main factor: groups; $F_{(2, 62)}$ = 11.67, p < 0.001]), suggesting a

rewarding effect induced by 20 Hz stimulation. Additionally, comparisons between the

laser and No-laser in Baseline condition in the D2$^{ChR2}$ group showed significant

differences for sucrose licks (L_s) (paired t-test; $t_{(44)}$ = 2.72, p = 0.0092) and water licks

(L_w) (paired t-test; $t_{(36)}$ = 3.71, p = 0.0007). A deeper analysis of the licking

microstructure revealed that D2 neuron activation disrupted normal licking patterns

(**Table 1**). Heatmaps of licking behavior (**Figs 1f-h**) revealed consistent licking patterns

across all groups during baseline conditions (**Fig 1f**). However, optogenetic stimulation

paired with sucrose consumption (**Fig 1g**) induced a significant decrease in lick bout

durations, specifically within the D2$^{ChR2}$ group [one-way ANOVA; main factor: groups

(within Laser sucrose condition) $F_{(1, 44)}$ = 35.90, p < 0.001] and reduced bout size [one-

way ANOVA $F_{(1, 44)}$ = 13.30, p < 0.001], accompanied by increased number of licking

bouts. A similar fragmentation in licking microstructure was observed when water was

paired with stimulation (**Fig 1h**) for bout durations [one-way ANOVA; groups; $F_{(1, 36)}$ =

7.73, p < 0.001] and bout size [one-way ANOVA; groups; $F_{(1, 36)}$ = 10.10, p < 0.001].

Further analysis of licking microstructural parameters (**Table 1**) confirmed that

optogenetic activation of D2 neurons significantly reduced bout size and duration while

increasing bout frequency for water in the D2[ChR2] group. These findings indicate that

optogenetic activation of NAc D2-expressing neurons disrupts normal licking behavior

and induces self-stimulation, suggesting a rewarding effect. This suggests that the

observed decrease in sucrose preference likely stems from the combined rewarding and

lick-impairing effects of optogenetic stimulation.

## Optogenetic frequency-dependent modulation of licking behavior by NAc D2-expressing neurons

To assess the impact of stimulation frequency on licking behavior, we used a freely licking

task with a laser scanning frequency test, as previously reported, to examine frequency-

dependent activation of D2 NAc neurons.[14] The task operates in a cycle of three

epochs: Before (4 licks), Laser (1 s), and Timeout (1 s). A new cycle is triggered by four

consecutive licks immediately after the Timeout epoch. In this task, control animals

(D2[tdTomato]) exhibited consistent lick rates across all laser frequencies (**Fig 2b**). In contrast,

D2[ChR2] activation reduced the lick rate compared to the No-laser trials, with stronger

reductions observed at higher frequencies, especially 8, 15, and 20 Hz (**Figs 2c, 2f, and**

**2i**). Cumulative lick distributions during laser epochs revealed significant reductions in the

D2[ChR2] group across all frequencies compared to No-laser trials [# one-way ANOVA; main

factor: types of trials; $F_{(4, 625)}$ = 38.94, p < 0.001, all Tukey- Kramer; p<0.001],  and the

control group (D2[tdTomatto]) with the most pronounced reductions at 8 Hz [* one-way

ANOVA; main factor: groups; $F_{(4, 354)} = 33.98$, $p < 0.001$], 15 Hz [* one-way ANOVA; main

factor: groups; $F_{(4, 354)} = 39.93$, $p < 0.001$], and 20 Hz [* one-way ANOVA; main factor:

groups; $F_{(4, 354)} = 52.72$, $p < 0.001$] (all [#,@] p's $< 0.001$) (**Fig 2i**). Additionally, the 5 Hz

stimulation allowed significantly more licks than higher frequencies (8, 15, and 20 Hz)

within the D2$^{ChR2}$ group [one-way ANOVA; 5 Hz vs. other frequencies; $F_{(3, 187)} = 8.79$, p <

0.001]. Inhibition of D2 neurons (D2$^{ArchT}$) at 5 Hz stimulation generated significantly more

licks compared to the control group (**Figs 2j**) [*one-way ANOVA; main factor: groups; $F_{(2,}$

$_{354)} = 18.69$, $p < 0.001$], while there were no significant differences at 8, 15, or 20 Hz

compared to the control group (**Fig 2j**). During the Laser epoch, D2$^{ChR2}$ animals displayed

significantly fewer licks at 8, 15, and 20 Hz compared to No-laser trials [#one-way ANOVA;

main factor: trial types; $F_{(4, 130)} = 14.07$, $p < 0.001$] (**Fig 2k**). Inhibition of D2 neurons

(D2$^{ArchT}$) at 20 Hz resulted in significant suppression of licking behavior [#one-way

ANOVA; main factor: trial types; $F_{(4, 160)} = 3.44$, $p = 0.009$] (**Fig 2k**). Stimulation effects

persisted into the post-laser timeout period, with a tendency for lick frequency to decrease

with increasing stimulation frequency (**Fig 2l**). Although this change was less pronounced

during the Timeout epoch, the tendency suggests a continued stimulation influence.

These results suggest that NAc D2 neuron activity modulates licking behavior in a

frequency-dependent manner, acting as a "brake" on consummatory actions and

decreasing sucrose intake. Higher stimulation frequencies, particularly at 15 and 20 Hz,

produce stronger suppression and have long-lasting effects on reinitiating reward-seeking

behavior.

**Self-paced (unlike experimenter-imposed frequency) optogenetic stimulation of D2 neurons attenuates disruption of licking in a brief access taste test**

We used the brief access taste test (BATT) to investigate whether the frequency-dependent effects of stimulation on licking behavior persisted under a self-paced lick-driven rhythmic stimulation pattern. In this task, mice could lick for a 10% sucrose (Suc) solution during a 4 s reward period with different optogenetic stimulation trials (No-laser, lick-paired (0 ms delay), 8 and 20 Hz). The D2$^{tdTomato}$ (control) (**Figs 3b, 3e**) group maintained consistent licking behavior across all trials, showing no significant disruption by these modulations. In contrast, based on the optogenetic stimulation pattern, the D2$^{ChR2}$ (activation) group exhibited a significant difference in licking behavior (**Figs 3c, 3f**). Notably, 20 Hz stimulation significantly reduced the licking rate [#one-way ANOVA; main factor; trial types $F_{(4, 416)} = 22.3$, $p < 0.001$], while lick-paired stimulation preserved a licking rate similar to the No-laser condition, highlighting the importance of self-timing in optogenetic stimulation and its effects on licking behavior.

These effects were also reflected in cumulative licks across the session, where the D2$^{ChR2}$ group exhibited significantly fewer licks during stimulation at 20 Hz than the other trial types (**Fig 3f** [#one-way ANOVA; main factor: trial types; $F_{(3, 416)} = 22.63$, $p < 0.001$]). A *post hoc* analysis confirmed that the No-laser and lick-paired conditions significantly differed in the D2$^{ChR2}$ group across trial types [#$p < 0.001$], underscoring the impact of stimulation frequency on licking behavior. The 8 Hz imposed stimulation also reduced cumulative licks compared to the No-laser condition [$p < 0.001$], although this reduction

was less pronounced than that induced by the 20 Hz stimulation [[1]p<0.001]. While the

D2$^{ArchT}$ (inhibition) group significantly increased the cumulative licks in the total licks and

in the lick paired, 8, 20 Hz trials (**Figs 3d, 3g**) compared to the control group (all *p's<0.01)

(**Figures 3e, 3g**).

We plotted the total licks per session to observe the overall impact on licking behavior

(**Fig 3h**). This plot further emphasized these differences across groups [*one-way

ANOVA; main factor: groups $F_{(2, 65)}$ = 7.78, p <0.001]. In the D2$^{ChR2}$ group [*one-way

ANOVA; main factor: trial types $F_{(3, 80)}$ = 7.668, p <0.001], total licks during the 20 Hz

condition were significantly reduced compared to the No-laser([#]p <0.001) and lick-paired

conditions([@]p <0.001) ). Conversely, total licks in the D2$^{tdTomato}$ and D2$^{ArchT}$ groups

showed no significant differences across stimulation types, confirming that optogenetic

activation of D2-expressing neurons, rather than inhibition, more strongly affected licking

behavior.

Importantly, the increased licking behavior observed during control trials in the D2$^{ChR2}$

group may be explained by the sequential nature of the task. Given that each trial must

be completed before passing to the next one, the animals could display an enhanced

motivation effect from previous trials where the D2 neurons were activated (panel **h**).

Microstructural analysis of licking behavior (**Fig 3i**) revealed, at 8 [*one-way ANOVA;

main factor: groups; $F_{(2, 65)}$ = 12.50, p <0.001]  and 20 Hz [*one-way ANOVA; main factor:

groups; $F_{(2, 65)}$ = 10.03, p <0.001] frequencies, more fragmented licking in the D2$^{ChR2}$

group characterized by shorter bout duration and increased bout number [*one-way

ANOVA; group; $F_{(2, 65)}$ = 16.95, p < 0.001]. Interestingly, despite the disrupted licking

behavior, the D2$^{ChR2}$ group initiated more trials than the other groups (**Fig 3j** [*one-way

ANOVA; groups; $F_{(2, 65)}$ = 14.14, p < 0.001]), indicating that 20 Hz stimulation disrupted

consummatory licking but increased the persistence in the task. **Table 3** shows that high-

frequency stimulation (20 Hz) significantly reduced lick rate, bout size, and bout duration

in the D2$^{ChR2}$ group while increasing the number of licking bouts. This suggests that while

high-frequency stimulation disrupts the typical licking behavior, it paradoxically enhances

the animal's incentive motivation to persist in seeking the reward, as evidenced by

increased trial initiation.

## 222 Lick-paired optogenetic activation with delayed timing of NAc
## 223 D2 neurons slightly disrupts the licking microstructure but
## 224 enhances sucrose intake

Given the advantage of lick-paired over imposed stimulation, we next assessed the effect

of lick-paired but using multiple delays relative to the lick tongue protrusion-retraction

cycle. To this aim, we used a variation of the brief access test, where optogenetic

activation was applied with different delays relative to the tongue contacting the sipper (0,

50, and 100 ms) after each lick, alongside no-laser trials that served as controls. Licking

rates remained relatively stable along the reward epoch across all trial types and groups

(**Figs 4b-d**), indicating that delayed activation or inhibition had minimal effects on the

overall licking behavior. However, despite these stable rates, the D2$^{ChR2}$ group exhibited

a significantly higher total number of cumulative licks than the D2$^{tdTomato}$ group across all

trial types (**Figs. 4e-f** [*one-way ANOVA; main group: No-laser $F_{(2, 312)}$ = 11.05; p < 0.001;

main group: 0 ms $F_{(2, 312)}$ = 3.29, p = 0.04; main group: 50 ms $F_{(2, 312)}$ = 12.49, p <0.001;

main group: 100 ms $F_{(2, 312)}$ = 18.67, p <0.001; main group: Total cumulative licks $F_{(2, 1257)}$

= 40.15, p <0.001]. The D2$^{ArchT}$ group also showed significant increases in total

cumulative licks (*p = 0.002), particularly in the 100 ms delay trials (*p < 0.001). These

findings indicate a significant increase in the total number of cumulative licks for the

D2$^{ChR2}$ group, suggesting greater overall licking activity than the other groups.

Although overall, the lick-paired protocol was more 'physiological' than imposed

frequency stimulation, we found that the inter-lick interval was significantly longer in the

D2$^{ChR2}$ compared to the D2$^{tdTomato}$ group across all conditions (**Fig 4h** [*one-way ANOVA;

main factor: group: No-laser $F_{(2, 30)}$ = 8.35, p = 0.01;  0 ms $F_{(2, 30)}$ = 13.95, p < 0.001; 50

ms $F_{(2, 30)}$ = 7.57, p = 0.002;  100 ms $F_{(2, 30)}$ = 13.55, p <0.001]). This behavioral protocol

also revealed the disruptive effect of optogenetic activation on rhythmic oromotor

components of licking behavior, even with lick-paired stimulation patterns.

Regarding licking bouts, the D2$^{ChR2}$ group showed a significantly higher number of bouts

than the D2$^{tdTomato}$ group across all conditions (**Fig 4i** [*all p's < 0.001]). However, the size

of each lick bout was smaller in the D2$^{ChR2}$ group compared to the D2$^{tdTomato}$ group under

the 0 ms delay condition (**Fig 4j** [*one-way ANOVA; main factor: group $F_{(2,30)}$ = 6.52, p =

0.01]).

Finally, despite the evident fragmentation of licking behavior, the D2$^{ChR2}$ group initiated

significantly more trials compared to both the D2$^{tdTomato}$ and D2$^{ArchT}$ groups (**Fig 4k** [one-

way ANOVA; main factor: group; $F_{(2, 30)}$ = 15.41, p < 0.001]), underscoring a paradoxical

increase in task engagement and reward driven by D2 neuron activation. These findings

indicate that although self-paced rhythmic optogenetic activation of NAc D2-expressing

neurons affects licking microstructure, this effect is less pronounced than when the

experimenter imposes the stimulation frequency. Interestingly, this self-paced activation

paradoxically increases motivation and task persistence, leading to greater sucrose

intake, especially when the activation is paired with physiological licking rhythms.

## 262 Optogenetic activation of NAc D2-expressing neurons at 20 Hz 263 (but not 8 Hz) drives place preference and enhances 264 exploratory behavior

Next, we explored how optogenetic activation of NAc D2-expressing neurons modulates

spatial contextual rewards beyond licking behavior. We assessed the effects of different

stimulation frequencies on place preference using a real-time place preference (RTPP)

assay. Optogenetic stimulation was delivered at 8 or 20 Hz in D2$^{ChR2}$ and D2$^{tdTomato}$

groups in this paradigm.

Heatmaps illustrating the time spent in a spatial location (in yellow; **Figs 5b-e**) revealed

a clear preference for the stimulated side in the D2$^{ChR2}$ group when stimulated at 20 Hz

(**Fig 5c**). Mice in this group spent significantly more time on the stimulated side during

both sessions compared to the D2$^{tdTomato}$ control and the 8 Hz stimulation condition (**Fig**

**5f** [*one-way ANOVA; main factor: group; $F_{(3, 22)}$ = 7.19, p < 0.001]). In contrast, no

significant place preference was observed in the D2$^{tdTomato}$, D2$^{ChR2}$ 8 Hz, and D2$^{tdTomato}$

20 Hz groups.

The number of transitions between the two sides of the chamber did not differ significantly

between groups (**Fig 5g**). However, the D2$^{ChR2}$ 20 Hz group exhibited significantly higher

velocity on the non-stimulated side compared to other groups(**Fig 5h** [*one-way ANOVA;

main factor: group; $F_{(3, 22)}$ = 8.85, p < 0.001]), and conditions ($^{\#}$p =0.01) while the total

distance traveled was greater on the stimulated side (**Fig 5i** [# t-test (paired); t(6) = 3.26,

p = 0.01]) and against other groups [*one-way ANOVA; main factor: group, $F_{(3, 22)}$ = 5.36,

p = 0.006]. These findings indicate that the increased locomotion was specific to the

stimulated chamber, suggesting that the 20 Hz stimulation enhanced both place

preference and movement within the stimulated area. These results highlight that 20 Hz

stimulation of D2-expressing neurons in the NAc induces a robust rewarding effect, as

evidenced by place preference, and modulates locomotor behavior, leading to increased

movement within the stimulated environment.

**NAc D2 neuron inhibition induces a negative hedonic state**

**(but not aversion) that is unmasked by an HFD choice**

**paradigm**

To investigate the role of NAc D2-expressing neurons in context-dependent behaviors,

we first examined their influence on place preference using the RTPP task. Mice freely

explored a chamber divided into two distinct halves, with one side paired with optogenetic

stimulation. In this task, optogenetic inhibition (continuous stimulation) was delivered to

D2$^{ArchT}$ mice, while D2$^{tdTomato}$ controls received 20 Hz stimulation (**Fig 6a**, left; D2$^{tdTomato}$

data reused in **Fig 5**). Heatmaps representing animal spatial exploration showed no

significant preference for the stimulated side in either group (**Fig 6a**, middle).

Quantitatively, the percentage of time spent on the stimulated side was similar across

both sessions, with no significant differences between groups (**Fig 6a**, right). These data

suggest that silencing NAc D2-expressing neurons generated, *per se,* neither aversion

nor preference under this standard RTPP protocol.

Next, we assessed the effects of D2 neuron modulation on feeding behavior using an

open-loop paradigm and HFD alone in one spatial context. In this protocol, mice were

placed in an arena with HFD pellets on both opposite sides, while optogenetic stimulation

was applied in 4 s on, 2 s off intervals for D2$^{tdTomato}$ and D2$^{ChR2}$ (20 Hz, 10 ms pulse width)

and continuous pulse stimulation for D2$^{ArchT}$ (**Fig 6b**, left). In this context, total HFD intake

did not significantly differ between the control (D2$^{tdTomato}$), activation (D2$^{ChR2}$), and

inhibition (D2$^{ArchT}$) groups (**Fig 6b**, right), indicating that D2 neuron activation or inhibition

*per se* was not sufficient to influence HFD feeding behavior in a neutral environment.

Finally, in a variation of the RTPP task, we explored whether NAc D2-expressing neurons

modulate place preference and HFD consumption in a conditioned context, where one

side of the chamber was paired with optogenetic stimulation, and the mice could choose

whether or not to eat HFD pellets on the stimulated side on the box (**Fig 6c**). During

conditioning, D2$^{ChR2}$ mice developed a significant preference for the stimulation-paired

side, which persisted into the post-conditioning test (**Fig 6d** [all $^{*}$p's <0.006, see

Supplementary Statistical Table]). Conversely, D2$^{ArchT}$ mice initially avoided the

stimulation-paired side during acquisition [$^{#}$one-way ANOVA; main factor: sessions; $F_{(4, 20)} = 18.78$, $p < 0.02$], but this behavior did not persist in the test phase (**Fig 6d**).

Analysis of total HFD intake revealed a tendency for reduced intake in the D2$^{ChR2}$ group

during activation, while the D2$^{ArchT}$ group exhibited increased consumption during

inhibition (**Fig 6e**). In particular, the D2$^{tdTomato}$ (**Fig 6f**) and D2$^{ChR2}$ (**Fig 6g**) groups had no

change in the HFD intake between the stimulated and non-stimulated side, while D2$^{ArchT}$

mice significantly increased HFD intake on the non-stimulated side during the final

acquisition days, an effect that persisted into the test phase (**Fig 6h** [#one-way ANOVA;

main factor: side; $F_{(4,25)}$ = 4.14, p = 0.01] [t-test; Adq2 $t_{(5)}$ = -4.40, p = 0.007; Adq3 $t_{(5)}$ = -

4.39, p = 0.007]). This suggests that inhibiting NAc D2-expressing neurons induces

negative hedonics, increasing palatable food consumption on the non-stimulated side.

These findings indicate that NAc D2 neuron inhibition induces a negative hedonic value

that an HFD choice test can unmask.

## 331    Inhibition of NAc D2 neuron drives negative reinforcement

After establishing that optogenetic activation of NAc D2-expressing neurons induces

place preference, we sought to determine whether inhibition of these neurons elicits a

negative hedonic value using a negative reinforcement task. In this more sensitive

paradigm, mice had to enter an 'off' zone to end optogenetic stimulation delivered in the

"on" zone. Stimulation was applied at 20 Hz for the D2$^{ChR2}$ and D2$^{tdTomato}$ groups and

continuously applied for the D2$^{ArchT}$ group (**Fig 7a**). Despite the natural tendency of mice

to explore, which led to most of their time being spent in the 'on' zone, the D2$^{ArchT}$ group

showed a significant shift in behavior, with increased time spent in the 'off' zone compared

to the D2$^{tdTomato}$ and D2$^{ChR2}$ groups (**Figs 7b, 7c** [*one-way ANOVA; main factor: groups;

$F_{(2, 33)}$ = 4.27, p = 0.03]), indicating that sustained inhibition of NAc D2-expressing neurons

generated a negative valence, prompting mice to avoid the area of stimulation.

Representative heatmaps from the third session (**Figs 7d-f**) further illustrate this behavior,

emphasizing the negative reinforcement effect of D2 neuron inhibition. These results

show that inhibiting NAc D2 neurons elicits a negative hedonic value, with mice actively

avoiding areas where stimulation occurs. This active avoidance behavior highlights the

role of NAc D2-expressing neurons in driving negative reinforcement.

# Discussion

This study demonstrates that dopamine D2 receptor-expressing neurons in the NAc play

a dual role in reward processing and ingestive behavior. Optogenetic activation of these

neurons at higher frequencies disrupts the licking microstructure, decreasing sucrose

preference while paradoxically increasing self-stimulation. This seemingly contradictory

effect highlights the complex influence of D2 neuron activity, which can simultaneously

enhance incentive motivation while disrupting specific oromotor behaviors. Interestingly,

synchronizing stimulation with the animal's natural licking rhythm minimized licking

disruptions and promoted sucrose intake. Regarding spatial contextual reward, high-

frequency stimulation (20 Hz) was sufficient to drive place preference, confirming the

rewarding properties of D2 neuron activation. Conversely, inhibiting these neurons led to

a negative hedonic state (but not complete aversion), affecting food choices in specific

contexts, particularly when a high-fat diet (HFD) was involved. These findings

demonstrate the dual role of NAc D2-expressing neurons in regulating reward and

consumption, highlighting the potential of entrained, self-paced stimulation that leverages

the naturalistic licking theta rhythm for precise activation of these neurons.

## Activation of NAc D2 neurons reduces sucrose preference and increases self-stimulation

Optogenetic activation of NAc D2 neurons at 20 Hz disrupted sucrose preference by fragmenting the licking microstructure (**Fig. 1**). This finding is consistent with previous studies emphasizing the role of the NAc in orchestrating consummatory actions and processing palatability-related information.[8] In this regard, electrophysiological and calcium imaging studies in rats and mice have shown that a subset of NAc neurons encodes sucrose concentration and palatability, while other subsets exhibit oromotor responses entrained to the licking rhythm,[1,2,22,23] highlighting that the NAc is embedded in dynamic circuits that regulate hedonic processing and goal-directed behaviors.[7] While D1 neurons have traditionally been associated with feeding,[8,9] our findings indicate that experimentally imposed 20 Hz activation of D2 neurons particularly disrupts ingestive behavior.

## Frequency-dependent disruption of licking behavior by NAc D2-expressing neurons

Prado et al. (2016) showed for the first time that optogenetic manipulations of glutamatergic inputs to the NAc are rewarding but also stopped licking behavior in a frequency-dependent manner, reducing lick duration and frequency, resembling our observation of D2 neurons activation shortened lick bout durations and reduced bout sizes (**Fig 2**).[14] Interestingly, these disruptions were frequency-dependent, with higher frequencies (15 and 20 Hz) producing more pronounced effects, supporting the notion

that NAc D2-expressing neurons would be, at least, one target of glutamatergic inputs to

fine-tune the intensity and timing of consummatory actions.[14]

In a detailed frequency test, we found that higher optogenetic stimulation frequencies (15

and 20 Hz) produced a significant suppression of licking behavior compared to lower

frequencies. These results support that D2-expressing neurons act as modulators of

consummatory actions depending on the stimulation pattern, as highlighted by Soares-

Cunha et al. (2019). In their study, brief stimulation was rewarding, but prolonged

stimulation became aversive and induced more pronounced behavioral impairments.[5]

Our study suggests that D2-expressing neurons play a dual role in regulating

consummatory behavior. While they enhance the incentive motivation to self-stimulate,

they simultaneously impair the execution of licking, as shown by increased self-

stimulation despite impaired licking (**Fig 4**). These results also indicate that NAc D2-

expressing neurons can modulate ingestive behavior through frequency-dependent

mechanisms, aligning with the concept of the NAc as a central hub integrating reward

and feeding behaviors, where the activity of D2-expressing neurons can either promote

or hinder consumption depending on the specific stimulation pattern.[8,14,15]

Our data can also reconcile some differences in literature. For example, O'Connor et al.

(2015) found that 20 Hz stimulation of D1 neurons abruptly impaired licking, [9] whereas

Sandoval et al. (2023) did not observe licking impairment using a lick-paired self-

stimulation protocol of D1 neurons, like the one used in this study. Surprisingly, like us,

they also observed a slight licking impairment using lick-paired self-stimulation of D2-

expressing neurons, [11] suggesting a unique property of self-rhythmic licking to stimulate

the NAc. These results also support the notion that rhythmic licking acts as an internal

clock to synchronize brain activity across different brain regions.[24]  While our findings

show that D2-expressing neurons impair rhythmic licking, particularly at high frequencies,

the precise circuit mechanisms underlying this disruption remain unclear. This is because

previous research has shown that lesioning D1, but not D2, neurons impairs licking

microstructure,[11] and rabies tracing studies indicate that, unlike D1 neurons, D2

neurons do not directly connect to taste buds or the CPG for licking.[11] One possibility

is that D2-expressing neurons disrupt licking behavior indirectly by modulating D1 neuron

activity, potentially through lateral inhibition.[25,26] This hypothesis warrants further

investigation, which could employ cell-type-specific manipulations and circuit-tracing

techniques to delineate the precise pathways and mechanisms involved.

## Self-paced stimulation enhances task engagement and reduces licking disruption

One of the key insights from the present study is the better preservation of licking behavior

and enhanced task engagement when stimulation was synchronized with natural licking

rhythms (**Figs 3,4**), highlighting the flexible nature of reward-seeking behaviors governed

by the NAc. This finding aligns with previous research showing that mice will self-stimulate

glutamatergic inputs to the NAc and that these glutamatergic inputs are suppressed

during feeding.[14,27] This suggests that NAc circuits dynamically adapt to regulate both

consumption and the pursuit of rewards.[13,14,27] Similarly, Cole et al. (2018) reported

that allowing animals to self-stimulate D2 neurons by licking a spout, mirroring their

natural behavior, led to increased engagement and fewer disruptions compared to

externally imposed stimulation triggered by entering a specific spatial location.[17] These

findings resonate with the concept that D2 neuron activity can modulate reward by

integrating motor information into reward-seeking,[28] promoting goal-directed behaviors

while maintaining hedonic value.[1] This supports the view that stimulation parameters,

particularly self-paced stimulation, are crucial in determining the behavioral outcomes of

D2 neuron activation.[4]

## High-frequency activation of NAc D2 neurons drives place preference and enhances exploratory behavior

In addition to modulating licking behaviors, our study revealed that high-frequency

activation of NAc D2 neurons (20 Hz) also induced robust place preference and

significantly enhanced exploratory behavior, indicating a strong, rewarding effect (**Fig 5**).

While place preference has traditionally been associated with D1 neurons,[20] our results

extend these findings to D2-expressing neurons, indicating that these neurons play a

more significant role in reward-seeking behaviors than previously assumed. Our data also

supports the observation of Cole et al. (2018) using a lick-spout self-stimulation task. They

found that mice self-stimulated both D1 and D2 neurons in NAc, although D1 excitation

was more rewarding than D2 neurons. They also suggest that the excitation of D2

neurons in NAc supports self-stimulation under some tasks (using a lick-spout) but fails

under others.[17] Oginsky et al. (2016) showed that NAc D2-expressing neurons are

involved in both place preference and exploratory behaviors, particularly in contexts

involving palatable food, supporting our findings of enhanced exploratory drive during 20

Hz stimulation.[29]

The enhanced exploratory behavior observed in the stimulated context suggests that D2-

expressing neurons may contribute to a broader behavioral repertoire beyond reward.[15]

This indicates a potential role for D2-expressing neurons in modulating spatial exploration

and context-dependent behaviors, contributing to the animal's overall motivational state

during task engagement.[26]

## Inhibition of NAc D2 neurons decreases palatable food intake and drives context-dependent negative reinforcement

Interestingly, while optogenetic activation produced place preference, inhibition of D2

neurons also revealed a role in processing hedonic value (**Fig 6**). Specifically, inhibition

led to a preference for HFD consumption on the non-stimulated side of a conditioned

chamber, suggesting that D2 neuron suppression induces a negative hedonic value that

palatable food can partially mask. Yang et al. (2020) found that inhibition of D2 neurons

unmasked negative hedonic value in specific contexts, mirroring our findings that D2

inhibition during HFD reveals a negative hedonic signal that palatable food can

override.[3] Cole et al. (2018) reported that optogenetic activation of D2 neurons leads to

a motivational state where both reward-seeking and sometimes negative avoidance

responses coexist, suggesting that D2-expressing neurons exhibit a context-dependent

ambivalence response.[17] These findings are consistent with earlier studies

demonstrating that D2 neurons contribute to positive and negative valence processing

depending on the context and task structure.[4,5] This result emphasizes the context-

dependent nature of D2 neuron function, where inhibition generates a negative hedonic

state, further supporting the idea that D2-expressing neurons can induce a negative

reinforcement (**Fig 7**). Our findings extend the current understanding of NAc D2-

expressing neurons, showing their critical role in balancing reward and consummatory

behavior across diverse contexts of motivation and motor control. This duality highlights

their complex contribution to reward circuits, aligning with the broader literature on their

dual functions.[15,17,30]

## Limitation of the study

Although our findings provide new insights into the role of NAc D2-expressing neurons in

reward and consummatory behaviors, it is important to acknowledge some limitations.

Firstly, while this study focused on D2-expressing neurons, future research should

investigate the role of D1-expressing neurons with similar paradigms to understand better

the network dynamics governing these behaviors. Future work should explore how D1

and D2 neurons interact to regulate both homeostatic and hedonic feeding, building upon

work by Walle et al. (2024), using simultaneous manipulations of D1 and D2 expressing

neurons that highlighted distinct but complementary roles for these neurons in energy

balance and reward-seeking.[15] Secondly, while we observed significant frequency-

dependent effects, the precise circuits and downstream targets mediating these effects

require further investigation. It is important to recognize that our use of optogenetics to

modulate NAc D2-expressing neurons (or any other cell type) while providing valuable

insights may not fully capture the 'physiological' role of endogenous neural activity and

naturalistic reward behaviors.[31] This study found that behavioral responses are highly

dependent on optogenetic stimulation parameters. Therefore, chemogenetic

manipulations are also expected to produce varying behavioral effects as different

stimulation approaches engage neuronal circuits in distinct ways.[15] The observation

that different stimulation parameters of D2-expressing neurons result in different

behavioral outputs aligns with the concept of optoception. We recently reported the

optoception phenomenon, suggesting that mice could perceive each stimulation pattern

slightly differently, providing a basis for mice's ability to identify and discriminate between

various optogenetic stimulation frequencies.[31] Finally, because cholinergic

interneurons in the NAc also express D2 receptors, future studies employing

intersectional genetic techniques are needed to differentiate the contribution of these

interneurons from those of D2 MSNs. This will provide a more precise understanding of

the specific roles of these distinct cell populations in reward and consumption.

## Conclusion

In summary, our findings demonstrate the dual role of NAc D2-expressing neurons in

modulating reward and consummatory behavior, where activation can simultaneously

enhance incentive motivation to self-stimulate and disrupt licking behavior. These results

highlight the importance of stimulation parameters, frequency, timing, and agency of

actions in shaping the behavioral outcomes of D2 neuron activity. Inhibition of these

neurons induces a negative hedonic state that is (un)masked in specific contexts, such

as during HFD consumption. Thus, NAc D2-expressing neurons act as critical

modulators, balancing reward and licking disruption through rhythmic optogenetic

stimulation, providing new insights into their role in reward circuits. These findings deepen

our understanding of the NAc role in reward circuits and may provide potential avenues

for exploring therapeutic interventions for disorders related to reward processing, such as

addiction[32] or obesity.[33]

# Materials and methods

## Ethical Considerations

521 All experimental procedures were approved by the Cinvestav Animal Care and Use

522 Committee (CINVESTAV) under permit number [0080-14] and conducted the Guide for

523 the Care and Use of Laboratory Animals of the National Institutes of Health.

## Animals

Drd2-Cre (B6.FVB(Cg)-Tg(Drd2-cre)ER44Gsat/Mmcd) mice of both sexes (7-10 weeks

old, 20-25 g at the beginning of the experiment; MMRC, stock number: 032108) were

single-housed in a temperature-controlled room (22 ± 1°C) with a 12-h light/dark cycle.

Mice had *ad libitum* access to a standard chow diet (PicoLab Rodent Diet 20, MO, USA)

and water unless otherwise specified for a particular behavioral paradigm. After surgery,

mice were single-housed and maintained in a temperature-controlled environment with a

12-h light/dark cycle. Post-surgical care included the administration of analgesic

(lidocaine hydrochloride) and antibiotic (enrofloxacin). Mice were monitored daily for signs

of pain, distress, or discomfort to ensure animal welfare. Efforts were made to reduce the

number of animals used by implementing habituation protocols and non-invasive

techniques, such as video tracking systems for behavioral monitoring.

## Viral constructs

For cell-type-specific manipulation of D2-expressing neurons in the NAc, mice were injected with Cre-dependent adeno-associated viral (AAV) vectors carrying genes encoding Channelrhodopsin (ChR2) for optogenetic activation and Archaerhodopsin (ArchT) for neuronal inhibition, as well as control vectors like tdTomato. AAV vectors were obtained from Addgene ([www.addgene.org](www.addgene.org)): ChR2-eYFP (AAV5-Efla-DIOhChR2(E123T/T159C)-EYFP, #35509) at a titer of 1 x 1013 vector genome/ml (vg/ml); ArchT-tdTomato (AAV5/FLEX-ArchT-tdTomato, #28305) at 7.0 x 1012 vg/ml; eYFP-vector (AAV5-Efla-DIO EYFP, #27056) at 1.0 x 1013 vg/ml; and AAV5 pCAG-FLEX-tdTomato-WPRE (#51503) at 3.8 x 1013 vg/ml. Viruses were aliquoted and stored at -80°C until use.

## Stereotaxic surgery, viral infection, and fiber implantation

Mice were anesthetized with 2% isoflurane (flow rate 0.5 L/min) and prior to surgery, mice received a subcutaneous analgesic (lidocaine hydrochloride; 5 mg/kg) on the scalp and topical ophthalmic ointment for eye protection. The scalp was shaved and disinfected with aseptic cotton swabs. Mice were then placed in a stereotaxic frame, and a midline sagittal scalp incision (~1 cm) was made to expose the skull, which was leveled in the bregma-lambda plane. Two stainless steel holding screws were implanted bilaterally onto the skull. A two-step procedure involving viral infection and fiber implantation was performed to achieve targeted gene expression and subsequent optical stimulation within the NAcSh. A microinjection needle (30-G) connected to a 10-µl Hamilton syringe was used for bilateral injections of AAV targeting the NAcSh at the following coordinates from

bregma (mm): AP +1.4, ML ± 0.65, DV -4.4. The injection volume was 250 nL at a rate of

30 nL/min, and the injector was left in position for an additional 5-min period to allow for

virus diffusion. Although the intended region was the NAcSh, some virus spread occurred

in the core region. Therefore, we refer to this manipulation as targeting the anterior medial

NAc. After AAV injection, a zirconia ferrule with a multimode optical fiber (200 µm,

Thorlabs) was implanted into the NAcSh at the following coordinates from bregma (mm):

AP +1.4, ML ± 1.35, DV -3.9, with a 10° angle (Stoelting stereotaxic instrument Model

Number 51730). The antibiotic enrofloxacin (5 mg/kg, i.p.) was administered once daily

for 3 days post-surgery. Mice were allowed to recover for 3 weeks to obtain stable

expression of ChR2, ArchT, EYFP, or tdTomato. In the D2$^{tdTomato}$ control group,

stimulation parameters were pseudo-randomly selected to match the protocols used for

the D2$^{ChR2}$ and D2$^{ArchT}$ groups, ensuring that control mice were exposed to similar

conditions without active modulation of D2 neuron activity.

## Optogenetic parameters

Mice expressing ChR2, ArchT opsins, or tdTomato were stimulated with a diode-pumped

solid-state laser system with a light output intensity of 10 mW at the tip of the optical fiber

patch cord, as measured with an optical power meter (PM20A, Thorlabs). For the D2$^{ChR2}$

group, the laser output wavelength was 473 nm (OEM Laser) with a pulse duration of 10

ms delivered at 20 Hz (unless otherwise indicated for each behavioral paradigm). For the

D2$^{ArchT}$ group, the laser output wavelength was 532 nm (Laserglow Technologies), and

continuous light pulses were delivered (unless otherwise indicated). The tdTomato-

expressing mice received random stimulation parameters from either the ChR2 or ArchT

protocols.

## Gustatory stimuli

Liquid solutions: Reagent-grade sucrose (Sigma-Aldrich) was diluted in distilled water to

yield 3%, 10%, and 18% w/v sucrose solutions. Solutions were prepared fresh every other

day, stored overnight at 4°C, and brought to room temperature (~23°C) before use. Solid

diets: For solid food consumption experiments, two types of food were provided: a

standard chow diet (PicoLab Rodent Diet 20) and a HFD with 45% kcal fat (Research

Diets).

## Behavioral tasks

All behavioral tasks were automated using an operant conditioning controller (Med

Associates Inc., VT, USA). Lick events were recorded with a contact lickometer (Med

Associates Inc., VT, USA). Mice were water-restricted for 23 h before the session and

given access to water for 1 h afterward for tasks involving licking behavior. For food

consumption tasks, mice were allowed to eat only a standard chow diet corresponding to

10% of their body weight (2-3 g) between sessions. For tasks requiring training, mice

were habituated to the patch cord during training sessions by connecting them to the

patch cord without laser stimulation.

## Two-bottle free-licking closed-loop assay

Mice were habituated to the behavioral box (43 x 23 cm) for two days before training. The training session aimed to familiarize mice with the two-bottle setup and establish similar licking behavior from both bottles. The box was equipped with two water bottles, with gravity-fed sippers, and connected lickometers on each side of the frontal wall. The entire test lasted six days, with one 20-min session per day. The positions of the water and sucrose bottles were randomized and counterbalanced between sessions across subjects. The procedures implemented each day were: a) Baseline (Days 1-2): Mice freely licked from both bottles without optogenetic stimulation to establish baseline sucrose preference. b) Laser Sucrose (Days 3-4): Licking the sucrose bottle triggered optogenetic stimulation. c) Laser Water (Days 5-6): Licking the water bottle triggered optogenetic stimulation. Optogenetic stimulation was delivered in a 4-s-on and 2-s-off cycle as long as the mice continued licking. The sucrose preference score was calculated as the ratio of licks received by the sucrose bottle divided by the total licks from both bottles for each day.

## Sucrose freely licking with optogenetic frequency scan

On the day before the experiment, mice were habituated to a behavioral box (22 x 18 cm, Med Associates). The task comprised five consecutive daily 20 min sessions. Each trial included three epochs: Before (laser off): The mouse had to complete three licks to initiate a new trial and enter the laser epoch. Laser epoch: On the fourth lick, the laser was activated for 1 s at different stimulation frequencies, presented in pseudo-random order: no stimulation (control) or 10 ms pulse width at 5, 8, 15, or 20 Hz. This ensured that the

animal could not predict the stimulation frequency. Time out (laser off): During this 1-s

epoch following laser stimulation, the laser could not be reactivated. The next trial began

after the mouse completed three additional licks following the time-out epoch. Each lick

was rewarded with a drop of sucrose solution (10% w/v, ~2 µL) from a licking spout

throughout the task, and the frequency-dependent effects of optogenetic stimulation on

licking behavior were analyzed.

## Brief access taste test with different optogenetic stimulation frequencies

Building upon earlier investigations, this variation of the brief access task examined the

effect of different stimulation frequencies (during the reward window) on motivated licking

behavior. The same behavioral box (22 x 18 cm, Med Associates) was used. The task

consisted of five consecutive daily 20 min sessions, each involving trials initiated by a lick

on the sipper. Following the lick, a 4-s reward epoch began, signaled by an auditory cue.

Every lick triggered the delivery of 10% w/v sucrose solution during this reward window.

Simultaneously with sucrose reward delivery, a laser was activated at different

frequencies in a pseudorandom order: no stimulation (control), 10 ms pulse width lick-

paired, 8 Hz, or 20 Hz. After the reward window ended, subsequent licks were not

rewarded. To initiate a new trial, mice had to stop licking for a randomly chosen inter-trial

interval (ITI) between 1 and 3 s.

## Brief access test with varying delays of optogenetic stimulation

This variation of the brief access task assessed the effect of the timing of lick-paired

optogenetic stimulation on behavior. The overall structure was like the previous brief

access task, with the key difference being the timing of the laser stimulation.

Simultaneously with sucrose reward delivery, a laser was activated at different delay

times in a pseudo-random order: no stimulation (control) or a 10 ms pulse delivered at

specific delays after the lick (lick-evoked), with no delay (0 ms), a 50 ms delay, or a 100

ms delay.

**Real-time place preference (RTPP)**

Mice were placed in a rectangular acrylic arena (42 x 20 x 20 cm) divided into two halves

differentiated by context (visual cues: one side with black/white stripes, the other with

black/white circles; Fig 5a). Mice freely explored both sides of the chamber. The

experiment included two 10-min sessions separated by approximately 5 h on the same

day. During each session, one side of the chamber was paired with optogenetic

stimulation. Each Time mice crossed to the paired chamber; they received 4 s of

optogenetic stimulation followed by 2 s without stimulation. This cycle (4 s on, 2 s off)

could repeat as long as the mouse remained on the paired side. The stimulation

parameters were: D2$^{tdTomato}$ and D2$^{ChR2}$, 10 ms pulse at 20 Hz and 8 Hz stimulation (each

frequency used on a separate day); D2$^{ArchT}$: Continuous pulse stimulation. The preference

index for the conditioned and unconditioned sides was calculated by dividing the total

time spent on each side by the total exploration time of the session. The chamber side

paired with optogenetic stimulation was counterbalanced across subjects.

**Open-loop palatable (HFD) consumption test**

This task investigated how mice respond to HFD consumption without context

conditioning elements. Mice were placed in a white arena with the same dimensions as

in the RTPP task, with an HFD pellet on each side (pellet weight measured before and

after). Mice freely explored for a single 20-min session, receiving stimulation: 4 s on, 2 s

off (D2$^{tdTomato}$ and D2$^{ChR2}$, 20 Hz, 10 ms pulse; D2$^{ArchT}$, Continuous pulse). Using the same

stimulation parameters as RTPP isolated the effects of optogenetic stimulation on HFD

consumption from context conditioning. Food consumption was calculated by dividing the

pellet weight difference by session time.

## RTPP with palatable (HFD) consumption

Similar to the RTPP task, mice were placed in a rectangular acrylic arena (same

dimensions) with an HFD pellet placed on each side. The pellet weight was determined

before and after each session. Mice freely explored both sides. The experiment consisted

of five consecutive daily sessions: a) Pre-Test (Day 1, 10 min): Mice explored with HFD

pellets but no optogenetic stimulation; b) Acquisition (Days 2-4, 20 min): Based on the

pre-test exploration time, mice were assigned a conditioned side (more Time for D2$^{tdTomato}$

and D2$^{ArchT}$ or less preferred for D2$^{ChR2}$). When crossing to and remaining on the

conditioned side, they received optogenetic stimulation: D2$^{tdTomato}$ and D2$^{ChR2}$: 4 s on (20

Hz, 10 ms pulse), 2 s off; D2$^{ArchT}$: Continuous pulse; c) Test (Day 5, 10 min): Mice

explored with HFD pellets without stimulation. The preference index was calculated by

dividing the time spent on each side by the total exploration time. Food consumption was

calculated by dividing the pellet weight difference by session time.

## Negative reinforcement test

Mice were tested for negative reinforcement in a white open circular arena (50 cm diameter) where they could freely explore during daily 30-min sessions for 3 consecutive days. Behavior was monitored from above using a video camera, and the mouse's position was analyzed in real-time using custom MATLAB code. The code calculated the centroid of the mouse and sent this information to the operant conditioning controller (Med Associates Inc., VT, USA). The arena was virtually divided into two zones: an inner circle (10 cm diameter) designated as the 'off zone' and an outer annulus surrounding the off zone defined as the 'on zone.' The on zone was paired with optogenetic stimulation, and when a mouse entered the on zone, it received 4 s of optogenetic stimulation followed by 2 s without stimulation. This cycle could be repeated if the mouse remained in the on zone. The stimulation parameters were $D2^{tdTomato}$ and $D2^{ChR2}$, 10 ms pulse at 20 Hz; $D2^{ArchT}$, optogenetic stimulation was delivered as continuous pulses. To stop optogenetic stimulation, the mouse had to enter the off zone and remain there. The analysis software determined zone entry and exit based on the position of the calculated centroid provided by the MATLAB code. The percentage of time spent in the stimulated and unstimulated zones was calculated by dividing the total time spent in each zone by the total exploration time of the session.

# Histology

After the experimental sessions, mice received a pentobarbital sodium overdose and were transcardially perfused with a phosphate-buffered saline solution followed by 4%

paraformaldehyde (PFA). Brains were removed, stored for 1 day in 4% PFA, and then

transferred to 30% sucrose in phosphate-buffered saline solution. Brains were sectioned

into 40-µm coronal slices and mounted in a medium (Dako) for confocal microscopy to

confirm viral expression and fiber placement. Mice with incorrect expression or fiber

placement were excluded from the analysis. Images were taken with a confocal

microscope, and contrast was improved with Adobe Photoshop CS5.1 software.

## Quantification and statistical analysis
711

Data were analyzed in MATLAB R2024a (The MathWorks Inc.) and GraphPad Prism 9.

Data were expressed as means ± SEM. Statistical analyses were performed by first

assessing normality distribution, homoscedasticity, and sample size to decide the use of

parametric or non-parametric tests. As most of the data met the assumptions for

parametric tests, one-way ANOVA was used, followed by a Tukey-Kramer post hoc test

with an $\alpha$ level of 0.05. Sample sizes were decided based on previous studies using

similar behavioral paradigms and are reported in figure legends. Potential confounders,

such as variations in individual animal behavior and environmental factors, were

controlled by randomizing the order of experimental conditions and balancing the groups

by sex and age.

## Data Availability Statement
722

723 All relevant data are within the paper and its Supporting Information files. Additional data

724 is available on request from the corresponding author.

# Acknowledgments

All behavioral task schemes were created using BioRender. We thank Elvi Gil Lievana for her help with histology, Fabiola Olvera Hernandez for her valuable assistance with animal care, and Jorge Luis-Islas for his collaboration in the early stages of this project. We also appreciate the comments from Luis Téllez on a previous version of this manuscript.

# Funding Statement

This research was partially supported by CONAHCyT grant number CF-2023-G-518 under the Ciencia de Frontera initiative to RG.

# Competing Interests

The authors have declared that no competing interests exist.

# References

1.    Taha SA, Fields HL. Encoding of Palatability and Appetitive Behaviors by Distinct Neuronal Populations in the Nucleus Accumbens. J Neurosci. 2005;25: 1193–1202. doi:10.1523/JNEUROSCI.3975-04.2005

2.    Villavicencio M, Moreno MG, Simon SA, Gutierrez R. Encoding of Sucrose's Palatability in the Nucleus Accumbens Shell and Its Modulation by Exteroceptive Auditory Cues. Front Neurosci. 2018;12. doi:10.3389/fnins.2018.00265

3.    Yang AK, Mendoza JA, Lafferty CK, Lacroix F, Britt JP. Hippocampal Input to the Nucleus Accumbens Shell Enhances Food Palatability. Biol Psychiatry. 2019;87: 597–608. doi:10.1016/j.biopsych.2019.09.007

4.    Yao Y, Gao G, Liu K, Shi X, Cheng M, Xiong Y, et al. Projections from D2 Neurons in Different Subregions of Nucleus Accumbens Shell to Ventral Pallidum Play Distinct Roles in Reward and Aversion. Neurosci Bull. 2021;37: 623–640. doi:10.1007/s12264-

021-00632-9

5.    Soares-Cunha C, Vasconcelos NAP de, Coimbra B, Domingues AV, Silva JM,
Loureiro-Campos E, et al. Nucleus accumbens medium spiny neurons subtypes signal
both reward and aversion. Mol Psychiatry. 2019; 1–15. doi:10.1038/s41380-019-0484-3

6.    Berthoud H-R. Metabolic and hedonic drives in the neural control of appetite: who
is the boss? Curr Opin Neurobiol. 2011;21: 888–896. doi:10.1016/j.conb.2011.09.004

7.    Berridge KC, Kringelbach ML. Pleasure systems in the brain. Neuron. 2015;86.
doi:10.1016/j.neuron.2015.02.018

8.    Kelley A, Baldo B, Pratt W, Will M. Corticostriatal-hypothalamic circuitry and food
motivation: Integration of energy, action and reward. Physiol Behav. 2005;86: 773–795.
doi:10.1016/j.physbeh.2005.08.066

9.    O'Connor EC, Kremer Y, Lefort S, Harada M, Pascoli V, Rohner C, et al. Accumbal
D1R Neurons Projecting to Lateral Hypothalamus Authorize Feeding. Neuron. 2015;88:
553–564. doi:10.1016/j.neuron.2015.09.038

10.    Gendelis S, Inbar D, Kupchik YM. The role of the nucleus accumbens and ventral
pallidum in feeding and obesity. Prog Neuropsychopharmacol Biol Psychiatry. 2021;111:
110394. doi:10.1016/j.pnpbp.2021.110394

11.    Sandoval-Rodríguez R, Parra-Reyes JA, Han W, Rueda-Orozco PE, Perez IO,
Araujo IE de, et al. D1 and D2 neurons in the nucleus accumbens enable positive and
negative control over sugar intake in mice. Cell Rep. 2023;42.
doi:10.1016/j.celrep.2023.112190

12.    Olds J. Pleasure Centers in the Brain. Sci Am. 1956;195: 105–117.

13.    Britt JP, Benaliouad F, McDevitt RA, Stuber GD, Wise RA, Bonci A. Synaptic and
Behavioral Profile of Multiple Glutamatergic Inputs to the Nucleus Accumbens. Neuron.
2012;76: 790–803. doi:10.1016/j.neuron.2012.09.040

14.    Prado L, Luis-Islas J, Sandoval OI, Puron L, Gil MM, Luna A, et al. Activation of
Glutamatergic Fibers in the Anterior NAc Shell Modulates Reward Activity in the aNAcSh,
the Lateral Hypothalamus, and Medial Prefrontal Cortex and Transiently Stops Feeding.
J Neurosci. 2016;36: 12511–12529. doi:10.1523/JNEUROSCI.1605-16.2016

15.    Walle R, Petitbon A, Fois GR, Varin C, Montalban E, Hardt L, et al. Nucleus
accumbens D1- and D2-expressing neurons control the balance between feeding and
activity-mediated energy expenditure. Nat Commun. 2024;15: 2543. doi:10.1038/s41467-
024-46874-9

16.    Lobo MK, Covington HE, Chaudhury D, Friedman AK, Sun H, Damez-Werno D, et
al. Cell Type–Specific Loss of BDNF Signaling Mimics Optogenetic Control of Cocaine
Reward. Science. 2010;330: 385–390. doi:10.1126/science.1188472

17.    Cole SL, Robinson MJF, Berridge KC. Optogenetic self-stimulation in the nucleus
accumbens: D1 reward versus D2 ambivalence. McCutcheon JE, editor. PLOS ONE.
2018;13: e0207694. doi:10.1371/journal.pone.0207694

18.    Durieux PF, Bearzatto B, Guiducci S, Buch T, Waisman A, Zoli M, et al. D2R
striatopallidal neurons inhibit both locomotor and drug reward processes. Nat Neurosci.
2009;12: 393–395. doi:10.1038/nn.2286

19.    Soares-Cunha C, Vasconcelos NAP de, Coimbra B, Domingues AV, Silva JM,
Loureiro-Campos E, et al. Nucleus accumbens medium spiny neurons subtypes signal
both reward and aversion. Mol Psychiatry. 2019; 1–15. doi:10.1038/s41380-019-0484-3

20.    Baik J-H. Dopamine Signaling in reward-related behaviors. Front Neural Circuits.

2013;7. doi:10.3389/fncir.2013.00152

21.   Oleson EB, Gentry RN, Chioma VC, Cheer JF. Subsecond Dopamine Release in
the Nucleus Accumbens Predicts Conditioned Punishment and Its Successful Avoidance.
J Neurosci. 2012;32: 14804–14808. doi:10.1523/JNEUROSCI.3087-12.2012

22.   Arroyo B, Hernandez-Lemus E, Gutierrez R. The flow of reward information
through neuronal ensembles in the accumbens. bioRxiv; 2024. p. 2024.02.15.580379.
doi:10.1101/2024.02.15.580379

23.   Coss A, Suaste E, Gutierrez R. Lateral NAc Shell D1 and D2 Neuronal Ensembles
Concurrently Predict Licking Behavior and Categorize Sucrose Concentrations in a
Context-dependent   Manner.   Neuroscience.   2022;493:   81–98.
doi:10.1016/j.neuroscience.2022.04.022

24.   Gutierrez R, Simon SA, Nicolelis MAL. Licking-Induced Synchrony in the Taste–
Reward Circuit Improves Cue Discrimination during Learning. J Neurosci. 2010;30: 287–
303. doi:10.1523/JNEUROSCI.0855-09.2010

25.   Francis TC, Yano H, Demarest TG, Shen H, Bonci A. High-Frequency Activation
of Nucleus Accumbens D1-MSNs Drives Excitatory Potentiation on D2-MSNs. Neuron.
2019;103: 432-444.e3. doi:10.1016/j.neuron.2019.05.031

26.   Soares-Cunha C, Coimbra B, Domingues AV, Vasconcelos N, Sousa N, Rodrigues
AJ. Nucleus Accumbens Microcircuit Underlying D2-MSN-Driven Increase in Motivation.
eNeuro. 2018;5. doi:10.1523/ENEURO.0386-18.2018

27.   Reed SJ, Lafferty CK, Mendoza JA, Yang AK, Davidson TJ, Grosenick L, et al.
Coordinated Reductions in Excitatory Input to the Nucleus Accumbens Underlie Food
Consumption. Neuron. 2018;99: 1260-1273.e4. doi:10.1016/j.neuron.2018.07.051

28.   Mogenson G, Jones D, Yim C. From motivation to action: Functional interface
between the limbic system and the motor system. Prog Neurobiol. 1980;14: 69–97.
doi:10.1016/0301-0082(80)90018-0

29.   Oginsky MF, Goforth PB, Nobile CW, Lopez-Santiago LF, Ferrario CR. Eating
'Junk-Food' Produces Rapid and Long-Lasting Increases in NAc CP-AMPA Receptors:
Implications   for   Enhanced   Cue-Induced   Motivation   and   Food   Addiction.
Neuropsychopharmacology. 2016;41: 2977–2986. doi:10.1038/npp.2016.111

30.   Yang H, Jong JW de, Tak Y, Peck J, Bateup HS, Lammel S. Nucleus Accumbens
Subnuclei Regulate Motivated Behavior via Direct Inhibition and Disinhibition of VTA
Dopamine   Subpopulations.   Neuron.   2018;97:   434-449.e4.
doi:10.1016/j.neuron.2017.12.022

31.   Luis-Islas J, Luna M, Floran B, Gutierrez R. Optoception: Perception of
Optogenetic Brain Perturbations. eNeuro. 2022;9. doi:10.1523/ENEURO.0216-22.2022

32.   Swinford-Jackson SE, Huffman PJ, Knouse MC, Thomas AS, Rich MT, Mankame
S, et al. High frequency DBS-like optogenetic stimulation of nucleus accumbens
dopamine D2 receptor-containing neurons attenuates cocaine reinstatement in male rats.
Neuropsychopharmacology. 2023;48: 459–467. doi:10.1038/s41386-022-01495-y

33.   Johnson PM, Kenny PJ. Dopamine D2 receptors in addiction-like reward
dysfunction and compulsive eating in obese rats. Nat Neurosci. 2010;13: 635–641.
doi:10.1038/nn.2519

# Figures

Figure 1. Optogenetic activation of NAc D2 neurons at 20 Hz reduces sucrose preference, disrupts licking microstructure, and increases self-stimulation trials.

(**a**) Coronal brain sections depicting Cre recombinase expression in D2-expressing neurons. Below, the viral constructs used: pAAV-Ef1a-DIO-hChR2-EYFP (activation), AAV-FLEX-ArchT-tdTomato (inhibition), and AAV-FLEX-tdTomato (control). (**b**) Two-bottle test protocol: baseline (No-laser, Days 1-2), laser triggered by licking the sucrose

bottle (Days 3-4), and laser triggered by the water bottle (Days 5-6). (**c**) Sucrose

preference (%) across conditions in each group. (**d**) Total licks during baseline, laser

sucrose, and laser water conditions. (**e**) Total trials (number of trials with self-activated

laser). Dashed lines represent the baseline, calculated from the licks given per each

solution in the off condition and the trial time (4 seconds on, 2 seconds off), which reflects

the expected times the mice would start a new trial if laser stimulation were present. (**f-h**)

Heatmaps of licking bouts. Water-licking bouts are shown in blue, and sucrose-licking

bouts in red. (**f**) Baseline condition. (**g**) 4 s laser triggered with sucrose bottle. (**h**) Laser

triggered with a water bottle. Licking rate (licks per second) is shown on the right.

* significant differences with the control group (D2$^{tdTomato}$) within the same condition; #

significant differences compared to baseline within the same group. @ significant

difference for total licks (water + sucrose) against control group within the same condition.

All significant differences had $p < 0.05$. Data are mean ± SEM. The statistical analyses

were conducted with one-way ANOVA followed by Tukey-Kramer post hoc tests.

D2$^{tdTomato}$ n=5, D2$^{ChR2}$ n=11, D2$^{ArchT}$ n=9. Licking microstructure details are provided in

**Table 1**.

**Figure 2. Optogenetic activation of NAc D2-expressing neurons disrupts licking behavior in a frequency-dependent manner.**

(**a**) Schematic representation of the freely licking test with 10% sucrose and an optogenetic scanning frequency. The freely licking task is composed of three epochs: Before (four licks), Laser (1 s), and Time out (1 s). Every four licks given after the time-out epoch starts a new Laser epoch. (**b-d**) Representative raster plots showing individual licks (red ticks) and laser pulses (blue for activation, green for inhibition) during a single session in D2$^{tdTomato}$ (control), D2$^{ChR2}$ (activation), and D2$^{ArchT}$ (inhibition) groups. The figures below correspond to the Peri-Stimulus Time Histogram (PSTH) of the single session. (**e-g**) Population PSTHs display mean lick rates for all mice aligned to laser stimulation onset (Time = 0 s). Note that the licking structure is disrupted in a frequency-dependent manner. (**h-j**) Cumulative lick distributions during laser epochs across the session confirmed the disruption of licking for the D2$^{ChR2}$ group. (**k**) Total licks during the 1 s laser epoch reveal frequency-dependent disruption in the D2$^{ChR2}$ group. (**l**) Licks during the 1 s timeout phase (post-laser) show recovery. (**m**) The time to start licking after the first laser pulse (Time = 0 s) shows delayed re-initiation in the D2$^{ChR2}$ group, increasing with frequency. * significant differences from the control group (D2$^{tdTomato}$) within the same condition; # significant differences compared to No-laser, @ significant differences compared to 5 Hz frequency within the same group. All significant differences had $p < 0.05$. Data are mean ± SEM. The statistical analyses were conducted with one-way ANOVA followed by Tukey-Kramer post hoc tests. D2$^{tdTomato}$ n=3, D2$^{ChR2}$ n=8, D2$^{ArchT}$ n=7. Licking microstructure details are provided in **Table 2**.

**Figure 3. Frequency-dependent disruption of licking behavior in a sucrose brief access test demonstrates the advantage of self-paced (lick-paired) activation over imposed optogenetic stimulation and exhibits enhanced task engagement through D2 neuron activation.**

(**a**) Schematic representation of the sucrose brief access (4 s) taste test with optogenetic stimulation, including No-laser (gray) and 10 ms pulses at 0 ms delay (blue), 8 Hz (yellow),

and 20 Hz (red) frequencies for 4  s. Each trial included the delivery of sucrose in each

lick for 4 s (red ticks); after that, additional licks were empty (gray ticks). To start a new

trial, mice must stop licking for a 1-3 s inter-trial interval (ITI). The experiment comprises

5 sessions, each lasting 20 min. (**b-d**) PSTH displaying mean sucrose lick rates during

the 6 s of trial for all D2$^{tdTomato}$ (**b**), D2$^{ChR2}$ (**c**), and D2$^{ArchT}$ mice (**d**). The shaded area

represents the 4 s laser stimulation period. (**e-g**) Cumulative licks over the 20-minute

session for all stimulation patterns within the same group are shown on the left y-axis,

with total cumulative licks (in purple) on the right y-axis. (**h**) Total licks per session for

each group under different stimulation conditions. (**i**) Heatmaps plotting licking bouts

across stimulation patterns for all groups, with the x-axis representing time in s (0-4 s)

from lick bout onset and the y-axis showing individual bouts (the number in white indicates

lick bout duration in s). Dark gray edges (bar line at left) represent No-laser bouts, blue

edges lick-paired laser stimulation, yellow edges 8 Hz imposed stimulation, and red edges

20 Hz stimulation. Light gray intensity indicates lick rate. (**j**) Total trials given by each

group under different stimulation conditions. The activation of D2 neurons in the NAc led

to significantly more trials started in the D2$^{ChR2}$ group than in the D2$^{tdTomato}$ group. Colored

symbols: * significant difference compared to D2$^{tdTomato}$ at the same frequency, #

significant difference compared to the No-laser condition within the same group, and @

significant difference between the lick-paired condition and the 8 Hz or 20 Hz conditions

within the same group.  The symbol ! indicates a significant difference between the 8 Hz

and 20 Hz conditions within the same group. All significant differences had $p < 0.05$. Data

are mean ± SEM. The statistical analyses were conducted with one-way ANOVA followed

by Tukey-Kramer post hoc tests. D2$^{tdTomato}$ n=6, D2$^{ChR2}$ n=5, D2$^{ArchT}$ n=5. Further analysis

of the licking microstructure is presented in **Table 3**.

**Figure 4. Lick-paired optogenetic stimulation of D2 NAc neuron fragments licking microstructure but paradoxically promotes sucrose intake through its rewarding effects.**

(**a**) Schematic representation of the brief access test with optogenetic stimulation. Mice were subjected to varying delays (0, 50, and 100 ms) of 10 ms pulses of laser stimulation entrained to licking during a 4 s reward period alongside a No-laser condition. (**b-d**) PSTHs display the mean sucrose lick rates during the 4 s laser stimulation period for D2$^{tdTomato}$ (**b**), D2$^{ChR2}$ (**c**), and D2$^{ArchT}$ (**d**). (**e-g**)Cumulative licks over the 20 min session are shown for all conditions within the same group on the left y-axis, while total cumulative licks (in purple) are shown on the right y-axis. (**h**) Inter-lick interval (ms) for each group under different stimulation conditions. In the D2$^{ChR2}$ group, inter-lick intervals across all conditions were significantly longer compared to the D2$^{tdTomato}$ group in the same condition. In the D2$^{ArchT}$ group, the 0 ms delay condition showed a significantly shorter inter-lick interval than the D2$^{tdTomato}$ group. (**i**) Number of licking bouts for each group across conditions. The D2$^{ChR2}$ group exhibited significantly more bouts than the D2$^{tdTomato}$ group across all frequencies. (**j**) Bout size (licks) for each group. The D2$^{ChR2}$ group had significantly smaller bout sizes than the D2$^{tdTomato}$ group in the No-laser, 0 ms delay, and 50 ms delay conditions. (**k**) Total trials initiated by each group under different stimulation conditions. The activation of D2 neurons in the NAc led to significantly more trials initiated in the D2$^{ChR2}$ group than in the D2$^{tdTomato}$ control group. Colored symbols: * significant difference compared to D2$^{tdTomato}$ (control) at the same frequency, significant difference compared to the No-laser condition within the same group. All significant differences had $p < 0.05$. Data are mean ± SEM. The statistical analyses were conducted with one-way

943 ANOVA followed by Tukey-Kramer post hoc tests. D2$^{tdTomato}$ n=6, D2$^{ChR2}$ n=5, and D2$^{ArchT}$

944 n=5. Further analysis of the licking microstructure is presented in **Table 4**.

**Figure 5. Optogenetic activation of NAc D2 neurons at 20 Hz (but not 8 Hz) is rewarding and consistently drives place preference.**

(**a**) Schematic representation of the Real-Time Place Preference (RTPP) task: Mice explored a rectangular acrylic chamber divided into two visually distinct halves (one with black/white stripes and the other with black/white circles). Upon entering the paired side, mice received optogenetic stimulation for 4  s(10 ms pulse width at 20 Hz or 8 Hz for D2$^{tdTomato}$ and D2$^{ChR2}$), followed by 2  s off, as long as mice remained on that side. Mice performed two 10 min sessions, separated by approximately 5 h. The side paired with stimulation was counterbalanced across subjects. (**b-e**) Heatmaps of mouse locations during the task under different conditions: (**b**) D2$^{tdTomato}$ with 20 Hz stimulation, (**c**) D2$^{ChR2}$ with 20 Hz stimulation, (**d**) D2$^{tdTomato}$ with 8 Hz stimulation, and (**e**) D2$^{ChR2}$ with 8 Hz stimulation (**f**) Percentage of time spent on the stimulated side during Session 1 and Session 2. Mice expressing D2$^{ChR2}$ at 20 Hz spent significantly more time on the stimulated side compared to D2$^{tdTomato}$ controls and the 8 Hz condition. No significant preference was observed for 8 Hz groups. (**g**) Number of side crossings between the two halves of the chamber. No significant differences were found across conditions. (**h**) Velocity of the mice across conditions. The D2$^{ChR2}$ 20 Hz group had significantly higher velocity on the non-stimulated side compared to the stimulated side and all other groups and conditions on the non-stimulated side. (**i**) Total distance traveled during stimulation sessions. The D2$^{ChR2}$ 20 Hz group traveled significantly more on the stimulated side than D2$^{tdTomato}$ controls and other conditions. * significant difference compared to other groups in the same session or side, # significant difference in the other side within the same group. All significant differences had $p < 0.05$. Data are mean ± SEM. The statistical

analyses were conducted with one-way ANOVA followed by Tukey-Kramer post hoc

tests. D2$^{tdTomato}$ 8 Hz n=5, D2$^{ChR2}$ 8 Hz n=4, D2$^{tdTomato}$ 20 Hz n=9 and D2$^{ChR2}$ 20 Hz n=7.

**Figure 6. Context-dependent effects of silencing NAc D2 neurons: inhibition only**

**alters HFD consumption based on environmental context**

(**a**) Left panel. Schematic representation of the RTPP task: Mice explored a rectangular

acrylic chamber divided into two visually distinct sides. Upon entering the laser paired

side, mice received optogenetic stimulation for 4 s (10 ms pulses at 20 Hz for D2$^{tdTomato}$

or continuous stimulation for D2$^{ArchT}$), followed by 2 s off. Mice experienced two 10 min

sessions, separated by approximately 5 h. The side paired with stimulation was

counterbalanced across subjects. Middle panels Heatmaps of mouse locations during the

task under different conditions: (middle-top) D2$^{ArchT}$ with continuous stimulation, (middle-

bottom) D2$^{tdTomato}$ with 20 Hz stimulation. Right panel percentage of time spent on the

stimulated side during session 1 and session 2. No significant preference was observed.

D2$^{tdTomato}$ 20 Hz n=9 (same data from **Fig 5**) and D2$^{ArchT}$ n=12. (**b**) Left panel. Schematic

of the open-loop paradigm used to assess the effects of optogenetic manipulation of D2-

NAc neurons on HFD intake. optogenetic stimulation was delivered in 4 s on / 2 s off

intervals, with continuous stimulation applied to the D2$^{ArchT}$ group, while a 20 Hz

frequency was used for the D2$^{ChR2}$ and D2$^{tdTomato}$ groups. Right panel. Total HFD intake

(mg/min) for D2$^{tdTomato}$ (control), D2$^{ChR2}$ (activation), and D2$^{ArchT}$ (inhibition) groups during

the 20 min session. D2$^{tdTomato}$ n=5, D2$^{ChR2}$ n=6, D2$^{ArchT}$ n=4. (**c**) Schematic representation

of the RTPP paradigm associated with palatable food intake. Mice were conditioned to

associate one side of the chamber with optogenetic stimulation (4 s on, 2 s off). D2$^{ChR2}$

and D2$^{tdTomato}$ groups received 10 ms pulses of 20 Hz stimulation, while the D2$^{ArchT}$ group

received continuous pulse stimulation. (**d**) Time course of the percentage of time spent

on the stimulation-paired side during pre-conditioning (PreTest), conditioning (Acq1,

Acq2, Acq3), and post-conditioning test (Test). Note that the D2$^{ChR2}$ group significantly preferred the stimulated side during conditioning, consistently promoting place preference in the test without stimulation. In contrast, silencing D2 neurons induces a tendency to avoid the laser-paired side (**e**) Total HFD intake across conditioning days for all groups. There is a tendency for reduced intake during the activation of NAc D2 neurons (D2$^{ChR2}$) and an opposite tendency for increased intake during inhibition (D2$^{ArchT}$). (**f-h**) Comparison of HFD intake on the stimulated versus non-stimulated sides for D2$^{tdTomato}$ (**f**), D2$^{ChR2}$ (**g**), and D2$^{ArchT}$ (**h**) groups. The D2$^{ArchT}$ group showed a significant increase in HFD consumption on the non-stimulated side during the last two conditioning days compared to pre-conditioning. Gradient rectangles depict the condition or side paired with optogenetic stimulation. D2$^{tdTomato}$ n=7, D2$^{ChR2}$ n=7, D2$^{ArchT}$ n=5: * significant difference compared to D2$^{tdTomato}$ in the same session, # significant difference compared to the PreTest session within the same group, and @ significant difference between the stimulated and non-simulated side within the same group. All significant differences had $p < 0.05$. Data are mean ± SEM. The statistical analyses were conducted with one-way ANOVA followed by Tukey-Kramer post hoc tests.

1011

**Figure 7. Silencing of NAc D2 neuron activity is hedonically negative and drives**

**negative reinforcement**

(**a**) Schematic illustration of the negative reinforcement paradigm. Mice were placed in a

circular arena with distinct 'on' (optogenetic stimulation; 4 s on, 2 s off) and 'off' zones

that turn off the stimulation. D2$^{ChR2}$ and D2$^{tdTomato}$ groups received pulsed stimulation in

the 'on' zone, while D2ArchT received continuous stimulation. (**b**) Percentage of time spent in the 'on' and 'off' zones during the test for D2$^{\text{tdTomato}}$ (gray), D2$^{\text{ChR2}}$ (blue), and D2$^{\text{ArchT}}$ (green) groups. (**c**) Time course of the percentage of time spent in the initially 'off' zone across the 3-day sessions. Mice expressing D2$^{\text{ArchT}}$ showed increased time spent in the 'off' zone on day 3, indicative of negative reinforcement behavior. (**d-f**) Representative heatmaps showing the spatial distribution of time spent by mice in the D2$^{\text{tdTomato}}$ (**d**), D2$^{\text{ChR2}}$ (**e**), and D2$^{\text{ArchT}}$ (**f**) groups during the post-conditioning test. * significant difference compared to other groups. All significant differences had $p < 0.05$. Data are mean ± SEM. The statistical analyses were conducted with one-way ANOVA followed by Tukey-Kramer post hoc tests. D2$^{\text{tdTomato}}$ n=6, D2$^{\text{ChR2}}$ n=4, D2$^{\text{ArchT}}$ n=4.

Tables

**Table 1. Analysis of licking microstructure for the two-bottle sucrose test (Fig 1).**

| | | D2^tdTomato | | D2^ChR2 | | D2^ArchT | |
|---|---|---|---|---|---|---|---|
| | | Water | Sucrose | Water | Sucrose | Water | Sucrose |
| **Condition 1** (Base line no-laser) | Preference (%) | 21.69 ± 4.35 | 78.31 ± 4.35 | 11.96 ± 1.68 | 88.04 ± 1.68 | 14.53 ± 3.49 | 85.48 ± 3.49 |
| | Inter lick interval (ms) | 135.08 ± 3.35 | 130.32 ± 2.58 | 146.00 ± 5.99 | 134.43 ± 4.41 | 133.64 ± 2.90 | 128.94 ± 1.85 |
| | Lick rate (Hz) | 10.04 ± 0.47 | 9.64 ± 0.38 | 9.75 ± 0.39 | 9.50 ± 0.13 | 10.15 ± 0.52 | 9.40 ± 0.27 |
| | Bout size (Licks) | 14.10 ± 3.53 | 23.18 ± 5.33 | 9.89 ± 1.61 | 19.41 ± 2.03 | 12.96 ± 2.46 | 28.38 ± 4.57 |
| | Bout duration (s) | 1.71 ± 0.44 | 2.88 ± 0.69 | 1.23 ± 0.21 | 2.40 ± 0.26 | 1.60 ± 0.33 | 3.55 ± 0.61 |
| | Inter bout interval (s) | 82.49 ± 11.96 | 39.96 ± 3.88 | 101.50 ± 25.40 | 22.18 ± 2.45 | 101.34 ± 24.55 | 29.76 ± 2.88 |
| | Number of bouts | 14.31 ± 2.09 | 35.54 ± 3.52 | 17.86 ± 2.54 | 70.14 ± 7.58 | 14.94 ± 2.77 | 48.82 ± 6.89 |
| | Total licks | 182.92 ± 39.79 | 714.00 ± 119.42 | 152.41 ± 24.99 | 1140.09 ± 94.84 | 154.24 ± 34.42 | 1198.82 ± 154.07 |
| | Total licks sum | 896.92 ± 132.45 | | 1292.50 ± 101.87 | | 1353.06 ± 150.57 | |
| **Condition 2** (Laser sucrose) | Preference (%) | 16.32 ± 3.70 | 83.68 ± 3.70 | #*40.43 ± 4.28 | #*59.57 ± 4.28 | 20.84 ± 4.48 | 79.16 ± 4.48 |
| | Inter lick interval (ms) | 137.22 ± 3.40 | 130.23 ± 1.83 | 136.46 ± 2.75 | #*160.19 ± 4.25 | 137.23 ± 4.54 | 128.70 ± 3.65 |
| | Lick rate (Hz) | 9.59 ± 0.63 | 9.08 ± 0.25 | 9.17 ± 0.29 | *9.86 ± 0.23 | #8.86 ± 0.36 | 9.57 ± 0.26 |
| | Bout size (Licks) | 20.14 ± 4.61 | 32.34 ± 3.65 | #22.68 ± 2.05 | # 5.66 ± 0.57 | 29.22 ± 7.40 | 25.74 ± 4.02 |
| | Bout duration (s) | 2.59 ± 0.61 | 4.09 ± 0.47 | #2.93 ± 0.29 | # 0.72 ± 0.08 | 3.65 ± 0.95 | 3.20 ± 0.53 |
| | Inter bout interval (s) | 172.61 ± 85.84 | 45.11 ± 5.54 | 52.88 ± 5.41 | #*14.78 ± 1.66 | 64.72 ± 24.94 | *28.36 ± 3.43 |
| | Number of bouts | 9.00 ± 1.68 | 32.78 ± 4.03 | *18.25 ± 1.66 | #*121.21 ± 14.65 | 13.91 ± 2.80 | *64.04 ± 10.03 |
| | Total licks | 172.61 ± 44.05 | 939.39 ± 123.44 | #*414.42 ± 45.60 | #659.25 ± 91.50 | 243.65 ± 45.51 | 1298.61 ± 166.53 |
| | Total licks sum | 1112.00 ± 122.96 | | #1073.67 ± 103.77 | | #*1542.26 ± 159.17 | |
| **Condition 3** (Laser water) | Preference (%) | 20.63 ± 5.18 | 79.37 ± 5.18 | 13.34 ± 3.34 | 86.66 ± 3.34 | 9.57 ± 1.80 | 90.43 ± 1.80 |
| | Inter lick interval (ms) | 145.94 ± 12.51 | 132.34 ± 2.61 | #166.96 ± 7.05 | 134.50 ± 1.89 | 139.97 ± 10.09 | 128.62 ± 2.22 |
| | Lick rate (Hz) | 10.50 ± 0.72 | 9.09 ± 0.35 | 9.38 ± 0.38 | #8.71 ± 0.21 | 9.93 ± 0.57 | 9.71 ± 0.28 |
| | Bout size (Licks) | 17.44 ± 4.60 | 31.50 ± 5.04 | #*5.15 ± 1.49 | #30.51 ± 3.55 | 14.14 ± 6.83 | 22.69 ± 3.98 |
| | Bout duration (s) | 2.18 ± 0.59 | 4.02 ± 0.69 | *0.65 ± 0.23 | #3.93 ± 0.44 | 1.70 ± 0.87 | 2.76 ± 0.50 |
| | Inter bout interval (s) | 168.50 ± 42.56 | 61.76 ± 5.59 | #*24.90 ± 2.91 | *28.28 ± 4.61 | 78.05 ± 12.98 | #*20.22 ± 2.93 |
| | Number of bouts | 9.42 ± 1.53 | 23.92 ± 2.80 | #*58.38 ± 14.21 | *61.69 ± 8.02 | 17.82 ± 4.70 | *78.82 ± 17.14 |
| | Total licks | 161.00 ± 41.41 | 647.25 ± 93.14 | 245.38 ± 66.20 | #*1588.12 ± 137.07 | 132.73 ± 24.25 | *1475.27 ± 223.91 |
| | Total licks sum | 808.25 ± 89.56 | | #1833.50 ± 142.29 | | #*1608.00 ± 231.36 | |

1036

1037 Bold font indicates statistically significant differences (p<0.05):

1038 *Significant difference compared to the D2^tdTomato (control group) in the same condition.

1039 #Significant difference compared to Baseline (condition 1) within the same group.

1040 **Table 2. Analysis of licking microstructure for sucrose freely licking with**

1041 **optogenetic frequency scan test (Fig 2).**

| | | D2$^{tdTomato}$ | D2$^{ChR2}$ | D2$^{ArchT}$ |
|---|---|---|---|---|
| **No-laser** | Inter lick interval (ms) | 159.75 ± 7.40 | **#168.31 ± 5.45** | **#162.87 ± 5.26** |
| | Lick rate (Hz) | 8.06 ± 0.35 | 8.28 ± 0.29 | 7.93 ± 0.16 |
| | Bout size (Licks) | 5.36 ± 0.26 | 4.71 ± 0.23 | 5.37 ± 0.17 |
| | Bout duration (s) | 0.68 ± 0.02 | 0.61 ± 0.03 | 0.69 ± 0.01 |
| | Inter bout interval (s) | 154.56 ± 21.92 | 194.52 ± 20.23 | 145.27 ± 13.77 |
| | Number of bouts | 7.50 ± 0.96 | 6.63 ± 0.88 | 9.00 ± 1.01 |
| | Total licks | 41.00 ± 6.44 | 32.81 ± 5.71 | 52.48 ± 7.32 |
| **5 Hz** | Inter lick interval (ms) | 176.21 ± 16.76 | **#202.76 ± 8.79** | **#165.96 ± 5.21** |
| | Lick rate (Hz) | 7.62 ± 0.37 | 7.70 ± 0.34 | 7.81 ± 0.20 |
| | Bout size (Licks) | 4.77 ± 0.28 | **#*3.64 ± 0.23** | 5.19 ± 0.16 |
| | Bout duration (s) | 0.62 ± 0.03 | **#*0.50 ± 0.03** | 0.67 ± 0.01 |
| | Inter bout interval (s) | 148.77 ± 23.21 | 211.43 ± 25.26 | 157.95 ± 20.53 |
| | Number of bouts | 7.67 ± 0.99 | 6.67 ± 0.85 | 8.73 ± 1.01 |
| | Total licks | 37.92 ± 5.97 | 25.85 ± 4.78 | 48.64 ± 6.99 |
| **8 Hz** | Inter lick interval (ms) | 168.78 ± 9.86 | **#*234.59 ± 11.70** | **#171.09 ± 6.46** |
| | Lick rate (Hz) | 8.15 ± 0.44 | #7.11 ± 0.33 | 7.78 ± 0.20 |
| | Bout size (Licks) | 4.63 ± 0.32 | **#*2.81 ± 0.19** | **#4.75 ± 0.19** |
| | Bout duration (s) | 0.58 ± 0.03 | **#*0.39 ± 0.03** | **#0.61 ± 0.02** |
| | Inter bout interval (s) | 122.57 ± 18.60 | 204.67 ± 24.27 | 146.39 ± 19.22 |
| | Number of bouts | 7.75 ± 1.05 | 5.89 ± 0.77 | 8.64 ± 1.01 |
| | Total licks | 38.33 ± 6.85 | **#18.00 ± 3.34** | 45.18 ± 6.61 |
| **15 Hz** | Inter lick interval (ms) | 156.65 ± 7.71 | **#*221.38 ± 12.27** | **#171.81 ± 5.07** |
| | Lick rate (Hz) | 8.36 ± 0.36 | 7.53 ± 0.41 | 7.70 ± 0.20 |
| | Bout size (Licks) | 4.97 ± 0.36 | **#*2.63 ± 0.28** | 4.89 ± 0.19 |
| | Bout duration (s) | 0.60 ± 0.05 | **#*0.32 ± 0.05** | **#0.64 ± 0.02** |
| | Inter bout interval (s) | 121.84 ± 16.57 | 173.98 ± 19.82 | 181.09 ± 37.38 |
| | Number of bouts | 7.92 ± 1.01 | **5.19 ± 0.74** | 8.67 ± 1.00 |
| | Total licks | 40.83 ± 6.86 | **#15.44 ± 3.30** | 46.64 ± 7.08 |
| **20 Hz** | Inter lick interval (ms) | 168.95 ± 9.95 | **#*214.47 ± 10.03** | **#166.83 ± 5.49** |
| | Lick rate (Hz) | 7.92 ± 0.41 | 8.24 ± 0.45 | 8.20 ± 0.22 |
| | Bout size (Licks) | 4.78 ± 0.22 | **#*2.43 ± 0.27** | **#4.58 ± 0.19** |
| | Bout duration (s) | 0.62 ± 0.03 | **#*0.28 ± 0.05** | **#0.57 ± 0.02** |
| | Inter bout interval (s) | 140.67 ± 25.52 | 229.92 ± 28.06 | 159.46 ± 18.53 |
| | Number of bouts | 7.58 ± 1.00 | 4.96 ± 0.77 | 8.70 ± 0.93 |
| | Total licks | 37.08 ± 5.61 | **#*13.00 ± 2.56** | 44.15 ± 6.46 |

*(Left vertical label spanning rows: Stimualtion Frequencies)*

1042

1043    Bold font indicates statistically significant differences (p<0.05):

1044     *Significant difference compared to the D2$^{tdTomato}$ (control group) at the same frequency.

1045    #Significant difference compared to the Control (no laser) frequency within the same

1046    group.

1047

**Table 3. Analysis of licking microstructure for sucrose BATT with frequency scan**

1049 **test (Fig 3).**

| | | | D2$^{tdTomato}$ | D2$^{ChR2}$ | D2$^{ArchT}$ |
|---|---|---|---|---|---|
| Stimulation pattern | No-laser | Inter lick interval (ms) | 171.05 ± 7.13 | 168.30 ± 8.88 | *150.41 ± 4.78 |
| | | Lick rate (Hz) | 4.02 ± 0.26 | 4.10 ± 0.47 | *5.04 ± 0.32 |
| | | Bout size (licks) | 15.73 ± 1.04 | 16.00 ± 1.97 | *20.00 ± 1.28 |
| | | Bout duration (s) | 2.40 ± 0.12 | 2.33 ± 0.25 | 2.82 ± 0.17 |
| | | Inter bout interval (s) | 129.94 ± 11.21 | *83.82 ± 9.69 | 178.04 ± 33.50 |
| | | Number of bouts | 8.00 ± 1.02 | *16.10 ± 2.14 | 6.80 ± 0.62 |
| | | Total licks | 124.77 ± 22.91 | *214.71 ± 23.36 | 122.36 ± 7.36 |
| | Lick paired | Inter lick interval (ms) | 174.78 ± 9.90 | 190.00 ± 8.97 | *148.58 ± 3.94 |
| | | Lick rate (Hz) | 4.13 ± 0.38 | 3.54 ± 0.32 | 4.55 ± 0.29 |
| | | Bout size (licks) | 16.04 ± 1.56 | 13.53 ± 1.43 | 18.07 ± 1.14 |
| | | Bout duration (s) | 2.37 ± 0.17 | 2.24 ± 0.22 | 2.49 ± 0.15 |
| | | Inter bout interval (s) | 152.61 ± 18.14 | *79.15 ± 9.33 | 150.76 ± 16.10 |
| | | Number of bouts | 8.68 ± 1.29 | *16.14 ± 1.89 | 7.16 ± 0.52 |
| | | Total licks | 127.09 ± 24.92 | 187.14 ± 18.81 | 121.16 ± 7.88 |
| | 8 Hz | Inter lick interval (ms) | 172.28 ± 8.50 | 187.64 ± 9.91 | *143.06 ± 3.34 |
| | | Lick rate (Hz) | 4.00 ± 0.37 | @#*2.54 ± 0.29 | *5.29 ± 0.31 |
| | | Bout size (licks) | 15.13 ± 1.62 | #*9.89 ± 1.18 | *20.88 ± 1.26 |
| | | Bout duration (s) | 2.24 ± 0.20 | @#*1.55 ± 0.17 | *2.81 ± 0.16 |
| | | Inter bout interval (s) | 129.70 ± 14.18 | *68.95 ± 8.01 | 159.22 ± 15.26 |
| | | Number of bouts | 8.82 ± 1.22 | *18.38 ± 2.15 | 6.48 ± 0.58 |
| | | Total licks | 119.91 ± 21.86 | #150.19 ± 15.43 | 125.20 ± 8.88 |
| | 20 Hz | Inter lick interval (ms) | 171.02 ± 8.56 | 180.40 ± 9.14 | *144.91 ± 4.74 |
| | | Lick rate (Hz) | 4.16 ± 0.36 | #*2.81 ± 0.40 | 5.13 ± 0.40 |
| | | Bout size (licks) | 16.44 ± 1.46 | @#*8.64 ± 1.66 | 18.80 ± 1.64 |
| | | Bout duration (s) | 2.46 ± 0.16 | @#*1.30 ± 0.28 | 2.51 ± 0.19 |
| | | Inter bout interval (s) | 142.75 ± 13.16 | *87.00 ± 9.26 | 175.73 ± 30.39 |
| | | Number of bouts | 7.36 ± 0.88 | *14.57 ± 1.58 | 6.72 ± 0.56 |
| | | Total licks | 119.64 ± 22.17 | !@#97.14 ± 14.57 | 121.04 ± 10.41 |

1050

Bold font indicates statistically significant differences ($p < 0.05$):

* Significant difference compared to the D2$^{tdTomato}$ (control group) at the same frequency.

# Significant difference compared to the Control (no laser) frequency within the same

group.

@ Significant difference between the Lick-paired condition and the 8 Hz or 20 Hz

conditions within the same group

! Significant difference between the 8 Hz and 20 Hz conditions within the same group

1058

1059 **Table 4. Analysis of licking microstructure analysis of licking for the sucrose BATT**

1060 **with varying optogenetic stimulation delay (Fig 4).**

| | | | D2$^{tdTomato}$ | D2$^{ChR2}$ | D2$^{ArchT}$ |
|---|---|---|---|---|---|
| Stimulation pattern | No-laser | Inter lick interval (ms) | 150.95 ± 5.76 | *179.48 ± 8.71 | 138.06 ± 4.97 |
| | | Lick rate (Hz) | 5.04 ± 0.36 | *3.48 ± 0.57 | 5.26 ± 0.45 |
| | | Bout size (licks) | 19.89 ± 1.50 | *13.85 ± 2.30 | 20.75 ± 1.80 |
| | | Bout duration (s) | 2.80 ± 0.22 | 2.18 ± 0.36 | 2.74 ± 0.28 |
| | | Inter bout interval (s) | 191.64 ± 21.62 | *84.71 ± 11.81 | 200.87 ± 38.04 |
| | | Number of bouts | 5.60 ± 0.72 | *15.70 ± 2.97 | 5.50 ± 1.09 |
| | | Total licks | 110.07 ± 19.52 | 181.00 ± 36.21 | 111.12 ± 21.37 |
| | 0 ms delay (lick paired) | Inter lick interval (ms) | 156.21 ± 6.95 | *206.89 ± 13.27 | 134.84 ± 5.75 |
| | | Lick rate (Hz) | 4.76 ± 0.44 | *2.68 ± 0.46 | 5.23 ± 0.64 |
| | | Bout size (licks) | 18.74 ± 1.76 | *10.53 ± 1.85 | 20.78 ± 2.55 |
| | | Bout duration (s) | 2.65 ± 0.19 | *1.81 ± 0.27 | 2.58 ± 0.26 |
| | | Inter bout interval (s) | 206.87 ± 27.54 | *79.25 ± 12.82 | 174.32 ± 20.18 |
| | | Number of bouts | 6.20 ± 0.94 | *18.20 ± 3.54 | 4.88 ± 0.74 |
| | | Total licks | 110.73 ± 19.11 | 159.10 ± 32.62 | 96.75 ± 14.28 |
| | 50 ms delay | Inter lick interval (ms) | 151.28 ± 8.05 | *190.58 ± 8.62 | 149.19 ± 5.76 |
| | | Lick rate (Hz) | 4.57 ± 0.43 | *2.88 ± 0.45 | 5.12 ± 0.75 |
| | | Bout size (licks) | 18.16 ± 1.74 | *11.36 ± 1.83 | 20.22 ± 3.03 |
| | | Bout duration (s) | 2.48 ± 0.21 | 1.87 ± 0.29 | 2.76 ± 0.34 |
| | | Inter bout interval (s) | 209.14 ± 29.89 | *82.81 ± 12.84 | 171.57 ± 35.85 |
| | | Number of bouts | 6.27 ± 0.75 | *18.00 ± 3.81 | 5.88 ± 0.79 |
| | | Total licks | 105.87 ± 16.64 | 167.70 ± 34.42 | 114.12 ± 19.19 |
| | 100ms delay | Inter lick interval (ms) | 147.26 ± 5.72 | *188.82 ± 7.84 | 139.84 ± 7.25 |
| | | Lick rate (Hz) | 4.82 ± 0.44 | *3.32 ± 0.57 | 6.39 ± 0.71 |
| | | Bout size (licks) | 18.87 ± 1.83 | 13.26 ± 2.30 | 25.14 ± 2.84 |
| | | Bout duration (s) | 2.53 ± 0.23 | 2.21 ± 0.38 | 3.25 ± 0.28 |
| | | Inter bout interval (s) | 212.17 ± 21.39 | *84.43 ± 13.25 | 148.41 ± 20.66 |
| | | Number of bouts | 5.47 ± 0.72 | *17.00 ± 3.47 | 4.88 ± 0.52 |
| | | Total licks | 97.00 ± 14.35 | *178.50 ± 32.34 | 119.88 ± 17.51 |

1061

1062    Bold indicates statistically significant differences ($p < 0.05$):

1063    * Significant difference compared to the D2$^{\text{tdTomato}}$ (control group) in the same condition

1064    (delay time).